Defining the spatial distribution of extracellular adenosine revealed a myeloid-dependent immunosuppressive microenvironment in pancreatic ductal adenocarcinoma

Vincenzo Graziano[1,2], Andreas Dannhorn[3*], Heather Hulme[3*], Kate Williamson[4*], Hannah Buckley[1], Saadia A Karim[5], Sheng Y. Lee[1], Brajesh P. Kaistha[1#], Sabita Islam[1,2,6], James E. D. Thaventhiran[4], Frances M. Richards[1°] Richard Goodwin[3], Rebecca Brais[7], Jennifer P Morton[5,8], Simon J. Dovedi[9], Alwin G. Schuller[10], Jim Eyles[9], Duncan I. Jodrell[1,6]

[1]Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge, UK

[2]Cancer Research UK Cambridge Centre, University of Cambridge, Cambridge, UK

[3]Imaging and Data Analytics, Clinical Pharmacology and Safety Sciences (CPSS), AstraZeneca, Cambridge, UK

[4]Medical Research Council Toxicology Unit, University of Cambridge, Cambridge, CB2 1QW, UK

[5]CRUK Beatson Institute, Switchback Rd, Glasgow G61 1BD, UK

[6]Department of Oncology, University of Cambridge, Cambridge UK

[7]Department of Pathology, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

[8]School of Cancer Sciences, University of Glasgow, Glasgow, UK

[9]Oncology R&D, Research and Early Development, AstraZeneca, Cambridge, UK

[10]Oncology R&D, Research and Early Development, AstraZeneca, Waltham, USA

[*] These authors contributed equally to this work

22 # Brajesh P. Kaistha current address: Oncology R&D, Research and Early Development,

23 AstraZeneca Cambridge, UK

24 ° Frances M. Richards current address: Oncology R&D, Translational Medicine, AstraZeneca,

25 Cambridge, UK

Running title

28 Extracellular adenosine shapes PDAC immune microenvironment

Keywords

30 Pancreatic cancer, adenosine pathway, immunotherapy, pro-tumorigenic macrophages, tumour

31 microenvironment

Financial support:

All CRUK CI authors received research funding from Cancer Research UK (Nos.

C14303/A17197 and C9545/A29580). The Li Ka Shing Centre where this work was performed

was generously funded by CK Hutchison Holdings Limited, the University of Cambridge,

CRUK, The Atlantic Philanthropies and others. This work was supported by Cancer Research

UK (C9685/A27444) funding for VG. This study was also funded by Cancer Research UK

Precision Panc grant C96/A25238. HB is funded by the University of Cambridge Experimental

Medicine Training Initiative programme in partnership with AstraZeneca (EMI-AZ) and SYL

is funded by the Cambridge Trust (Cambridge International Scholarship). Work by JT and KW

is funded by a core grant award from the MRC (MC_UU_00025/12). S.A.K. and. J.P.M. were

supported by Cancer Research UK core funding to the Beatson Institute (A17196 and A31287)

and to J.P.M. Lab (A29996).

Corresponding authors:

Vincenzo Graziano: Cancer Research UK Cambridge Institute, University of Cambridge, Li

Ka Shing Centre, Robinson Way, Cambridge, CB2 0RE, United Kingdom. Email:

vincenzo.graziano@cruk.cam.ac.uk

Duncan Jodrell: Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka

Shing Centre, Robinson Way, Cambridge, CB2 0RE, United Kingdom. Email:

dij21@cam.ac.uk

Authors' disclosures:

AD, HH, BPK, FMR, RG, SJD, AGS and JE are AstraZeneca employees and own company

stocks and shares. HB holds a studentship partially funded by AstraZeneca.

Synopsis: The clinical development of anti-CD73 antibodies has demonstrated their utility, but

the mechanism for this effect has not been fully elucidated.   We now show that targeting the

adenosine pathway reduces extracellular adenosine in myeloid rich, hypoxic areas of pancreatic

ductal   adenocarcinoma,   abrogating   the   pro-tumorigenic,   immunosuppressive

microenvironment and improving the efficacy of anti-tumour interventions.

Abstract

The prognosis for patients with pancreatic ductal adenocarcinoma (PDAC) remains extremely

poor. It has been suggested that the adenosine pathway contributes to the ability of PDAC to

evade the immune system and its resistance to immunotherapies (Immuno-Oncology Therapy,

IOT), by generating extracellular adenosine (eAdo).

Using genetically engineered allograft models of PDAC in syngeneic mice with differential immune infiltration and response to IOT, we showed enrichment of the adenosine pathway in tumour-infiltrating immune cells (in particular, myeloid populations). Using MS-Imaging, we showed that extracellular adenosine distribution is heterogeneous in tumours, with high concentrations in hypoxic margins that surround necrotic areas, associated with a rich myeloid infiltration, demonstrated using Imaging Mass Cytometry (IMC). Pro-tumorigenic M2 macrophages express high levels of the Adora2a receptor; particularly in the IOT resistant model. Blocking the *in vivo* formation and function of eAdo (Adoi), using a combination of anti-CD73 antibody and an Adora2a inhibitor slowed tumour growth and reduced metastatic burden. In addition, blocking the adenosine pathway improved the efficacy of combinations of cytotoxic agents or immunotherapy. Finally, Adoi remodelled the tumour microenvironment (TME), as evidenced by reduced infiltration of M2 macrophages and Tregs. RNAseq analysis showed that genes related to immune modulation, hypoxia and tumour stroma were downregulated following Adoi and a specific adenosine signature derived from this is associated with a poorer prognosis in patients with PDAC.

The formation of eAdo appears to promote the development of the immunosuppressive TME in PDAC, contributing to its resistance to conventional and novel therapies. Therefore, inhibition of the adenosine pathway may represent a strategy to modulate the stroma and improve therapy response in patients with PDAC.

Introduction

Survival for patients with pancreatic ductal adenocarcinoma (PDAC) has not changed significantly in the last 50 years and remains poor (https://www.cancerresearchuk.org/health-professional/cancer-statistics-for-the-uk). There is a need for new treatments, given that current standard of care for patients with metastatic disease is associated with poor outcomes, with less than 10% of patients living for more than 2 years [1]. In addition to relative resistance to conventional therapies, cancer immunotherapy (Immuno-Oncology Therapy, IOT) is also ineffective in this disease, except in a small group of patients (1-2%) with microsatellite instability/mismatch repair deficient (MSI-H/dMMR) tumours [2]. Several authors consider that the reason for this resistance can be ascribed to the low mutational burden of this neoplasm, which leads to lymphocyte exclusion and anergy [3, 4]. However, the tumour microenvironment in PDAC has also been shown to be populated by a rich variety of immune cells, most of which demonstrate immune suppressive features, which contribute to the resistance to immunotherapy [5].

The adenosine pathway is an immunosuppressive axis which has gained much attention in cancer immunology for its role in suppressing the immune activation associated with cytotoxic treatments (chemotherapy, targeted therapy and radiotherapy) [6]. This has led to the clinical evaluation of inhibitors of the pathway in combination with more conventional approaches [7]. The adenosine pathway involves conversion of extracellular ATP (eATP), a powerful immune activator, to extracellular adenosine (eAdo) by the ectonucleotidases CD39 and CD73 [8]. eADO has been linked to cancer in several studies that have demonstrated that its concentration in different tumour tissues is several folds higher than in normal tissues [6, 9]. CD39 is overexpressed in a subpopulation of exhausted tumour-infiltrating T-cells [10, 11] and its expression correlates with another marker of immunosuppression (PD1 expression) [11]. CD39 and CD73 have roles in the aggressiveness of adult glioblastoma [12], where they are expressed

on infiltrating macrophages [13, 14]. The adenosine signature recently published by Sidders and

colleagues [15] shows that this pathway correlates with resistance to immunotherapies and is

associated with other genetic features of tumour aggressiveness, such as p53 mutations. The

abundant presence of eAdo in the microenvironment can dampen immune activation through

the stimulation of a pro-tumorigenic stroma. This is mostly orchestrated by macrophages and

myeloid derived suppressive cells (MDSCs) [16, 17], favouring a tolerogenic function of dendritic

cells (DCs) [18, 19, 20] which results in inhibition of T-cells/NK cell activation [21, 22].

The myeloid populations play a pivotal role in the aggressiveness of many cancer types

and in particular, PDAC. For instance, the presence of pro-tumorigenic populations of

macrophages [23, 24] and myeloid-derived suppressive cells (MDSCs) [25] infiltrating the

microenvironment, is associated with poor survival and correlates with immune exclusion of

PDAC. Macrophages can elicit the secretion of cytokines which can on the one hand, favour

the proliferation and invasiveness of cancer cells while interacting with cancer-associated

fibroblasts [26, 27, 28], and on the other hand induce anergy and physical exclusion of adaptive

immune cells [29, 30, 31]. Targeting macrophages in a pre-clinical pancreatic cancer model has

been demonstrated to be effective to obtain tumour regression and reduce metastatic formation

[32]. Unfortunately, this approach has not translated into clinical benefit, which in part can be

explained by the fact that the global reduction of the tumour-infiltrating macrophages can be

biologically different from reprogramming distinct tumour associated macrophage (TAM)

subtypes [28].

Some recent publications link the adenosine pathway to the biology and aggressiveness

of PDAC. PDAC has been shown to have an increased adenosine pathway RNA signature

associated with a worse prognosis [15], and genes encoding for the receptors for eAdo as well as

CD73 have been found to be overexpressed in bulk-RNA sequencing (RNAseq) when

comparing tumours to normal pancreatic tissue [33]. However, little is known about the complex

mechanism generated by the adenosine pathway resulting in the immunosuppressive

characteristics of pancreatic cancer microenvironment and stroma, in particular the role that

the adenosine pathway has in shaping the immune infiltration of this disease.

Here, we propose a model where the tumour-infiltrating immune cell populations of

PDAC generate an axis of immunosuppression, where extracellular adenosine produced mostly

in hypoxic regions of the tumour (identified using Mass Spectrometry Imaging), enriched for

the myeloid cell infiltration, stimulates pro-tumorigenic M2 macrophages. The axis described

is expressed preferentially by the IOT-resistant model when compared to the IOT-responsive

one. Therefore, blocking the adenosine pathway in the IOT-resistant PDAC model, strongly

suppresses the formation of extracellular adenosine and reshapes the immune

microenvironment, favouring disease control when combined with cytotoxic treatments and

immunotherapies. Bulk RNAseq gene analysis confirms the role of the myeloid-dependent

adenosine pathway in PDAC survival, underpinning the importance of our results for

understanding the biological complexity and the clinical utility of the adenosine pathway

inhibition.

Methods and materials

Cell lines and chemicals

$Kras^{LSL-G12D/+}$; $Trp53^{LSL-R172H/+}$; $Pdx1$-Cre; $Rosa26^{YFP/YFP}$ (KPCY)-derived cell lines

2838c3, 6499c4, 6620c1 (IOT-responsive), 6419c5, 6694c2 and 6422c1 (IOT-resistant) were

a kind gift from Ben Stanger (University of Pennsylvania). The cell lines were obtained from

single cell cloning strategy, as described previously, and were generated from tumours

developed in KPCY mice on a C57BL/6 background [34]. Cells were grown up to 20 passages

in DMEM (with pyruvate, L-glutamine and D-glucose; Gibco, #41966029) supplemented with

5% FBS (Gibco, #10270106). All the cell lines were analysed for STR fingerprinting and

tested for mycoplasma routinely.

Mice and *in vivo* experiments

Tumour allograft experiments were performed in the animal facility (Biological

Resource Unit, BRU) of the CRUK Cambridge Institute, in accordance with the UK Animals

Scientific Procedures Act 1986, with approval from the CRUK Cambridge Institute Animal

Ethical Review and Welfare Body. 8-12 week old female C57BL/6 mice were used for *in vivo*

experiments and were purchased from Charles River (UK).

Tumour allograft studies were performed with technical assistance from CRUK-CI BRU staff.

Mice were subcutaneously injected in the right flank with 1×10^6 KPCY-derived cells in 50%

PBS and 50% Matrigel basement membrane matrix (#354234, Corning). In the interventional

experiments, mice were treated as indicated, starting 12-14 days from tumour cell implantation,

to allow the microenvironment to establish. Tumour volume was calculated using the formula;

$(\pi/6)*(\text{width})^2*\text{length}$. Tumour response was defined based on the % of change of the longest

diameter from start of therapy (stable disease < 20% increase and < 30% decrease of target

lesion RECIST v. 1.1). Mice were then killed at specific endpoints (e.g. 14 days from start of

treatment) or when the tumour reached 2000 mm^3 (or before in case of appearance of clinical

signs).

Kras$^{\text{LSL-G12D/+}}$; *Trp53*$^{\text{LSL-R172H/+}}$; *Pdx1*-Cre (KPC) mice for tumour phenotyping, were obtained

from a breeding colony maintained by the CRUK-CI Genome Editing Core team. Tumours

were detected by palpation followed by ultrasound imaging by the Genome Editing Core.

Tumour tissues, spleens and mesenteric and inguinal lymph nodes from KPC mice were

provided once tumour dimensions or health status rendered them unsuitable for therapeutic

studies. KPC mice were killed when showing clinical signs of the disease (swollen abdomen,

loss of body conditioning resembling cachexia, reduced mobility).

KPC mice used for interventional study were bred at the CRUK Beatson Institute and

maintained on a mixed background. All work was performed under UK Home Office license

and approved by the University of Glasgow Animal Welfare and Ethical Review Board. Mice

of both sexes, in similar proportions, were used in all cohorts. Mice suspected to have PDAC

following palpation were anesthetised in 0.2L/min medical air and isoflurane and then

underwent 3D ultrasound imaging using the VisualSonics Vevo 3100 ultrasound system with

MX550D 40μm resolution transducer (FujiFilm). Mice were randomly assigned to 1 of the 4

treatment groups (A: vehicles + isotype; B: AZD6738 + gemcitabine; C: AZD4635 +

2c5mIgG1; D: AZD6738 + gemcitabine + AZD4635 + 2c5mIgG1) once tumours were

confirmed by imaging, and follow-up scans were performed weekly until endpoint was

reached. Schedule of the treatments are specified in the results and figures sections. Mice were

culled when exhibiting moderate symptoms of pancreatic cancer (see above). Statistical

assessment of survival was carried out by Kaplan-Meier and Log-Rank analysis. Analysis of

ultrasound images was performed using VevoLab software (version 3.1.1) from VisualSonics.

In 3D mode, stacked images of the tumour were imported and the tumour border annotated,

allowing a 3D construct to be formed.

When indicated the following drugs were used: AZD6738 (ATRi; 25 mg/kg daily for 4 days),

AZD4635 (Adora2ai; 50 mg/kg bid), 2c5mIgG1 (anti-CD73; 10 mg/kg twice weekly),

AB740080 D265A (anti-PD-L1: 10 mg/kg twice weekly), NIP228 mouse IgG1 control kappa

(isotype; 10 mg/kg twice weekly) and NIP228 muIgG1 D265A (isotype; 10 mg/kg twice

weekly) were provided by AstraZeneca; gemcitabine hydrochloride (Tocris, 3259) was used at

100 mg/kg twice weekly; inVivoPlus anti-CD40 (clone FGK4.5/FGK45; bioxcell BE0016-2)

and InVivoPlus rat IgG2a isotype control, anti-trinitrophenol (clone 2A3; bioxcell BE0089)

were used as a single injection of 100 µg. InVivoPlus anti-CTLA-4 (clone 9H10; bioxcell

BP0131) or InVivoPlus isotype control polyclonal Syrian hamster IgG (bioxcell BP0087): 200

µg/dose x 3 times.

Clonogenic assay

KPCY-derived cells were seeded at 200 cells/well in a 6-well plate. After 24 hours,

cells were treated with different concentrations of anti-CD73 (2c5mIgG1) antibody (1, 10, 100

µg/ml) or isotype (NIP228 mouse IgG1 control kappa) for 8 days. Antibodies were added every

3 days. On day 8, colonies were stained with SRB protocol previously described (36). Images

were taken and analysed using GelCount (Oxford Optronix). Colony forming efficiency was

calculated as a ratio between the number of colonies and number of plated cells. Surviving

fractions were calculated as the ratio between wells treated with anti-CD73 antibody and the

ones treated with isotype. At least 3 wells per condition were plated for each of the 3 replicates

per experiment.

IncuCyte time lapse imaging

KPCY-derived cells were plated at 2500 cells/well density in a 96-well black-wall plate

(at least 3 wells per condition). Cells were grown in cell culture medium supplemented with

the indicated concentration of anti-CD73 (2c5mIgG1) or isotype (NIP228 mouse IgG1 control

kappa) antibody in triplicate. Images were acquired with 10x objective, every 3 hours from 3

different fields per well using Incucyte Live cells imaging microscope (Essen Bioscience).

Confluence was calculated as the average of the 3 fields using the Incucyte algorithm.

Experiments were repeated at least 3 times.

Single cell suspension preparation

For experiments in subcutaneous allografts, tumours were weighed and placed in RPMI

and finely minced with scissors in a 2 ml tube, which was then washed with up to 2.5 ml of

digestion buffer (Tumour dissociation kit, Miltenyi, 130-096-730) plus Deoxyribonuclease I

(300 µg/ml, Sigma, DN25-1G). Dissociation was performed using the protocol suggested by

Miltenyi. For KPC tumours, a trypsin inhibitor (250 µg/ml, Sigma, T6522) was added to the

digestion buffer. Following the digestion, the samples were passed through a 70 µM strainer

filter (Grainer Bio-one, 542-070), washing with MACS buffer (PBS + 0.1% FBS and 2nM

EDTA).

Mouse spleens, inguinal and mesenteric lymph nodes were mashed on a 100 µM filter

(Grainer Bio-one, 542-000) over a 50 ml tube, using a syringe plug and the filter was washed

with MACS buffer and centrifuged (at 4° C as for all the following centrifugation). Red cell

lysis buffer (1 ml; 0.15M Ammonium Chloride; 10mM Potassium hydrogen carbonate; 0.1mM

EDTA; pH 7.4 (adjusted with KOH) was then used to resuspend splenocytes, 3 minutes at room

temperature and then washed with MACS buffer and centrifuged at 300g for 5 minutes.

Samples were eventually resuspended in 200-400 µl of MACS buffer.

Flow cytometry

Single cell suspensions were aliquoted in a round-bottom 96 well/plate (Costar, 3879)

and stained with live/dead fixable stain (Invitrogen, L34962; 1:100 in PBS) for 10 minutes at

room temperature. After washing in MACS buffer, cells were FC-blocked with anti-CD16/32

antibody (BioLegend Cat# 101320, RRID:AB_1574975; 1:100) for 5 minutes. Then,

antibodies for surface staining were added and incubated at 4° C degrees for 30 minutes. After

washing, cells were fixed with FACS fix buffer [PBS + 1% Formaldehyde + 0.02 g/ml Glucose

+ 0.02% Sodium Azide] for 10 minutes and then washed and resuspended in MACS buffer for

FACS analysis. For intracellular staining, cells were fixed with fixation/permeabilization

buffer (Invitrogen 00-5123-43 and 00-5223-56) for 15 minutes, then washed in perm buffer

and stained with the relevant antibody in perm buffer for 60 minutes. Cells were then washed

and resuspended in MACS buffer for FACS analysis. Samples were acquired using BD-Symphony flow cytometer and the generated FCS files were analysed using FlowJo V10 software (RRID:SCR_008520). The following antibodies were used and gating strategies are shown in the supplementary figures(myeloid populations and lymphoid population; suppl. fig. 1A and 1B respectively): BV786-CD45 (BD Biosciences Cat# 564225, RRID:AB_2716861; 1:200), APC/Fire750-CD3 (BioLegend Cat# 100248, RRID:AB_2572118; 1:50), BV650-CD8 (BioLegend Cat# 100741, RRID:AB_11124344; 1:100), BV711-CD4 (BioLegend Cat# 100549, RRID:AB_11219396; 1:200), APC-Foxp3 (Invitrogen Cat# 17-5773-82, RRID:AB_469457; 1:100) , FITC-CD19 (BioLegend Cat# 115505, RRID:AB_313640; 1:200) , BV510-CD11b (BioLegend Cat# 101245, RRID:AB_2561390; 1:200), PerCP/Cy5.5-CD44 (BioLegend Cat# 103031, RRID:AB_2076206; 1:200), BV421-PD1 (BioLegend Cat# 135218, RRID:AB_2561447; 1:100), BV421-PD-L1 (BioLegend Cat# 124315, RRID:AB_10897097; 1:100), PE/Cy7-CD39 (BioLegend Cat# 143805, RRID:AB_2563393; 1:100), PE-CD73 (BioLegend Cat# 127206, RRID:AB_2154094; 1:100), FITC-F4/80 (BioLegend Cat# 123108, RRID:AB_893502; 1:200), PE/Cy7-CD206 (BioLegend Cat# 141719, RRID:AB_2562247; 1:100), PerCP/Cy5.5-Ly6C (BioLegend Cat# 128012, RRID:AB_1659241; 1:100), APC/Cy7-Ly6G (BioLegend Cat# 127624, RRID:AB_10640819; 1:100), AF700-MHCII (BioLegend Cat# 107629, RRID:AB_2290801; 1:200), PE-CD11c (eBioscience Cat# 12-0114-83, RRID:AB_465553; 1:200) , BV605-NKp46 (BioLegend Cat# 137619, RRID:AB_2562452; 1:25), APC-Adora2a (Novus Biotech Cat# NBP1-39474APC; 1:150) , APC-CD86 (BioLegend Cat# 105012, RRID:AB_493342; 1:100). Gating strategy for immune subpopulations is shown in supplementary figure 1A-B. For CD73 *in vitro* staining of KPCY-derived cell lines, cells were stained as described above with live/dead fixable stain and PE-CD73 antibody and analysed with BD-Symphony flow cytometer. For in vitro treatment, cells were treated with anti-CD73 (2c5mIgG1) or isotype (NIP228) antibody at a concentration of 10 µg/ml for 24

hours. Competitive staining was performed before this experiment to confirm there was no

competition between 2c5mIgG1 and TY11.8 (PE-CD73) clones (data not shown).

Tissue preparation for Mass Spectrometry Imaging (MSI) and Imaging Mass Cytometry

(IMC) analysis.

PDAC mouse tumours were snap frozen in liquid nitrogen immediately after resection

and the tissues were embedded in a HPMC/PVP hydrogel as previously described [35].

Sectioning was performed on a CM3050 S cryostat (Leica Biosystems, Nussloch, Germany) at

a section thickness of 10 µm and the tissue sections were immediately thaw mounted and dried

under a stream of nitrogen and sealed in vacuum pouches to preserve the metabolic integrity

of the sections. Tissue sections for DESI-MSI and IMC were thaw-mounted onto Superfrost

microscope slides (Thermo Scientific Waltham, MA, USA), whilst sections prepared for

MALDI-MSI were thaw mounted onto conductive ITO coated slides (Bruker Daltonik,

Bremen, Germany). Polyvinylpyrrolidone (PVP) (MW 360 kDa) and (Hydroxypropyl)-

methylcellulose (HPMC) (viscosity 40-60 cP, 2 % in H2O (20 C) were purchased from Merck

(Darmstadt, Germany). Methanol, water, iso-pentane and isopropyl alcohol were obtained from

Fisher Scientific (Waltham, MA, USA).

Mass Spectrometry Imaging (MSI)

DESI-MSI analysis was performed on a Q-Exactive mass spectrometer (Thermo

Scientific, Bremen, Germany) equipped with an automated 2D-DESI ion source (Prosolia Inc.,

Indianapolis, IN, USA) operated in negative ion mode, covering the applicable mass range up

to a m/z of 1000, with a nominal mass resolution of 70,000. The injection time was fixed to

150 ms resulting in a scan rate of 3.8 pixel/s. The spatial resolution was adapted between

experiments to allow acquisition of the data for all directly compared samples within a single

experiment of 48 h, with pixel sizes ranging from 65-75 µm. A home-built Swagelok DESI

sprayer was operated with a mixture of 95% methanol, 5% water delivered with a flow rate of

1.5 µL/min and nebulized with nitrogen at a backpressure of 6 bar. The resulting .raw files

were converted into .mzML files using ProteoWizard msConvert [36] (version 3.0.4043) and

subsequently compiled to an .imzML file (imzML converter [37] version 1.3). All subsequent

data processing was performed in SCiLS Lab (version 2021b, Bruker Daltonik, Bremen,

Germany).

MALDI-MSI analysis was performed on a RapifleX Tissuetyper instrument (Bruker Daltonik,

Bremen, Germany) operated in negative detection mode. 9-Aminoacridine (9-AA) prepared in

80:20 methanol:water was used as a MALDI matrix and spray deposited using an automated

spray system (M3-Sprayer, HTX technologies, Chapel Hill, NC, USA). MALDI experiments

were performed with a spatial resolution of 50 µm. A total of 400 laser shots were summed up

per pixel to give the final spectra. For all experiments the laser was operated with a repetition

rate of 10 kHz. All raw data were directly uploaded and processed in SCiLS lab (Version

2021b) software packages. All DESI and MALDI data and images were normalised to the total

ion current (TIC) to compensate for signal variation across the course of the experiments. Data

segmentation pipeline is shown in supplementary materials and methods.

Imaging Mass Cytometry (IMC)

Imaging mass cytometry was performed on a slide which had already been analysed by DESI-

MSI. Antibodies used for IMC staining are shown in Table 1. Untagged antibodies were tagged

in house, using Fluidigm Maxpar Antibody Labelling Kit, according to manufacturer's

instructions. Following DESI-MSI analysis, the slide was fixed with 4% paraformaldehyde in

phosphate-buffered saline (PBS) for 10 minutes. The slide was washed 3 x 5 minutes in PBS,

permeabilized for 5 minutes with 1:1000 dilution of Triton X-100 in Casein Solution, washed

3 x 5 minutes in PBS, and blocked for 30 minutes with Casein Solution. Antibodies were

diluted to an appropriate concentration in Casein Solution and the slide incubated overnight

with the antibody solution at 4 ˚C. The slide was washed 3 x 5 minutes in PBS and nuclei were

stained with DNA Intercalator-iridium at a dilution of 1:400 in PBS for 30 minutes. The slide

was washed 3 x 5 minutes in PBS, 30 seconds in deionized water, then dried for storage at

room temperature until analysis. A region for IMC analysis was selected using consecutive H

and E stained sections and the DESI-MSI results. A box with approximately 2 x 1.8 mm area

was selected for analysis to include necrotic, necrotic margin and viable tumour regions. IMC

analysis was performed using a Hyperion instrument (Fluidigm Corporation, San Francisco,

CA, USA) with an ablation energy of 6 db an ablation frequency of 200 Hz. IMC images were

produced using MCD viewer (Version 1.0, Fluidigm) and analysis was performed using HALO

(Indica labs). Tissue regions were classified using random forest with all markers included.

Cells positive for each marker were manually optimised by setting a cell intensity threshold.

Values for the numbers of positive cells for markers of interest were exported for analysis in

Graphpad Prism v.8 (RRID:SCR_002798)

Immunohistochemistry (IHC)

Immunohistochemistry was performed as previously described[38] in the histopathology

core at the CRUK CI. Briefly, tissues were removed from the mouse at the endpoint and

immediately formalin-fixed for 24 hours. Fixed tissues were then processed, embedded in

paraffin and sectioned (3 µm sections). Following dewaxing and rehydration, as standard,

antigen retrieval was performed using Leica's Epitope Retrieval Solution 2 (Tris EDTA) at

100° C for 20 minutes. Additional protein block from Dako (X090930-2) was applied. The

staining using anti-mouse CD8 (Cell signalling, #98941), anti-mouse Foxp3 (Affymetrix. #14-

5773) and anti-mouse p53 (Novocastra; #NCL-L-p53-CM5p) antibodies, was performed on

Leica's automated Bond-III platform in conjunction with their Polymer Refine Detection

System (DS9800) and a modified version of their standard template. Slides were dehydrated

and cleared in xylene on Leica's automated ST5020 before sections were mounted on Leica's

coverslipper, CV5030 (mounting media: DPX Mountant for Histology; Sigma Aldrich, 06522-

500ML) and scanned using a ScanScopeAT2 (Aperio Leica Biosystems). Quantification of

viable tumour tissue was performed after exclusion of necrotic area using the Halo software v.

3.3.2541.405 (Indica Labs). Cell density was calculated as the number of positive cells x mm2

of tumour tissue analysed. Sections of mouse spleen were used on each slide as internal control.

For the analysis of the lung metastatic burden of any individual mouse, the 4 right lobes and

the left lobe were cut into multiple pieces and together fixed and then embedded, then treated

as above. p53 staining was used for helping the detection of smaller lesions (min. of 5 cells).

Analysis was performed using Halo software and expressed as % of metastatic areas/total lung

area analysed. Mice with intra-abdominal/thoracic organs direct infiltration were excluded

from the analysis.

RNA sequencing

RNA was extracted from twelve subcutaneous allograft tumour tissues (6 mice of the

vehicle + isotype group or control and 6 mice of the AZD4635 + 2c5mIgG1 group or Adoi)

weighing up to 30 mg. Tissues were firstly disrupted and homogenised using TissueLyser II

and then RNA was extracted using a Qiagen RNAeasy kit, according to the manufacturer's

instructions. RNA was then quantified using a Qubit 3.0 (life technologies) and purity and

quality were assessed using an Agilent 4150 (G2992AA) TapeStation system (Agilent).

Library construction was followed by paired-end 50 bp sequencing on a Novaseq 6000

sequencer.

Bioinformatics analysis

Sequencing files in FASTQ format were aligned against the GRCh38 mouse genome

using HISAT2 (RRID:SCR_015530) with default parameters. Samtools (RRID:SCR_002105)

was used to create, index and merge BAM files of reads from different lanes belonging to

individual samples. FeatureCounts (RRID:SCR_012919) was utilized to quantify gene-level

expression of transcripts. All downstream analyses were completed in R version 4.1.2. Prior to

analysis, MSI data for sequenced samples were examined. From the vehicle/isotype-treated

arm (control group), sample 23729 showed minimal necrosis, low peri-necrotic adenosine and

a high ATP/AMP ratio suggesting a very high energetic state. This identified the sample as a

potential outlier which was confirmed upon visual inspection of a PCA plot (Supplementary

fig. 1C). It was excluded prior to downstream analysis.

For the remaining 11 samples, initially genes were filtered to maintain only genes that were

expressed at a reasonable level in >5 treatment conditions using the filterByExpr() command

from the edgeR package (ver 3.36.0) .

Differential gene expression analyses were performed on raw read counts of the combined data

object of all 11 samples. To identify significantly expressed genes between Adoi and control

groups, we utilised a Wald test within the DESeq2 package (RRID:SCR_000154, ver 1.34.0).

Genes were considered differentially expressed when the analysis resulted in an adjusted P-

value (corrected for multiple testing using the Benjamini and Hochberg method) below 0.05.

The volcano plot was generated using the EnhancedVolcano package (ver 1.12.0) with the

addition of custom code.

Gene set enrichment analysis (GSEA) of 712 genes identified as differentially expressed with

padj≤0.05 and log2(fold change)≤-0.58 was performed via the Enrichr (RRID:SCR_001575)

(suppl. table 2) server database for Kyoto Encyclopedia of Genes and Genomes (KEGG)

(https://www.genome.jp/kegg/) pathways and Gene Ontology (GO): Biological Processes

(http://geneontology.org/). Subsequently, enriched terms ranked for significance for each

database were downloaded and are reported in Supplementary Table 4-5. Terms of interest

were selected from the top 15 ranks in each table. Genes from this study which were shown to

be enriched in these terms of interest were then selected to be displayed in a heatmap. Raw

counts were normalised with DESeq2 (RRID:SCR_000154) prior to visualisation of gene

expression levels with pheatmap (ver 1.0.12). Please refer to the supplementary methods and

materials references section for all of the above.

Analysis of human PDAC available datasets and generation of PDAC-specific adenosine

signature

In order to the evaluate the correlation of the adenosine-related gene expression profile

to human PDAC we analysed 712 genes which had at least a 50% decreased expression

(Log2FC <-0.58) following adenosine inhibition treatment, of which 561 had a human ortholog

(suppl. table 2-3).

For the analysis of the adenosine-related gene expression in Bailey [39] PDAC subtypes (ADEX,

Immunogenic, Squamous and Pancreatic progenitor) we derived z-scores for the 517 genes

analysed in the dataset (out of the 561 genes) for the 97 patients with RNAseq data and subtype

information (https://www.cbioportal.org/, RRID:SCR_014555). The z-score of all genes were

summed per patient and the total number represented as the adenosine pathway gene score as

previously shown [15] and in supplementary materials and methods.

For the generation of a PDAC-specific adenosine signature and application of this to PDAC

survival, from the list of 561 human ortholog genes, we manually curated the ones related

without ambiguity to the major biological processes implicated in PDAC pathogenesis and

indicated by pathway analysis (hypoxia, immunity and extracellular matrix organisation). Of

these genes, only those that correlated positively or negatively to survival in PDAC

(https://kmplot.com) and were significantly co-expressed with CD73 and/or Adora2a in public

datasets (Bailey et al. or TCGA) were selected. A final list of 52 genes was analysed (suppl.

table 6).

Using PDAC specific data from TCGA [40] available in https://www.cbioportal.org/, we derived

the z-score of these 52 genes for each patient with known disease-specific-survival (DSS),

Progression free survival (PFS) and disease-free-survival (DFS). The z-scores for all genes

were summed up for each patient and was deemed high adenosine score if >0 or low adenosine

score if <0, as previously shown [15].

Statistics

Graphpad Prism v.8 (RRID:SCR_002798) was used for statistical analyses. Analysis

and comparisons of two groups was performed with two-tailed unpaired Student's T-test when

assuming Gaussian distribution or Mann Whitney test. Analysis of three or more groups was

performed with one-way ANOVA with Tukey's multiple comparisons post-test analysis unless

otherwise specified. Kaplan-Meier analysis with log-rank Mantel-Cox test was used to evaluate

difference in survivals. Differences were considered significant when $p < 0.05$.

Data Availability Statement

442 The data generated in this study are available within the article and its supplementary data files

443 or from the corresponding authors upon reasonable request. Code for differential expression

444 analysis and visualisation of RNAseq data is available via Github https://github.com/ka-

445 lw/AdenoPDAC .

Results

Adenosine pathway expression on KPCY-derived cell lines

Human PDAC cells have been shown to express for CD73 expression and demonstrate

sensitivity, although weakly, to the targeting of CD73 *in vitro* [41]. For this reason, we sought to

investigate whether murine KPCY-derived cell lines (which are associated with contrasting

ability to generate IOT- resistant or responsive tumours when re-implanted in syngeneic mice)

express the proteins of the canonical adenosine pathway (CD39, CD73 and Adora2a). We

found that, as in human cells, mouse PDAC cell lines express CD73 (from 72% to 99% of cells;

fig. 1A and suppl. fig. 2A) but demonstrate negligible or no expression of CD39 and Adora2a

(suppl. fig. 2B-C). Exposing cells to an anti-CD73 antibody (2c5mIgG1) reduced significantly

the detection of CD73 in all the cells after only 24-hour treatment ($p<0.05$ in all cell lines, fig.

1B), but this did not translate into inhibition of cell growth after short or long exposure at high

concentrations. In addition, confluency experiments showed that the treatment did not affect

the proliferation of any of the cell lines over a period of 72 hours (fig. 1C and suppl. fig. 2D-

E), and colony-forming experiments performed on 2838c3 (IOT-responsive) and 6419c5 (IOT-

resistant) demonstrated no differences in terms of number or size of the colonies formed after

8 days of continuous treatment (fig. 1D-E and suppl. fig. 2F). In order to evaluate whether a

direct effect of anti-CD73 exposure affects cell proliferation, reducing adenosine formation,

we cultured KPCY-derived cell lines with increasing concentrations of AMP and 5'-N-

(Ethylcarboxamido)adenosine (NECA, a stable form of adenosine). We were able to

demonstrate that adenosine and AMP have no effect on the proliferation capacity of these cell

lines (suppl. fig. 2G-H), corroborating the hypothesis of a non-cancer cell direct effect of anti-

CD73 therapeutic approaches.

Comparison between KPCY-derived cell lines allograft and KPC autochthonous tumours

Despite the lack of activity in our cell line experiments, preliminary reports from clinical

trials suggest activity in patients with PDAC treated with an anti-CD73 antibody (oleclumab)

[42]. As we hypothesised that this might be the result of impact on the tumour microenvironment

(TME), we investigated the expression of this pathway on the tumour-infiltrating immune cells

for different murine PDAC models, including KPCY-derived cell line allografts (with

differential response to IOT) and autochthonous KPC tumours. In order to understand the

complexity and similarities of the immune system in these models we first compared the

immune infiltration of the cell line allografts to the KPC model.

The immunosuppressive characteristics identified in autochthonous KPC tumours, appear

to be more aligned with those of the IOT-resistant model. In particular regarding lymphocyte

populations, KPC tumours are usually infiltrated by a low number of $CD8^+$ T cells (mean

number of $CD45^+$ cells for 2838c3 is 6.9%, 0.9% for 6419c5 and 1.9% for KPC tumours) which

bear fewer features of activation/exhaustion (mean 70% vs 13% vs 10%) as shown in suppl.

figure 3A and have a similar total $CD4^+$ T-cells (suppl. fig. 3C) and regulatory T-cell (Tregs)

infiltration (mean 3,2% vs 1% vs 0.7%, suppl. fig 3B) to IOT-resistant tumours. Moreover,

KPC tumours showed greater heterogeneity regarding myeloid infiltrating populations (suppl.

fig 3D-H). These results suggest that our IOT- resistant and responsive models stand out as the

extreme clonotypes which can arise from the complex and heterogeneous biology found in

KPC autochthonous tumours.

The adenosine pathway is enriched in immune cells infiltrating PDAC models

We hypothesized that the adenosine pathway might have a more impactful role in the

TME, as opposed to a cell autonomous effect. For this reason, we investigated the expression

of the adenosine pathway components on tumour-infiltrating immune cells which represent

significant proportion cells seen in PDAC lesions. We showed a highly significant enrichment

in both the IOT-resistant and IOT-responsive models for $CD39^+CD73^+$ double-expressing

immune cells, when compared to secondary lymphoid organs (spleen and nodes). In particular,

the majority (65-91% in tumour vs 36-56% in spleen) of tumour-infiltrating $CD11b^+$ myeloid

cells express the two receptors, due to an increase in expression of CD73, given that those cells

are normally $CD39^+$ (fig. 2A-B). Similar results were shown for Tregs and $CD8^+$ T-cells,

which are normally $CD73^+$ and displayed an increase in expression of CD39 in tumour,

compared to the secondary lymphoid organs counterparts (suppl. fig. 4A; p<0.05). There was

no significant difference in these findings when comparing the two models, despite their

differential response to IOT. We then confirmed these findings in KPC autochthonous tumours

and found similar results, with a significant increase of CD39-CD73 double expressing CD11b$^+$

myeloid cells infiltrating the tumours compared to spleens (mean 77% vs 36% p<0.0001, fig.

2C-D), harvested from the same mice. Of note, 4 KPC mice had synchronous metastases (3

liver and 1 spleen), and in 3 of these, myeloid cells infiltrating the metastases were also

enriched for CD73$^+$CD39$^+$ double expression (suppl. fig. 4 C) when compared to secondary

lymphoid organs. Significant increased co-expression was also observed for Tregs and CD8$^+$

T-cells in KPC tumours when compared to mesenteric or inguinal lymph nodes (suppl. fig.

4B). An increased percentage of CD39/CD73 double expressing CD8 T-cells and Tregs was

noted in the spleens (suppl. fig. 4B) if compared to what was found in non-tumour bearing

mice (not shown) or subcutaneous tumour bearing mice, suggesting trafficking of immune

lymphoid populations between primary lesions and closer lymphoid organs.

Adenosine distributes primarily in the areas surrounding necrosis

Given the enrichment of the adenosine pathway in the TME of PDAC models, we

anticipated that extracellular adenosine might have been abundant in the tumour

microenvironment. Using Mass Spectrometry Imaging (MSI) we evaluated the presence and

the distribution of the purinergic system in the TME of IOT-resistant tumours. The tissue

classification and segmentation approaches were driven by tissue-defining metabolic pattern.

Areas characterised by a high energetic state defined by a high abundance of ATP and ADP

and a low abundance of depleted high energy phosphates such as AMP, were called *viable*

tumour. In contrast, areas of tumour adjacent to necrosis (termed *necrotic margin)* were

characterised by high abundance of lactate, products of ribonucleotide catabolism (i.e. xanthine

and hypoxanthine) and other metabolites associated with tissue hypoxia and an overall energy-

deprived state (fig. 3A-B). We found that adenosine is present in high concentrations in the

microenvironment of IOT-resistant tumours, although showing a heterogenous distribution,

with high abundance in the necrotic margin areas (fig. 3A,C). When we investigated the cell

population distribution in the different areas using Imaging Mass Cytometry (IMC) in IOT-

resistant tumours, we noted in the necrotic margin areas a 2.7-fold increase in the number of

infiltrating CD11b[+] myeloid cells (mean 1970 in the *necrotic margin* vs 730 in *viable tumour*

CD11b[+] cells), that led to a significant decrease of the ratio between cancer cells/myeloid cells

(fig. 3D). This again suggests that myeloid cells have an instrumental role in the generation of

adenosine in this aggressive model of PDAC.

Expression of Adora2a receptor on myeloid subpopulations of pancreatic cancer models

Having shown in these models that in the PDAC microenvironment, immunosuppressive

adenosine is present abundantly, we then investigated which cells within the microenvironment

might be responsive to this. We investigated the expression of the adenosine A2a receptor

(Adora2a, the receptor with the highest affinity for adenosine) that has been found to be

frequently overexpressed in human tumours, by flow cytometry We found that Adora2a was

highly expressed by tumour-infiltrating myeloid population when compared to the spleen

(suppl. fig. 4D-E) and this expression was significantly higher in the IOT-resistant model in

terms of MFI (10000 vs 6700, p<0.0001) and % of Adora2a[+] myeloid cells (15% vs 11%,

p=0.009). When comparing different subpopulations, pro-tumorigenic M2 macrophages,

infiltrating both IOT-resistant and IOT-responsive PDAC showed high positivity for the

receptor. The IOT-resistant model had higher expression of Adora2a compared to the IOT-

responsive model (fig. 3E-F; p<0.0001) and percentage of Adora2a[+] M2 positivity [72% vs

43%; p<0.0001) (fig. 3I)]. Once more, these findings were confirmed in KPC tumours where

Adora2a was found to be increased in M2 macrophages infiltrating the lesions when compared

to matched spleens (fig. 3G-H). The KPC model demonstrated once again the heterogeneity of

pancreatic lesions, which in terms of M2 macrophages, positive for the Adora2a receptor,

covers the entire range of expression seen in the two subcutaneous models used (fig. 3I).

Notably, of three KPC mice where metastatic nodules were found, Adora2a expression was

found retained in the M2 macrophages infiltrating the secondary lesions (suppl. fig. 4F).

In addition to pro-tumorigenic macrophages, Adora2a expression was found enriched in

other myeloid immune populations infiltrating the tumours. In particular CD11b$^-$ dendritic

cells, CD11b$^+$ dendritic cells (suppl. fig. 4G), M1 macrophages (suppl. fig. 4H), and mo-

MDSCs (suppl. fig. 4I) for both models and gMDSCs for IOT-resistant tumours (suppl. fig. 4J)

express significantly higher Adora2a amount when compared to matched spleens. This

expression differs significantly between IOT- responsive and resistant models in CD11b$^+$

dendritic cells (mean MFI 2080 ± 174 vs 3190 ± 636 respectively; p=0.007), M1 macrophages

(4520 ± 983 vs 6940 ± 1690; p=0.02) and mo-MDSCs (1890 ± 479 vs 6110 ± 1870; p=0.001)

(suppl. fig. 4G-I).

Targeting adenosine pathway delays tumour growth of murine pancreatic cancer
representing a combinational therapeutic opportunity

Our data suggest a mechanism by which the myeloid population contributes to the pro-

tumorigenic functionality of the pancreatic cancer microenvironment, where eAdo generated

by the myeloid cell populations and cancer cells would target and stimulate further the myeloid

cell subpopulations, in particular the pro-tumorigenic M2 macrophages. Therefore, we

inhibited *in vivo* eAdo formation and function, using an antibody against CD73 (2c5mIgG1)

and a small molecule inhibitor of Adora2a (AZD4635), a combination (Adoi) which would

maximise the inhibition of the axis. The 14-day treatment was started after the

microenvironment was allowed to establish (12-14 days after implantation) in the IOT-resistant

allografts (fig 4A). The anti-CD73 was extremely effective in reducing the expression of CD73

on the surface of all live cells (suppl. fig. 5A). MSI data confirmed marked reduction in

adenosine formation in the TME (Fig. 4 B-C).   In particular, adenosine was completely

abolished in the viable tumour areas, while a small amount remained in the necrotic margins,

accounting for a 95% decrease (fig. 4C), highlighting the importance to inhibit not only the

formation of adenosine through CD73 inhibition, but also targeting adenosine receptors. The

effectiveness of the treatment on the extracellular purinergic pathway was also supported by

the decrease of molecules downstream of adenosine (adenine and inosine in viable tumour and

necrotic margin areas) (suppl. fig 5B-C), and the increase of upstream and alternative pathway

molecules as AMP (in the necrotic margin, suppl. fig 5D) and xanthine (in both viable tumour

and necrotic margin, suppl. fig 5E) respectively. There was no change in the distribution of

ATP, ADP and hypoxanthine (suppl. fig 5F-H).

The Adoi approach led to a 30% reduction of tumour growth ratio (mean of 8 vs 5.5 -folds

increase from the baseline, p=0.003, fig. 4D) and tumour weight (median 1.24 vs 0.78 grams,

0.039 fig. 4A). Single agent AZD4635 induced a similar tumour control to the full combination.

However, the therapeutic effect was delayed (suppl. fig. 6B), with a 30% growth reduction   in

the combination arm when compared to Adora2ai inhibition alone with single agent AZD4635,

over the first 4 days of treatment (p=0.01). For biological reasons, given also that AZD4635 is

a competitive inhibitor and its efficacy is dependent on the amount of extracellular adenosine,

we chose to use it in combination with anti-CD73 antibody to maximise the blocking on

extracellular adenosine effects. These data support previous findings showing the same

combination had greater anti-tumour immune effect [43]. The adenosine pathway has been shown

to control the metastatic process and several authors have shown that inhibiting this axis can

reduce the metastatic burden in mouse models [44, 45]. However, we were able to show for the

first time that blocking adenosine generation and function can significantly reduce the

occurrence of spontaneous metastasis. The 6419c5 s.c. model spontaneously develops lung

metastases in 100% of the mice and blocking adenosine strongly reduced the metastatic burden

(median of % mets/lung area 0.77% vs 2.6%; p=0.016; fig. 4E).

These data suggest that targeting myeloid related, extracellular adenosine formation and

effect would have an effect on tumour growth, making this approach a candidate for

combinatorial therapeutic studies. Indeed, when combined with cytotoxic treatment (AZD6738

and gemcitabine; fig. 4F-G) or IOT (anti-CD40 agonist, anti-PD-L1 and anti-CTLA4 (FCP);

suppl. fig. 6C), the adenosine modulation reduced further the tumour growth rate of the

aggressive IOT-resistant 6419c5 tumour model. AZD6738/gemcitabine alone was able to

significantly slow the growth of the IOT-resistant model (2/7 stable disease, SD [28.5%]), but

the addition of the adenosine blocking (AZD4635 + 2c5mIgG1) led to further stabilization of

the tumour growth in a 2-week regimen (p=0.003 vs AZD6738/gem alone; 4/7 SD [57.1%];

fig. 4G-H). These data supported the investigation of the same combination in autochthonous

tumours in KPC mice (fig. 4I) to assess whether the addition of Adoi to cytotoxic treatment

(AZD6738+gemcitabine) prolonged survival in this model, considered a gold standard in this

disease. As figure 4J shows, the combination of Adoi and ATRi/gem induced a 3-fold increase

in median overall survival (mOS) in KPC mice compared to control groups (vehicles + isotype

12.5 days vs combo 41 days, p=0.0006). The 4-drug regimen is significantly better than single

schedule arms (mOS: Adoi 19 days and AZD6738 + gemcitabine 13 days). A weekly

ultrasound revealed a tendency for tumour stabilisation in mice treated with cytotoxic

therapeutics plus adenosine inhibition, when compared to the control arm during the first 2

weeks of treatment (fig. 4K).

Combining adenosine blockade (Adoi) and immunotherapy (FCP) reduced significantly

tumour growth of the IOT-resistant allograft model when compared to control treatment (65%

tumour growth reduction; p=0.002) and FCP (35% tumour growth reduction; p=0.021) arms

(suppl. fig. 6 C-D). Data from two separate experiments with the same controls showed that

adding Adoi to FCP remains the best combination in controlling tumour growth (suppl. fig.

6E-G). These data suggest again that targeting the adenosine pathway in PDAC offers a new

strategy to modulate the anti-tumour immune response.

Targeting the adenosine pathway results in reprogramming of the TME in the IOT-

resistant pancreatic cancer mouse model.

To understand the role of adenosine modulation in reprogramming the TME, given the effect

on tumour growth alone or in combination we analysed the changes in immune infiltration

following anti-CD73 and AZD4635 treatment.

The TME of PDAC is an intricate structure that relies on the presence of multiple non-

malignant cells. This TME is recapitulated in pre-clinical models of PDAC [34]. In order to

investigate the broader effect of Adoi in the 6419c5 PDAC model, given the effect we have

seen on tumour growth when used alone or in combination, we performed a bulk RNAseq

analysis of the 6419c5 model treated with Adoi or control (5 vs 6 mice, see methods).

Following the 14-day treatment with AZD4635 and anti-CD73 the relative expression of 712

genes decreased by ≥50% (logfold <-0.58; fig. 5A). KEGG and GO Biological process pathway

analysis revealed a dependency of the TME on the presence of extracellular adenosine (fig. 5B

and suppl. fig. 7A). Genes associated with hypoxia response and vasculogenesis (e.g. *Hif1a*,

Slc2a1, *Hilpda*, *Adm*, *Vegfa*, *Vegfd*), immunity and immune suppression (e.g. *Cd274*, *Cd209*,

Mrc1, *Cd200*, *Il1a*, *Il6*, *Ptgs2*) and tumour stroma/ECM organisation (e.g. *Col5a3*, *Col6a3*,

Itga2, *Mmp13*, *Mmp3*, *Mmp9*, *Ereg*, *Pthlh*) were significantly downregulated by the treatment

(fig. 5B-C).

Of note, several functional and structural downregulated genes were associated with

the innate immune system (fig. 5D left), particularly with M2 polarised macrophages (e.g.

Cd209, *Mrc1*, *Mrc2*, *CD163*). RNAseq analysis also showed a significant decrease of the

expression of *Cd274* (PD-L1) gene, suggesting a strong rationale for the use of adenosine

inhibition as combination for immunotherapy studies involving immune checkpoint inhibitors.

The downregulation of *Mrc1* (CD206) and *Cd274* (PD-L1) was validated analysing their

protein expression on tumour-infiltrating live cells following Adoi treatment. As shown in fig

5D (right), the proteins of these genes were strongly downregulated in the adenosine inhibition

arm (38% and 41% reduction of CD206 and PD-L1 on live cells respectively).

Furthermore, the adenosine signalling has previously been associated with the presence

of hypoxia [46, 47]; here we show for the first time that the presence of a functional extracellular

adenosine pathway is responsible for the expression of several genes related to hypoxia

(including *Hif1a*) suggesting the presence of a positive feedback loop.

Considering that the RNAseq analysis indicated that the innate immune system was

affected by adenosine inhibition, we analysed the changes in immune infiltration following

anti-CD73 and AZD4635 treatment. Flow cytometry analysis confirmed the findings

highlighted by the RNAseq analysis. Following treatment, tumours were less likely to be

infiltrated by M2 macrophages (median approximately 35000 vs 23000 cells/100mg; p=0.004;

fig. 6 A-B), in particular PD-L1$^+$ ones (median 79% vs 65%; p=0.004; fig. 6 B, right). IMC

analysis revealed that the reduction of M2 macrophages either F4/80$^+$ CD206$^+$ (fig. 6C, E) or

CD68$^+$ CD206$^+$ (fig. 6D, F) cells, was more prominent in viable tumour areas. Notably this

reduced infiltration is present in areas other than those where adenosine is abundant, suggesting

that adenosine is stimulating the production of factors affecting recruitment of macrophages in

the viable tumour areas. Blocking the axis, also led to a reduced frequency of infiltrating Tregs

(mean 42% vs 27%; p=0.03, suppl. fig. 7B-C) and a reduction of PD-L1 expression for all live

cells within the tumour, more prominently CD45$^+$ cells (p=0.008) as shown in suppl. fig. 7D.

However, the expression of PD-L1 however declined in F4/80$^+$ macrophages (p=0.02) but not

in dendritic cells following treatment (suppl. fig. 7E). Of note, the combination of 2c5mIgG1 and AZD4635 is required to reduce the M2 macrophage infiltration (p=0.03, suppl. fig. 7F) and the ratio of M2/M1 macrophages (p=0.04; suppl. fig. 7G) in the tumour, while single agents fail to do so. Finally, IMC data also confirmed a reduction of total macrophages in the treated tumours as shown by flow, again more evident in the viable tumour regions (fig. 6C,D, F4/80 and CD68 panels and suppl. fig 7H). IMC also showed a global reduction of M1 macrophages (F4/80$^+$ CD206$^-$ MHCII$^+$) (suppl. fig. 7 I-J) which was not apparent in the flow analysis. Further studies would further shed light on the relevance of these results, analysing the contribution of adenosine abundance and spatial distribution to modelling of tumour infiltration by distinct macrophage subpopulations beyond M1-M2 dichotomy.

Considering the reduction of the immune suppression in the TME following Adoi, we explored changes associated with the adaptive immune system following combination of adenosine inhibition and cytotoxic or immunotherapy therapeutics. To assess whether the combination of Adoi plus ATRi/gemcitabine had an effect on the immune infiltration of the TME of IOT-resistant model (6419c5 allografts), we performed IHC staining for CD8 and Foxp3 and showed that the quadruple combination almost tripled the ratio CD8/Tregs (median 0.70 vs 0.25 of vehicles + isotype group; p=0.04; fig. 6G-H). The 4-drug combination produced an enhanced increase of the CD8$^+$ cell infiltration (median 4.5 cells/mm^2 vs 2.7 in the control group, 3.9 in the AZD6738/gem group and 2.8 in the AZD4635/αCD73 group) and reduction of Foxp3$^+$ cell infiltration (median 6.5 cells/mm^2 vs 8.1 in the control group, 9.3 in the AZD6738/gem group and 7.2 in the AZD4635/αCD73 group) (suppl. fig. 7K). Similarly, combination with IOT (Adoi + FCP) determined a further increase of the intra-tumoral CD8/Tregs ratio induced by FPC treatment (ratio means control 0.28 vs FCP/Adoi 6.10 p=0.008; suppl. fig. 7L) mostly due to the repression of the Treg recruitment induced by IOT alone.

Adenosine-related gene expression is associated with phenotype and survival in human pancreatic cancer

To evaluate the importance of this pathway in the context of human PDAC and highlight the role in the formation of TME we scored our adenosine-related gene expression set against the PDAC subtypes published by Bailey [39]. Of our 712 genes with a 50% decrease, 561 had a human ortholog and of these the z-scores of 517 genes were summed in each of the 97 patients with RNAseq data in Bailey et al. to obtain a score. We were able to show that the adenosine-related gene expression is mostly expressed in the aggressive squamous subtype (fig. 7A), that has been associated with a poorer prognosis. The score remained significantly higher even when comparing the squamous subtypes with the grouped non-squamous (fig. 7B).

In order to create a PDAC-specific adenosine signature and to evaluate its performance in PDAC-specific outcomes, we created a signature using the 561 human orthologous genes with a $\geq$ 50% decrease in expression following Adoi treatment. Of these, 52 genes were selected for the signature (fig. 7C), that according to our RNAseq dataset were clearly associated with the pathway areas dependent on adenosine (hypoxia response, immunity, tumour stroma), were associated positively or negatively with PDAC prognosis, and were significantly co-expressed with CD73 and/or Adora2a in the PDAC genome datasets (TCGA or Bailey). Of the 176 and 170 patients with available progression free survival (PFS) and disease-specific survival (DSS) outcome respectively, we found that the presence of a high adenosine signature is associated with higher probability of PDAC progression (mPFS high Ado 13.05 vs low Ado 18.25, p=0.02; fig. 7D) and a poorer PDAC-specific survival (mDSS high Ado 19.66 vs low Ado *NR*, p=0.01; fig. 7E), suggesting again that the presence of a functional adenosine pathway has a detrimental role in human PDAC. These data should be

considered in light of the fact that patients in the TCGA dataset are predominantly non-metastatic, and the presence of the adenosine signature seems to become relevant for PDAC associated death 20 months after diagnosis. The presence of a high adenosine signature appears to be associated with shorter disease-free-survival, (mDFS high Ado 23.5 months vs low Ado 49.7 months, p=0.12; suppl. fig. 7M).

Discussion

Overall our data highlight that the generation of the extracellular adenosine is instrumental for the innate immune system in shaping a pro-tumorigenic, immune suppressive microenvironment in PDAC, in the context of a hypoxic milieu. We can speculate that this unfavourable environment may create the condition for a more aggressive PDAC phenotype which would then translate into the ability to escape the immune system, be resistant to cytotoxic treatment and metastasize readily.

Pancreatic ductal adenocarcinoma (PDAC) is projected to become the second highest cause of cancer-related death in the US within 10 years [48], and represents one of the major unmet needs of cancer treatment. Despite extensive efforts by laboratory and clinical scientists in the last 50 years, only 1% of patients diagnosed with PDAC today will survive for 10 years (https://www.cancerresearchuk.org/health-professional/cancer-statistics-for-the-uk). The response rate following standard treatments is poor, usually short-lasting, and associated with significant treatment related toxicity [1]. In the past few years, immunotherapy has provided new hope in the treatment of several types of cancer, and has dramatically changed the life expectancy of many patients with metastatic disease [49]. However, this has not been true for patients with PDAC which is associated with a very low response rate to immunotherapy, usually confined to MSI-H/dMMR tumours, found only rarely in this disease [2].

The role of the innate immune system in the generation of an immune suppressive/pro-
tumorigenic microenvironment in PDAC is well known. The presence of marked infiltration
of macrophages has been identified as an independent predictive factor of the aggressiveness
and prognosis of PDAC, in patients [23, 24]. Only recently, three phase I clinical trials in patients
with PDAC have shown that targeting the innate immune system can have impact in patients
with PDAC. A phase I trial published on Lancet Oncology, showed that a combination of anti
PD-1 and CD40 agonist, added to gemcitabine and nab-paclitaxel, led to 60% of response rate,
with some durable responses [50]. In addition, the inhibition of the CXCL12/CXCR4 axis has
been demonstrated to modify the immunosuppressive TME of PDAC and CRC patients [51, 52].

The extracellular adenosine pathway has also been shown to influence the TME
fostering the immune suppression provided by some innate immune subpopulations (as
myeloid and NKs) and inhibiting the function of the adaptive immune system, in particular, T-
cells [6, 33, 44]. By activating its receptors, eAdo is able to increase the intracellular concentration
of cAMP which leads to the induction of a M2 phenotype of macrophages and block the
secretion of IL1β increasing the release of CXCL1, IL-6, IL-10 and IL-8 among others from
myeloid population which are known to orchestrate immune exclusion [8, 53]. eAdo also favours
the formation and maintenance of Treg cells [8], which are known to favour cancer progression
and IOT resistance .

A recent publication, shows that genomic targeting on mouse PDAC cells of CD73
leads to a reduced in vivo tumour formation and change in the circulating and infiltrating
immune system[54].

However, to date, little was known about the expression of the adenosine pathway in
the context of the innate and adaptive immune system in PDAC, how the extracellular

adenosine is generated and what are the targets of adenosine also in regard to its spatial

distribution and formation of adenosine.

Our results show for the first time, that the mechanism of generation of extracellular

adenosine in pancreatic cancer TME is finely orchestrated by tumour infiltrating myeloid cells

and tumour cells, due to the expression of high level of CD39 in infiltrating myeloid cells and

CD73 on both cell types. We have also demonstrated that the pathway can be overexpressed

in T-cells infiltrating the tumours, regardless of their activation status (fig. 1,2 and suppl. fig.

2,4). The distribution of extracellular adenosine is spatially heterogeneous and a high level of

extracellular adenosine correlates with the presence of a hypoxic environment and is favoured

by the presence of necrosis, where the myeloid population is enriched (fig.3). Necrosis is

common in human PDAC and related to poor prognosis for all stages [55]. The enrichment of a

CD39$^+$ CD73$^+$ double population, potentially able to independently produce adenosine, does

not seem to correlate with IOT- resistant or responsive tumour models, but there is a difference

when the target of adenosine (Adora2a receptor), is considered. Adora2a on myeloid

populations, in particular in pro-tumorigenic M2 macrophages but also in antigen presenting

cells, is differentially expressed in regard of IOT response phenotype, with the resistant

tumours abundantly overexpressing the receptor in these populations (fig.3 and suppl. fig. 4).

The bulk RNAseq analysis of tumour treated with adenosine inhibition revealed indeed a

broader role for adenosine in PDAC TME (fig. 5). Genes related to immunosuppression and

innate immunity recruitment (*Cd274*, *Csf2*, *Cxcl2*, *Ccl3*, *Ccl4*, *Ccl12*, *Il1a*, *Osm*, *Il6*),

angiogenesis (*Vegfa*, *Vegfd*, *Adm2*, *Flt1*, *Pgf*, *Egln1*) and cell-ECM interaction (*Adam19*,

Adamts14, *Adamts4*, *Adamts5*, *Col5a3*, *Col6a*, *Itga2*, *Itga7*, *Mmp3*, *Mmp9*, *Mmp12*) indicated

the targeting of population of cells responsible for the acquisition of a pro-tumorigenic, pro-

metastatic, pro-fibrotic and immune resistant phenotype. Further, the downregulation of genes

associated with hypoxia (*Hif1a*, *Hilpda*, *Nos2*, *Hk1*, *Hk2*, *Egln3*) following treatment shows

not only that the adenosine pathway is induced during hypoxia (e.g. CD73 and Adora2b), but

that the hypoxia response is also dependent on the presence of the adenosine pathway in what

we can speculate is a positive feedback loop. The inhibition of adenosine led to a reduced

infiltration of M2 macrophages further from the hypoxic regions where adenosine is most

abundant, suggesting that the effect of adenosine is stimulating the secretion of factors that are

recruiting monocytes into the tumour microenvironment, replenishing macrophage infiltration

(fig. 6 and suppl. fig. 7). Notably, targeting the pathway can reduce tumour growth in an IOT-

resistant model, improving the response to cytotoxic and immunotherapy combinations (fig. 4

and suppl. fig. 6) even in historically therapy-resistant models as the KPC mice.

Targeting adenosine would represent an alternative strategy to reduce the infiltration of

pro-tumorigenic macrophages in PDAC lesions. One approach has been the administration of

a CSF1R inhibitor [32], which has shown promising pre-clinical data, that have not been

translated in human so far. A recent publication shows that in CRC mouse models, the use of

anti CSF1R treatment spares a subpopulation of macrophages characterised by the expression

of *Cd274* (PD-L1), *Vegfa*, *Hilpda*, *Bhlhe40*, *Mmp12*, *Cebpb*, *Hmox1* among others [28]. Given

that all of these genes are among the immune suppressive and vasculogenic molecules that

seem to be strongly downregulated by adenosine inhibition and our data show a reduction of

some subpopulation of PD-L1$^+$ macrophages, we can speculate that adenosine inhibition could

potentially target these populations of pro-angiogenic, immune suppressive macrophages.

Information provided by the PDAC specific adenosine signature, indicate that the

adenosine pathway appears to play a role in progression and survival of human PDAC, due to

the ability of the adenosine inhibition to profoundly reprogram the tumour microenvironment

in PDAC models (fig. 5-6 and suppl. fig. 7). Gathering more information on the role of this

pathway in human cancers should be a priority, retrospectively evaluating and then

prospectively stratifying the patients on the basis of histopathological/radiological features and

the spatial distribution of adenosine.

In summary, we have shown for the first time that tumour-infiltrating myeloid immune

cells contribute to the generation of extracellular adenosine in the context of PDAC, in

correlation with the presence of hypoxia. Macrophages in particular, express high levels of

Adora2a receptor in PDAC models and targeting the adenosine/myeloid axis remodels the

TME. Data from IMC, flow cytometry and RNAseq suggest that the adenosine pathway is

fundamental to the formation of a pro-tumorigenic, immunosuppressive TME, and its

expression is associated with an aggressive phenotype and poor survival in human PDAC. The

re-modelling of the TME caused by the inhibition of the adenosine pathway, translate into a

delay of PDAC tumour growth. Finally, targeting extracellular adenosine should be considered

as an alternative to improve the efficacy of cytotoxic and IOT in future PDAC-specific clinical

trials.

Author contributions

V.G. and D.I.J conceptualised the project. V.G., A.D., H.H., K.W., J.P.M., J.E.

and DIJ designed the experiments. V.G., A.D., H.H., K.W., H.B., S.A.K., S.Y.L.,

B.P.K. conducted the experiments. V.G., A.D., H.H., K.W., S.I., F.M.R., R.B.,

J.P.M, S.J.D. J.E. and D.I.J analysed the data. J.E.D.T. R.G., J.P.M., S.J.D.,

A.G.S, J.E., D.I.J. provided supervision and V.G., A.D., H.H., K.W., S.I., J.P.M.,

S.J.D., A.G.S, J.E., D.I.J. discussed the results. V.G. and D.I.J. wrote the

manuscript. All authors read, review, edited and approved the final manuscript.

Acknowledgments

All CRUK CI authors received research funding from Cancer Research UK (Nos.

C14303/A17197 and C9545/A29580). The Li Ka Shing Centre where this work was performed

was generously funded by CK Hutchison Holdings Limited, the University of Cambridge,

CRUK, The Atlantic Philanthropies and others. This work was supported by Cancer Research

UK (C9685/A27444) funding to VG. The authors wish to thank all the CRUK Cambridge

Institute core facilities, in particular Research Instrumentation & Cell Services (RICS), Flow

Cytometry, Histopathology, Biological Resource Unit (BRU), Genomics and Genome editing

cores. VG would like to thank Maike de la Roche (CRUK Cambridge Institute - University of

Cambridge) and Klaus Okkenhaug (Department of Pathology - University of Cambridge) for

helpful suggestions and advices. This study was also supported by Cancer Research UK

Precision Panc grant C96/A25238. HB holds a PhD studentship at the University of Cambridge

which is supported jointly by the University of Cambridge Experimental Medicine Training

Initiative (EMI) programme in partnership with AstraZeneca (EMI-AZ) and the NIHR

Biomedical Research Centre and SYL is founded by the Cambridge Trust (Cambridge

International Scholarship). Work by JT / KW is funded by a core grant award from the MRC

(MC_UU_00025/12). S.A.K. and. J.P.M. were supported by Cancer Research UK core

funding to the Beatson Institute (A17196 and A31287) and to J.P.M. Lab (A29996). The

KPCY-derived cell lines were a kind gift of Ben Stanger (University of Pennsylvania).

AZD4635, AZD6738 and the antibodies anti-CD73 (2C5, murine IgG1-Fc), anti-PD-L1

(AB740080 D265A), NIP228 muIgG1 isotype, NIP228 muIgG1 D265A isotype, were kindly

provided by AstraZeneca. The results shown here are in part based upon data generated by the

TCGA Research Network: https://www.cancer.gov/tcga.

Figure legends

Figure 1. Expression of CD73 on KPCY-derived cell lines and response to anti-CD73 in

vitro inhibition.

(A) Representative histogram of CD73 expression on KPCY-derived cell line (2838c3) in flow

cytometry. (B) CD73 expression was evaluated on KPCY-derived cell lines after 24 hours

incubation with 10 μg/ml of anti-CD73 or isotype.

(C) 2838c3 (left) and 6419c5 (right) cell lines were grown with increasing concentration of

anti-CD73 or isotype (100 μg/ml) and confluency was evaluated using IncuCyte time lapse

imaging for up to 72 hours. For each experiment, 3 different wells per condition were used per

experiment.

(D-E) Representative images (D) and graphs (E) showing survival fraction of cells (2838c3

left, 6419c5 right) from the colony-forming experiment following 8-day treatment with anti-

CD73 or isotype. For each experiment, 3 different wells per condition were used.

All data are presented as mean ± SEM from experiments repeated 3 times. Statistical analysis

was performed with two-tailed unpaired Student's t-test (B), mixed-effect model (C) and one-

way ANOVA with post-test analysis for multiple comparisons; p values are shown in the

graphs when considered significant (p<0.05).

Figure 2. The adenosine pathway members are expressed on PDAC-infiltrating immune

cells.

(A) Representative flow cytometry plots showing expression of CD39 and CD73 on myeloid

population (left), Tregs (middle) and CD8+ T-cells (right) from KPCY-cell line derived tumour

(upper) and matched spleen (lower) (N= 5 mice per group).

(B) Box and whisker graph showing CD39$^+$CD73$^+$ double expression on CD11b$^+$ cells for

2838c3 (N=5) and 6419c5 (N=5) in tumours, matched spleens and tumour draining lymph

nodes.

(C-D) Representative flow cytometry plots (C) and box and whisker graph (D) showing

$CD39^+CD73^+$ double expression on $CD11b^+$ cells infiltrating autochthonous KPC tumours

(N=13). All data are presented as interleaved box and whiskers.

Statistical analysis was performed using one-way ANOVA with post-hoc test analysis for

multiple comparisons (B) and two-tailed unpaired Student's t-test (D); p values are shown in

the graphs when considered significant (p<0.05).

Figure 3. Adenosine distribution is spatially heterogeneous and targets myeloid

subpopulations.

(A) Mass spectrometry Imaging (MSI) representative images showing adenosine expression

and distribution in PDAC allografts (4 mice per group) at day 21 post-implantation.

Classification was obtained based on metabolites expression and is represented as follows:

viable tumour (red), necrotic margins (green).

(B) Relative tissue composition differences of viable tumour, necrotic margin and necrotic

areas for 6419c5 (N=4) and 2838c3 (N=4) allografts at day 21 post-implantation. As shown,

necrosis was present in only one 2838c3 sample at 21 days post-implantation. Error bars

represent Standard Deviation.

(C) MSI analysis showing relative abundance (a.u.) of adenosine in the different areas in

6419c5 and 2838c3 allografts at day 21 post-implantation. Bars represent means.

(D) The plot (upper left) shows cell density of the ratio cancer cells ($PanCK^+$, purple) / myeloid

cells ($CD11b^+$, blue) on a Log2 scaled plot from Imaging Mass cytometry (IMC). The results

of the comparison between the whole area analysed and the other tissue regions are not shown.

N = 6. Asterisks ****, show adjusted p value of <0.0001. The IMC and tissue segmentation

images (upper right and lowers) are from one representative 6419c5 allograft replicate at day

28 post-implantation. The tissue segmentation of the IMC image was performed by Random

Forest Classification using all markers analysed. Scale bar on the IMC image is 200 µm. Segmentation shows viable tumour (green), necrosis (yellow), necrotic margin (blue) and off-tissue (red).

(E-F) Representative plots (E) and summary graph (F) from flow cytometry analysis showing Adora2a expression on pro-tumorigenic M2 macrophages in allografts (upper) derived from 6419c5 (left) and 2838c3 (right) implantation. Same expression is shown in M2 macrophages in matched spleens (lower) (8-9 mice per group were used).

(G-H) Flow cytometry plots (G) and graph (H) showing expression of Adora2a in M2 macrophages comparing KPC (n=8) autochthonous tumours and matched spleens.

(I) Box and whisker plot of the percentage of M2 macrophages positive for Adora2a comparing two allografts (6419c5 and 2838c3) and KPC tumours.

Statistical analysis was performed using one-way ANOVA with post-hoc test analysis for multiple comparisons (D,F,I) and two-tailed unpaired Student's t-test (H); p values are shown in the graphs when considered significant ($p<0.05$).

Figure 4. *In vivo* modulation of the adenosine pathway reduces tumour growth and metastasis and improves the efficacy of cytotoxic treatment.

(A) Schedule of adenosine inhibition (Adoi) treatment. Treatment was started following 12-14 days from implantation and continued for 2 weeks. Antibody anti-CD73 (2c5mIgG1, murine IgG1) was dosed twice per week intraperitoneally at 10 mg/kg. Adora2a inhibitor (AZD4635) was given by oral gavage twice daily at 50 mg/kg.

(B-C) Mass spectrometry Imaging (MSI) representative images (B) and MSI analysis graph with relative abundance (a.u.) (C) showing adenosine expression and distribution in PDAC allografts treated with vehicle + isotype or Adoi, at day 14 from treatment start (6 mice per

treatment group). Classification was obtained based on metabolites expression and is represented as follows: viable tumour (green line), necrotic margins (yellow line).

(D) Tumour growth ratio (left) and weight (right) of 6419c5 allografts in C57Bl/6 mice treated with anti-CD73 + AZD4635 (N=16) or vehicle + isotype (N=15).

(E) Representative image (left) and graph (right) showing % of area of the lung analysed occupied by metastasis (met/lung areas x 100) in vehicle + isotype (N=12) and anti-CD73+AZD4635 (N=13), evaluated for the presence of spontaneous occurrence of lung metastases. Every dot represents a single mouse. Tissues were stained with an anti-p53 antibody to highlight the presence of cancer cells. A group of more than 5 p53-positive cells was counted as metastasis.

(F) Schedule of 6419c5 tumour allografts 14-day treatment as following (N=7 mice per group): vehicles + isotype, AZD6738 + gemcitabine, anti-CD73 + AZD4635, AZD6738 + gemcitabine + anti-CD73 + AZD4635.

(G) Tumour growth ratio (14 days) of 6419c5 tumour allografts treated as above

(H) Percentage change in the long diameter length following 14 days of treatment per group. Number of mice with stable disease (SD, <20% increase and <30% decrease) are shown at the bottom.

(I) Schedule of KPC mice treatment as following: vehicles + isotype (12 mice), AZD6738 + gemcitabine (11 mice), anti-CD73 + AZD4635 (7 mice), AZD6738 + gemcitabine + anti-CD73 + AZD4635 (combo, 12 mice). Adenosine inhibition was administered until endpoint, while AZD6738 + gemcitabine was allowed up to 3 weeks.

(J) Survival analysis of KPC mice treated as above, including median overall survival (mOS).

(K) Percentage of tumour volume increased in vehicles vs combo group at day 7 and 14 of treatment.

All data are presented as mean ± SEM. Statistical analysis was performed using Mann-Whitney test (D-E,K), mixed-effect model (F) and one-way ANOVA with post-hoc test analysis for multiple comparisons (C). Log-rank Mantel-Cox test to evaluate difference in survivals (J); p values are shown in the graphs and considered significant when $p < 0.05$.

Figure 5. Adenosine inhibition remodels the PDAC tumour microenvironment

(A) Volcano plot related to 6419c5 allografts. Genes overexpressed in vehicle + isotype arm are on the left side (blue dots), and genes overexpressed in the anti-CD73 + AZD4635 arm are on the right side (Adoi, orange dots).

(B) Enrichment bar plot of significantly overexpressed pathways in the vehicle + isotype group (adenosine high), according to KEGG and GO Biological processes.

(C) Heatmap showing genes regulated during treatment (light grey for controls, dark grey for Adoi) which are part of the significantly different pathways according to KEGG.

(D) Table (left) showing genes associated with innate immunity that are downregulated by Adoi treatment. Validation of RNAseq data (right graphs) through flow analysis of PD-L1 and CD206 expression (MFI ratio) on intra-tumoral live cells (8 mice in the control and 10 in the Adoi arms). The samples analysed for validation were from different mice from the ones used for RNAseq.

Figure 6. Immunosuppressive immune subpopulations are modulated following adenosine pathway blockade.

(A) Representative flow cytometry plots showing tumour associated macrophages (TAM) infiltrating 6419c5 allografts following 14 day of treatment with vehicle + isotype (left panel) or anti-CD73 + AZD4635 (right panel).

(B) M2 macrophage allograft infiltration (left, number of cells/100 mg of tumour) and (right) percentage of M2 macrophages positive for PD-L1 (8 and 10 mice per group analysed).

(C-D) Representative IMC image of F4/80 and CD206 (C) and CD68 and CD206 (D) positive

cells infiltrating a 6419c5 allograft merged (1st panel of C and D) or not with segmentation

following Adoi (lower panels) or control (upper). The tissue segmentation of the IMC image

was performed by Random Forest Classification using all markers analysed. The scale bar on

the IMC image is 200 µm. Segmentation shows viable tumour (green), necrosis (yellow),

necrotic margin (blue) and off-tissue (red).

(E-F) The bar plots show cell density (number of cells per mm2) of $F480^+CD206^+$ cells (E) and

$CD68^+CD206^+$ (F) per segment area (6 mice per treatment group).

(G-H) Immunohistochemistry analysis showing 6419c5 tumour-infiltrating $CD8^+/Foxp3^+$ ratio

(G) at day 14 of the following treatments: Vehicles + isotype (N=7), AZD6738 + gemcitabine

(N=7), aCD73 + AZD4635 (N=6) and AZD6738 + gemcitabine + aCD73 + AZD4635 (N=7).

(H) Representative IHC images of the latter.

All data are represented as box and whisker plots. Statistical analysis was performed using

Mann-Whitney test (B-H) and one-way ANOVA with post-hoc test analysis for multiple

comparisons (E-H); p values are shown in the graphs when considered significant ($p<0.05$).

1001

Figure 7. RNA gene expression demonstrates a crucial role for the adenosine pathway in

correlation to human PDAC prognosis.

(A-B) Human ortholog genes with a 50% downregulation following 14 days of Adoi, were

scored using z-score derived from Bailey's[39] subtypes dataset. Genes scores comparing Adex,

immunogenic, squamous and pancreatic progenitor (A) or squamous vs non-squamous (B) are

shown. Each dot represents a single patient.

(C) PDAC-specific gene signature of 52 genes used for analysis of the TCGA human PDAC

dataset, related to the main pathways implied (immunity, hypoxia response and tumour

stroma).

(D-E) Gene signature applied to PDAC TCGA dataset (G) for progression free survival (PFS, 176 patients) and (H) disease specific survival (DSS, 170 patients) Kaplan-Meier. All data are presented as mean ± SEM when applicable.

Statistical analysis was performed using one-way ANOVA with post-hoc test analysis for multiple comparisons (A), two-tailed unpaired Student's t-test (B) and log-rank Mantel-Cox test to evaluate difference in survivals (D-E); p values are shown in the graphs when considered significant ($p < 0.05$).

References

Cancer Research UK, https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/pancreatic-cancer, Accessed March 2022

1. Conroy T, *et al.* FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med* **364**, 1817-1825 (2011).

2. Le DT, *et al.* Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* **357**, 409-413 (2017).

3. Hu ZI, *et al.* Evaluating Mismatch Repair Deficiency in Pancreatic Adenocarcinoma: Challenges and Recommendations. *Clin Cancer Res* **24**, 1326-1336 (2018).

4. Balachandran VP, *et al.* Identification of unique neoantigen qualities in long-term survivors of pancreatic cancer. *Nature* **551**, 512-516 (2017).

5. Ho WJ, Jaffee EM, Zheng L. The tumour microenvironment in pancreatic cancer - clinical challenges and opportunities. *Nat Rev Clin Oncol* **17**, 527-540 (2020).

6. Antonioli L, Blandizzi C, Pacher P, Hasko G. Immunity, inflammation and cancer: a leading role for adenosine. *Nat Rev Cancer* **13**, 842-857 (2013).

7. Augustin RC, Leone RD, Naing A, Fong L, Bao R, Luke JJ. Next steps for clinical translation of adenosine pathway inhibition in cancer immunotherapy. *J Immunother Cancer* **10**, (2022).

8. Cekic C, Linden J. Purinergic regulation of the immune system. *Nat Rev Immunol* **16**, 177-192 (2016).

1046

1047   9.   Goodwin KJ*, et al.* Development of a quantification method for adenosine in tumors by LC-
1048   MS/MS with dansyl chloride derivatization. *Anal Biochem* **568**, 78-88 (2019).

1049

1050   10.   Duhen T*, et al.* Co-expression of CD39 and CD103 identifies tumor-reactive CD8 T cells in
1051   human solid tumors. *Nat Commun* **9**, 2724 (2018).

1052

1053   11.   Simoni Y*, et al.* Bystander CD8(+) T cells are abundant and phenotypically distinct in human
1054   tumour infiltrates. *Nature* **557**, 575-579 (2018).

1055

1056   12.   Ott M*, et al.* Profiling of patients with glioma reveals the dominant immunosuppressive axis is
1057   refractory to immune function restoration. *JCI Insight* **5**, (2020).

1058

1059   13.   Takenaka MC*, et al.* Control of tumor-associated macrophages and T cells in glioblastoma via
1060   AHR and CD39. *Nat Neurosci* **22**, 729-740 (2019).

1061

1062   14.   Goswami S*, et al.* Immune profiling of human tumors identifies CD73 as a combinatorial target
1063   in glioblastoma. *Nat Med* **26**, 39-46 (2020).

1064

1065   15.   Sidders B*, et al.* Adenosine Signaling Is Prognostic for Cancer Outcome and Has Predictive
1066   Utility for Immunotherapeutic Response. *Clin Cancer Res* **26**, 2176-2187 (2020).

1067

1068   16.   Cekic C, Day YJ, Sag D, Linden J. Myeloid expression of adenosine A2A receptor suppresses T
1069   and NK cell responses in the solid tumor microenvironment. *Cancer Res* **74**, 7250-7259 (2014).

1070

1071   17.   Sorrentino C, Miele L, Porta A, Pinto A, Morello S. Myeloid-derived suppressor cells contribute
1072   to A2B adenosine receptor-induced VEGF production and angiogenesis in a mouse melanoma
1073   model. *Oncotarget* **6**, 27478-27489 (2015).

1074

1075   18.   Li L*, et al.* Dendritic cells tolerized with adenosine A(2)AR agonist attenuate acute kidney
1076   injury. *J Clin Invest* **122**, 3931-3942 (2012).

1077

1078   19.   Borodovsky A*, et al.* Small molecule AZD4635 inhibitor of A2AR signaling rescues immune cell
1079   function including CD103(+) dendritic cells enhancing anti-tumor immunity. *J Immunother*
1080   *Cancer* **8**, (2020).

1081

1082   20.   Wilson JM*, et al.* The A2B adenosine receptor promotes Th17 differentiation via stimulation
1083   of dendritic cell IL-6. *J Immunol* **186**, 6746-6752 (2011).

1084

1085   21.   Romio M, Reinbeck B, Bongardt S, Huls S, Burghoff S, Schrader J. Extracellular purine
1086   metabolism and signaling of CD73-derived adenosine in murine Treg and Teff cells. *Am J*
1087   *Physiol Cell Physiol* **301**, C530-539 (2011).

1088

1089 22. Cekic C, Linden J. Adenosine A2A receptors intrinsically regulate CD8+ T cells in the tumor
1090 microenvironment. *Cancer Res* **74**, 7239-7249 (2014).

1091
1092 23. Ino Y*, et al.* Immune cell infiltration as an indicator of the immune microenvironment of
1093 pancreatic cancer. *Br J Cancer* **108**, 914-923 (2013).

1094
1095 24. Steele NG*, et al.* Multimodal Mapping of the Tumor and Peripheral Blood Immune Landscape
1096 in Human Pancreatic Cancer. *Nat Cancer* **1**, 1097-1112 (2020).

1097
1098 25. Zhang Y*, et al.* Myeloid cells are required for PD-1/PD-L1 checkpoint activation and the
1099 establishment of an immunosuppressive environment in pancreatic cancer. *Gut* **66**, 124-136
1100 (2017).

1101
1102 26. Raghavan S*, et al.* Microenvironment drives cell state, plasticity, and drug response in
1103 pancreatic cancer. *Cell* **184**, 6119-6137 e6126 (2021).

1104
1105 27. Zhu Y*, et al.* Tissue-Resident Macrophages in Pancreatic Ductal Adenocarcinoma Originate
1106 from Embryonic Hematopoiesis and Promote Tumor Progression. *Immunity* **47**, 597 (2017).

1107
1108 28. Zhang L*, et al.* Single-Cell Analyses Inform Mechanisms of Myeloid-Targeted Therapies in
1109 Colon Cancer. *Cell* **181**, 442-459 e429 (2020).

1110
1111 29. Beatty GL*, et al.* Exclusion of T Cells From Pancreatic Carcinomas in Mice Is Regulated by
1112 Ly6C(low) F4/80(+) Extratumoral Macrophages. *Gastroenterology* **149**, 201-210 (2015).

1113
1114 30. Ruffell B*, et al.* Macrophage IL-10 blocks CD8+ T cell-dependent responses to chemotherapy
1115 by suppressing IL-12 expression in intratumoral dendritic cells. *Cancer Cell* **26**, 623-637 (2014).

1116
1117 31. Daley D*, et al.* NLRP3 signaling drives macrophage-induced adaptive immune suppression in
1118 pancreatic carcinoma. *J Exp Med* **214**, 1711-1724 (2017).

1119
1120 32. Candido JB*, et al.* CSF1R(+) Macrophages Sustain Pancreatic Tumor Growth through T Cell
1121 Suppression and Maintenance of Key Gene Programs that Define the Squamous Subtype. *Cell*
1122 *Rep* **23**, 1448-1460 (2018).

1123
1124 33. Allard B, Allard D, Buisseret L, Stagg J. The adenosine pathway in immuno-oncology. *Nat Rev*
1125 *Clin Oncol* **17**, 611-629 (2020).

1126
1127 34. Li J*, et al.* Tumor Cell-Intrinsic Factors Underlie Heterogeneity of Immune Cell Infiltration and
1128 Response to Immunotherapy. *Immunity* **49**, 178-193 e177 (2018).

1129
1130 35. Dannhorn A*, et al.* Universal Sample Preparation Unlocking Multimodal Molecular Tissue
1131 Imaging. *Anal Chem* **92**, 11080-11088 (2020).

36. Adusumilli R, Mallick P. Data Conversion with ProteoWizard msConvert. *Methods Mol Biol* **1550**, 339-368 (2017).

37. Race AM, Styles IB, Bunch J. Inclusive sharing of mass spectrometry imaging data requires a converter for all. *J Proteomics* **75**, 5111-5112 (2012).

38. Dunlop CR*, et al.* Complete loss of ATM function augments replication catastrophe induced by ATR inhibition and gemcitabine in pancreatic cancer models. *Br J Cancer* **123**, 1424-1436 (2020).

39. Bailey P*, et al.* Genomic analyses identify molecular subtypes of pancreatic cancer. *Nature* **531**, 47-52 (2016).

40. Liu J*, et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. *Cell* **173**, 400-416 e411 (2018).

41. Zhou L*, et al.* The distinct role of CD73 in the progression of pancreatic cancer. *J Mol Med (Berl)* **97**, 803-815 (2019).

42. Overman MJ*, et al.* Safety, efficacy and pharmacodynamics (PD) of MEDI9447 (oleclumab) alone or in combination with durvalumab in advanced colorectal cancer (CRC) or pancreatic cancer (panc). *Journal of Clinical Oncology* **36**, 4123-4123 (2018).

43. Young A*, et al.* Co-inhibition of CD73 and A2AR Adenosine Signaling Improves Anti-tumor Immune Responses. *Cancer Cell* **30**, 391-403 (2016).

44. Yan J*, et al.* Control of Metastases via Myeloid CD39 and NK Cell Effector Function. *Cancer Immunol Res* **8**, 356-367 (2020).

45. Sun X*, et al.* CD39/ENTPD1 expression by CD4+Foxp3+ regulatory T cells promotes hepatic metastatic tumor growth in mice. *Gastroenterology* **139**, 1030-1040 (2010).

46. Synnestvedt K*, et al.* Ecto-5'-nucleotidase (CD73) regulation by hypoxia-inducible factor-1 mediates permeability changes in intestinal epithelia. *J Clin Invest* **110**, 993-1002 (2002).

47. Philip K*, et al.* HIF1A up-regulates the ADORA2B receptor on alternatively activated macrophages and contributes to pulmonary fibrosis. *FASEB J* **31**, 4745-4758 (2017).

48. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res* **74**, 2913-2921 (2014).

1175 49.   Mellman I, Coukos G, Dranoff G. Cancer immunotherapy comes of age. *Nature* **480**, 480-489
1176   (2011).

1177
1178 50.   O'Hara MH*, et al.* CD40 agonistic monoclonal antibody APX005M (sotigalimab) and
1179   chemotherapy, with or without nivolumab, for the treatment of metastatic pancreatic
1180   adenocarcinoma: an open-label, multicentre, phase 1b study. *Lancet Oncol* **22**, 118-131
1181   (2021).

1182
1183 51.   Biasci D*, et al.* CXCR4 inhibition in human pancreatic and colorectal cancers induces an
1184   integrated immune response. *Proc Natl Acad Sci U S A* **117**, 28960-28970 (2020).

1185
1186 52.   Bockorny B*, et al.* BL-8040, a CXCR4 antagonist, in combination with pembrolizumab and
1187   chemotherapy for pancreatic cancer: the COMBAT trial. *Nat Med* **26**, 878-885 (2020).

1188
1189 53.   Fong L*, et al.* Adenosine 2A Receptor Blockade as an Immunotherapy for Treatment-
1190   Refractory Renal Cell Cancer. *Cancer Discov* **10**, 40-53 (2020).

1191
1192 54.   King RJ*, et al.* CD73 induces GM-CSF/MDSC-mediated suppression of T cells to accelerate
1193   pancreatic cancer pathogenesis. *Oncogene* **41**, 971-982 (2022).

1194
1195 55.   Hiraoka N*, et al.* Tumour necrosis is a postoperative prognostic marker for pancreatic cancer
1196   patients with a high interobserver reproducibility in histological evaluation. *Br J Cancer* **103**,
1197   1057-1065 (2010).

1198

1199

Figure 1

Figure 2

A

B

C

D

Figure 3

Figure 4

Figure 5

C

KEGG pathway genes

D

GENES ASSOCIATED WITH INNATE IMMUNITY			
Gene	**Log2FC**	**pvalue**	**padj**
Nos2	-3.1079	1.15E-12	1.39E-09
Cd209d	-2.4151	0.00376	0.037288082
Hilpda	-2.3894	7.45E-25	1.02E-20
Cd209a	-2.0564	0.00198	0.024647181
Hmox1	-1.9360	7.25E-19	4.40E-15
Cxcl2	-1.7851	1.98E-05	0.001103048
Ccl8	-1.7496	0.00224	0.026325184
Ptx3	-1.7342	0.00012	0.003792336
Mmp9	-1.7005	0.00184	0.023332748
Bhlhe40	-1.6830	1.39E-15	4.23E-12
Cd274	-1.6819	3.68E-17	1.34E-13
Il1a	-1.5816	3.37E-05	0.001606959
Osm	-1.5691	5.55E-06	0.000422935
Ptgs2	-1.4432	8.51E-07	9.76E-05
Ccl3	-1.4289	0.00046	0.009160474
Csf2	-1.3760	0.00501	0.04459598
Ccl4	-1.2580	0.00115	0.017056477
Siglecg	-1.2162	0.0052	0.045382971
Il6	-1.2093	0.00327	0.033948487
Treml2	-1.0966	0.00528	0.045784691
Cd163	-1.0852	0.00182	0.023059989
Ccrl2	-1.0704	2.53E-06	0.000232561
Mmp12	-1.05495	1.78E-05	0.001018296
Lilrb4a	-1.0335	6.59E-05	0.002501365
Vegfa	-0.96557	0.00035	0.007515902
Mrc1	-0.81427	0.00276	0.030275915
Cd93	-0.78978	0.00078	0.013175012
Cd53	-0.76056	0.0005	0.00967682
Mrc2	-0.75703	0.00014	0.00411241
Ereg	-0.74863	0.00454	0.041930646

PDL1 (CD274)

0.0001

Log2FC= -0.48

CD206 (MRC1)

0.01

Log2FC= -0.46

Figure 6

Figure 7

Adenosine gene score

A

Ado score

○ ADEX
○ Immunogenic
○ Squamous
○ Pancreatic progenitor

B

Ado score

○ Squamous
○ Non-squamous

PDAC-specific Adenosine signature

C

Immunity			Hypoxia response		Tumour stroma		

Immune response / immune suppression:
Adora2b
Bhlhe40
Cd109
Cd200
Cd274
Ifit2
Ltbp2
P2ry2
Prdm1
Rarres2
Serpinb2
Serping1
Tnfrsf10a
Tnfsf9
Treml2

Inflammation:
Il1a
Il6
Nos2
Osm
Ptgs2

Hypoxia:
Hif1a
Hilpda
Hk2
Slc2a3
Slc2a1
Trarg1

Angiogenesis/vascular remodelling:
Adm
Pgf
Tfpi
Vegfd

ECM:
Cadm3
Col5a3
Col6a3
Itga2
Itga7
Itgb7
Mmrn2
Pecam1
Plau
Tnc

Growth / secreted factors:
Adipoq
Areg
Chrdl1
Ereg
Grem1
Pthlh

Invasion / Metastasis:
Adamts14
Adamts4
Mmp13
Mmp3
Mmp9
Serpine1

D

PDAC Progression Free Survival

high eADO signature: mPFS 13.05 months
low eADO signature: mPFS 18.25 months
HR (*logrank*): 1.54 (95% CI; 1.04-2.27)

p=0.02

Number at risk (n= 176 total)

high eADO sign.	103	19	3	0	0	0
low eADO sign.	73	21	12	6	2	0

E

PDAC Disease-specific survival

high eADO signature: mDSS 19.66 months
low eADO signature: mDSS *NR*
HR (*logrank*): 1.84 (95% CI; 1.16-2.93)

p=0.01

Number at risk (n= 170 total)

high eADO sign.	100	28	6	2	0	0
low eADO sign.	70	28	12	8	2	0