Dissociation of reliability, predictability, and heritability in fine- and coarse-scale functional connectomes during development

Erica L. Busch[1][†], Kristina M. Rapuano[1], Kevin Anderson[1], Monica D. Rosenberg[2], Richard Watts[1], BJ Casey[1], James Haxby[3], Ma Feilong[3][†]

[1]Department of Psychology, Yale University
[2]Department of Psychology, University of Chicago
[3]Department of Psychological and Brain Sciences, Dartmouth College

[†]Corresponding authors:
Erica L. Busch: Erica.Busch@yale.edu
Ma Feilong: feilong.ma@dartmouth.edu

Abstract

The functional connectome supports information transmission through the brain at various

spatial scales, from exchange between broad cortical regions to finer-scale connections that

underlie specific information processing mechanisms. In adults, while both the coarse- and fine-

scale functional connectomes predict cognition, the fine-scale can predict twice the variance.

Yet, past brain-wide association studies, particularly using large developmental samples, limit

their focus to the coarse connectome to understand the neural underpinnings of individual

differences in cognition. We examine resting-state fMRI (n=1,115 children, 389 twin pairs) and

use hyperalignment to access reliable information in the fine-scale connectome. Though

individual differences in the fine-scale connectome are more reliable than those in the coarse-

scale, they are less heritable. Further, the alignment and scale of connectomes influence their

ability to predict behavior: both scales equally predict more heritable cognitive traits, but the

fine-scale connectome predicts less heritable traits significantly better than the coarse-scale.

Introduction

The building blocks of the functional connectome are an interplay of genetics[1–3] and experience[4,5]. Functional connections at various spatial scales, ranging from coarse, global regional oscillations to fine, vertex-wise patterns of information transfer, account for related yet distinct types of information encoding and transmission[6,7]. While variation in functional connectivity can predict individual differences in aspects of memory[8], attention[9], and intelligence[10], among other cognitive traits[10–15], it is unclear how the spatial scales of functional connectomes differentially relate to genetic and experience-based cognitive traits during periods of biological and experiential development. To understand how this is reflected at the population level in the developing brain requires consideration of how we define functional connectivity information at both fine and coarse scales. In the current study, we disentangle more experience-based attributes of the functional connectome from genetic influences by estimating the heritability, reliability, and predictive validity of resting-state functional connectivity (RSFC) during development[16,17] in twins and unrelated children at different levels of the granularity of connectome organization.

Past studies using structural and functional magnetic resonance imaging (MRI) of twin samples have revealed strong genetic control over various measures of brain morphology[4,18–21] and RSFC[2,3,22–26]. Genetic influences on brain structure and function are associated with cognitive functioning[18], increase with age[18,27], and interact with the environment to sculpt an individual's functional connectome[1,18,27,28]. RSFC is a powerful tool for investigating individual differences in brain–behavioral associations, especially in clinical and developmental populations where task-based fMRI is often difficult[11,13,14,29]. Of neuroimaging phenotypes, recent studies indicate

that structural MRI is more heritable than resting-state fMRI, which is in turn more heritable than

task-based fMRI[19]. For this reason, we focus on resting-state functional connectivity (RSFC) to

understand the nature of individual differences in the brain and behavior in childhood.

Prior work has established heritability of RSFC between specific regions of interest[30] and whole-

brain oscillations between large cortical regions or networks[3,5,22,24,25]. In whole-brain studies,

functional connectivity is commonly calculated by dividing the cortex into parcels and collapsing

over the hundreds of vertices within a parcel, then correlating the average timeseries between

each pair of parcels. This approach aims to mitigate high-dimensionality, measurement

instabilities, and inter-individual differences in the brain's functional organization, and has been

used in prior studies relating RSFC and cognition in development[11,15,31]. Within developmental

samples, parcellations are believed to account for developmental differences in cortical anatomy

by coarsely aligning fMRI signals across subjects. However, group parcellations are defined on

average adult brains and assume that the same boundaries around brain regions exist in the same

location across all brains, regardless of inter-subject or developmental differences[32]. Individual-

defined parcellations better capture individualized functional topography[24,33–35], but still, typical

parcels may cover a larger surface area than is most informative and effects that may be present

for a subset of vertices in a parcel can be obscured by aggregating into a single parcel[32]. More

generally, both types of parcellations collapse over local information processing at the fine-scale,

which has been shown to reliably represent behaviorally-relevant information[12].

A useful alternative to group-average parcellations, functional alignment of fMRI data

maximizes the intersubject correspondence of cortical information processing while resolving

variation in cortical anatomy at the fine-scale. Reliable, predictive individual differences in fine-

scale signals are most apparent after resolving idiosyncrasies of individual functional

topographies with methods like hyperalignment[6,36]. In this study, modeling the developing

functional connectome at the fine-scale with hyperalignment allows two main affordances. First,

functional alignment improves between-subject correspondence of both cortical responses and

functional connectivity[36–40] and highlights reliable[41] and behaviorally-relevant[12] individual

differences. Second, hyperalignment allows us to disentangle idiosyncratic anatomical

topography from functional topography[6], and enables us to ask how individual differences in

RSFC relate to genetics and behavior, independent of anatomical similarity.

Cortical architecture changes rapidly and idiosyncratically across development[42], which prompts

important concerns about anatomical differences confounding our understanding of brain–

behavior relationships in children. Accessing the cortex's fine-scale functional topography

mitigates these concerns by retaining high dimensionality in a representational space

disentangled from brain anatomy. To date, genetic contributions to the fine-scale functional

connectome remain to be quantified, as well as their reliability and behavioral relevance during

development. The current study seeks to address the following questions: Does accessing the

fine-scale connectome tighten the link between functional connectivity, cognition, and genetics

shown in the coarse-scale connectome? Or is there a dissociation of genetic control over these

two scales of connectivity? Here, we use connectivity hyperalignment[38] to access and dissociate

fine-scale and coarse-scale RSFC in a cohort of 1,115 9–10 year-olds from the Adolescent Brain

Cognitive Development (ABCD) Study[®16] ([Figure 1]). We leverage machine learning approaches

to understand how connectome granularity relates to reliability, heritability, and neurocognitive

abilities in children.

Figure 1: Methods

A. Connectivity hyperalignment training procedure. For each parcel in the Glasser

parcellation (here, region 329 with 430 cortical vertices), connectivity matrices are computed as

the strength of connectivity (Pearson's correlation) between the activity timeseries of vertices

within the parcel (seeds; u) and the average activation across vertices in all other parcels (targets;

v). The connectivity matrices for 200 held-out subjects are used to train the hyperalignment

common model space for a given parcel. The model space is of dimensions $\mathbf{D} = [d_1...d_n]$ which

are weighted combinations of $\mathbf{U} = [u_1...u_n]$, the dimensions of each subject's connectivity

matrices in their native anatomical space. $\mathbf{D}$ represents the space where connectivity targets are

best aligned across training subjects. Then, connectomes for the 915 test subjects are projected

into **D** via subject-specific transformation matrices **R**, which detail a mapping from given subject

i's anatomical space U_i into **D**. This procedure is repeated to derive **R** for each test subject and

parcel. Then, the regional activation pattern for each parcel in anatomical space is mapped via **R**

into the region's common space to compute the hyperaligned timeseries for each subject, which

is then used to recompute connectivity matrices between hyperaligned seeds *v* and model

dimensions *d*. **B. Connectome granularity.** Fine-scale connectomes are multivariate patterns of

the connectivity weights between the time-series responses of all individual vertices in a given

parcel and the average time-series response of each other parcel in the brain. Coarse-scale

connectomes are vectors (univariate), representing the connectivity weight between the average

time-series response of a given parcel, averaged across all of its vertices, and those of each other

parcel in the brain.

Results

Establishing the reliability of individual differences

Variability in RSFC can reflect idiosyncrasies driven by cortical communication patterns

indicative of neurocognitive variability, but it can also reflect artifacts due to data acquisition

noise, movement[43,44], and functional–anatomical mismatch. In adult populations, brain

parcellations (e.g., Glasser parcellation[33]) average adjacent areas of cortex to improve signal-to-

noise and reduce the dimensionality of the data, and adult-defined parcellations are frequently

applied to development datasets[15,31]. This approach assumes functional–anatomical

correspondence will improve at the group level and that functional boundaries are constant

across development, and comes at the cost of lower granularity.

Here, we analyzed RSFC in a large cohort of 1,115 children, aged nine to ten, including 389 twin

pairs (see Table S1 for subject breakdown) using two methods of group alignment. The first uses

a standard approach to align data in a common space based on anatomical location (anatomical

alignment [AA]), while the second approach uses connectivity hyperalignment (CHA)[38] to learn

a common information space based on patterns of functional connectivity via an adapted

Generalized Procrustes Analysis. CHA allows for local remixing of vertex-wise functional

connectivity profiles (vertices' vectors of connectivity to targets) within an anatomically-

constrained region[36], thereby projecting idiosyncratic patterns of vertex connectivities to the same

target into a common template connectome[6]. In contrast with coarse alignment, which averages

time-series responses across a region's vertices to calculate a single connectivity strength with

each target region, CHA retains the vertex-by-vertex variation on connectivity strength to each

target region (Figure 1B). Within this common, fine-grained space, individual differences in

functional connectivity become more pronounced and reliable[41] and are more predictive of

cognitive traits[12] in adults.

To address the question of individual differences in fine-scale functional connectivity among

children, we present (to our knowledge) the first derivation of a child-defined and applied model

of shared brain function, which negates concerns about differences in cortical development or

anatomy that may confound modeling of functional connectivity patterns. To do this, we used

CHA to build a template common connectome for each parcel on a training sample of 200

unrelated children, counterbalanced across included sites and representative of demographics

within site (Table S1; Figures S1, S2, S3). We then hyperaligned each of the 915 remaining

connectomes to this common connectome. Note that this common information space is based on

children of the same age and demographics as the test sample, and that the test sample's data

played no role in deriving the common space. We addressed the relative effects of hyperalignment

versus simply higher-dimensional measurements by averaging the functionally-aligned CHA data

and by retaining the high-dimensional, anatomically aligned data, to have a full comparison of

connectomes that are aligned anatomically, at both the coarse-grained and fine-grained level, and

connectomes that are CHA-aligned, at both the coarse-grained and fine-grained level.

Since this is the first time a functional alignment algorithm has been trained and tested on a

developmental cohort, we first established that idiosyncrasies in cortical activity patterns were,

indeed, more reliable after fine-scale hyperalignment, as has been shown in adult cohorts[41]. We

analyzed the reliability of idiosyncrasies in functional connectivity patterns across connectome

alignment and granularity. Here, we define idiosyncrasies in terms of individual differences

matrices (IDMs), or the pairwise correlation distance between all pairs of subjects. IDMs capture

the dissimilarity structure (or idiosyncrasies) of functional connectivity patterns across subjects.

We assessed the reliability of these idiosyncrasies by splitting each subject's RS timeseries in half

to create two "pseudo-sessions" and calculating independent connectomes each split-half, then

computing an IDM for each pseudo-session. We then correlated the two pseudo-sessions IDMs to

measure the reliability of individual differences. This process was performed for anatomically-

aligned coarse-grained and fine-grained connectomes and for CHA-aligned coarse-grained and

fine-grained connectomes separately (see Individual differences matrix (IDM) reliability for more

information).

CHA, particularly at the fine scale, improved the reliability of individual differences in functional

connectivity over anatomical alignment alone (Figure 2). Across all parcels, reliability of coarse

connectomes increased from an average of $r = 0.34 \pm 0.1$ for AA to $r = 0.58 \pm 0.08$ for CHA.

Reliability of fine connectomes increased from an average of $r = 0.43 \pm 0.11$ for AA to $r = 0.86 \pm$

0.07 for CHA. After CHA, 100% of parcels at both scales showed greater reliability, and 99% of

parcels showed greater reliability at the fine scale than the coarse scale. Notably, all four types of

connectomes showed high split-half reliability, with CHA affording the greatest reliability. This

result establishes that hyperalignment improves access to reliable information within an

individual's functional connectome. Particularly at the fine scale, individual differences in

functional connectivity were highly reliable, affording confidence in subsequent results based on

these connectomes.

Reliability of individual differences in functional connectivity profiles

Figure 2. Reliability of individual differences in functional connectivity profiles

Procedure: The reliability of individual differences in functional connectivity profiles between

regions (defined with the Glasser parcellation[33]) was analyzed with data from 526 unrelated

subjects. Subjects' timeseries data for each parcel were split into two halves, which were

hyperaligned into the trained common space separately. For each region, split-half timeseries

data pre- and post-hyperalignment were used to compute functional connectivity profiles at the

fine and coarse scales. Then, individual differences matrices (IDMs) were computed across all

pairs of subjects for each independent split half. The IDMs for each pair of split halves were then

correlated (Pearson's r) for each region. Reliabilities of individual differences captured by

regional functional connectivity profiles are presented on the cortex for the coarse and fine scales

before and after hyperalignment. This approach is a variation of the "fingerprinting"[10,45]

approach, whereby measured individual differences are more reliable if a subject is more similar

to themselves across split-halves than to other subjects. **A.** Spatial distribution of reliability

scores before and after hyperalignment at the fine and coarse scales, as visualized on the cortical

surface for each parcel. **B.** The reliability of individual differences increases after

hyperalignment for all parcels. 99% of fine connectivity profiles (darker) are more reliable than

coarse ones (lighter) regardless of alignment type. **C.** Box-and-whisker plots showing the

distribution of IDM reliability scores across parcels within connectome type. Box shows first

quartile, median, and third quartile of distribution. Whiskers represent 1.5 times the interquartile

range, and points represent outlier values.

Genetics contribute most strongly to the coarse-scale connectome

We asked whether the more reliable fine-scale connectome simply reflected the same

information about cortical functional architecture at higher magnification or a different class of

information that may be more or less heritable or more or less reflective of experience. We

investigated this question by calculating the heritability of the two coarse-grained and the two

fine-grained connectomes. We hypothesized that heritable factors determine the skeletal

attributes of the connectome and are more evident at the coarse spatial scale[4,27,28], whereas the

detailed configurations of connectomes may develop under the influence of less heritable factors,

such as experience, and are reflected at a finer spatial scale.

The heritability analysis used RSFC data from 778 subjects comprising 219 dizygotic twin pairs

and 170 monozygotic twin pairs (see breakdown in Table S1). As in prior analyses, the

connectivity hyperalignment model was trained on 200 unrelated subjects, then the connectomes

of the 778 twin subjects were hyperaligned to this model. Functional connectomes of vertex-

granularity are an inherently high-dimensional phenotype. To explicitly model the heritability of

this phenotype, we used a multidimensional estimate of heritability[24] to model both the

phenotypic similarity matrix between participants as well as the genetic kinship matrix (see

Multidimensional heritability estimation for more details).

We find that both the anatomically-aligned and CHA-aligned coarse-grained connectomes were

more heritable than the more reliable fine-grained connectomes ($p < 0.0001$) (Figure 3). At the

coarse scale, anatomically-aligned functional connectomes are significantly more heritable than

hyperaligned connectomes (AA $h^2 = 0.198 \pm 0.05$; CHA $h^2 = 0.161 \pm 0.04$; $p < 0.0001$), with

96% of parcels showing greater heritability. These results indicate that connectivity information

aligned coarsely based on anatomical features are under strong genetic control, consistent with

previous studies showing genetic control over cortical anatomy in children[20] and network-level

RSFC[2].

Figure 3. Heritability of functional connectivity profiles

Procedure: Heritability of functional connectivity profiles between regions (defined with the

Glasser parcellation[33]) was estimated using a multidimensional heritability model[24] using resting-

state fMRI data from 778 9–10-year-olds (170 MZ pairs and 219 DZ pairs). Functional

connectivity between a given parcel and all other parcels in the brain was calculated at fine and

coarse granularities, before and after connectivity hyperalignment. The coarse-grained

connectivity profiles aligned based on cortical anatomy are the most heritable. **A.** Spatial

distribution of h^2-multi scores before and after hyperalignment at the fine and coarse scales, as

visualized on the cortical surface for each parcel. **B.** For both anatomically-aligned and

hyperaligned data, connectivity profiles are more heritable when calculated at the coarse scale

than the fine scale (99.6 percent of the time). **C.** At the coarse scale, the anatomically aligned

connectivity profiles are more heritable than the hyperaligned ones, but not at the fine scale,

which is under comparatively minuscule genetic control. Box-and-whisker plots showing the

distribution of h^2-multi across parcels within connectome type. Box shows first quartile, median,

and third quartile of distribution. Whiskers represent 1.5 times the interquartile range, and points

represent outlier values.

Prediction of neurocognitive abilities

We next asked whether the reliable individual differences in the fine-scale connectomes that

have low heritability reflect differences in cognitive abilities that are fostered by experience.

Compared with brain structure, complex psychological traits are less heritable, but cognitive

abilities are among the most heritable dimensions of human behavior[46]. Heritability (h^2) of

cognitive abilities varies by domain and increase with age, with higher estimates for general

cognitive ability ($h^2 = 0.54$–0.85 at 10–12 years[47] and $h^2 = 0.60$–0.73 at 8–21 years[48]) than for

specific cognitive abilities like learning/memory[49] ($h^2 = 0.18$–0.55 at 8–21 years[48] and $h^2 = 0.39$

at 18–67 years[50]), which tend to be more sensitive to experiential factors. In this proof-of-

principle analysis, we leveraged the variation in heritability between learning/memory and

general cognitive ability to examine the functional significance of our heritability and reliability

findings. Previous work[51] using data from the first release of baseline ABCD data to determine

components of higher-cognitive abilities during development calculated composite

neurocognition component scores using a Bayesian Probabilistic Principal Components Analysis

(BPPCA) model over the NIH Cognitive Toolbox tasks administered in the ABCD study and

released as a single score for each component per participant in the data repository. Here, we

focused on the composite scores for general cognitive ability and learning/memory. Since

learning/memory is regarded as less heritable than general cognitive ability, we leveraged the

variability of heritability in connectivity to predict variably-heritable cognitive phenotypes[48,49].

Given that CHA improves reliability of individual differences in the fine-scale connectome, we

predicted that CHA fine connectomes would capture more variance in both cognitive traits than

the CHA coarse, AA coarse, or AA fine connectomes. Moreover, we predicted that the most

heritable connectome (AA coarse) would reflect more variance in a more heritable cognitive trait

(general cognitive ability) relative to a less heritable one (learning/memory).

We used a principal components analysis ridge regression to predict scores on these two

cognitive domains from connectivity profiles at each scale. Models were trained using the

individual differences matrices (IDMs) across 526 unrelated subjects, computed for each scale,

alignment, and parcel. Crucially, predictions based on IDMs ensure that all models are receiving

data with the same number of features, regardless of connectome type, since IDMs are 526

subjects by 526 subjects where each cell is a pairwise correlation distance between the

connectomes of two subjects. We used a nested cross-validation procedure to tune

hyperparameters (number of PCs and α) then apply the model to predict loadings from unseen

subjects (see Neurocognition PC prediction analysis for more details). Model scores are

presented as R2 between predicted scores and true values, and score significance was assessed

by repeating this analysis with permuting behavioral scores (1,000 iterations per model) to create

a null distribution. Model scores were compared across connectome types using 10,000 bootstrap

iterations with resampling.

Overall, hyperaligned fine-scale connectomes best predicted individual differences in cognitive

performance (i.e., both general cognitive ability and learning and memory), and the degree to

which the coarse connectome also contributed to prediction was associated with the heritability

of the trait. The more heritable psychological trait of general cognitive ability was similarly

predicted by CHA fine, AA coarse, and AA fine connectomes (average R^2 ± s.d. across parcels:

CHA fine = 0.0133 ± 0.021; AA fine = 0.0134 ± 0.020; AA coarse = 0.0124 ± 0.018;

Bonferroni-corrected p-values reported from permutation test with 10,000 iterations: CHA fine

vs. AA fine p= 1.0; CHA fine vs. AA coarse p = 1.0; AA fine vs. AA coarse p = 0.39) (Figure 4),

indicating that the information content carried in each of these connectomes predicts general

cognitive ability. Compared with the other three connectomes, CHA coarse connectomes

predicted significantly less variance in general cognitive ability (R^2 = 0.006 ± 0.02; CHA coarse

vs. CHA fine p < 0.0001; CHA coarse vs. AA fine p < 0.0001; CHA coarse vs. AA coarse p <

0.0001), suggesting that functionally aligning fine-scale information then smoothing over the

functional information space actually diminishes the specificity of the information. After

performing permutation tests at each parcel and thresholding at p < 0.01, 42%, 42.5%, 30.5%,

and 43% of parcels predicted general cognitive ability significantly greater than chance at the

AA coarse, AA fine, CHA coarse, and CHA fine scales, respectively (Figure S6).

Learning/memory is regarded as a less heritable psychological trait ($h^2 = 0.36–0.56$)[48], so we

hypothesized that the information content of the coarse connectomes (the more heritable

information) would contribute less to the prediction of this trait relative to the fine connectomes

(the more reliably idiosyncratic information). Indeed, prediction of learning/memory scores were

significantly higher for the CHA fine connectome than the other connectome types (average R^2

± s.d. across parcels: CHA fine = 0.018 ± 0.012; CHA coarse = 0.011 ± 0.014; AA fine 0.008 ±

0.014; AA coarse 0.007 ± 0.014; Bonferroni-corrected p-values: CHA fine vs. CHA coarse p <

0.0001; CHA fine vs. AA fine p < 0.0001; CHA fine vs. AA coarse p < 0.0001; CHA coarse vs.

AA fine p = 0.007; CHA coarse vs. AA coarse p < 0.0001; AA fine vs. AA coarse p = 0.529)

(Figure 4). The gap in the percentage of parcels capturing a significant amount of variance was

also more pronounced for this trait; 30%, 33%, 39%, and 61% for AA coarse, AA fine, CHA

coarse, and CHA fine.

Figure 4: Prediction of neurocognitive PC loadings.

Distribution of R^2 values across all parcels for prediction of general cognitive ability (left) and

learning/memory (right). Histogram heights correspond to density of scores and lines represent

the mean of the score distribution. For the purpose of distribution matching, R^2 values are not

318   thresholded here; thresholded R^2 values mapped to the cortical surface are displayed in Figure

319   S6.

Discussion

In this study, we investigated whether heritable and experiential factors have dissociable effects

on the development of coarse- and fine-grained levels of functional organization in the

developing connectome. We hypothesized that the coarse-grained structure of connectivity

between cortical regions — the scaffolding of the connectome — would be under stronger

genetic control, whereas the fine-grained structure of local variations in connectivity would be

shaped more by individual-specific learning and experience. We improved access to individual

differences in fine-grained connectivity by applying connectivity hyperalignment (CHA) to

resting-state fMRI data in a large sample of children from the ABCD® dataset. Our study

presents the first example of defining and applying a functional common space to a

developmental cohort. This represents an important advancement in developmental

neuroimaging as it disentangles meaningful individual differences from misaligned cortical

anatomy or measurement noise.

Using a normative set of 200 children's brains, we built a fine-grained model connectome and

hyperaligned a separate set of 915 children to that template. Before hyperalignment, the

connectivity profiles of anatomically-aligned vertices exist in distinct, individual representational

spaces and are, therefore, both less reliable and less predictive of behavioral traits.

Hyperalignment resamples an individual's vertex-wise connectivity profiles into the model

connectivity profiles, which affords meaningful comparisons of fine-grained profiles across

individuals. Prior literature has relied upon anatomical alignment, or matching children's data to

a common anatomical template brain, then applying a parcellation (commonly defined on adult

brains) to extract an average (coarse) timeseries for each predefined region[11,15,31,52]. By averaging

connectivity profiles across neighboring vertices in a region, one can factor out misaligned fine-

grained variation to reveal the scaffolding of the coarse-grained connectome, but this obscures

the information contained in fine-grained variations. We considered multiple connectome types

— coarse- and fine-grained, pre- and post-CHA — to understand the effects of genetics and

experience at different scales of connectivity.

Using CHA, we revealed reliably idiosyncratic patterns of fine-scale functional connectivity. The

coarse scale connectome (AA coarse) shows only moderately reliable idiosyncrasies (mean

Pearson's $r = 0.34$), versus the CHA fine connectome's mean reliability of $r = 0.86$ ([Figure 2](#)).

Across the cortex, individual differences in fine-scale functional connectivity were more reliable

than in coarse-scale, suggesting that traditional, coarse alignment diminishes spatially-resolved

topographies of information transfer. By estimating the heritability of functional connectivity at

each scale and alignment in monozygotic and dizygotic twins, we showed that idiosyncrasies in

the fine-scale connectomes were dissociable from idiosyncrasies in the coarse scale connectome.

Multiscale heritability analysis of coarse scale, anatomically aligned connectomes attributed

about 20% (5.47–33.06%) of variance to genetic relatedness, which converged with our

expectations based on adult coarse RSFC estimates[2,3,24] and how neuroimaging phenotypes

increase in heritability with age[4,19]. At the fine scale, h^2 estimated dropped to around 10%

regardless of alignment technique (CHA: 3.1–19.2%, AA: 2.0–16.67%), suggesting that at the

fine scale, idiosyncrasies in RSFC were likely more associated with unique experience than

genetics ([Figure 3](#)). These results supported our hypothesis: whereas individual differences in

fine-grained connectivity after hyperalignment were dramatically *more* reliable than differences

in coarse-grained connectivity, they were dramatically *less* heritable, indicating a dissociation

between the spatial scales of heritable and reliable patterns of functional connectivity.

Our final analysis examined the behavioral relevance of this dissociation. The most reliably

idiosyncratic (CHA fine) and most heritable (AA coarse) connectomes were similarly predictive

of general cognitive ability, a canonically heritable cognitive trait. In contrast, the most reliably

idiosyncratic connectome (CHA fine) was most predictive of a less heritable cognitive trait

(learning/memory) ([Figure 4](#)). Past studies using coarse RSFC from the ABCD dataset were

limited in their ability to predict general cognitive ability and not learning/memory scores[11]. By

applying CHA, we uncovered individual-specific patterns of functional connectivity that better

predict a presumably less heritable, more experience-related cognitive trait (learning/memory).

Taken together, we demonstrate a link between reliable, heritable, and predictive scales of

functional connectivity and highlight the importance of nuanced, multi-scale connectivity

analyses for holistically understanding individual differences during development. Our findings

dovetail with the interactive specialization framework of functional brain development, which

states that the development of the functional connectome is not solely determined by a blueprint,

but the interaction of a heritable template of connections and individualized experience[53]. We

provide neural evidence in support of this framework: the coarse connectome serves as a

heritable blueprint which represents more heritable cognitive phenotypes. Fine-scale

connectomes capture *similar* information as the coarse (as AA coarse never out-predicts CHA

fine), but the organization of fine-scale information is remixed in a way that diverges from the

heritable blueprint and individualized, experience-driven phenotypes.

Recent work has recognized the need for massive numbers of subjects and samples to reliably

associate brain phenotypes with behavior, specifically in RSFC of developmental cohorts[31]. The

need for massive cohorts could be attributed to the specifics of neuroimaging phenotypes.

However, leveraging machine learning advances including multivariate models, functional

alignment, or individual-specific functional atlases as opposed to group-average atlases is one

approach to improving prediction[12,15,54,55,34,35,56]. Our study demonstrates the power of

hyperalignment as a tool for understanding developing individual differences in brain and

behavior. The hyperalignment model was trained on RSFC from 200 children and tested on a

separate cohort, using under 20 minutes of resting-state data per subject. The prediction results

assure that the increased reliability of individual differences was driven by generalizable, shared

signals in the hyperalignment model rather than overfitting to subject-specific signals or simply

higher-dimensional data, since CHA improved reliability over AA at the fine-scale as well. By

performing all subsequent analyses on pairwise (dis)similarity matrices, we further ascertain that

our results are driven by the information content represented in different connectomes, rather

than the dimensionality difference between coarse- and fine-grained connectomes.

This study should be interpreted in light of potential limitations. Though statistically significant,

our predictions accounted for relatively low variance in behavioral measures relative to similar

studies in adults[10,12,57]. While task-based FC is more stable and predictive of behavior than RSFC

among adults[9,12,38,58], this requires sufficient amounts of high-quality data, notably difficult to

collect in children[29,59]. RSFC measures appear to be reliable in the ABCD dataset[52], whereas the

task fMRI has known reliability limitations[29]. Furthermore, RSFC has been shown to be more

heritable than task-based FC[19], suggesting that the added cognitive load of a task introduces

idiosyncrasies in information processing less attributable to genetics similarity.

Our choice to focus on two composite scores of neurocognition (general cognitive ability and

learning/memory), previously identified in Thompson et al. (2019)[51], should be viewed as a

proof-of-principle analysis relating the heritability and reliability of individual differences in

neuroimaging phenotypes to differentially heritable behavioral phenotypes. H^2 of general

cognitive ability and learning/memory in the present cohort (Figure S4) were both higher and

more similar to one another than the values reported in the literature[48–50,60]. This could be

because our sample has a diverse racial background and restricted age range, in contrast to

previous research based on predominantly European descent[60–63] with a larger age range[47–50].

Our predictions align most closely with Mollon et al. (2021)[48], which considered a more racially-

diverse cohort, potentially enhancing generalizability beyond Caucasians, and an overall younger

sample than prior studies, overlapping with our sample demographics. We also replicated our

heritability results using data from the entire cohort (n = 9,519) to rule out potential concerns on

sampling bias (Figure S4). Across the tasks, those loading more strongly onto the general

cognitive ability component (*pic vocab, reading*) showed stronger heritability compared to tasks

loading more strongly onto the learning/memory component (*card sort, picture*) (Figure S5).

Notably, even with overlap in the two components (i.e., *list sorting* task loading on both), we still

see distinct brain-based predictions associated with heritability.

This paper presents several novel findings. First, hyperalignment can be used as a tool to

improve the reliability and behavioral relevance of functional connectivity in children. After

hyperalignment, reliability of individual differences in RSFC is improved at the fine-scale

relative to the coarse scale, but the heritability of these signals is weakened. Finally, we present a

potential application of this dissociation in the fine and coarse scale connectomes to show how

the heritability of RSFC relates to ability to predict behavior based on the heritability of

behavioral traits. With the substantial twin cohort, behavioral testing, and longitudinal

neuroimaging data included in the ABCD study, future work could investigate the connection

between heritability and predictivity of functional connectivity over time, as the heritability of

both neurocognitive measures and neuroimaging-based phenotypes increase with age[1,4,21,64].

Moreover, this approach could be used to investigate how functional topographies reflect

cognitive development over time, by hyperaligning a subject to themselves at multiple

timepoints. In conclusion, the current study shows that by breaking apart the pieces of the

functional connectome into their coarse and fine structures, we can better understand how these

scales interact to scaffold and instantiate reliable idiosyncrasies in brain and behavior during

development.

Materials and methods

Participants

We considered data from the 11,875 children included in the Adolescent Brain Cognitive

Development (ABCD) Study[®16] , data release 2.0.1. The ABCD Study is a longitudinal study

with 21 sites around the United States and aims to characterize cognitive and neural development

with measures of neurocognition, physical and mental health, social and emotional function, and

culture and environment. The ABCD Study obtained centralized institutional review board (IRB)

approval from the University of California, San Diego, and each site obtained local IRB

approval. Ethical regulations were followed during data collection and analysis. Parents or

caregivers provided written informed consent, and children gave written assent. Four leading

twin research centers at the University of Minnesota, Virginia Commonwealth University,

University of Colorado-Boulder, and Washington University in St. Louis comprise the ABCD

Twin Hub. Each site enrolled approximately 200 same-sex monozygotic or dizygotic twin pairs

as well as singletons[17]. Their inclusion in the ABCD study affords unique access to the causal

interrelation between genetics, environment, brain function, and cognition during

development[1,4,17].

The present study used imaging and behavioral data from a subset of n=1,115 subjects from the

original 11,875 subjects. Included subjects were enrolled in the University of Minnesota,

Washington University in St. Louis, and University of Colorado-Boulder sites, which all use 3T

Siemens MRI scanners. We excluded Virginia Commonwealth University, the fourth Twin Site,

as it uses a GE scanner, a potential confound in the current study[16]. ABCD study-wide exclusion

criteria include a diagnosis of schizophrenia, moderate to severe autism spectrum disorder,

intellectual disabilities, major and persistent neurological disorders, multiple sclerosis, sickle cell

disease, seizure disorders like Lennox-Gastaut Syndrome, Dravet Syndrome, and Landau

Kleffner Syndrome, or substance abuse disorders at time of recruitment. Subjects with mild

autism spectrum diagnosis, history of epilepsy, traumatic brain injury, and MR incidental

findings were excluded from the present analysis.

Resting state fMRI data collection

Structural and functional data from the three included sites were acquired using Siemens Prisma

3T scanners with a 32-channel head coil. Detailed acquisition parameters are previously

described in the literature[16,65]. Scan sessions consisted of a high-resolution (1 mm^3) T1-weighted

image, diffusion weighted images, T2-weighted spin echo images (1 mm^3), resting-state fMRI

(rs-fMRI), and task-based fMRI. We utilized rs-fMRI solely in this study. rs-fMRI data were

collected with an echo-planar imaging sequence with 2.4 mm voxels, TR = 800 ms, TE = 30 ms,

and multiband slice acceleration factor = 6. Participants completed up to four runs of 5-minute

resting state scans. Framewise integrated real-time MRI monitoring (FIRMM[66]) was used to

monitor subject head motion during data collection, and scan operators may have stopped

resting-state data collection after three runs if 12.5 min of low-motion resting-state data had been

collected.

Additional exclusions were made based on poor structural scan quality (determined with curated

data release 2.0.1 sheet *freesqc01.txt*) or low quality/high motion resting state fMRI (determined

as a score of zero for *fsqc_qc* or greater than one for *fsqc_qu_motion, fsqc_qu_pialover,*

fsqc_qu_wmunder, or *fsqc_qu_inhomohgeneity*), as reported in prior studies[54]. Participants with

fewer than 900 TRs of rs-fMRI data passing these measures for FreeSurfer reconstruction were

also excluded. ABCD Data Release 2.0.1 used FreeSurfer version 5.3 for quality control

checks[65]. After filtering, we used resting state fMRI and behavioral data from 1,115 unique

subjects (552 male), including 170 pairs of monozygotic (MZ) twins, 219 pairs of dizygotic (DZ)

twins, and 337 singletons. All twin pairs were same-sex, and all subjects were between 9 and 10

years old. Included singletons were also balanced for ancestry, ethnicity, sex, and pubertal

development with twins (Table S1).

Preprocessing

rs-fMRI data were downloaded via the DCAN Labs ABCD-BIDS Community Collection[67]

(NDA Collection 3165). This is a regularly updated dataset of ABCD Brain Imaging Data

Structure (BIDS)[68] version 1.2.0 pipeline inputs and derivatives, using source data from the

ABCD Study participants baseline year 1 arm 1 DICOM imaging data that passed initial

acquisition quality control from the ABCD Data Analysis and Informatics Center (DAIC)[65] and

retrieved from the NIMH Data Archive (NDA) share of ABCD fast-track data (NDA Collection

2573). DICOMs were converted to BIDS input data using Dcm2Bids[69], which reorganizes

NiftiImages produced with dcm2niix[70]. Raw images were preprocessed using the DCAN Labs

ABCD-BIDS MRI processing pipeline[71] (for details, see github.com/DCAN-Labs/abcd-hcp-

pipeline; osf.io/89pyd), which is based on the Human Connectome Project (HCP) Minimal

Preprocessing Pipeline[72] with additional modifications specifically for the ABCD dataset and

summarized below.

The first stage of the pipeline, PreFreeSurfer, performed brain extraction, alignment, and N4 bias

field correction on the T1w and T2w images. The second stage, FreeSurfer[73,74], segmented the

resulting T1w images and identified tissue boundaries to register to a FreeSurfer template and

produce brain masks. The third stage, PostFreeSurfer, used the brain masks to register T1w

images to MNI space using ANTs symmetric image normalization method[75]. Surfaces were then

transformed to standard space using spherical registration and converted to CIFTI format along

with the standardized brain volumes. Multimodal surface registration was performed with MSM-

sulc. The fMRIVolume stage performed functional image distortion correction using reverse

phase-encoded spin echo images to correct for local field inhomogeneities. Eta squared values

were computed for each image to a participant-level average of all field maps, and the pair with

the highest value (i.e., most representative of the average) was selected to avoid potential motion

confounds. Finally, the fMRISurface stage performed 2-mm full-width half-max spatial

smoothing.

DCAN BOLD Processing[76] was used to perform standard data processing. First, fMRI data were

demeaned and detrended with respect to time. Covariates of no interest were regressed from the

data, including mean white matter, cerebrospinal fluid, overall global signal, mean grayordinate

timeseries, and 6-movement variables (X,Y,Z translation and roll, pitch, yaw rotation).

Timeseries data were then band-pass filtered between 0.008 and 0.09 Hz using a 2nd order

Butterworth filter. Data were then filtered for respiratory motion for the frequencies (18,582 to

25.726 breaths per minute) of the respiratory signal, which has been shown to improve the

stability of framewise displacement (FD) estimates[77]. "Bad" frames, where motion exceeds FD

of 0.3 mm, were removed when demeaning and detrending such that denoising betas were only

calculated for "good" frames. For the band-pass filtering, interpolation was used to replace those

frames, so that pre-processing of the timeseries only includes "good" data but avoids aliasing due

to missing timepoints. Finally, the filtered timeseries were normalized and concatenated across

runs. The derivatives used in this analysis are *ses-baselineYear1Arm1_task-rest_bold_desc-

filtered_timeseries.dtseries.nii.

Connectivity hyperalignment

We used connectivity hyperalignment[38] to functionally-align fine-scale connectivity information

across subjects. Given the small quantity (1480 ± 142 TRs; mean ± std) of resting state fMRI

data available per subject, we chose to train the hyperalignment model common space using a

cohort of 200 subjects who were then excluded from further analysis. These 200 training subjects

were singletons enrolled at one of the three included sites and were matched with twin subjects

on gender, pubertal development, age, and race/ethnicity (Table S1). The remaining 915 subjects

(including the 389 twin pairs and 137 singletons) were used for subsequent analyses. Crucially,

the hyperalignment model space never saw data from any subjects included in analyses during

training.

We followed the same connectivity hyperalignment procedure for whole-cortex parcellation data

as in our previous study[12]. We used the average timeseries for each of the 360 regions in the

Glasser cortical parcellation[33] as connectivity targets and the vertices in each parcel as

connectivity seeds. For each parcel we computed a fine-scale connectivity matrix as the

correlation between each of the vertices within the parcel and the average regional timeseries of

the other 359 regions in the parcellation. For example, for a parcel with 430 vertices, its fine-

scale connectivity matrix would be a matrix of dimensions 430 by 359 correlation coefficients,

whereas its coarse-scale connectivity profile would be a vector of 359 correlation coefficients

(Figure 1B).

For each region[36], we trained a separate high-dimensional common model space based on the

fine-scale functional connectomes of the 200 training subjects. The dimensionality of the model

space is equal to the number of vertices, but the model dimensions correspond to shared

functional connectivity properties across individuals instead of anatomical locations[6]. After the

model dimensions were learned, we discarded training subjects' data from downstream analyses.

We derived invertible transformation matrices for the remaining subjects' connectomes into the

model space to map vertex timeseries from the anatomically-defined dimensions into the

functionally-defined dimensions.

Functional connectivity profiles

After aligning the test data into a common functional space, we used functional connectivity

profiles to summarize a brain region's correlation with the rest of the cortex. We defined fine-

scale functional connectivity profiles as above, as the correlation between the timeseries of all

59,412 vertices in the cortex with the average timeseries of each parcel, resulting in a matrix of

59,412 by 360 correlation coefficients per subject. We also defined coarse-scale functional

connectivity profiles as the correlation between the average timeseries for all pairs of parcels,

resulting in a matrix of 360 by 360 correlation coefficients per subject (Figure 1B). The fine-

scale and coarse-scale connectomes were computed on data pre- and post-hyperalignment for

each subject to evaluate the effect of functional alignment and connectome granularity on

heritability, reliability, and predictivity.

Individual differences matrix (IDM) reliability

Given the high levels of noise and low reliability associated with developmental neuroimaging

data[29,43,59,78,79] as well as smaller (vertex-resolution) connectivity seeds[7], we used hyperalignment

to improve the reliability of individual differences in our data. That is, by finding a space that

aligns patterns of functional connectivity across subjects (within anatomically constrained

parcels), we correct more thoroughly for anatomical variability and improve access to reliable

individual differences[6,56]. We calculated the reliability of functional connectomes before and

after hyperalignment as follows. For 526 unrelated subjects, we computed fine-scale functional

connectomes from split-half timeseries (742 ± 71 TRs). We derived transformation matrices to

rotate these connectomes into the common space separately, then aligned the split timeseries data

using those transformations. We then computed functional connectivity profiles, as explained

above, for each half separately. For each parcel at the fine and coarse scale separately, we used

the vectorized functional connectivity profiles before and after hyperalignment and computed

individual differences matrices (IDM) for each data split. These are subject-by-subject pairwise

dissimilarity matrices, where each value in the matrix is the correlation distance (1 - Pearson's r)

between two subjects' regional connectivity profiles. After computing IDMs, we took the upper

triangle of the IDM and correlated them (Pearson's r) to measure the reliability of the individual

differences in the functional connectomes across the split halves[41]. This metric reflects the

reliability of the idiosyncrasies in a subject's functional connectomes, where subjects are more

similar to themselves across split halves than they are to any other subject. We used a Mantel

test[80,81] to perform statistical tests on IDMs with 10,000 permutations.

Multidimensional heritability estimation

Functional connectomes at vertex-granularity are an inherently high-dimensional phenotype;

when dimensionality of the phenotype is collapsed, the reliability of individual differences

decreases[41]. To offer greater statistical power in analyzing the heritability of this high-

dimensional phenotype, we used a multidimensional estimate of heritability[24,25] to summarize the

degree to which functional connectivity profiles for each parcel are under genetic control. We

also investigated whether the alignment and granularity of the connectomes impact their

heritability, which would indicate that the information highlighted by that type of connectome

calculation is differentially impacted by genetics. Genetic kinship was estimated with SOLAR

Kinship2[82].

Neurocognition PC prediction analysis

Neurocognition principal components (neurocog PCs)[51] were derived from the neurocognitive

battery included in the ABCD study. The neurocognitive battery includes the NIH Toolbox

cognition measures covering episodic memory, executive function, attention, working memory,

processing speed, and language abilities. The neurocog PCs were derived with a Bayesian

Probabilistic Principal Components Analysis, which revealed a three-component solution. The

three components roughly corresponded to general cognitive ability, executive function, and

learning/memory. Since the NIH Toolbox tasks that canonically measure executive function did

not load onto the executive function component in this solution, we excluded this component

from our analyses, instead focusing on general cognitive ability and learning/memory. Tasks

loading onto the general cognitive ability component include Toolbox Picture Vocabulary,

Toolbox Oral Reading test, List Sort Working Memory task, and Little Man task. Tasks loading

onto the learning/memory component include the Toolbox Picture Sequence Memory task, the

RAVLT total number correct, and the List Sort Working Memory task. Heritability of composite

scores were estimated with SOLAR Polygenic[82] including sex, age, ethnicity, and site as

covariates.

We predicted individual subjects' scores onto the neurocognition principal components based on

individual differences in RSFC from our cohort of 526 unrelated subjects using a principal

component ridge regression adapted from our previous study[12]. For each parcel, scale, and

alignment, we computed subject-wise individual differences matrices (IDMs) of functional

connectivity over the entire resting-state timeseries. Next, we ran a principal components

analysis (PCA) over the IDM to decompose the IDM into main orthogonal dimensions along

which individuals' connectivity profiles differ from one another. We then trained a ridge

regression model to predict neurocog scores from the decomposed IDM. Models were trained

and tested using nested cross-validation, where the inner fold optimized regression model

parameters (including the number of IDM PCs retained and α) across three sub-folds and the

outer fold scored the selected model on a held-out subject. Candidate regression model

parameters were distributed evenly on a logarithmic scale, from 10 to 320 PCs and no PCs (no

dimensionality reduction), and α chosen from 121 values between 10^{-20} to 10^{40}. Models were

evaluated using the cross-validated coefficient of determination R^2, which shows the variance in

neurocog scores accounted for by the prediction models. To assess statistical significance of

model performance against chance, we used permutation testing with 1000 iterations of shuffling

the behavioral scores.

Data availability

The ABCD study is longitudinal, so the data repository changes over time and can be found here:

https://nda.nih.gov/abcd. In the current study, we used the ABCD data release 2.0.1, downloaded

via ABCD-BIDS Community Collection[67] NDA Collection 3165.

Code availability

ABCD data preprocessing code can be found here: https://github.com/DCAN-Labs/abcd-hcp-

pipeline. ABCD data processing code can be found here: https://github.com/DCAN-

Labs/dcan_bold_processing. Code for running the multiscale heritability analysis can be found

here: https://github.com/kevmanderson/h2_multi. Code for running the nested PCA ridge

regression was based off: https://github.com/feilong/IDM_pred. All analysis code specific to this

study will be released upon publication.

Acknowledgements

This project was supported by the National Science Foundation Graduate Research Fellowship

Award # 2139841 to Erica L. Busch.

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive

Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA).

This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and

follow them over 10 years into early adulthood. The ABCD Study is supported by the National

Institutes of Health and additional federal partners under award numbers U01DA041048,

U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037,

U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028,

U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025,

U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089,

U24DA041123, U24DA041147. A full list of supporters is available at

https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing

of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD

consortium investigators designed and implemented the study and/or provided data but did not

necessarily participate in analysis or writing of this report. This manuscript reflects the views of

the authors and may not reflect the opinions or views of the NIH or ABCD consortium

investigators.

This research also benefited from the ABCD Workshop on Brain Development and Mental

Health, supported by the National Institute of Mental Health of the National Institutes of Health

under Award Number R25MH120869 and by UG3-DA045251 from the National Institute of

Drug Abuse.

Author Contributions

Conceptualization, ELB, KMR, MDR, MF, JH, RW, and BJC; Methodology, ELB, MF, JH, and

KMA; Software, ELB, KMA, and MF; Validation, ELB, KMA, MF, and JH; Investigation, ELB;

Writing - Original Draft, ELB, MF, and BJC; Writing - Review & Editing, ELB, KMR, MDR,

RW, KMA, MF, JH, and BJC; Visualization, ELB and KMR; Supervision, MF, JH, and BJC;

Funding acquisition, BJC.

Declaration of interests

KMA is an employee of Neumora Therapeutics.

Supplemental tables and figures

Table S1: Cohort Breakdown

	Total Included	Monozygotic Pairs	Dizygotic Pairs	Unrelated Test	Unrelated Train
Total	1,115	170	219	526	200
Site 20	387	75	86	184	60
Site 21	371	34	81	166	90
Site 22	357	75	68	176	50
Age (mo) (*interview_age*)	121.6	122.5	122.0	121.6	120.7
% Female (*gender*)	50%	52%	50%	50%	50%
Ancestry: African (*AFR*)	0.127	0.092	0.118	0.113	0.204
Ancestry: European (*EUR*)	0.805	0.825	0.826	0.814	0.733
Ancestry: East Asian (*EAS*)	0.020	0.016	0.016	0.021	0.025
Ancestry: (*AMR*)	0.049	0.067	0.040	0.052	0.038
Ethnicity (*race_ethnicity*)	1.667	1.659	1.63	1.73	1.58

Figure S1: Participant information.

Left: Number of participants included from each of the three selected sites, separated by gender.

Right: Participants were broken into two cohorts for training/testing the CHA model. 200 total

unrelated participants reflective of within-site age, demographics, and ancestry (see below

figures; 60, 90, and 50 from each respective site) were selected to train the CHA model. The

remaining participants were used in subsequent tests.

Figure S2: Participant demographics.

The three sites significantly differed in mean demographic attributes, including a four-level race

model, parental education level, parental income, and Hispanic identity. To respect this diversity

and sample a wide range of demographics, we selected CHA training subjects to reflect sample

demographics along these four dimensions. Shown here are the distributions of participants'

scores, scaled to a 0–1 range for visualization, for the participants allocated for CHA train or test

cohorts. Significance of differences along each dimension between CHA cohorts were tested

with a two-tailed independent samples t-test and * denotes $p < 0.05$ after Bonferroni correction.

Figure S3: Participant genetic ancestry.

The three sites significantly differed in mean genetic ancestry. To respect this diversity and add

generalizability to our heritability analyses, we selected CHA training subjects to reflect site

sample demographics as best as possible. Shown here are the distributions of participants' scores

for the participants allocated for CHA train or test cohorts. Significance of differences along

each dimension between CHA cohorts were tested with a two-tailed independent samples t-test

and * denotes $p < 0.05$ after Bonferroni correction.

Figure S4: Heritability of neurocognition PC loadings

Heritability of neurocognition PC solution loadings for general cognitive ability and

learning/memory.

Left: H2r of general cognitive ability and learning/memory were computed using SOLAR

polygenic across participants in our twin cohort (full sample: n = 778; with complete data: n =

748). h^2 of general cognitive ability = 0.884 ± 0.01; h^2 of learning/memory = 0.837 ± 0.01

(reported as SOLAR polygenic h2r +/- standard error).

Right: To evaluate whether the behavioral heritability effect was due to sampling bias toward

the subjects included in the neuroimaging analysis, we recalculated H2r of the same components

using the full ABCD study sample (full sample: n = 11,875; with complete data n = 9,519).

Estimate of general cognitive ability = 0.830 ± 0.02, learning/memory = 0.800 ± 0.02.

Heritability of NIH toolbox tasks

Figure S5: Heritability of individual tasks loading onto neurocognition PCs

Heritability analysis of the individual NIH Toolbox tasks loading onto the general cognitive

ability and learning/memory components. Tasks loading more strongly onto the general

cognitive ability component (*pic vocab, reading*) showed stronger heritability whereas tasks

loading more strongly onto the learning/memory component (*card sort, picture*) showed weaker

heritability and a task loading onto both components evenly (*list sorting*) showed intermediate

heritability.

Abbreviation: PicVocab = Toolbox Picture Vocabulary Task (language skills and verbal

intellect); Reading = Toolbox Oral Reading Recognition Task (read and pronounce single

words), List = Toolbox List Sorting Working Memory Test; CardSort = Toolbox Dimensional

Change Card Sort Task (cognitive flexibility); Picture = Toolbox Picture Sequence Memory

Test.

Figure S6: Prediction of neurocognition displayed on cortical surface.

Prediction of individual differences in the general cognitive ability component (top) and the learning/memory component (bottom). R^2 values displayed for each parcel and significance assessed with 1000 permutations. Nonsignificant R^2 values ($p > 0.01$) displayed in gray.

References

1. Jansen, A. G., Mous, S. E., White, T., Posthuma, D. & Polderman, T. J. What twin studies tell us about the heritability of brain development, morphology, and function: a review. *Neuropsychol. Rev.* **25**, 27–46 (2015).

2. Glahn, D. C. *et al.* Genetic control over the resting brain. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 1223–1228 (2010).

3. Miranda-Dominguez, O. *et al.* Heritability of the human connectome: A connectotyping study. *Network Neuroscience* vol. 02 175–199 (2018).

4. Blokland, G. A. M., de Zubicaray, G. I., McMahon, K. L. & Wright, M. J. Genetic and Environmental Influences on Neuroimaging Phenotypes: A Meta-Analytical Perspective on Twin Imaging Studies. *Twin Res. Hum. Genet. Off. J. Int. Soc. Twin Stud.* **15**, 351–371 (2012).

5. Yang, Z. *et al.* Genetic and Environmental Contributions to Functional Connectivity Architecture of the Human Brain. *Cereb. Cortex N. Y. N 1991* **26**, 2341–2352 (2016).

6. Haxby, J. V., Guntupalli, J. S., Nastase, S. A. & Feilong, M. Hyperalignment: Modeling shared information encoded in idiosyncratic cortical topographies. *eLife* **9**,.

7. Haak, K. V., Marquand, A. F. & Beckmann, C. F. Connectopic mapping with resting-state fMRI. *NeuroImage* **170**, 83–94 (2018).

8. Lin, Q., Yoo, K., Shen, X., Constable, T. R. & Chun, M. M. Functional Connectivity during Encoding Predicts Individual Differences in Long-Term Memory. *J. Cogn. Neurosci.* **33**, 2279–2296 (2021).

9. Rosenberg, M. D. *et al.* A neuromarker of sustained attention from whole-brain functional connectivity. *Nat. Neurosci.* **19**, 165–171 (2016).

10. Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat. Neurosci.* **18**, 1664–1671 (2015).

11. Sripada, C. *et al.* Prediction of neurocognition in youth from resting state fMRI. *Mol. Psychiatry* **25**, 3413–3421 (2020).

12. Feilong, M., Guntupalli, J. S. & Haxby, J. V. The neural basis of intelligence in fine-grained cortical topographies. *eLife* vol. 10 e64058 (2021).

13. Cui, Z. *et al.* Individual Variation in Functional Topography of Association Networks in Youth. *Neuron* vol. 106 340-353.e8 (2020).

14. Dosenbach, N. U. F. *et al.* Prediction of Individual Brain Maturity Using fMRI. *Science* **329**, 1358–1361 (2010).

15. Chen, J. *et al.* Shared and unique brain network features predict cognitive, personality, and mental health scores in the ABCD study. *Nat. Commun.* **13**, 2217 (2022).

16. Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Dev Cogn Neurosci* vol. 32 43–54 (2018).

17. Iacono, W. G. *et al.* The utility of twins in developmental cognitive neuroscience research: How twins strengthen the ABCD research design. *Dev. Cogn. Neurosci.* **32**, 30–42 (2017).

18. Brouwer, R. M. *et al.* Heritability of brain volume change and its relation to intelligence. *NeuroImage* **100**, 676–683 (2014).

19. Elliott, L. T. *et al.* Genome-wide association studies of brain imaging phenotypes in UK Biobank. *Nature* **562**, 210–216 (2018).

20. Peper, J. S. *et al.* Heritability of regional and global brain structure at the onset of puberty: A magnetic resonance imaging study in 9-year-old twin pairs. *Hum. Brain Mapp.* **30**, 2184–2196 (2009).

21. Thompson, P. M. *et al.* Genetic influences on brain structure. *Nat. Neurosci.* **4**, 1253–1258 (2001).

22. Colclough, G. L. *et al.* The heritability of multi-modal connectivity in human brain activity. *eLife* vol. 6 e20178 (2017).

23. Barber, A. D., Hegarty, C. E., Lindquist, M. & Karlsgodt, K. H. Heritability of Functional Connectivity in Resting State: Assessment of the Dynamic Mean, Dynamic Variance, and Static Connectivity across Networks. *Cereb. Cortex N. Y. NY* **31**, 2834–2844 (2021).

24. Anderson, K. M. *et al.* Heritability of individualized cortical network topography. *Proc. Natl. Acad. Sci.* **118**, e2016271118 (2021).

25. Ge, T., Holmes, A. J., Buckner, R. L., Smoller, J. W. & Sabuncu, M. R. Heritability analysis with repeat measurements and its application to resting-state functional connectivity. *Proc. Natl. Acad. Sci.* **114**, 5521–5526 (2017).

26. Fu, Y. *et al.* Genetic influences on resting-state functional networks: A twin study. *Hum. Brain Mapp.* **36**, 3959–3972 (2015).

27. Schmitt, J. E. *et al.* The dynamic role of genetics on cortical patterning during childhood and adolescence. *Proc. Natl. Acad. Sci.* **111**, 6774–6779 (2014).

28. Lenroot, R. K. *et al.* Differences in genetic and environmental influences on the human cerebral cortex associated with development during childhood and adolescence. *Hum. Brain Mapp.* **30**, 163–174 (2009).

29. Kennedy, J. T. *et al.* Reliability and stability challenges in ABCD task fMRI data. *NeuroImage* **252**, 119046 (2022).

30. Achterberg, M. *et al.* Distinctive heritability patterns of subcortical-prefrontal cortex resting state connectivity in childhood: A twin study. *NeuroImage* **175**, 138–149 (2018).

31. Marek, S. *et al.* Reproducible brain-wide association studies require thousands of individuals. *Nature* **603**, 654–660 (2022).

32. Bandettini, P. A. *et al.* The challenge of BWAs: Unknown unknowns in feature space and variance. *Med* (2022).

33. Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* vol. 536 171–178 (2016).

34. Kong, R. *et al.* Spatial Topography of Individual-Specific Cortical Networks Predicts Human Cognition, Personality, and Emotion. *Cereb. Cortex* **29**, 2533–2551 (2019).

35. Kong, R. *et al.* Individual-Specific Areal-Level Parcellations Improve Functional Connectivity Prediction of Behavior. *Cereb. Cortex* **31**, 4477–4500 (2021).

36. Haxby, J. V. *et al.* A Common, High-Dimensional Model of the Representational Space in Human Ventral Temporal Cortex. *Neuron* **72**, 404–416 (2011).

37. Guntupalli, J. S. *et al.* A Model of Representational Spaces in Human Cortex. *Cereb. Cortex* **26**, 2919–2934 (2016).

38. Guntupalli, J. S., Feilong, M. & Haxby, J. V. A computational model of shared fine-scale structure in the human connectome. *PLOS Comput. Biol.* **14**, e1006120 (2018).

39. Nastase, S. A., Liu, Y.-F., Hillman, H., Norman, K. A. & Hasson, U. Leveraging shared connectivity to aggregate heterogeneous datasets into a common response space. *NeuroImage* **217**, 116865 (2020).

40. Busch, E. L. *et al.* Hybrid hyperalignment: A single high-dimensional model of shared information embedded in cortical patterns of response and functional connectivity. *NeuroImage* **233**, 117975 (2021).

41. Feilong, M., Nastase, S. A., Guntupalli, J. S. & Haxby, J. V. Reliable individual differences

in fine-grained cortical functional architecture. *NeuroImage* **183**, 375–386 (2018).

42. Paus, T., Keshavan, M. & Giedd, J. N. Why do many psychiatric disorders emerge during adolescence? *Nat. Rev. Neurosci.* **9**, 947–957 (2008).

43. Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *NeuroImage* **59**, 2142–2154 (2012).

44. Van Dijk, K. R. A., Sabuncu, M. R. & Buckner, R. L. The influence of head motion on intrinsic functional connectivity MRI. *NeuroImage* **59**, 431–438 (2012).

45. Noble, S. *et al.* Influences on the Test–Retest Reliability of Functional Connectivity MRI and its Relationship with Behavioral Utility. *Cereb. Cortex* **27**, 5415–5429 (2017).

46. Plomin, R. & Deary, I. J. Genetics and intelligence differences: five special findings. *Mol. Psychiatry* **20**, 98–108 (2015).

47. Bouchard, T. J. Genetic Influence on Human Psychological Traits: A Survey. *Curr. Dir. Psychol. Sci.* **13**, 148–151 (2004).

48. Mollon, J. *et al.* Genetic influence on cognitive development between childhood and adulthood. *Mol. Psychiatry* **26**, 656–665 (2021).

49. Plomin, R. & Spinath, F. M. Intelligence: Genetics, Genes, and Genomics. *J. Pers. Soc. Psychol.* **86**, 112 (20040112).

50. Fletcher, J. M., Marks, A. D. G., Hine, D. W. & Coventry, W. L. Heritability of Preferred Thinking Styles and a Genetic Link to Working Memory Capacity. *Twin Res. Hum. Genet.* **17**, 526–534 (2014).

51. Thompson, W. K. *et al.* The structure of cognition in 9 and 10 year-old children and associations with problem behaviors: Findings from the ABCD study's baseline

neurocognitive battery. *Developmental Cognitive Neuroscience* vol. 36 100606 (2019).

52. Marek, S. *et al.* Identifying reproducible individual differences in childhood functional brain networks: An ABCD study. *Dev. Cogn. Neurosci.* **40**, (2019).

53. Johnson, M. H. Interactive Specialization: A domain-general framework for human functional brain development? *Dev. Cogn. Neurosci.* **1**, 7–21 (2011).

54. Rosenberg, M. D. *et al.* Behavioral and Neural Signatures of Working Memory in Childhood. *J. Neurosci.* **40**, 5090–5104 (2020).

55. DeYoung, C. G. *et al.* Reproducible between-person brain-behavior associations do not always require thousands of individuals. Preprint at https://doi.org/10.31234/osf.io/sfnmk (2022).

56. Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI. *Trends Cogn Sci* vol. 20 425–443 (2016).

57. Dubois, J., Galdi, P., Paul, L. K. & Adolphs, R. A distributed brain network predicts general intelligence from resting-state human neuroimaging data. *Philos. Trans. R. Soc. B Biol. Sci.* **373**, 20170284 (2018).

58. Greene, A. S., Gao, S., Scheinost, D. & Constable, R. T. Task-induced brain state manipulation improves prediction of individual traits. *Nat. Commun.* **9**, 2807 (2018).

59. Satterthwaite, T. D. *et al.* Impact of in-scanner head motion on multiple measures of functional connectivity: Relevance for studies of neurodevelopment in youth. *NeuroImage* **60**, 623–632 (2012).

60. Need, A. C. & Goldstein, D. B. Next generation disparities in human genomics: concerns and remedies. *Trends Genet.* **25**, 489–494 (2009).

61. Savage, J. E. *et al.* Genome-wide association meta-analysis in 269,867 individuals identifies

new genetic and functional links to intelligence. *Nat. Genet.* **50**, 912–919 (2018).

62. Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* **538**, 161–164 (2016).

63. Grasby, K. L. *et al.* The genetic architecture of the human cerebral cortex. *Science* **367**, eaay6690 (2020).

64. Haworth, C. M. A. *et al.* The heritability of general cognitive ability increases linearly from childhood to young adulthood. *Mol. Psychiatry* **15**, 1112–1120 (2010).

65. Hagler, D. J. *et al.* Image processing and analysis methods for the Adolescent Brain Cognitive Development Study. *NeuroImage* **202**, 116091 (2019).

66. Dosenbach, N. U. *et al.* Real-time motion analytics during brain MRI improve data quality and reduce costs. *Neuroimage* **161**, 80–93 (2017).

67. Feczko, E. *et al.* Adolescent Brain Cognitive Development (ABCD) Community MRI Collection and Utilities. 2021.07.09.451638 Preprint at https://doi.org/10.1101/2021.07.09.451638 (2021).

68. Gorgolewski, K. J. *et al.* The brain imaging data structure, a format for organizing and describing outputs of neuroimaging experiments. *Sci. Data* **3**, 160044 (2016).

69. dcm2bids. (2022).

70. Li, X., Morgan, P. S., Ashburner, J., Smith, J. & Rorden, C. The first step for neuroimaging data analysis: DICOM to NIfTI conversion. *J. Neurosci. Methods* **264**, 47–56 (2016).

71. dsturge, Earl, E. & kathy-snider. DCAN-Labs/abcd-hcp-pipeline: release v0.0.0. (2019) doi:10.5281/zenodo.2587210.

72. Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage* **80**, 105–124 (2013).

73. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *NeuroImage* **9**, 179–194 (1999).

74. Fischl, B. FreeSurfer. *NeuroImage* **62**, 774–781 (2012).

75. Avants, B. B. *et al.* A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage* **54**, 2033–2044 (2011).

76. dcan signal processing. (2022).

77. Fair, D. A. *et al.* Correction of respiratory artifacts in MRI head motion estimates. *NeuroImage* **208**, 116400 (2020).

78. Birn, R. M. *et al.* The effect of scan length on the reliability of resting-state fMRI connectivity estimates. *NeuroImage* **83**, 550–558 (2013).

79. Fair, D. *et al.* Distinct neural signatures detected for ADHD subtypes after controlling for micro-movements in resting state functional connectivity MRI data. *Front. Syst. Neurosci.* **6**, (2013).

80. Mantel, N. Ranking Procedures for Arbitrarily Restricted Observation. *Biometrics* **23**, 65–78 (1967).

81. Carr, J. mantel. (2021).

82. Almasy, L. & Blangero, J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am. J. Hum. Genet.* **62**, 1198–1211 (1998).

83. Falconer, D. S. *Introduction to quantitative genetics*. (Pearson Education India, 1996).