How important is budding speciation for comparative studies?

Daniel S. Caetano[1,2,*] and Tiago Bosisio Quental[2]

[1] *Department of Biological Sciences, Towson University, Towson, MD, 21252, USA*

[2] *Department of Ecology, University of São Paulo, São Paulo, SP, 05508-090, Brazil*

* Author for correspondence: dcaetano@towson.edu

Abstract: The acknowledgment of evolutionary dependence among species has fundamentally

changed how we ask biological questions. Phylogenetic models became the standard approach for

studies with three or more lineages, in particular those using extant species. Most phylogenetic

comparative methods (PCMs) translate relatedness into covariance, meaning that evolutionary

changes before lineages split should be interpreted together whereas after the split lineages are

expected to change independently. This clever realization has shaped decades of research. Here we

discuss one element of the comparative method often ignored or assumed as unimportant: if nodes

of a phylogeny represent the dissolution of the ancestral lineage into two new ones or if the ancestral

lineage can survive speciation events (i.e., budding). Budding speciation is often reported in

paleontological studies, due to the nature of the evidence for budding in the fossil record, but it is

surprisingly absent in comparative methods. Here we show that many PCMs assume that divergence

happens as a symmetric split, even if these methods don't explicitly mention this assumption. We

discuss the properties of trait evolution models for continuous and discrete traits and their adequacy

under a scenario of budding speciation. We discuss the effects of budding speciation under a series

of plausible evolutionary scenarios and show when and how these can influence our estimates. We

also propose that long-lived lineages that have survived through a series of budding speciation

events and given birth to multiple new lineages can produce evolutionary patterns that challenge our

intuition about the most parsimonious history of trait changes in a clade. We hope our discussion

can help bridge comparative approaches in paleontology and neontology as well as foster awareness

about the assumptions we make when we use phylogenetic trees.

Keywords: budding speciation; diversification; fossil record; model adequacy; phylogenetic

comparative methods; speciation mode.

Phylogenetic trees are the main representation of evolutionary relationships among lineages

and stand as a symbol of evolutionary thought. However, their relatively simple structure is not

capable of informing us about all aspects of evolution. Processes such as horizontal gene transfer,

hybridization, and introgression can produce complex evolutionary relationships that challenge the

explanatory power of bifurcating phylogenies (Philippe and Douady, 2003; Mallet et al., 2016;

Bastide et al., 2018). Interestingly, the graphical representation of bifurcating trees could even

influence how we think (Baum et al., 2005). A node connecting one ancestral branch to two new

ones can suggest evolutionary histories much simpler than what we observe in nature and potentially

downplay important aspects of macroevolution. For instance, if read literally, bifurcating

phylogenies can be seen as speciation events that happened due to the split of an ancestral lineage

into two new ones, coincident with the extinction, or dissolution, of the ancestral lineage (Meier and

Willmann, 2000; Bokma, 2008). This view makes the concept of a lineage synonymous with a

branch of a phylogenetic tree. However, evidence from empirical systems shows the mode of

speciation can be varied and often complex (Rosenblum et al., 2012), including frequent instances of

budding speciation (Wagner, 1998; Funk and Omland, 2003; Gottlieb, 2004; Crawford, 2010;

Anacker and Strauss, 2014; Otero et al., 2019; Patsis et al., 2021).

The paleontological literature often adopts the representation of budding trees (e.g., Raup

and Gould, 1974; Raup, 1985; Foote, 1996; Wagner, 1998; Benton and Pearson, 2001) that inform

which branches are new lineages and which are the continuation of the ancestral lineage. Budding is

recognized in the fossil record as a cladogenetic event in which a new lineage appears as a

branching of an older lineage that can still be found after the speciation event (e.g., Foote, 1996).

Although budding speciation is commonly reported in paleontology, it is rarely incorporated into

phylogenetic comparative models (PCMs). Bokma and colleagues have developed a series of PCMs

incorporating budding and the effect of punctuated equilibrium (Bokma 2002; Bokma, 2008; Matilla

and Bokma, 2008; Monroe and Bokma, 2009; Bokma, 2010; Jansen et al., 2022). Unfortunately,

these and other similar methods (Bartoszek, 2014; Bartoszek, 2020, Pagel et al., 2022) have not been

widely used in the PCM literature, perhaps due to the perception that bifurcating molecular

phylogenies show no evidence of budding speciation. Here we bring a different, and perhaps

controversial, point of view; that budding speciation can be common, that it might affect inferences

of trait evolution, that it can be detected in molecular phylogenies, and it should be considered in

PCMs even when there is no information from the fossil record evidencing its role in the

diversification of the group.

There has not been a consensus about the effect of budding speciation on our estimates of the

tempo and mode of trait evolution using PCMs. Bokma (2008) implemented a trait evolution model

to estimate the contribution from saltational and anagenetic changes, however, there is no

investigation of the impact of the saltational process in inferences using PCMs that do not

accommodate such effects. De Lisle and colleagues (2021) explored the effect of shifts in the

adaptive optima on extinction rates using a model based on population-level dynamics and showed

that lineages would rarely survive peak shifts and those occupying stable optima are expected to

have a higher chance of survival. In turn, Duchen and colleagues (2021) studied how cladogenesis

changes the average species phenotype using individual-level simulations and showed that new

lineages budding off from an ancestral population can show a significant phenotypic deviation due

to neutral processes. Combined, these results point to the idea that long-lived evolutionary lineages

might occupy stable optima (Eldredge and Gould, 1972; Gould and Eldredge, 1993; also see

Goldberg and Foo, 2020) whereas new lineages may bud off with distinct phenotypes, due to the

effect of cladogenesis on the species trait and selection to occupy a distinct region of the

morphospace (Eldredge and Gould, 1972; Gould and Eldredge, 1993; Bokma, 2008; Matilla and

Bokma, 2008).

Here we discuss in which circumstances budding speciation can affect our estimates of the

tempo and mode of trait evolution. More specifically, we review the properties of PCMs of trait

evolution that dictate whether or not budding speciation can influence our conclusions. In our view,

most PCMs of trait evolution were developed with the strong assumption that speciation is

symmetric, lineages cannot continue after speciation events, and there is no effect of budding in

molecular phylogenies. Here we discuss how budding speciation can bias our estimates using

simulations as an argumentative guide to our narrative. Distinct from previous studies (Matilla and

Bokma, 2008; De Lisle et al., 2021; Duchen et al., 2021; Crouch et al., 2021 among others), our

discussion focuses on cases in which budding speciation is not considered when using PCMs.

What is budding and how to recognize it?

Budding is defined as a speciation event in which the new species co-occurs in time with its

ancestral lineage (Foote, 1996), meaning the ancestral lineage continues to exist after speciation. We

use the term progenitor lineage to help differentiate the parent from the daughter lineage (see

Gottlieb, 2004). A progenitor lineage is a lineage that has given birth to one or more new lineages

through budding. Budding speciation is inherent in Mayr's (1942) concept of speciation via

peripatric speciation. Also, as discussed by Eldredge and Gould (1972) and Grant (1981), drift likely

has an important impact on small founder populations and, naturally, will have consequences for

trait evolution (see De Lisle et al., 2021). Perhaps due to its intrinsic role in diversification, the

neontological literature has attributed different nomenclature to what is fundamentally budding

speciation. In this section, we discuss how budding speciation has been recognized in the literature,

which patterns might be the result of past budding events, and how budding can be detected using

data from extant species. The reader will note that we attribute a variety of processes to the effect of

budding speciation. Indeed, one of the main goals of this discussion is to bring awareness to the role

of budding speciation in studies of macroevolution and how it connects to multiple patterns we

observe in phylogenetic trees.

In the absence of the fossil record, budding has been recognized as a new lineage formed

within or at the edge of the ancestral lineage (Anacker and Strauss, 2014) or as a biological cause of

paraphyletic species (Funk and Omland, 2003; also see Fig. S1). This has also been associated with

the hypothesis of Punctuated Equilibrium (Eldredge and Gould, 1972; Gould and Eldredge, 1993),

since peripheral populations can become isolated and speciate, leading to fast trait evolution due to a

shift in the optima for the traits (Mayr 1942; Eldredge and Gould, 1972; Grant, 1981; Gould and

Eldredge, 1993; Bokma, 2008; Matilla and Bokma, 2008; De Lisle et al., 2021). Budding speciation

is considered to be opposed by bifurcation—the split of an ancestral lineage into two new ones (Fig.

S1). Hagen and colleagues (2015), for example, utilized the term *symmetrical* speciation to capture

the role of allopatric speciation and opposed it to *asymmetrical* speciation which represents

peripatric speciation with the continuation of the progenitor species—thus, budding. Although these

authors use distinct nomenclature, each is an example of budding. The paleontological literature

suggests budding is a common evolutionary pattern and some argue it represents the majority of

speciation events observed in deep time. For example, Wagner (1998) used budding speciation to

estimate a phylogeny of hyenas that provided a better fit to the stratigraphic history of the fossil

record. Aze and colleagues (2011) reconstructed a large phylogeny of macroperforate foraminifera

in which most cladogenetic events were recognized as budding through analysis of morphological

characters. Bapst and Hopkins (2017) applied an explicit probabilistic model to date a phylogeny of

trilobites and also show that budding events are often supported by the fossil record. Similarly,

Parins-Fukuchi (2021) re-evaluated the diversification of hominins and suggests the occurrence of

budding speciation events. In contrast, there is little to no mention of budding speciation in the

neontological literature which, in our view, creates an undesirable disconnect between paleontology

and neontology. One could argue that this absence is due to the impossibility of detecting budding

using molecular phylogenies, however, as we discuss below, we disagree with this statement.

If budding speciation is frequent, we expect to recover recent events of budding using

molecular data. When a new lineage buds off from its ancestral, the progenitor species becomes

paraphyletic (Funk and Omland, 2003; see Fig. S1B). The advantage of neontological data is that

molecular phylogenies can show evidence of budding independent of the use of morphological

divergence to estimate the tree, which is necessary to both estimate phylogenies and detect

speciation based on fossil remains (Foote, 1996; Wagner, 1998; Bapst, 2013). If the new lineage

maintains cohesion and does not go extinct (see De Lisle et al., 2021) or is not reabsorbed via

hybridization with the progenitor (Taylor et al., 2006; Richmond and Jockush, 2007; Behm et al.,

2010; Lackey and Boughman, 2017), gene flow among the populations of the progenitor should

complete sorting and the progenitor and daughter lineages will eventually become sister species in

estimated molecular phylogenies—erasing the signal of budding. Thus, budding speciation can be

detected using molecular phylogenies, but its signal disappears over time whereas, in the fossil

record, the information is preserved if the record is reasonably complete. Otero and colleagues

(2019) show an interesting case in *Iberodes* plants which underwent two events of budding within

the last 5 million years. In both instances, the new lineage evolved distinct morphological and

ecological traits (Otero et al., 2019). *Iberodes* has inland and coastal species and the potential

change in selective pressure together with the peripheral distribution of the younger coastal lineages

likely were key factors for budding divergence. Similarly, Papuga and colleagues (2018) show

peripheral plant populations that have lower niche breadth (i.e., are more specialized) than central

populations as well as divergence in ecological traits (i.e., soil parameters), both factors that can

cause budding by ecological speciation. Strong evidence for budding speciation was also detected

from molecular phylogenies by Baldwin (2005), showing that *Layia glandulosa* (a Compositae

plant) is the progenitor species for *L. discoidea*. Anacker and Strauss (2014) tested 71 sister pairs

and demonstrated that young divergences frequently show overlapping and asymmetrical ranges—

another indication of budding speciation. This asymmetry was not detected among older clades,

suggesting the signal of budding on the geographic distribution of sister pairs is lost as lineages get

older. We also note that taxonomic revisions that re-name paraphyletic species into several new

species, also erase the signal of budding.

If budding is frequent, and we suspect it is, it can be an important factor in understanding

trait evolution because peripheral populations can show distinct mean phenotypic values (Papuga et

al., 2018) and divergence through budding can generate new lineages with distinct average

phenotypes (Gottlieb, 2004; Duchen et al., 2021) and evolutionary trajectories (De Lisle et al.,

2021). If we assume molecular phylogenies are literal bifurcating trees, despite the evidence for

budding speciation, then PCMs might be based on inadequate assumptions. In the next two sections,

we visit the most popular PCMs and discuss scenarios in which the presence of budding would, or

would not, affect our estimates.

When budding doesn't matter

Raup (1985) stated that budding should not influence estimates of net diversification rate

because the addition or subtraction of lineages at any given time would be perceived similarly if we

represent a phylogeny either by budding or bifurcation. This question has been revisited by Bapst

and Hopkins (2017) and Crouch et al. (2021), both showing that budding can change divergence

time estimation and alter estimates of the accumulation of lineages through time (also see Wagner,

1998). Thus, budding should not influence the net diversification rate only if the true dated tree is

known, otherwise, changes in divergence time estimation can potentially impact estimates of

diversification down the line.

With respect to models of trait evolution, budding should not influence our estimates if

changes happening at any point in time, and at any branch of the phylogeny, are independent of the

prior history of the lineage and their ancestors. Two important simplifications were introduced when

Felsenstein described the method of independent contrasts (1981; 1985); trait changes happen

independently in each branch of the phylogeny and evolutionary changes at each point along a

branch are independent and identically distributed (iid). Most models of trait evolution share these

assumptions (see review in O'Meara, 2012; Pennell and Harmon, 2013). However, most, if not all,
PCMs were not created with the intent to accurately describe evolution in a mechanistic way, and
the use of simplifications does not mean we assume evolution follows these rules.

If models of trait evolution that assume a homogeneous process across all branches of the
tree are adequate representations of macroevolution, the incorporation of budding speciation will not
change our estimates. This is because differentiating lineages in the phylogenetic tree will have no
influence on the underlying model—lineages become effectively interchangeable. However, this is
not the trend that we are currently observing in PCMs. Extensions allowing heterogeneity in the
process, often associated with some predictor, have been shown to better capture the variation of
empirical data (e.g., Eastman, 2011; Rabosky et al., 2014; Uyeda and Harmon, 2014; Caetano et al.,
2018; Pagel et al., 2022). More recently, studies have demonstrated that rate heterogeneity should be
taken into account even when no a priori predictors are present (e.g., Rabosky and Goldberg, 2015;
Beaulieu and O'Meara, 2016; Caetano et al., 2018; May and Moore, 2020). Development of more
adequate models often means the increase in model complexity to reflect the dynamic nature of
macroevolution and, as a result, hint that the condition of homogeneous and memoryless
evolutionary changes with interchangeable lineages—under which budding would not matter—is
unlikely across the tree of life. Below we discuss how budding could be generating heterogeneity in
the phylogenetic history of phenotypes and in which ways the results affect our conclusions about
trait evolution.

When budding matters
Budding is expected to be important in any evolutionary scenario in which the identity of
evolutionary lineages is relevant. This might be the case if lineage age influences the tempo and/or
mode of trait evolution (Hagen et al., 2018; Goldberg and Foo, 2020) or if the age of competing
lineages is important to predict their competitive strength and/or risk of extinction (Ezard et al.,
2011; Rosenblum et al., 2012; Carrillo et al., 2020; Januario and Quental, 2021). Although there are
other evolutionary processes under which the identity of lineages might be important, here we focus
on these two scenarios for simplification. In contrast, there are special cases that generate
heterogeneity in trait evolution but under which budding likely is not relevant. For example, if shifts
in rates of trait evolution are due to abiotic causes equally affecting all lineages concurrent with the

event, such as response to climatic changes or mass extinctions, then, everything else being equal,

we would expect responses to be independent of lineage identity (e.g., Clavel and Morlon, 2017).

Below we enumerate scenarios in which we argue that budding speciation could influence our

conclusions about the tempo and mode of trait evolution when using PCMs.

1) When evolutionary changes are concentrated at or near lineage origination

The central distinction between budding and bifurcation is the age contrast between

progenitor and daughter lineage immediately after divergence. The daughter species will usually

have a smaller population size and geographic distribution (Foote et al., 2007; Liow and Stenseth,

2007) and might undergo quick phenotypic change as they move towards a new adaptive peak

(Eldredge and Gould, 1972; Gould and Eldredge, 1993; Hunt et al., 2008; De Lisle et al., 2021). In

contrast, progenitor lineages might show a slowdown in trait evolution due to prolonged time under

a stable adaptive zone (Goldberg and Foo, 2020; De Lisle et al., 2021). If lineage age is related to

the tempo of trait evolution, such that younger lineages are expected to show faster rates of trait

change, we would expect relatively more evolution to happen in a daughter lineage when compared

to its progenitor. Thus, the disparity between two descendants of a budding node in a phylogenetic

tree should not be attributed to equal amounts of change at each branch because budding suggests

evolution will be concentrated in the daughter lineage (Fig. 1 top left panel).

Figure 1: Approach used to illustrate and simulate budding speciation and trait evolution under lineage-age dependent rates. Top-left: Interaction between age-dependent trait evolution and speciation with and without budding. Stronger shades of color indicate faster rates of trait change. Bottom-left: Rate slowdown as a function of lineage age under the three treatments used. Right: Example of budding history with 20 extant species showing progenitor lineages in light and dark blue.

2) *When daughter lineage survival depends on being ecologically different from its progenitor*

The asymmetry in age generated by budding speciation can influence the competitive strength of daughter lineages relative to their progenitors and, as a result, also the extinction risk of the younger lineage (Ezard et al., 2011; Rosenblum et al., 2012; Carrillo et al., 2020; Januario and Quental, 2021). Progenitor lineages are expected to have larger population sizes and geographic ranges (Anacker and Strauss, 2014) which, everything else being equal, improves their chance of survival in interspecific competition with newly formed species. When competition between

progenitor and daughter lineages is present, daughter lineages that have lived enough to be sampled,

either in the fossil record or still living today, are expected to be sufficiently distinct from their

progenitors to have escaped competitive exclusion (De Lisle et al., 2021). Of course, competition is

not exclusive to budding. However, budding could potentially intensify the effect of interspecific

competition, and eventually increase heterogeneity in trait evolution.

Although we predict an intensifying effect of budding speciation on interspecific

competition, other natures of interactions might be more complex. Nuismer and Harmon (2015)

demonstrated mathematically the effect of the mode of trait evolution and phylogenetic diversity

(PD) in the outcome of interspecific interactions in communities of closely related taxa. They

showed that PD is a good predictor of interspecific interactions if these are dependent on phenotypic

matching, such as competition, with more closely related lineages showing stronger interspecific

interactions. Budding could change the relationship between PD and expected trait similarity,

because long-lived progenitor species would accumulate fewer evolutionary changes than expected

under a homogeneous trait evolution model, such as Brownian motion, causing the role of the

phylogeny as a predictor to become less prevalent. In contrast, Nuismer and Harmon (2015) show

that under stabilizing coevolution the phylogeny is a poor predictor of interactions, and we do not

expect that budding would influence this result.

How do budding speciation and lineage-age-dependent processes influence estimates of trait

evolution?

We use simple simulations to illustrate different scenarios in which budding speciation

should impact trait evolution and, more importantly, discuss if these deviations hinder our

understanding of phenotypic evolution using phylogenetic trees. We explored the impact of budding

on the parameter estimates and adequacy of PCMs for continuous and discrete traits. We also

investigated how likely is budding speciation to produce erroneous estimates of ancestral states. We

focused our attention on phylogenies of extant species, which are most often estimated using

molecular data, and in the absence of fossil tips.

Simulation of trait evolution under budding

We simulated 50 phylogenetic trees using a constant rate birth-death model ($\lambda = 0.2$, $\mu = 0.1$)

with root age set to 40 My and excluding all extinct lineages. To reduce variation in tree size we

used rejection sampling to keep only phylogenies with 250 to 350 extant lineages. We used the same

pool of 50 trees to perform all simulation replicates and conducted pairwise comparisons across

scenarios. We simulated a single continuous trait using a Brownian motion model ($\sigma^2 = 0.2$) and a

discrete trait with three states using an equal rates Markov model (transition rates = 0.02). To

emulate a scenario of fast evolution in younger lineages we introduced a rate slowdown process.

Rates of change, for both continuous and discrete traits, varied along the branches of the tree

following a scaling factor (s) computed as a function of lineage-age (a), such that

$$s_{(a_i)} = \frac{1}{\exp(z\,a_i)} \qquad (1)$$

where a_i is the average lineage-age at time interval i and z is the parameter controlling the rate

slowdown. In order to compute a, we divided the branches of the tree into i time intervals of length

1×10^{-3} of the tree height. At lineage-age of 0 My, for instance, s is equal to 1 and it decays as a

function of z (Fig. 1). We simulated three scenarios of lineage-age dependent rates of trait evolution:

a mild effect ($z = 0.042$); a medium effect ($z = 0.279$); and a strong effect ($z = 0.925$). The parameter

values were chosen to produce a rate reduction of 10%, 50%, and 90% of the base rate when a

lineage becomes 2.5 My old, respectively. Because the base rate is scaled by s, which depends on

the lineage age and the budding history of each phylogeny, the average rate of trait evolution can

vary. We standardized the average rate across the tree (for both discrete and continuous traits) to

differentiate the rate slowdown within lineages from the confounding effect of an overall change in

the average rate. For that, we computed the weighted average as

$$\bar{r} = \sum_i r\,s_{(a_i)}\,t_i / \sum_i t_i \qquad (2)$$

where r is the base rate of trait evolution (i.e., the σ^2 for the BM model and the transition rate for the

equal rates Markov model), and $s_{(a_i)}$ is the slowdown scale factor at a time interval t_i (see Equation

(1)). Then we chose r values that minimized the distance of $\bar{r}$ among replicates.

Figure 2: Conflict between ancestral state estimation and the true history of lineage evolution under budding.
New species appeared through budding and converged three times into a different niche, associated with the
coastal distribution (black). The progenitor lineage is the sister to the outgroup and carries the inland
distribution (yellow). Neither likelihood nor parsimony recovered the true history of the lineages. Left:
Marginal ancestral state estimation for the best fit Markov model. Branches are painted following a posterior
distribution of 100 stochastic maps. Right: Most parsimonious ancestral estimation following Sankoff's
(1975) algorithm. Center: True history of the trait. Phylogeny in the bottom left shows the continuation of the
progenitor lineage through multiple budding events.

We simulated budding speciation using an independent binary variable to control the

presence of budding on each node of the tree. As a result, long-lived progenitor lineages are

produced by random events of successive budding events. We produced four scenarios of budding

speciation, with frequencies of 0%, 25%, 50%, and 100% of the nodes. We also explored the effect

of cladogenetic changes on discrete traits (associated or not with budding). Cladogenetic changes

were simulated as a change with equal probability to any state immediately after speciation. When

budding is present, cladogenetic changes were restricted to daughter lineages whereas it could

happen to any lineage in the absence of budding. We also explored a scenario in which cladogenetic

changes restricted to daughter lineages (thus dependent on budding) produce convergence among all

daughters of the same long-lived progenitor lineage (see examples in Figs. 2 and 3). A detailed

report of the simulation is available in the Supplementary Materials.

Progenitor carrying the state
Progenitor not carrying the state

Figure 3: Evolution of a truly irreversible trait. Long-lived progenitor lineages carry the ancestral state
whereas daughter lineages have lost the trait independently several times. Fossil lineages show evidence of
trait homology. PCMs in the absence of the fossil record would wrongly estimate two independent origins of
the "black" trait.

Evaluating model adequacy and errors in ancestral estimation

We used the method described by Pennell and colleagues (2015) to evaluate the adequacy of

PCMs for continuous traits in the presence of budding speciation and lineage-age-dependent rates of

trait evolution. This method computes a pool of summary statistics (see Table 1 in Pennell *et al.*,

2015) and compares each with an expected distribution estimated from the data. If the model is

adequate, the observed summary statistics should fall around the mean of the null distribution

whereas values outside the 95% highest density interval indicate that the PCM is inadequate. To

evaluate the effect of budding in the ancestral estimation of discrete states we used an index of how

incorrect the estimate at a node is with respect to the true history of the trait. We measured the

highest marginal probability among all states excluding the true state for the nodes as an estimate of

"wrongness". This metric reflects how likely the state of a node would be estimated as the wrong

ancestral state. Note that this is distinct from uncertainty because wrongness is maximized when we

have certainty of the ancestral state but it is incongruent with the true (simulated) history.

Uncertainty is a lesser problem than wrongness because we will not, or at least should not, support

or refute evolutionary hypotheses based on uncertain estimates. Wrongness, on the other hand, can

result in misleading interpretations. We then used linear mixed models to test for the association

between node age and wrongness across all simulation scenarios and selected the best model using

the Akaike Information Criterion (AIC).

Effects of budding and lineage-age dependent processes on the adequacy of continuous trait
evolution models

After simulating continuous traits under 12 scenarios, varying the strength of the lineage-age
dependent process and the frequency of budding speciation, we estimated parameters for single rate
Brownian motion (Felsenstein, 1973), variable rate BM (Eastman et al., 2011), single optimum
Ornstein-Uhlenbeck (Butler and King, 2004), and Early-Burst (Harmon et al., 2010) models as
implemented in *geiger* (Pennell et al., 2014). Note that none of those models is the true model that
generated the data. Our goal is to evaluate which is the preferred model among the suite of PCMs
most used in the literature and to better understand the potential effects of budding speciation on our
inferences. We also hope that this simple illustration through the use of simulations motivates
further research in model development. Overall, variable rate BM models showed significant
improvement in model adequacy under budding.

Model adequacy tests for a homogeneous rate BM model (Pennell et al., 2015) detected a
negative slope of the linear fit between node depth and the size of the phylogenetic independent
contrasts (S_{hgt}) indicating that larger trait changes are estimated to have happened closer to the tips
(Fig. S3). However, a variable rate BM model does not show evidence for such deviation (Fig. S3),
suggesting this is an effect of underestimating the rate variation introduced by budding (which
introduces heterogeneity in a different way than the variable rate BM models). A regression of
phylogenetic independent contrasts (PICs) and their expected variance (S_{var}) shows that nodes
connected by short branches are associated with more trait change (Fig. S2), independently if the
fitted BM model was homogeneous or not. Inadequacy of S_{var} is expected due to the concentration of
rates early in the history of lineages and the stronger effect of slowdown on the more longevous
lineages when compared with short-lived ones. A pattern that is expected under budding speciation
and punctuated equilibrium. The deviances for S_{hgt} and S_{var} are only detectable when the rate
slowdown is strong, meaning that a relaxed rates model (Eastman et al., 2011) seems to be able to
adequately describe trait variation if lineage-age effects are mild, but not if they are strong.
Inadequacies in both S_{hgt} and S_{var} point to trait changes concentrated close to the tips, which is
expected since molecular trees have an accumulation of nodes near the present, some of these

generated by budding, producing new lineages with higher rates of trait evolution. Model

inadequacy could be an artifact of unobserved speciation events deeper in the tree but results remain

constant when we replicate analyses including extinct lineages. Deviations of S_{hgt}, S_{var}, and C_{var}

(coefficient of variation of PICs, a measure of rate heterogeneity) get stronger as the intensity of the

lineage-age-dependent slowdown factor increases. On the other hand, changing the frequency of

budding speciation, while controlling for the strength of the slowdown factor, did not change the

patterns of model adequacy across all summary statistics we investigated (Figs. S2-4). We did not

verify any deviation from the remaining summary statistics adopted by Pennell and colleagues

(2015). In summary, if budding speciation is frequent and there is strong age-dependent trait

evolution (punctuated equilibrium representing an extreme version of this), current inference

methods would have trouble adequately capturing patterns of trait evolution.

With respect to the support for alternative trait evolution models (i.e., Brownian motion,

Ornstein-Uhlenbeck, and Early-Burst) as a function of lineage-age dependent rate variation, the OU

model shows a marked increase in AIC weights in response to stronger slowdown factors (Figs. S5

and S6). In the majority of cases, the phylogenetic half-life was estimated to be multiple times

longer than the age of the clade (40 My) indicating processes indistinguishable from Brownian

motion (Cooper et al. 2016). Average phylogenetic half-life estimated across replicates was only

shorter than clade age when the strongest slowdown factor was applied (Fig. S7). This means that an

OU process is only supported when progenitor lineages are practically in stasis whereas virtually all

trait evolution is concentrated on the origination of new lineages following budding speciation. In

other words, if budding speciation produces a pattern congruent with Punctuated Equilibrium we

expect support for OU models. These results remain constant regardless of the frequency of budding

speciation used to simulate the data (Fig. S4) or the inclusion of extinct lineages.

When we simulate continuous traits under a lineage-age dependent process, the BM model

with varying rates adequately describes most characteristics of the data but fails to capture the

concentration of trait changes on shorter branches (S_{var}). If lineage-age dependent processes happen

in nature, our results reinforce the cautionary note that parameter estimates can be more informative

than model choice alone (Cooper et al. 2016). Although the majority of scenarios supported OU

models, only the strongest case of rate slowdown resulted in patterns distinguishable from Brownian

motion. Fortunately, the deviance of the slope of absolute contrasts as a function of their expected

variance (S_{var} - Pennel et al. 2015) can help to detect a concentration of trait change towards shorter

branches, even when controlled for rate variation, which is one of the expectations of lineage-age-

dependent rates of evolution. Those simulations are far from being comprehensive (our goal was not

to be exhaustive but to provide examples to support our narrative), but they emphasize the potential

effects of not explicitly considering budding speciation, in particular when age-dependent trait

evolution is present. Future simulation studies should more deeply focus on the different aspects

superficially touched here as well as on others not discussed.

Effects of budding speciation and lineage-age dependent processes on ancestral estimation

Here we investigate how budding affects our estimations of ancestral state for discrete traits.

As expected, all fitted models show a strong positive association between node age and wrongness

(Fig. 4), meaning that ancestral estimation of nodes closer to the root of the tree is more likely to be

misleading. The best-ranked linear mixed model using AIC (Table S1) suggests that budding

speciation has a significant effect on wrongness when cladogenetic trait changes are also present

(see example in Fig. 3). Without cladogenetic changes, there is no detectable difference between the

null model (homogeneous rates and bifurcating speciation) and the model with budding speciation

(Fig. 4). Budding associated with cladogenetic changes increases the chance of misleading ancestral

state estimation, especially for younger nodes. This result is somewhat unexpected and important

because younger nodes are often expected to have more information than older ones (Schultz et al.,

1996; Boyko and Beaulieu, 2021).

Figure 4: Relationship between the presence of nodes with wrongly reconstructed states and node age. Nodes were considered wrongly reconstructed if the marginal ancestral state probability for any state distinct from the correct state were > 0.5 at that node. Logistic regressions were performed with the pooled results from the 50 simulation replicates per group and independent for each type of node and study group (see **Table S1**). Dashed lines represent nodes with budding speciation whereas solid lines show nodes without budding. Dots on the top and bottom represent pooled nodes across simulation replicates correctly and incorrectly estimated, respectively.

Can budding help us understand complex histories of trait evolution?

Here we used simulations to exemplify the effect of budding speciation in PCMs of trait evolution. Our initial results show that budding has an impact but does not completely hinder the utility of the most popular models of trait evolution. Some of the effects we report here, for the most part, can be translated as heterogeneity in trait evolution among lineages. Different from other sources of heterogeneity (e.g., Uyeda and Harmon, 2014; Boyko and Beaulieu, 2021), budding produces variation dependent on lineage identity, requiring the identification of progenitor lineages potentially comprising multiple contiguous branches of a phylogeny. From this perspective, we are optimistic about incorporating budding speciation into PCMs, and we hope our initial discussion on the subject motivates further research on how we can properly incorporate budding speciation and

age-dependent trait evolution in PCMs. In fact, it is plausible that a portion of the intrinsic

heterogeneity captured by rate-varying models, such as hidden rates models for discrete traits

(Beaulieu and O'Meara, 2016; Caetano et al., 2018; Boyko and Beaulieu, 2021) and those applying

reversible jump MCMC for continuous traits (Eastman et al., 2011; Rabosky et al., 2014; Uyeda and

Harmon, 2014), is due to the effect of budding speciation.

Distinct from scenarios in which some predictor trait is responsible for rate shifts, budding is

expected to affect trait evolution dependent on the mode of speciation. This introduces a

complication because we need to reconstruct the budding history of lineages, which does not easily

leave a trace on molecular phylogenies (e.g., it needs samples of multiple populations of recently

diverged lineages). One potential solution is to use data augmentation (e.g., Quintero and Landis,

2020) to co-estimate budding history and trait evolution model parameters using simulations. This

approach could be challenging because both the frequency of budding speciation and the location of

the progenitor lineages would need to be sampled. However, our inability to pinpoint the location of

progenitor lineages should not be used as an argument for ignoring its effect on trait evolution.

Today we have several PCMs that are able to recover the signal of shifts in the tempo and mode of

trait evolution without a priori hypotheses (Eastman, 2011; Rabosky et al., 2014; Uyeda and

Harmon, 2014; Pagel et al., 2022) and, more importantly, there is evidence that such methods

improve model adequacy (Rabosky and Goldberg, 2015; Beaulieu and O'Meara, 2016; Caetano et

al., 2018). We suggest that budding speciation should be considered as a confounding factor akin to

rate heterogeneity, which needs to be taken into account when estimating the history of trait

evolution using molecular phylogenies—even if budding is not the focus of the study.

Another challenge is that progenitor lineages can produce scenarios incongruent with the

most parsimonious history for a trait (e.g. Figs. 2 and 3). For example, ancestral estimates of the

scenario shown in Figure 3 in the absence of fossil information would suggest, with confidence, that

the trait is a convergence. This is a scenario in which PCM estimates can conflict with external

evidence of homology. For instance, Pyron (2015) discusses the inference of multiple transitions

from viviparity back to oviparity in snakes, based on PCMs, despite the external evidence based on

development and physiology against so (Griffith et al., 2015). Pyron (2015) suggests that

comparative approaches should not ignore external evidence but also that findings from

phylogenetic inferences should be further investigated integratively. However, an unlikely ancestral

reconstruction of parity might simply mean that the model is inadequate. For instance, budding
speciation could help explain oviparous lineages nested deep into viviparous clades as descendants
of long-lived progenitor lineages (see discussion in Pyron 2015). The budding speciation scenario
would require many additional evolutionary transitions, but it would support the extensive
knowledge about genetics, development, and physiology of snakes (see discussion in Griffith et al.,
2015). In our view, when there is a clash between model estimates based on projections into millions
of years in the past and biological knowledge, it is wise to review our models and ponder which
important processes the model might be failing to capture, including the possibility of budding
speciation.

Closing remarks
Budding speciation might indirectly or directly impact both estimates of lineage
diversification and trait evolution under PCMs widely used in the literature. Our results suggest that
ignoring budding speciation when age-dependent trait evolution operates might lead to incorrect
inferences such as inferring the wrong ancestral state for younger nodes. We also suggest that it
might be possible, although challenging, to incorporate budding speciation into PCMs for both
discrete and continuous traits. The introduction of budding speciation in comparative approaches,
however, depends on the departure from the parsimony paradigm which we suspect is a barrier to
the development of macroevolutionary models that can fully integrate external biological
information about trait evolution. When we intuitively imagine a parsimonious trait history, we are
doing so independently of what is known about the evolutionary history of the system. Reflecting on
the role of comparative approaches and recognizing their limitations, especially when testing
scenarios of complex trait evolution, is key to the development of alternative models that help the
study of macroevolution to become a more integrative endeavor. The incorporation of budding
speciation is one example of the direction we can take in improving comparative studies, and we
hope our discussion motivates researchers to explore further some of these possibilities.

References

Anacker, B. L., and S. Y. Strauss. 2014. The geography and ecology of plant speciation: range overlap and niche divergence in sister species. *Proceedings of the Royal Society B: Biological Sciences* 281:20132980.

Aze, T., T. H. G. Ezard, A. Purvis, H. K. Coxall, D. R. M. Stewart, B. S. Wade, and P. N. Pearson. 2011. A phylogeny of Cenozoic macroperforate planktonic foraminifera from fossil data. *Biological Reviews* 86:900–927.

Baker, J., A. Meade, M. Pagel, and C. Venditti. 2015. Adaptive evolution toward larger size in mammals. *Proceedings of the National Academy of Sciences* 112(16):5093–5098.

Baldwin, B. G. 2005. Origin of the serpentine-endemic herb *Layia discoidea* from the widespread *L. glandulosa* (Compositae). *Evolution* 59:2473.

Bapst, D. W. 2013. When can clades be potentially resolved with morphology? *PLOS ONE* 8:e62312.

Bapst, D. W., and M. J. Hopkins. 2017. Comparing cal3 and other a posteriori time-scaling approaches in a case study with the pterocephaliid trilobites. *Paleobiology* 43:49–67.

Bartoszek, K. 2014. Quantifying the effects of anagenetic and cladogenetic evolution. *Mathematical Biosciences* 254:42-57.

Bartoszek, K. 2020. A central limit theorem for punctuated equilibrium. *Stochastic Models* 36:473-517.

Bastide, P., C. Solís-Lemus, R. Kriebel, K. W. Sparks, and C. Ané. 2018. Phylogenetic comparative methods on phylogenetic networks with reticulations. *Systematic Biology* 67:800–820.

Baum, D. A., S. D. Smith, and S. S. S. Donovan. 2005. The Tree-Thinking Challenge. *Science* 310:979–980.

Beaulieu, J. M., and B. C. O'Meara. 2016. Detecting Hidden Diversification Shifts in Models of Trait-Dependent Speciation and Extinction. *Systematic Biology* 65:583–601.

Behm, J. E., Ives, A. R., and J. W. Boughman. 2010. Breakdown in postmating isolation and the collapse of a species pair through hybridization. *The American Naturalist* 175:11–26.

Benton, M. J., and P. N. Pearson. 2001. Speciation in the fossil record. *Trends in Ecology & Evolution* 16:405–411.

Bokma, F. 2002. Detection of punctuated equilibrium from molecular phylogenies. *Journal of Evolutionary Biology* 15:1048–1056.

Bokma, F. 2008. Detection of "Punctuated Equilibrium" by Bayesian estimation of speciation and extinction rates, ancestral character states, and rates of anagenetic and cladogenetic evolution on a molecular phylogeny. *Evolution* 62:2718–2726.

Bokma, F. 2010. Time, species, and separating their effects on trait variance in clades. *Systematic Biology* 59:602–607.

Boyko, J. D., and J. M. Beaulieu. 2021. Generalized hidden Markov models for phylogenetic comparative datasets. *Methods in Ecology and Evolution* 12:468–478.

Butler, M. A., and A. A. King. 2004. Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *American Naturalist* 164:683–695.

Caetano, D. S., B. C. O'Meara, and J. M. Beaulieu. 2018. Hidden state models improve state-dependent diversification approaches, including biogeographical models. *Evolution* 72:2308–2324.

Carrillo, J. D., S. Faurby, D. Silvestro, A. Zizka, C. Jaramillo, C. D. Bacon, and A. Antonelli. 2020. Disproportionate extinction of South American mammals drove the asymmetry of the Great American Biotic Interchange. *Proceedings of the National Academy of Sciences* 117:26281–26287.

Clavel, J., and H. Morlon. 2017. Accelerated body size evolution during cold climatic periods in the Cenozoic. *Proceedings of the National Academy of Sciences* 114:4183–4188.

Cooper, N., G. H. Thomas, C. Venditti, A. Meade, and R. P. Freckleton. 2016. A cautionary note on the use of Ornstein Uhlenbeck models in macroevolutionary studies. *Biological Journal of the Linnean Society* 118:64–77.

Crawford, D. J. 2010. Progenitor-derivative species pairs and plant speciation. *Taxon* 59:1413–1423.

Crouch, N. M. A., S. M. Edie, K. S. Collins, R. Bieler, and D. Jablonski. 2021. Calibrating phylogenies assuming bifurcation or budding alters inferred macroevolutionary dynamics in a densely sampled phylogeny of bivalve families. *Proceedings of the Royal Society B: Biological Sciences* 288:20212178.

De Lisle, S. P., D. Punzalan, N. Rollinson, and L. Rowe. 2021. Extinction and the temporal distribution of macroevolutionary bursts. *Journal of Evolutionary Biology* 34:380–390.

Duchen, P., M. L. Alfaro, J. Rolland, N. Salamin, and D. Silvestro. 2021. On the effect of asymmetrical trait inheritance on models of trait evolution. *Systematic Biology* 70:376–388.

Eastman, J. M., M. E. Alfaro, P. Joyce, A. L. Hipp, and L. J. Harmon. 2011. A novel comparative method for identifying shifts in the rate of character evolution on trees. *Evolution* 65:3578–3589.

Eldredge, N., and S. J. Gould. 1972. Punctuated equilibria: An alternative to phyletic gradualism, pp. 82-115. In: Schopf, T. J. M., ed. Models in Paleobiology. Freeman, Cooper and Co.; San Francisco, California.

Elliot, M. G., and A. Ø. Mooers. 2014. Inferring ancestral states without assuming neutrality or gradualism using a stable model of continuous character evolution. *BMC Evolutionary Biology* 14.

Ezard, T. H. G., T. Aze, P. N. Pearson, and A. Purvis. 2011. Interplay between changing climate and species' ecology drives macroevolutionary dynamics. *Science* 332:349–351.

Felsenstein, J. 1973. Maximum likelihood estimation of evolutionary trees from continuous characters. *American Journal of Human Genetics* 25:471–492.

Felsenstein, J. 1981. Evolutionary trees from DNA sequences: A maximum likelihood approach. *Journal of Molecular Evolution* 17:368–376.

Felsenstein, J. 1985. Phylogenies and the Comparative Method. *American Naturalist* 125:1–15.

Foote, M. 1996. On the Probability of ancestors in the fossil record. *Paleobiology* 22:141–151.

Foote, M., J. S. Crampton, A. G. Beu, B. A. Marshall, R. A. Cooper, P. A. Maxwell, and I. Matcham. 2007. Rise and Fall of Species Occupancy in Cenozoic Fossil Mollusks. *Science* 318:1131–1134.

Funk, D. J., and K. E. Omland. 2003. Species-Level Paraphyly and Polyphyly: Frequency, Causes, and Consequences, with Insights from Animal Mitochondrial DNA. *Annual Review of Ecology, Evolution, and Systematics* 34:397–423.

Goldberg, E. E., and J. Foo. 2020. Memory in Trait Macroevolution. *The American Naturalist* 195:300–314.

Gottlieb, L. D. 2004. Rethinking classic examples of recent speciation in plants. *New Phytologist* 161:71–82.

Gould, S. J., and N. Eldredge. 1993. Punctuated equilibrium comes of age. *Nature* 366:223–227.

Grant, V. 1981. Plant speciation, 2nd ed. New York, NY, USA: Columbia University Press.

Griffith, O. W., D. G. Blackburn, M. C. Brandley, J. U. Van Dyke, C. M. Whittington, and M. B.
Thompson. 2015. Ancestral state reconstructions require biological evidence to test
evolutionary hypotheses: A case study examining the evolution of reproductive mode in
squamate reptiles. *Journal of Experimental Zoology Part B: Molecular and Developmental*
Evolution 324:493–503.

Hagen, O., K. Hartmann, M. Steel, and T. Stadler. 2015. Age-dependent speciation can explain the
shape of empirical phylogenies. *Systematic Biology* 64:432–440.

Harmon, L. J., J. B. Losos, T. J. Davies, R. G. Gillespie, J. L. Gittleman, W. B. Jennings, K. H.
Kozak, M. A. McPeek, F. Moreno-Roark, T. J. Near, A. Purvis, R. E. Ricklefs, D. Schluter,
J. A. Schulte II, O. Seehausen, B. L. Sidlauskas, O. Torres-Carvajal, T. J. Weir, and A. Ø.
Mooers. 2010. Early bursts of body size and shape evolution are rare in comparative data.
Evolution 64: 2385–2396.

Helmstetter, A.J., Zenil-Ferguson, R., Sauquet, H., Otto, S.P., Méndez, M., Vallejo-Marin, M. et al.
(2023) Trait-dependent diversification in angiosperms: Patterns, models and data. *Ecology*
Letters 26:640– 657.

Huelsenbeck, J. P., R. Nielsen, and J. P. Bollback. 2003. Stochastic mapping of morphological
characters. *Systematic Biology* 52:131–158.

Hunt, G., M. A. Bell, and M. P. Travis. 2008. Evolution toward a new adaptive optimum:
phenotypic evolution in a fossil stickleback lineage. *Evolution* 62:700–710.

Janzen, T., F. Bokma, and R. S. Etienne. 2022. Nucleotide substitutions during speciation may
explain substitution rate variation. *Systematic biology* 71:1244-1254

Januario, M., and T. B. Quental. 2021. Re-evaluation of the "law of constant extinction" for
ruminants at different taxonomical scales. *Evolution* 75:656–671.

Lackey, A. C, J. W. Boughman. 2017. Evolution of reproductive isolation in stickleback fish.
Evolution 71:357-372

Liow L.H. & Stenseth N.C. 2007 The rise and fall of species: implications for macroevolutionary
and macroecological studies. *Proceedings of the Royal Society of London, Series B*
274:2745–2752.

Mallet, J., N. Besansky, and M. W. Hahn. 2016. How reticulated are species? *BioEssays* 38:140–
149.

Mattila, T. M., and F. Bokma. 2008. Extant mammal body masses suggest punctuated equilibrium.
Proceedings of the Royal Society B: Biological Sciences 275:2195–2199.

May, M. R., and B. R. Moore. 2020. A bayesian approach for inferring the impact of a discrete
character on rates of continuous-character evolution in the presence of background-rate
variation. *Systematic Biology* 69:530–544.

Mayr, E. 1942. Systematics and the origin of species. New York, NY, USA: Columbia University
Press.

Meier, R., and R. Willmann. 2000. The Hennigian species concept. In Wheeler, Q. D., and R. Meier
(eds). Species concepts and phylogenetic theory: a debate (pp. 30-43). Columbia University
Press, New York.

Monroe, M. J., and F. Bokma. 2009. Do speciation rates drive rates of body size evolution in
mammals? *The American Naturalist* 174:912–918.

Nuismer, S. L., and L. J. Harmon. 2015. Predicting rates of interspecific interaction from
phylogenetic trees. *Ecology Letters* 18:17–27.
O'Meara, B. C. 2012. Evolutionary Inferences from Phylogenies: A Review of Methods. *Annual*
Review of Ecology, Evolution, and Systematics 43:267–285.
Otero, A., P. Vargas, M. Fernández-Mazuecos, P. Jiménez-Mejías, V. Valcárcel, I. Villa-Machío,
and A. L. Hipp. 2022. A snapshot of progenitor–derivative speciation in *Iberodes*
(Boraginaceae). *Molecular Ecology* 31:3192–3209.
Pagel, M., C. O'Donovan, and A. Meade. 2022. General statistical model shows that
macroevolutionary patterns and processes are consistent with Darwinian gradualism. *Nature*
Communications 13:1113.
Papuga, G., P. Gauthier, V. Pons, E. Farris, and J. D. Thompson. 2018. Ecological niche
differentiation in peripheral populations: a comparative analysis of eleven Mediterranean
plant species. *Ecography* 41:1650–1664.
Parins-Fukuchi, C. 2021. Morphological and phylogeographic evidence for budding speciation: an
example in hominins. *Biology Letters* 17:20200754.
Patsis, A., R. P. Overson, K. A. Skogen, N. J. Wickett, M. G. Johnson, W. L. Wagner, R. A. Raguso,
J. B. Fant, and R. A. Levin. 2021. Elucidating the evolutionary history of oenothera sect.
Pachylophus (onagraceae): A phylogenomic approach. *Systematic Botany* 46:799–811.
Pennell, M. W., and L. J. Harmon. 2013. An integrative view of phylogenetic comparative methods:
Connections to population genetics, community ecology, and paleobiology. *Annals of the*
New York Academy of Sciences 1289:90–105.
Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E. Alfaro,
and L. J. Harmon. 2014. geiger v2.0: An expanded suite of methods for fitting
macroevolutionary models to phylogenetic trees. *Bioinformatics* 30:2216–2218.
Pennell, M. W., R. G. FitzJohn, W. K. Cornwell, and L. J. Harmon. 2015. Model adequacy and the
macroevolution of angiosperm functional traits. *American Naturalist* 186:E33–E50.
Philippe, H., and C. J. Douady. 2003. Horizontal gene transfer and phylogenetics. *Current Opinion*
in Microbiology 6:498-505.
Pyron, R. A. 2015. Advancing perspectives on parity-mode evolution. *Journal of Experimental*
Zoology Part B: Molecular and Developmental Evolution 324:562–563.
Quintero, I., and M. J. Landis. 2020. Interdependent phenotypic and biogeographic evolution driven
by biotic interactionst. *Systematic Biology* 69:739–755.
Rabosky, D. L., S. C. Donnellan, M. Grundler, and I. J. Lovette. 2014. Analysis and visualization of
complex macroevolutionary dynamics: An example from australian scincid lizards.
Systematic Biology 63:610–627.
Rabosky, D. L., and E. E. Goldberg. 2015. Model inadequacy and mistaken inferences of trait-
dependent speciation. *Systematic Biology* 64(2):340–355.
Raup, D. M. 1985. Mathematical models of cladogenesis. *Paleobiology* 11:42–52.
Raup, D. M., and S. J. Gould. 1974. Stochastic simulation and evolution of morphology-towards a
nomothetic paleontology. *Systematic Zoology* 23(3):305–322.
Richmond, J. Q. and E. L. Jockusch. 2007. Body size evolution simultaneously creates and collapses
species boundaries in a clade of scincid lizards. *Proceedings of Biological Sciences*
274:1701–1708.
Rosenblum, E. B., B. A. J. Sarver, J. W. Brown, S. Des Roches, K. M. Hardwick, T. D. Hether, J.
M. Eastman, M. W. Pennell, and L. J. Harmon. 2012. Goldilocks meets Santa Rosalia: An

653   ephemeral speciation model explains patterns of diversification across time scales.
654   *Evolutionary Biology* 39:255–261.
655 Sankoff, D. 1975. Minimal mutation trees of sequences. *SIAM Journal on Applied Mathematics*
656   28:35–42.
657 Schultz T.R., Cocroft R.B., Churchill G.A. 1996. The reconstruction of ancestral character states.
658   *Evolution* 50:504–511.
659 Taylor, E. B., J. W. Boughman, M. Groenenboom, M. Sniatynski, D. Schluter, and J. L. Gow. 2006.
660   Speciation in reverse: Morphological and genetic evidence of the collapse of a threespined
661   stickleback (*Gasterosteus aculeatus*) species pair. *Molecular Ecology* 15:343–355.
662 Uyeda, J. C., and L. J. Harmon. 2014. A novel Bayesian method for inferring and interpreting the
663   dynamics of adaptive landscapes from phylogenetic comparative data. *Systematic Biology*
664   63:902–918.
665 Wagner, P. J. 1998. A likelihood approach for evaluating estimates of phylogenetic relationships
666   among fossil taxa. *Paleobiology* 24:430–449.