1 **Par3/Bazooka binds NICD and promotes Notch signalling during *Drosophila***

2 **development**

5 Jun Wu, Neeta Bala Tannan, Linh T. Vuong, Yildiz Koca, Giovanna M. Collu,

6 and Marek Mlodzik

Affiliations

Dept. of Cell, Developmental, and Regenerative Biology,

Graduate School of Biomedical Sciences,

Icahn School of Medicine at Mount Sinai,

One Gustave L. Levy Place, New York, NY 10029.

Abstract

The conserved *bazooka (baz/par3*) gene acts as a key regulator of asymmetrical cell divisions

across the animal kingdom. Associated Par3/Baz-Par6-aPKC protein complexes are also well

known for their role in the establishment of apical/basal cell polarity in epithelial cells. Here we

define a novel, positive function of Baz/Par3 in the Notch pathway. Using *Drosophila* wing and

eye development, we demonstrate that Baz is required for Notch signaling activity and optimal

transcriptional activation of Notch target genes. Baz appears to act independently of aPKC in

these contexts, as knockdown of *aPKC* does not cause *Notch* loss-of-function phenotypes.

Using transgenic Notch constructs, our data positions Baz activity downstream of activating

Notch cleavage steps and upstream of Su(H)/CSL transcription factor complex activity on Notch

target genes. We demonstrate a biochemical interaction between NICD and Baz, suggesting

that Baz is required for NICD activity before NICD binds to Su(H). Taken together, our data

define a novel role of the polarity protein Baz/Par3, as a positive and direct regulator of Notch

signaling through its interaction with NICD.

Introduction

Cell polarity is a critical feature of most cell types, particularly epithelial cell sheets, which display obligate apical-basal (A/B) polarity that is required for their barrier function and vectorial secretion. Many proteins and mechanisms that govern the establishment and maintenance of cell polarity are evolutionarily conserved and were discovered genetically in *C. elegans* and *Drosophila* (reviewed in (Goldstein and Macara, 2007; St Johnston and Sanson, 2011; Tepass, 2012)). Similarly, stem cells often display polarized features using the same factors, which are critical for stem cell maintenance and for progenitor cell fates, with cell contents being asymmetrically inherited by the two daughter cells (e.g. rev. in (Neumuller and Knoblich, 2009; Schweisguth, 2015)). The protein complexes that mediate polarization are also shared among these distinct instances of cell polarization.

The conserved *par3* gene, known as *bazooka/baz* in *Drosophila*, was originally identified in *C. elegans*, where it acts as a key regulator of asymmetrical cell divisions in the early embryo (Etemad-Moghadam et al., 1995). It generally acts together with Par6 and aPKC as a regulatory complex of many asymmetric cell divisions across the animal kingdom (reviewed in (Neumuller and Knoblich, 2009; Schweisguth, 2015)). This Par3/Baz-Par6-aPKC complex is also well known for its conserved role in the establishment of A/B polarity in epithelial cells (Suzuki and Ohno, 2006). Our interest in the A/B epithelial cell polarity proteins originates from the fact that several of these have also been linked to planar cell polarity (PCP), that is polarity within the epithelial sheet, orthogonal to the A/B axis (Humphries and Mlodzik, 2018). For example aPKC can associate with Frizzled, and Scribble (another A/B polarity protein) associates with the PCP protein Van Gogh/Vang (Vangl in vertebrates) (Courbard et al., 2009; Djiane et al., 2005). aPKC and Scribble thus contribute to the stability and asymmetric segregation of the core PCP complexes and establishment of planar polarity within epithelial cell sheets. To systematically examine whether additional A/B epithelial polarity factors can play a role in PCP establishment,

we tested members of these protein families for PCP associated knockdown phenotypes during

Drosophila wing development. Surprisingly when we examined phenotypes of *baz/par3*

knockdown in this context, we observed a *Notch* mutant-like phenotype of loss of wing margin

tissue. Here we focus on investigating a specific role for Baz in regulating Notch signaling.

Notch (N) was originally identified in *Drosophila* based on its haplo-insufficient, dominant wing-

notching phenotype of loss of tissue at the dorso-ventral border of the wing (Guruharsha et al.,

2012; Mohr, 1919; Southgate et al., 2015). *Notch* encodes a single-pass transmembrane

receptor protein containing EGF-like repeats in its extracellular domain (Wharton et al., 1985)

and a conserved intracellular domain (NICD) that, upon cleavage, can translocate to the

nucleus and function as transcriptional co-activator (reviewed in (Andersson et al., 2011; Bray,

2016; Henrique and Schweisguth, 2019; Kopan and Ilagan, 2009; Wilson and Radtke, 2006)).

Many studies demonstrated that the highly conserved Notch receptor family plays important

roles during the development and homeostasis of most tissues through cell-cell communication

and cellular signaling, controlling many target genes that can be either common or context

dependent (reviewed in (Andersson et al., 2011; Bray, 2016; Henrique and Schweisguth, 2019;

Kopan and Ilagan, 2009; Wilson and Radtke, 2006) (Siebel and Lendahl, 2017)). Misregulation

of N-signaling is associated with many human diseases, including cancer (Siebel and Lendahl,

2017). The Notch receptor is activated by the binding of ligands of the Delta (Dl) or

Serrate/Jagged (Ser) families from neighboring cells to trigger juxtamebrane cleavage events to

release NICD (Andersson et al., 2011; Bray, 2016; Struhl and Greenwald, 1999). NICD can then

translocate into the nucleus, where it interacts with transcription factors of the Suppressor of

Hairless (Su(H)/CSL) family, and co-activators, including Mastermind, to activate target gene

expression (Bray, 2016). Notch signaling is subject to many regulatory steps that have been and

continue to be identified and characterized in *Drosophila*, ranging from receptor and ligand

trafficking, receptor glycosylation, membrane lipid composition, receptor cleavage, and protein

stability regulators (Bala Tannan et al., 2018; Kandachar and Roegiers, 2012; Kidd et al., 1998;

Medina-Yáñez et al., 2020; Rana and Haltiwanger, 2011; Shilo and Schejter, 2011; Singh and

Mlodzik, 2012; Struhl and Greenwald, 1999; Weber et al., 2003) to specific nuclear events at the

transcriptional end of pathway activation (rev. in (Andersson et al., 2011; Bray, 2016; Henrique

and Schweisguth, 2019; Kopan and Ilagan, 2009; Wilson and Radtke, 2006)).

Previous work proposed an indirect role for Baz/Par3 in regulating Notch signaling during

asymmetric cell divisions in *Drosophila*. During asymmetric cell division of the sensory organ

precursors (SOPs), specific protein complexes segregate to opposite sides of the cell cortex in

each dividing SOP cell, on one side the Baz/Par3-Par6-aPKC complex and on the other the

Gai-Pins-Mud-Dlg-Numb-Neur complex. The two daughter cells that result from this cell division

adopt distinct cell fates due to the asymmetric distribution of the above complexes: with one

daughter cell inheriting Baz/Par3-Par6-aPKC, and the other the Gai-Pins-Mud-Dlg-Numb-Neur

complex (Schweisguth, 2015). Notably, the asymmetric segregation of these complexes causes

an asymmetric activation of Notch signaling, as Numb is known to repress N-signaling by

reducing Notch levels at the cell surface, either through internalization or inhibition of recycling

(Schweisguth, 2015). In a similar scenario, in the context of intestinal stem cell (ISC) number

regulation, Baz is also required within the asymmetric apical-basal polarity complex segregation

to regulate Notch activity (Wu et al., 2023). As such, an indirect role of Baz/Par3 in the

regulation of N-signaling has been proposed, acting through the promotion of the asymmetric

segregation of the respective apical-basal complexes to allow Notch activity in the Baz-

containing daughter cells. However, due to this indirect role of Baz/Par3 in the pathway, such

model systems are not suited to address potential direct roles of Baz/Par3 in Notch-signaling.

Using *Drosophila* wing and eye development, we define a novel, direct and positive function of

Baz/Par3 in Notch signaling. We demonstrate that Baz is required for Notch signaling activity

and activation of Notch transcriptional target genes. Baz appears to act independently of aPKC

in these contexts, as knockdown of *aPKC* causes PCP defects and not Notch-associated

phenotypes. Our data indicate that Baz acts the level of NICD, the cleaved active Notch

fragment, but upstream of the Su(H)-NICD-Mam activation complex. We further demonstrate a

direct biochemical interaction between NICD and Baz. Taken together, our data define a novel

role of the polarity protein Baz/Par3, as a positive regulator of Notch signaling through its

interaction with NICD.

Results

Reduction of *baz/par3* causes defects similar to *Notch* loss-of-function phenotypes

To initially assess potential functions of Baz/Par3-Par6-aPKC factors in PCP we expressed two

independent non-overlapping RNAi (IR) lines for each gene in the posterior compartment of

developing wing discs using the *en-Gal4* driver (Figure 1 and Suppl. Figure S1). Surprisingly,

with individual components of the A/B polarity complex containing Baz/Par3, Par6, and aPKC

we detected distinct outcomes. While knock-down of aPKC showed classical PCP defects, as

expected based on previous work (Djiane et al., 2005) (see also below), both *baz-IR* lines

caused wing margin loss in the posterior compartment (Figure 1A-C; Baz protein levels were

markedly reduced and barely detectable in the *baz-IR* expressing domain: Figure 1E, and

Suppl. Figure S1B-C). The loss of the wing margin is associated with a reduction in Notch

signaling or in effects on the Wg signaling pathway, as *wg* is a direct target of Notch signaling

along the dorsal/ventral compartment border. Besides the posterior compartment of the wing

being slightly reduced, no other patterning defects were observed in the *baz* knockdowns

(Figure 1A-C). In contrast to *baz*, knockdown of either *aPKC* or *par6* caused cell lethality with

loss of large areas of the posterior wing compartment at 25°C or 29°C (Suppl. Figure S1). When

aPKC-IR was expressed at reduced levels (driven at 18°C), causing a milder knock-down, the

posterior wing compartment was still markedly smaller (Figure 1D; note that wing hairs were

also often very short, Fig. 1D-D', orange arrows), but the wing margin appeared largely normal

(Figure 1B'-D'). PCP defects were also observed in the *aPKC-IR* expressing domain (Figure

1D', green arrow indicating PCP hair orientation), consistent with the previously identified role of

aPKC in PCP (Djiane et al., 2005). *Par6-IR* mediated knock-down caused severe loss of wing

tissues even at 18°C, making it impossible to correlate these defects with a specific cellular

mechanism or signaling pathway (Suppl. Figure S1D-D').   Nevertheless, the wing margin

appeared relatively normal in areas where there was no tissue loss (Suppl. Figure S1D-D',

magenta arrows).

Taken together, knockdown of *baz* caused a distinct loss-of-function (LOF) phenotype from

aPKC and *Par6*, affecting wing margin development. To confirm this effect, we also analyzed

wings containing mutant clones of baz^4 or baz^{815-8} alleles (Suppl. Figure S1A for allele

information). Consistent with the RNAi knock-downs, *baz* LOF clones also displayed wing

margin loss (Suppl. Figure S2A-C). On the notum, *baz* mutant LOF clones displayed loss of

sensory bristles (Suppl. Figure S2D-F), in accordance with previous work (Roegiers et al., 2001)

confirming the behavior of the *baz* alleles. Taken together with the RNAi phenotypes above, the

baz LOF clonal phenotypes in the wing suggest a potential role for Baz/Par3 in Notch-signaling.

***Notch* and *baz* interact genetically**

To corroborate a relationship between *Notch* and *baz* phenotypes, we performed genetic

interaction analyses. In heterozygous conditions of the null allele N^{55e11}/+, approx. 30% of wings

displayed distal margin notching (near vein 3 and 4; Figure 2A, B, and F). We compared these

wings to double heterozygous combinations with the LOF *baz* alleles: baz^4 or baz^{815-8}.

Strikingly, the double heterozygous animals displayed a marked enhancement of wing notching

with 68% and 79% of wing margin defects in N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} allelic combinations,

respectively (Figure 2C-D, and F; see also Suppl. Figure S3). Moreover, in addition to the

enhanced distal wing notch phenotype, a novel defect appeared in the double heterozygous

animals with wing notching in the proximal region of the posterior compartment (Figure 2B', D'

and quantified in 2G; see also Suppl. Figure S3B'-C'). Whereas N^{55e11}/+ wings did not display

any margin defects in this proximal region (Figure 2B' and 2G), 100% of the double

heterozygous N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} wings carried such margin defects (quantified in

Figure 2G; $p<0.0001$). The equivalent baz^4/+ and baz^{815-6}/+ heterozygous control wings did not

display any margin notching phenotypes (Figure 2E-E', also Suppl. Figure S3C-C'). These data

suggest that *Notch* and *baz* genetically interact during wing margin patterning in a synergistic

manner.

We next asked whether interactions between *Notch* and *baz* are also detected in other tissues

and with a gain-of-function (GOF) N genotype. The specification of cell fates in the developing

eye requires Notch input at multiple steps, from early patterning of the presumptive eye field to

the specification of individual photoreceptor (R cell) and accessory cell fates. A classical assay

for N-signaling during eye patterning is the expression of constitutively active (ca)-N molecules

under control of the *sevenless* (*sev*) enhancer (expressed in R3-4 and R1,6, and R7). Such ca-

Notch molecules are either N-terminally truncated molecules that lack the inhibitory region in the

extracellular domain preventing g-secretase-mediated cleavage (NDECD, (Fortini et al., 1993)),

or the intracellular domain itself, which lacks the membrane targeting sequence (NICD, (Kidd et

al., 1998)). Depending upon the strength of activation, ectopic Notch activity induces the

transformation of R3 to R4 and/or the R1/R6 photoreceptors to R7 (Cooper and Bray, 1999;

Fanto and Mlodzik, 1999; Tomlinson and Struhl, 1999, 2001). The induction of R7 is a classical

Notch-dependent cell fate and ectopic activation of N causes the transformation of R1/R6

photoreceptors to R7 independently of any PCP input (Tomlinson and Struhl, 2001). In the *sev-*

NICD background a large fraction of ommatidia display a multi-R7 phenotype (Figure 2H, and

(Tomlinson and Struhl, 2001)), while the presence of one copy of the strong baz^{815-8} allele in this

background suppresses the GOF Notch effect (Figure 2H). Taken together with the above

genetic interactions with *N* LOF features, these results suggest that Baz/Par3 is generally

required for Notch signaling function in different tissues and contexts, as seen in both, LOF and

GOF, scenarios.

baz loss-of-function leads to reduction of Notch target gene expression

To test whether *baz-IR* affects Notch signaling directly, we examined the expression of the

endogenous Notch signaling targets *wg* and *cut* (Micchelli et al., 1997; Neumann and Cohen,

1996). To monitor *wg* expression we analyzed the expression of *wg-lacZ*, an enhancer trap

insertion in the *wg* locus, which faithfully reproduces *wg* expression patterns (Kassis et al.,

1992); and to monitor *cut* expression we used an antibody to Cut. Knock-down of *baz* in the

posterior compartment caused reduction of both *wg-lacZ* expression and Cut protein levels, as

compared to control levels in the anterior compartment (Figure 3A-C"). Both *wg-lacZ* and Cut

levels were significantly reduced in quantifications of their intensity ratio between the anterior

(control) and posterior (experimental) compartment (Figure 3D-E). Since the reduction of *wg*

and *cut* expression is detected at the third instar larval stage in wing discs, before sensory

development occurs at the pupal stages (Jack et al., 1991), the *baz* effect on their expression

precedes sensory cell differentiation and is thus clearly separated in time from the known Baz

function in asymmetric SOP cell division.

To confirm these results, we also examined the expression of a synthetic reporter containing the

Notch response element (NRE), *NRE-GFP* (Furriols and Bray, 2001; Saj et al., 2010). In wing

discs, *NRE-GFP* is expressed along the dorsal/ventral compartment border in pattern

indistinguishable from endogenous Notch target genes such as *wg* and *cut* (Saj et al., 2010)

(Suppl. Figure S4A). Importantly, reduced expression of *NRE-GFP* was observed in the

posterior compartment in *enGal4>baz-IR(s)* wings (Suppl. Figure S4B-C', and E), indicating that

a reduction of *baz* directly correlated with reduced Notch-signaling and target gene expression.

To corroborate these data and to test for a general effect of *baz* LOF on Notch signaling

outcomes in different experimental set ups, we analyzed the expression of Notch targets in *baz*

mutant clones. Consistent with the *baz-IR* knockdown in wing discs, Cut expression levels were

reduced in *baz$^{815-8}$* mutant cells (clones were positively labeled by RFP expression, Figure 4A-C,

examples highlighted with white arrows; quantified in Figure 4D). Next we tested for effects on

Notch signaling in eye discs. To test an independent Notch target gene, we asked whether

expression of *atonal* (*ato*), a N-signaling target during early eye patterning prior to R-cell

specification (Freeman, 2001) also requires *baz* for its full expression. Ato levels were reduced

in *baz$^{815-8}$* mutant cells (orange arrows in Figure 4E-E", F-F", mutant clone marked by lack of

GFP). Importantly, apical-basal polarity as apparent in optical z-sections was normal in *baz$^-$*

clones (Figure 4G-G"; note Patj as A/B marker being localized normally in *baz$^{815-8}$* clones – see

Discussion), whereas a reduction of Ato expression was also easily detected in this optical

plane (Figure 4D-F, orange arrows in E" and F"; quantified in Figure 4H),

Taken together, these results are consistent with the notion that Baz functions as a positive

regulator within the Notch-signaling pathway, promoting the expression of Notch target genes.

Baz affects Notch-signaling at the NICD level

To mechanistically dissect Baz function in Notch signaling, we used N-terminal Notch

truncations in GOF assays to determine at what level within the pathway Baz was required.

First, we used NDECD (Fortini et al., 1993), which lacks the extracellular domain and is

constitutively active independent of ligand binding (see above). As such, if NDECD were

affected by *baz* LOF, Baz would be required downstream of the N-ligands and the subsequent

extracellular cleavages of Notch, but upstream of the g-secretase-mediated cleavage

(Andersson et al., 2011; Struhl and Greenwald, 1999). When NDECD was clonally expressed in

larval wing discs, 43% of NDECD clones induced the Notch target *wg* (Figure 5A, F; as detected

by Wg antibody). When *baz-IR(*s) were co-expressed with NDECD, the frequency of clones

expressing Wg was reduced to 20% (*baz-IR*-1) and 27% (*baz-IR*-2), respectively (Figure 5B, F).

These data suggested that Baz was required at the level of the g-secretase cleavage or further

downstream within the Notch-signaling pathway. Consistent with this hypothesis *baz-IR* knock-

down displayed no detectable effect on Notch protein levels at the membrane or cellular

distribution of Notch (Suppl. Figure S1E-F).

To further investigate the position of Baz within the Notch pathway, we tested the NICD

fragment, which is fully active and does not associate with the membrane or require any

cleavage and is thought to directly translocate to the nucleus to activate Notch-signaling target

genes as transcriptional co-activator (Andersson et al., 2011; Bray, 2016; Struhl and Greenwald,

1999). In control wing disc clones expressing NICD, Wg expression was readily detected in 63%

of clones (or 71% with a control IR co-expression) (Figure 5C-D, and G). In experimental

clones, with *baz-IR*(s) being co-expressed with NICD, the NICD effect was markedly

suppressed: to 41% with *baz-IR*-1 and 32% with *baz-IR*-2, respectively (Figure 5E, G). These

data are consistent with the notion that Baz acts downstream of membrane tethered Notch and

that it is required for the NICD fragment to activate Notch-signaling target gene expression.

In the nucleus, NICD functions as a co-activator interacting with Su(H) (the *Drosophila* member

of the CSL transcription factor family), Mastermind (Mam), and other co-activators to form a

complex (or complexes) at transcriptional enhancer sites to activate Notch target gene

expression (Bray, 2016). To gain insight into whether Baz/Par3 promotes the formation of the

Su(H)-NICD-Mam complex, we asked *whether baz* LOF could reduce the activity of an active

Su(H)-Ank fusion protein. A direct fusion of the Ankyrin repeats of Notch to Su(H), was shown in

Xenopus to bypass a requirement for endogenous Notch itself by directly recruiting Mam, and

thus it is acting even further "downstream" relative to NICD in the pathway (Wettstein et al.,

1997). To this end, we generated a *Drosophila* version of the construct, fusing the Notch ankyrin

repeat region to the Su(H) protein [yielding Su(H)-Ank]. We used the clonal expression assay in

the wing pouch, as used above for NICD for example (Figure 5), and expressed Su(H)-Ank in

equivalent clones with *baz-RNAi*. Strikingly, clonal expression of Su(H)-Ank alone or together

with *baz-RNAi* showed similar percentages of clones inducing expression of *wg* (Suppl. Figure

S5). This result demonstrates that Su(H)-Ank activates Notch target gene expression

independently of Baz. Taken together with the clonal behavior of NICD, these data suggested

that Baz acted on NICD, upstream of the Su(H) based transcription complex, prior to NICD

association with Su(H) and Mam (see Discussion).

To complelement the above *in vivo* studies we next addressed whether Baz and NICD could

physically interact. This was tested biochemically via co-immunoprecipitation (co-IP) in S2 cells

that were co-transfected with HA-tagged NICD and either Baz-myc (full length Baz tagged with

myc) or myc-Hippo (Hpo, as negative control). Protein extracts were immunoprecipitated with

anti-myc antibody and analyzed by immunoblotting with anti-HA antibody. NICD was readily co-

immunoprecipitated with Baz-myc (Figure 6A), but not by the Hpo negative control (Figure 6A-B;

see figure legend for additional controls), suggesting that NICD and Baz/Par3 can physically

associate.

Taken together with the functional *in vivo* data, these results suggested that Baz and NICD were

associated in the same protein complex, and that the presence of Baz is required for normal

NICD function in Notch target gene induction. However, Baz appeared to act upstream of the

Su(H) based transcription factor complex, as it did not have an effect in the context of the

Su(H)-Ank fusion, which bypassed a requirement for NICD.

Baz and Numb are required at different levels within the Notch pathway

To gain further mechanistic insight into how Baz might affect NICD, we tested the relationship of

Baz to Numb, which has itself a complex relationship to Notch-signaling (see also Introduction).

While Numb has been shown to regulate Notch-signaling at the level of trafficking of the

receptor (Couturier et al., 2013; Houssin et al., 2021; Schweisguth, 2015), it has also been

reported to directly interact with NICD and inhibit Notch signaling at the NICD level (Frise et al.,

1996; Guo et al., 1996). To address these possibilities, we wished to test whether *numb* and

baz could interact genetically with each other in the above wing margin specification context.

First we asked, whether reduction of *baz* function (via *baz-RNAi* equivalent to experiments

shown in Figure 1) could be modified by Numb levels. Indeed, loss of wing margin bristle cells

(Figure 7A, C, D, F) was suppressed upon removal of one *numb* gene copy, in a *numb^{15}/+*

heterozygous background (Figure 7B, C, E, F), displaying a dosage sensitive relationship

between the two proteins. To confirm this, we compared phenotypes induced by overexpression

of *numb-GFP* with or without *baz-Flag* in developing wings. Overexpression of *numb-GFP*

suppressed Notch signaling, as detected by margin bristle loss (Suppl. Figure S6A-B and D,

(Frise et al., 1996); of note, Numb-GFP also inhibited Notch function in the context of wing vein

formation). Co-overexpression of *baz-Flag* with *numb-GFP* reduced the inhibitory effect of

Numb on Notch signaling at the wing margin (Suppl. Figure S6B-D). These experiments support

the notion that *numb* and *baz* antagonize each other genetically during wing margin cell

induction with Baz promoting Notch function and Numb inhibiting it.

We next wished to test whether they could act at the same level in Notch-signaling, at the level

of NICD. To this end, we asked whether the NICD clonal expression phenotype, induction of

target gene expression, which is suppressed by *baz-IR* (see Figure 5 in Results above), could

also be affected by Numb levels. Again, clonal NICD expression in the wing pouch induced

Notch-signaling target genes, for example *wg* (Figure 5) and *cut* (Figure 7G-G') and the removal

of *baz* (via RNAi) in such clones reduced the ability of NICD to induce target gene expression

(see Figure 5E,G). We thus asked whether co-overexpression of *numb-GFP* in such NICD

expression clones could also suppress NICD's function in target gene activation. However, we

did not observe an effect of Numb-GFP overexpression on NICD function in target gene

expression (Figure 7G-H, quantified in 7I, as shown for Cut). Taken together with the genetic

interactions above, these data suggested that Numb and Baz act at different levels in the Notch

pathway, with Numb acting further upstream and Baz/Par3 acting at the level of NICD.

Discussion

In this study we define a novel, direct, and positive role of the polarity protein Baz/Par3 in the

Notch pathway. This stands in contrast to its previously established role in Notch signaling

regulation during asymmetric SOP cell divisions, where Baz/Par3 affects N-signaling indirectly

by asymmetrically preventing the inheritance of the Notch inhibitor Numb (Schweisguth, 2015).

In the SOP scenario, the Baz/Par3-Par6-aPKC protein complex localizes to one side within the

dividing cell, and a Numb-Neuralized containing complex localizes to the other side of the cell.

As Numb inhibits Notch signaling by regulating Notch-Sanpodo trafficking and thus reducing

Notch levels at the cell surface, this asymmetry causes Notch inhibition in the Numb containing

daughter cell (reviewed in (Schweisguth, 2015)). In the SOP context, Baz/Par3 has been thus

proposed to have a passive role in N-signaling activation, via inhibiting an inhibitor, Numb. Our

data are consistent with the notion that Baz/Par3 is also required positively for the Notch

signaling response and full transcriptional activation of Notch targets in *Drosophila* in multiple

cell types. Genetic epistasis studies indicate that Baz acts at the level of cleaved active Notch,

NICD, or in parallel to it. Consistent with this notion Baz and NICD co-immunoprecipitated,

suggesting that Baz promotes Notch signaling at the level of NICD.

A novel Baz/Par3 function independent of the A/B polarity complex

Baz/Par3 is generally associated with Par6-aPKC at early to intermediate stages of A/B polarity

establishment, but later in mature epithelia it is no longer essential for their maintenance

(Morais-de-Sa et al., 2010; St Johnston and Sanson, 2011; Tepass, 2012). Indeed, the fact that

we can recover *baz* LOF clones at all, whereas Par6 and aPKC null alleles are cell lethal,

suggests that *baz* can behave differently to the 'core' A/B factors. In our experimental set-up we

observed no evidence for a Par6 or aPKC involvement in Notch signaling. Our data suggest that

Baz/Par3 regulates NICD independently of its role in the A/B polarity complex. Similar to A/B

polarity establishment, in the context of planar cell polarity (PCP) patterning, Baz/Par3 functions

together with aPKC to regulate Frizzled levels (and hence the PCP outcomes) in the *Drosophila*

eye (Djiane et al., 2005), and thus, while Par6 could not be assayed in this context, these data

suggested that in PCP determination Baz is functioning with established A/B polarity factors. As

such, the discovery that in Notch signaling Baz/Par3 appears to function independently of its

partners in A/B establishment within the same tissue is unexpected.

Par3 homologues have however been shown to act independently of the aPKC-Par6 complex in

several scenarios in mammalian cells, including Hippo/Yap signaling and during epithelial-

mesenchymal transition (Zhang et al., 2016). Par3 has even been suggested to act with Ku70

and Ku80 as part of the DNA-dependent protein kinase complex to regulate double-strand

breaks (Fang et al., 2007)**.**

Why was Baz/Par3 not identified earlier as a component of Notch signaling?

The Notch pathway genes were initially identified through classic genetic collections of

spontaneous mutants showing morphological defects in *Drosophila* and *C. elegans* (Fleming et

al., 1990; Greenwald, 1985; Kopczynski et al., 1988; Lindsley and Zimm, 1992). Subsequent

genetic screens and molecular studies have added many components of the Notch pathway

and many regulatory mechanisms that affect Notch signaling at all levels, ranging from receptor

and ligand trafficking and membrane stability and its membrane associated proteolytic

cleavages to nuclear co-factors and chromatin associated target gene regulation (reviewed in

(Bray, 2016; Henrique and Schweisguth, 2019; Kopan and Ilagan, 2009). In particular, many of

the screens were performed using the notum Notch associated phenotypes and score for

asymmetric division-dependent SOP/bristle patterning defects. It has thus been technically

difficult, if not impossible, to separate potential additional Baz/Par3 involvements in Notch

signaling. Similarly, classical loss-of-function screens in the *Drosophila* embryo would have

missed Baz/Par3 due to its initial requirement in epithelial polarization that could have caused

earlier lethality (Harris and Peifer, 2004; Muller and Wieschaus, 1996), which obscures a

potential role in subsequent signaling events. Indirectly through our screen asking whether A/B

polarity factors can also play a role in PCP establishment in the posterior wing disc

compartment we used a unique experimental scenario: first the margin loss is due to tissue loss

and not asymmetric division of innervated SOPs, and, second, at this later stage of the epithelial

sheet is fully formed and Baz/Par3 is not required for maintenance of A/B polarity. This

combination thus allowed the identification of *baz* as a novel factor that positively regulates

Notch signaling in general.

Baz/Par3 has multiple independent functions during Notch signaling.

Baz/Par3 and Numb have both been implicated in the regulation of Notch signaling via

trafficking of the Notch receptor (Couturier et al., 2013; Houssin et al., 2021; Schweisguth,

2015). In addition, independent of trafficking, Numb has been suggested to inhibit Notch

signaling directly by binding to NICD (Frise et al., 1996; Guo et al., 1996). Overexpression of

Numb indeed can interfere with Notch signaling at the wing margin, causing significant Notch

loss-of-function type defects ((Frise et al., 1996) and this work). To address this directly, we

asked whether Numb overexpression could interfere with NICD function in the context Notch

target gene activation. However, NICD activity was not affected by Numb overexpression,

suggesting that it acted further upstream, likely as suggested at the level of regulating Notch

trafficking, which is consistent with previous studies (Couturier et al., 2013; Houssin et al.,

2021). In contrast to *numb*, knockdown of *baz/par3* directly affected NICD function and

reduction of Baz/Par3 levels significantly reduced NICD function in target gene activation.

Therefore, Baz/Par3 can act at two independent levels within Notch signaling, affecting (i)

membrane trafficking of the Notch receptor (Couturier et al., 2013; Houssin et al., 2021;

Schweisguth, 2015) and (ii) serving a positive function downstream in the pathway at the NICD

level (see below).

415 Baz acts at the NICD level affecting expression of N-signaling targets positively

The Notch-signaling associated DNA binding complex is centered around the Notch-pathway

dedicated DNA-binding factor CSL, Su(H) in *Drosophila*, and it switches from a repressor

complex to a transcriptional activator complex via NICD (Bray, 2016). NICD binds to Su(H), and

recruits co-activators including Mastermind (Mam), and its binding to Su(H)/CSL displaces the

"repressor complex" and induces N-signaling target gene transcription. Several factors have

been identified that associate with Su(H)/CSL and NICD (see reviews (Borggrefe and Oswald,

2009; Bray, 2016)). Genetic epistasis data, both in the wing and the eye, position the function of

Baz/Par3 at the level of NICD after its last cleavage, suggesting that Baz might act in the

nucleus together with NICD or alternatively it might promote stability, nuclear translocation,

function, or retention of NICD. The biochemical immunoprecipitation results support the genetic

data and are consistent with the notion that Baz is a direct regulator of Notch signaling, acting at

the level of NICD. However, the Su(H)-Ank fusion appears to not require Baz/Par3 for its

capability to activate Notch target gene expression, with *baz* knockdown having no effect on

Su(H)-Ank mediated activation, unlike with NICD, suggesting that Baz is acting upstream of the

Su(H)-Ank association. The Ank repeats of NICD are critical within the Notch protein for its

nuclear response as part of the Su(H) transcription factor complex by recruiting Mastermind,

and as such a Su(H)-Ank fusion protein is sufficient to activate Notch signaling target gene

expression (Collu et al., 2012; Wettstein et al., 1997). As *baz* mutations affect the levels of

target gene expression and the Baz protein acts genetically at the level of, or in parallel to,

NICD, Baz/Par3 might be a factor participating in regulating NICD localization, stability, or

modifications to mediate its association with Su(H).

Interestingly, the human Baz/Par3 homologue, PARD3, has been suggested to act at a similar

level in the YAP/TAZ pathway. PARD3, which itself is found in the nucleus in addition to the

junction-bound aPKC/Par6 complex, has been suggested to regulate TAZ phosphorylation and

its nuclear localization (Zhang et al., 2016), as a scaffold for YAP-binding proteins. The Notch

signaling pathway is simpler than the Hippo-YAP/TAZ pathway, as Notch/NICD acts as a

"membrane-tethered" transcriptional co-activator, without the complex cascade of kinase-

dependent steps as in YAP/TAZ regulation. Nonetheless, NICD is still regulated by

phosphorylation events, for example to limit prolonged pathway activity, through the action of

Mastermind and CycC:CDK8 (Fryer et al., 2004). It is thus possible that Baz/Par3 may regulate

NICD phosphorylation and nuclear localization as well.

Generally, the strength of activation of target gene expression depends on NICD stability within

the resulting activator complexes, which also contain histone modifying/remodeling proteins,

and accessibility of enhancers of any given target gene (Borggrefe and Oswald, 2009; Bray,

2016). N-signaling acts in many different contexts and can activate many distinct target genes,

and as such the presence of context-determining factors will determine whether an enhancer is

primed to react to a specific signal, which may result in a gene having different responses

towards the same signals (Borggrefe and Oswald, 2009). Our data suggest that Baz/Par3 acts

rather generally on NICD mediated target gene activation, as we see effects on Notch target

expression in multiple tissues and on different target genes. It is likely that Baz/Par3 is directly

or indirectly involved in promoting NICD association with Su(H), and the formation of the Su(H)-

NICD-Mam complex. The precise function of Baz/Par3 in the context of NICD regulation will

require transcription specific assays to be explored in future studies.

Materials and Methods

Drosophila genetics and strains

w-IR (w^{JF01540}, P{TRiP.JF01540}), baz-IR-1 (baz^{JF01078}, P{TRiP.JF01078}), baz-IR-2 (baz^{JF01079}, P{TRiP.JF01079}), aPKC-IR (aPKC^{JF01966}, P{TRiP. JF01966}), UAS-mCD8RFP, baz^4 FRT9-2 and baz$^{815-8}$ FRT9-2 fly stocks were obtained from Bloomington Drosophila Stock Center. baz$^{815-8}$ was recombined with FRT19A and then used for baz$^{815-8}$ mutant clonal analysis with FRT19A MARCM stock (genotype: w, tub-gal80, hs-flp, FRT19A; act-Gal4, UAS-mCD8RFP / CyO). UAS-numb-GFP was a gift from Bingwei Lu (Song and Lu, 2012).

N^{55e11}/baz^4 and N^{55e11}/baz$^{815-8}$ were generated by intercrossing via several crosses: ubi-baz-mCherry (Bosveld et al., 2012) was recombined with Sb^1 to generate ubi-baz-mcherry Sb^1 flies. baz^4 (or baz$^{815-8}$) males with ubi-baz-mCherry Sb^1 which rescued baz$^-$ mutations (baz^4 or baz$^{815-8}$; ubi-baz-mCherry Sb^1/+) were crossed to N^{55e11}/FM7 females to generate N^{55e11}/baz^4 or N^{55e11}/baz$^{815-8}$ females (selected against Sb^1 phenotype).

UAS-NICD (Zaffran and Frasch, 2000) flip-on clones were analyzed and induced together with either UAS-GFP, baz-IR1, or baz-IR2, UAS-numbGFP, and UAS-Su(H)Ank (full genotypes listed in the figure legends). Third instar larvae were dissected 44-49 hours after clone inductions (which were induced via brief heat-shocks). Control samples from the same set of experiments were heat-shocked, processed and stained together to reduce experimental variations.

DNA constructs:

The baz-pENTR was kindly provided by Tony Harris (McKinley et al., 2012). Gateway cloning (Invitrogen) recombined pPWF (from Drosophila Genomics Resource Center) for UASP-baz (C-terminal 3X Flag tagged baz). The construct was sequenced with baz oligos to confirm correct recombination insertion and the Flag tag sequence. Transgenic flies were injected with UASP-baz-Flag ppWF plasmid by BestGene.

A fusion of *D. melanogaster* Su(H) and the Notch Ankyrin repeats was designed along the same

principle as X-Su(H)-ANK (Wettstein et al., 1997). *Su(H)* coding sequence was cloned from

GH10914 (DGRC Stock 5973) into pAc5.1/V5-His A (Invitrogen) using the following primers:

Fwd: ACCGGAATTCACCatgaagagctacagccaatt and

Rev: ACGCGGATCCggataagccgctaccatgac to remove the stop codon. The sequence encoding

Notch Ankyrin repeats were cloned from pAc5.1/N[IC]-HA (gift of Ram DasGupta) using the

following primers:

Fwd: ACGCGGATCCgatgtggatgcacgtggacct and

Rev: CTTTGCGGCCGCtcacagatcttcttcagaaataagtttttgttcatgctcgtccagcagcctga

to introduce a stop codon and myc epitope. The Su(H)-ANK-myc sequence was then subcloned

into pUASt-AttB, (and injected into UAS-AttP flies (RRID:BDSC_9724) by BestGene. Multiple

transgenic lines were sequenced and phenotypically tested to confirm insertion of the UAS-

Su(H)-ANK-myc sequence.

Immunohistochemistry and microscopy:

The following antibodies were used in the study: guinea pig Anti-Baz 1:200 (Blankenship et al.,

2007), rabbit anti-Dlg 1:3000-4000 (Izaddoost et al., 2002), mouse anti-Wg 1:10-20 (Brook and

Cohen, 1996), mouse Anti-Cut 1:20-1:40 (Blochlinger et al., 1990), rabbit anti-RFP 1:300

(Rockland Immunochemicals), rabbit anti-Galactosidase 1:2500 (Cappel). Mouse anti-Cut and

anti-Wg antibodies were obtained from Developmental Studies Hybridoma Bank at the

University of Iowa.

Adult wings and nota were imaged with conventional Zeiss Axioplan microscope. Immuno-

fluorescent samples were imaged with Leica TCS SP5 or SP8 confocal microscope. Images

were projected and quantified with Fiji-Image J. Cut expression levels were measured in cell

nuclei marked by Hoechst staining. *baz*[815-8] mutant cells were outlined by RFP expression. Only

first row of *baz*[815-8] mutant cells right at D/V border were measured for "average Cut channel

pixel intensity" minus "average background pixel intensity outside Cut expression domain"

(similar to Figure S4D1) for each nucleus. Neighboring control A/P border cells were also

measured for "average Cut pixel intensity" minus "average background pixel intensity".

Data visualization and Statistics:

Data were plotted via Microsoft Excel, LibreOffice Calc. Fisher's exact test (two tailed) and

student t-test (unpaired) were used for statistical analyses to calculate probabilities of null

hypothesis with the web version of Graphpad (Fisher's exact test) and scipy.stats (t-test).

Cell Culture and co-immunoprecipitation

NICD-HA pAc and myc-Hpo pAc were generous gifts from Ramanuj DasGupta and Duojia Pan

(Wu et al., 2003). Baz-myc pAWM was generated from pAWM gateway vector (Drosophila

Genomics Resource Center) and *baz* pEntr 2B (a gift from Tony Harris).

1mg of NICD-HA pAc and Baz-myc pAWM or myc-Hpo pAc plasmids each were transfected

into *Drosophila* S2 cells in a 6-well plate with Qiagen Effectene. Cells were lysed one day after

transfection with IP buffer (20 mM HEPES pH 7.5, 100 mM NaCl, 0.05% Triton X100, 1 mM

EDTA, 5 mM DTT, 10mM NaVO4, 10% glycerol and protease inhibitor cocktail). The cell

lysates (1mg of total protein) were pre-cleared by incubating with protein A-sepharose beads

(Thermo Scientific 20333) for 1 hour at 4°C, followed by centrifugation to remove the beads.

Mouse anti-Myc antibody (sc-40, Santa Cruz technology) was used at 1/1000 dilution for

immunoprecipitation. The A-sepharose beads and antibody were incubated together to form

pre-immuno-precipitate complexes at 4°C for 1 hour, then incubated with lysates. Myc-tagged

protein and associated proteins were captured by protein A-sepharose/antibody complexes after

overnight incubation at 4°C. Samples were washed 4 times with IP buffer. Immunoprecipitates

were resuspended in SDS sample buffer, boiled for 5 min, separated by SDS-PAGE, and

transferred to nitrocellulose membrane for immunoblotting. The blots were incubated with

mouse anti-myc (Santa Cruz technology) or rat anti-HA (Roche) at 1:1000, subsequently with

anti-mouse or anti-rat IgG-HRP. Proteins were then detected by enhanced chemiluminescence

kit (Millipore).

Acknowledgements

We are grateful to Kwang-Wook Choi, Kyung-ok Cho, Ramanuj DasGupta, Tony Harris, Duojia

Pan, Bingwei Lu, and Jennifer Zallen for *Drosophila* strains and antibodies, and to Cathie

Pfleger for helpful suggestions on the manuscript. We also thank the Bloomington Drosophila

Stock Center for fly strains, and the Developmental Studies Hybridoma Bank at the University of

Iowa for antibodies. We would also like to thank the ISMMS Microscopy CoRE, where confocal

microscopy was performed and was in part supported by the Tisch Cancer Institute P30

CA196521 grant from the NCI. This work was supported by National Institutes of Health grants

R01 EY013256 and R35 GM127103 (to MM).

References:

Andersson, E.R., Sandberg, R., Lendahl, U., 2011. Notch signaling: simplicity in design,
versatility in function. Development (Cambridge, England) 138, 3593-3612.

Bala Tannan, N., Collu, G., Humphries, A.C., Serysheva, E., Weber, U., Mlodzik, M., 2018.
AKAP200 promotes Notch stability by protecting it from Cbl/lysosome-mediated degradation in
Drosophila melanogaster. PLoS genetics 14, e1007153.

Blankenship, J.T., Fuller, M.T., Zallen, J.A., 2007. The Drosophila homolog of the Exo84
exocyst subunit promotes apical epithelial identity. Journal of cell science 120, 3099-3110.

Blochlinger, K., Bodmer, R., Jan, L.Y., Jan, Y.N., 1990. Patterns of expression of cut, a protein
required for external sensory organ development in wild-type and cut mutant Drosophila
embryos. Genes & development 4, 1322-1331.

Borggrefe, T., Oswald, F., 2009. The Notch signaling pathway: transcriptional regulation at
Notch target genes. Cell Mol Life Sci 66, 1631-1646.

Bosveld, F., Bonnet, I., Guirao, B., Tlili, S., Wang, Z., Petitalot, A., Marchand, R., Bardet, P.L.,
Marcq, P., Graner, F., Bellaiche, Y., 2012. Mechanical control of morphogenesis by
Fat/Dachsous/Four-jointed planar cell polarity pathway. Science (New York, N.Y 336, 724-727.

Bray, S.J., 2016. Notch signalling in context. Nat Rev Mol Cell Biol 17, 722-735.

Brook, W.J., Cohen, S.M., 1996. Antagonistic interactions between wingless and
decapentaplegic responsible for dorsal-ventral pattern in the Drosophila Leg. Science (New
York, N.Y 273, 1373-1377.

Collu, G.M., Hidalgo-Sastre, A., Acar, A., Bayston, L., Gildea, C., Leverentz, M.K., Mills, C.G.,
Owens, T.W., Meurette, O., Dorey, K., Brennan, K., 2012. Dishevelled limits Notch signalling
through inhibition of CSL. Development (Cambridge, England) 139, 4405-4415.

Cooper, M.T., Bray, S.J., 1999. Frizzled regulation of Notch signalling polarizes cell fate in the
Drosophila eye. Nature 397, 526-530.

Courbard, J.R., Djiane, A., Wu, J., Mlodzik, M., 2009. The apical/basal-polarity determinant
Scribble cooperates with the PCP core factor Stbm/Vang and functions as one of its effectors.
Developmental biology 333, 67-77.

Couturier, L., Mazouni, K., Schweisguth, F., 2013. Numb localizes at endosomes and controls
the endosomal sorting of notch after asymmetric division in Drosophila. Curr Biol 23, 588-593.

Djiane, A., Yogev, S., Mlodzik, M., 2005. The apical determinants aPKC and dPatj regulate
Frizzled-dependent planar cell polarity in the Drosophila eye. Cell 121, 621-631.

Etemad-Moghadam, B., Guo, S., Kemphues, K.J., 1995. Asymmetrically distributed PAR-3
protein contributes to cell polarity and spindle alignment in early C. elegans embryos. Cell 83,
743-752.

Fang, L., Wang, Y., Du, D., Yang, G., Tak Kwok, T., Kai Kong, S., Chen, B., Chen, D.J., Chen,
Z., 2007. Cell polarity protein Par3 complexes with DNA-PK via Ku70 and regulates DNA
double-strand break repair. Cell research 17, 100-116.

Fanto, M., Mlodzik, M., 1999. Asymmetric Notch activation specifies photoreceptors R3 and R4
and planar polarity in the Drosophila eye. Nature 397, 523-526.

Fleming, R.J., Scottgale, T.N., Diederich, R.J., Artavanis-Tsakonas, S., 1990. The gene Serrate
encodes a putative EGF-like transmembrane protein essential for proper ectodermal
development in Drosophila melanogaster. Genes & development 4, 2188-2201.

Fortini, M.E., Rebay, I., Caron, L.A., Artavanis-Tsakonas, S., 1993. An activated Notch receptor
blocks cell-fate commitment in the developing Drosophila eye. Nature 365, 555-557.

Freeman, A.B.a.M., 2001. Notch signalling and the initiation of neural development in the
Drosophila eye. Development (Cambridge, England) 128, 3889-3898.

Frise, E., Knoblich, J.A., Younger-Shepherd, S., Jan, L.Y., Jan, Y.N., 1996. The Drosophila
Numb protein inhibits signaling of the Notch receptor during cell-cell interaction in sensory organ
lineage. Proceedings of the National Academy of Sciences of the United States of America 93,
11925-11932.

Fryer, C.J., White, J.B., Jones, K.A., 2004. Mastermind recruits CycC:CDK8 to phosphorylate
the Notch ICD and coordinate activation with turnover. Molecular cell 16, 509-520.

Furriols, M., Bray, S., 2001. A model Notch response element detects Suppressor of Hairless-
dependent molecular switch. Curr Biol 11, 60-64.

Goldstein, B., Macara, I.G., 2007. The PAR proteins: fundamental players in animal cell
polarization. Developmental cell 13, 609-622.

Greenwald, I., 1985. lin-12, a nematode homeotic gene, is homologous to a set of mammalian
proteins that includes epidermal growth factor. Cell 43, 583-590.

Guo, M., Jan, L.Y., Jan, Y.N., 1996. Control of daughter cell fates during asymmetric division:
interaction of Numb and Notch. Neuron 17, 27-41.

Guruharsha, K.G., Kankel, M.W., Artavanis-Tsakonas, S., 2012. The Notch signalling system:
recent insights into the complexity of a conserved pathway. Nature reviews 13, 654-666.

Harris, T.J., Peifer, M., 2004. Adherens junction-dependent and -independent steps in the
establishment of epithelial cell polarity in Drosophila. The Journal of cell biology 167, 135-147.

Henrique, D., Schweisguth, F., 2019. Mechanisms of Notch signaling: a simple logic deployed in
time and space. Development (Cambridge, England) 146, dev172148 doi:
172110.171242/dev.172148.

Houssin, E., Pinot, M., Bellec, K., Le Borgne, R., 2021. Par3 cooperates with Sanpodo for the
assembly of Notch clusters following asymmetric division of Drosophila sensory organ precursor
cells. eLife 10.

Humphries, A.C., Mlodzik, M., 2018. From instruction to output: Wnt/PCP signaling in
development and cancer. Curr Opin Cell Biol 51, 110-116.

Izaddoost, S., Nam, S.C., Bhat, M.A., Bellen, H.J., Choi, K.W., 2002. Drosophila Crumbs is a
positional cue in photoreceptor adherens junctions and rhabdomeres. Nature 416, 178-183.

Jack, J., Dorsett, D., Delotto, Y., Liu, S., 1991. Expression of the cut locus in the Drosophila
wing margin is required for cell type specification and is regulated by a distant enhancer.
Development (Cambridge, England) 113, 735-747.

Kandachar, V., Roegiers, F., 2012. Endocytosis and control of Notch signaling. Curr Opin Cell
Biol 24, 534-540.

Kassis, J.A., Noll, E., VanSickle, E.P., Odenwald, W.F., Perrimon, N., 1992. Altering the
insertional specificity of a Drosophila transposable element. Proceedings of the National
Academy of Sciences of the United States of America 89, 1919-1923.

Kidd, S., Lieber, T., Young, M.W., 1998. Ligand-induced cleavage and regulation of nuclear
entry of Notch in Drosophila melanogaster embryos. Genes & development 12, 3728-3740.

Kopan, R., Ilagan, M.X., 2009. The canonical Notch signaling pathway: unfolding the activation
mechanism. Cell 137, 216-233.

Kopczynski, C.C., Alton, A.K., Fechtel, K., Kooh, P.J., Muskavitch, M.A., 1988. Delta, a
Drosophila neurogenic gene, is transcriptionally complex and encodes a protein related to blood
coagulation factors and epidermal growth factor of vertebrates. Genes & development 2, 1723-
1735.

Lindsley, D.L., Zimm, G.G., 1992. The Genome of Drosophila melanogaster.

McKinley, R.F., Yu, C.G., Harris, T.J., 2012. Assembly of Bazooka polarity landmarks through a
multifaceted membrane-association mechanism. Journal of cell science 125, 1177-1190.

Medina-Yáñez, I., Olivares, G.H., Vega-Macaya, F., Mlodzik, M., Olguín, P., 2020. Phosphatidic
acid increases Notch signalling by affecting Sanpodo trafficking during Drosophila sensory
organ development. Scientific reports 10, 21731.

Micchelli, C.A., Rulifson, E.J., Blair, S.S., 1997. The function and regulation of cut expression on
the wing margin of Drosophila: Notch, Wingless and a dominant negative role for Delta and
Serrate. Development (Cambridge, England) 124, 1485-1495.

Mohr, O.L., 1919. Character changes caused by mutation of an entire region of a chromosome
in Drosophila. Genetics 4, 275-282.

Morais-de-Sa, E., Mirouse, V., St Johnston, D., 2010. aPKC phosphorylation of Bazooka
defines the apical/lateral border in Drosophila epithelial cells. Cell 141, 509-523.

Muller, H.A., Wieschaus, E., 1996. armadillo, bazooka, and stardust are critical for early stages
in formation of the zonula adherens and maintenance of the polarized blastoderm epithelium in
Drosophila. The Journal of cell biology 134, 149-163.

Neumann, C.J., Cohen, S.M., 1996. A hierarchy of cross-regulation involving Notch, wingless,
vestigial and cut organizes the dorsal/ventral axis of the Drosophila wing. Development
(Cambridge, England) 122, 3477-3485.

Neumuller, R.A., Knoblich, J.A., 2009. Dividing cellular asymmetry: asymmetric cell division and
its implications for stem cells and cancer. Genes & development 23, 2675-2699.

Rana, N.A., Haltiwanger, R.S., 2011. Fringe benefits: functional and structural impacts of O-
glycosylation on the extracellular domain of Notch receptors. Current opinion in structural
biology 21, 583-589.

Roegiers, F., Younger-Shepherd, S., Jan, L.Y., Jan, Y.N., 2001. Bazooka is required for
localization of determinants and controlling proliferation in the sensory organ precursor cell
lineage in Drosophila. Proceedings of the National Academy of Sciences of the United States of
America 98, 14469-14474.

Saj, A., Arziman, Z., Stempfle, D., van Belle, W., Sauder, U., Horn, T., Durrenberger, M., Paro,
R., Boutros, M., Merdes, G., 2010. A combined ex vivo and in vivo RNAi screen for notch
regulators in Drosophila reveals an extensive notch interaction network. Developmental cell 18,
862-876.

Schweisguth, F., 2015. Asymmetric cell division in the Drosophila bristle lineage: from the
polarization of sensory organ precursor cells to Notch-mediated binary fate decision. Wiley
Interdiscip Rev Dev Biol 4, 299-309.

Shilo, B.Z., Schejter, E.D., 2011. Regulation of developmental intercellular signalling by
intracellular trafficking. Embo j 30, 3516-3526.

Siebel, C., Lendahl, U., 2017. Notch Signaling in Development, Tissue Homeostasis, and
Disease. Physiol Rev 97, 1235-1294.

Singh, J., Mlodzik, M., 2012. Hibris, a Drosophila nephrin homolog, is required for presenilin-
mediated Notch and APP-like cleavages. Developmental cell 23, 82-96.

Song, Y., Lu, B., 2012. Interaction of Notch signaling modulator Numb with α-Adaptin regulates
endocytosis of Notch pathway components and cell fate determination of neural stem cells. The
Journal of biological chemistry 287, 17716-17728.

Southgate, L., Sukalo, M., Karountzos, A.S.V., Taylor, E.J., Collinson, C.S., Ruddy, D., Snape,
K.M., Dallapiccola, B., Tolmie, J.L., Joss, S., Brancati, F., Digilio, M.C., Graul-Neumann, L.M.,
Salviati, L., Coerdt, W., Jacquemin, E., Wuyts, W., Zenker, M., Machado, R.D., Trembath, R.C.,
2015. Haploinsufficiency of the NOTCH1 Receptor as a Cause of Adams-Oliver Syndrome With
Variable Cardiac Anomalies. Circ Cardiovasc Genet 8, 572-581.

St Johnston, D., Sanson, B., 2011. Epithelial polarity and morphogenesis. Curr Opin Cell Biol
23, 540-546.

Struhl, G., Greenwald, I., 1999. Presenilin is required for activity and nuclear access of Notch in
Drosophila. Nature 398, 522-525.

Suzuki, A., Ohno, S., 2006. The PAR-aPKC system: lessons in polarity. Journal of cell science
119, 979-987.

Tepass, U., 2012. The apical polarity protein network in Drosophila epithelial cells: regulation of
polarity, junctions, morphogenesis, cell growth, and survival. Annual review of cell and
developmental biology 28, 655-685.

Tomlinson, A., Struhl, G., 1999. Decoding vectorial information from a gradient: sequential roles
of the receptors Frizzled and Notch in establishing planar polarity in the Drosophila eye.
Development (Cambridge, England) 126, 5725-5738.

Tomlinson, A., Struhl, G., 2001. Delta/Notch and Boss/Sevenless signals act combinatorially to
specify the Drosophila R7 photoreceptor. Molecular cell 7, 487-495.

Weber, U., Eroglu, C., Mlodzik, M., 2003. Phospholipid membrane composition affects EGF
receptor and Notch signaling through effects on endocytosis during Drosophila development.
Developmental cell 5, 559-570.

Wettstein, D.A., Turner, D.L., Kintner, C., 1997. The Xenopus homolog of Drosophila
Suppressor of Hairless mediates Notch signaling during primary neurogenesis. Development
(Cambridge, England) 124, 693-702.

Wharton, K.A., Johansen, K.M., Xu, T., Artavanis-Tsakonas, S., 1985. Nucleotide sequence
from the neurogenic locus notch implies a gene product that shares homology with proteins
containing EGF-like repeats. Cell 43, 567-581.

Wilson, A., Radtke, F., 2006. Multiple functions of Notch signaling in self-renewing organs and
cancer. FEBS letters 580, 2860-2868.

Wu, S., Huang, J., Dong, J., Pan, D., 2003. hippo encodes a Ste-20 family protein kinase that
restricts cell proliferation and promotes apoptosis in conjunction with salvador and warts. Cell
114, 445-456.

Wu, S., Yang, Y., Tang, R., Zhang, S., Qin, P., Lin, R., Rafel, N., Lucchetta, E.M., Ohlstein, B.,
Guo, Z., 2023. Apical-basal polarity precisely determines intestinal stem cell number by
regulating Prospero threshold. Cell Rep 42, 112093.

Zaffran, S., Frasch, M., 2000. Barbu: an E(spl) m4/m(alpha)-related gene that antagonizes
Notch signaling and is required for the establishment of ommatidial polarity. Development
(Cambridge, England) 127, 1115-1130.

Zhang, P., Wang, S., Wang, S., Qiao, J., Zhang, L., Zhang, Z., Chen, Z., 2016. Dual function of
partitioning-defective 3 in the regulation of YAP phosphorylation and activation. Cell Discov 2,
16021.

 Figures

Figure 1. Reduction of *bazooka/baz function* causes wing margin bristle loss.

(A-D) Overview and high magnification view (A'-D') of wings of the indicated genotypes.

(A, A') Wild-type wing; note the associated margin bristles in high magnification of area of

posterior wing margin (A'). Cellular hairs point towards the distal margin (green arrow).

(B, B') *enGal4, UAS-baz-IR-1* wing, with *baz* knock-down in the posterior compartment; note

loss of margin bristles (magenta arrows). Cellular hair orientation was not significantly affected.

(Genotype: *UAS-dcr2; enGal4, UAS-RFP, NRE-GFP / UAS-Baz-IR-1 at 29°C*)

(C, C') *enGal4, UAS-baz-IR2* wing (independent RNAi, see Fig. S1), displaying similar loss of

margin bristles phenotype with hair orientation not significantly affected. (Genotype: *UAS-dcr2;*

enGal4, UAS-RFP, NRE-GFP / UAS-Baz-IR-2 at 29°C)

(D, D') *enGal4, UAS-aPKC-IR* wing, displaying largely normal margin bristle development. Note

wing size reduction in posterior compartment with some areas displaying abnormally short wing

hairs (orange arrow) and other areas with cellular hair orientation defects, suggesting that

reduction of aPKC affected several aspects of wing development, yet had no effect on margin

bristle development (Genotype: *enGal4* /+; *UAS-aPKC-IR (aPKC[JF01966])* /+ at 19°C)

(E-E") RNAi-based knock down of Baz protein in posterior compartment. All panels are showing

max-projections of a stack of confocal images (~40mm to cover the apical surface of all of the

wing pouch area), including composite (E) and individual channels. Anti-Dlg (Discs large, green

and E') staining, serves as positive control to outline cells in both anterior (labeled "A") and

posterior compartments (labeled "P"). Anti-Baz (magenta) levels were markedly reduced in the

posterior *baz-IR* expressing compartment, suggesting an effective knock-down via the *baz* RNAi

transgene. (Genotype: *dcr2; enGal4/+; /UAS-baz-IR-2 29°C*).

Figure 2. *Notch* and *baz* interact genetically during wing margin patterning.

(A-B) Heterozygous N^{55e11}/+ wings display wing margin notching phenotypes; approx. 2/3 of N/+

wings are normal with no margin loss (example in A), while 1/3 displayed distal margin notches

(B, arrow). Proximal posterior wing margin was always normal (boxed area in B, and B') in all

animals. Quantified in panels (F) and (G).

(C-D) N^{55e11}/baz^{815-8} double heterozygous wings. Note increased margin notching in distal area

(D, arrows) and proximal-posterior wing margin in all wings (C, D and D', arrows). (D') Higher

magnification of boxed area in D. Quantified in panels (F) and (G).

(E) baz^{815-8}/+ single heterozygous wings show no notching/tissue loss at margin, neither distally

nor in the proximal-posterior region (E', boxed area from E shown at higher magnification).

(F) Quantification of distal wing margin phenotypes as percentage of wings containing notching

(genotypes as indicated). Note that N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} flies show enhanced defects

(higher percentage) of margin notching as compared to N^{55e11}/+ alone (p=0.0062 and p=0.0033,

respectively (Fisher's exact test). The control baz^4/+ and baz^{815-8}/+ single heterozygous flies

have no wing margin phenotypes.

(G) Quantification of proximal wing margin phenotypes (posterior compartment) as percentage

of wings with margin loss (genotypes as indicated). 100% of N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} had

proximal wing margin notches. N^{55e11}/+, baz^4/+ and baz^{815-8}/+ heterozygous flies have no

proximal wing margin defects. p<0.0001 (Fisher's exact test) for phenotypic differences between

both double heterozygous combinations and N^{55e11}/+.

(H) baz^{815-8}/+ heterozygosity suppresses the sev-$NICD$/+ gain-of-function eye phenotype. sev-

$NICD$ induces several phenotypes, and most prominently a transformation of the R1/R6 outer

photoreceptors to R7s, which results in ommatidia classified as 4:3 (4 outer R-cells and 3 R7s)

or 5:2 (5 outer R-cells and 2 R7s), in contrast to wt, which has a 6:1 appearance (6 outer R-cells

and 1 R7). Note that heterozygosity of baz suppresses the effect of sev-NICD and reverts the

ommatidial appearance partially towards 6:1 (the wt appearance).

Figure 3. Notch target gene expression is affected in *baz* mutant cells.

(A-C") Confocal images of third instar wing discs with expression of Notch target genes: *wg-lacZ*

(detected with anti-bGal, red; monochrome in A', B', and C') and *cut* (anti-Cut, blue;

monochrome in A", B", and C") are shown in wild type (A-A"), *baz-IR-1* (B-B"), and *baz-IR-2* (C-

C") genotypes. *wg-lacZ* and Cut expression are both reduced within the *enGal4>IR* expression

domain (posterior compartment, marked by GFP in green). Genotypes: (A) *UAS-dcr2; enGal4,*

UAS-GFP / wg-lacZ. (B) *UAS-dcr2; enGal4, UAS-GFP / wg-lacZ; baz-IR-1* (*BL31522)/+.* (C)

UAS-dcr2; enGal4, UAS-GFP / baz-IR-2 (BL31523)/+ (all at 29°C).

(D) Quantification of intensity ratio of Cut expression between anterior and posterior

compartments (see Materials and Methods for details). Note reduction with both independent

baz-IR transgenes (*p*-values 0.0003 and 0.0001 (student t-test), respectively). (E) Quantification

of intensity ratio of *wg-lacZ* expression (anti-bgal staining) between anterior and posterior

compartments (*p*-values 0.002, and 0.001, respectively).

Figure 4. Clones mutant for *baz* display markedly reduced Notch target gene expression in eye

and wing discs.

(A-A") Confocal optical section of *baz$^{815-8}$* wing disc clone (positively labeled by co-expression of

RFP), and (B-B") cross-section of same *baz$^{815-8}$* clone at position indicated by yellow line in (A).

Cut expression is reduced in *baz* mutant cells. Yellow line traverses Cut expression domain in

two regions indicated by white and blue arrows. Cut levels in *baz$^{815-8}$* mutant cells (white arrow)

are reduced, as compared to heterozygous control cells (blue arrows). Genotype: *w, tub-gal80,*

hs-flp, FRT19A/ baz$^{815-8}$ FRT19A; act-Gal4, UAS-mCD8RFP /+. (C-C''') Projection of 3 optical

sections at clonal region in D/V border cells, showing Cut (green, monochrome in C''')

expressing cells. Cut is normally expressed in two rows, each flanking the D/V-boundary on the

respective side. Note that *baz$^{815-8}$* mutant cells (RFP+, in red) in the cells abutting the D/V-

boundary express Cut at markedly reduced levels or not at all (examples highlighted by white

arrows), nuclei are marked by Hoechst (blue in C, or turquoise in C"). (D) Quantification of Cut

expression levels of D/V border cells (measured by average pixel intensity for each cell, see

Materials and Methods for details), plotted to compare Cut levels between control and baz^{815-8}

mutant cells. Note higher Cut expression in neighboring wild-type cells, as compared to baz^{815-8}

mutant cells.

(E-E") Atonal (Ato) expression at and near the morphogenetic furrow (MF) in eye imaginal discs.

containing baz^{815-8} mutant clone (marked by absence of GFP expression, red, highlighted by

white outlines). Note markedly reduced Ato expression (green) in many mutant cells (orange

arrows in E', E"), although some remained unaffected (E' and E", white arrow). All cells,

including those with low Ato expression levels, stained equally well with nuclear marker

(Hoechst, blue in E, magenta in E'). (F-G") Two z-section views of baz^{815-8} mutant clone in

furrow region, with (F) scanned through the middle of Ato expression stripe. Note that many

baz^{815-8} mutant cells displayed markedly reduced or lost Ato expression (orange arrows in F")

with Hoechst (Hoechst, blue in F,G and magenta in F',G') staining serving as control. (G) z-

scan through posterior region of Ato expression stripe, where scan-plane intersected with the

apical domain (due to tissue shape of the MF and pseudo-stratified epithelium of imaginal disc)

reflecting junctional marker Patj (blue in G, magenta in G"). Note that apical junctional region

remains intact in baz^{815-8} mutant cells, indicating that apical-basal polarity was not affected.

Genotype: baz^{815-8} FRT9-2/hsflp ubi-GFP FRT9-2. (H) Quantification of Ato expressing cells in

baz^{815-} clones and neighboring wild-type control cells revealed a decrease in Ato positive cells in

baz^{815-} mutant clone vs wild-type control. Distributions were significantly different based on

Fisher's exact test.

Figure 5. Baz acts at the level of NICD or further downstream within Notch signaling.

(A-A') NDECD expressed via flip-out clones induces ectopic Wg expression within wing blade

precursor cells (Wg in red, monochrome in A'; clones labeled by co-expression of GFP in A).

Endogenous Wg expression stripe at DV-boundary serves as control (blue arrow in A') here and

in other genotypes, clones overlapping with endogenous Wg stripe were not scored in

quantification. Arrowheads mark examples of NDECD expressing clones (white in A, red in A'),

many of which induce Wg expression. Only clones in the wing pouch proper (pouch outlined by

green lines) were scored for Wg expression. Genotype: *hsflp/UAS-dcr2; act>y$^+$>Gal4, UAS-*

GFP/+; UAS-NDECD /+.

(B-B') Clonal co-expression of NDECD with *baz-IR-2* markedly reduced the number of Wg

expressing clones (Wg in red, monochrome in B'; clones labeled by co-expression of GFP).

Arrowheads mark examples of NDECD expressing clones (white in B, red in B'); note reduction

in such clones expressing Wg (compare to A,A') Genotype: *hsflp/ UAS-dcr2; act5c>y$^+$>Gal4,*

UAS-GFP/+; UAS-NDECD /baz-IR-2.

(C-C') NICD expressing clones induce ectopic Wg expression with high frequency (Wg in red,

monochrome in C'; clones labeled by GFP). Arrowheads highlight examples of NICD expressing

clones (white in C, red in C'). Genotype: *hsflp/UAS-dcr2: act5c>y$^+$>Gal4, UAS-GFP/+; UAS-*

NICD /+.

(D-D') *IR* control (*white*/*w-IR*) co-expressed in NICD in clones. Ectopic Wg expression was

induced at similar frequencies to NICD alone (Wg in red, monochrome in D'; clones labeled by

GFP). Arrowheads highlight examples of NICD expressing clones (white in D, red in D').

Genotype: *hsflp/UAS-dcr2: act5c>y$^+$>Gal4, UAS-GFP/ UAS-NICD /+*; *w-IR/+.*

(E-E') Clonal co-expression of NICD with *baz-IR-2* markedly reduced the number of Wg

expressing clones (Wg in red, monochrome in E'; clones labeled by GFP). Arrowheads highlight

examples of NICD expressing clones (white in E, red in E'). Genotype: *hsflp/ UAS-dcr2;*

act5c>y$^+$>Gal4, UAS-GFP/UAS-NICD; baz-IR-2/+.

(F) Quantification of NDECD clones with induced Wg expression with or without *baz-IR(s)*. The

frequency of Wg activation in >*NDECD* clones was markedly reduced by the presence of *baz-*

IR-1 or baz-IR-2 (*p*=0.0008 or *p*=0.011, respectively). Only clones inside wing pouch region

were counted, due to *wg* expression being regulated by other pathways outside pouch region.

(G) Quantification of NICD expressing clones with induced Wg expression with genotypes as

indicated (with *w-IR* (control), *baz-IR-1*, and *baz-IR-2*). Wg expression frequency in NICD clones

was markedly reduced when *baz-IR-1 or baz-IR-2* were co-expressed (both *p*<0.001 comparing

to NICD alone or NICD with *w-IR*; as calculated by Fisher's exact test).

Figure 6. Baz is detected in the same protein complex as NICD.

Baz (myc-tagged) or myc-Hippo (Hpo, negative control) and HA-tagged NICD were co-

transfected into S2 cells. Lysates of transfected cells were immnunoprecipitated (IP-ed) with

anti-myc antibody.

(A) Upper panel: Western blot showing NICD-HA detected in lysates of cells co-transfected with

myc-Hpo or Baz-myc. Lower panel: Anti-myc co-IP experiments, note NICD-HA was co-IPed

with Baz-myc, whereas it was not IP-ed by myc-Hpo (specificity control).

(B) Both Baz-myc and myc-Hpo were efficiently pulled down with anti-Myc in cell lysates co-

transfected with NICD-HA.

Figure 7. *baz* and *numb* act antagonistically to each other, but at different levels within the

Notch pathway

(A-F) Wings expressing *baz RNAi* in the posterior compartment in wild-type (*baz-IR-1* in A,A'

and *baz-IR-2* in D,D') causing wing margin bristle cell loss (also compare to Figure 1) or the

equivalent *baz-IRs* in a *numb* heterozygous background, *numb^{15}/+* (in B,B' and E,E',

respectively). Quantifications, as bristle number, in (C) and (F); wild-type posterior wing margin

has on average 230 bristle cells (Suppl. Figure S6). Note that removing a copy of *numb*

suppresses the effect of either *baz-RNAi*. Experiments were carried out at 29°C. Genotypes: (A,

A') *UAS-dcr2/+; enGal4/+; baz-IR-1/+*. (B,B') *UAS-dcr2/+; enGal4/numb^{15}; baz-IR-1/+, .* (D,D')

UAS-dcr2/+; enGal4/+; baz-IR-2/+. (E,E'') *UAS-dcr2/+; enGal4/ numb[15]; baz-IR-2/+,.* Note that

baz-IR-2 causes stronger defects, and that the phenotypic suppression caused by *numb[15]*

heterozygosity was significant (t-test) in both cases (C and F, respectively). "n" = number of

wings analyzed. A', B', D', and E' show higher magnification of the corresponding wings shown.

(G-I) Induction of Notch target gene expression by NICD expressing clones was not affected by

Numb co-overexpression. NICD was expressed via flip-on clones, as in Figure 5, with clones

labeled with GFP or numb-GFP co-expression. (G) Confocal optical projection with NICD and

GFP expressing clones (control). Note ectopic induction of the Notch target gene *cut*, visualized

via anti-Cut immunohistochemical staining (red and monochrome in G'), in many NICD clones.

Cut is normally expressed in a stripe at the dorsal/ventral boundary (arrow, in panel G).

Genotype: *hsFlp/+; act5c>y>Gal4, UAS-GFP/UAS-NICD*. (H) Projection of wing disc confocal

optical sections containing NICD expressing clones (marked in green by Numb-GFP co-

expression) with Numb-GFP overexpressed together with NICD. Note ectopic induction of *cut*

expression in many clones within the wing pouch. Genotype: *hsFlp/+; act5c>y>Gal4, UAS-*

numb-GFP/UAS-NICD. (I) Quantification of the percent of Cut expression positive clones in

NICD/GFP and NICD/numb-GFP expressing clones. Note similar ratio of expressing clones in

both genotypes with no significant difference (p = 0.702 based on Fisher's exact test; "n"

indicates the number of clones).

Figure 1:

enGal4, UAS-aPKC-IR (19°C)

Figure 2:

918
919
920

921 **Figure 3:**

924   **Figure 4:**

Figure 5:

act > NΔECD   act>NΔECD, >baz-IR-2   act > NICD   act >NICD, >w-IR   act >NICD, >baz-IR-2

930 **Figure 6:**

A

NICD-HA + myc-Hpo
NICD-HA + Baz-myc

lysate input
(10%)

Blot: Anti-HA

150 –
100 –
75 –
50 –

150 –
100 –
75 –

← NICD-HA

IP: Anti-Myc
Blot: Anti-HA

B

150 –
100 –
75 –

← Baz-myc

← myc-Hpo

IP: Anti-Myc
Blot: Anti-Myc

933 **Figure 7:**