Par3/Bazooka promotes Notch pathway target gene activation

Jun Wu, Neeta Bala Tannan, Linh T. Vuong, Yildiz Koca, Giovanna M. Collu,
and Marek Mlodzik

Affiliations

[1] Dept. of Cell, Developmental, and Regenerative Biology,

[2] Graduate School of Biomedical Sciences,

Icahn School of Medicine at Mount Sinai,

One Gustave L. Levy Place, New York, NY 10029.

Abstract

The conserved *bazooka (baz/par3*) gene acts as a key regulator of asymmetrical cell divisions

across the animal kingdom. The associated Par3/Baz-Par6-aPKC protein complex is also well

known for its conserved role in the establishment of apical/basal cell polarity in epithelial cells.

Here we define a novel, positive function of Baz/Par3 in the Notch pathway. Using *Drosophila*

wing and eye development, we demonstrate that Baz is needed for Notch-signaling activity and

in general for correct transcriptional activation of Notch target genes. Baz appears to act

independently of aPKC in these contexts, as knockdown of *aPKC* causes unrelated phenotypes.

Our genetic *in vivo* data suggest that Baz acts the level of active NICD, the cleaved Notch

intracellular domain, which upon pathway activation translocates to the nucleus and acts as a

transcriptional co-activator. We also demonstrate a biochemical interaction between NICD and

Baz. Taken together, our data define a novel role of the polarity protein Baz/Par3, as a positive

and direct regulator of Notch signaling through its interaction with NICD.

35   **Key Words**:

36   Notch signaling, Par3, transcriptional target regulation, cell polarity, Drosophila, tissue

37   patterning

Introduction

Cell polarity is a critical feature of most cell types and particularly so in epithelial cell sheets, which display a mandatory apical-basal (A/B) polarity that is required for their barrier function and vectorial secretion behavior. Similarly, stem cells often display polarized features, which are critical for stem cell maintenance and for progenitor cell fates. Here, when such polarized cells divide, the cell contents are asymmetrically inherited by the two daughter cells (e.g. rev. in [1, 2]). The mechanisms that govern the establishment and maintenance of cell polarity are evolutionarily conserved. The protein complexes that mediate polarization are also shared among distinct instances of cell polarization, for example between maintaining A/B polarity and asymmetric division of progenitor cells. Generally, the process is regulated by a set of protein complexes that were discovered genetically in *C. elegans* and *Drosophila* [3].

The conserved *par3* gene, known as *bazooka/baz* in *Drosophila*, was originally identified in *C. elegans*, where it acts as a key regulator of asymmetrical cell divisions in the very early embryo [4]. It generally acts together with Par6 and aPKC as a regulatory complex of many asymmetric cell divisions across the animal kingdom (reviewed in [1, 2]). This Par3/Baz-Par6-aPKC complex is also well known for its conserved role in the establishment of A/B polarity in epithelial cells [5]. Our interest in the A/B epithelial cell polarity proteins originates from the fact that several of these have also been linked to planar cell polarity (PCP), that is polarity within the epithelial sheet, orthogonal to the A/B axis [6], for example aPKC and Scribble [7, 8]. Both aPKC and Scribble can associate with core PCP factors, aPKC with Frizzled and Scribble with Van Gogh/Vang (Vangl in vertebrates), and they thus contribute to the stability and asymmetric segregation of the core PCP complexes and PCP establishment within an epithelial cell sheet [9]. To systematically examine whether additional A/B epithelial polarity factors can play a role in PCP establishment, we tested members of these protein families for PCP associated knockdown phenotypes during *Drosophila* wing development. Surprisingly when we examined

phenotypes of *baz/par3* knockdown in this context, we observed predominantly notching

features of the wing margin, a *Notch* mutant-like phenotype. Here we focus on investigating a

specific role for Baz in regulating Notch signaling.

Notch (N), originally identified in *Drosophila* based on its haplo-insufficient, dominant wing-

notching phenotype [10-12], encodes a single-pass transmembrane receptor protein containing

EGF-like repeats in its extracellular domain [13] and a conserved intracellular domain (NICD)

that can translocate to the nucleus and function as transcriptional co-activator (reviewed in [14-

18]). Many subsequent studies demonstrated that the highly conserved Notch receptor family

plays important roles during almost all biological processes through cell-cell communication and

cellular signaling, controlling many target genes that can be either shared or context dependent

(reviewed in [14-19]. Misregulation of N-signaling is associated with many human diseases,

including cancer [19]. The Notch receptor is activated by the binding of ligands of the Delta (Dl)

or Serrate/Jagged (Ser) families from neighboring cells to trigger two cleavage events to release

NICD [15, 16, 20]. NICD can then translocate into the nucleus, where it interacts with

transcription factors of the Suppressor of Hairless (Su(H)/CSL) family, and co-activators,

including Mastermind, to activate target gene expression [16]. Notch signaling is subject to

many regulatory steps, ranging from receptor and ligand trafficking and protein stability

regulators to specific nuclear events at the transcriptional end of pathway activation, many of

which were initially identified in *Drosophila* [16].

Previous work has proposed an indirect role for Baz/Par3 in regulating Notch signaling outcome

during asymmetric cell divisions in *Drosophila*. During asymmetric cell division of the sensory

organ precursors (SOPs), specific protein complexes segregate to opposite sides of the cell

cortex in each dividing SOP cell, on one side the Baz/Par3-Par6-aPKC complex and on the

other the Gαi-Pins-Mud-Dlg-Numb-Neur complex. The two daughter cells that result from this

cell division adopt distinct cell fates due to the asymmetric distribution of the above complexes:

with one daughter cell inheriting Baz/Par3-Par6-aPKC, and the other the Gαi-Pins-Mud-Dlg-

Numb-Neur complex [2]. Notably, the asymmetric segregation of these complexes causes an

asymmetric activation of Notch -signaling, as Numb is known to repress N-signaling by reducing

Notch levels at the cell surface, either through internalization or inhibition of recycling [2]. As

such, an indirect role of Baz/Par3 in the regulation of N-signaling has been proposed, acting

through the promotion of the asymmetric segregation of the inhibitory Numb away from the Baz-

containing daughter. However, due to this indirect role of Baz/Par3 in the pathway, this SOP

model system is not suited to address a more direct role of Baz/Par3 in Notch-signaling.

Using *Drosophila* wing and eye development, we define a novel, direct and positive function of

Baz/Par3 in Notch-signaling. We demonstrate that Baz is needed for full Notch-signaling activity

and activation of Notch transcriptional target genes. Baz appears to act independently of aPKC

in these contexts, as knockdown of *aPKC* causes PCP defects rather than Notch-associated

phenotypes. Our data indicate that Baz may act the level of NICD, the cleaved active Notch

fragment, which upon pathway activation translocates to the nucleus and acts as a

transcriptional co-activator, or further downstream. We also demonstrate a biochemical

interaction between NICD and Baz. Taken together, our data define a novel role of the polarity

protein Baz/Par3, as a positive regulator of Notch signaling through its interaction with NICD.

Results

Reduction of *bazooka* causes phenotypes similar to *Notch* loss of function phenotypes

To assess potential functions of A/B-polarity factors in PCP, or other signaling pathways, we

expressed two independent non-overlapping RNAi (IR) lines in the posterior compartment of

developing wing discs with the *en-Gal4* driver (Figure 1 and Suppl. Figure S1). Surprisingly, with

components of the Baz/Par3-Par6-aPKC complex we detected distinct outcomes. While knock-

down of aPKC showed classical PCP defects, as expected based on previous work [7], see also

below), both *baz-IR* lines caused wing margin loss in the posterior compartment (Figure 1A-C;

Baz protein levels were markedly reduced and barely detectable in the *baz-IR* expressing

domain: Figure 1E, and Suppl. Figure S1B-C). The loss of the wing margin is associated with a

reduction in Notch signaling or in effects on the Wg signaling pathway, as *wg* is a direct target of

Notch signaling along the dorsal/ventral compartment border (see also below); the gene name

Notch is associated with its haplo-insufficient wing notching phenotype (see above). Besides the

posterior compartment of the wing being slightly reduced, no other patterning defects were

observed in the *baz* knockdowns (Figure 1A-C). In contrast to *baz*, knockdown of either *aPKC*

or *par6* caused either cell lethality with loss of large areas of the posterior wing compartment at

25°C or 29°C. When *aPKC-IR* was expressed at reduced levels (driven at 18°C), causing a

milder knock-down, the posterior wing compartment was still markedly smaller (Figure 1D; note

that wing hairs were also often very short, Fig. 1D-D', orange arrows), but the wing margin

appeared largely normal (Figure 1B'-D'). PCP defects were also observed in the *aPKC-IR*

expressing domain (Figure 1D', green arrow indicating PCP hair orientation), consistent with the

previously identified role of aPKC in PCP [7]. *Par6-IR* mediated knock-down caused severe loss

of wing tissues even at 18°C, making it impossible to correlate these defects with a specific

cellular mechanism or signaling pathway (Suppl. Figure S1D-D'). Nevertheless, the wing

margin appeared relatively normal in areas where there was no tissue loss (Suppl. Figure S1D-

D', magenta arrows).

Taken together, knock down of *baz* caused a distinct loss-of-function (LOF) phenotype from

aPKC and *Par6*, affecting wing margin development. To confirm this effect, we also analyzed

wings containing mutant clones of *baz⁴* or *baz⁸¹⁵⁻⁸* alleles (Suppl. Figure S1A for allele

information). Consistent with the RNAi knock-downs, *baz* LOF clones also displayed wing

margin loss (Suppl. Figure S2A-C). On the notum, *baz* mutant LOF clones displayed loss of

sensory bristles (Suppl. Figure S2D-F), in accordance with previous work [21] confirming the

correct behavior of the *baz* alleles. Taken together with the RNAi phenotypes above, the *baz*

LOF clonal phenotypes in the wing suggest a potential role for Baz/Par3 in Notch-signaling.

***Notch* and *baz* interact genetically**

To corroborate a relationship between *Notch* and *baz* phenotypes, we performed genetic

interaction analyses. In heterozygous conditions of the null allele N^{55e11}/+, approx. 30% of wings

displayed distal margin notching (near vein 3 and 4; Figure 2A, B, and F). We compared these

wings to double heterozygous combinations with the LOF *baz* alleles: baz^4 and baz^{815-8}.

Strikingly, the double heterozygous animals displayed a marked enhancement of wing notching

with 68% and 79% of wing margin defects in N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} allelic combinations,

respectively (Figure 2C-D, and F; see also Suppl. Figure S3). Moreover, in addition to the

enhanced distal wing notch phenotype, a novel defect appeared in the double heterozygous

animals with wing notching in the proximal region of the posterior compartment (Figure 2B', D'

and quantified in 2G; see also Suppl. Figure S3B'-C'). Whereas N^{55e11}/+ wings did not display

any notches in this proximal region (Figure 2B' and 2G), 100% of the double heterozygous

N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} wings carried such margin defects (quantified in Figure 2G;

p<0.0001). The equivalent baz^4/+ and baz^{815-6}/+ heterozygous control wings did not display any

margin notching phenotypes (Figure 2E-E', also Suppl. Figure S3C-C'). These data suggest that

Notch and *baz* genetically interact during wing margin patterning in a synergistic manner.

We next asked whether interactions between *Notch* and *baz* are also detected in other tissues

and with a gain-of-function (GOF) N genotype. The specification of cell fates in the developing

eye requires Notch input at multiple steps, from early patterning of the presumptive eye field to

the specification of individual photoreceptor (R cell) and accessory cell fates. A classical

example of activation of N-signaling during eye patterning is the expression of active N

molecules under control of the *sevenless* (*sev*) enhancer (expressed in R3-4 and R1,6, and R7).

Elevated levels of Notch pathway activity can be induced by expressing N-terminally truncated

molecules that lack the inhibitory region in the extracellular domain that prevents γ-secretase-

mediated cleavage (NΔECD, [22]), or the intracellular domain itself, lacking any membrane

targeting sequence (NICD, [23]). Depending upon the strength of activation, elevated Notch

signaling induces the transformation of R3 to R4 or the R1/R6 photoreceptors to R7 [24, 25, 26,

27]. While the N-signaling requirement in R4 specification [24-26] cannot be used in this

context, as Baz is required to maintain high levels of Fz during PCP signaling and thus indirectly

affects R4 [7], the induction of R7 is a classical N-dependent cell fate and ectopic activation of N

causes the transformation of R1/R6 photoreceptors to R7 [27]. In the *sev-NICD* background a

large fraction of ommatidia display a multi-R7 phenotype (Figure 2H, and [27]), while removal of

one copy of the strong *baz$^{815-8}$* allele in this background suppresses the GOF Notch effect

(Figure 2H). Taken together with the above genetic interactions with *N* LOF features, these

results suggest that Baz/Par3 is generally required for Notch signaling function in different

tissues and both in LOF and GOF scenarios.

baz$^-$ loss of function caused reduction of Notch target gene expression

To test whether *baz-IR* affects Notch-signaling directly, we examined the expression of

endogenous Notch-signaling targets *wg* and *cut* [28, 29]. To monitor *wg* expression we

analyzed the expression of *wg-lacZ*, an enhancer trap insertion in the *wg* locus, which faithfully

reproduces *wg* expression patterns; and to monitor *cut* expression we used an antibody to Cut.

Knock-down of *baz* in the posterior compartment caused reduction of both *wg-lacZ* expression

and Cut protein levels, as compared to control levels in the anterior compartment (Figure 3A-

C"). Both *wg-lacZ* and Cut levels were significantly reduced in quantifications, by comparing

their intensity ratio between the anterior (control) and posterior (experimental) compartment

(Figure 3D-E). Since the reduction of *wg* and *cut* expression is detected at the third instar larval

stage in wing discs, before wing margin sensory development occurs at the pupal stages [30],

the *baz* effect on their expression precedes sensory cell differentiation and is thus clearly

separated in time from the known Baz function in asymmetric SOP cell division.

To confirm these results, we also examined the expression of the Notch response element

(NRE) reporter, *NRE-GFP* [31, 32]. In wing discs, *NRE-GFP* is expressed along the

dorsal/ventral compartment border in an indistinguishable pattern to endogenous Notch target

genes, as for example *wg* and *cut* [31] (Suppl. Figure S4A). Importantly, reduced expression of

NRE-GFP was observed in the posterior compartment in *enGal4>baz-IR(s)* wings (Suppl.

Figure S4B-C', and E), indicating that a reduction of *baz* directly correlated with reduced Notch-

signaling and target expression.

To corroborate these data and to test for a general effect of *baz* LOF on Notch signaling

outcomes in different tissues, we analyzed the expression of Notch targets in *baz* mutant clones.

Consistent with the *baz-IR* knockdown in wing discs, Cut expression levels were reduced in

baz$^{815-8}$ mutant cells (clones were positively labeled by RFP expression, Figure 4A-C, examples

highlighted with white arrows; quantified in Figure 4D). Next we tested for effects on Notch

signaling targets in eye discs. To test an independent Notch target gene, we asked whether

expression of *atonal* (*ato*), a N-signaling target during early eye patterning prior to R-cell

specification [33] also requires *baz* for its expression. Ato levels were reduced in *baz$^{815-8}$* mutant

cells (orange arrows in Figure 4E-E", F-F", mutant clone marked by lack of GFP). Importantly,

apical-basal polarity as apparent in optical z-sections was normal in *baz$^-$* clones (Figure 4G-G";

note Patj as A/B marker being localized normally in *baz$^{815-8}$* clones), whereas a reduction of Ato

expression was also easily detected in this optical plane (Figure 4D-F, orange arrows in E" and

F"; quantified in Figure 4H),

Taken together, these results are consistent with the notion that Baz functions as a positive

regulator within the Notch-signaling pathway, promoting the expression of Notch target genes.

Baz affects Notch-signaling at the NICD level

To mechanistically dissect Baz function in Notch signaling, we used N-terminal Notch

truncations in GOF assays to determine at what level within the pathway Baz was required.

First, we used N∆ECD[22], which lacks the extracellular domain and is constitutively active

independent of ligand binding. As such, if N∆ECD were affected by *baz* LOF, Baz would be

required downstream of the N-ligands and the subsequent extracellular cleavages of Notch, but

upstream of the γ-secretase-mediated cleavage [15, 20]. When N∆ECD was clonally expressed

[34, 35] in larval wing discs, 43% of N∆ECD clones induced the Notch target *wg* (Figure 5A, F;

as detected by Wg antibody).   When *baz-IR(*s) were co-expressed in N∆ECD clones, the

frequency of Wg expression, was reduced to 20% (*baz-IR*-1) and 27% (*baz-IR*-2) clones (Figure

5B, F). These data suggested that Baz was required at the level of the γ-secretase cleavage or

further downstream within the Notch-signaling pathway. Consistent with this hypothesis *baz-IR*

knock-down displayed no detectable effect on Notch protein levels at the membrane or cellular

distribution of Notch (Suppl. Figure S1E-F).

To further investigate the position of Baz within the Notch pathway, we tested the NICD

fragment, which is fully active and does not associate with the membrane or require any

cleavage and is thought to directly translocate to the nucleus to activate Notch-signaling target

genes as transcriptional co-activator [15, 16, 20, 23]. In control wing disc clones expressing

NICD, Wg expression was readily detected in 63% or 71% (with a control IR co-expression)

clones (Figure 5C-D, and G). In experimental clones, with *baz-IR*(s) being co-expressed with

NICD, the NICD effect was markedly suppressed: to 41% with *baz-IR*-1 and 32% with *baz-IR*-2,

respectively (Figure 5E, G). These data are consistent with the notion that Baz acts downstream

of membrane tethered Notch and that it is required for the ability of the NICD fragment to

activate Notch-signaling target gene expression.

The above *in vivo* studies suggested that Baz might either act in the nucleus or affect NICD

nuclear translocation. We therefore asked whether Baz and NICD physically interacted with

each other. This was tested biochemically via co-immunoprecipitation (co-IP) from S2 cell

samples that were co-transfected with HA-tagged NICD and either Baz-myc (full length Baz

tagged with myc) or myc-Hippo (Hpo, as negative control). Proteins extracted from S2 cell

samples were immunoprecipitated with anti-myc and analyzed by immunoblotting with anti-HA

antibody. NICD was readily co-immunoprecipitated with Baz-myc (Figure 6A), but not by the

Hpo control (Figure 6A-B; see figure legend for additional controls).

Taken together with the functional *in vivo* data, these results suggested that Baz and NICD were

associated in the same protein complex, and that the presence of Baz is required for normal

NICD function in signaling activation and Notch target gene expression.

Discussion

In this study we define a novel, direct, and positive role of the polarity protein Baz/Par3 in the

Notch pathway. This stands in contrast to its previously established role in Notch signaling

regulation during asymmetric SOP cell divisions, where Baz/Par3 affects N-signaling indirectly

by asymmetrically sequestering the Notch inhibitor Numb [2]. In the SOP scenario, the

Baz/Par3-Par6-aPKC protein complex localizes to one side within the dividing cell, and a Numb-

Neuralized containing complex localizes to the other side of the cell. As Numb inhibits Notch

signaling by regulating Notch-Sanpodo trafficking and thus reducing Notch levels at the cell

surface, this asymmetry causes Notch inhibition in the Numb containing daughter cell (reviewed

in [2]. In the SOP context, Baz/Par3 has been thus proposed to have a passive role in N-

signaling activation, via the asymmetric segregation of an inhibitor, Numb. Our data are

consistent with the notion that Baz/Par3 is also required positively for the Notch signaling

outcome and full transcriptional activation of Notch targets in *Drosophila* in many cell types.

Genetic epistasis studies and biochemical experiments indicate that Baz acts the level of

cleaved active Notch, NICD, or in parallel to it. Consistent with this notion Baz and NICD co-

immunoprecipitated, suggesting that Baz promotes Notch signaling at the level of NICD.

A novel Baz/Par3 function independent of the A/B polarity complex

Baz/Par3 is generally associated with Par6-aPKC at early to intermediate stages of A/B polarity

establishment, but later in mature epithelia it is not essential for their maintenance [36-38].

Indeed, the fact that we can recover *baz* LOF clones at all, whereas Par6 and aPKC are cell

lethal in null alleles, suggests that *baz* can behave differently to the 'core' A/B factors. In our

experimental set-up we observed no evidence for a Par6 or aPKC involvement in Notch

signaling. Our data suggest that Baz/Par3 regulates NICD independently of its role in the A/B

polarity complex. Similar to A/B polarity establishment, in the context of planar cell polarity

(PCP) patterning, Baz/Par3 functions together with aPKC to regulate Frizzled levels (and hence

the PCP outcomes) in the *Drosophila* eye [7], and thus, while Par6 could not be assayed in this

context, these data suggested that in PCP determination Baz is functioning with established A/B

polarity factors. As such, the discovery that in Notch signaling Baz/Par3 appears to function

completely independently of its partners in A/B establishment within the same tissue is

unexpected.

Par3 homologues have however been shown to act independently of the aPKC-Par6 complex in

several scenarios in mammalian cells, including Hippo/Yap signaling and during epithelial-

mesenchymal transition [39]. Par3 has even been suggested to act with Ku70 and Ku80 as part

of the DNA-dependent protein kinase complex to regulate double-strand breaks [40].

Why was Baz/Par3 not identified earlier as a component of Notch signaling?

The initial core Notch pathway genes were identified through classic genetic collections of

spontaneous mutants showing morphological defects in *Drosophila* and *C. elegans* [41-44].

Subsequent genetic screens and molecular studies have added many components of the Notch

pathway and many regulatory mechanisms that affect Notch signaling at all levels, ranging from

receptor and ligand trafficking and membrane stability and its membrane associated proteolytic

cleavages to nuclear co-factors and chromatin associated target gene regulation (reviewed in

[16-18]. In particular, many of the screens were performed using the notum Notch associated

phenotypes and score for asymmetric division-dependent SOP/bristle patterning defects. It has

thus been technically difficult, if not impossible, to separate potential additional Baz/Par3

involvements in Notch signaling. Similarly, classical loss-of-function screens in the *Drosophila*

embryo would have missed Baz/Par3 due to its initial requirement in epithelial polarization that

could have caused earlier lethality [45, 46], which obscures a potential role in subsequent

signaling events. Indirectly through our screen asking whether A/B polarity factors can also play

a role in PCP establishment or other pathways in the developing posterior wing disc

compartment we created a unique experimental scenario: first the posterior margin bristles are

not innervated SOPs and thus do not require N-signaling during asymmetric SOP divisions, and,

second, at this later stage of epithelial patterning Baz/Par3 is not required for maintenance of

apical/basal polarity. The combination of this thus allowed the identification of *baz* as a novel

factor that positively regulates Notch signaling outcomes.

Baz acts at the NICD level affecting expression of N-signaling targets positively

Genetic epistasis data, both in the wing and the eye, position the function of Baz/Par3 at the

level of NICD after it last cleavage, suggesting that Baz acts either in the nucleus together with

NICD or alternatively it might promote nuclear translocation or retention of NICD. The

biochemical immunoprecipitation results support the genetic data and are consistent with the

notion that Baz as a direct regulator of Notch signaling, acting at the level of NICD.

Interestingly, the human Baz/Par3 homologue, PARD3, has been suggested to act at a similar

level in the YAP/TAZ pathway. PARD3, which itself is found in the nucleus outside of the

junction-bound aPKC/Par6 complex, has been suggested to regulate TAZ phosphorylation and

its nuclear localization [39], as a scaffold for YAP-binding proteins. The Notch signaling pathway

is simpler than the pathway(s) that can activate YAP/TAZ proteins, as Notch acts as a

membrane-tethered transcriptional co-activator, without the complex cascade of kinase-

dependent steps as in YAP/TAZ regulation. Nonetheless, NICD is still regulated by

phosphorylation events, for example to limit prolonged pathway activity, through the action of

Mastermind and CycC:CDK8 [47]. It is possible that Baz/Par3 may also regulate NICD

phosphorylation and nuclear localization in a similar manner.

During Notch signaling, the associated DNA binding complex, centered around the Notch-

pathway dedicated DNA-binding factor CSL, Su(H) in *Drosophila*, switches from a repressor

complex to a transcriptional activator complex via NICD, which binds to Su(H) and also recruits

co-activators like Mastermind (Mam) [16]. The binding of NICD and Mam to Su(H)/CSL

displaces the repressing factors of the "repressor complex" and allows N-signaling target gene

transcription. Several factors have been identified that associate with Su(H)/CSL and NICD (see

reviews [16, 48]). Generally, the strength of activation of target gene expression depends on

NICD stability within the resulting activator complexes, which also contain histone

modifying/remodeling proteins, and accessibility of enhancers of any given target gene. As *baz*

mutations affect the levels of target gene expression and the Baz protein acts genetically at the

level of or in parallel to NICD, Baz/Par3 might be a factor participating in either scenario. N-

signaling acts in many different contexts and can activate many distinct target genes, and as

such the presence of context-determining factors will determine whether an enhancer is primed

to react to a specific signal, which may result in a gene having different responses towards the

same signals [49]. Our data suggest that Baz/Par3 acts rather generally on N-signaling target

gene activation, as we see effects on Notch target expression in multiple tissues and on

different target genes. The precise function of Baz/Par3 in the context of N-signaling target

activation will require transcription specific assays to be explored in future studies.

Materials and Methods

Fly genetics and stocks.

w-IR ($w^{JF01540}$, P{TRiP.JF01540}), baz-IR-1 ($baz^{JF01078}$, P{TRiP.JF01078}), baz-IR-2 ($baz^{JF01079}$, P{TRiP.JF01079}), $aPKC$-IR ($aPKC^{JF01966}$, P{TRiP. JF01966}), UAS-$mCD8RFP$, baz^4 $FRT9$-2 and baz^{815-8} $FRT9$-2 fly stocks were obtained from Bloomington Drosophila Stock Center. baz^{815-8} was recombined with FRT19A and then used for baz^{815-8} mutant clonal analysis with FRT19A MARCM stock (genotype: w, tub-$gal4$, hs-flp, $FRT19A$; act-$Gal4$, UAS-$mCD8RFP$ / CyO).

N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} were generated by intercrossing via several crosses: ubi-baz-$mcherry$ [50] was recombined with Sb^1 to generate ubi-baz-$mcherry$ Sb^1 flies. baz^4 (or baz^{815-8}) males with ubi-baz-$mcherry$ Sb^1 which rescued baz^- mutations (baz^4 or baz^{815-8}; ubi-baz-$mcherry$ Sb^1/+) were crossed to $N^{55e11}/FM7$ females to generate N^{55e11}/baz^4 or N^{55e11}/baz^{815-8} females (selected against Sb^1 phenotype).

Immunohistochemistry and microscopy:

The following antibodies were used in the study: guinea pig Anti-Baz 1:200 [51], rabbit anti-Dlg 1:3000-4000 [52], mouse anti-Wg 1:10-20 [53], mouse Anti-Cut 1:20-1:40 [54], rabbit anti-RFP 1:300 (Rockland Immunochemicals), rabbit anti-Galactosidase 1:2500 (Cappel). Mouse anti-Cut and anti-Wg antibodies were obtained from Developmental Studies Hybridoma Bank at the University of Iowa.

Adult wings and nota were imaged with conventional Zeiss Axioplan microscope. Immuno-fluorescent samples were imaged with Leica TCS SP5 confocal microscope. Images were projected and quantified with Fiji-Image J. Cut expression levels were measured in cell nuclei marked by Hoechst staining. baz^{815-8} mutant cells were outlined by RFP expression. Only first row of baz^{815-8} mutant cells right at D/V border were measured for "average Cut channel pixel intensity" minus "average background pixel intensity outside Cut expression domain" (similar to

Figure S4D1) for each nucleus. Neighboring control A/P border cells were also measured for

"average Cut pixel intensity" minus "average background pixel intensity".

Data visualization and Statistics:

Data were plotted via Microsoft Excel and LibreOffice Calc. Fisher's exact test (two tailed) and

student t-test (unpaired) were used for statistical analyses to calculate probabilities of null

hypothesis with the web version of Graphpad.

Cell Culture and co-immunoprecipitation

NICD-HA pAc and myc-Hpo pAc were generous gifts from Ramanuj DasGupta and Duojia Pan

[55]. Baz-myc pAWM was generated from pAWM gateway vector (Drosophila Genomics

Resource Center) and *baz* pEntr 2B (a gift from Tony Harris).

1μg of NICD-HA pAc and Baz-myc pAWM or myc-Hpo pAc plasmids each were transfected into

Drosophila S2 cells in a 6-well plate with Qiagen Effectene. Cells were lysed one day after

transfection with IP buffer (20 mM HEPES pH 7.5, 100 mM NaCl, 0.05% Triton X100, 1 mM

EDTA, 5 mM DTT, 10mM NaVO4, 10% glycerol and protease inhibitor cocktail). The cell

lysates (1mg of total protein) were pre-cleared by incubating with protein A-sepharose beads

(Thermo Scientific 20333) for 1 hour at 4°C, followed by centrifugation to remove the beads.

Mouse anti-Myc antibody (sc-40, Santa Cruz technology) was used at 1/1000 dilution for

immunoprecipitation. The A-sepharose beads and antibody were incubated together to form

pre-immuno-precipitate complexes at 4°C for 1 hour, then incubated with lysates. Myc-tagged

protein and associated proteins were captured by protein A-sepharose/antibody complexes after

overnight incubation at 4°C. Samples were washed 4 times with IP buffer. Immunoprecipitates

were resuspended in SDS sample buffer, boiled for 5 min, separated by SDS-PAGE, and

transferred to nitrocellulose membrane for immunoblotting. The blots were incubated with

mouse anti-myc (Santa Cruz technology) or rat anti-HA (Roche) at 1:1000, subsequently with

anti-mouse or anti-rat IgG-HRP. Proteins were then detected by enhanced chemiluminescence

kit (Millipore).

Acknowledgements

We are grateful to Kwang-Wook Choi, Kyung-ok Cho, Ramanuj DasGupta, Tony Harris, Duojia

Pan, and Jennifer Zallen for Drosophila strains and antibodies, and to Cathie Pfleger for helpful

suggestions on the manuscript. We also thank the Bloomington Drosophila Stock Center for fly

strains, and the Developmental Studies Hybridoma Bank at University of Iowa for antibodies.

We would also like to thank the ISMMS Microscopy CoRE, where confocal microscopy was

performed and was in part supported by the Tisch Cancer Institute P30 CA196521 grant from

the NCI. This work was supported by National Institutes of Health grants R01 EY013256 and

R35 GM127103 and (to MM).

References:

1.   Neumuller RA, Knoblich JA. Dividing cellular asymmetry: asymmetric cell division and its

implications for stem cells and cancer. Genes & development. 2009;23(23):2675-99. doi:

10.1101/gad.1850809. PubMed PMID: 19952104; PubMed Central PMCID: PMC2788323.

2.   Schweisguth F. Asymmetric cell division in the Drosophila bristle lineage: from the

polarization of sensory organ precursor cells to Notch-mediated binary fate decision. Wiley

Interdiscip Rev Dev Biol. 2015;4(3):299-309. doi: 10.1002/wdev.175. PubMed PMID: 25619594;

PubMed Central PMCID: PMC4671255.

3.   Goldstein B, Macara IG. The PAR proteins: fundamental players in animal cell

polarization. Developmental cell. 2007;13(5):609-22. doi: 10.1016/j.devcel.2007.10.007.

PubMed PMID: 17981131; PubMed Central PMCID: PMC2964935.

4.   Etemad-Moghadam B, Guo S, Kemphues KJ. Asymmetrically distributed PAR-3 protein

contributes to cell polarity and spindle alignment in early C. elegans embryos. Cell.

1995;83(5):743-52. Epub 1995/12/01. doi: 10.1016/0092-8674(95)90187-6. PubMed PMID:

8521491.

5.   Suzuki A, Ohno S. The PAR-aPKC system: lessons in polarity. Journal of cell science.

2006;119(Pt 6):979-87. doi: 10.1242/jcs.02898. PubMed PMID: 16525119.

6.   Humphries AC, Mlodzik M. From instruction to output: Wnt/PCP signaling in

development and cancer. Curr Opin Cell Biol. 2018;51:110-6. doi: 10.1016/j.ceb.2017.12.005.

PubMed PMID: 29289896; PubMed Central PMCID: PMC5949250.

7.   Djiane A, Yogev S, Mlodzik M. The apical determinants aPKC and dPatj regulate

Frizzled-dependent planar cell polarity in the Drosophila eye. Cell. 2005;121(4):621-31. PubMed

PMID: 15907474.

8.   Courbard JR, Djiane A, Wu J, Mlodzik M. The apical/basal-polarity determinant Scribble

cooperates with the PCP core factor Stbm/Vang and functions as one of its effectors.

Developmental biology. 2009;333(1):67-77. Epub 2009/07/01. doi: S0012-1606(09)00993-2 [pii]

10.1016/j.ydbio.2009.06.024. PubMed PMID: 19563796.

9.   Wang M, Marco P, Capra V, Kibar Z. Update on the Role of the Non-Canonical

Wnt/Planar Cell Polarity Pathway in Neural Tube Defects. Cells. 2019;8(10). doi:

10.3390/cells8101198. PubMed PMID: 31590237; PubMed Central PMCID: PMC6829399.

10.   Mohr OL. Character changes caused by mutation of an entire region of a chromosome in

Drosophila. Genetics. 1919;4:275-82.

11.   Southgate L, Sukalo M, Karountzos ASV, Taylor EJ, Collinson CS, Ruddy D, et al.

Haploinsufficiency of the NOTCH1 Receptor as a Cause of Adams-Oliver Syndrome With

Variable   Cardiac   Anomalies.   Circ   Cardiovasc   Genet.   2015;8(4):572-81.   doi:

10.1161/CIRCGENETICS.115.001086. PubMed PMID: 25963545; PubMed Central PMCID:

PMC4545518.

12.   Guruharsha KG, Kankel MW, Artavanis-Tsakonas S. The Notch signalling system:

recent insights into the complexity of a conserved pathway. Nature reviews. 2012;13(9):654-66.

doi: 10.1038/nrg3272. PubMed PMID: 22868267; PubMed Central PMCID: PMC4369923.

13.   Wharton KA, Johansen KM, Xu T, Artavanis-Tsakonas S. Nucleotide sequence from the

neurogenic locus notch implies a gene product that shares homology with proteins containing

EGF-like repeats. Cell. 1985;43(3 Pt 2):567-81. doi: 10.1016/0092-8674(85)90229-6. PubMed

PMID: 3935325.

14.   Wilson A, Radtke F. Multiple functions of Notch signaling in self-renewing organs and

cancer. FEBS letters. 2006;580(12):2860-8. doi: 10.1016/j.febslet.2006.03.024. PubMed PMID:

16574107.

15.   Andersson ER, Sandberg R, Lendahl U. Notch signaling: simplicity in design, versatility

in function. Development (Cambridge, England). 2011;138(17):3593-612. doi:

10.1242/dev.063610. PubMed PMID: 21828089.

16.   Bray SJ. Notch signalling in context. Nat Rev Mol Cell Biol. 2016;17(11):722-35. doi:

10.1038/nrm.2016.94. PubMed PMID: 27507209.

17.   Kopan R, Ilagan MX. The canonical Notch signaling pathway: unfolding the activation

mechanism. Cell. 2009;137(2):216-33. doi: 10.1016/j.cell.2009.03.045. PubMed PMID:

19379690; PubMed Central PMCID: PMC2827930.

18.   Henrique D, Schweisguth F. Mechanisms of Notch signaling: a simple logic deployed in

time and space. Development (Cambridge, England). 2019;146(3):dev172148 doi: 10.1242/dev.

doi: 10.1242/dev.172148. PubMed PMID: 30709911.

19.   Siebel C, Lendahl U. Notch Signaling in Development, Tissue Homeostasis, and

Disease. Physiol Rev. 2017;97(4):1235-94. doi: 10.1152/physrev.00005.2017. PubMed PMID:

28794168.

20.   Struhl G, Greenwald I. Presenilin is required for activity and nuclear access of Notch in

Drosophila. Nature. 1999;398(6727):522-5. doi: 10.1038/19091. PubMed PMID: 10206646.

21.   Roegiers F, Younger-Shepherd S, Jan LY, Jan YN. Bazooka is required for localization

of determinants and controlling proliferation in the sensory organ precursor cell lineage in

Drosophila. Proceedings of the National Academy of Sciences of the United States of America.

2001;98(25):14469-74. Epub 2001/12/06. doi: 10.1073/pnas.261555598. PubMed PMID:

11734647; PubMed Central PMCID: PMC64705.

22.   Fortini ME, Rebay I, Caron LA, Artavanis-Tsakonas S. An activated Notch receptor

blocks cell-fate commitment in the developing Drosophila eye. Nature. 1993;365(6446):555-7.

Epub 1993/10/07. doi: 10.1038/365555a0. PubMed PMID: 8413612.

23.   Kidd S, Lieber T, Young MW. Ligand-induced cleavage and regulation of nuclear entry of

Notch in Drosophila melanogaster embryos. Genes & development. 1998;12(23):3728-40. Epub

1998/12/16. doi: 10.1101/gad.12.23.3728. PubMed PMID: 9851979; PubMed Central PMCID:

PMCPMC317253.

24.   Cooper MT, Bray SJ. Frizzled regulation of Notch signalling polarizes cell fate in the

Drosophila eye. Nature. 1999;397(6719):526-30. doi: 10.1038/17395.

25.   Fanto M, Mlodzik M. Asymmetric Notch activation specifies photoreceptors R3 and R4

and planar polarity in the Drosophila eye. Nature. 1999;397(6719):523-6. Epub 1999/02/24. doi:

10.1038/17389. PubMed PMID: 10028968.

26.   Tomlinson A, Struhl G. Decoding vectorial information from a gradient: sequential roles

of the receptors Frizzled and Notch in establishing planar polarity in the Drosophila eye.

Development (Cambridge, England). 1999;126(24):5725-38.

27.   Tomlinson A, Struhl G. Delta/Notch and Boss/Sevenless signals act combinatorially to

specify the Drosophila R7 photoreceptor. Molecular cell. 2001;7(3):487-95. doi: 10.1016/S1097-

2765(01)00196-4.

28.   Neumann CJ, Cohen SM. A hierarchy of cross-regulation involving Notch, wingless,

vestigial and cut organizes the dorsal/ventral axis of the Drosophila wing. Development

(Cambridge, England). 1996;122(11):3477-85.

29.   Micchelli CA, Rulifson EJ, Blair SS. The function and regulation of cut expression on the

wing margin of Drosophila: Notch, Wingless and a dominant negative role for Delta and Serrate.

Development (Cambridge, England). 1997;124(8):1485-95. Epub 1997/04/01. PubMed PMID:

9108365.

30.   Jack J, Dorsett D, Delotto Y, Liu S. Expression of the cut locus in the Drosophila wing

margin is required for cell type specification and is regulated by a distant enhancer.

Development (Cambridge, England). 1991;113(3):735-47. PubMed PMID: 1821846.

31.   Saj A, Arziman Z, Stempfle D, van Belle W, Sauder U, Horn T, et al. A combined ex vivo

and in vivo RNAi screen for notch regulators in Drosophila reveals an extensive notch

interaction network. Developmental cell. 2010;18(5):862-76. doi: 10.1016/j.devcel.2010.03.013.

PubMed PMID: 20493818.

32.   Furriols M, Bray S. A model Notch response element detects Suppressor of Hairless-

dependent molecular switch. Curr Biol. 2001;11(1):60-4. Epub 2001/02/13. doi: 10.1016/s0960-

9822(00)00044-0. PubMed PMID: 11166182.

33.   Freeman ABaM. Notch signalling and the initiation of neural development in the

Drosophila eye. Development (Cambridge, England). 2001;128:3889-98.

34.   Ito K, Awano W, Suzuki K, Hiromi Y, Yamamoto D. The Drosophila mushroom body is a

quadruple structure of clonal units each of which contains a virtually identical set of neurones

and glial cells. Development (Cambridge, England). 1997;124(4):761-71.

35.   Pignoni F, Zipursky SL. Induction of Drosophila eye development by decapentaplegic.

Development (Cambridge, England). 1997;124(2):271-8.

36.   Morais-de-Sa E, Mirouse V, St Johnston D. aPKC phosphorylation of Bazooka defines

the apical/lateral border in Drosophila epithelial cells. Cell. 2010;141(3):509-23. doi:

10.1016/j.cell.2010.02.040. PubMed PMID: 20434988; PubMed Central PMCID: PMC2885938.

37.   St Johnston D, Sanson B. Epithelial polarity and morphogenesis. Curr Opin Cell Biol.

2011;23(5):540-6. doi: 10.1016/j.ceb.2011.07.005. PubMed PMID: 21807488.

38.   Tepass U. The apical polarity protein network in Drosophila epithelial cells: regulation of

polarity, junctions, morphogenesis, cell growth, and survival. Annual review of cell and

developmental biology. 2012;28:655-85. doi: 10.1146/annurev-cellbio-092910-154033. PubMed PMID: 22881460.

39.   Zhang P, Wang S, Wang S, Qiao J, Zhang L, Zhang Z, et al. Dual function of partitioning-defective 3 in the regulation of YAP phosphorylation and activation. Cell Discov. 2016;2:16021. Epub 2016/07/28. doi: 10.1038/celldisc.2016.21. PubMed PMID: 27462467; PubMed Central PMCID: PMCPMC4932730.

40.   Fang L, Wang Y, Du D, Yang G, Tak Kwok T, Kai Kong S, et al. Cell polarity protein Par3 complexes with DNA-PK via Ku70 and regulates DNA double-strand break repair. Cell research. 2007;17(2):100-16. Epub 2007/02/09. doi: 10.1038/sj.cr.7310145. PubMed PMID: 17287830.

41.   Lindsley DL, Zimm GG. The Genome of Drosophila melanogaster.1992.

42.   Greenwald I. lin-12, a nematode homeotic gene, is homologous to a set of mammalian proteins that includes epidermal growth factor. Cell. 1985;43(3 Pt 2):583-90. Epub 1985/12/01. doi: 10.1016/0092-8674(85)90230-2. PubMed PMID: 3000611.

43.   Fleming RJ, Scottgale TN, Diederich RJ, Artavanis-Tsakonas S. The gene Serrate encodes a putative EGF-like transmembrane protein essential for proper ectodermal development in Drosophila melanogaster. Genes & development. 1990;4(12A):2188-201. Epub 1990/12/01. doi: 10.1101/gad.4.12a.2188. PubMed PMID: 2125287.

44.   Kopczynski CC, Alton AK, Fechtel K, Kooh PJ, Muskavitch MA. Delta, a Drosophila neurogenic gene, is transcriptionally complex and encodes a protein related to blood coagulation factors and epidermal growth factor of vertebrates. Genes & development. 1988;2(12B):1723-35. Epub 1988/12/01. doi: 10.1101/gad.2.12b.1723. PubMed PMID: 3149249.

45.   Muller HA, Wieschaus E. armadillo, bazooka, and stardust are critical for early stages in

formation of the zonula adherens and maintenance of the polarized blastoderm epithelium in

Drosophila. The Journal of cell biology. 1996;134(1):149-63. doi: 10.1083/jcb.134.1.149.

PubMed PMID: 8698811; PubMed Central PMCID: PMC2120925.

46.   Harris TJ, Peifer M. Adherens junction-dependent and -independent steps in the

establishment of epithelial cell polarity in Drosophila. The Journal of cell biology.

2004;167(1):135-47. PubMed PMID: 15479740.

47.   Fryer CJ, White JB, Jones KA. Mastermind recruits CycC:CDK8 to phosphorylate the

Notch ICD and coordinate activation with turnover. Molecular cell. 2004;16(4):509-20. doi:

10.1016/j.molcel.2004.10.014. PubMed PMID: 15546612.

48.   Borggrefe T, Oswald F. The Notch signaling pathway: transcriptional regulation at Notch

target genes. Cell Mol Life Sci. 2009;66(10):1631-46. doi: 10.1007/s00018-009-8668-7. PubMed

PMID: 19165418.

49.   Housden BE, Terriente-Felix A, Bray SJ. Context-dependent enhancer selection confers

alternate modes of notch regulation on argos. Molecular and cellular biology. 2014;34(4):664-

72. Epub 2013/12/11. doi: 10.1128/mcb.01045-13. PubMed PMID: 24324007; PubMed Central

PMCID: PMCPMC3911485.

50.   Bosveld F, Bonnet I, Guirao B, Tlili S, Wang Z, Petitalot A, et al. Mechanical control of

morphogenesis by Fat/Dachsous/Four-jointed planar cell polarity pathway. Science (New York,

NY. 2012;336(6082):724-7. doi: 10.1126/science.1221071. PubMed PMID: 22499807.

51.   Blankenship JT, Fuller MT, Zallen JA. The Drosophila homolog of the Exo84 exocyst

subunit promotes apical epithelial identity. Journal of cell science. 2007;120(Pt 17):3099-110.

doi: 10.1242/jcs.004770. PubMed PMID: 17698923.

52.   Izaddoost S, Nam SC, Bhat MA, Bellen HJ, Choi KW. Drosophila Crumbs is a positional

cue in photoreceptor adherens junctions and rhabdomeres. Nature. 2002;416(6877):178-83.

doi: 10.1038/nature720. PubMed PMID: 11850624.

53.   Brook WJ, Cohen SM. Antagonistic interactions between wingless and decapentaplegic

responsible for dorsal-ventral pattern in the Drosophila Leg. Science (New York, NY.

1996;273(5280):1373-7. Epub 1996/09/06. PubMed PMID: 8703069.

54.   Blochlinger K, Bodmer R, Jan LY, Jan YN. Patterns of expression of cut, a protein

required for external sensory organ development in wild-type and cut mutant Drosophila

embryos. Genes & development. 1990;4(8):1322-31. doi: 10.1101/gad.4.8.1322. PubMed

PMID: 1977661.

55.   Wu S, Huang J, Dong J, Pan D. hippo Encodes a Ste-20 Family Protein Kinase that

Restricts Cell Proliferation and Promotes Apoptosis in Conjunction with salvador and warts.

Cell. 2003;114(4):445-56. doi: 10.1016/s0092-8674(03)00549-x.

56.   Blair SS. Genetic mosaic techniques for studying Drosophila development. Development

(Cambridge, England). 2003;130(21):5065-72. PubMed PMID: 12975340.

Figures

Figure 1. Reduction of *bazooka/baz function* causes wing margin bristle loss.

(A-D) Overview and high magnification view (A'-D') of wings of the indicated genotypes.

(A, A') Wild-type wing; note the associated margin bristles in high magnification of area of

posterior wing margin (A'). Cellular hairs point towards the distal margin (green arrow).

(B, B') *enGal4, UAS-baz-IR-1* wing, with *baz* knock-down in the posterior compartment; note

loss of margin bristles (magenta arrows). Cellular hair orientation was not significantly affected.

(Genotype: *UAS-dcr2; enGal4, UAS-RFP, NRE-GFP / UAS-Baz-IR-1 at 29°C*)

(C, C') *enGal4, UAS-baz-IR2* wing (independent RNAi, see Fig. S1), displaying very similar loss

of margin bristles phenotype with hair orientation not significantly affected. (Genotype: *UAS-*

dcr2; enGal4, UAS-RFP, NRE-GFP / UAS-Baz-IR-2 at 29°C)

(D, D') *enGal4, UAS-aPKC-IR* wing, displaying largely normal margin bristle development. Note

wing size reduction in posterior compartment with some areas displaying abnormally short wing

hairs (orange arrow) and other areas with cellular hair orientation defects, suggesting that

reduction of aPKC affected several aspects of wing development, yet had no effect on margin

bristle development (Genotype: *enGal4 /+; UAS-aPKC-IR (aPKC^{JF01966}) /+ at 19°C*)

(E-E") RNAi-based knock down of Baz protein in posterior compartment. All panels are showing

max-projections of a stack of confocal images (~40μm to cover the apical surface of all of the

wing pouch area), including composite (E) and individual channels. Anti-Dlg (Discs large, green

and E') staining, serves as positive control to outline cells in both anterior (labeled "A") and

posterior compartments (labeled "P"). Anti-Baz (magenta) levels were markedly reduced in the

posterior *baz-IR* expressing compartment, suggesting an effective knock-down via the *baz* RNAi

transgene. (Genotype: *dcr2; enGal4/+; /UAS-baz-IR-2 29°C*).

Figure 2. *Notch* and *baz* interact genetically during wing margin patterning.

(A-B) Heterozygous N^{55e11}/+ wings display wing margin notching phenotypes; approx. 2/3 of *N*/+

wings are normal with no margin loss (example in A), while 1/3 displayed distal margin notches

(B, arrow). Proximal posterior wing margin was always normal (boxed area in B, and B') in all

animals. Quantified in panels (F) and (G).

(C-D) N^{55e11}/baz^{815-8} double heterozygous wings. Note increased margin notching in distal area

(D, arrows) and proximal-posterior wing margin in all wings (C, D and D', arrows). (D') Higher

magnification of boxed area in D. Quantified in panels (F) and (G).

(E) baz^{815-8}/+ single heterozygous wings show no margin notching, neither distally nor in the

proximal-posterior region (E', boxed area from E shown at higher magnification).

(F) Quantification of distal wing margin phenotypes as percentage of wings containing notching

(genotypes as indicated). Note that N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} flies show enhanced defects

(higher percentage) of margin notching as compared to N^{55e11}/+ alone ($p=0.0062$ and $p=0.0033$,

respectively (Fisher's exact test). The control baz^4/+ and baz^{815-8}/+ single heterozygous flies

have no wing margin phenotypes.

(G) Quantification of proximal wing margin phenotypes (posterior compartment) as percentage

of wings with margin loss (genotypes as indicated). 100% of N^{55e11}/baz^4 and N^{55e11}/baz^{815-8} had

proximal wing margin notches. N^{55e11}/+, baz^4/+ and baz^{815-8}/+ heterozygous flies have no

proximal wing margin defects. $p<0.0001$ (Fisher's exact test) for phenotypic differences between

both double heterozygous combinations and N^{55e11}/+.

(H) baz^{815-8}/+ heterozygosity suppresses the *sev-NICD*/+ gain-of-function eye phenotype. *sev-*

NICD induces several phenotypes, and most prominently a transformation of the R1/R6 outer

photoreceptors to R7s, which results in ommatidia classified as 4:3 (4 outer R-cells and 3 R7s)

or 5:2 (5 outer R-cells and 2 R7s), in contrast to *wt*, which has a 6:1 appearance (6 outer R-cells

and 1 R7). Note that heterozygosity of baz suppresses the effect of sev-NICD and reverts the

ommatidial appearance partially towards 6:1 (the *wt* appearance).

Figure 3. Notch target gene expression is affected in *baz* mutant cells.

(A-C") Confocal images of third instar wing discs with expression of Notch target genes: *wg-lacZ*

(detected with anti-βGal, red; monochrome in A', B', and C') and *cut* (anti-Cut, blue;

monochrome in A", B", and C") are shown in wild type (A-A"), *baz-IR-1* (B-B"), and *baz-IR-2* (C-

C") genotypes. *wg-LacZ* and Cut expression are both reduced within the *enGal4>IR* expression

domain (posterior compartment, marked by GFP in green). Genotypes: (A) *UAS-dcr2; enGal4,*

UAS-GFP / wg-lacZ. (B) *UAS-dcr2; enGal4, UAS-GFP / wg-lacZ; baz-IR-1* (*BL31522)/+*. (C)

UAS-dcr2; enGal4, UAS-GFP / baz-IR-2 (BL31523)/+ (all at 29°C).

(D) Quantification of intensity ratio of Cut expression between anterior and posterior

compartments (see Materials and Methods for details). Note reduction with both independent

baz-IR transgenes: *p*-values 0.0003 and 0.0001 (student t-test), respectively. (E) Quantification

of intensity ratio of *wg-lacZ* expression (anti-βgal staining) between anterior and posterior

compartments (*p*-values 0.002, and 0.001, respectively).

Figure 4. (A-A") Confocal optical section of *baz$^{815-8}$* wing disc clone (positively labeled by co-

expression of RFP), and (B-B") cross-section of same *baz$^{815-8}$* clone at position indicated by

yellow line in (A). Cut expression is reduced in *baz* mutant cells. Yellow line traverses Cut

expression domain in two regions indicated by white and blue arrows. Cut levels in *baz$^{815-8}$*

mutant cells (white arrow) are reduced, as compared to heterozygous control cells (blue

arrows). Genotype: *w, tub-gal4, hs-flp, FRT19A/ baz$^{815-8}$ FRT19A; act-Gal4, UAS-mCD8RFP /+.*

(C-C"') Projection of 3 optical sections at clonal region in D/V border cells, showing Cut (green,

monochrome in C"') expressing cells. Cut is normally expressed in two rows, each flanking the

D/V-boundary on the respective side. Note that *baz$^{815-8}$* mutant cells (RFP+, in red) in the cells

abutting the D/V-boundary express Cut at markedly reduced levels or not at all (examples

highlighted by white arrows), nuclei are marked by Hoechst (blue in C, or turquoise in C"). (D)

Quantification of Cut expression levels of D/V border cells (measured by average pixel intensity

for each cell, see Materials and Methods for details), plotted to compare Cut levels between

control and *baz*[815-8] mutant cells. Note higher Cut expression in neighboring wild-type cells, as

compared to *baz*[815-8] mutant cells.

(E-E") Ato expression at and near the morphogenetic furrow (MF) in eye imaginal discs.

containing *baz*[815-8] mutant clone (marked by absence of GFP expression, red, highlighted by

white outlines). Note markedly reduced Ato expression (green) in many mutant cells (orange

arrows in E', E"), although some remained unaffected (E' and E", white arrow). All cells,

including those with low Ato expression levels, stained equally well with nuclear marker

(Hoechst, blue in E, magenta in E'). (F-G") Two z-section views of *baz*[815-8] mutant clone in

furrow region, with (F) scanned through the middle of Ato expression stripe. Note that many

baz[815-8] mutant cells displayed markedly reduced or lost Ato expression (orange arrows in F")

with Hoechst (Hoechst, blue in F,G and magenta in F',G') staining serving as control. (G) z-

scan through posterior region of Ato expression stripe, where scan-plane intersected with the

apical domain (due to tissue shape of the MF and pseudo-stratified epithelium of imaginal disc)

reflecting junctional marker Patj (blue in G, magenta in G"). Note that apical junctional region

remains intact in *baz*[815-8] mutant cells, indicating that apical-basal polarity was not affected.

Genotype: *baz*[815-8] *FRT9-2/hsflp ubi-GFP FRT9-2*. (H) Quantification of Ato expressing cells in

baz[815-] clones and neighboring wild-type control cells revealed a decrease in Ato positive cells in

baz[815-] mutant clone vs wild-type control. Distributions were significantly different based on

Fisher's exact test.

Figure 5. Baz acts at the level of NICD or further downstream within Notch-signaling pathway.

(A-A') NΔECD expressed via flip-out clones; it can induce ectopic Wg expression within wing

blade precursor cells (Wg in red, monochrome in A'; clones labeled by co-expression of GFP in

A). Endogenous Wg expression stripe at DV-boundary serves as control (blue arrow in A') here

and in other genotypes, clones overlapping with endogenous Wg stripe were not scored in

quantification. Arrowheads mark examples of NΔECD expressing clones (white in A, red in A'),

many of which induce Wg expression. Only clones in the wing pouch proper (pouch outlined by

green lines) were scored for Wg expression. Genotype: *hsflp/UAS-dcr2; act>y⁺>Gal4, UAS-*

GFP/+; UAS-NΔECD /+.

(B-B') Clonal co-expression of NΔECD with *baz-IR-2* markedly reduced the number of Wg

expressing clones (Wg in red, monochrome in B'; clones labeled by co-expression of GFP).

Arrowheads mark examples of NΔECD expressing clones (white in B, red in B'); note reduction

in such clones expressing Wg (compare to A,A') Genotype: *hsflp/ UAS-dcr2; act5c>y⁺>Gal4,*

UAS-GFP/+; UAS-NΔECD /baz-IR-2.

(C-C') NICD expressing clones induce ectopic Wg expression with high frequency (Wg in red,

monochrome in C'; clones labeled by GFP). Arrowheads highlight examples of NICD expressing

clones (white in C, red in C'). Genotype: *hsflp/UAS-dcr2: act5c>y⁺>Gal4, UAS-GFP/+; UAS-*

NICD /+.

(D-D') *IR* control (*white*/*w-IR*) co-expressed in NICD in clones. Ectopic Wg expression was

induced at similar frequencies to NICD alone (Wg in red, monochrome in D'; clones labeled by

GFP). Arrowheads highlight examples of NICD expressing clones (white in D, red in D').

Genotype: *hsflp/UAS-dcr2: act5c>y⁺>Gal4, UAS-GFP/ UAS-NICD /+; w-IR/+.*

(E-E') Clonal co-expression of NICD with *baz-IR-2* markedly reduced the number of Wg

expressing clones (Wg in red, monochrome in E'; clones labeled by GFP). Arrowheads highlight

examples of NICD expressing clones (white in E, red in E'). Genotype: *hsflp/ UAS-dcr2;*

act5c>y⁺>Gal4, UAS-GFP/UAS-NICD; baz-IR-2/+.

(F) Quantification of NΔECD clones with induced Wg expression with or without *baz-IR(s).* The

frequency of Wg activation in *>NΔECD* clones was markedly reduced by the presence of *baz-*

IR-1 or baz-IR-2 (*p*=0.0008 or *p*=0.011, respectively). Only clones inside wing pouch region

were counted, due to *wg* expression being regulated by other pathways outside pouch region.

(G) Quantification of NICD expressing clones with induced Wg expression with genotypes as

indicated (with *w-IR* (control), *baz-IR-1*, and *baz-IR-2*). Wg expression frequency in NICD clones

was markedly reduced when *baz-IR-1 or baz-IR-2* were co-expressed (both *p*<0.001 comparing

to NICD alone or NICD with *w-IR*; as calculated by Fisher's exact test).

Figure 6. Baz is detected in the same protein complex as NICD.

Baz (myc-tagged) or myc-Hippo (Hpo, negative control) and HA-tagged NICD were co-

transfected into S2 cells. Lysates of transfected cells were immnunoprecipitated (IP-ed) with

anti-myc antibody.

(A) Upper panel: Western blot showing NICD-HA detected in lysates of cells co-transfected with

myc-Hpo or Baz-myc. Lower panel: Anti-myc co-IP experiments, note NICD-HA was co-IPed

with Baz-myc, whereas it was not IP-ed by myc-Hpo (specificity control).

(B) Both Baz-myc and myc-Hpo were efficiently pulled down with anti-Myc in cell lysates co-

transfected with NICD-HA.

Figure 1:

Figure 2:

Figure 3:

757 **Figure 4:**

760 **Figure 5:**

763 **Figure 6:**