A high throughput screen with a clonogenic endpoint to identify radiation modulators of cancer

Nathan P. Gomes*[1], Barbara Frederick*[1], Jeremy R. Jacobsen[4], Doug Chapnick[4], Tin Tin Su[1,2,.3,5]

[1]SuviCa, Inc. PO Box 3131, Boulder CO, 80307-3131

[2] Department of Molecular, Cellular and Developmental Biology, University of Colorado Boulder, CO,

80309-0347

[3] Molecular and Cellular Oncology Program, University of Colorado Cancer Center, Aurora, CO 80045

[4]BioLoomics, 4665 Nautilus Ct. S., Ste. 501, Boulder, CO 80301

[5]Corresponding author tin.su@colorado.edu

*Equal Contribution

Running Title: A clonogenic HTS for radiation modulators

Abstract

Gomes, N. P., Frederick, B., Jacobsen, J. R., Chapnick. D. and Su, T. T. A high throughput screen with a clonogenic endpoint to identify radiation modulators of cancer. *Radiat. Res.*

Clonogenic assays evaluate the ability of single cells to proliferate and form colonies. This process approximates the regrowth and recurrence of tumors after treatment with radiation or chemotherapy, and thereby provides a drug discovery platform for compounds that block this process. However, because of their labor-intensive and cumbersome nature, adapting canonical clonogenic assays for high throughput screening (HTS) has been challenging. We overcame these barriers by developing an integrated system that automates cell- and liquid-handling, irradiation, dosimetry, drug administration, and incubation. Further, we developed a fluorescent live-cell based automated colony scoring methodology that identifies and counts colonies precisely based upon actual nuclei number rather than colony area, thereby eliminating errors in colony counts caused by radiation induced changes in colony morphology. We identified 13 cell lines from 7 cancer types, where radiation is a standard treatment module, that exhibit identical radiation and chemoradiation response regardless of well format and are amenable to miniaturization into small-well HTS formats. We performed pilot screens through a 1584 compound NCI Diversity Set library using two cell lines representing different cancer indications. Radiation modulators identified in the pilot screens were validated in traditional clonogenic assays, providing proof-of-concept for the screen. The integrated methodology, hereafter 'clonogenic HTS', exhibits excellent robustness (Z' values >0.5) and shows high reproducibility (>95%). We propose that clonogenic HTS we developed can function as a drug discovery platform to identify compounds that inhibit tumor regrowth following radiation therapy, to identify new efficacious pair-wise combinations of known oncologic therapies, or to identify novel modulators of approved therapies.

Introduction

More than half of US cancer patients undergo radiation therapy (RT) alone, or in combination with other oncologic therapies (*1*). Cancer recurrence following RT is common, resulting from proliferation of tumor clonogens that survived initial RT modalities (*2*). Identification of therapies that specifically inhibit clonogenic proliferation following radiation would be a significant advancement for RT-based cancer treatment, likely to reduce recurrence rates, and improve overall outcomes. Several therapeutic agents are approved for combinatorial use with RT for the treatment cancer, but none were initially identified as specific inhibitors of clonogenic growth following RT. Rather, mostly cytotoxic agents approved as discrete therapies are combined with RT empirically (*3*). It remains to be determined if direct screening for compounds that inhibit clonogenic proliferation can objectively identify novel and more potent radiation modulators. Advancements in cheminformatics, chemical synthesis, and improved ability to extract natural product compounds provide abundant and diverse chemical libraries from which to search. To systematically screen for inhibitors of clonogenic growth following RT, canonical clonogenic assays would have to be adapted to high-throughput screen (HTS) platforms, hereafter 'clonogenic HTS'.

The tradiational clonogenic assay is the only methodology that measures growth delay and total clonogenic death, whether it be it by mitotic catastrophe, apoptosis, necrosis, terminal differentiation, reproductive death, or other modes of cellular death (*4*). It is the assay of choice for monitoring celluar growth in response to RT (*5*). Relative to simpler MTT-type HTS assays, clongenic HTS assays, in principle, would be better suited for identifing novel radiation modulators, as well as more accurately testing known-drug/RT combinations. Resulting hits from this methodology should improve success in subsequent pre-clinical studies and accelerate translation into clinically relevant therapeutics. However, traditional clonogenic assays are extremely labor-, resource- and time-intensive. The traditional requirement to laboriously manually count a minimium of 50 well-seperated colonies with greater than 50 cells per colony, for each treatment condidtion has traditionally necessitated use of lager well formats, making the assay imcompatable with known HTS

methodolgies. To our knowledge no HTS system exists that follows the traditional clonogenic

methodology, despite description of the original assay 6 decades ago (*6*).

Several recent papers describe methodologies that approximate clonongic assays in HTS

formats by miniaturizing to small well formats and utilizing automated image analysis to score fixed

and stained colonies (*7-11*). The closest approximation of a clonogenic HTS among this body of work

is a 'High Content' screen of Non Small Cell Lung Cancer cells in 96-well plates (*10*). Cells were

seeded at non-clonogenic desities that yield somewhat similar survival and radiation response as in

traditional 6-well clonogenic assays. Cells were treated with radiation in the absence or presence of

serially diluted drugs, and "colonies" were counted after 4 days. Assay processing was as for the

conventional methodology: "colonies" were fixed, stained, destained, and scored by automatated

microscopy aided image analysis. A subset of hits from the screen of 146 compounds were validated

in conventional clonogenic assays, and xenograft models. While this work indicates that clongenic-like

HTS screens are possible, the assay does not have time/growth periods representative of true

clonogenic endpoints, and the protocol is as labor-intensive as conventional clonogenic assays,

including sample processing, and the necessity for multiple drug and radiation doses. The data

analysis methodology was identical to that of the conventional assay, and not amenable to large

datasets produced from an HTS. In a related study, short term (5-day) survival at a single radiation

dose and its modification by 18 drugs was assessed in 32 cancer cell lines (*12*). Survival was found to

correlate with dose-modifying factors in traditional clonogenic assays, suggesting that a simpler,

shorter-term assay in a high-throughput format may be used as the primary screen to identify potential

radiation modulators, later to be validated with traditional clonogenic assays. We again note that this

technology does not represent an HTS with a true clonogenic endpoint. A more recent paper reports a

High Content Screen (HCS) based on the growth of cancer cells in 2D and in 3D (spheroids) assays

(*8*). Colony growth of SW620 colorectal cancer cells was monitored following serial seeding dilutions

to achieve 1-100 colonies per well, in the presence or absence of drug. This method allows for the

testing of a small number of molecules at different plating densities, but which densities in 96-well

exhibits similar growth characteristics and drug-response as cells seeded as for traditional clonogenic

assays was not addressed.

The goal of HTS is to use automation to rapidly screen large numbers of compounds while

reducing time, reagents, and sample manipulation compared to conventional assays. Further,

automation inherently increases experimental consistency by reproducibly implementing standardized

protocols. Assay formats should be amenable to handling many-fold more compounds per unit time

than conventional assays. A survey of the literature indicates that most components of conventional

clonogenic assays have been successfully modified for use in HTS formats, and key technological

components for the assay are commercially available. However, a true clonogenic HTS remains

absent; its development required the functional combination of these known

methodologies/technolgoies, development of missing components, and integration into a discrete HTS

sytem, where scored hits validate in traditional assays. We have accomplished these tasks and

describe here an integrated high-throughput system with a clonogenic endpoint. Our system

integrates and automates cell- and liquid-handling, irradiation, dosimetry, drug-addition, and

incubation. We report a fluorescent live-cell based automated colony scoring methodology that

identifies and counts colonies based upon precise nuclei counting rather than colony area. We report

the identification of 13 cell lines, from 7 cancer types where radiation is a standard treatment module,

that exhibit equivalent radiation and chemoradiation response regardless of well format and are

amenable to miniaturization into small-well HTS formats. We performed pilot screens in two cell lines

representing two different cancer indications with a 1584 compound library, identifying radiation

modulators that validated in traditional clonogenic assays. Finally, our clonogenic HTS methodology

exhibits excellent robustness (Z'>0.5) and high reproducibility (>95%).

Results

Identification of cell lines amenable to miniaturized clonogenic growth.

Clonogenic assays are traditionally performed in 6-well plates or larger vessels, allowing for

growth of single cells into colonies without significant influence from neighboring clones. The large

growth area allows for well separated single cell seeding, while also achieving a minimum of 50

colonies per well, as is recommended in the literature (*13*). High throughput screens require assays to

be performed in smaller well formats to allow for many drugs to be screened simultaneously. To

develop a true clonogenic HTS, we set out to identify cell lines that: 1) could be grown in smaller well

formats, yielding well separated colonies that number >50/well over the traditional clonogenic growth

period of ~10 days and 2) exhibit equivalent radiation response between smaller well and traditional

6-well clonogenic formats. We focused on 7 cancer indications where radiation is a main treatment

modality: HNSCC, NSCLC, metastatic melanoma, glioma, soft tissue sarcoma, esophageal, and

rectal cancers. Not all cancer cell lines are expected to meet all two criteria. As shown in Figure 1A,

four different cell lines were seeded at clonogenic density in 6-well plates and allowed to grow for 10

days. Each colony contains >50 cells yet A2068 melanoma cells form large diffuse colonies while

SW837 rectal cancer cells form discrete compact colonies. An extensive literature search for

clonogenic images was performed to identified cell lines that display similar colony morphology to

SW837 and were commercially available with limited use restrictions. We obtained 2 to 5 cell lines for

each of the 7 cancer indications, seeded them at equivalent growth densities in 6-, 24- and 96-well

plates, and grew then over a period of 7-10 days to confirm the published colony morphology (Figure

1B). Of the four representative cell lines shown, the first two formed into well-separated colonies of

>50 per well in 24-well plates, while the final two did so in both 24- and 96-well plates. In general,

about half of the cell lines tested met this criterion in 24-well or smaller formats.

We next compared clonogenic radiation response between different well formats, using 3

doses of radiation as was done for validation of other high-content clonogenic assay formats (*10*).

Figure 1C-E shows representative data for two cell lines. Cells were seeded at identical density

(number of cells per growth area in the well) in the different well formats. In the traditional 6-well

clonogenic assay, cells are typically grown in 4-5 ml of media per well. This volume is equivalent to

1.7-2.2 ml per well in 24-well plates and 0.7-0.9 ml per well in 96-well plates. At these equivalent

media volumes, cells exhibit greater radiation resistance consistently in smaller well formats

compared to 6-well plates. For example, compare '6-well 4ml' (red arrow) to '24-well 1.5 ml' and '24-

well 2 ml' (bracket) in Figure 1C-D. Systematic analysis shows that radiation survival is exquisitely

sensitive to culture volume regardless of well format, with smaller relative volumes correlated with

increased radiation sensitivity. We note that this is not an artifact of smaller well formats, as increased

radiation sensitivity in smaller culture volumes is also seen in the traditional 6-well plates for all cell

lines examined (Figure 1C, compare 6-well 2, 3, and 4 ml). One possible explanation is RT-induced

production of secreted growth inhibitory factors that would be, by definition, more concentrated in

smaller volumes. Importantly, as this parameter is typically not reported, variable culture volumes

could partially explain lab-to-lab clonogenic variability widely observed in literature (*14*).

Regardless, consistent variation of radiation survival relative to culture volume allowed for the

identification of conditions where survival in smaller well formats are equivalent to that of traditional 6-

well formats. Specifically, cells grown in 4 ml in 6-well plates, 1 ml in 24-well plates and 150 μl in 96-

well plates produced equivalent radiation survival curves, with 'equivalent' defined as average survival

from technical replicates that fall within one standard deviation at each radiation dose tested (red

arrows in Figure 1C-E). Similar results were obtained for all cell lines tested (thirteen total), suggesting

that this phenomenon is specific to well format and seeding volume but independent of the cell line

used. More specifically, nine of thirteen cell lines showed equivalent radiation survival in 6-well and

24-well formats when culture volumes were controlled while the remaining four showed equivalent

radiation survival in 6-well, 24-well and 96-well formats when culture volumes were controlled (Table 1

and Supplemental Figure 1). We identified two cell lines for each of the seven cancer indications of

interest, except for glioma where only one of more than ten cell lines we examined was amenable to

miniaturization.

Confirmation of equivalent clonogenic chemo-radiation response regardless of plate format.

The goal of the clonogenic HTS system is to identify compounds that modulate radiation

responses. Therefore, it was necessary to confirm that cell lines exhibiting equivalent radiation

responses in miniaturized formats also exhibit equivalent chemo-radiation responses. This was

assessed using chemotherapeutics that represent standard-of-care for each of four cancer

indications, specifically, Cisplatin for HNSCC; 5-FU, Oxaliplatin and Irinotecan for rectal cancer;

Dacarbazine for metastatic melanoma; and Temozolomide for glioma. Using clinically relevant doses

for each chemotherapeutic and the same 3-dose radiation design as in Figure 1, dose-modifying

factors (DMF's) were calculated using linear-quadratic plots for clonogenic survival for each well

format, cell line and appropriate chemo-radiation combinations (*15*). Importantly, the miniaturized

clonogenic formats exhibit equivalent chemo-radiation responses, in addition to radiation responses

(Figure 2 and Supplemental Fig. 2; 'equivalent' defined as averages that are within one standard

deviation from each growth condition).

Automated colony scoring by nuclear position yields equivalent results as manual counting

The most laborious and time-consuming step in clonogenic assays is the counting of colonies,

with discrimination between true clones and those that experienced clonogenic death (below the 50-

cell/colony threshold). While typically done manually, recent publications describe several automated

counting methodologies. A common feature amongst these is the use of area 'masks' to identify

colonies that have at least 50 cells (Figure 3A, pink circle). While likely accurate under control

conditions, this methodology is error-prone in the context of irradiation as radiation can induce cell

spreading in some cell lines, thereby changing the area occupied by 50 cells between control and

irradiated samples. The extent of cellular spreading is cell type and radiation dose-dependent and can

increase the area of a 50-cell colony by as much as 50% (Figure 3B). While this problem may be

partially addressed by quantifying the area of a 50-cell colony at different radiation doses and using

masks of differing sizes, this methodology would be further confounded by morphological changes

induced by drug and/or drug-radiation combinations in a compound library screen. A robust HTS

system must accommodate these variables; therefore, we employed an alternative method.

We developed a nuclear fluorescence-based, automated colony counting method that utilizes

cell line specific parameters to segment colonies and individual nuclei from images of live cells. It is

comprised of three steps: image preprocessing to eliminate background and spurious fluorescence,

colony segmentation (Figure 3C), and nuclei segmentation to determine nuclear number for each

colony (see Methods for details). A delimited text file is generated containing each colony's relative

position within the well along with the number of nuclei in it. A 50-nuclei threshold then identifies

clonogens. Fluorescent nuclear dyes are amenable to this methodology but, we found that these have

poor reproducibility, low signal to noise ratios, and require stain/de-stain steps (not shown). Instead,

we generated stable nuclear RFP (Nuclight Red)-expressing variants for cell lines in Table 1 and

validated their use by fixing and staining as in traditional clonogenic assays after live fluorescence

images have been acquired (Figure 3C). This allowed us to compare directly automated to manual

counts for the same wells, which were found to be equivalent (Figure 3D,F). Survival curves were also

equivalent between parental and nuclear RFP variants, suggesting that stable transfection of nuclear

RFP did not alter radiation responses (Figure 3E,G). Similar results were observed for all cell lines in

Table 1 (Supplemental Figure 3) but we chose to use CCL-30 and SW837 in pilot screens because of

their robust growth, well-defined and compact clonogens, and amenability for 96-well miniaturization

(Figure 1A-B, D-E and Supplemental Fig. 2).

Automation of the clonogenic HTS screen

With cell lines amenable to clonogenic miniaturization identified and a validated methodology

for automated clonogenic counting in place, we next developed a customized instrument that

integrates automated cell- and liquid-handling, irradiation with dosimetry monitoring, and cell plate

incubation (Supplemental Figure 4A). Automated seeding of SW837 cells at clonogenic density and

radiation survival following RT is consistent across 96-well plates (Supplemental Figure 4B-C).

Clonogenic survival with varying drug treatment volumes up to 25 μl is consistent, providing significant

flexibility for HTS drug administration (Supplemental Fig 4D). Clonogenic survival is unaffected at up

to 0.75% DMSO (drug solvent), more than an order of magnitude greater than the concentration

expected to be used in the clonogenic HTS (Supplemental Fig 4E). Similar results were observed

when using CCL-30 cells (data not shown). Finally, radiation dosing as measured via robotically

placed nanoDot (Landauer) dosimetry is consistent amongst plates (for example, 4.118±0.096 Gy

received by 6 plates irradiated with 4 Gy).

As stated previously, CCL-30 and SW837 cells were chosen for pilot clonogenic HTS screens,

with the overall goal to identify compounds that modulate clonogenic survival following RT. Therefore,

for the primary screen radiation doses that induce partial growth inhibition (30-40%; 3 Gy for CCL-30,

9Gy for SW837) were chosen as this provides conditions for identifying modulators that increase or

decrease clonogenic radiation survival. With these conditions, we performed the clonogenic HTS in

each cell line with the 1584 compound Diversity Set VI library from the Developmental Therapeutics

Program of the National Cancer Institute. The Diversity Set library is assembled from a larger

(140,000) molecule library by grouping chemicals according to structure and selecting representative

members. Thus, we would be sampling many structurally diverse chemical families with this library.

Robustness of the clonogenic HTS screen

Cells were seeded in 150 μl, irradiated 24 hr post-seeding, and then treated to a final drug

concentration of 1 μM in total volume of 175 μl, with a final DMSO concentration of 0.01%. Cells were

seeded to average clonogenic densities of 66.0±7.7 (CCL30) and 110.2±13.2 (SW837) per well.

Each screen was conducted in seventeen 96-well plates, with a minimum of six vehicle (DMSO)

controls per plate. Live fluorescent images were obtained 10-days post-treatment with an IncuCyte

S3. Automated clonogenic segmentation was performed for each cell line in batch, and average

growth and standard deviations (Z-score) for each cell line were calculated (Figure 4A-B). To assess

the dynamic range of the primary screen, we examined the responses between controls and

compounds exhibiting significant growth effects. Vehicle (DMSO) wells function as negative controls,

while for positive controls we selected compounds that are 4 (CCL-30) or 4.5 (SW837) Z-scores

below the mean. Z'-factor was computed from the average (μ) and standard deviation (σ) of the

positive (+) and negative (-) controls, according to the formula $Z' = 1-[(3\sigma_{c+} + 3 \sigma_{c-})/|\mu_{c+} - \mu_{c-}|]$ exceed

0.5 in both cell lines (Figure 4C), indicative of an HTS with 'excellent' robustness and dynamic range

(*16*).

Reproducibility of the clonogenic HTS hits

A common threshold for calling hits in robust HTS screens are effects greater than 2 standard deviations from the mean (*17*). In our primary clonogenic HTS screen, 54 compounds in SW837 and 62 compounds in CCL-30 increased the efficacy of radiation by greater than two standard deviations (>2 Z-score), while no compounds for either cell line exhibited significant radiation protective or growth enhancement effects (Figure 4A-B, shaded areas). Importantly, 41 compound hits were shared between the two cell lines (Figure 5A). To access reproducibility of the identified shared hits, we retested each under the same conditions as the primary screen (1 μM drug plus 3Gy in CCL-30, or 9Gy in SW837). Since the reproducibility test does not include the rest of the library as a control, we assessed growth inhibition for both screens relative to vehicle (DMSO) controls. 39 of the 41 hits (95%) reproducibly exhibited growth inhibition greater than 2 standard deviations from the control in both cell lines (Figure 5B-C). The two compounds that did not reproduce were shared between the cell lines and exhibited no activity in the re-test. Based upon these data we conclude that the clonogenic HTS is highly reproducible.

A secondary screen differentiates radiation modulators from single agent activity alone

The clonogenic HTS screen was performed on irradiated cells, and it is therefore expected that some hits will result from robust single agent activity alone rather than radiation modulatory effects. Furthermore, radiation modulation could result in a simple sum of the drug and radiation effects (additive interaction) or greater than the simple sum (synergistic interaction). To identify true radiation modulators in the 39-compound hit pool, each compound was retested with and without radiation (3Gy for CCL-30, 9Gy for SW837) at varying concentrations (10 nM, 0.1 μM, and 1 μM). In SW837 cells, 6 compounds exhibited potent single agent activity, producing growth inhibition of greater than 90% at 10 nM drug (for example NSC330770 in Supplemental Figure 5A). 29 compounds exhibited what is expected if combination with radiation is additive (for example, NSC250594 in Supplemental Figure 5B). And 4 compounds exhibited greater than additive interaction with radiation, which is what is expected of true radiation modulators (Figure 6A). Two of

these compounds (NSC177365 and NSC228155) exhibited greater than additive activity at 1 μM, the

drug concentration used in the primary screen. The other two compounds exhibited complete single

agent growth inhibition at 1 μM, but partial single agent activity when reduced to 10 nM (NSC94600

and NSC317003). And at 10 nM both showed greater than additive effects when combined with

radiation. Similar results were obtained with CCL30 cells; three of 39 compounds showing greater

than additive effects with radiation. All three showed robust single agent activity at 1 μM, with radiation

modulation becoming apparent at 1-100 nM depending on the compound (Figure 6B and

Supplemental Table 1A). Of these hits, two are shared between both cell lines, resulting in a total of 5

potential radiation modulators identified (Supplemental Table 1B).

Validation of pilot HTS hits in traditional clonogenic assays

To this point, the data from the primary clonogenic HTS and confirmatory experiments were

obtained using compounds supplied as 10 mM DMSO stocks in multi-well plates from the NCI. Our

previous experience indicates that plated compounds in solution can exhibit reduced activity relative

to freshly acquired and prepared solutions of the same compound. Therefore, we acquired from the

NCI vialed solid aliquots of the five potential radiation modulators and prepared fresh DMSO stocks.

As expected, freshly prepared compounds show varying degrees of activity as single agents relative

to plated drug, ranging from similar (NSC94600, NSC317003, NSC149286) to 10-fold more active

(NSC177365, NSC228155) (Supplemental Figure 5C). Therefore, to test for radiation modulation, we

titrated each fresh compound to induce 25-30% growth inhibition. We then used fresh dilutions at

these respective concentrations to generate survival curves with a broad range of 9 radiation doses,

and calculated the resulting DMF for each hit. These experiments were performed first in 96-well

plates as with the clonogenic HTS (Supplemental Figure 6), and then validated with traditional 6-well

clonogenic assays (Figure 7).

In the 96-well format, three of four hits found in the SW837 screen recapitulated radiation

modification with DMF's >1.4 (Supplemental Figure 6A-C), with the exception being NSC228155 (not

shown, see Discussion). In CCL-30 cells, all three hits recapitulated (Supplemental Figure 6E-G).

Interestingly, NSC149286, a hit in CCL-30 screen but not SW837 screen, was found also to modulate

radiation in SW837 when tested with 9 radiation doses; a possible explanation for this result is

provided in the Discussion. More important, all compounds that exhibited radiation dose modification

in miniaturized formats exhibited nearly identical modification in traditional formats, with DMFs ranging

from 1.375 to 1.581 in SW837 (Figure 7A-D) and from 1.220 to 1.411 in CCL-30 (Figure 7E-G). We

conclude that the clonogenic HTS methodology can robustly identify compounds that function as

radiation modulators in traditional clonogenic formats, and therefore can serve as a functional novel

drug discovery platform with a true clonogenic endpoint.

Discussion

A validated HTS screen with a clonogenic endpoint

We describe here an integrated system capable of performing HTS screens with a clonogenic

endpoint to identify radiation modulators. This system differs significantly from previously described

methodologies in key aspects: 1) while miniaturized, this is a true clonogenic assay wherein cells are

seeded at clonogenic densities and grown over traditional time periods, with traditional endpoints and

statistical practice; 2) the ability to perform clonogenic HTS in multiple cell lines for seven cancer

indications where RT is a standard treatment module; 3) automated colony scoring based upon

nuclei count rather than colony size masks, ensuring accurate colony counts regardless of

radiation/drug induced changes in colony morphology; and 4) integrated radiation dosimetry ensuring

consistency over HTS runs. Our data demonstrate that this clonogenic HTS system can identify

compounds that exhibit robust radiation Dose Modifying Factors (DMF) in traditional clonogenic

assays. Throughput is markedly increased due to miniaturization and automation of colony counting.

The significance of these innovations cannot be overstated: the traditional clonogenic assay is

laborious, including manual cell seeding, irradiation, drug treatment, fixing and staining, that ultimately

culminates in exceedingly tedious manual counting of colonies. Simple human limitations reduce

screening capability to no more than a dozen compounds per day. In contrast, our automated

clonogenic HTS is capable of processing over 2000 compounds a day, with throughput capacity that

can be scaled linearly with additional incubator space, off-the-shelf imaging units, and computer

processing.

Clonogenic assay harmonization

There is increasing recognition that disparate clonogenic assay protocols make comparing

data across research labs nearly impossible (*5, 18*). Worse, some agents identified as radiation

sensitizers appear not to be so, and vice versa depending upon the research group and assay

protocol used (*14*). The clonogenic HTS described here address several parameters of concern. We

assessed the role of plate format (well size) in relation to clonogenic survival and identified conditions

wherein clonogenic radiation survival is equivalent between miniaturized and traditional 6-well formats

(Figure 1). Our automated colony segmentation methodology controls for radiation induced variations

in cell and colony shape/size by assessing nuclear position/number rather than size masks (Figure 3).

The resulting positional identification of actual nuclear counts allows for the accurate quantification of

colonies greater than 50 cells, in agreement with the parameters of the original clonogenic assay

colony (*6*). We standardized protocols to achieve a minimum colony number of 50 per well, as

indicated by the literature (*13*), and exceed this parameter in the pilot screens (Figure 4C, the second

column). We validate candidate radiation modulators with a broad range of radiation doses (9 total),

allowing for monitoring of the initial "shoulder" and exponential portions of survival curves (Figure 7

and Supplemental Figure 6). We use DMF to quantify radiation modulation here but have developed

automated programs capable of computing additional parameters including, survival fraction at a

given radiation dose and α/β ratio from linear-quadratic plots. Finally, we address inter-person

variation by fully automating the clonogenic process, and conducting the pilot screens independently

with two researchers, in two different cell lines.

Radiation modulators identified in the pilot screen

The pilot screen and reproducibility tests identified five potential radiation modulators (Figure

5A): two shared between the SW837 and CCL-30 screens (NSC94600, NSC317003), two exclusive

to SW837 (NSC177365, NSC228115), and one exclusive to CCL-30 (NSC149286). Of these, only

NSC228155, an EGFR agonist (*19*), did not fully validate through subsequent experiments.

Specifically, radiation modulation was reproducibly observed with plated drug (Figure 6A), however

fresh compound exhibited 10-fold greater single agent activity (Supplemental Figure 5C) and showed

no modulation under any condition tested. The other four exhibited reproducible radiation modulation

in traditional 6-well clonogenic formats, regardless of variables between plated and fresh drug single

agent activity. Available data on these four compounds illustrate how our clonogenic HTS can identify

known as well as new radiation modulators.

One hit shared between the cell lines is Camptothecin (CPT, NSC94600) a well described

DNA intercalator and potent topoisomerase I inhibitor and a known radiation sensitizer (*20*). Clinically

relevant derivatives of Camptothecin include Irinotecan (Camptosar), a known radiation modulator

and standard of care for colorectal cancers which SW837 cells represent (*20, 21*). Our clonogenic

HTS was performed blind, using a library based on chemical structure and not for anti-cancer activity.

Therefore, identification of CPT is proof-of-concept that the screen is capable of accurately identifying

compounds that modulate radiation effects.

NSC317003, was shared between cell lines and is a structural analog of lucanthone and its

metabolite hycanthone (Supplemental Figure 8A). Lucanthone, a known DNA intercalator and

topoisomerase I/II inhibitor, was historically used to treat parasitic diseases, but more recently has

been studied as a radiation sensitizer (*22*). Lucanthone inhibits post-radiation DNA repair in human

cancer cells (*23*), accelerates the regression of brain metastases in patients receiving radiation

therapy (*24*), and has been in clinical trials as a radiation sensitizer for the treatment of Glioblastoma

(NCT01587144) and brain metastases of NSCLC (NCT02014545). Lucanthone and hycanthone are

not in Diversity Set VI but identification of a structural analog in our clonogenic HTS further validates

its ability to identify radiation modulators.

NSC149286 is an analog of curcumin (Supplemental Fig. 8B), which is has multiple reported

activities including radiosensitization of cancer cells and radio-protection of normal cells [(*25-27*);

reviewed in (*28*)]. NSC149286 exhibited slightly greater than additive effect with radiation in CCL-30

cells (Figure 6B) and an additive effect in SW837 cells (not shown). Fresh compound, however,

showed robust radiation modulation in both cell lines when tested with 7 radiation doses (Figure 7D

and G). The survival curves of SW837 suggest an explanation for the discrepancy between the

secondary screen and radiation dose range response data. Specifically, survival with or without drug

was equivalent at low doses of radiation, began to diverge at 9Gy, and increased significantly at

higher radiation doses (Figure 7D, Supplemental Fig. 6D). The level of divergence at 9Gy in the

secondary screen was considered to be within error and therefore scored as additive for SW837.

Greater divergence at >9Gy, however, resulted in a DMF>1. These data suggest that while use of a

single radiation dose in the pilot and secondary screens maximizes the number of compounds

assessed, we may miss modulators that show effect at other doses (false negatives). We believe this

tradeoff to be acceptable as compounds that modulate RT under many conditions are a preferred

outcome. While curcumin is not in the NCI Diversity Set VI library, identification of a closely related

analog further validates the ability of the clonogenic HTS to identify radiation modulators.

The pilot screens also identified a compound with known antineoplastic activity but previously

undescribed radiation modulation effects. This hit, specific to SW837 cells, was NSC177365 which

has been demonstrated to partially restore folding of the G-quadruplex in human telomerase reverse

transcriptase (hTERT) promoters, thereby silencing hTERT transcription and leading to cell death in

melanoma cells (*29*). Further, NSC177365 and related compounds have been demonstrated to inhibit

cystathionine β-synthase, reducing intratumor production of hydrogen sulfide which plays important

roles in cancer proliferation and migration [reviewed in (*30*)]. Finally, a patent for use of NSC177365

as a bone morphogenetic protein (BMP) signaling modulator has been granted (US-10434220-B2),

which has direct relation to cancer therapeutics [reviewed in (*31*)]. The ability of NSC177365 to

modulate radiation in SW837 cells significantly expands its potential role as a cancer therapeutic.

Interestingly, NSC177365 exhibited exquisitely robust single agent activity in CCL-30 cells even at the

lowest doses tested, to the point that no relevant compound/RT data could be obtained.

Standard Operating Procedure and looking forward

The streamlined clonogenic HTS procedure includes 4 critical steps: (1) a primary screen with a single LD_{30-40} radiation dose for the cell line and 1 μM of drug library; (2) the use of Z>2 from the population mean to call hits; (3) a secondary screen with 0 and LD_{30-40} Gy and each hit at 10 nM, 0.1 μM and 1 μM; and (4) validation in 6-well plates using fresh compound titrated to achieve 20-30% growth inhibition on its own. We believe this methodology is a major step forward for the unbiased identification of novel radiation modulators, but represents only the initial uses of this technology. Clonogenic HTS may be adapted to screen for compounds that inhibit regrowth of tumors following any standard therapies, as well as identify efficacious pair-wise combinations among existing therapeutic agents. For example, an IC_{50} dose of a single agent may be applied to all wells, much like single dose radiation, and used as the basis to screen for compounds that modulate that activity towards a clonogenic endpoint. Further adaptions to this methodology in development include 3D tumor growth, co-culture with somatic cells to recreate tumor microenvironments, and assays for immune-cell infiltration in the context of irradiation.

Materials and Methods

Cell culture

All cell lines were obtained from ATCC (American Type Culture Collection) except for FLO-1, A204 and KYSE270 which were obtained from Sigma-Aldrich. Cell line identities were confirmed by STR fingerprinting at the University of Colorado Cancer Center Cell Technologies Shared Resource. All cell lines were maintained in DMEM with 10% FBS supplemented with Penicillin (50 units/ mL) and Streptomycin (50 µg/mL).

Transduction with nucRFP lentivirus

Cells with low passage number (20 or fewer) were labelled with Nuclight Red (Sartorius) according to manufacturer's instructions. Briefly, cells were plated to 15-35% confluence and allowed to adhere overnight. Nuclight Red lentivirus was added at 1 multiplicity of infection (MOI)/cell along

with Polybrene (8 µg/mL). After 24 hours, media was replaced with fresh media containing 1 µg/mL

puromycin. Media with puromycin was relaced every 48 hours until stable clones were selected.

Plating efficiency, cell seeding and colony formation

Cells were stained with trypan blue (1:5 in PBS) and manually counted with a

hematocytometer. Cell populations with less than 10% trypan blue-stained (dead) cells were serially

diluted in media and plated in 24 or 96 well plates to determine plating efficiency (PE) and optimum

cell seeding for clonogenic assays. PE varied widely among the 13 cell lines (>90% for A375 to <16%

for KYSE270) and were consistent in additional assays for each cell line. Colony formation was

monitored daily over 7-10 days to determine an endpoint where colonies were still distinguishable.

Radiation survival curves of parental and nucRFP-labelled cells

Parental and nucRFP cells were plated in 6, 24 or 96 well formats to give equivalent colony

densities and irradiated 24 hours later. After 10 days, parental cells were fixed and stained with 0.1%

crystal violet or 0.4% Sulforhodamine B for manual counting whereas nucRFP derivatives were

imaged live using the IncuCyte S3 for colony segmentation.

High throughput screen

The SuviCa high throughput system consists of 4 functional instruments: a 37°C CO_2 incubator

(LiCONiC), a Fluent 780 drug distribution workstation (Tecan), a TEMPEST Liquid Handler

(FORMULATRIX), and a custom irradiator (Hopewell Designs) calibrated to deliver 3.2 Gy/min. Cells

were counted using trypan blue exclusion described above and plated according to previously

determined cell seeding (250 cells/well for CCL30, 200 cells for SW837) in 150 µL/well in 96-well

plates. To prevent edge effects of prolonged incubation, outside wells were filled with 200 µL sterile

H_2O. Cells were incubated overnight before irradiation, and drugs were added immediately after

irradiation.

The NCI Diversity Set VI is arrayed in 96 well plates at 10 mM in 20 μL DMSO. Upon arrival,

compounds were partitioned into duplicate plates at 5 μL per drug and stored at -20°C. For all assays,

a replicate plate was thawed only once and diluted 1:40 in complete media 1-2 hours before

irradiation. 1.4 μL of diluted compound was added to 200 μL of complete media. The Fluent was

programmed to add 25 μL/well to give a final concentration of 1 μM in a total volume of 175 μL/well.

Drugs were added after irradiation to avoid X-ray damage to the drugs. Plates were incubated for 10

days before imaging in IncuCyte S3.

Automated nuclear fluorescent based live-cell colony counting

This is comprised of three steps: image preprocessing, colony segmentation, and nuclei

segmentation. Parameters for each step are empirically optimized for cell line and growth format

specific conditions, then subject to images from a given experiment in batch. Image preprocessing

involves reducing spurious signal from culture well walls (Crop Perc.) and defining the maximum and

minimum signal values within the image to be analyzed (Thresh min/max). Colony segmentation

involves implementing Gaussian blur functions to merge proximal nuclear signals into discrete

colonies (Blur Factor), followed by iterated erode functions to further refine Gaussian colonies and

remove other spurious signals (erode iter.), finally implementation of colony prominence thresholds

and seeded watershedding functions define actual colonies from the pool of putative Gaussian

colonies (colony prominence). Cell segmentation involves similar methodologies, wherein for each

identified colony individual nuclear signals are defined by prominence thresholds and seeded

watershedding functions (cell prominence). Further parameters allow for previewing of defined

portions of a given well image, allowing for optimization of the parameters for image processing and

colony/cell segmentation. Colony segmentation output for each well is provided in delimited text

format (CSV) containing each clonogens relative position within the well, and the nuclei count for all

colonies. Nuclei threshold (>50) is then applied to identify clonogens. A JAVA applet employing this

methodology can be obtained here: download link (to be available upon manuscript acceptance).

Acknowledgements

This work was supported by Small Business Innovation Research phase I contract No.

HHSN261201700045C and Phase II contract No. 75N910196C00038 from the National Cancer

Institute to SuviCa (PIs: TTS and BF). Work in the Su lab is supported by National Institutes of Health

grant R35 GM130374 to TTS.

  Figure and Table Legends

Figure 1. Identification of cell lines that are amenable to clonogenic growth in a small well

format.

(A) Five cell lines that show the variation in colony size. Each was grown for 10 days to reach

colonies of 50 or more cells.

(B) Growth in 6, 24 and 96 well formats. 24 well images are shown at 2.3x magnification to

display the same final area as 6 well. 96 well growth is magnified 5.4x times to show equivalent area

as the 6 well format.

(C-E) A375 metastatic melanoma and SW837 rectal adenocarcinoma cells were seeded at the

same relative density (cells/surface area) but in different volumes per well in 6-, 24- or 96-well plates.

Cells were irradiated with 3, 6 or 9 Gy 24 hours after seeding. Clonogenic survival was quantified 10

days (SW837) or 7 days (A375) post radiation.

Figure 2. Chemoradiation responses in 6-, 24- and 96-well plates.

(A-D) CCL30 HNSCC and SW837 rectal adenocarcinoma cells were seeded at the same

relative density (cells/surface area) in 6-well (4 ml/ well), 24-well (1ml/well) and 96-well (150 µl/well)

plates and irradiated 24 hours after seeding and drugs were added immediately after irradiation.

Clonogenic survival was determined 10 days post radiation. Dose modifying factor (DMF) was

computed according to (*15*).

(E) Summary of additional known radiation sensitizers with SW837, A375 and LN-18 in

different plate formats.

Figure 3. Colony segmentation allows automated colony counting.

(A) A diagram to illustrate why a constant area mask cannot be used when cells spread after

irradiation.

(B) The area of a 50-cell colony increases with radiation for Cal27 HNSCC cells.

(C) The same well of a 6-well plate is shown stained for manual counting or imaged for nuclear

RFP. 'Colony segmentation' panel shows the RFP image after automated digital segmentation. Three

colonies are shown magnified to illustrate how the raw nuclear images are grouped or 'segmented'

into separate colonies based on the position of the nuclei.

(D-G) Manual counting and automated segmentation show similar results in clonogenic

survival after irradiation as do parental and RFP-expressing derivatives of CCL30 HNSCC and

SW837 rectal cancer cells.

Figure 4. Results of NCI Diversity Set VI screen for two cell lines.

(A-B) Data from the pilot screen through the Diversity Set of compounds using SW837 (A) and

CCL30 (B) cells. For both cell lines, the data for individual drugs are arranged along the x-axis in

same order that they occur on the plates of the drug library.

(C) Screen parameters and Z' table.

Figure 5. Results of NCI Diversity Set VI screen for two cell lines.

(A-B) Retest of 41 hits shared between the two cell lines from the Diversity set screen. Percent

(%) clonogenic survival relative to DMSO vehicle controls from the pilot screen (1st screen) are

plotted against the same values in the re-test (2nd screen) for SW837 (A) and CCL30 (B). Thirty nine

of 41 were within two STD from the vehicle average in both screens, demonstrating reproducibility.

Note that the data point at the coordinates 0,0 is comprised of multiple over-laid points; these are

wells with no growth in both screens.

(C) A summary of screening steps and the results.

Figure 6. A secondary screen differentiates radiation modulators from cytotoxic agents.

(A-B) Clonogenic survival after treatment with DMSO (control), radiation alone, drug alone or

the combination with drug with radiation in SW837 (A) and CCL30 (B) cells. Dashed lines show the

expected survival if the drug and radiation acted in an additive manner. Survival below the dashed

lines of combined treatments are likely caused by a synergistic interaction between the drug and

radiation. Control and radiation alone samples are an average of 6 replicate wells each. The others

are from one well per condition.

Figure 7. Radiation modulators found in the HTS were validated in 6-well clonogenic assays.

(A-D) Four compounds that showed greater than additive effect with radiation in the secondary

screen using SW837 cells were assessed in traditional 6-well clonogenic assays using 8 radiation

doses. Dose Modifying Factors indicate that four enhanced the effect of radiation. Drug

concentrations used with SW837 cells were 8 nM (NSC94600), 15 nM (NSC317003), 250 nM (NSC

177365) and 150 nM (NSC149286). Drug concentrations used with CCL-30 cells were 1 nM

(NSC94600), 10 nM (NSC317003) and 100 nM (NSC149286).

(E-G) Three compounds that showed greater than additive effect with radiation in the

secondary screen using CCL30 cells were assessed in traditional 6-well clonogenic assays using 7

radiation doses. Dose Modifying Factors indicate that three enhanced the effect of radiation.

Table 1. 13 cell lines that are amenable to miniaturization. Cancer type and available

information on mutation status and patient origin are shown. NSCLC = Non-Small Cell Lung Cancer;

HNSCC = Head and Neck Squamous Cell Carcinoma. The information is compiled from (*32-37*) and

the following online sources:

https://www.cbioportal.org/

https://www.atcc.org/

https://seandavi.github.io/UCCCReporter/reference/cell_line_ancestry.html

https://web.expasy.org/cellosaurus/

References

1.   Baskar R, Lee KA, Yeo R, Yeoh KW, Cancer and radiation therapy: current advances and future directions. Int J Med Sci 2012; 9, 193-9.

2.   Barker HE, Paget JT, Khan AA, Harrington KJ, The tumour microenvironment after radiotherapy: mechanisms of resistance and recurrence. Nat Rev Cancer 2015; 15, 409-25.

3.   Hennequin C, Guillerm S, Quero L, Combination of chemotherapy and radiotherapy: A thirty years evolution. Cancer Radiother 2019; 23, 662-65.

4.   Colevas AD, Brown JM, Hahn S, Mitchell J, Camphausen K, Coleman CN, et al., Development of investigational radiation modifiers. J Natl Cancer Inst 2003; 95, 646-51.

5.   Coleman CN, Higgins GS, Brown JM, Baumann M, Kirsch DG, Willers H, et al., Improving the Predictive Value of Preclinical Studies in Support of Radiotherapy Clinical Trials. Clin Cancer Res 2016; 22, 3138-47.

6.   Puck TT, Marcus PI, Action of x-rays on mammalian cells. J Exp Med 1956; 103, 653-66.

7.   Ye R, Qiao Y, Singh PK, Wang Y, He J, Li N, et al., High-Content Clonogenic Survival Screen to Identify Chemoradiation Sensitizers. Int J Radiat Oncol Biol Phys 2021; 111, e27-e37.

8.   Esquer H, Zhou Q, Abraham AD, LaBarbera DV, Advanced High-Content-Screening Applications of Clonogenicity in Cancer. SLAS Discov 2020; 25, 734-43.

9.   Georgantzoglou A, Merchant MJ, Jeynes JC, Mayhead N, Punia N, Butler RE, et al., Applications of High-Throughput Clonogenic Survival Assays in High-LET Particle Microbeams. Front Oncol 2015; 5, 305.

10.   Lin SH, Zhang J, Giri U, Stephan C, Sobieski M, Zhong L, et al., A high content clonogenic survival drug screen identifies mek inhibitors as potent radiation sensitizers for KRAS mutant non-small-cell lung cancer. J Thorac Oncol 2014; 9, 965-73.

11.   Katz D, Ito E, Lau KS, Mocanu JD, Bastianutto C, Schimmer AD, et al., Increased efficiency for performing colony formation assays in 96-well plates: novel applications to combination therapies and high-throughput screening. Biotechniques 2008; 44, ix-xiv.

12.   Liu Q, Wang M, Kern AM, Khaled S, Han J, Yeap BY, et al., Adapting a drug screening

platform to discover associations of molecular targeted radiosensitizers with genomic biomarkers. Mol

Cancer Res 2015; 13, 713-20.

13.   Munshi A, Hobbs M, Meyn RE, Clonogenic cell survival assay. Methods Mol Med 2005; 110,

21-8.

14.   Stone HB, Bernhard EJ, Coleman CN, Deye J, Capala J, Mitchell JB, et al., Preclinical Data on

Efficacy of 10 Drug-Radiation Combinations: Evaluations, Concerns, and Recommendations. Transl

Oncol 2016; 9, 46-56.

15.   Pike MC, Alper T, A Method for Determining Dose-Modification Factors. Br J Radiol 1964; 37,

458-62.

16.   Zhang JH, Chung TD, Oldenburg KR, A Simple Statistical Parameter for Use in Evaluation and

Validation of High Throughput Screening Assays. J Biomol Screen 1999; 4, 67-73.

17.   Malo N, Hanley JA, Cerquozzi S, Pelletier J, Nadon R, Statistical practice in high-throughput

screening data analysis. Nat Biotechnol 2006; 24, 167-75.

18.   Falls KC, Sharma RA, Lawrence YR, Amos RA, Advani SJ, Ahmed MM, et al., Radiation-Drug

Combinations to Improve Clinical Outcomes and Reduce Normal Tissue Toxicities: Current

Challenges and New Approaches: Report of the Symposium Held at the 63rd Annual Meeting of the

Radiation Research Society, 15-18 October 2017; Cancun, Mexico. Radiat Res 2018; 190, 350-60.

19.   Chen S, Yang L, Li Z, Zhuo S, Yan B, Zhang Z, et al., EGFR/EGFRvIII partly regulates the

tumourigenesis of glioblastoma through the SOX9-GLUT3 axis. Am J Transl Res 2021; 13, 6055-65.

20.   Rich TA, Kirichenko AV, Camptothecin radiation sensitization: mechanisms, schedules, and

timing. Oncology (Williston Park) 1998; 12, 114-20.

21.   Liu YQ, Li WQ, Morris-Natschke SL, Qian K, Yang L, Zhu GX, et al., Perspectives on

biologically active camptothecin derivatives. Med Res Rev 2015; 35, 753-89.

22.   Bases RE, Mendez F, Topoisomerase inhibition by lucanthone, an adjuvant in radiation

therapy. Int J Radiat Oncol Biol Phys 1997; 37, 1133-7.

23.   Bases R, Enhancement of x-ray damage in HeLa cells by exposure to lucanthone (Miracil D)

following radiation. Cancer Res 1970; 30, 2007-11.

24.   Del Rowe JD, Bello J, Mitnick R, Sood B, Filippi C, Moran J, et al., Accelerated regression of

brain metastases in patients receiving whole brain radiation and the topoisomerase II inhibitor,

lucanthone. Int J Radiat Oncol Biol Phys 1999; 43, 89-93.

25.   Walker BC, Mittal S, Antitumor Activity of Curcumin in Glioblastoma. Int J Mol Sci 2020; 21.

26.   Sak K, Radiosensitizing Potential of Curcumin in Different Cancer Models. Nutr Cancer 2020;

72, 1276-89.

27.   Momtazi-Borojeni AA, Ghasemi F, Hesari A, Majeed M, Caraglia M, Sahebkar A, Anti-Cancer

and Radio-Sensitizing Effects of Curcumin in Nasopharyngeal Carcinoma. Curr Pharm Des 2018; 24,

2121-28.

28.   Verma V, Relationship and interactions of curcumin with radiation therapy. World J Clin Oncol

2016; 7, 275-83.

29.   Kang HJ, Cui Y, Yin H, Scheid A, Hendricks WPD, Schmidt J, et al., A Pharmacological

Chaperone Molecule Induces Cancer Cell Death by Restoring Tertiary DNA Structures in Mutant

hTERT Promoters. J Am Chem Soc 2016; 138, 13673-92.

30.   Hellmich MR, Coletta C, Chao C, Szabo C, The therapeutic potential of cystathionine beta-

synthetase/hydrogen sulfide inhibition in cancer. Antioxid Redox Signal 2015; 22, 424-48.

31.   Bach DH, Park HJ, Lee SK, The Dual Role of Bone Morphogenetic Proteins in Cancer. Mol

Ther Oncolytics 2018; 8, 1-13.

32.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al., The cBio cancer

genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer

Discov 2012; 2, 401-4.

33.   Contino G, Eldridge MD, Secrier M, Bower L, Fels Elliott R, Weaver J, et al., Whole-genome

sequencing of nine esophageal adenocarcinoma cell lines. F1000Res 2016; 5, 1336.

34.   Dutil J, Chen Z, Monteiro AN, Teer JK, Eschrich SA, An Interactive Resource to Probe Genetic

Diversity and Estimated Ancestry in Cancer Cell Lines. Cancer Res 2019; 79, 1263-73.

35.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al., Integrative analysis of

complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal 2013; 6, pl1.

36.   Klijn C, Durinck S, Stawiski EW, Haverty PM, Jiang Z, Liu H, et al., A comprehensive

transcriptional portrait of human cancer cell lines. Nat Biotechnol 2015; 33, 306-12.

37.   Tratnjek L, Sibinovska N, Kralj S, Makovec D, Kristan K, Kreft ME, Standardization of

esophageal adenocarcinoma in vitro model and its applicability for model drug testing. Sci Rep 2021;

11, 6664.

Figure 1.

Figure 2.

Cell type	Cell line	Drug	Cell format	Ave DMF	Std Dev
Rectal Cancer	SW837	5-FU	6-well	1.379	0.153
			24-well	1.487	0.155
			96-well	1.494	0.133
		Irinotecan	6-well	2.222	0.574
			24-well	1.937	0.613
		Oxaliplatin	6-well	1.690	0.153
			24-well	1.523	0.187
Melanoma	A375	Dacarbazine	6-well	1.431	0.216
			24-well	1.624	0.384
Glioblastoma	LN-18	Temozolomide	6-well	0.921	0.168
			24-well	0.909	0.183

Figure 3.

A. Same cell # — 50 cell mask

B. The area of a 50-cell colony of Cal27 cells in 6-well plates (X10⁴ µm²)

	0 Gy	2 Gy	4 Gy	6 Gy
Avg	4.15	5.46	5.62	6.20
STD	1.29	0.95	0.82	1.34

C. Manual — Nuclear fluorescence — Colony Segmentation — Raw data — Digitally segmented

D. Manual vs. segmentation — CCL30 HNSCC

E. Parental vs. RFP — CCL30 HNSCC

F. SW837 Rectal Cancer

G. SW837 Rectal Cancer

Figure 4.

A.

B.

C.

Cell line	m_{c-} (AVE -ctrl)	s_{c-} (STD -ctrl)	m_{c+} (AVE +ctrl)	s_{c+} (STD +ctrl)	+ctrl used	$Z' = 1 - \dfrac{(3\sigma_{c+} + 3\sigma_{c-})}{|\mu_{c+} - \mu_{c-}|}$
CCL30	66.0	7.7	1.2	0.9	Z -4 or less	**0.60**
SW837	110.2	13.2	0.5	1.1	Z -4.5 or less	**0.61**

Figure 5.

A.

| 1584 compounds in Diversity Set IV |

SW837   CCL30

Primary Screen:
3 or 9 Gy, 1 μM drug

13   41   21   with Z<2

Reproducibility test:
3 or 9 Gy, 1 μM drug

39   39   of 41

Secondary Screen:
0 Gy and 3 or 9 Gy; 1nM, 0.1 μM, 1 μM drug

2   2   1

Validate using fresh compound in traditional 6-well clonogenics:
7 radiation doses, one drug concentration

✓**4 validated radiation modulators**

B.

SW837

C.

CCL30

Figure 6.

A.

SW837

B.

CCL30

Figure 7.

Table 1.

	Cancer type	Cell lines	Plate format	Mutations	Origin
1	NSCLC	NCI-H358	24 96	TP53 homozygous del, KRAS G12C	Caucasian male, metastatic site before chemotherapy
		NCI-H460	24	TP53 WT, CDKN2A del, KRAS G61H, PIK3CA 1633G>A	Male of European ancestry, Lung effusion
2	HNSCC (HPV⁻)	CCL30	24 96	STK11 A205T, BRD4-NUTM1 fusion	Caucasian male, nasal septum, metastatic pleural effusion
		Cal27	24	TP53 H193L, CDKN2A del E69*	Caucasian male, tongue
3	Esophageal cancer	FLO1	24	TP53 C277F	Caucasian male, primary oesophageal adenocarcinoma
		KYSE270	24	EGFR L861Q, TP53 X187_splice, CDKN2A del	Japanese male, primary oesophageal squamous cell
4	Soft tissue sarcoma	A204	24	Dip/tetraploid, EGFR L747_S752del, TP53 WT	Female of European ancestry, rhabdomyosarcoma, muscle
		SW872	24	Hyper-triploid, BRAF V600E, CDKNA R80*, TP53 I251N	Caucasian male, liposarcoma
5	Metastatic melanoma	A375	24	EGFR L858R D256Y, TERT AMP, TP53 WT	Female of European ancestry, malignant melanoma
		G-361	24	BRAF V600E, MET AMP, CDKN2A del	Caucasian male, malignant melanoma
6	Rectal cancer	SW837	24 96	TP53 R248W, KRAS G12C, APC R213* R1450*	Caucasian male, grade IV, adenocarcinoma
		SW1463	24 96	TP53 743G>A, APC V452fs*7, FBXW7 1436G>A, KRAS 34G>T	Caucasian female, adenocarcinoma
7	Glio-blastoma (only 1 fits criteria)	LN-18	24	TP53 C238S, PIK3CB E1051K, PTEN WT, CDKN2A del	Caucasian male, right temporal lobe