Research Article

Intranasal vaccination induced cross-protective secretory IgA antibodies against

SARS-CoV-2 variants with reducing the potential risk of lung eosinophilic

immunopathology

Takuya Hemmi[1, 7], Akira Ainai[1, 7*], Takao Hashiguchi[2, 5], Minoru Tobiume[1], Takayuki

Kanno[1], Naoko Iwata-Yoshikawa[1], Shun Iida[1], Yuko Sato[1], Sho Miyamoto[1], Akira Ueno[1,

6], Kaori Sano[1], Shinji Saito[1], Nozomi Shiwa-Sudo[1], Noriyo Nagata[1], Koji Tamura[7],

Ryosuke Suzuki[3, 7], Hideki Hasegawa[4], Tadaki Suzuki[1]

[1] Department of Pathology, National Institute of Infectious Diseases.

[2] Laboratory of Medical Virology, Institute for Life and Medical Sciences, Kyoto

University.

[3] Department of Virology II, National Institute of Infectious Diseases.

[4] Research Center for Influenza and Respiratory Virus, National Institute of Infectious

Diseases.

[5] Department of Virology, Faculty of Medicine, Kyushu University

[6] Department of Life Science and Medical Bioscience, Waseda University.

[7] Department of Biological Science and Technology, Tokyo University of Science.

* Correspondence to: Akira Ainai, PhD. Department of Pathology, National Institute of

Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo, 162-8640 Japan. Tel:

+81-3-4582-2701; Fax: +81-3-5285-1189; E-mail: ainai@niid.go.jp

Abstract

To control the coronavirus disease 2019 (COVID-19) pandemic, there is a need to

develop vaccines to prevent infection with severe acute respiratory syndrome coronavirus

2 (SARS-CoV-2) variants. One candidate is a nasal vaccine capable of inducing secretory

IgA antibodies in the mucosa of the upper respiratory tract, the initial site of infection.

However, regarding the development of COVID-19 vaccines, there is concern about the

potential risk of inducing lung eosinophilic immunopathology as a vaccine-associated

enhanced respiratory disease as a result of the T helper 2 (Th2)-dominant adaptive

immune response. In this study, we investigated the protective effect against virus

infection induced by intranasal vaccination of recombinant trimeric spike protein derived

from SARS-CoV-2 adjuvanted with CpG oligonucleotides, ODN2006, in mouse model.

The intranasal vaccine combined with ODN2006 successfully induced not only systemic

spike-specific IgG antibodies, but also secretory IgA antibodies in the nasal mucosa.

Secretory IgA antibodies showed high protective ability against SARS-CoV-2 variants

(Alpha, Beta and Gamma variants) compared to IgG antibodies in the serum. The nasal

vaccine of this formulation induced a high number of IFN-γ-secreting cells in the

draining cervical lymph nodes and a lower spike-specific IgG1/IgG2a ratio compared to

that of subcutaneous vaccination with alum as a typical Th2 adjuvant. These features are

consistent with the induction of the Th1 adaptive immune response. In addition, mice

intranasally vaccinated with ODN2006 showed less lung eosinophilic immunopathology

after viral challenge than mice subcutaneously vaccinated with alum adjuvant. Our

findings indicate that intranasal vaccine adjuvanted with ODN2006 could be a candidate

that can prevent the infection of antigenically different variant viruses, reducing the risk

of vaccine-associated enhanced respiratory disease.

Keywords: intranasal vaccination, secretory IgA antibody, COVID-19, SARS-CoV-2,

lung eosinophilic immunopathology

1. Introduction

The coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has spread worldwide since December 2019 and is presently a major public health concern [1, 2]. Vaccines could be the most promising approach to protect us from the threat of this infectious disease. In fact, several new vaccines have already been in use, showing high vaccine effectiveness [3-6]. Typical examples include the mRNA vaccines [3, 4] and the viral vector vaccines [5, 6]. Particularly in the UK, it is noteworthy that mRNA vaccine, BNT162b2 (Pfizer-BioNTech), was approved for practical use on December 2, 2020, less than a year after the epidemic began [7]. The advantage of the mRNA vaccine is that if the formulation of the vaccine has already been determined, a new vaccine can be launched for practical use in the shortest time, as long as the nucleic acid information of the target antigen is available. However, the long-term adverse events associated with the new modality of vaccines approved for emergency use are currently unknown; therefore, this issue should be adequately evaluated in the future. On the other hand, although the development of inactivated virus vaccines and subunit vaccines as conventional vaccine

formulations has been promoted, the development tends to be delayed because of time

taken to prepare the antigen. Unlike vaccines of new modality, these conventional

vaccines may have the advantage of being less burdensome to the vaccinees because

reactogenicity can be assumed to some extent.

Currently, most vaccines are administered intramuscularly or subcutaneously,

resulting in the induction of systemic antigen-specific IgG antibodies. In studies on nasal

influenza vaccines, we have already shown that a systemic antibody response is effective

for reducing mortality and morbidity associated with influenza but insufficient to prevent

infection, and that mucosal secretory IgA antibodies induced by intranasal vaccination

are highly protective against not only the vaccine-homologous virus but also

antigenically different viruses from the vaccine antigen [8-13]. The World Health

Organization (WHO) has recommended the development of a COVID-19 vaccine to

prevent infection with SARS-CoV-2 variants [14], therefore the study of vaccines

inducing mucosal immunity should be accelerated. In fact, 11 candidate intranasal

vaccines against SARS-CoV-2 have already been in clinical trials as of May 13, 2022 [15,

16].

However, there may be a risk of lung eosinophilic immunopathology caused by viral

infection among COVID-19 vaccinators, a phenomenon known as vaccine-associated

enhanced respiratory disease (VAERD) [17-23]. This phenomenon was first observed in

the 1960s in a clinical trial of formalin-inactivated respiratory syncytial virus (FI-RSV)

and measles vaccines. Two children died of severe pneumonia with eosinophilic

infiltration due to natural infection after vaccination in the clinical trial of the FI-RSV

vaccine [24]; thus, VAERD cannot be ignored in vaccine studies. Histological analysis of

postmortem lung sections revealed immune complex formation and complement

activation in the smaller airways. Similar eosinophilic immunopathology was observed in

a mouse experiment with SARS-CoV and Middle East respiratory syndrome

(MERS)-CoV vaccines. Some studies have suggested that lung eosinophilic

immunopathology is due to the induction of T helper 2 (Th2)-shifted immune responses

with high levels of antibody responses and insufficient neutralizing ability [21-23].

Although several studies on nasal COVID-19 vaccines in mouse models have already

been reported, these studies did not analyze vaccine-induced eosinophilic

immunopathology [25, 26]. Using a mouse model, we recently revealed that the

immunopathology of pneumonia with eosinophilic infiltration was induced by the

infection with SARS-CoV-2 among mice immunized with recombinant spike (S) protein

of the virus combined with alum adjuvant [23]. Thus, the U.S. Food and Drug

Administration (FDA) recommended that the balance of T-cell responses should be

properly assessed to avoid the risk of VAERD for vaccine candidates to be practically

used in the future[27].

In this study, we used a mouse model to evaluate both the protective effect and

neutralizing antibodies induced by intranasal administration of recombinant trimeric

spike protein derived from SARS-CoV-2 combined with synthetic oligodeoxynucleotides

containing the CpG motif, ODN2006 [28]. In addition, T-cell responses and the risk of

VAERD were examined after viral challenge in immunized mice. Our data showed that

intranasal administration of recombinant spike protein with ODN2006, rather than

subcutaneous administration in the presence of alum adjuvant, induced neutralizing

antibodies with well-balanced T-cell responses, resulting in the protection against

homologous or heterologous virus infection without lung eosinophilic immunopathology.

2. Materials and Methods

2.1. Purification of recombinant SARS-CoV-2 Spike protein

Recombinant trimeric ectodomain of SARS-CoV-2 S protein with two proline

mutations (rtS-ecto2P) as a vaccine antigen was produced using a *Drosophila* expression

system (Thermo Fisher Scientific, Grand Island, NY). The protein sequence was

modified to remove the furin cleavage site (RRAR to GSAG), and two stabilizing

mutations were introduced (K986P and V987P; wild-type numbering) [29, 30]. Protein

expression and purification were performed as previously described [31]. Recombinant

trimeric ectodomain of S protein with six proline mutations (rtS-ecto6P) for

enzyme-linked immunosorbent assay (ELISA) was produced using Expi293F cells

(Thermo Fisher Scientific). In addition to the mutation at the furin cleavage site, six

stabilizing mutations were introduced (F817P, A892P, A899P, A942P, K986P, and

V987P; wild-type numbering) [32].

2.2. Immunization and sampling

Female BALB/c mice (20-24 weeks old) (Japan SLC Inc., Hamamatsu, Shizuoka,

Japan) were maintained in specific pathogen-free facilities. The mice were either

intranasally or subcutaneously vaccinated with 3 μg rtS-ecto2P three times at 2-week

intervals. Intranasal vaccination was performed by instillation of 6 μL of vaccine solution

with or without 10 μg of ODN2006 into each nostril (total, 12 μL/mouse). Subcutaneous

vaccination was performed by inoculation with 100 μL of vaccine solution containing 10

μg of ODN2006 (InvivoGen, San Diego, CA) or 1 mg of Imject Alum adjuvant (Thermo

Fisher Scientific) in the dorsal part of the cervical region. ODN2006 and Imject Alum

were used as adjuvants to induce Th1- and Th2-dominant immune responses, respectively.

For the time-course evaluation of the antibody response, partial blood sampling from the

orbit was performed at 2-week intervals from the final vaccination. Serum, nasal and lung

wash, and cervical lymph nodes were collected from mice for the evaluation of antibody

and cellular immune responses, respectively, one week after the final vaccination.

All immunizations and partial blood sampling were performed under anesthesia. All

animal experiments were performed in accordance with the Guide for Animal

Experiments Performed at the National Institute of Infectious Diseases (NIID) and were

approved by the Animal Care and Use Committee of NIID.

2.3 Virus challenge and sampling

To evaluate effectiveness of protection against virus infection, vaccinated mice were

inoculated intranasally with the mouse adapted SARS-CoV-2 strain, QHmusX (GenBank

Accession No.: LC605054) [23], into the lungs (40 LD_{50} per mouse) and nasal cavity (6

LD_{50} per mouse) 2 weeks after the final immunization. To protect against SARS-CoV-2

variants—Alpha variant QHN001 (lineage B.1.1.7, GISAID: EPI_ISL_804007), Beta

variant TY8-612 (lineage B.1.351, GISAID: EPI_ISL_1123289), and Gamma variant

TY7-501 (lineage P.1, GISAID: EPI_ISL_833366)—were intranasally challenged with

3.5×10^5 $TCID_{50}$ into the lungs and nasal cavity. Intranasal challenge into the upper and

lower respiratory tracts was performed by instillation of 30 µL and 4 µL (2 µL in each

nostril), respectively. To determine the viral titer in the nasal mucosa and lungs, nasal and

lung washes were collected 3 days after virus challenge, and the lungs were collected for

the evaluation of eosinophilic immunopathology at 6 days post-infection. In addition,

body weight was monitored for 10 days after the virus challenge. The humane endpoint

was defined as the appearance of clinical diagnostic signs of respiratory stress, including

respiratory distress and > 25% weight loss. The SARS-CoV-2 challenge was performed

in a biosafety level 3 facility according to the Guidelines for Animal Experiments

performed at NIID.

2.4. Estimation of SARS-CoV-2 S-specific antibody responses

SARS-CoV-2 S-specific antibodies were estimated using ELISA. Half-area

flat-bottomed microtiter plates (Corning Inc., NY) were coated with 50 ng/well

rtS-ecto6P, followed by blocking with PBS containing 5% skim milk and 0.05% Tween

20. Serial dilutions of serum samples from vaccinated mice were added to each well of

microtiter plates. IgG antibodies were detected using biotin-conjugated goat anti-mouse

IgG antibody (Jackson Immunoresearch, West Grove, PA), followed by alkaline

phosphatase-conjugated streptavidin (Invitrogen, CA, USA). The enzymatic reaction was

initiated by the addition of the substrate p-nitrophenylphosphate (Sigma-Aldrich,

Burlington, MA). The absorbance at 405 nm was measured using an iMark microplate

reader (Bio-Rad, Hercules, CA). All procedures were performed at room temperature.

The S-specific IgG antibody titer was defined as the reciprocal of the highest dilution of

the test sample, giving a higher absorbance than the cut-off value obtained as 2-fold mean

absorbance of serial dilutions of control naive mouse serum set in each plate.

Quantification of S-specific IgG1 or IgG2a antibodies in the serum and IgA

antibodies in nasal or lung washes was performed as previously described [23]. Chimeric

human-mouse monoclonal IgG1, IgG2a, and IgA antibodies bearing variable regions of

the S-specific human monoclonal antibody S309 [33] were used as standard antibodies

for quantification. Horseradish peroxidase (HRP)-conjugated polyclonal anti-mouse

IgG1 antibody (Bethyl Laboratories, Montgomery, TX), anti-mouse IgG2a antibody

(Bethyl Laboratories), or polyclonal anti-mouse IgA antibody (Bethyl Laboratories) were

used as detection antibodies. The enzymatic reaction was obtained by adding ABTS

substrate (Roche, Basel, Switzerland), and the absorbance of 405 nm was measured.

2.5. SARS-CoV-2 neutralization assay

The neutralization assay was performed as previously described [23, 34]. Briefly,

50uL of QHmusX (100TCID50) and 50uL of heat-inactivated serum serially diluted by

two-fold were mixed and incubated in 96-well microtiter plates for 1h at 37□, followed

by the addition of 100 μL of VeroE6-TMPRSS2 cells (JCRB1819, Japanese Collection of

Research Bioresources Cell Bank) [35, 36]. After five days of cultivation, samples were

examined for viral cytopathic effects (CPEs). Neutralizing antibody titers were

determined as the reciprocal of the highest dilution rate at which no CPEs were observed.

The neutralization assay was performed in a biosafety level 3 laboratory at the NIID,

Japan.

2.6. Enzyme-linked immunospot assay

Cells secreting interferon (IFN)-γ, interleukin (IL)-4, or IL-5 were determined using

a mouse enzyme-linked immunospot (ELISpot) assay kit (Mabtech, Cincinnati, OH)

according to the manufacturer's instructions. Briefly, in plates pre-coated with

anti-mouse IFN-γ, IL-4, or IL-5 antibodies, 3×10^5 cells harvested from the spleen or

cervical lymph nodes were incubated for 16 h in the presence of a peptide pool derived

from the S protein of SARS-CoV-2 (a mixture of PepTivator SARS-CoV-2 Prot_S, S1,

and S+; Miltenyi Biotec, Bergisch Gladbach, Germany). After washing the cells with

PBS, biotin-conjugated anti-IFN-γ, IL-4, or IL-5 detection antibodies were added and

incubated at RT for 2 h, followed by incubation with ALP-conjugated streptavidin at RT

for 1 h. The enzymatic reaction was initiated by addition of BCIP/NBT. Each experiment

was performed in duplicate. Spots formed by cytokine-secreting cells were counted and

analyzed using ELISpot reader S6 Universal with ImmunoSpot 7.0 software (Cellular

Technology, Ltd., Shaker Heights, OH).

2.7. Flow cytometric analysis

T follicular helper (Tfh) cells, germinal center B (GCB) cells, and eosinophils were

evaluated by flow cytometry. Single cell suspensions were obtained from the cervical

lymph nodes, spleen and lungs of immunized mice. One million cells were stained with

FVD506 (Thermo Fisher Scientific) for dead cell removal and blocked with anti-mouse

CD16/CD32 monoclonal antibody (BD Pharmingen, San Jose, CA). Cell surface markers

of Tfh cells were defined as $CD4^+$ $CD8^-$ $PD-1^+$ $CXCR5^+$ among $TER119^-$ $Ly-6G/Ly-6C^-$

$CD11b^-$ $CD19^-$ populations, and those of GCB cells were defined as $CD19^+$ $GL7^+$ $CD95^+$

cells among $TER119^-$ $Ly-6G/Ly-6C^-$ $CD11b^-$ $CD3^-$ populations [37]. Eosinophils are

defined as $CD45^+$ $CD11b^+$ $CD11c^-$ $Ly6G^+$ $Siglec-F^-$ cells [38]. The antibodies used in

flow cytometric analysis are summarized in Supplementary Table 1. Samples were

analyzed with CantoII (BD Biosciences), and data were analyzed using FlowJo software

version 10.8.0 (Tree Star Inc., Ashland, OR).

2.8. Quantification of SARS-CoV-2 subgenomic RNA

Total RNA was extracted from 125 µL of nasal or lung wash using ISOGEN-LS

(Nippon gene, Toyko, Japan) and purified using a Maxwell RSC 48 Instrument (Promega,

Madison, WI) with a Maxwell RSC miRNA Plasma and Serum Kit (Promega).

Quantification of subgenomic RNA was performed by real-time reverse transcription

PCR (RT-PCR) using a QuantiTect Probe RT-PCR Kit (QIAGEN, Hilden, Germany) with

primers and probes as previously described [39]. Real-time RT-PCR was performed using

Mx3005P (Stratagene, La Jolla, CA, USA).

2.9. Immunohistochemistry

The lungs collected from mice were fixed in phosphate buffer containing 10% formalin.

The fixed lungs were embedded in paraffin and sectioned. Eosinophils were identified

using Astra Blue/Vital New Red staining (C.E.M. Stain Kit, Diagnostic Biosystems,

Pleasanton, CA). The lung tissue sections were observed for eosinophil infiltration in the

peribronchiolar areas using an optical microscope.

2.10. Statistical analysis

Data analysis and visualization were performed using GraphPad Prism 7.0 software

(GraphPad Software Inc., San Diego, CA). For statistical analysis, the Kruskal-Wallis test

with Dunn's multiple comparison test was used for comparisons between groups.

Comparison of body weight and survival was performed using Dunnett's multiple

comparisons test following the mixed-effects model or log-rank (Mantel-Cox) test,

respectively. Statistical significance was set at $P < 0.05$.

3. Results

3.1. Intranasal vaccines induced S-specific antibodies in nasal mucosa and lung as

well as serum

To evaluate the S-specific antibody responses, serum, nasal and lung wash

specimens were collected from mice that received three doses of either intranasal or

subcutaneous vaccines (Fig. 1A). Intranasal vaccination was performed with or without

ODN2006 as a mucosal adjuvant and subcutaneous vaccination was performed with

ODN2006 or alum adjuvant. Naïve mice were used as negative controls. Serum

S-specific IgG antibody titers and neutralization titers were determined using ELISA and

microneutralization assays, respectively. The concentration of S-specific IgA antibodies

in the nasal or lung wash samples was quantified using ELISA. Intranasal administration

without mucosal adjuvant induced low levels of serum S-specific IgG antibodies. In

contrast, intranasal vaccination adjuvanted with ODN2006 successfully induced

S-specific IgG antibodies in serum at a level similar to that induced by subcutaneous

vaccination in the presence of ODN2006 or alum adjuvant (Fig. 1B). The results for the

neutralizing antibody titer were similar to those of the S-specific IgG antibody titer (Fig.

1C). Although intranasal vaccination in the absence of mucosal adjuvant failed to induce

local antibody responses, nasal and lung S-specific IgA antibodies were detected among

intranasally immunized mice in the presence of ODN2006 (Fig. 1D and 1E). No secretory

IgA antibodies were detected in samples from subcutaneously vaccinated mice.

These results indicated that intranasal vaccination with ODN2006 as a mucosal

adjuvant induced not only secretory IgA antibodies in the nasal mucosa and lungs but also

systemic IgG antibodies at the same level as those obtained from subcutaneous

vaccination with ODN2006 or alum adjuvant.

283   3.2. Intranasal vaccine adjuvanted with ODN2006 effectively protect mice from

284   SARS-CoV-2 infection

To evaluate the protection against viral challenge, intranasal inoculation of the

mouse-adapted SARS-CoV-2 strain, QHmusX, was performed on mice vaccinated under

the same conditions as described above (Fig. 2A). Nasal and lung wash samples were

collected at three days post-infection (dpi), and the amount of subgenomic RNA (sgRNA)

derived from SARS-CoV-2 in these samples was evaluated by real-time RT-PCR to

assess the protection of mice against infection. The number of sgRNA copies in the nasal

wash was significantly reduced in mice with S-specific IgA in the nasal mucosa induced

by intranasal vaccination adjuvanted with ODN2006 (Fig. 1D and 2B). No significant

decrease in sgRNA in the nasal mucosa was observed in mice that received subcutaneous

vaccine or mice intranasally vaccinated with antigen only. A significant decrease in

sgRNA copies in the lung wash was observed in mice vaccinated intranasally or

subcutaneously in the presence of ODN2006 or alum, respectively, which showed high

neutralizing antibody titers in serum (Fig. 1C and 2C). Simultaneously, the mice were

monitored for body weight and survival for 10 days after the challenge (Fig. 2D and 2E).

Naïve mice and those intranasally vaccinated with antigen only died by 6 dpi. Among

mice that received subcutaneous vaccine with ODN2006 or alum, although few mice died,

others recovered from the apparent decrease in body weight post challenge and survived

until 10 dpi; in contrast, all mice intranasally vaccinated with ODN2006 survived without

remarkable body weight change during the observation period.

In addition, protection against viral challenge with SARS-CoV-2 variants (Alpha,

Beta and Gamma variants) was assessed (Fig. 3A). Significant reductions in nasal sgRNA

derived from each variant virus were achieved in mice intranasally vaccinated in the

presence of ODN2006 when compared to naïve mice (Fig. 3B-3D). In the lungs, sgRNA

of Alpha and Gamma variants significantly decreased by both intranasal and

subcutaneous vaccination with antigen together with ODN2006, while that of Beta

variant presented a significant decrease only by intranasal but not subcutaneous

vaccination (Fig. 3E-3G).

These findings indicate that IgA antibodies in the nasal mucosa and lung induced by

intranasal, but not subcutaneous, vaccination combined with ODN2006 were

cross-protective against SARS-CoV-2 variants.

3.3. Formation of germinal center and maintenance of antibody responses by

intranasal vaccination

The induction of memory B cells and long-lasting humoral immune responses

derived from long-lived plasma cells is important for successful vaccine-evoking

protective antibody responses [40]. The formation of a germinal center is required for the

affinity maturation of antibodies and determination of the B cell life span [37]. Hence,

vaccination-induced changes in the percentage of Tfh cells and GCB cells were evaluated

in lymphocytes from draining cervical lymph nodes using flow cytometry. The

proportions of both Tfh and GCB cells in cervical lymph nodes were definitely increased

by intranasal vaccination with antigen plus ODN2006 (Fig. 4A and 4B). In addition,

systemic S-specific IgG antibody responses were evaluated in serum samples collected

for 20 weeks at 2-week intervals after the initial vaccination (Fig. 4C). The highest

responses were achieved two weeks after the final vaccination, and mice that received

subcutaneous vaccine adjuvanted with alum or ODN2006 showed the highest S-specific

IgG antibodies, followed by mice intranasally vaccinated in the presence of ODN2006.

Although systemic S-specific IgG antibodies slightly declined over time during the

observation period, IgG antibodies were well held in mice subcutaneously vaccinated

with alum compared to mice that received intranasal or subcutaneous vaccination with

ODN2006 (Fig. 4D). At 20 weeks after the initial vaccination, S-specific nasal IgA

antibodies were detected in five out of six mice with intranasal vaccination combined

with ODN2006 but not in mice with other vaccinations (Fig. 4E).

Overall, these results suggest that intranasal vaccination with ODN2006 induced the

formation of GCs in the draining lymph nodes and the maintenance of local secretory IgA

antibody responses in the nasal mucosa, despite a slight decline in systemic IgG

responses.

3.4. Induction of Th1 response by using ODN2006 as an adjuvant

Lung eosinophilic immunopathology as a phenomenon of VAERD was observed in a clinical trial of FI-RSV and measles vaccine, and in a mouse model of SARS- or MERS-CoV vaccine. This phenomenon is suspected to be dependent on the Th2 dominant immune response of vaccine candidates [21-23]. Therefore, the Th response induced by intranasal vaccination with recombinant S protein in the presence of ODN2006 should be carefully examined. In mice, Th1 cells produce IFN-γ resulting in IgG2a induction, in contrast, Th2 cells produce IL-4 and IL-5 inducing IgG1 responses [41].

Cells isolated from the spleen and cervical lymph nodes one week after the final immunization were evaluated for cytokine production under stimulation of the peptide pool of S protein by ELISpot assay (Fig. 5A). Significant induction of IFN-γ-secreting cells was observed in the splenocytes of mice subcutaneously vaccinated with ODN2006 (Fig. 5B). In contrast, a significant induction of IL-4-secreting cells was observed in splenocytes obtained from mice subcutaneously vaccinated with alum adjuvant, and a similar tendency was observed in IL-5-secreting cells (Fig. 5C and 5D). In the case of

splenocytes, each cytokine-secreting cell significantly increased in mice that received

subcutaneous vaccination with ODN2006 or alum but not in those intranasally vaccinated.

Therefore, cytokine-secreting cells were evaluated using cells isolated from the draining

cervical lymph nodes of intranasally vaccinated or naive mice. Intranasal vaccination in

the presence of ODN2006 significantly increased the number of IFN-γ-secreting cells,

while IL-4 and IL-5 secreting cells decreased compared to intranasal vaccines with only

antigen (Fig. 5E). In contrast, intranasal vaccination with antigen in the absence of

mucosal adjuvant significantly induced IL-4 and IL-5 secreting cells, but not

IFN-γ-secreting cells, in draining lymph nodes (Fig 5F and 5G).

The dominant T cell response was estimated using an S-specific IgG antibody

subclass (Fig. 5H and 5I). Large amounts of S-specific IgG1 antibodies were obtained in

mice subcutaneously vaccinated with alum adjuvant compared to mice immunized in the

presence of ODN2006; however, S-specific IgG2a antibodies in mice intranasally or

subcutaneously vaccinated in the presence of ODN2006 were higher than those obtained

in mice vaccinated with alum. When the S-specific IgG1/IgG2a ratio was calculated, the

IgG1/IgG2a ratio after vaccination with ODN2006 was significantly lower than that after

vaccination with alum (Fig. 5J).

These results, obtained from the estimation of cytokine-secreting cells and

IgG1/IgG2a ratio, suggested that ODN2006, used as an adjuvant, induced Th1 dominant

immune responses regardless of the administration route.

3.5. Lung eosinophilic immunopathology is reduced by vaccines that induce a

remarkable Th1 response

The correlation between vaccine-induced Th2 dominant immune responses and lung

eosinophilic immunopathology was evaluated in a lethal challenge model in mice (Fig.

6A). S-specific IgG1/IgG2a ratios were calculated by ELISA in sera obtained one week

prior to challenge, and eosinophilic infiltration into the lungs was examined at 6 dpi by

histopathological and flow cytometric analyses. As shown in Fig. 6B, histological

analysis revealed that mice immunized in the presence of ODN2006, regardless of the

route of vaccination, showed small lesions with infiltration of inflammatory cells,

including neutrophils and mononuclear cells around the blood vessels and bronchi, but

little eosinophil infiltration. In contrast, eosinophilic infiltration around the bronchi and

blood vessels was observed in mice intranasally vaccinated with only antigen or

subcutaneously vaccinated with alum. Similar results were obtained in the flow

cytometric analysis. Although there were no significant differences in the percentages of

eosinophils induced among the different vaccines, these values correlated well with the

S-specific IgG1/IgG2a ratios (r = 0.779, p = 0.0015) (Fig. 6C and 6D).

Our results showed that Th2 dominant immune response, suspected by large values

of IgG1/IgG2a, caused lung eosinophilic immunopathology. In contrast, in immunization

combined with ODN2006, Th1 shifted immune responses correlating with low values of

IgG1/IgG2a ratio alleviated the risk of eosinophilic infiltration.

4. Discussion

In the current study, we revealed that intranasal vaccination with S protein together

with ODN2006, a toll-like receptor 9 agonist [28], could induce cross-protective

secretory IgA antibodies against SARS-CoV-2 variants in the nasal mucosa, which is the

initial site of infection. Furthermore, we demonstrated that this vaccine could reduce the

potential risk of lung eosinophilic immunopathology in the case of post-vaccination

infection.

In our previous studies on intranasal influenza vaccine, it has been revealed that

intranasal vaccination induces not only IgG antibodies in the serum but also

cross-protective secretory IgA antibodies on the surface of mucosal epithelial cells in the

upper respiratory tract [8-13]. Here, we evaluated serum and mucosal antibody responses

and protective effects induced in mice by intranasal vaccination with recombinant

SARS-CoV-2 S protein. Only mice immunized intranasally with antigen combined with

ODN2006 induced mucosal IgA as well as systemic IgG antibody responses

accompanied by a significant reduction in viral load in both the upper respiratory tract

and lungs. All individuals receiving this vaccine survived a lethal challenge with the

mouse adapted SARS-CoV-2 strain without significant weight loss. In addition, the

cross-protective ability by nasal vaccine was evaluated against SARS-CoV-2 Alpha, Beta

or Gamma variant. Results of neutralization assays using human sera collected from

mRNA vaccinees or individuals who suffered from breakthrough infections suggest that

the antigenicity of the Beta variant differs from those of the Alpha and Gamma variants.

[42-44]. When SARS-CoV-2 variants were challenged into the lungs, infections of Alpha

or Gamma variants were suppressed in mice intranasally or subcutaneously vaccinated,

whereas infection with the Beta variant could not be prevented by either vaccination. On

the other hand, all variants challenged into the nasal cavity were significantly prevented

in mice possessing mucosal secretory IgA antibody induced by intranasal vaccination, but

not in those subcutaneously vaccinated. These results indicate that, compared to systemic

IgG antibodies which are primarily responsible for protection against infection in the

lungs [11], secretory IgA antibodies that can be induced by intransal vaccination possess

higher cross-protective activity against SARS-CoV-2 variantviruses with different

antigenicities. Thus, a nasal vaccine that could induce highly cross-protective secretory

IgA antibodies in the nasal mucosa, which is the gateway to respiratory infection, would

be a more effective, reasonable vaccine candidate. It has been shown that the Omicron

variant, which is currently prevalent around the world, could replicate more effectively in

the bronchi than in the lungs, compared with other variants and the ancestor [45-48].

Therefore, there is high chance that a nasal vaccine which could induce cross-protective

IgA antibodies in the upper respiratory tract would be effective against the Omicron

variant as well.

When considering the COVID-19 vaccine, there is a concern about the potential risk

of lung eosinophilic immunopathology in post-vaccination infections as VAERD. The

FDA recommends addressing the potential risk of VAERD in animal models [27]. Since it

has been considered that a Th2-dominant response increases the risk of VAERD, we

investigated the T-cell response induced by intranasal vaccination combined with

ODN2006 in detail. While IL-4- or IL-5-secreting cells were highly observed in spleen

collected from mice subcutaneously vaccinated with alum as a typical Th2 adjuvant,

individuals who received subcutaneous vaccination with ODN2006 showed high

amounts of IFN-γ-secreting cells. Although nasal vaccination had no significant impact

on spleen cells compared to subcutaneous vaccination, IFN-γ-secreting cells were

significantly detected in draining cervical lymph nodes of intranasally vaccinated

individuals with ODN2006. It was suggested that ELISpot assay using spleen cells might

be unsuitable for the evaluation of T cell responses induced by intranasal vaccines. In

contrast, the IgG1/IgG2a ratios determined from the quantification of S-specific IgG1

and IgG2 subclasses in serum were likely able to assess T cell responses reflecting

neutralizing antibody titers, because the tendency to show similar levels of the

IgG1/IgG2a ratio among mice subcutaneously or intranasally vaccinated with ODN2006

was consistent with that of neutralizing antibody titers. Interestingly, eosinophil

infiltration into the lung correlated well with the serum IgG1/IgG2a ratio in our mouse

model. At this time, although the threshold for eosinophil infiltration that causes

eosinophilic pneumonia is unknown, analyzing immunization-induced IgG subclasses

could be an indicator for estimating whether there is a potential risk of VAERD. Although

several substances have been reported as potential mucosal adjuvants (e.g., cholera toxin

B subunit and synthetic double-stranded RNA), the potential risk of lung eosinophilic

immunopathology should be adequately investigated when designing a COVID-19

vaccine [8, 13].

In addition, the induction of B-cell memory and long-lived plasma cells by

vaccination is noteworthy, since germinal center formation is essential not only for the

production of high-affinity antibodies, but also for the determination of the B cell life

span.. The induction of Tfh and GCB cells by vaccination is essential to address this issue

[47, 49, 50]. In the current study, intranasal vaccination in the presence of ODN2006

successfully induced Tfh and GCB cells, long-lasting IgG antibodies in the serum, and

IgA antibodies in the nasal mucosa up to 16 weeks after the final vaccination.

Considering the emergence of new SARS-CoV-2 variants, nasal vaccines inducing a

secretory IgA antibody with high cross-protective ability on the mucosal epithelium of

the upper respiratory tract, which is the site of infection, could be highly useful, avoiding

repeated vaccinations using newly manufactured antigens. In conclusion, our study

showed that an intranasal COVID-19 vaccine of recombinant spike protein combined

with an adjuvant inducing a Th1-shifted response would be a safe and effective vaccine

not only for preparing cross-protective secretory IgA antibodies, but also to reduce the

potential risk of VAERD, that is, lung eosinophilic immunopathology.

Acknowledgements

We thank A. Kojima and A. Sataka (Department of Pathology, NIID) and N. Kurisaki

(Department of Virology, Kyushu Univ) for excellent support, and S. Takatsuka

(Department of Fungal Infection, NIID) for technical suggestions and helpful

discussions. Additionally, we would like to thank Editage (www.editage.com) for English

language editing.

Author Contributions

Conceptualization: A.A, T.He., and T. S

Methodology: T.He., A.A., T.Ha., N.I.-Y., S.I., Y.S. A.U., K.S., N.S.-S., N.N. and T.S.

Investigation: T.He, T.Ha, M.T., T.K., N.I.-Y., Y.S., S.M., A.U., K.S., S.S. and A.A.

Visualization: T.He. and A.A.

Funding acquisition: H.H. and T.S.

Project administration: H.H. and T.S.

Supervision: R.S., K.T., H.H., and T.S.

Writing—Original draft: T.H. and A.A.

Writing—Review and editing: T.He., A.A., T.Ha., M.T., T.K., N.I.-Y., S.I., Y.S., S.M.,

A.U., K.S., S.S., N.S.-S., N.N., K.T., R.S., H.H. and T.S.

Funding

This work was supported in part by the Naito foundation, and by a Grant-in-Aid for

Research on Emerging and Reemerging Infectious Diseases from the Japanese Ministry

of Health, Labor and Welfare and the Japan Agency for Medical Research and

Development (AMED) under grant numbers JP20nk0101603, JP20nk0101604,

JP20nk0101626, JP20fk0108411 and JP21fk0108083. The funding agencies had no role

in the study design, data collection and analysis, decision to publish, or manuscript

preparation.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal

relationships that could have influenced the work reported in this study.

References

[1] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical

characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a

descriptive   study.   Lancet.   2020;395:507-13.

https://doi.org/10.1016/S0140-6736(20)30211-7

[2] Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from

Patients   with   Pneumonia   in   China,   2019.   N   Engl   J   Med.   2020;382:727-33.

https://doi.org/10.1056/NEJMoa2001017

[3] Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety

and   Efficacy   of   the   BNT162b2   mRNA   Covid-19   Vaccine.   N   Engl   J   Med.

2020;383:2603-15. https://doi.org/10.1056/NEJMoa2034577

[4] Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. Efficacy and

Safety of the mRNA-1273 SARS-CoV-2 Vaccine. N Engl J Med. 2021;384:403-16.

https://doi.org/10.1056/NEJMoa2035389

[5] Knoll MD, Wonodi C. Oxford-AstraZeneca COVID-19 vaccine efficacy. Lancet.

2021;397:72-4. https://doi.org/10.1016/S0140-6736(20)32623-4

[6] Sadoff J, Gray G, Vandebosch A, Cardenas V, Shukarev G, Grinsztejn B, et al.

Safety and Efficacy of Single-Dose Ad26.COV2.S Vaccine against Covid-19. N Engl J

Med. 2021;384:2187-201. https://doi.org/10.1056/NEJMoa2101544

[7] Medicines and Healthcare products Regulatory Agency. UK medicines regulator

gives approval for first UK COVID-19 vaccine.,

https://www.gov.uk/government/news/uk-medicines-regulator-gives-approval-for-first-u

k-covid-19-vaccine; 2020 [accessed 15 May 2022]

[8] Tamura S, Samegai Y, Kurata H, Nagamine T, Aizawa C, Kurata T. Protection

against influenza virus infection by vaccine inoculated intranasally with cholera toxin B

subunit. Vaccine. 1988;6:409-13. https://doi.org/10.1016/0264-410x(88)90140-5

[9] Tamura S, Funato H, Hirabayashi Y, Kikuta K, Suzuki Y, Nagamine T, et al.

Functional role of respiratory tract haemagglutinin-specific IgA antibodies in protection

against influenza. Vaccine. 1990;8:479-85.

https://doi.org/10.1016/0264-410x(90)90250-p

[10] Tamura S, Funato H, Hirabayashi Y, Suzuki Y, Nagamine T, Aizawa C, et al.

Cross-protection against influenza A virus infection by passively transferred respiratory

tract IgA antibodies to different hemagglutinin molecules. Eur J Immunol.

1991;21:1337-44. https://doi.org/10.1002/eji.1830210602

[11] Asahi Y, Yoshikawa T, Watanabe I, Iwasaki T, Hasegawa H, Sato Y, et al.

Protection against influenza virus infection in polymeric Ig receptor knockout mice

immunized intranasally with adjuvant-combined vaccines. J Immunol. 2002;168:2930-8.

https://doi.org/10.4049/jimmunol.168.6.2930

[12] Asahi-Ozaki Y, Yoshikawa T, Iwakura Y, Suzuki Y, Tamura S, Kurata T, et al.

Secretory IgA antibodies provide cross-protection against infection with different strains

of influenza B virus. J Med Virol. 2004;74:328-35. https://doi.org/10.1002/jmv.20173

[13] Ichinohe T, Watanabe I, Ito S, Fujii H, Moriyama M, Tamura S, et al. Synthetic

double-stranded RNA poly(I:C) combined with mucosal vaccine protects against

influenza virus infection. J Virol. 2005;79:2910-9.

https://doi.org/10.1128/JVI.79.5.2910-2919.2005

[14] World Health Organization. Interim Statement on COVID-19 vaccines in the

context of the circulation of the Omicron SARS-CoV-2 Variant from the WHO

Technical Advisory Group on COVID-19 Vaccine Composition (TAG-CO-VAC),

https://www.who.int/news/item/11-01-2022-interim-statement-on-covid-19-vaccines-in-

the-context-of-the-circulation-of-the-omicron-sars-cov-2-variant-from-the-who-technica

l-advisory-group-on-covid-19-vaccine-composition; 2022 [accessed 6 May 2022]

[15] World Health Organization. The COVID-19 vaccine tracker and landscape

compiles detailed information of each COVID-19 vaccine candidate in development by

closely   monitoring   their   progress   through   the   pipeline.,

https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccin

es; 2022 [accessed 6 May 2022]

[16] Alu A, Chen L, Lei H, Wei Y, Tian X, Wei X. Intranasal COVID-19 vaccines: From

bench   to   bed.   EBioMedicine.   2022;76:103841.

https://doi.org/10.1016/j.ebiom.2022.103841

[17] Bolles M, Deming D, Long K, Agnihothram S, Whitmore A, Ferris M, et al. A

double-inactivated severe acute respiratory syndrome coronavirus vaccine provides

incomplete protection in mice and induces increased eosinophilic proinflammatory

pulmonary   response   upon   challenge.   J   Virol.   2011;85:12201-15.

https://doi.org/10.1128/JVI.06048-11

[18] Tseng CT, Sbrana E, Iwata-Yoshikawa N, Newman PC, Garron T, Atmar RL, et al.

Immunization with SARS coronavirus vaccines leads to pulmonary immunopathology

on challenge with the SARS virus. PLoS One. 2012;7:e35421.

https://doi.org/10.1371/journal.pone.0035421

[19] Yasui F, Kai C, Kitabatake M, Inoue S, Yoneda M, Yokochi S, et al. Prior

immunization with severe acute respiratory syndrome (SARS)-associated coronavirus

(SARS-CoV) nucleocapsid protein causes severe pneumonia in mice infected with

SARS-CoV. J Immunol. 2008;181:6337-48.

https://doi.org/10.4049/jimmunol.181.9.6337

[20] Sekimukai H, Iwata-Yoshikawa N, Fukushi S, Tani H, Kataoka M, Suzuki T, et al.

Gold nanoparticle-adjuvanted S protein induces a strong antigen-specific IgG response

against severe acute respiratory syndrome-related coronavirus infection, but fails to

induce protective antibodies and limit eosinophilic infiltration in lungs. Microbiol

Immunol. 2020;64:33-51. https://doi.org/10.1111/1348-0421.12754

[21] Iwata-Yoshikawa N, Uda A, Suzuki T, Tsunetsugu-Yokota Y, Sato Y, Morikawa S,

et al. Effects of Toll-like receptor stimulation on eosinophilic infiltration in lungs of

BALB/c mice immunized with UV-inactivated severe acute respiratory

syndrome-related coronavirus vaccine. J Virol. 2014;88:8597-614.

https://doi.org/10.1128/JVI.00983-14

[22] Honda-Okubo Y, Barnard D, Ong CH, Peng BH, Tseng CT, Petrovsky N. Severe

acute respiratory syndrome-associated coronavirus vaccines formulated with delta

inulin adjuvants provide enhanced protection while ameliorating lung eosinophilic

immunopathology. J Virol. 2015;89:2995-3007. https://doi.org/10.1128/JVI.02980-14

[23] Iwata-Yoshikawa N, Shiwa N, Sekizuka T, Sano K, Ainai A, Hemmi T, et al. A

lethal mouse model for evaluating vaccine-associated enhanced respiratory disease

during SARS-CoV-2 infection. Sci Adv. 2022;8:eabh3827.

https://doi.org/10.1126/sciadv.abh3827

[24] Kim HW, Canchola JG, Brandt CD, Pyles G, Chanock RM, Jensen K, et al.

Respiratory syncytial virus disease in infants despite prior administration of antigenic

inactivated vaccine. Am J Epidemiol. 1969;89:422-34.

https://doi.org/10.1093/oxfordjournals.aje.a120955

[25] An X, Martinez-Paniagua M, Rezvan A, Sefat SR, Fathi M, Singh S, et al.

Single-dose intranasal vaccination elicits systemic and mucosal immunity against

SARS-CoV-2. iScience. 2021;24:103037. https://doi.org/10.1016/j.isci.2021.103037

[26] Du Y, Xu Y, Feng J, Hu L, Zhang Y, Zhang B, et al. Intranasal administration of a

recombinant RBD vaccine induced protective immunity against SARS-CoV-2 in mouse.

Vaccine. 2021;39:2280-7. https://doi.org/10.1016/j.vaccine.2021.03.006

[27] U.S. Food and Drug Administration. Development and licensure of vaccines to

prevent COVID-19,

https://www.fda.gov/regulatory-information/search-fda-guidance-documents/developme

nt-and-licensure-vaccines-prevent-covid-19; 2020 [accessed 6 May 2022]

[28] Nakagawa T, Tanino T, Onishi M, Tofukuji S, Kanazawa T, Ishioka Y, et al.

S-540956, a CpG Oligonucleotide Annealed to a Complementary Strand With an

Amphiphilic Chain Unit, Acts as a Potent Cancer Vaccine Adjuvant by Targeting

Draining Lymph Nodes. Front Immunol. 2021;12:803090.

https://doi.org/10.3389/fimmu.2021.803090

[29] Amanat F, Stadlbauer D, Strohmeier S, Nguyen THO, Chromikova V, McMahon M,

et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. Nat Med.

2020;26:1033-6. https://doi.org/10.1038/s41591-020-0913-5

[30] Onodera T, Kita S, Adachi Y, Moriyama S, Sato A, Nomura T, et al. A

SARS-CoV-2 antibody broadly neutralizes SARS-related coronaviruses and variants by

coordinated recognition of a virus-vulnerable site. Immunity. 2021;54:2385-98 e10.

https://doi.org/10.1016/j.immuni.2021.08.025

[31] Hashiguchi T, Fukuda Y, Matsuoka R, Kuroda D, Kubota M, Shirogane Y, et al.

Structures of the prefusion form of measles virus fusion protein in complex with

inhibitors.   Proc   Natl   Acad   Sci   U   S   A.   2018;115:2496-501.

https://doi.org/10.1073/pnas.1718957115

[32] Hsieh CL, Goldsmith JA, Schaub JM, DiVenere AM, Kuo HC, Javanmardi K, et al.

Structure-based   design   of   prefusion-stabilized   SARS-CoV-2   spikes.   Science.

2020;369:1501-5. https://doi.org/10.1126/science.abd0826

[33] Pinto D, Park YJ, Beltramello M, Walls AC, Tortorici MA, Bianchi S, et al.

Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody.

Nature. 2020;583:290-5. https://doi.org/10.1038/s41586-020-2349-y

[34] Moriyama S, Adachi Y, Sato T, Tonouchi K, Sun L, Fukushi S, et al. Temporal

maturation of neutralizing antibodies in COVID-19 convalescent individuals improves

potency and breadth to circulating SARS-CoV-2 variants. Immunity. 2021;54:1841-52

e4. https://doi.org/10.1016/j.immuni.2021.06.015

[35] Nao N, Sato K, Yamagishi J, Tahara M, Nakatsu Y, Seki F, et al. Consensus and

variations in cell line specificity among human metapneumovirus strains. PLoS One.

2019;14:e0215822. https://doi.org/10.1371/journal.pone.0215822

[36] Matsuyama S, Nao N, Shirato K, Kawase M, Saito S, Takayama I, et al. Enhanced

isolation of SARS-CoV-2 by TMPRSS2-expressing cells. Proc Natl Acad Sci U S A.

2020;117:7001-3. https://doi.org/10.1073/pnas.2002589117

[37] Tian JH, Patel N, Haupt R, Zhou H, Weston S, Hammond H, et al. SARS-CoV-2

spike glycoprotein vaccine candidate NVX-CoV2373 immunogenicity in baboons and

protection in mice. Nat Commun. 2021;12:372.

https://doi.org/10.1038/s41467-020-20653-8

[38] Misharin AV, Morales-Nebreda L, Mutlu GM, Budinger GR, Perlman H. Flow

cytometric analysis of macrophages and dendritic cell subsets in the mouse lung. Am J

Respir Cell Mol Biol. 2013;49:503-10. https://doi.org/10.1165/rcmb.2013-0086MA

[39] Ishii H, Nomura T, Yamamoto H, Nishizawa M, Thu Hau TT, Harada S, et al.

Neutralizing-antibody-independent SARS-CoV-2 control correlated with

intranasal-vaccine-induced CD8(+) T cell responses. Cell Rep Med. 2022;3:100520.

https://doi.org/10.1016/j.xcrm.2022.100520

[40] Hassan AO, Shrihari S, Gorman MJ, Ying B, Yaun D, Raju S, et al. An intranasal

vaccine durably protects against SARS-CoV-2 variants in mice. Cell Rep.

2021;36:109452. https://doi.org/10.1016/j.celrep.2021.109452

[41] Toellner KM, Luther SA, Sze DM, Choy RK, Taylor DR, MacLennan IC, et al. T

helper 1 (Th1) and Th2 characteristics start to develop during T cell priming and are

associated with an immediate ability to induce immunoglobulin class switching. J Exp

Med. 1998;187:1193-204. https://doi.org/10.1084/jem.187.8.1193

[42] Kotaki R, Adachi Y, Moriyama S, Onodera T, Fukushi S, Nagakura T, et al.

SARS-CoV-2 Omicron-neutralizing memory B cells are elicited by two doses of

BNT162b2 mRNA vaccine. Sci Immunol. 2022;7:eabn8590.

https://doi.org/10.1126/sciimmunol.abn8590

[43] Mantus G, Nyhoff LE, Edara VV, Zarnitsyna VI, Ciric CR, Flowers MW, et al.

Pre-existing SARS-CoV-2 immunity influences potency, breadth, and durability of the

humoral response to SARS-CoV-2 vaccination. Cell Rep Med. 2022;3:100603.

https://doi.org/10.1016/j.xcrm.2022.100603

[44] Miyamoto S, Arashiro T, Adachi Y, Moriyama S, Kinoshita H, Kanno T, et al.

Vaccination-infection interval determines cross-neutralization potency to SARS-CoV-2

Omicron after breakthrough infection by other variants. Med (NY). 2022;3:249-61 e4.

https://doi.org/10.1016/j.medj.2022.02.006

[45] Halfmann PJ, Iida S, Iwatsuki-Horimoto K, Maemura T, Kiso M, Scheaffer SM, et

al. SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters. Nature.

2022;603:687-92. https://doi.org/10.1038/s41586-022-04441-6

[46] Hui KPY, Ho JCW, Cheung MC, Ng KC, Ching RHH, Lai KL, et al. SARS-CoV-2

Omicron variant replication in human bronchus and lung ex vivo. Nature.

2022;603:715-20. https://doi.org/10.1038/s41586-022-04479-6

[47] Haque A, Pant AB. Mitigating Covid-19 in the face of emerging virus variants,

breakthrough infections and vaccine hesitancy. J Autoimmun. 2022;127:102792.

https://doi.org/10.1016/j.jaut.2021.102792

[48] Uraki R, Kiso M, Iida S, Imai M, Takashita E, Kuroda M, et al. Characterization

and antiviral susceptibility of SARS-CoV-2 Omicron/BA.2. Nature. 2022.

https://doi.org/10.1038/s41586-022-04856-1

[49] Crotty S. T follicular helper cell differentiation, function, and roles in disease.

Immunity. 2014;41:529-42. https://doi.org/10.1016/j.immuni.2014.10.004

[50] Quast I, Tarlinton D. B cell memory: understanding COVID-19. Immunity.

2021;54:205-10. https://doi.org/10.1016/j.immuni.2021.01.014

Figure legends

Fig. 1. The induction of S-specific antibodies in serum, nasal mucosa and lungs of

mice vaccinated intranasally.

(**A**) Each of six mice were vaccinated three times at 2-week intervals. One week after the

final vaccination, serum, nasal wash (NW) and lung wash (LW) were collected for the

evaluation of antibody responses. (**B**, **C**) SARS-CoV-2 S-specific IgG titers and

neutralizing antibody titers in sera were measured by ELISA and microneutralization

assay, respectively. Graphs shown as the geometric mean titers ± the geometric standard

deviation (SD). The dashed line indicates the detection limit of measurement. (**D**, **E**) The

concentration of S-specific IgA antibodies in NW and LW was estimated by ELISA. Data

shown as the means ± SD. Each dotted line indicates the detection limit of measurement.

The p-values were calculated by Kruskal-Wallis test followed by Dunn's multiple

comparison test (*P < 0.05, **P < 0.01). in: intranasally, sc: subcutaneously.

Fig. 2. Intranasal vaccination adjuvanted with ODN2006 protected mice from the

lethal challenge with SARS-CoV-2.

(**A**) Each of 12 mice were vaccinated three times at 2-week intervals. Two weeks after the

final vaccination, mice were intranasally challenged with mouse adopted SARS-CoV-2

strain, QHmusX, into both lungs and nasal cavity (40 and 6 LD_{50} per mouse, respectively).

NW and LW were collected from each of six mice for the evaluation of antibody

responses at 3 dpi. Body weight and survival of six mice were monitored 10 days after

viral challenge. (**B**, **C**) Copy numbers of SARS-CoV-2 subgenomic RNA (sgRNA) in

NW and LW were evaluated by real-time RT-PCR. Data shown as the geometric mean ±

the geometric SD. The p-values were calculated by Kruskal-Wallis test followed by

Dunn's multiple comparison test (*$P < 0.05$, *** $P < 0.001$). (**D**, **E**) Body weight changes

and survival rates during 10 days of observation after challenge with QHmusX. Data

shown as the means ± SD. The p-values of body weight and survival were compared with

mice intranasally vaccinated with ODN2006 by mixed-model analysis followed by

Dunnett's multiple comparisons test and log-rank (Mantel-Cox) test (**$P < 0.01$).

Fig. 3. Secretory IgA antibodies were protective against SARS-CoV-2 variants.

(**A**) Mice were vaccinated three times at 2-week intervals. Two weeks after the final

vaccination, each of six mice were intranasally challenged with 3.5×10^5 TCID$_{50}$ of Alpha

(**B**, **E**), Beta (**C**, **F**) or Gamma (**D**, **G**) variant into both lungs and nasal cavity. At 3 dpi,

NW and LW were collected. Copy numbers of sgRNA in NW (**B-D**) and LW (**E-G**) were

evaluated by real-time RT-PCR. Data shown as the geometric mean ± the geometric SD.

The p-values were calculated by Kruskal-Wallis test followed by Dunn's multiple

comparison test (*P < 0.05, *** P < 0.001).

Fig. 4. Germinal center formation and long-lasting antibody responses.

(**A**, **B**) The frequency of Tfh cells (CXCR5$^+$ PD-1$^+$) among CD4$^+$ cells and GCB cells

(GL7$^+$ CD95$^+$) among CD19$^+$ B cells was evaluated by flow cytometry in cervical lymph

nodes collected from mice one week after third vaccination. Results were shown as the

mean ± SD. (**C**) To evaluate time-course of antibody responses induced by three doses of

vaccination, serum samples were collected every two weeks. After 20 weeks from the

initial vaccination, serum and NW samples were collected (six mice per group). (**D**) The

changes of optical density (405 nm) of S-specific IgG antibodies were shown as the mean

± SEM. (**E**) S-specific IgA concentration in NW collected at 20 weeks after the initial

vaccination was evaluated by ELISA. Results were shown as the mean ± SD. Each dotted

line indicates the detection limit of measurement. The p-values were calculated by

Kruskal-Wallis test followed by Dunn's multiple comparison test (*P < 0.05, **P < 0.01,

***P < 0.001).

Fig. 5. Induction of significant Th1 response by using ODN2006 as an adjuvant.

(**A**) Spleen and cervical lymph nodes were collected from individuals shown in

Figure.1(**A**). Single cell suspensions obtained from each tissue was cultured under the

stimulation of peptide pool of SARS-CoV-2 S protein. Spleen (**B-D**) or cervical lymph

node cells (**E-G**) were counted on the production of IFN-r (**B**, **E**), IL-4 (**C**, **F**) and IL-5 (**D**,

G) by ELISpot assay. Data shown as the means ± SD. (**H**, **I**) SARS-CoV-2 S-specific

IgG1 and IgG2a antibodies were quantified by ELISA, and (**J**) IgG1/IgG2a ratio was

calculated. Data shown as the geometric means ± geometric SD. The p-values were

calculated by Kruskal-Wallis test followed by Dunn's multiple comparison test (*P < 0.05,

**P < 0.01, ***P < 0.001).

Fig. 6. Eosinophilic infiltration into the lungs was suppressed in mice vaccinated

with ODN2006 inducing a remarkable Th1 response.

(**A**) Each of six mice were vaccinated three times at 2-week intervals. Two weeks after the

final vaccination, mice were intranasally challenged with mouse adopted SARS-CoV-2

strain, QHmusX, into both lungs and nasal cavity (40 LD_{50} and 6 LD_{50} per mouse,

respectively). At 6 dpi, infiltration of eosinophils was evaluated by histopathological or

flow cytometric analysis, on lungs collected from surviving 1~2 or 2~4 mice in each

group, respectively. (**B**) Histopathological findings of mouse lungs by eosinophil staining

using the combined eosinophil-mast cell staining (C.E.M.) kit. Green arrow heads point

to eosinophils. The images in the lower panels are enlargements of area boxed in the

upper images. The scale bar is 200 µm for low magnification and 20 µm for high

magnification. (**C**) The percentage of eosinophils ($CD11b^+$ $CD11c^-$ $Siglec-F^+$ $Ly-6G^-$)

among CD45+ cells was analyzed by flow cytometry. Results were shown as the mean ±

SD. The p-values were calculated by Kruskal-Wallis test followed by Dunn's multiple

comparison test. (**D**) Correlation between S-specific IgG1/IgG2a ratio and the frequency

777   of eosinophils in lung cells was analyzed by Spearman correlation. S-specific

778   IgG1/IgG2a ratio was evaluated using serum samples collected seven days before the

779   virus challenge.

Vaccination		
Antigen	Adjuvant	Route
-	-	-
+	-	in
+	ODN2006	in
+	ODN2006	sc
+	Alum	sc

A Vaccination

1st · 2nd · 3rd

0 · 14 · 28 · 35 day

Collect serum, NW and LW

20~24 weeks old female BALB/c mice

B S-specific serum IgG titer

C Serum neutralization titer

D S-specific nasal IgA (ng/mL)

E S-specific lung IgA (ng/mL)

A Vaccination
1st 2nd 3rd
Challenge
0 14 28 42 45 52 day

20~24 weeks old female BALB/c mice

Collect NW and LW
Sacrifice at the end-point or 10 dpi

Vaccination		
Antigen	Adjuvant	Route
-	-	-
+	-	in
+	ODN2006	in
+	ODN2006	sc
+	Alum	sc

B Nasal sgRNA (copies/mL)

C Lung sgRNA (copies/mL)

D

E

Days Post Infection

Body weight (%)

Survival rate (%)

naive — Ag (in)

Ag+ODN (in) — Ag+ODN (sc) — Ag+Alum (sc)

A

Vaccination
1st   2nd   3rd   Challenge

20~24 weeks old
female
BALB/c mice

0	14	28	35	42	48	day

Collect
serum

Collect
lung

C

Eosinophils/CD45+ cells(%)

Antigen	+	+	+	+
Adjuvant	-	ODN	ODN	Alum
Route	in	in	sc	sc

D

S-specific IgG1/IgG2a ratio

r=0.779
P=0.0015

Eosinophils/CD45+cells(%)

B

	Ag (in)	Ag+ODN2006 (in)	Ag+ODN2006 (sc)	Ag+Alum (sc)
x10				
x40				