Genomic surveillance of carbapenem-resistant *Klebsiella* in Wales reveals persistent spread of *K. pneumoniae* ST307 and adaptive evolution of pOXA-48-like plasmids

Sophia David[1], Massimo Mentasti[2], Kirsty Sands[3,4], Edward Portal[4], Lee Graham[5], Joanne Watkins[5], Catie Williams[5], Brendan Healy[6], Owen B. Spiller[4], David M. Aanensen[1], Mandy Wootton[2] & Lim Jones[2]

[1]Centre for Genomic Pathogen Surveillance, Big Data Institute, University of Oxford, Oxford, OX3 7LF, UK

[2]Specialist Antimicrobial Chemotherapy Unit, Public Health Wales Microbiology Cardiff, CF14 4XW, UK

[3]Department of Zoology, University of Oxford, Oxford OX1 3SZ, UK

[4]Division of Infection and Immunity, Cardiff University, Cardiff, CF14 4XN, UK

[5]Pathogen Genomics Unit, Public Health Wales Microbiology Cardiff, CF14 4XW, UK

[6]Public Health Wales Microbiology Swansea, SA2 8QA, UK

Corresponding authors: Sophia David, sophia.david@cgps.group; Lim Jones, lim.jones@wales.nhs.uk

Keywords: genomic surveillance, *Klebsiella*, carbapenem resistance, ST307, ST1788, pOXA-48-like plasmids
Repositories: All raw short read sequence data and hybrid assemblies are available in the European Nucleotide Archive (ENA) under project accession PRJEB48990.

Abstract

Rising rates of multi-drug resistant *Klebsiella* infections necessitate a comprehensive understanding of the major strains and plasmids driving spread of resistance elements. Here we analysed 540 *Klebsiella* isolates recovered from patients across Wales between 2007 and 2020 using combined short- and long-read sequencing approaches. We identified resistant clones that have spread within and between hospitals including the high-risk strain, sequence type (ST) 307, which acquired the $bla_{OXA-244}$ carbapenemase gene on a pOXA-48-like plasmid. We found evidence that this strain, which caused an acute outbreak largely centred on a single hospital in 2019, had been circulating undetected across South Wales for several years prior to the outbreak. In addition to clonal transmission, our analyses revealed evidence for substantial plasmid spread, mostly notably involving bla_{KPC-2} and $bla_{OXA-48-like}$ (including $bla_{OXA-244}$) carbapenemase genes that were found among many species and strain backgrounds. Two thirds (20/30) of the bla_{KPC-2} genes were carried on the Tn*4401a* transposon and associated with IncF plasmids. These were mostly recovered from patients in North Wales, reflecting an outward expansion of the plasmid-driven outbreak of bla_{KPC-2}-producing *Enterobacteriaceae* in North-West England. 92.1% (105/114) of isolates with a $bla_{OXA-48-like}$ carbapenemase carried the gene on a pOXA-48-like plasmid. While this plasmid family is highly conserved, our analyses revealed novel accessory variation including integrations of additional resistance genes. We also identified

46 multiple independent deletions involving the *tra* gene cluster among pOXA-48-like plasmids in
47 the ST307 outbreak lineage. These resulted in loss of conjugative ability and signal adaptation of
48 the plasmids to carriage by the host strain. Altogether, our study provides the first high resolution
49 view of the diversity, transmission and evolutionary dynamics of major resistant clones and
50 plasmids of *Klebsiella* in Wales and forms an important basis for ongoing surveillance efforts.
51
52

53 # Data Summary

55 All raw short read sequence data and hybrid assemblies are available in the European
56 Nucleotide Archive (ENA) under project accession PRJEB48990.
57
58
59

Introduction

The *Klebsiella* genus, which belongs to the *Enterobacteriaceae* family, includes
several species that are important opportunistic pathogens of humans. In particular,
Klebsiella pneumoniae is a leading cause of healthcare-associated infections globally (Magill
et al. 2014; Ling et al. 2015; Vincent et al. 2020). *Klebsiella* species cause a range of
disease types including pneumonia, skin and wound infections, urinary tract infections, and
sepsis. These occur most frequently in the elderly, neonates and immunocompromised
individuals (Meatherall et al. 2009).
Klebsiella are commonly carried in the gastrointestinal tract of healthy humans
(Podschun & Ullmann, 1998). Analysis using whole genome sequencing (WGS) performed
on paired screening swabs and clinical samples has demonstrated that gastrointestinal
carriage is a major reservoir of *K. pneumoniae* infections (Gorrie et al. 2017). In addition,
outbreaks are frequently reported in healthcare settings, driven by transmission via person-
to-person contact, contaminated environmental surfaces and medical devices (e.g.
ventilators, catheters) (Snitkin et al. 2012; Gu et al. 2018; Tavoschi et al. 2020).
Across the world, studies have reported an increasing prevalence of infections with
multidrug-resistant *Klebsiella* (Cassini et al. 2019; Castanheira et al. 2019). A particular
concern has been the rise of infections that are resistant to carbapenems, which has been
listed as a "critical" threat by the World Health Organisation (WHO, 2017). Carbapenems are
a class of antibiotics that are vital for treatment of severe *Klebsiella* infections due to their
broad spectrum of activity (e.g. against extended-spectrum beta lactamase (ESBL)
producers) and limited adverse effects. Carbapenem resistance in *Klebsiella* is usually
conferred by plasmid-encoded carbapenemase genes including bla_{KPC}, $bla_{OXA-48-like}$, bla_{VIM},
bla_{NDM} and bla_{IMP} (Queenan & Bush, 2007). These are often associated with mobile genetic
elements (MGEs) such as transposons, which commonly insert into plasmid backbones.
A new cohort of drugs has become available in recent years to treat carbapenem-
resistant infections including novel beta lactam-beta lactamase inhibitor (BL-BLI)
combinations (e.g. ceftazidime-avibactam, meropenem-vaborbactam) and the siderophore
cephalosporin, cefiderocol. However, some of these drugs are not effective against certain
types of carbapenemase producer, including the BL-BLIs which have no activity against

metallo-betalactamases (bla_{VIM}, bla_{NDM} and bla_{IMP}). These also may not work reliably against
non-carbapenemase-producing carbapenem resistant strains (Bradley & Lee, 2019).
As a result, a multi-pronged approach is required for the control of multi-drug
resistant *Klebsiella*, including new therapeutics, heightened infection control and enhanced
surveillance. The latter involves acquiring a detailed understanding of the diversity and
transmission routes of different resistance mechanisms. Integration of genome sequencing
into surveillance systems, which is ongoing across many public health laboratories, is
providing a vital role in achieving this. However, while the current focus is largely on
monitoring important strains and detecting clonal transmission, there is still a strong need for
bioinformatic workflows that provide data on the role of specific plasmid vectors in resistance
spread.
The Specialist Antimicrobial Chemotherapy Unit (SACU) based at Public Health
Wales (PHW) Microbiology in Cardiff provides a reference laboratory service for the
identification and confirmation of antimicrobial resistance mechanisms in bacterial isolates
including *Klebsiella*, referred by microbiology laboratories across and beyond Wales. It also
collects *Klebsiella (*and other bacterial) isolates from blood cultures for surveillance purposes
and receives isolates from suspected outbreaks as part of the development of PHW's Whole
Genome Sequencing (WGS) based typing service, in partnership with the Pathogen
Genomics Unit (PenGU). This study describes the retrospective genomic analysis of a
collection of *Klebsiella* isolates (*n*=540) gathered by SACU between 2007 and 2020. Using a
combination of short- and long-read sequencing, our analysis revealed detailed insights into
the evolutionary dynamics of the major *K. pneumoniae* clones and plasmids driving spread
of antibiotic resistance genes in Wales.
114
115
116

Methods

Sample collection
540 *Klebsiella* isolates were included in this study (**Supplementary Tables 1 & 2**).
Isolates were obtained from clinical, screen or environmental samples between 2007 and
2020 and bead-stored at -80°C within the SACU culture collection at PHW. They were
cultured overnight from beads on blood agar (OXOID, UK) at 35 ± 1°C prior to sequencing.
124

Short read sequencing
Isolates were subbed onto 500μL of Tryptone Soya Broth (TSB) (E&O Laboratories,
UK) and incubated overnight at 35 ± 1°C. Growth from 125μL of TSB culture was harvested
at 12,000*g* for 2 min and the supernatant discarded. A lysis step was performed by adding
190μL of Buffer G2 (Qiagen, Germany) and 10μL of Proteinase K (Qiagen, Germany)
followed by incubation at 56°C for 30 min. DNA was extracted from the lysates using a
generic 3.0.4 protocol on the EMAG platform (bioMerieux, France) according to the
manufacturer's instructions and eluted in 100μL final volume. DNA was quantified using the
FLUOstar Omega fluorometer (BMG Labtech, Germany) and normalised to approximately
0.4ng/μL. Sequencing libraries were prepared using the Nextera XT DNA library preparation
kit and combinations of IDT index plates (Illumina, USA). The final libraries were cleaned
through AMPure SPRI bead (Beckman Coulter, USA) fragment size selection, and quantified

by the Collibri™ Library Quantification Kit (Invitrogen, USA) before normalising to
approximately 0.5nM. The normalised final libraries were pooled and loaded onto MiSeq or
NextSeq platforms (Illumina, USA).
140

Long read sequencing
Isolates were subbed onto 2mL of LB broth (Becton Dickinson, UK) and incubated
overnight at 35 ± 1°C. DNA was extracted using the automated QIAcube (Qiagen) platform
with an additional RNAse step and then quantified using the Qubit 4.0 (Thermofisher) using
HS and BR dsDNA kits as appropriate. DNA was purified and concentrated using SPRI
beads (Mag-Bind TotalPure, Omega) at a 1:1 ratio with a final 15µL elution in nuclease-free
water. Genomic libraries were prepared using the Rapid Barcoding Kit (SQK-RBK004;
Oxford Nanopore Technology, UK). Eight to twelve isolates were loaded onto a FLO-MIN106
R9.4 flow cell and sequenced on a MinION device (Oxford Nanopore Technology, UK).
Sequencing was performed on an Intel i7-6700 desktop computer over a 72 h period and
basecalling was performed within MinKnow using Guppy v4.5.4 (Oxford Nanopore
Technology, UK).
153

Short-read and hybrid assembly
Short sequence reads from all isolates were assembled using SPAdes v3.10.0
(Bankevich et al. 2012) using the "--careful" flag and the "--cov-cutoff" flag set to "auto". Long
sequence reads were de-multiplexed within Minknow and assembled with the corresponding
short reads using Unicycler v0.4.7 (Wick et al. 2017) with default parameters. Assembly
statistics are available in **Supplementary Tables 1 and 2**. Assemblies were annotated
using Prokka v1.14.5 (Seemann, 2014).
161

162   **Characterisation of *Klebsiella* genomes**
163   Kleborate v2.0 (Lam et al. 2021) was used to determine the species and sequence type
164   (ST) (for species with available schemes) from the assemblies, and identify virulence and
165   resistance genes. Kaptive (Wyres et al. 2016), integrated into Kleborate, was used to type
166   the capsule and O antigen biosynthesis loci.
167

Species-wide phylogenetic analyses
Core genes (i.e. those present in ≥99% of isolates) were identified among assemblies of
each *Klebsiella* species using Panaroo v1.2.4 (Tonkin-Hill et al. 2020) in the "sensitive"
mode (in species with ≥5 isolates only). Variable positions were extracted from each core
gene alignment using SNP-sites (Page et al. 2016). These were used to construct a
maximum likelihood phylogenetic tree for each species using IQ-TREE v1.6.10 (Nguyen et
al. 2015).
175

Phylogenetic analyses of individual STs
Additional public genomes from *K. pneumoniae* ST307 (*n*=964) and ST1788 (*n*=2),
the two most frequent STs in the PHW collection, were identified using the Pathogenwatch
database (Argimon et al. 2021). Using the available accession numbers, all corresponding
short-read data were downloaded from the ENA while the associated metadata was
obtained from Pathogenwatch (https://pathogen.watch/).
Short-read sequencing data from ST307 and ST1788 isolates from both the PHW
and public collections (including also single-locus variants of ST307 from the PHW

collection) were mapped to ST-specific reference genomes. For ST307, this was the complete chromosome from the hybrid assembly of ARGID_32304, while the short-read assembly of ARGID_33748 (62 contigs) was used for ST1788. Mapping was performed using Burrows Wheeler Aligner (Li & Durbin, 2009) with SNPs identified using SAMtools v1.2 mpileup and BCFtools v1.2 (Li et al. 2009). Gubbins v2.4.1 (Croucher et al. 2015) was used to remove recombined regions from the resulting pseudo-genome alignments and generate maximum likelihood phylogenies based on the remaining variable sites. Pairwise SNP differences between isolates were determined from the pseudo-genome sequences using pairsnp (https://github.com/gtonkinhill/pairsnp).

Visualisation of trees and metadata

Phylogenetic trees were visualised together with metadata using the Interactive Tree of Life tool v6 (Letunic & Bork, 2021) and further annotated using Adobe Illustrator v2017.1.0. Interactive visualisations were also created using Microreact v178 (Argimon et al. 2016).

Characterisation of plasmids and MGEs

Plasmid replicon types carried by all isolates were determined from the short-read sequence data using Ariba v2.14.6 with the PlasmidFinder database (Carattoli et al. 2014). The PlasmidFinder tool (https://bitbucket.org/genomicepidemiology/plasmidfinder/src/master/) (Carattoli et al. 2014) was used to identify plasmid replicons in completely sequenced plasmids. oriTfinder v1.1 (Li et al. 2018) was used to identify the origin of transfer site (*oriT*) among complete pOXA-48-like plasmid sequences.

Linear comparisons between multiple pairs of complete plasmids were generated using EasyFig (Sullivan et al. 2011) and further annotated using Adobe Illustrator v2017.1.0. The percentage nucleotide similarities between individual plasmid genes and those from a reference plasmid were determined using BLASTn v2.6.0 (Camacho et al. 2009). The number of SNPs between pairs of aligned plasmids were determined using NUCmer v3.1 from the MUMmer package (Kurtz et al. 2004).

The presence of the Tn*4401* transposon among bla_{KPC}-encoding isolates, together with the Tn*4401* variant type, was determined from the short sequence reads using TETyper v1.1 (Sheppard et al. 2018).

Identification and comparison of pOXA-48-like plasmid sequences from short-read data

Sequence reads from all isolates were mapped to the 61.8kb IncL plasmid, pOXA48a (Poirel et al. 2012) (accession JN626286), using Burrows Wheeler Aligner (Li & Durbin, 2009) with SNPs identified using SAMtools v1.2 mpileup and BCFtools v1.2 (Li et al. 2009). The binary alignment map (BAM) file from each isolate was used to determine the length of the reference plasmid that was mapped by at least one sequence read. A SNP matrix, calculated from pseudo-plasmid sequences of isolates with ≥70% mapping to the reference, was used to construct a minimum spanning tree using the standalone version of GrapeTree v1.5.0 (Zhou et al. 2018). Pairwise SNP differences between all plasmid pseudo-sequences were also determined using pairsnp (https://github.com/gtonkinhill/pairsnp).

To determine the short-read plasmid coverage over the pOXA48a reference relative to the chromosomal coverage, we mapped all isolates to the NTUH-K2044 chromosome

(accession AP006725.1) as described above. We used the "genomecov" function within
bedtools v2.29.0 (Quinlan & Hall, 2010) to determine the chromosomal coverage using
sorted BAM files as input, and determined the median value for each isolate. We then
calculated the plasmid coverage, relative to the median chromosomal coverage, across
100bp windows of each mapped plasmid sequence (sliding every 20bp) and visualised the
resulting matrix using the "heatmap.2" function in R v.4.0.2.
237
Bacterial conjugation
Transfer of the pOXA-48-like plasmid-encoded $bla_{OXA-244}$ gene by six *K. pneumoniae*
ST307 isolates was determined using a liquid-mating conjugation method. All mating
experiments were performed using both the *E. coli* J53 strain (200mg/L sodium azide) and a
K. pneumoniae strain, 71.1 (passaged over 10d to generate a rifampicin MIC of 4,096mg/L)
as separate recipients. Two mating culture ratios were tested, 1:3 and 5:1, for both donor
and recipient respectively. Briefly, donor and recipient isolates were grown to log phase in
LB broth and mixed according to the appropriate mating-culture ratio, then incubated at 37°C
for 16-20h. Following serial dilution of the mating cultures, transconjugant colonies were
selected for using appropriate chromogenic agar (Sigma Aldrich) and antibiotic combination
(J53, 200mg/L sodium azide and 4mg/L meropenem; 71.1, 256mg/L rifampicin). To
differentiate donor-recipient in the *K. pneumoniae-K. pneumoniae* experiment, the ST307
donor colonies were selected for using the chromogenic agar antibiotic combination (4mg/L
meropenem and ampicillin at 32mg/L) as 71.1 was sensitive to ampicillin. Transfer of the
$bla_{OXA-244}$ was confirmed via PCR, and transconjugates from the *K.pneumoniae-*
K.pneumoniae experiments were further confirmed by short-read (Illumina) sequencing.
Briefly, gDNA from transconjugates was extracted using the automated QIAcube (Qiagen).
gDNA libraries were prepared using the Nextera XT v2 kit with bead-based normalisation,
and sequenced using V3 chemistry on an Illumina Miseq. The conjugation frequency was
calculated as a ratio of the CFU/mL of transconjugate CFU/mL to the donor isolate.
258
259
260 **Data availability**
261   All raw short read sequence data and hybrid assemblies are available in the ENA
262 under project accession PRJEB48990. Individual accession numbers for samples are
263 available in **Supplementary Tables 1 and 2**.
264
265
266
267 # Results
268
269
270 **High diversity of carbapenemase-producing *Klebsiella* among clinical isolates**
271 **in Wales**
272
273   We first used short-read sequencing to analyse 540 *Klebsiella* isolates collected
274 between 2007 and 2020 that had largely been referred to SACU at PHW for investigation of
275 antimicrobial resistance or obtained from blood cultures (regardless of the resistance
276 phenotype) (**Supplementary Table 1**). They also included three contextual isolates from UK

NEQAS. Overall, 387/540 (71.7%) isolates were from clinical samples (including 238 from blood, 93 from urine, 23 from a wound site, 33 other/unknown), 143/540 (26.5%) from screen samples, 6/540 (1.1%) from environmental samples and 4/540 (0.7%) from an unknown source. The clinical and screen isolates were recovered from 405 patients in healthcare facilities across Wales, of whom 72 (17.8%) contributed two or more isolates (up to eight).

Among our collection, *K. pneumoniae* was the dominant species, accounting for 421/540 (78.0%) isolates. A further eight species were also identified: *K. aerogenes* (33/540; 6.1%), *K. michiganensis* (27/540; 5.0%), *K. variicola* (23/540; 4.3%), *K. oxytoca* (17/540; 3.1%), *K. quasipneumoniae* (11/540; 2.0%), *K. grimontii* (5/540; 0.9%), *K. planticola* (2/540; 0.4%) and *K. pasteurii* (1/540; 0.2%). A phylogenetic tree of the 421 *K. pneumoniae*, constructed from an alignment of 4034 core genes, demonstrated a high number of distinct lineages (**Figure 1;** https://microreact.org/project/aSJqpc9MZVZcurTWtoadW8-k-pneumoniae-phw-n421). We identified 130 STs across this species, with 98 (75.4%) of these restricted to a single patient. However, some STs were observed frequently including ST307 (88 isolates; 55 patients), ST1788 (43 isolates; 30 patients), ST20 (19 isolates; 10 patients), ST15 (19 isolates; 16 patients) and ST14 (17 isolates; 12 patients). Phylogenetic analyses of all other *Klebsiella* species also revealed a high diversity among each, with most isolates occurring as singletons or in small clusters (see **Supplementary Table 3** for Microreact URLs).

Inspection of the *Klebsiella* genomes for antimicrobial resistance markers revealed that 282/540 (52.2%) isolates carried an ESBL gene, which was largely $bla_{CTX-M-15}$ (228/282; 80.9%) (**Figure 1**). 207/540 (38.3%) isolates carried a carbapenemase gene. The most frequently observed carbapenemases were $bla_{OXA-244}$ (58/207; 28.0%), bla_{OXA-48} (46/207; 22.2%), bla_{KPC-2} (31/207; 15.0%), bla_{NDM-1} (29/207; 14.0%), bla_{IMP-4} (16/207; 7.7%) and bla_{VIM-4} (12/207; 5.8%). Carbapenemase-encoding isolates were found in six of the nine observed species, although *K. pneumoniae* accounted for the vast majority (181/207; 87.4%). Among *K. pneumoniae*, isolates carrying carbapenemase genes were found in 52 STs. However, $bla_{OXA-244}$ was mostly carried by ST307 (49/58; 84.5%), bla_{IMP-4} by ST20 (15/16; 93.8%), and bla_{VIM-4} by ST15 (11/12; 91.7%) or their related variants. Other carbapenemases such as bla_{KPC-2} and bla_{OXA-48} were more widely dispersed across different species and STs.

Persistent spread of globally- and locally-important clones of *K. pneumoniae*

Despite the overall high diversity among isolates, our phylogenetic analyses revealed numerous small to large clusters, representing clonal transmission of *Klebsiella* within and between healthcare facilities in Wales. Below we discuss features of the two largest clusters involving *K. pneumoniae* ST307 and ST1788.

ST307

The most frequently observed ST in our collection, ST307, is an important clone that has been identified in diverse locations worldwide and associated with the ESBL gene, $bla_{CTX-M-15}$ (Peirano et al. 2020). In 2019, an outbreak involving ST307 was identified in an acute tertiary hospital in South West Wales (Hospital A), based on epidemiological and resistance profiling data. The isolates showed a multi-drug resistant phenotype (resistance to third-generation cephalosporins, gentamicin, ciprofloxacin +/- ertapenem). The

carbapenemase gene, $bla_{OXA-244}$, was detected in the subset of isolates that exhibited
carbapenem resistance (ertapenem +/- meropenem +/- imipenem). Recognition of the
outbreak prompted regular faecal screening for carbapenemase-producing
Enterobacteriaceae (CPE) carriage in the initial affected ward, and this was extended to
additional wards where clinical isolates of $bla_{OXA-244}$-producing *K. pneumoniae* were later
identified. Here we analysed genome data of 88 isolates from ST307 (or single-locus
variants) collected between 2017 and 2020 from ten hospitals and additional GP surgeries in
Wales, with particular emphasis on understanding the emergence, evolution and
transmission of the outbreak strain.

We constructed a phylogenetic tree of these 88 isolates, also including a further 964
geographically-diverse public ST307 genomes to aid contextualisation. The phylogeny
demonstrated that 82/88 (93.2%) of the ST307s from Wales formed a monophyletic cluster
(**Figure 2A**; https://microreact.org/project/1ZRxgto9HVsaV6qRHAuf6P-st307s-phw-n1052).
This is indicative of a single major introduction of this strain into Wales followed by local
spread. The cluster incorporated clinical isolates from 19 patients (of whom 7 had
bloodstream infections), as well as screen isolates from a further 31 patients.

Almost all of the 82 clustered isolates carried $bla_{CTX-M-15}$ (80/82; 97.6%) while a
descendent subclade of the cluster later acquired $bla_{OXA-244}$ (49/82; 59.8%). A single isolate
also carried the combination of $bla_{CTX-M-15}$ and bla_{OXA-48}, the latter of which has a single amino
acid substitution relative to $bla_{OXA-244}$. We searched for alterations (including truncations) in
major outer membrane porin genes that have been shown to reduce susceptibility to
carbapenems when acting in concert with ESBL, AmpC or carbapenemase enzymes
(Martinez-Martinez et al. 1999; Hamzaoui et al. 2018). We found that all 82 isolates carried a
truncated *ompK35* gene while 12/82 (14.6%) also possessed a truncated *ompK36* gene.

Isolates in the $bla_{OXA-244}$ subclade were characterised by low pairwise SNP
differences (range 0-116; mean 15.5) and mostly collected from patients (both clinical and
screen) or environmental samples in Hospital A (40/49; 81.6%) over the outbreak period
(**Figures 2B & 2C**). This is consistent with heightened within-hospital transmission. The
remaining nine $bla_{OXA-244}$-encoding isolates were recovered from other nearby hospitals and
GP surgeries in the South Wales region. Notably, the first sampled isolates from this
subclade were obtained from a single patient in 2017 (outside of Hospital A). This
demonstrates the acquisition of $bla_{OXA-244}$ by this strain and emergence of this subclade at
least two years prior to its later expansion within Hospital A.

The non-$bla_{OXA-244}$-encoding isolates in the Welsh ST307 cluster had been referred to
SACU from 2019 onwards, largely prompted by the similarity of their phenotypic profiles to
the $bla_{OXA-244}$-encoding outbreak isolates rather than by epidemiological links. Despite their
collection over a limited timeframe, we found that they were considerably more divergent
than the $bla_{OXA-244}$-encoding isolates (pairwise SNP differences from 0-134; mean 54.4)
(**Figure 2B**). They were also recovered from a wider number of hospitals in South Wales
(nine in total) although included isolates from Hospital A. These findings are further
consistent with the undetected circulation of this ST307 strain across South Wales over
several years prior to the emergence and expansion of the $bla_{OXA-244}$-encoding outbreak
subclade in Hospital A in 2019.

The $bla_{OXA-244}$-encoding ST307 outbreak was declared formally closed in May 2020
with no further isolates of this subclade identified since March 2020.

369
370
371 ***ST1788***

ST1788 is the second most common ST of *K. pneumoniae* in our collection. In contrast to ST307, this has rarely been observed elsewhere to our knowledge. SACU first detected ST1788 during the 2019 ST307 outbreak investigation and it continued to be recovered across South Wales up until the end of the study period (November 2020). The 43 ST1788 isolates in this sequenced collection are derived from 30 patients (including 18 with clinical isolates and 3 with blood isolates) across seven hospitals and additional GP surgeries. They include 20 isolates from 13 patients obtained from a rehabilitation unit (Hospital B) in South East Wales from early 2020 onwards.

To investigate the evolution and spread of this strain, we constructed a phylogenetic tree of the 43 ST1788 isolates from Wales, together with two publicly available ST1788 genomes from Nigeria. This revealed that the Welsh isolates formed a distinct cluster (**Figure 3;** https://microreact.org/project/rMY2fakRxkNKXqqBPs3is6-st1788-phw-n45). Although additional contextual data are required to draw strong conclusions on the routes of strain spread, these findings are consistent with a single introduction into South Wales followed by local transmission within and between healthcare facilities. This is further supported by the low SNP differences between our isolate pairs (0-36 SNPs, mean 17.9), calculated after recombination removal, suggestive of a recent clonal expansion.

Our phylogenetic analysis also revealed that the isolates obtained from patients in Hospital B clustered together, also with a bloodstream isolate from an inpatient at an acute tertiary care hospital in South East Wales (Hospital C) who was transferred from Hospital B (**Figure 3**). This demonstrated the occurrence of a single introduction of the strain into Hospital B followed by within-hospital spread. Apart from the isolate obtained from the Hospital C patient, all isolates obtained from Hospital B were screen isolates, obtained via an enhanced screening programme at this site.

Almost all ST1788 isolates from Wales carried $bla_{CTX-M-15}$ (42/43; 97.7%) while three isolates from one patient also carried bla_{OXA-48} (**Figure 3**). The majority (39/43; 90.7%) had intact *ompK35* and *ompK36* genes. Notably, all ST1788 isolates possessed the KL2 capsule type, which has been associated with increased virulence (Shon et al. 2013), and most (39/43; 90.7%) also possessed a yersiniabactin locus.

At the time of writing, transmission of ST1788 remains an ongoing concern, in particular in Hospital B. We also have evidence of ongoing secondary cases at nearby hospitals within the health board, linked by epidemiology and genomic typing, and have recently recovered ST1788 isolates from patients in other regions of Wales.

Plasmid epidemiology of bla_{KPC-2}-producing *Klebsiella* reflects distinct hospital networks and referral patterns between North and South Wales

We next aimed to elucidate the spread of particular carbapenemase genes in Wales where we could not identify clonal transmission as a key driver. We first investigated bla_{KPC-2} genes, which were found in 30 isolates from five different species among our collection (excluding a NEQAS isolate with bla_{KPC-2}). While the majority were from *K. pneumoniae* (21/30; 70.0%), 13 different STs from this species were represented and no single ST accounted for more than four isolates.

Using our short-read data with the TETyper tool (Sheppard et al. 2018), we first determined whether the bla_{KPC-2} genes were carried on the Tn*4401* transposon, which has previously been associated with bla_{KPC} gene spread (Naas et al. 2008; Cuzon et al. 2011). In

ten isolates (10/30; 33.3%), TETyper did not identify any known Tn*4401* variant. However,
the remaining isolates (20/30; 66.7%), from a variety of species and STs, carried bla_{KPC-2} on
the Tn*4401a* variant. These were recovered from 16 patients between 2013 and 2020.
Tn*4401a* is the same variant associated with endemic spread of bla_{KPC-2}-encoding
Enterobacteriaceae in North-West England recognised from 2008 onwards (Stoesser et al.
2020). Notably, the majority (17/20; 85.0%) of our isolates with bla_{KPC-2} carried on Tn*4401a*
were sampled from patients in North Wales. These findings likely reflect acquisition of bla_{KPC-2}-encoding strains and/or plasmids in North-West England hospitals, consistent with the
2-encoding strains and/or plasmids in North-West England hospitals, consistent with the
typical referral of patients in North Wales to this region for specialist treatment.
To further elucidate the diversity of plasmids carrying bla_{KPC-2} on Tn*4401a,* we long-
read sequenced 13 of these isolates (including the NEQAS isolate) (**Supplementary Table
2**). These were selected to represent the diversity of species and STs observed. We
obtained complete plasmids carrying bla_{KPC-2} from hybrid (combined short- and long-read)
assemblies in all 13 isolates. One *K. pneumoniae* ST11 isolate (ARGID_32407) also carried
an additional copy of bla_{KPC-2} on the chromosome. Of the 13 bla_{KPC-2} plasmids, all possessed
either one or two IncF replicons, sometimes in combination with other replicon types (e.g.
IncR). They ranged in size from 113.5-210.3kb. In particular, we found that seven had a
similar structure and size to pKpQIL-like plasmids, a family of plasmids represented by
pKpQIL (Leavitt et al. 2010).
A visual comparison of pKpQIL-like plasmids from our collection, together with two
published pKpQIL-like plasmids from North-West England (Stoesser et al. 2020) and
pKpQIL (Leavitt et al. 2010), showed the high structural conservation overall (**Figure 4**).
However, two plasmids (from ARGID_31921 and ARGID_24826) had lower nucleotide
similarity with pKpQIL over one half, likely as a result of recombination. All others showed
high nucleotide similarity to pKpQIL, differing by 11-32 SNPs. We also found some support
for recent common ancestry between plasmids from North Wales and North-West England.
For example, the plasmids from ARGID_18508 and ARGID_31165 (both North Wales)
differed from trace384 (North-West England) by only 4 SNPs.
We next assessed the remaining ten bla_{KPC-2}-carrying isolates with no known Tn*4401*
variant. These isolates were from five patients and one environmental sample, and from a
mixture of species and STs. They were recovered between 2017 and 2020 from three
hospitals in South Wales, with seven isolates from Hospital A. Inspection of the short-read
genome assemblies demonstrated a shared genetic context around bla_{KPC-2}. We also found
that all ten isolates had an IncN replicon in common. To investigate the possibility of bla_{KPC-2}-
IncN plasmid spread, we long-read sequenced six of these isolates from different species
and/or STs (**Supplementary Table 2**).
We obtained a complete IncN plasmid from the hybrid assemblies of all six isolates,
and confirmed carriage of bla_{KPC-2} in a conserved genetic environment (next to bla_{TEM-1}) on
these plasmids. The IncN plasmids varied considerably in size from 48.2kb to 137.4kb, and
the largest also carried an additional IncFII replicon. Visual comparison of the plasmids
showed that they shared high nucleotide similarity across shared "core" regions (**Figure 5**).
This is despite frequent recombination events leading to alternate structural arrangements
and a large variation in "accessory" sequence. We also found high nucleotide similarity
among the shared regions of these plasmids and a previously described IncN plasmid from
an ST307 isolate recovered in the UK (accession KY271414) (Villa et al. 2017) (**Figure 5**).
These findings point to the recent emergence and spread of this bla_{KPC-2}-encoding IncN
backbone between different *Klebsiella* species and STs in South Wales and the potentially
wider UK region.

467
468
469 **Frequent carriage and horizontal dissemination of $bla_{\text{OXA-48-like}}$ genes via pOXA-**
470 **48-like plasmids and related variants**
471

The most common type of carbapenemase genes among our *Klebsiella* collection
were $bla_{\text{OXA-48-like}}$ genes, which comprised $bla_{\text{OXA-244}}$ (*n*=58), $bla_{\text{OXA-48}}$ (*n*=46), $bla_{\text{OXA-181}}$ (*n*=7)
and $bla_{\text{OXA-232}}$ (*n*=3) variants. While most $bla_{\text{OXA-244}}$-encoding isolates (49/58; 84.5%)
belonged to the ST307 lineage, other $bla_{\text{OXA-48-like}}$ genes were identified in diverse species
and STs. In particular, $bla_{\text{OXA-48}}$ genes were found in three species (*K. pneumoniae*, *K.*
quasipneumoniae, *K. oxytoca*) and 15 different STs of *K. pneumoniae*, and recovered
between 2013 and 2020.
We investigated a possible association of $bla_{\text{OXA-48-like}}$ genes with pOXA-48-like
plasmids, which are highly conserved ~61-63kb IncL plasmids shown previously to
frequently carry these carbapenemase genes (Poirel et al. 2012; David et al. 2020). To do
this, we mapped the short reads of all isolates in our collection (including those without
$bla_{\text{OXA-48-like}}$ genes) to the 61.8kb pOXA48a reference plasmid (Poirel et al. 2012). 113/540
(20.9%) isolates mapped to ≥70% of the plasmid length (including 95 that mapped to ≥90%),
suggesting that they harboured the same or a similar plasmid. Most of the remaining isolates
mapped to ≤10%. The 113 isolates with ≥70% plasmid mapping were from 61 different
patients in 15 hospitals and additional GP surgeries, largely from across South Wales. They
accounted for most isolates with $bla_{\text{OXA-48-like}}$ genes, including all of those with $bla_{\text{OXA-244}}$
(58/58) and $bla_{\text{OXA-48}}$ (46/46), and 1/3 (33.3%) of those with $bla_{\text{OXA-232}}$. Only 8/113 (7.1%)
possessed no $bla_{\text{OXA-48-like}}$ gene. Among the 113 isolates, 94 (83.2%) possessed an IncL
replicon (the same as pOXA48a). We did not find an IncL replicon in the remaining 19/113
(16.8%) isolates, but instead found either an IncM1 (*n*=18) or IncM2 (*n*=1) replicon. IncL and
IncM plasmids are known to be closely related and indeed were previously classified in the
same family, IncL/M (Carattoli et al. 2015). However, the latter have not been implicated as
vectors of $bla_{\text{OXA-48-like}}$ genes to date.
Analysis of SNP diversity among the 113 mapped plasmid sequences demonstrated
0-5 differences among many (≥75%) of the sequences, despite their recovery from diverse
locations and times. This high similarity was reflected in a minimum spanning tree (**Figure**
6A). Only some sequences were considerably more divergent, including those associated
with IncM replicons (**Figure 6B**). While the overall low amount of SNP diversity among these
plasmid sequences makes inference of transmission events difficult, we did find identical
plasmid sequences shared among different species and STs, suggestive of recent horizontal
spread. For example, we found 0 SNPs among 55 sequences (all associated with $bla_{\text{OXA-244}}$),
which included 48 from the *K. pneumoniae* ST307 outbreak isolates, two from ST1430, two
from ST48, two from ST353 and one from *K. oxytoca* (**Figure 6C**). The vast majority of
these, including the non-ST307s, were recovered from Hospital A, supporting the possibility
of horizontal transmission. Notably, five of these non-ST307s were recovered in 2017-2018,
implying local circulation of this plasmid variant prior to the expansion of $bla_{\text{OXA-244}}$-encoding
ST307 in 2019.
Finally, we used the short-read data to investigate structural differences among the
113 mapped plasmid sequences. To do this, we calculated and visualised the short-read
coverage across the length of pOXA48a, relative to the median chromosomal coverage of
the host strain (**Figure 7**). This revealed that nine of the $bla_{\text{OXA-244}}$-encoding ST307 outbreak

isolates lacked coverage across parts of the plasmid (8.6-16.6kb) that include the *tra* region
in most cases, with a conserved breakpoint at one end of the deletion. This suggests that
these plasmids may have lost the ability to self-conjugate and/or be mobilised. Notably, they
did not cluster into a single clade in the core genome-based phylogenetic tree, suggesting
that the deletions occurred on multiple independent occasions (**Figure 2**).
519
520

521 **Structural evolution of pOXA-48-like plasmids including repeated loss of**
522 **conjugative ability in *K. pneumoniae* ST307**
523

We next used our short-read analysis of the SNP and structural diversity among
pOXA48-like plasmids to guide selection of 19 isolates for further investigation with long-
read sequencing (**Supplementary Table 2**). These were chosen to investigate: a) the
possible carriage of $bla_{OXA-48-like}$ genes on IncM plasmids and their structural differences with
IncL plasmids; b) other divergent variants of pOXA-48-like plasmids and c) the evolution of
pOXA-48-like plasmids in the ST307 outbreak lineage. We constructed hybrid assemblies
and obtained complete IncL or IncM1 plasmids in all 19 isolates, which ranged in size from
46.8kb to 73.0kb.
Among the 19 complete plasmids, we obtained three with an IncM1 replicon and
confirmed carriage of bla_{OXA-48} genes on these. Overall they showed high structural
homology to the IncL plasmids albeit with key differences (**Figure 8**). These included a
section with the replicon typing determinants that lacks homology, as well as lower
sequence similarity over the *tra* gene cluster region. However, large parts of the IncL and
IncM1 plasmids shared 100% nucleotide similarity, including the region containing bla_{OXA-48},
suggestive of recent recombination.
Among other isolates selected for long-read sequencing based on the high structural
and/or SNP divergence of their pOXA-48-like plasmids from pOXA48a, we found several
with novel integrations (**Figure 8**). These included an island harbouring $bla_{CTX-M-15}$ in one
plasmid recovered from a *K. pneumoniae* ST307 strain (unrelated to the ST307 outbreak
strain). Furthermore, a genetic island with another ESBL gene, $bla_{CTX-M-14b}$, as well as
aminoglycoside resistance genes, was found in both IncL and IncM1 plasmids recovered
from *K. pneumoniae* ST101, ST383 and ST750 isolates.
Finally, we analysed complete pOXA-48-like plasmids from six ST307 outbreak-
associated isolates to investigate the putative deletions involving the *tra* region. These
included ARGID_31642, in which no deletion had been observed, and five isolates with
putative deletions. As expected, we found that the plasmid from ARGID_31642 had a similar
structure and size to pOXA48a (**Figure 9**). All others possessed a deletion with a conserved
breakpoint at one end adjacent to an IS*1R* element. The variable size and nature of these
deletions further confirmed that they had occurred on multiple independent occasions. This
suggests that the deletions confer a fitness advantage. The deletions involved one or more
tra genes and the *oriT* region in four of the five cases, with one plasmid (from
ARGID_33503) retaining these regions despite a deletion. We used conjugation experiments
to test the ability of the six isolates with complete pOXA-48-like sequences to transfer the
plasmid to both *E. coli* (J53 strain) and *K. pneumoniae* (71.1 strain), using two different
ratios of donor/recipient (1:3 and 5:1). These confirmed that both ARGID_31642 and
ARGID_33503 could transfer the plasmid, while no transfer of plasmids with deletions in the
tra/*oriT* regions was observed (**Supplementary Table 4**).

Discussion

In line with other geographic regions worldwide, *Klebsiella* infections represent a
substantial healthcare burden in Wales and are the third most common cause of
bloodstream infections (PHW, 2018). While rates of resistance to carbapenems remain low
(<1% among bloodstream isolates in 2017) (PHW, 2018), the rapid global rise of resistance
necessitates vigilant monitoring (Cassini et al. 2019; Castanheira et al. 2019). Here we
describe the major strains and plasmids driving spread of multi-drug resistant *Klebsiella* in
Wales between 2007 and 2020, providing a detailed view of resistance dynamics in a largely
non-endemic setting.
We found that more than half of carbapenemase-producing *Klebsiella* isolates
(124/207; 59.9%) referred to our laboratory at PHW belonged to just nine STs from *K.*
pneumoniae. This finding echoes numerous other studies reporting a concentration of
resistant *Klebsiella* in "high-risk" STs (David et al. 2019; Bonnin et al. 2020; Di Pilato et al.
2021). The major carbapenemase-producing clones detected in our region include the
globally important *K. pneumoniae* strains, ST307, ST20 and ST15, which carried $bla_{OXA-244}$,
bla_{IMP-4} and bla_{VIM-4}, respectively. We also identified spread of a strain which has rarely been
reported elsewhere, ST1788, carrying $bla_{CTX-M-15}$ and which also acquired bla_{OXA-48} in one
patient. Detailed phylogenetic analyses of the ST307 and ST1788 clones revealed that a
single major introduction of each into Wales, followed by local spread within and between
hospitals, accounted for the dominance of these clones among patient isolates.
Our phylogenetic analyses also suggested that the ST307 strain had been circulating
in Wales for several years prior to its acquisition of $bla_{OXA-244}$ and an outbreak involving the
$bla_{OXA-244}$-encoding subclade centred on Hospital A in 2019. The lack of recognition of ST307
by SACU until the 2019 outbreak reflects the lack of active surveillance or typing of third-
generation cephalosporin-resistant *Klebsiella* infections in Wales over this time period, and
the absence of clear epidemiological links between infections. Since the outbreak involving
the $bla_{OXA-244}$-encoding subclade was declared closed, it is possible that the ST307 strain
(without $bla_{OXA-244}$) continues to disseminate and be responsible for a significant proportion of
invasive infections. However, in the absence of epidemiological links and active surveillance
of all isolates, this once more remains undetected. Yet past reports demonstrate the ability
of ST307 to readily acquire additional resistance determinants and expand rapidly (Lowe et
al. 2019; Haller et al. 2019). Thus, as our laboratory and others move towards using WGS
data for routine surveillance of *Klebsiella*, this experience demonstrates the importance of
urgently tailoring our systems to detect and control the spread of high-risk clones, ideally
prior to acquisition of additional resistance determinants.
Our analysis also revealed the dominant plasmid vectors driving spread of bla_{KPC-2}
and $bla_{OXA-48-like}$ genes in Wales, which were found among diverse species and strain
backgrounds. We found that bla_{KPC-2} genes from North and South Wales were largely
encoded on different plasmid backgrounds, consistent with the referral of patients from these

regions to different hospitals for their tertiary medical care (i.e. to North-West England and
South Wales, respectively). Notably, bla_{KPC-2} genes from isolates in North Wales collected
from 2013 onwards were typically encoded within the Tn*4401a* transposon on IncF
plasmids, the same genetic background identified among bla_{KPC-2} isolates from North-West
England. This region has experienced endemic spread of bla_{KPC-2}-encoding
Enterobacteriaceae since 2008, driven largely by horizontal exchange of plasmids and
plasmid fragments between strains and species (Stoesser et al. 2020). Thus our findings
likely reflect the outward expansion of this complex, multi-species, plasmid-driven outbreak
to Wales via hospital networks.
By contrast, bla_{KPC-2} genes found among isolates from South Wales since 2017 were
typically harboured by IncN plasmids. bla_{KPC-2}-encoding IncN plasmids have been only rarely
reported (Villa et al. 2017) and likely represent an emerging plasmid background transmitting
these genes. Despite their large variation in size and structure, the high nucleotide similarity
found among their shared regions strongly suggests the existence of a recent common
ancestor and thereby the occurrence of local plasmid transmission events between different
species and STs in Welsh hospitals. This genomic evidence is further supported by the
spatial clustering of these plasmids (i.e. 7/10 were recovered from a single hospital). As
exemplified with these IncN plasmids, the large structural variability among some plasmid
types (despite recent common ancestry) heightens the challenge of routinely identifying
plasmid transmissions, even with complete plasmid sequences. We propose that calculating
nucleotide similarity across individual segments (or sliding windows) of plasmid sequences,
rather than determining an average over the entire sequence, is crucial for detecting
transmissions involving plasmids that can rapidly acquire or lose accessory regions, and/or
exchange modular regions with other backbone types.
We found that most $bla_{OXA-48-like}$ genes (105/114; 92.1%) encoded by isolates referred
to our laboratory were harboured by pOXA-48-like plasmids. These accounted for the clonal
spread of $bla_{OXA-244}$ by the ST307 outbreak strain and also contributed to the horizontal
spread of both bla_{OXA-48} and $bla_{OXA-244}$ variants across numerous species and STs. Other
countries and regions have reported a similar dominance of these plasmids, which have
been found to be remarkably conserved over time and space (Skalova et al. 2017; David et
al. 2020). Given their importance, we propose that the identification and typing of pOXA-48-
like plasmids by genomic surveillance systems would have large gains for detection and
control of resistance spread. Due to the high structural conservation of this plasmid family,
this is largely achievable using short-read approaches (e.g. mapping or plasmid MLST
(Brehony et al. 2019)).
Our analysis did, however, reveal a small proportion of novel divergent variants of
pOXA-48-like plasmids. These included those carrying additional resistance genes as well
as those associated with IncM, rather than IncL, replicons. We also detected deletions
among pOXA-48-like plasmids in the ST307 outbreak lineage which extended to the *oriT*
and *tra* regions in most cases. Experiments confirmed that the plasmids lacking these
regions could no longer be transferred via conjugation to a recipient *K. pneumoniae* or *E.*
coli strain. Since these deletions occurred multiple times independently, we propose that
they represent adaptation of the pOXA-48-like plasmid to carriage by ST307 strains, likely
via a reduction in the fitness cost imposed on the host.
Altogether this study provides a comprehensive picture of resistance gene spread
among *Klebsiella* species in Wales, mediated via both clonal and horizontal transmission. It
provides important contextual data for ongoing surveillance as well as key considerations for

the design and optimisation of routine genomic surveillance systems in public health
laboratories.
656
657

Authors and contributors

659
660 SD, MM, KS, MW and LJ conceived the study. SD performed the bioinformatic data analysis
661 and drafted the manuscript. MM, KS, EP, LG and JW performed the sequencing, laboratory
662 experiments and data analysis. OBS, DMA, MW and LJ supervised the study. All authors
663 reviewed and edited the manuscript.
664
665

Conflicts of interest

667
668 The authors declare no conflicts of interest.
669
670

Funding information

SD and DMA are supported by funding from the Centre for Genomic Pathogen Surveillance
and Li Ka Shing Foundation.
Illumina sequencing was funded as part of the Antimicrobial Resistance and Genomic
Typing Project (ARGENT) by the Welsh Assembly Government.
MinION sequencing was funded by the Public Health Wales R&D Pump-Priming Fund –
2018/19

679
680

Acknowledgements

We thank the Pathogen Informatics team at the Wellcome Sanger Institute for informatics
support; the Pathogen Genomics Unit laboratory and bioinformatics teams at Public Health
Wales; the team at the Specialist Antimicrobial Chemotherapy Unit at Public Health Wales;
the Infection Prevention & Control departments for Health Boards, especially those teams
which commented on the local outbreaks reported herein; and referring Microbiology
Laboratories from across Wales and their host Health Boards.

689
690

References

1. Magill SS, Edwards JR, Bamberg W, Beldavs ZG, Dumyati G, Kainer MA, et al.
Multistate point-prevalence survey of health care-associated infections. N Engl J Med.
2014;370(13):1198-208.

2. Ling ML, Apisarnthanarak A, Madriaga G. The Burden of Healthcare-Associated
Infections in Southeast Asia: A Systematic Literature Review and Meta-analysis. Clin
Infect Dis. 2015;60(11):1690-9.

3. Vincent JL, Sakr Y, Singer M, Martin-Loeches I, Machado FR, Marshall JC, et al.
Prevalence and Outcomes of Infection Among Patients in Intensive Care Units in 2017.
JAMA. 2020;323(15):1478-87.

4. Meatherall BL, Gregson D, Ross T, Pitout JD, Laupland KB. Incidence, risk factors, and
outcomes of *Klebsiella pneumoniae* bacteremia. Am J Med. 2009;122(9):866-73.

5. Podschun R, Ullmann U. *Klebsiella spp.* as nosocomial pathogens: epidemiology,
taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev.
1998;11(4):589-603.

6. Gorrie CL, Mirceta M, Wick RR, Edwards DJ, Thomson NR, Strugnell RA, et al.
Gastrointestinal Carriage Is a Major Reservoir of *Klebsiella pneumoniae* Infection in
Intensive Care Patients. Clin Infect Dis. 2017;65(2):208-15.

7. Snitkin ES, Zelazny AM, Thomas PJ, Stock F, Henderson DK, Palmore TN, et al.
Tracking a hospital outbreak of carbapenem-resistant *Klebsiella pneumoniae* with
whole-genome sequencing. Sci Transl Med. 2012;4(148):148ra16.

8. Gu D, Dong N, Zheng Z, Lin D, Huang M, Wang L, et al. A fatal outbreak of ST11
carbapenem-resistant hypervirulent *Klebsiella pneumoniae* in a Chinese hospital: a
molecular epidemiological study. Lancet Infect Dis. 2018;18(1):37-46.

9. Tavoschi L, Forni S, Porretta A, Righi L, Pieralli F, Menichetti F, et al. Prolonged
outbreak of New Delhi metallo-beta-lactamase-producing carbapenem-resistant
Enterobacterales (NDM-CRE), Tuscany, Italy, 2018 to 2019. Euro Surveill. 2020;25(6).

10. Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, et al.
Attributable deaths and disability-adjusted life-years caused by infections with
antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a
population-level modelling analysis. Lancet Infect Dis. 2019;19(1):56-66.

11. Castanheira M, Deshpande LM, Mendes RE, Canton R, Sader HS, Jones RN.
Variations in the Occurrence of Resistance Phenotypes and Carbapenemase Genes
Among *Enterobacteriaceae* Isolates in 20 Years of the SENTRY Antimicrobial
Surveillance Program. Open Forum Infect Dis. 2019;6(Suppl 1):S23-S33.

12. World Health Organisation. Global priority list of antibiotic-resistant bacteria to guide
research, discovery, and development of new antibiotics. 2017.

13. Queenan AM, Bush K. Carbapenemases: the versatile beta-lactamases. Clin Microbiol
Rev. 2007;20(3):440-58.

14. Bradley N, Lee Y. Practical Implications of New Antibiotic Agents for the Treatment of
Carbapenem-Resistant *Enterobacteriaceae*. Microbiol Insights.
2019;12:1178636119840367.

15. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a
new genome assembly algorithm and its applications to single-cell sequencing. J
Comput Biol. 2012;19(5):455-77.

16. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. PLoS Comput Biol.
2017;13(6):e1005595.

17. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068-9.

18. Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. A genomic
surveillance framework and genotyping tool for *Klebsiella pneumoniae* and its related
species complex. Nat Commun. 2021;12(1):4188.
19. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, et al. Identification
of *Klebsiella* capsule synthesis loci from whole genome data. Microb Genom.
2016;2(12):e000102.
20. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing
polished prokaryotic pangenomes with the Panaroo pipeline. Genome Biol.
2020;21(1):180.
21. Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, et al.: rapid efficient
extraction of SNPs from multi-FASTA alignments. Microb Genom. 2016;2(4):e000056.
22. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.
2015;32(1):268-74.
23. Argimón S, David S, Underwood A, Abrudan M, Wheeler NE, Kekre M, et al. Rapid
genomic characterisation and global surveillance of Klebsiella using Pathogenwatch.
Clinical Infectious Diseases. 2021;73:S325:S335.
24. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics. 2009;25(14):1754-60.
25. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
26. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid
phylogenetic analysis of large samples of recombinant bacterial whole genome
sequences using Gubbins. Nucleic Acids Res. 2015;43(3):e15.
27. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Res. 2021;49(W1):W293-W6.
28. Argimón S, Abudahab K, Goater RJ, Fedosejev A, Bhai J, Glasner C, et al. Microreact:
visualizing and sharing data for genomic epidemiology and phylogeography. Microb
Genom. 2016;2(11):e000093.
29. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, et al. *In
silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus
sequence typing. Antimicrob Agents Chemother. 2014;58(7):3895-903.
30. Li X, Xie Y, Liu M, Tai C, Sun J, Deng Z, et al. oriTfinder: a web-based tool for the
identification of origin of transfers in DNA sequences of bacterial mobile genetic
elements. Nucleic Acids Res. 2018;46(W1):W229-W34.
31. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
Bioinformatics. 2011;27(7):1009-10.
32. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST
plus: architecture and applications. Bmc Bioinformatics. 2009;10.
33. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al. Versatile
and open software for comparing large genomes. Genome Biology. 2004;5(2).
34. Sheppard AE, Stoesser N, German-Mesner I, Vegesana K, Sarah Walker A, Crook
DW, et al. TETyper: a bioinformatic pipeline for classifying variation and genetic
contexts of transposable elements from short-read whole-genome sequencing data.
Microb Genom. 2018;4(12).
35. Poirel L, Bonnin RA, Nordmann P. Genetic features of the widespread plasmid coding
for the carbapenemase OXA-48. Antimicrob Agents Chemother. 2012;56(1):559-62.

36. Zhou Z, Alikhan NF, Sergeant MJ, Luhmann N, Vaz C, Francisco AP, et al. GrapeTree:
visualization of core genomic relationships among 100,000 bacterial pathogens.
Genome Res. 2018;28(9):1395-404.
37. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2.
38. Peirano G, Chen L, Kreiswirth BN, Pitout JDD. Emerging Antimicrobial-Resistant High-
Risk *Klebsiella pneumoniae* Clones ST307 and ST147. Antimicrob Agents Chemother.
2020;64(10).
39. Martínez-Martínez L, Pascual A, Hernández-Allés S, Alvarez-Díaz D, Suárez AI, Tran
J, et al. Roles of beta-lactamases and porins in activities of carbapenems and
cephalosporins against *Klebsiella pneumoniae*. Antimicrob Agents Chemother.
1999;43(7):1669-73.
40. Hamzaoui Z, Ocampo-Sosa A, Maamar E, Fernandez Martinez M, Ferjani S, Hammami
S, et al. An Outbreak of NDM-1-Producing *Klebsiella pneumoniae*, Associated with
OmpK35 and OmpK36 Porin Loss in Tunisia. Microb Drug Resist. 2018;24(8):1137-47.
41. Shon AS, Bajwa RP, Russo TA. Hypervirulent (hypermucoviscous) *Klebsiella
pneumoniae*: a new and dangerous breed. Virulence. 2013;4(2):107-18.
42. Naas T, Cuzon G, Villegas MV, Lartigue MF, Quinn JP, Nordmann P. Genetic
structures at the origin of acquisition of the beta-lactamase blaKPC gene. Antimicrob
Agents Chemother. 2008;52(4):1257-63.
43. Cuzon G, Naas T, Nordmann P. Functional characterization of Tn*4401*, a Tn3-based
transposon involved in blaKPC gene mobilization. Antimicrob Agents Chemother.
2011;55(11):5370-3.
44. Stoesser N, Phan HTT, Seale AC, Aiken Z, Thomas S, Smith M, et al. Genomic
epidemiology of complex, multi-species, plasmid-borne blaKPC carbapenemase in
Enterobacterales in the United Kingdom from 2009 to 2014. Antimicrob Agents
Chemother. 2020;64(5):e02244.
45. Leavitt A, Chmelnitsky I, Carmeli Y, Navon-Venezia S. Complete nucleotide sequence
of KPC-3-encoding plasmid pKpQIL in the epidemic *Klebsiella pneumoniae* sequence
type 258. Antimicrob Agents Chemother. 2010;54(10):4493-6.
46. Villa L, Feudi C, Fortini D, Brisse S, Passet V, Bonura C, et al. Diversity, virulence, and
antimicrobial resistance of the KPC-producing *Klebsiella pneumoniae* ST307 clone.
Microb Genom. 2017;3(4):e000110.
47. David S, Cohen V, Reuter S, Sheppard AE, Giani T, Parkhill J, et al. Integrated
chromosomal and plasmid sequence analyses reveal diverse modes of
carbapenemase gene spread among *Klebsiella pneumoniae*. Proc Natl Acad Sci U S
A. 2020;117(40):25043-54.
48. Carattoli A, Seiffert SN, Schwendener S, Perreten V, Endimiani A. Differentiation of
IncL and IncM Plasmids Associated with the Spread of Clinically Relevant Antimicrobial
Resistance. PLoS One. 2015;10(5):e0123063.
49. Antibacterial resistance in Wales 2008-2017. Public Health Wales. 2018.
50. David S, Reuter S, Harris SR, Glasner C, Feltwell T, Argimon S, et al. Epidemic of
carbapenem-resistant *Klebsiella pneumoniae* in Europe is driven by nosocomial
spread. Nat Microbiol. 2019;4(11):1919-29.
51. Bonnin RA, Jousset AB, Chiarelli A, Emeraud C, Glaser P, Naas T, et al. Emergence of
New Non-Clonal Group 258 High-Risk Clones among *Klebsiella pneumoniae*
Carbapenemase-Producing *K. pneumoniae* Isolates, France. Emerg Infect Dis.
2020;26(6):1212-20.

52. Di Pilato V, Errico G, Monaco M, Giani T, Del Grosso M, Antonelli A, et al. The
changing epidemiology of carbapenemase-producing *Klebsiella pneumoniae* in Italy:
toward polyclonal evolution with emergence of high-risk lineages. J Antimicrob
Chemother. 2021;76(2):355-61.
53. Lowe M, Kock MM, Coetzee J, Hoosien E, Peirano G, Strydom KA, et al. *Klebsiella*
pneumoniae ST307 with blaOXA-181, South Africa, 2014-2016. Emerg Infect Dis.
2019;25(4):739-47.
54. Haller S, Kramer R, Becker K, Bohnert JA, Eckmanns T, Hans JB, et al. Extensively
drug-resistant *Klebsiella pneumoniae* ST307 outbreak, north-eastern Germany, June to
October 2019. Euro Surveill. 2019;24(50).
55. Skalova A, Chudejova K, Rotova V, Medvecky M, Studentova V, Chudackova E, et al.
Molecular Characterization of OXA-48-Like-Producing *Enterobacteriaceae* in the Czech
Republic and Evidence for Horizontal Transfer of pOXA-48-Like Plasmids. Antimicrob
Agents Chemother. 2017;61(2).
56. Brehony C, McGrath E, Brennan W, Tuohy A, Whyte T, Brisse S, et al. An MLST
approach to support tracking of plasmids carrying OXA-48-like carbapenemase. J
Antimicrob Chemother. 2019;74(7):1856-62.

853
854
855
856 **Figure legends**
857
Figure 1. Maximum likelihood phylogenetic tree of 421 *K. pneumoniae* isolates
demonstrates high diversity. The tree was constructed from an alignment of 4034 core
genes that were present in ≥99% of *K. pneumoniae* isolates referred to the SACU laboratory
at PHW. The most frequently observed STs are highlighted. Metadata columns show the
ESBL and carbapenemase variants found among the isolates (if applicable). The scale
indicates the number of SNPs per variable site. The tree is available to view interactively and
with additional metadata at https://microreact.org/project/aSJqpc9MZVZcurTWtoadW8-k-
pneumoniae-phw-n421
866
Figure 2. A) Maximum likelihood phylogenetic tree of 1052 *K. pneumoniae* ST307
isolates demonstrates a main introduction of ST307 into Wales. The tree includes 964
ST307 isolates from public archives (from diverse international locations) and 88 isolates
from the PHW collection. The tree was constructed from an alignment generated by
mapping all short sequence reads to an ST307 reference genome (ARGID_32304) and after
the removal of recombined regions. 82 ST307 isolates from the PHW collection form a
monophyletic cluster highlighted in red. Metadata columns show the carbapenemase, ESBL
and yersiniabactin variants among all isolates (if applicable). The tree is available to view
interactively and with additional metadata at
https://microreact.org/project/1ZRxgto9HVsaV6qRHAuf6P-st307s-phw-n1052 **B)** Cluster of
82 ST307 isolates from the PHW collection, as zoomed in from (A). Filled circles of the same
colour on the tree tips represent isolates from the same patient. Metadata columns show the
sampling hospital, sample type, carbapenemase variant and the presence/absence of the
pOXA-48-like plasmid. For the latter, the plasmid was scored as "present" if sequence reads
mapped to ≥90% of the pOXA48a reference plasmid (accession JN626286), "partial" if

between 70% and 90% and "absent" if <70%. The scale bars in (A) and (B) represent the
number of SNPs. **C)** Timeline showing the isolation dates of the 82 clustered ST307 isolates
from the PHW collection. Bars are coloured by the carbapenemase variant (if applicable).
885
Figure 3. Maximum likelihood phylogenetic tree of 45 ST1788 isolates demonstrates
high clonality among isolates from Wales. The tree includes two isolates from public
archives (both from Nigeria) and 43 from the PHW collection. It was constructed from an
alignment generated by mapping all short sequence reads to an ST1788 reference genome
(ARGID_33748) and after the removal of recombined regions. The 43 ST1788 isolates from
the PHW collection form a monophyletic cluster. Filled circles of the same colour on the tree
tips represent isolates from the same patient. Metadata columns show the sampling hospital,
sample type, and carbapenemase, ESBL and yersiniabactin variants (if applicable). The
scale bar represents the number of SNPs. The tree is available to view interactively and with
additional metadata at https://microreact.org/project/rMY2fakRxkNKXqqBPs3is6-st1788-
phw-n45
897
Figure 4. Pairwise comparisons between complete *bla*$_{KPC}$-encoding pKpQIL-like
plasmids demonstrate high structural and sequence conservation. The plasmids
include the original pKpQIL sequence (accession GU595196) (Leavitt et al. 2010), seven
from the PHW collection and two previously sequenced plasmids recovered in North-West
England hospitals (Stoesser et al. 2020). Open reading frames (ORFs) are depicted as
arrows with resistance genes (including *bla*$_{KPC}$) labelled. Grey blocks between plasmid pairs
indicate regions of homology. The colour of the ORFs represents the percentage nucleotide
similarity to pKpQIL via BLASTn.
906
Figure 5. A) Pairwise comparisons between complete *bla*$_{KPC-2}$-encoding IncN plasmids
demonstrate substantial plasticity despite high nucleotide similarity among shared
backbone regions. The plasmids include a previously published sequence (accession
KY271414) (Villa et al. 2017) and six from the PHW collection. ORFs are depicted as arrows
with resistance genes (including *bla*$_{KPC-2}$) labelled. Grey blocks between plasmid pairs
indicate regions of homology. The colour of the ORFs represents the percentage nucleotide
similarity to the plasmid from ARGID_28504 via BLASTn. **B)** Conserved genetic
environment of *bla*$_{KPC-2}$ among the IncN plasmids.
915
Figure 6. A) Minimum spanning tree of 113 pOXA-48-like plasmid sequences
demonstrates high nucleotide conservation as well as some divergent sequences.
Plasmid sequences included in the tree, which were obtained by mapping short reads to
pOXA48a (accession JN626286) (Poirel et al. 2012), are from isolates with ≥70% mapping
to this reference sequence. The tree was constructed from a SNP distance matrix. The
branch lengths are displayed using a log scale and indicated on the branches. The nodes
are partitioned by (and their sizes proportional to) the number of represented isolates.
Isolates are coloured by ST with the number of isolates from each ST indicated in square
brackets. Isolates marked with a black filled circle were selected for long-read sequencing to
enable complete plasmid sequencing. **B)** The same trees as in (A) coloured by species of
the host isolate or replicon type identified from the genome. **C)** The major node from the tree
in (A) coloured by ST, with annotations marking the sampling hospitals.
928

Figure 7. Heatmap showing the short-read coverage across the pOXA48a reference
sequence indicates deletions among some isolates, including several involving the
***tra* region in ST307.** The plasmid coverage is shown relative to the median chromosomal
coverage for 113 isolates with ≥70% mapping to pOXA48a (accession JN626286) (Poirel et
al. 2012). ORFs from the pOXA48a plasmid are depicted at the top, including bla_{OXA-48}.
Metadata columns indicate the host species, those belonging to frequently observed STs,
and replicon and $bla_{OXA-48-like}$ variants identified from the isolate genomes. The far-right
column indicates isolates selected for long-read sequencing to enable complete plasmid
assembly.
938
Figure 8. Pairwise comparisons between complete pOXA-48-like plasmids show
structural differences to the "classic" backbone of pOXA48a, including insertion of
islands carrying resistance genes. The plasmids include pOXA48a (accession JN626286)
(Poirel et al. 2012) and five from the PHW collection. They are from a range of STs and
include those with both IncL and IncM1 replicon types. ORFs are depicted as arrows with
resistance genes (including bla_{OXA-48}) labelled. Grey blocks between plasmid pairs indicate
regions of homology. The colour of the ORFs represents the percentage nucleotide similarity
to pOXA48a via BLASTn.
947
Figure 9. Pairwise comparisons between complete pOXA-48-like plasmids from ST307
demonstrate multiple independent deletions of regions adjacent to an IS*1R* element.
One plasmid (from ARGID_31642), which possesses the "classic" pOXA-48-like backbone
and which lacks these deletions, is included. ORFs are depicted as arrows with resistance
genes (including bla_{OXA-48}/$bla_{OXA-244}$) labelled. Grey blocks between plasmid pairs indicate
regions of homology. ORFs coloured blue are predicted to be involved in conjugation while
those in red have other predicted functions. The *oriT* region is indicated if identified in the
plasmid.
956
957
958

959 ## Supplementary Tables

960
961 **Supplementary Table 1.** Metadata, assembly statistics and genotyping results for 540
962 short-read sequenced *Klebsiella* isolates included in this study.
963
964 **Supplementary Table 2.** Metadata and hybrid assembly statistics for 38 long-read
965 sequenced *Klebsiella* isolates included in this study.
966
967 **Supplementary Table 3.** URLs for interactive Microreact projects containing phylogenetic
968 analyses for each *Klebsiella* species (those with ≥5 isolates only) together with all metadata
969 and genotypic data.
970
971 **Supplementary Table 4.** Conjugation frequencies determined from mating cultures
972 comprising ST307 donor isolates harbouring a $bla_{OXA-48-like}$-encoding pOXA-48-like plasmid
973 and a recipient *K. pneumoniae* or *E. coli* strain.

A

B

C

Carbapenemase

None	*bla*OXA-181
*bla*CTX-M-33	*bla*OXA-244
*bla*GES-5	*bla*OXA-48
*bla*KPC-2	*bla*KPC-2 & *bla*NDM-7
*bla*KPC-3	*bla*KPC-2 & *bla*OXA-48
*bla*NDM-1	*bla*KPC-3 & *bla*OXA-48
*bla*NDM-5	*bla*NDM-1 & *bla*OXA-48
*bla*NDM-7	*bla*NDM-5 & *bla*OXA-232
*bla*OXA-162	

ESBL

- None
- *bla*CTX-M-14
- *bla*CTX-M-15
- *bla*CTX-M-63
- *bla*CTX-M-156
- *bla*CTX-M-15 & *bla*TEM-19
- *bla*CTX-M-15 & *bla*TEM-116

Yersiniabactin

- None
- *ybt*9; ICE*Kp*3
- *ybt*10; ICE*Kp*4
- *ybt*13; ICE*Kp*2
- *ybt*14; ICE*Kp*5
- *ybt*15; ICE*Kp*11
- *ybt*16; ICE*Kp*12

Hospital

- GP surgery
- Hospital A
- Hospital B
- Hospital C
- Hospital D
- Hospital E
- Hospital F
- Hospital G
- Hospital H
- Hospital I
- Hospital J

Sample type

- Clinical
- Screen
- Environmental

pOXA-48-like plasmid

- Absent
- Present
- Partial

Wales cluster

50 SNPs

20 SNPs

x10

pKpQIL, 113.6kb (*Kp* ST258)
Israel, 2010

trace384, 113.6kb (*Kp* ST1828)
NW England, 2011

ARGID_31921, 115.6kb (*Kp* ST41-4LV)
Hospital M, Wales, 2019

ARGID_31680, 114.4kb (*K. michiganensis*)
Hospital N, Wales, 2019

ARGID_18508, 114.5kb (*Kp* ST25)
Hospital N, Wales, 2014

ARGID_31165, 116.1kb (*K. michiganensis*)
Hospital N, Wales, 2019

ARGID_22065, 113.5kb (*Kp* ST922)
Hospital M, Wales, 2015

ARGID_12502, 113.6kb (*Kp* ST258)
NEQAS strain

ARGID_24826, 116.1kb (*Kp* ST15)
Hospital C, Wales, 2017

trace41, 113.6kb (*Kp* ST588)
NW England, 2010

Nucleotide similarity
to pKpQIL

No BLAST hit <95 95-97 97-98 98-98.5 98.5-99 99-99.2 99.2-99.4 99.4-99.6 99.6-99.8 99.8-100 100

A

ARGID_32776, 109.0kb (*K. michiganensis*)
Hospital A, Wales, 2019

KY271414, 52.8kb (*Kp* ST307)
UK

ARGID_31501, 63.4kb (*Kp* ST54)
Hospital O, Wales, 2019

ARGID_28504, 62.6kb (*Kp* ST48)
Hospital A, Wales, 2018

ARGID_27030, 48.2kb (*Kp* ST48-3LV)
Hospital A, Wales, 2017

ARGID_25120, 137.4kb (*Kp* ST461)
Hospital E, Wales, 2017

ARGID_32889, 63.4kb (*K. variicola*)
Hospital A, Wales, 2020

Nucleotide similarity
to ARGID_28504

No BLAST hit | <95 | 95-97 | 97-98 | 98-98.5 | 98.5-99 | 99-99.2 | 99.2-99.4 | 99.4-99.6 | 99.6-99.8 | 99.8-100 | 100

B

tnpR Tn3 *bla*TEM-1 *bla*KPC-2 ΔIS*Kpn6* *tnpR* Tn5403

A

ST
- ST307 [48]
- NA [9]
- ST101 [9]
- ST14 [9]
- ST395 [7]
- ST48 [5]
- ST392 [4]
- ST1788 [3]
- ST307-1LV [3]
- ST1430 [2]
- ST290 [2]
- ST353 [2]
- ST392-1LV [2]
- ST750 [2]
- ST11 [1]
- ST16 [1]
- ST1845 [1]
- ST257-4LV [1]
- ST29 [1]
- ST383 [1]
- ST45 [1]

pOXA48a

B

Species
- *K. pneumoniae* [103]
- *K. oxytoca* [5]
- *K. aerogenes* [2]
- *K. michiganesis* [2]
- *K. quasipneumoniae* [1]
- *K. variicola* [1]

Replicon type
- IncL [95]
- IncM1 [18]
- IncM2 [1]

C

Hospital
- Hospital A
- Hospital F
- Hospital G
- Hospital I
- GP surgery

ARGID_32490 (*Kp* ST307)
IncL, 72.2kb
Hospital A, Wales, 2019

pOXA48a (*Kp* ST14)
IncL, 61.9kb
Turkey, 2001

ARGID_19108 (*Kp* ST395)
IncM1, 61.0kb
Hospital B, Wales, 2014

ARGID_31477 (*Kp* ST383)
IncL, 73.0kb
Hospital P, Wales, 2019

ARGID_27157 (*Kp* ST101)
IncM1, 72.1kb
Hospital C, Wales, 2019

ARGID_10727 (*Kp* ST750)
IncM1, 72.3kb
Hospital I, Wales

Nucleotide similarity
to pOXA-48a

No BLAST hit   <95 95-97 97-98 98-98.5 98.5-99 99-99.2 99.2-99.4 99.4-99.6 99.6-99.8 99.8-100 100