Ubiquitin-like cGAS chain formation by a super enzyme activates anti-phage

response

Yan Yan[1,#], Jun Xiao[2,#], Fengtao Huang[1,3,#,*], Bingbing Yu[1], Rui Cheng[1], Hui Wu[1], Xueling Lu[1],

Xionglue Wang[1], Greater Kayode Oyejobi[2], Carol V. Robinson[4,5], Hao Wu[6,7], Di Wu[4,5,*], Longfei

Wang[2,*], Bin Zhu[1,3,*]

[1]Key Laboratory of Molecular Biophysics, the Ministry of Education, College of Life Science and

Technology, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China

[2]School of Pharmaceutical Sciences, Wuhan University, Wuhan, China

[3]Shenzhen Huazhong University of Science and Technology Research Institute, Shenzhen 518063,

China

[4]Department of Chemistry, University of Oxford, Oxford, OX1 3QZ, UK

[5]Kavli Institute for Nanoscience Discovery, University of Oxford, Oxford, OX1 3QU, UK

[6]Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School,

Boston, MA 02115, USA

[7]Program in Cellular and Molecular Medicine, Boston Children's Hospital, Boston, MA 02115,

USA.

[#]These authors contribute equally to this work.

[*]To whom correspondence should be addressed. Email: huang_fengtao@126.com;

di.wu2@chem.ox.ac.uk; wanglf@whu.edu.cn; bin_zhu@hust.edu.cn

The cyclic oligonucleotide-based anti-phage signaling system (CBASS) is a family

of defense system in prokaryotes[1,2]. Composed of a cyclic GMP-AMP synthase

(cGAS) and CBASS-associated proteins, CBASS utilizes cyclic oligonucleotides to activate antiviral immunity[3-6]. One major group of CBASS-associated proteins are homologs of eukaryotic E2 ubiquitin-conjugating enzymes. However, the function of E2 in CBASS remains elusive. Here, we report that a bacterial E2 enzyme regulates cGAS by imitating the entire ubiquitination cascade. This includes the processing of the cGAS C-terminus, conjugation of cGAS to a cysteine residue, ligation of cGAS to a lysine residue, cleavage of the isopeptide bond, and poly-cGASylation. The poly-cGASylation fully activates cGAS to produce cGAMP, which acts as an antiviral signal and leads to cell death. Our findings reveal unique regulatory roles of E2 in CBASS and provide insights into the origin of the ubiquitin system.

Cyclic GMP-AMP synthase (cGAS) plays a key role in mammalian cGAS-STING innate immunity[7,8]. cGAS shares a conserved catalytic domain with *Vibrio cholerae* DncV[9] and they were classified as cGAS/DncV-like nucleotidyltransferases (CD-NTases)[3,10]. The prokaryotic CD-NTases and ancillary proteins, which are encoded by operons and involved in bacterial anti-phage defense, constitute the cyclic oligonucleotide-based anti-phage signaling system (CBASS)[2]. Certain CBASSs share evolutionary origins with the eukaryotic cGAS-STING innate immunity pathway[2,5,6], indicating similar underlying principles between bacterial anti-phage defense and mammalian innate immunity. Over 5,000 distinct CBASSs have been identified in more than 10% of bacteria with known genome sequences[1,3,11]. About half of CBASS operons

encode a CD-NTase that can generate signal molecules and an effector that induces

programed cell death; while the remaining CBASS operons encode ancillary genes in

addition to the CD-NTase and the effector[1,11]. Some of these ancillary components have

been reported to function as a threat sensor or a regulator of bacterial defenses[4-6];

however, most of their functions and mechanisms are unclear. Among over 5,000

predicted CBASSs, 2,199 CBASSs encode homologs of ubiquitin systems[11], which can

be divided into two main categories: one category (hereafter referred to as E1E2/JAB-

CBASS, Fig. 1a) contains an E1-E2 fusion protein, and a JAB deubiquitinating

peptidase, but no E3 ligase[12,13], while the other category (hereafter referred to as E2-

CBASS, Fig. 1a) consisting of 616 CBASSs that only contain an E2 protein[1,11], of

which the function is elusive.

Structure of the E2-cGAS complex

To elucidate the role of the CBASS-related E2 protein, we focused on a E2-CBASS

operon from *Serratia marcescens* (Fig. 1a). The E2-CBASS operon encodes a cGAS

(Extended Data Fig. 1), an E2, a putative exonuclease, and a nucleotide sensor domain

(SAVED) fused to two transmembrane (TM) helices (termed 2TM-SAVED) (Fig. 1a)[1].

Reversed-phase liquid chromatography-ultraviolet (RPLC-UV) and mass spectrometry

revealed that the product of cGAS is 3′,2′-cGAMP (Extended Data Fig. 1). Interestingly,

when cGAS and E2 were co-expressed and co-purified, an additional band with a

molecular weight of 64 kDa appears on the SDS-PAGE gel, suggesting that cGAS (46

kDa) and E2 (18 kDa) are fused (Fig. 1b). When separately-purified cGAS and E2 were

mixed at 1:1 molar ratio, the linkage still formed without any cofactors in the storage buffer (50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 0.1 mM EDTA, 0.1% Triton X-100, and 50% glycerol) (Extended Data Fig. 2a). This fusion of E2 and cGAS could not be separated by SDS, confirming a covalent linkage between the two.

To elucidate the structural basis of the covalent linkage between E2 and cGAS, we purified the cGAS-E2 complex and performed structural studies using cryo-EM (Extended Data Fig. 2d-f). Two cryo-EM datasets of cGAS-E2 complex were collected and processed (Extended Data Fig. 3a, b; Extended Data Table 3). The 3D reconstruction of cGAS-E2 complex was refined to 3.3 Å and 3.1 Å, respectively. Initial model of cGAS-E2 complex were generated using AlphaFold and ColabFold[14,15] and rigid-fitted in the electron density map (Extended Data Fig. 2b, c). Although the cryo-EM density map of cGAS-E2 was anisotropic as a result of preferred orientations, it was of sufficient quality to perform model refinement guided by the initial model (Extended Data Fig. 3c-e).

The refined cryo-EM structure of cGAS-E2 is consistent with the AlphaFold prediction (Extended Data Fig. 4), with E2 binding to the C-terminal region of cGAS (Fig. 1c, d). The C-terminus of cGAS is threaded into a pocket in E2 (Fig. 1e) and in close proximity with the side chains of one Cys and two Lys residues (C101, K151, and K159). The overall structure of *S. marcescens* E2 is similar to the structure of a human E2 (UBE2D2[16]) with a root mean square deviation (RMSD) of 2.5 (Fig. 1f). C101 of *S. marcescen* E2 is located in close proximity to the active site of human E2, indicating that it is the active site cysteine. Notably, *S. marcescen* E2 has a C-terminal loop

(residue 150-160) that covers the active site instead of the C-terminal α-helices (α3 and

α4) found in human E2. In contrast to the E1E2/JAB-CBASS system, where cGAS

interacts with dimerized E1-E2 fusion, cGAS of E2-CBASS forms extensive

interactions with E2 at three interfaces in addition to the C-terminus (Fig. 1g). Although

the core region of cGAS from E2-CBASS resembles human cGAS and cGAS from the

E1E2/JAB-CBASS system, the C-terminal region that mediates E2 interactions is

unique in cGAS from E2-CBASS (Fig. 1h, i).

Covalent linkage between E2 and cGAS

In eukaryotes, the ubiquitin forms a thioester bond between its C-terminal Gly and a

Cys in E1, E2, and E3, or forms an isopeptide bond between its C-terminal Gly and a

Lys in a target protein or another ubiquitin. Similarly, the C-terminal tail of *S.*

marcescens cGAS (residues 400-407) is inserted into the catalytic pocket of E2 (Fig.

1e) and the cryo-EM density of the C-terminal E407 after G406 is absent. Interestingly,

G406 of cGAS is in close proximity to C101, K151, and K159 of E2, suggesting that

the covalent bond between cGAS and E2 in this study is either a thioester bond between

G406 of cGAS and C101 of E2, or an isopeptide bond between G406 and K151 or K159

of E2.

Alignment of the C-terminal sequences of CD-NTase homologs associated with E2 in

various CBASSs showed that the glycine corresponding to cGAS G406 is highly

conserved (Extended Data Fig. 5a). Deletion of the C-terminal 8 residues (ΔP400–E407)

or 2 residues (ΔG406-E407) of cGAS abolished the covalent linkage between cGAS

and E2, while ΔE407 had no effect. These results confirmed that G406 participates in

the covalent linkage (Fig. 2a). Consistently, mutation of G406 to Val or Leu also abolished the covalent linkage, while mutation of G406 to Ala that has a shorter side chain still allowed the linkage (Fig. 2a). K401A and Y405A mutations also showed no effect (Fig. 2a).

C101 is the active site cysteine (Fig. 1f) and is conserved among prokaryotic E2 homologs (Extended Data Fig. 5b). C101A mutant failed to form the covalent complex with cGAS (Fig. 2a) in contrast to C69A that showed no effect. These results suggest that a thioester bond between cGAS G406 and E2 C101 exists (Fig. 2b).

To validate the thioester bond between G406 and E2 C101, we incubated cGAS-E2 fusion with 10 mM DTT, as the thioester bond is unstable in the presence of DTT[17]. Interestingly, the cGAS-E2 complexes are mostly resistant to DTT treatment and only a small portion was separated by 10 mM DTT (Fig. 2c), indicating that the majority of cGAS-E2 complexes are not linked by thioester bond but perhaps isopeptide bond.

To test if cGAS and E2 are linked by an isopeptide bond, we mutated K151 and K159 of E2 that are close to G406 of cGAS (Fig. 1e). Mutation of K159A but not K151A significantly reduced the cGAS-E2 complex in the presence of DTT (Fig. 2c), suggesting that the DTT-insensitive bond between cGAS and E2 is a G406-K159 isopeptide bond, which is similar to the isopeptide bond between a ubiquitin and its target protein (Fig. 2d). Interestingly, E2 K159A mutation not only reduced the DTT-insensitive isopeptide bond, but also increased the DTT-sensitive cGAS-E2 complex (Fig. 2c), indicating that the both types of covalent bonds exist and the E2 studied could catalyze the formation of both thioester and isopeptide bonds. Consistently, cryo-EM

structures of cGAS-E2 exhibit heterogeneity in the E2 region, revealing conformations corresponding to both types of covalent bonds (G406-C101 and G406-K159), as depicted in Fig. 1e and Extended Data Fig. 6. Thus, bacterial E2 mimics the function of both eukaryotic E2 and E3.

As evident from the ubiquitin cascade, it is likely that the unstable thioester bond between cGAS and E2 is transient and will convert to a more stable isopeptide bond. The rates of total bond (thioester+isopeptide, -DTT, 0.63 ± 0.17 min^{-1}) and isopeptide bond (+DTT, 0.50 ± 0.14 min^{-1}) formation *in vitro* were measured and shown in Fig. 2e.

Poly-cGASylation of cGAS by E2

Notably, the disruption of G406-K159 isopeptide bond by E2 K159A mutation not only increased the G406-C101 thioester bond, but also generated larger species insensitive to DTT (Fig. 2c, f). The molecular weight of these proteins indicates that they are polymers containing various numbers of cGAS. Either co-expressed with cGAS or mixed with cGAS *in vitro*, E2 K159A mutant generated the cGAS polymers (Fig. 2f). We excised the gel bands corresponding to the cGAS-cGAS dimer from both co-expressed and *in vitro* assembled cGAS-E2 (K159A) and analyzed them with mass spectrometry. The results showed that the majority of peptides detected were from cGAS, with minimal signals from E2 and *E. coli* protein contaminants (Fig. 2f), supporting the formation of cGAS-cGAS dimer. These cGAS polymers are insensitive to DTT, suggesting that they are linked by isopeptide bonds similar to poly-ubiquitin. Deletion of the C-terminus of E2 including K151 and K159 (ΔK151-R162) resulted in more significant poly-cGASylation, and less thioester bond (Fig. 2c).

Cysteine protease activity of E2

The formation of either the G406-C101 thioester or the G406-K159 isopeptide bond requires the removal of the last C-terminal residue E407 of cGAS in order to expose the carboxyl group of G406. In eukaryotes, the removal of C-terminal residues of ubiquitin is carried out by specific proteases[18]. Since the single E407 is too small to observe, we constructed a cGAS-MBP by attaching a maltose binding protein to the C-terminus of cGAS to monitor the process. When cGAS-MBP was mixed with E2 *in vitro*, the presence of a protein with the same molecular weight as cGAS-E2 (64 kDa) suggested that the covalent linkage between cGAS-MBP and E2 is formed and the C-terminus of cGAS-MBP was removed (Fig. 3a). The presence of a protein with the same molecular weight as MBP further confirmed that the C-terminus of cGAS was indeed removed during the covalent bonding (Fig. 3a).

The eukaryotic ubiquitin system[18] and the E1E2/JAB-CBASS system[12,13] both rely on deubiquitinating enzymes such as DUBs or JAB, which are mostly cysteine proteases. However, processing of cGAS requires only E2, suggesting that E2 itself may function as a cysteine protease. The C101 provides structural basis for an active site cysteine. Since low-pH-induced autocleavage is well-known for classical cysteine proteases like asparaginyl endopeptidase and cathepsin L[19,20], we examined the effect of low-pH on the cGAS-E2 complex. Interestingly, pH 5.0 induced the cleavage of the isopeptide bond between cGAS and E2, as shown by the reduction of cGAS-E2 complex in the presence of DTT (Fig. 3b). However, in the absence of DTT, we found that like the K159A mutation, pH 5.0 also increased the thioester bond; and pH 5.0 had

no effect on E2 K159A and ΔK151-R162 mutants (Fig. 3c). When the pH was adjusted

from 5.0 to 8.0, the thioester bond (-DTT) between cGAS and E2 were converted back

to isopeptide bond (+DTT, Extended Data Fig. 7a), confirming that the two types of

covalent bond are interconvertible. The rate of thioester bond formation at pH 5.0

$(0.50\pm0.07$ min$^{-1})$ is shown in Fig. 3d.

Another typical feature of cysteine proteases is a His residue near the catalytic Cys

for proton transfer during proteolytic catalysis[21,22]. Interestingly, our structure of cGAS-

E2 complex reveals two His (H87 and H152) residues that are very close to C101 (Fig.

3e). The E2 H87A mutation, like C101A (Fig. 2a), abolished the covalent linkage

between cGAS and E2 (Fig. 3f). Both mutants failed to cleave the MBP of cGAS-MBP

(Extended Data Fig. 7b), confirming the catalytic roles of C101 and H87 in E2 protease-

like activity. The E2 H152A mutation, although showed no effect on the complex

formation (Fig. 3f), significantly reduced the conversion from isopeptide bond to

thioester bond at pH 5.0 (Fig. 3g), suggesting that His152 is a pH-sensor at protease

active site as those reported previously[23,24]. Thus, the E2 in this study possesses both

the catalytic[25,26] and regulatory elements of proteases. The protease activity appears

unique for E2 from the E2-CBASS systems, as sequence alignment showed that the

histidine residues corresponding to H87 and H152 are only conserved in E2 from the

E2-CBASS systems, but not in E2 from the E1E2/JAB-CBASS systems (Extended

Data Fig. 7c). Also, no similar histidine residue was observed in eukaryotic E2

(Extended Data Fig. 7d). Altogether, the E2 is a "super" enzyme functionally

mimicking the entire eukaryotic ubiquitination system (Fig. 3h).

cGAS is activated by E2-mediated poly-cGASylation

Human cGAS is activated through oligomerization[27,28]. The poly-cGAS observed in this study is likely the active form of cGAS. We compared the cGAMP synthesis activities of frees cGAS, cGAS-E2, cGAS-E2 (K159A), and cGAS-E2 (ΔK151-R162). The concentration of cGAS in 4 samples were kept the same (800 nM), and the latter two samples contained more poly-cGAS (Fig. 4a). Interestingly, although the cGAS-E2 linkage inhibited the activity of cGAS, cGAS-E2 (K159A) and cGAS-E2 (ΔK151-R162) significantly increased the synthesis activity of cGAS when poly-cGAS were formed (Fig. 4a). When the concentration of ATP substrate is lowered by half, there was a more significant increase of synthesis activity by poly-cGAS (Fig. 4b), suggesting that the poly-cGASylation increased the affinity of cGAS to ATP.

Immunoblot using cGAS-specific antibody detected the poly-cGAS in the CBASS-expressing bacteria upon T4 phage infection (Fig. 4c), confirming that the phage-induced poly-cGASylation occurs *in vivo*. Recent work has demonstrated that CD-NTase+2TM-SAVED operons utilize antiviral nucleotide signals to activate CBASS TM effectors, which in turn induce inner-membrane disruption and cause cell death to limit phage replication[29]. With this effector, the antiviral signal producer cGAS, and a regulatory E2, the E2-CBASS operon confers strong immunity on *E. coli* against bacteriophage T4, as judged by plaque formation (Fig. 4d). This immunity was abolished upon: (1) deletion of either cGAS, or E2, or the effector 2TM-SAVED gene; (2) disruption of the second messenger synthesis by the cGAS D84A/D86A mutation; (3) disruption of the covalent linkage by the cGAS-G406R, E2-C101A, or E2-H87A

mutation; (4) disruption of the low pH-dependent bond conversion by the E2-H152A, or E2-ΔK151-R162 mutation (Fig. 4d). Although K159 is the preferred residue for stable isopeptide bond formation, its mutation did not show significant effect on the antiviral immunity (Fig. 4d), indicating that the partial releasing of poly-cGAS by K159A does not impede the antiviral immunity.

DISCUSSION

Origin of the eukaryotic ubiquitin system

The ubiquitin system is a highly conserved in eukaryotes and plays a critical role in a variety of cellular processes, including protein degradation, DNA repair, and signal transduction[18,30]. The ubiquitin machinery consists of at least 5 proteins: a protease, an E1, an E2, an E3, and a target protein. We demonstrate that the bacterial E2 in our study regulates cGAS by imitating the entire ubiquitination cascade, including the cleavage of the C-terminus of cGAS or the isopeptide bond (functions like a protease), conjugation of cGAS to a cysteine residue (functions like E1 and E2), and ligation of cGAS to a lysine residue (functions like E3 and a target protein) (Fig. 3h). The multifunctional E2 provides insights into the evolution of eukaryotic ubiquitin system. Our results suggest that the ubiquitin system may have arisen from a single E2-like protein that originally regulate antiviral synthase rather than ubiquitin. Over time, the function of E2 diverged, resulting in the emergency of E1, E2, and JAB. The E1E2/JAB-CBASS[12,13] may present this evolutionary stage. Finally, ubiquitin and E3 was incorporated for regulations of protein degradation in eukaryotes. One interesting aspect of the E2 protein described in this study is its protease activity, which appears to

be unique among all known E2s to date, suggesting that this particular E2 may have a

more primitive status in evolution. Our work also provides a compelling evidence for

the central role of E2 in ubiquitin systems[31,32].

Poly-cGASylation by E2 for cGAS activation

The poly-cGASylation for cGAS activation revealed in this study is unique. However,

it shares similarity with human cGAS and cGAS in the E1E2/JAB-CBASS system. The

human cGAS depends on oligomerization for activation[27,28]; and bacterial cGAS in the

E1E2/JAB-CBASS system is activated by conjugation to other proteins[12,13]. Activation

through inter-protein interaction is likely a common feature for cGAS. The poly-

cGASylation in this study was triggered by mutations of the E2 C-terminus. Although

the poly-cGASylation was also observed *in vivo* upon phage infection (Fig. 4c), the

exact signal from phages to E2 remains to be determined. Upon the phage signal

disrupting the C-terminus of E2, the sequential poly-cGASylation and cGAMP

production trigger membrane disruption and abortive infection (Fig. 4e).

REFERENCES

1   Burroughs, A. M., Zhang, D., Schaffer, D. E., Iyer, L. M. & Aravind, L. Comparative genomic
        analyses reveal a vast, novel network of nucleotide-centric systems in biological conflicts,
        immunity and signaling. *Nucleic acids research* **43**, 10633-10654, doi:10.1093/nar/gkv1267
        (2015).

2   Cohen, D. *et al.* Cyclic GMP-AMP signalling protects bacteria against viral infection. *Nature* **574**,
        691-695, doi:10.1038/s41586-019-1605-5 (2019).

3   Whiteley, A. T. *et al.* Bacterial cGAS-like enzymes synthesize diverse nucleotide signals. *Nature*
        567, 194-199, doi:10.1038/s41586-019-0953-5 (2019).

4   Lowey, B. *et al.* CBASS Immunity Uses CARF-Related Effectors to Sense 3'-5'- and 2'-5'-Linked
        Cyclic Oligonucleotide Signals and Protect Bacteria from Phage Infection. *Cell* **182**, 38-49 e17,
        doi:10.1016/j.cell.2020.05.019 (2020).

5   Ye, Q. *et al.* HORMA Domain Proteins and a Trip13-like ATPase Regulate Bacterial cGAS-like

Enzymes to Mediate Bacteriophage Immunity. *Molecular cell* **77**, 709-722 e707, doi:10.1016/j.molcel.2019.12.009 (2020).

Morehouse, B. R. *et al.* STING cyclic dinucleotide sensing originated in bacteria. *Nature* **586**, 429-433, doi:10.1038/s41586-020-2719-5 (2020).

Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z. J. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* **339**, 786-791, doi:10.1126/science.1232458 (2013).

Wu, J. *et al.* Cyclic GMP-AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* **339**, 826-830, doi:10.1126/science.1229963 (2013).

Davies, B. W., Bogard, R. W., Young, T. S. & Mekalanos, J. J. Coordinated regulation of accessory genetic elements produces cyclic di-nucleotides for V. cholerae virulence. *Cell* **149**, 358-370, doi:10.1016/j.cell.2012.01.053 (2012).

Kranzusch, P. J., Lee, A. S., Berger, J. M. & Doudna, J. A. Structure of human cGAS reveals a conserved family of second-messenger enzymes in innate immunity. *Cell reports* **3**, 1362-1368, doi:10.1016/j.celrep.2013.05.008 (2013).

Millman, A., Melamed, S., Amitai, G. & Sorek, R. Diversity and classification of cyclic-oligonucleotide-based anti-phage signalling systems. *Nature microbiology* **5**, 1608-1615, doi:10.1038/s41564-020-0777-y (2020).

Ledvina, H. E. *et al.* An E1-E2 fusion protein primes antiviral immune signalling in bacteria. *Nature*, doi:10.1038/s41586-022-05647-4 (2023).

Jenson, J. M., Li, T., Du, F., Ea, C.-K. & Chen, Z. J. Ubiquitin-like Conjugation by Bacterial cGAS Enhances Anti-phage Defence. *Nature*, doi:10.1038/s41586-023-05862-7 (2023).

Mirdita, M., Ovchinnikov, S. & Steinegger, M. ColabFold - Making protein folding accessible to all. *bioRxiv*, 2021.2008.2015.456425, doi:10.1101/2021.08.15.456425 (2021).

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

Dou, H., Buetow, L., Sibbet, G. J., Cameron, K. & Huang, D. T. BIRC7–E2 ubiquitin conjugate structure reveals the mechanism of ubiquitin transfer by a RING dimer. *Nature structural & molecular biology* **19**, 876-883 (2012).

Termathe, M. & Leidel, S. A. The Uba4 domain interplay is mediated via a thioester that is critical for tRNA thiolation through Urm1 thiocarboxylation. *Nucleic acids research* **46**, 5171-5181, doi:10.1093/nar/gky312 (2018).

Kerscher, O., Felberbaum, R. & Hochstrasser, M. Modification of proteins by ubiquitin and ubiquitin-like proteins. *Annual review of cell and developmental biology* **22**, 159-180, doi:10.1146/annurev.cellbio.22.010605.093503 (2006).

Zhao, L. *et al.* Structural analysis of asparaginyl endopeptidase reveals the activation mechanism and a reversible intermediate maturation stage. *Cell research* **24**, 344-358, doi:10.1038/cr.2014.4 (2014).

Lowther, J. *et al.* The importance of pH in regulating the function of the Fasciola hepatica cathepsin L1 cysteine protease. *PLoS neglected tropical diseases* **3**, e369, doi:10.1371/journal.pntd.0000369 (2009).

Verma, S., Dixit, R. & Pandey, K. C. Cysteine Proteases: Modes of Activation and Future Prospects as Pharmacological Targets. *Frontiers in pharmacology* **7**, 107, doi:10.3389/fphar.2016.00107 (2016).

Elsässer, B. *et al.* Distinct Roles of Catalytic Cysteine and Histidine in the Protease and Ligase Mechanisms of Human Legumain As Revealed by DFT-Based QM/MM Simulations. *ACS Catalysis* **7**, 5585-5593, doi:10.1021/acscatal.7b01505 (2017).

Feliciangeli, S. F. *et al.* Identification of a pH sensor in the furin propeptide that regulates enzyme activation. *The Journal of biological chemistry* **281**, 16108-16116, doi:10.1074/jbc.M600760200 (2006).

Williamson, D. M., Elferich, J., Ramakrishnan, P., Thomas, G. & Shinde, U. The Mechanism by Which a Propeptide-encoded pH Sensor Regulates Spatiotemporal Activation of Furin. *Journal of Biological Chemistry* **288**, 19154-19165, doi:10.1074/jbc.M112.442681 (2013).

Vernet, T. *et al.* Processing of the papain precursor. The ionization state of a conserved amino acid motif within the Pro region participates in the regulation of intramolecular processing. *The Journal of biological chemistry* **270**, 10838-10846, doi:10.1074/jbc.270.18.10838 (1995).

FUENTES-PRIOR, P. & SALVESEN, Guy S. The protein structures that shape caspase activity, specificity, activation and inhibition. *Biochemical Journal* **384**, 201-232, doi:10.1042/bj20041142 (2004).

Li, X. *et al.* Cyclic GMP-AMP synthase is activated by double-stranded DNA-induced oligomerization. *Immunity* **39**, 1019-1031, doi:10.1016/j.immuni.2013.10.019 (2013).

Yu, L. & Liu, P. Cytosolic DNA sensing by cGAS: regulation, function, and human diseases. *Signal transduction and targeted therapy* **6**, 170, doi:10.1038/s41392-021-00554-y (2021).

Duncan-Lowey, B., McNamara-Bordewick, N. K., Tal, N., Sorek, R. & Kranzusch, P. J. Effector-mediated membrane disruption controls cell death in CBASS antiphage defense. *Molecular cell* **81**, 5039-5051.e5035, doi:10.1016/j.molcel.2021.10.020 (2021).

Li, W. & Ye, Y. Polyubiquitin chains: functions, structures, and mechanisms. *Cellular and molecular life sciences : CMLS* **65**, 2397-2406, doi:10.1007/s00018-008-8090-6 (2008).

David, Y., Ziv, T., Admon, A. & Navon, A. The E2 ubiquitin-conjugating enzymes direct polyubiquitination to preferred lysines. *The Journal of biological chemistry* **285**, 8595-8604, doi:10.1074/jbc.M109.089003 (2010).

Stewart, M. D., Ritterhoff, T., Klevit, R. E. & Brzovic, P. S. E2 enzymes: more than just middle men. *Cell research* **26**, 423-440, doi:10.1038/cr.2016.35 (2016).

Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nature Methods* **14**, 290-296 (2017).

Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* **14**, 331-332, doi:10.1038/nmeth.4193 (2017).

Bepler, T. *et al.* Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. *Nature Methods* **16**, 1153-1160, doi:10.1038/s41592-019-0575-8 (2019).

Zivanov, J., Nakane, T. & Scheres, S. A Bayesian approach to beam-induced motion correction in cryo-EM single-particle analysis. *Cold Spring Harbor Laboratory* (2018).

Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction. *Nature Methods* **17**, 1214-1221 (2020).

Punjani, A. & Fleet, D. J. 3D Variability Analysis: Resolving continuous flexibility and discrete heterogeneity from single particle cryo-EM. *Journal of Structural Biology* **213**, 107702 (2021).

Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume post-processing. *Communications biology* **4**, 874 (2021).

Pettersen, E. F., Goddard, T. D., Huang, C. C., Meng, E. C. & Ferrin, T. E. UCSF ChimeraX: Structure

Visualization for Researchers, Educators, and Developers. *Protein Science* (2020).

41   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*

crystallographica.   Section   D,   Biological   crystallography   **66**,   486-501,

doi:10.1107/S0907444910007493 (2010).

42   Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and

electrons: recent developments in Phenix. *International Union of Crystallography* (2019).

43   Wang Ray, Y.-R. *et al.* Automated structure refinement of macromolecular assemblies from

cryo-EM maps using Rosetta. *eLife* **5**, doi:https://doi.org/10.7554/eLife.17219 (2016).

44   Leman, J. K. *et al.* Macromolecular modeling and design in Rosetta: recent methods and

frameworks. *Nat Methods* **17**, 665-680, doi:10.1038/s41592-020-0848-2 (2020).

45   Mazzocco, A., Waddell, T. E., Lingohr, E. & Johnson, R. P. Enumeration of Bacteriophages Using

the Small Drop Plaque Assay System. *Methods Mol Biol* **501**, 81-85, doi:10.1007/978-1-60327-

164-6_9 (2009).

Figure legends

Fig. 1 Cryo-EM structure of a cGAS-E2 complex. a, Domain organization of the

CBASS operons from *Serratia marcescens*, *E. coli*[13], *V. cholerae* and *Enterobacter cloacae*[12]. **b**, Left, Schematic diagram showing the constructs for co-expression of cGAS and E2 in different forms (I–III). Right, SDS-PAGE analysis of the purified proteins from various constructs as shown on the left. Data are representative of 3 independent experiments. **c**, Overlay of cryo-EM density and the atomic model of the cGAS-E2 complex. **d-e**, Ribbon diagram of the cGAS-E2 complex highlighting the C-terminal tail of cGAS (green) that is inserted into the E2 domain (orange). The potential interaction region (boxed) is enlarged in **e**. **f**, Superimposed structures of E2 used in this study (orange) and human E2 (gray, PDB ID 4AUQ). CT refers to the C-terminus. **g**, Ribbon diagram of the cGAS-E2 complex highlighting the interface between cGAS and E2. **h**, Superimposed structures of cGAS used in this study (green) and human cGAS (gray, PDB ID 6EDC). **i**, Superimposed structures of cGAS used in this study from E2-CBASS (green) and *Ec*CdnD[12] from E1E2/JAB-CBASS (gray, PDB ID 7TO3). CTD refers to the C-terminal domain.

Fig. 2 Covalent linkage between E2 and cGAS. a, SDS-PAGE analysis of cGAS

mutants (indicated on top of gel) co-purified with E2, or E2 mutants (indicated on top

of gel) co-purified with cGAS. The G406 residue of cGAS and the C101 residue of E2

were identified to be critical residues for the covalent linkage. Data are representative

of 3 independent experiments. b, Schematic diagram showing the thioester bond

between E2 and cGAS, and the thioester bond between E2 and ubiquitin. c, SDS-PAGE

analysis of the co-purified proteins consisting of cGAS with E2 or E2 mutants in the

presence (+DTT) or absence (-DTT) of 10 mM DTT (10 mM DTT was used in all

following analysis). The K159 residue of E2 was identified to be the key residue for the

isopeptide bonding. Data are representative of 3 independent experiments. d, Schematic

diagram showing the isopeptide bond between E2 and cGAS, and the isopeptide bond

between ubiquitin and substrate. **e**, Left, SDS-PAGE analysis showing the *in vitro*

formation of the cGAS-E2 complex in the presence (+DTT) or absence (-DTT) of 10

mM DTT. Right, Rate of the total bond (thioester+isopeptide, -DTT) and isopeptide

bond (+DTT) formation between cGAS and E2, as calculated based on gels in the left

panel. Data are mean ± s.d. for $n = 3$ independent replicates and are representative of 3

independent experiments. **f**, Left, SDS-PAGE analysis showing the *in vivo* and *in vitro*

formation of the covalent cGAS-E2 (K159A) protein. The *in vivo* formation of the

covalently-linked cGAS-E2 (K159A) protein was achieved by the co-purification of

His-tagged cGAS and non-tagged E2 (K159A) mutant. For the *in vitro* assembly, two

proteins were mixed at a molar ratio of 1:1 overnight. Right, Mass spectrometry

showing the score of proteins contained in the potential cGAS-cGAS extracted from

the gel analyzing the cGAS-E2 (K159A) as shown on the left (marked by green arrow).

Mass spectrometry signals corresponding to *E. coli* protein contaminants in cGAS-

cGAS dimer are detailed in Extended Data Table 4.

Fig. 3 E2 is a cysteine protease. a, Left, SDS-PAGE analysis showing the *in vitro*

processing and bonding of cGAS-MBP by E2. cGAS-MBP or cGAS and E2 were

mixed at a molar ratio of 1:1 overnight. Right, Schematic diagram showed that the C-

terminal cleavage of cGAS by E2 is similar to the C-terminal cleavage of ubiquitin by

eukaryotic DUBs. **b**, SDS-PAGE analysis showing the low-pH-induced autocleavage

of the covalent cGAS-E2 complex. The optimal pH for the autocleavage is 5.0. 10 mM

DTT is presented in all reactions. Data are representative of 3 independent experiments.

c, SDS-PAGE analysis of the low-pH-induced bond conversion in cGAS-E2 or cGAS-

E2 (K159A) or cGAS-E2 (ΔK151-R162) in the presence (+DTT) or absence of DTT (-

DTT). The covalent cGAS-E2 complex obtained by the co-purification of His-tagged

cGAS and non-tagged E2 or E2 mutants were incubated in the reaction buffer at pH 5.0

for 30 min. **d**, Rate of the thioester bond formation in cGAS-E2 at pH 5.0 (right)

calculated based on gel analysis (left). At each time interval, the amount of total bonds

(thioester+isopeptide, as quantification of cGAS-E2 in the absence of DTT) subtracting

the amount of isopeptide bond (as quantification of cGAS-E2 in the presence of DTT)

was calculated as the amount of thioester bond. Data are mean ± s.d. for $n = 3$

independent replicates and are representative of 3 independent experiments. **e**,

Magnified view of E2 showing Cys101, His87, and His152 and other key residues in

the active site. E2-CT refers to the C-terminus of E2. **f**, SDS-PAGE analysis of cGAS

co-purified with E2 or E2 mutants (as indicated on top of gel) in the presence (+DTT)

or absence of DTT (-DTT). The H87 residue of E2 was identified to be crucial for the

covalent linkage. Data are representative of 3 independent experiments. **g**, SDS-PAGE

analysis of the low-pH-induced autocleavage of the covalent cGAS-E2 complex in the

presence (+DTT) or absence of DTT (-DTT). The covalent cGAS-E2 complex obtained

by the co-purification of His-tagged cGAS and non-tagged E2 or E2 mutants were

incubated in the reaction buffer at pH 5.0 for 30 min. The H152 residue of E2 was

crucial to mediate the low-pH-induced cleavage of the isopeptide bond between cGAS

and E2. Data are representative of 3 independent experiments. **h**, Schematic diagram

showing the eukaryotic ubiquitin pathway (left) and E2-mediated cGAS regulation

(right).

Fig. 4 Poly-cGASylation stimulates cGAMP production and anti-phage immunity.

a, Left, SDS-PAGE analysis of cGAS and the co-purified proteins consisting of cGAS

with E2 or E2 mutants (indicated on top of gel). Right, Quantification of 3′,2′-cGAMP

(based on HPLC analysis shown in Extended Data Fig. 8a) synthesized by cGAS and

variants shown in left panel. Data are mean ± s.d. for $n = 3$ independent replicates and

are representative of 3 independent experiments. **b**, Quantification of 3′,2′-cGAMP

(based on HPLC analysis shown in Extended Data Fig. 8b) synthesized by cGAS and

variants shown in (**a**), with the concentration of ATP or GTP or Mn^{2+} in the reaction

reduced by half compared to those in (**a**), respectively. Data are mean ± s.d. for $n = 3$

independent replicates and are representative of 3 independent experiments. **c**, Western

blot analysis with cGAS-specific antibody showing the poly-cGAS in *E. coli* BL21

expressing E2-CBASS operon during T4 phage infection. Phage T4 was applied at a

multiplicity of infection (MOI) of 5. Data are mean ± s.d. for $n = 3$ biological replicates.

d, Phage T4 infection of cells expressing empty vector, the full native E2-CBASS

operon (WT system), WT system with deletion or mutations of cGAS, WT system with

deletion or mutations of E2, or WT system with deletion of the effector 2TM-SAVED.

Data are representative of 3 independent experiments. **e**, Model for the E2-CBASS

antiviral mechanism.

Methods

Plasmid construction

All experiments were performed with the CBASS system from *Serratia marcescens*

(NCBI RefSeq NZ_KE560335.1). For the expression and purification of cGAS

(WP_016928966.1) and E2 (WP_016928967.1), their coding sequences were

codon-optimized and commercially synthesized by Genscript. For the co-expression of

cGAS and E2, an N-terminal 6×His-tag sequence was constructed in pET28a vector.

Non-tagged expression vectors were constructed from pQE82L vector, with the T5

promoter replaced by a T7 promoter for E2. Three forms of co-expression present in

Fig. 1b were assessed for cGAS and E2: (1) the gene encoding cGAS with an

N-terminal 6×His-tag cloned into pET28a vector, the gene encoding E2 gene without

tag cloned into pQE82L vector, the T5 promoter in pQE82L replaced with a T7

promoter; (2) the gene encoding cGAS without tag cloned into pQE82L vector and the

gene encoding E2 with N-terminal 6×His-tag cloned into pET28a vector; (3) the native

sequence of cGAS and E2 in a polycistronic form derived from *S. marcescens* was

cloned into pET28a vector. For phage infections, the four-gene CBASS system operon

from *S. marcescens* spanning a nucleotide range of 26,427–29,783 were cloned into

plasmid pQE82L. Details on the cloning of cGAS and E2 mutants are listed in Extended

Data Table 1 and 2, respectively. Plasmids generated in this study were constructed by

Gibson from synthesized genes with ≥18 base pairs of homology flanking the insert

sequence using ClonExpress II One Step Cloning Kit (Vazyme Biotech). Plasmids were

transformed into *E. coli* DH5α cells and confirmed by Sanger sequencing.

Protein expression, co-expression and purification

Plasmids for protein expression were transformed into *E. coli* BL21 (DE3). To

determine the interaction between cGAS and E2, they were co-transformed into *E. coli*

BL21 (DE3) and selected with their respective antibiotic marker. Bacterial cells were

cultivated as a 4 mL starter culture in LB broth overnight at 37 ºC with shaking at 220

rpm. The cultures were then transferred into 200 mL LB broth and cultivated for

approximately 3 h until the optical density at 600 nm (OD_{600}) reached 0.8–1.0. After

cooling down to room temperature, 0.25 mM IPTG was added to the cultures, and

cultivation was continued overnight at 16 ºC.

Cultures were collected and lysed by sonication in Lysis buffer (20 mM Tris-HCl,

pH 8.0, 300 mM NaCl, 20 mM imidazole), and the lysates were clarified by

centrifugation at $21,000 \times g$, 4 ºC for 1 h and filtered through a 0.45-μm filter. Proteins

were purified by affinity chromatography using Ni-NTA resin (Qiagen) and a gravity

column (Biorad). Ni-NTA resin was pre-equilibrated with Lysis buffer, bound to target

proteins, washed with Wash buffer (20 mM Tris-HCl, pH 8.0, 300 mM NaCl, and 50

mM imidazole for cGAS and its mutants; 20 mM Tris-HCl, pH 8.0, 300 mM NaCl, and

50, 80, or 100 mM imidazole for E2, its mutants and the co-expressed proteins) and

eluted with Elution buffer (20 mM Tris-HCl, pH 8.0, 300 mM NaCl, and 100 mM

imidazole for cGAS, its mutants and the co-expressed proteins; 20 mM Tris-HCl, pH

8.0, 300 mM NaCl, and 200 mM imidazole for E2 and its mutants). Proteins were

filter-concentrated using centrifugation and a 10-kDa or 30-kDa cut-off column

(Millipore Sigma) and dialyzed at 4 ºC for ~24 h against Dialysis Buffer (50 mM

Tris-HCl pH 7.5, 100 mM NaCl, 0.1 mM EDTA, 0.1% Triton X-100 and 50% glycerol).

Proteins were analyzed by SDS-PAGE with Coomassie blue (Bio-Rad) staining.

Structure prediction

ColabFold was used to predict the structure of the cGAS-E2 complex. Multiple

sequence alignment (MSA) was performed using MMseqs2 (UniRef + Environmental).

Structural prediction was performed using AlphaFold on a virtual cluster node with a

Nvidia Tesla P100 GPU.

Cryo-EM sample preparation and data collection

cGAS-E2 complex was further purified by size-exclusion chromatography on a HiLoad

16/600 Superdex 200 pg column (Cytiva) in TBS buffer (20 mM Tris-HCl, pH 8.0, 100

mM NaCl). Proteins were concentrated to 3-7 mg/mL, flash-frozen in liquid nitrogen,

and stored at -80°C.

A 3 µL-drop of cGAS-E2 sample at 0.7 mg/mL was applied to glow-discharged

copper Quantifoil R1.2/R1.3 grids, and blotted for 3 s in 100% humidity at 8 °C, then

plunged into liquid ethane using a Vitrobot Mark IV. All grids were screened using a

Glacios microscope. The data were collected on a 300 keV Titan Krios G4 microscope

equipped with a Gatan K3 direct electron detector and a Gatan Quantum energy filter

at the Cryo-EM Unit of Core Facility of Wuhan University.

For the initial dataset, 5,883 movies were collected in super-resolution mode, with

40 frames per movie, 1.61 s exposure time, 50 e-/Å2 accumulated dose, and 0.335 Å

pixel size. For the final dataset, cGAS-E2 sample concentration was increased to 3.2

mg/mL and UltrAuFoil grids was used instead for plunging. 4,752 movies were

collected in super-resolution mode, with 40 frames per movie, 2.16 s exposure time, 70

e-/Å2 accumulated dose, and 0.335 Å pixel size.

Cryo-EM data processing and model building

For dataset I, movies were binned 2x (pixel size 0.67 Å) and motion-corrected using

patch motion correction in CryoSPARC[33,34], followed by patch contrast transfer

function (CTF) estimation performed in the same software. Particles were picked using

blob picker, binned 2x (pixel size 1.34 Å) and subjected to two rounds of 2D

classifications in CryoSPARC with number of online-EM iterations set to 40 and batch

size per class set to 400. 869,610 particles representing the binary complex were

selected, and were used for ab-initio 3D reconstruction and heterogeneous refinements

(Extended Data Fig. 3a). One class out of four were selected for 3D refinement using

homogeneous refinement in C1 symmetry, resulting in a 3.8 Å cryo-EM density map

of cGAS-E2. After global CTF refinement, a soft mask excluding the flexible terminal

loops of cGAS-E2 complex was generated and used for local refinement in CryoSPARC.

Particles were unbinned and used for another round of local refinement, and the final

map resolution of cGAS-E2 complex was 3.35 Å.

The data processing procedures of dataset II were similar to dataset I, with a few

differences (Extended Data Fig. 3b). (1) A general Topaz[35-38] model was trained with

the particles from the initial dataset and used for particle picking. Blob picker was also

used for picking and after the first round of 2D classification, particles picked from the

two pickers were merged and duplicate particles were removed. (2) Heterogeneous

refinement (3D classification) was performed on 512,003 particles from 2D

classification using the low-passed 3.35 Å map as initial models. (3) The final cryo-EM

map of cGAS-E2 complex reached a resolution of 3.0 Å. Some degree of preferred

orientation was present in both of the datasets, resulting in anisotropic maps (Extended

Data Fig. 3c, d). DeepEMhancer[39] was used for post-processing to improve the

anisotropic map, which helped with the interpretation of the map. The initial model of

cGAS-E2 was generated by rigid body fitting of the predicted cGAS-E2 structure into

the cryo-EM maps using Chimera[40]. Regions with no electron densities were removed

from the model, including residues 1-4, 191-210, 389-396 from cGAS and 1-3 from E2.

Linkage between cGAS G406 and E2 K159 or C101 were modeled based on the

densities with local adjustments. Initial B-factors for all the atoms of the model were

set to 94.7 based on the Guinier Plot from final local refinement job. Inspection, model

building, and manual adjustments were performed in Coot[41] and real-space refinements

were carried out in Phenix[42,43]. Representations of cryo-EM densities and structural

models were generated using ChimeraX[40] and PyMOL[44].

***In vitro* bonding assay**

To reconstitute the covalent attachment of cGAS to E2 *in vitro*, cGAS and E2 or their

mutants were mixed in 50 mM Tris-HCl (pH 7.0), unless otherwise noted. The mixture

was incubated at 4 ºC for the indicated periods. The reaction was quenched by adding

SDS sample buffer with DTT (Solarbio Life Sciences) or without DTT (Biosharp Life

Sciences) and heating at 95°C for 5 minutes. Proteins in the solutions were separated

by SDS-PAGE in the absence of reducing agent and stained with Coomassie blue.

GraphPad Prism was used for data analysis.

***In vitro* cleavage assay**

To examine the effect of low-pH on the cGAS-E2 complex, cGAS-E2 complex or its

mutant was incubated in citrate buffer with different pH (Shyuanye) at 4 ºC for the

indicated periods. The reaction was quenched by adding SDS sample buffer with DTT

(Solarbio Life Sciences) or without DTT (Biosharp Life Sciences) and heating at 95°C

for 5 minutes. Proteins in the solutions were separated by SDS-PAGE in the absence of

reducing agent and stained with Coomassie blue. GraphPad Prism was used for data

analysis.

HPLC analysis of enzymatic reactions

CD-NTase reactions were performed essentially as previously described[3]. Briefly,

20-μL reactions contained 2 μM enzyme and 250 μM NTPs (New England BioLabs) in

reaction buffer with 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 2.5 mM $MnCl_2$/$MgCl_2$,

and 1 mM DTT, unless otherwise noted. Reactions were incubated at 37 ºC for the

indicated periods, inactivated at 80 ºC for 10 min, and centrifuged for 15 min at 21,000

× *g* to remove precipitated protein. Reaction products were analyzed by HPLC with a

ZORBAX Eclipse XDB-C18 (4.6 × 250 mm) column and an Agilent 1260 Infinity II

Series LC system. Next, 10 µL of the reaction product was injected into the column and

eluted with solvent A (methanol, sigma) and solvent B (20 mM ammonium acetate,

sigma) at a flow rate of 1 mL/min using the following linear gradient: 0–5 min, 5% A;

5–15 min, 5–100% A; 15–20 min, 100% A. The column was re-equilibrated for 5 min

at 5% A. For analyzing the reaction products treated with CIP or nuclease P1, 20 µL of

the reaction product was treated with 1 µL of CIP or nuclease P1 (New England BioLabs)

at 37 ºC for 1 h and inactivated at 80 ºC for 10 min prior to centrifugation and analysis

by HPLC.

Western blots

Rabbit cGAS polyclonal antibody was generated by a commercial vendor (AtaGenix)

using a purified cGAS antigen. Serum was used at 1:1,000 for cGAS immunoblot

detection. GAPDH antibody (ab125247, Abcam) was used at 1:1,000 for use as a

loading control.

 For whole-cell lysate analysis, *E. coli* BL21 carrying the indicated plasmid were

grown to mid-logarithmic phase and protein expression was induced by 0.2 mM IPTG.

Cells were infected with phage T4 for the indicated periods at an MOI of 5. Cells were

lysed by FastBreak™ cell lysis reagent (Promega) at room temperature for 15 min,

followed by centrifugation at 4 ºC to remove cellular debris. 80 µL soluble lysates were

then mixed with 20 µL 5× SDS sample buffer with DTT, heated and concentrated at

100 ºC for 30 min followed by a 5-min centrifugation at 21,000× g. Proteins in the

solutions were separated by SDS-PAGE, then transferred to PVDF membranes

(Millipore Sigma) charged in methanol. Membranes were blocked in nonfat dry milk

for 1 h at 24 ºC, followed by incubation with primary antibodies diluted in antibody

dilution buffer at 1:1,000 overnight at 4 ºC. PVDF membranes were then incubated with

the HRP-conjugated goat anti-rabbit IgG (SA00001-2, Proteintech) or HRP-conjugated

goat anti-mouse IgG (SA00001-1, Proteintech) at 1:5,000 dilution in TBST for 1 h at

24 ºC and detected with an ECL Chemiluminescent Substrate Kit (YEASEN) and

Amersham Imager 680 (GE Health).

Plaque assays

Phages were propagated by infecting exponentially-growing *E. coli* B culture in LB

medium at an MOI of 1:100–1:1,000, incubated at 37 ºC with shaking (200 rpm) until

clearing of the culture, the lysate was centrifuged and the supernatant was filter-

sterilized through a 0.45-µm filter.

Bacteriophage infection assay was determined using the small drop plaque assay

method[45]. The E2-CBASS operon and its mutants, as indicated, were cloned into the

pQE82L vector and transformed into *E. coli* BL21, respectively. Cells containing an

empty vector (pQE82L) were used as control. A single bacterial colony was picked,

respectively, and grown in LB broth at 37 ºC to an OD_{600} of ~0.2. E2-CBASS operon

expression was induced through 50 µM IPTG, followed by further growth for 1 h to an

OD_{600} of 0.6–0.7. 800 µL of cells were mixed with 25 mL of LB with 0.75% agar

containing appropriate antibiotic selection and IPTG (50 µM) and the entire sample was

poured onto plates. The phage stock was 10-fold serially diluted and 4-μL drops of

diluted phage lysate were placed on the solidified agar. The plates were then incubated

at 37 ºC for 16–18 h before imaging.

Data analysis

Data analysis was performed using Microsoft Excel 2019, OriginPro version 9.1 and

GraphPad Prism version 8.0.2. Graph plotting and statistical analysis was performed

using GraphPad Prism version 8.0.2.

Acknowledgements We thank Profs. En-Duo Wang, Wei Yan, Hongyu Hu for their

advice; Prof. Ping Xu for providing materials; Xuemei Li for technical supports; and

all lab members for helpful discussions. We thank Danyang Li and Yi Zeng of the Core

Facility of Wuhan University for their assistance with cryo-EM grid screening and data

collection. This work was supported by National Natural Science Foundation of China

(grant 32150009 and 31870165 to B.Z., 31900032 to F.T.H.) and Fund from Science,

Technology and Innovation Commission of Shenzhen Municipality (grant

JCYJ20210324115811032 to B.Z.). Funding for open access charge: National Natural

Science Foundation of China.

Author contributions F.T.H. and B.Z. conceived the project. Y.Y., J.X., F.T.H., D.W.,

L.F.W. and B.Z. designed the experiments. Y.Y., J.X. and D.W. carried out the

experiments. J.X. and L.F.W. determined and analyzed structures. Y.Y., F.T.H., G.K.O.,

C.V.R., H.W., D.W., L.F.W. and B.Z. analyzed the data and wrote the manuscript. All

authors discussed the results and contributed to the final manuscript.

Competing interests The authors declared that they have no conflicts of interest to this

work.

Additional Information

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to Bin Zhu**.**

Peer review information *Nature* thanks the anonymous reviewers for their

contribution to the peer review of this work.

Reprints and permissions information is available at http://www.nature.com/reprints.

Extended data

Extended Data Table 1 Primers for construction of cGAS mutations.

Name	Sequence
cGAS-D84A/D864-F	GCAATACGCGATCGCGCTGATTTGTTATCTGC
cGAS-D84A/D864-R	CAAATCAGCGCGATCGCGTATTGCTCTTCGTTT
cGAS-ΔP400-E407-F	AGGCTGGCGTGTAAGAATTCGAGCTCCGTC
cGAS-ΔP400-E407-R	CTCGAATTCTTACACGCCAGCCTGGGCCGCCGT
cGAS-ΔG406-E407-F	CACCTTCTACTAAGAATTCGAGCTCCGTCGAC
cGAS-ΔG406-E407-R	CGAATTCTTAGTAGAAGGTGTTCTTCGGCACGC
cGAS-ΔE407-F	CTTCTACGGCTAAGAATTCGAGCTCCGTCGAC
cGAS-ΔE407-R	CGAATTCTTAGCCGTAGAAGGTGTTCTTCG
cGAS-K401A-F	TGGCGTGCCGGCGAACACCTTCTACGGCGA
cGAS-K401A-R	AGAAGGTGTTCGCCGGCACGCCAGCCTGGGCC
cGAS-Y405A-F	ACACCTTCGCGGGCGAGTAAGAATTCG
cGAS-Y405A-R	TACTCGCCCGCGAAGGTGTTCTTCGGC

cGAS-G406A-F	CACCTTCTACGCGGAGTAAGAATTCGAGCTCC
cGAS-G406A-R	ATTCTTACTCCGCGTAGAAGGTGTTCTTCGGC
cGAS-G406V-F	CACCTTCTACGTGGAGTAAGAATTCGAGCTCC
cGAS-G406V-R	ATTCTTACTCCACGTAGAAGGTGTTCTTCGG
cGAS-G406L-F	CACCTTCTACCTGGAGTAAGAATTCGAGCTCC
cGAS-G406L-R	ATTCTTACTCCAGGTAGAAGGTGTTCTTCGGC

2 **Extended Data Table 2 Primers for construction of E2 mutations.**

Name	Sequence
E2-C69A-F	ACCCCGCCCATCGCGGTTGTTACCGAACCG
E2-C69A-R	TCGGTAACAACCGCGATGGGCGGGGTGCCG
E2-C101A-F	GGTGCGCGTCTGGCGCTGTTCCTGCCGAGAT
E2-C101A-R	CGGCAGGAACAGCGCCAGACGCGCACCCGGAAT
E2-H87A-F	GCCATCCCGGCGCTGTATCCCGCCGATAA
E2-H87A-R	GGATACAGCGCCGGGATGGCCCGCGTG
E2-H152A-F	GGCGGTAAGGCGCCACGCCCCAGCGAGGTTAA
E2-H152A-R	TGGGGCGTGGCGCCTTACCGCCGCCTTCCCAGT
E2-K151A-F	GCGGCGGCGCGCACCCGCGCCCATCTGAGGT
E2-K151A-R	GTGCGCGCCGCCGCCTTCCCAATGACCGGTGT
E2-K159A-F	CTGAGGTCGCGAACGAACGTTAAGAATTCG
E2-K159A-R	AACGTTCGTTCGCGACCTCAGATGGGCGCG
E2-ΔK151-R162-F	GCGGCTAAGGATCCGCATGCGAGCTCGGTA
E2-ΔK151-R162-R	ATGCGGATCCTTAGCCGCCGCCTTCCCAATGAC

4 **Extended Data Table 3 Cryo-EM data collection, refinement and validation statistics.**
5

	cGAS-E2 (EMDB-34992) (PDB 8HSB)
Data collection and processing	
Magnification	130,000x
Voltage (kV)	300
Electron exposure (e–/Å^2)	70
Defocus range (μm)	- 0.8 ~ - 2.2
Pixel size (Å)	0.67
Symmetry imposed	C1
Initial particle images (no.)	1,381,6133
Final particle images (no.)	242,891
Map resolution (Å)	3.1
FSC threshold	0.143
Map resolution range (Å)	40-2.68

Refinement

Initial model used (PDB code)	AlphaFold 2
Model resolution (Å)	3.9
FSC threshold	0.5
Model resolution range (Å)	60-3.9
Map sharpening B factor (Å^2)	94.7
Model composition	
Non-hydrogen atoms	4,278
Protein residues	533
Ligands	0
B factors (Å^2)	
Protein	131.28
Ligand	
R.m.s. deviations	
Bond lengths (Å)	0.003
Bond angles (°)	0.715
Validation	
MolProbity score	1.83
Clashscore	10.62
Poor rotamers (%)	0
Ramachandran plot	
Favored (%)	95.81
Allowed (%)	3.81
Disallowed (%)	0.38

Extended Data Table 4 *E. coli* proteins identified in cGAS-cGAS sample by mass spectrometry.

Accessions	Gene names	Descriptions
A0A140NGZ5	ECBD_4014	Uncharacterized protein OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=ECBD_4014 PE=4 SV=1
A0A140NAN3	ECBD_2463	Isocitrate dehydrogenase [NADP] OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=ECBD_2463 PE=4 SV=1
A0A140NC10	ECBD_2941	Citrate synthase OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=ECBD_2941 PE=3 SV=1
A0A140NF15	ECBD_4022	Isocitrase OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=ECBD_4022 PE=4 SV=1
A0A140NA70	ECBD_2927	Cytochrome d ubiquinol oxidase, subunit II OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=ECBD_2927 PE=3 SV=1

A0A140NBQ4	ECBD_2131	HipA N-terminal domain protein OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=ECBD_2131 PE=3 SV=1
A0A140NCI6	tuf	Elongation factor Tu OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=tuf PE=3 SV=1
A0A140N6C1	gyrB	DNA gyrase subunit B OS=*Escherichia coli* (strain B / BL21-DE3) OX=469008 GN=gyrB PE=3 SV=1

Extended Data Fig. 1 *In vitro* characterization of the enzymatic activity of cGAS.

a, HPLC analysis of nucleotide second messenger synthesis by cGAS using various nucleotide substrates. cGAS synthesizes a product only in the presence of both ATP and GTP. Data are representative of 3 independent experiments. **b**, cGAS synthesis activity is dependent on Mn^{2+}. Data are representative of 3 independent experiments. **c**, Optimal Mn^{2+} concentration for cGAS. The products synthesized at various Mn^{2+} concentrations were quantified by HPLC. Data are mean ± s.d. for $n = 3$ independent replicates and are representative of 3 independent experiments. **d**, Comparison of the activities of

cGAS and its D84A/D86A mutant. Data are representative of 3 independent

experiments. **e**, cGAS product was degraded by nuclease P1 but not CIP, as analyzed

by HPLC. Data are representative of 3 independent experiments. **f**, Comparison of the

retention time of possible cGAMP variants (3′,3′-cGAMP, 2′,3′-cGAMP, and

3′,2′-cGAMP) and the cGAS product. Data are representative of 3 independent

experiments. **g**, MS/MS fragmentation spectra of the cGAS product (top) and the

standard 3′,2′-cGAMP (bottom), and the chemical structure of 3′,2′-cGAMP. **h**,

Michaelis-Menten kinetics plot of ATP and GTP for cGAS. Data are mean ± s.d. for *n*

= 3 independent replicates and are representative of 3 independent experiments.

Extended Data Fig. 2 Predicted structure of the cGAS-E2 complex. a, SDS-PAGE

analysis of the *in vitro* assembly of the covalent cGAS-E2 protein. Two proteins were

mixed at a molar ratio of 1:1 and incubated at 4 ºC for the indicated periods (cGAS+E2).

Since both cGAS and E2 were His-tagged during the *in vitro* reconstruction, the molecular weight of cGAS-E2 protein was slightly higher than that obtained *in vivo* from His-tagged cGAS and non-tagged E2. Data are representative of 3 independent experiments. **b**, Ribbon diagram of the top ranked structure prediction of cGAS-E2 complex colored by the per-residue LDDT scores. **c**, Ribbon diagram of the top ranked structure prediction of cGAS-E2 complex with cGAS colored in green and E2 colored in orange. **d-e**, Chromatography traces and SDS-PAGE analysis of purified cGAS-E2 complex. **f**, Representative cryo-EM micrograph of cGAS-E2 complex.

Extended Data Fig. 3 Cryo-EM workflows and the quality of reconstructed cryo-EM maps. a-b, Workflow of 3D reconstruction of the initial cGAS-E2 dataset (**a**) and the UltrAuFoil cGAS-E2 dataset (**b**). **c**, Directional FSC of the cGAS-E2 cryo-EM density map. **d**, Orientation distributions of cGAS-E2 3D reconstruction. **e,** Local resolutions of the cGAS-E2 cryo-EM map.

Extended Data Fig. 4 Structural comparison between experimental and predicted

cGAS-E2. Superimposed predicted and experimental structures of cGAS-E2.

Extended Data Fig. 5 Sequence alignment of cGAS and E2 homologs. a, Sequence

alignment of the C-terminal residues of cGAS and CD-NTase homologs from other E2-

CBASSs. A conserved C-terminal glycine is marked by a red triangle. **b**, Sequence

alignment of E2 and its orthologs shows two conserved cysteine residues (marked by

red triangles).

Dataset I: Continuous density between G406 (cGAS) and K159 (E2)

Dataset II: Continuous density between G406 (cGAS) and C101 (E2)

Extended Data Fig. 6 Magnified view of the E2 active site in cGAS-E2 complex, showing continuous density between G406 of cGAS and K159 or C101 of E2.

Extended Data Fig. 7 Two crucial histidine residues in E2 active site. a, SDS-PAGE analysis showed that the thioester bond (-DTT) and isopeptide bond (+DTT) in cGAS-E2 are interconvertible at various pH. Data are representative of 3 independent experiments. **b**, SDS-PAGE analysis of the *in vitro* processing of cGAS-MBP by E2 or E2 (C101A) or E2 (H87A). **c**, Sequence alignment of E2 homologs from E2-CBASSs and E1E2/JAB-CBASSs showed that they shared low homology. The two crucial

histidine residues (marked by red triangles) are conservative in E2 from E2-CBASSs

but not E1E2/JAB-CBASSs. **d**, Superimposed structures of E2 (orange) and human E2

(gray, PDB ID 4AUQ) showed that no similar catalytic site was found in human E2.

Extended Data Fig. 8 Regulation of cGAS activity by E2-mediated poly-

cGASylation. a, HPLC analysis and quantification of the 3′,2′-cGAMP synthesized by

cGAS and variants as shown in **Fig. 4a**. 20-μL standard reactions contained 800 nM

enzyme, 250 μM ATP and 250 μM GTP in reaction buffer with 50 mM Tris-HCl (pH

7.5), 100 mM NaCl, 2.5 mM MnCl$_2$, and 1 mM DTT. Reactions were incubated at 37

ºC for 30 min, inactivated at 80 ºC for 10 min, and centrifuged for 15 min at 21,000 ×

g to remove precipitated protein. **b**, HPLC analysis and quantification of the

3′,2′-cGAMP synthesized by cGAS and variants as shown in **Fig. 4b**, with

concentration of ATP or GTP or Mn^{2+} in the reaction reduced by half compared to those

in (**a**), respectively. Data are mean ± s.d. for *n* = 3 independent replicates and are

representative of 3 independent experiments.

Extended Data Methods

Multiple Sequence Alignments

Alignment of cGAS or E2 homologs was performed using Clustal Omega and Jalview

version 2.11.2.5. Matrix, BLOSUM; Open gap penalty, 10; Extending gap penalty, 0.05;

End gap penalty, 10; Separation gap penalty 0.05.

Steady-state kinetic measurement

For steady-state kinetic measurements of cGAS, the standard 20 μL-reaction mixture

included 50 mM Tris-HCl (pH 7.5), 2.5 mM $MnCl_2$, 100 mM NaCl, 1 mM DTT, 1 μM

cGAS, 1 mM ATP (or GTP), and various concentrations (15.6, 31.3, 62.5, 125, 250,

500 and 1,000 μM) of GTP (or ATP). For each nucleotide concentration, samples were

incubated at 37 ºC for 15 min. Reactions were immediately heat-inactivated at 80 ºC

for 10 min and the samples were centrifuged for 15 min at 17,000 × g to remove

precipitated protein. Of each sample, 10 μL was injected and reactions were analyzed

by HPLC, as detailed above. Absorbance units were converted to μmol/L by comparing

to a standard curve from 7.8 μM to 1 mM of chemically-synthesized 3′,2′-cGAMP

(Biolog Life Sciences). Data were fitted by linear regression, and non-linear curve

fitting Michalis-Menten kinetics and allosteric sigmoidal were calculated using

GraphPad Prism version 8.0.2.

Mass spectrometry

To characterize the product of cGAS, we performed LC-MS/MS on a Thermo

Scientific™ UltiMate™ 3000 system coupled to a Thermo Scientific Orbitrap LC/MS

(Q Exactive), using a Thermo Hypersil GOLD C18 column (100 mm × 2.1 mm, 3 μm)

maintained at 25 ºC with a flow rate of 0.25 mL/min. The mobile phase consisted of

methanol (A) and 20 mM ammonium acetate (B). The HPLC gradient was as follows:

0–7 min, 2% A; 7–12 min, 2–30% A. The column was re-equilibrated for 7 min at 2%

A. Detection was performed in positive ionization mode using an electrospray

ionization (ESI) source under the following parameters: spray voltage, 3.2 kV; sheath

and auxiliary gas flow rates, 40 and 15 arbitrary units, respectively; max spray current,

100.00 μA; S-Lens RF Level, 50%; capillary temperature, 300 ºC; probe heater

temperature, 350 ºC. Profile MS1 spectra were acquired with the following settings:

mass resolution, 70,000; AGC volume, 3×10^6; maximum IT, 100 ms; scan range, 300–

1,000 m/z. Acquisition of data-dependent MS/MS spectra was performed using

collision-induced dissociation (CID) with the following settings: mass resolution,

17,500; AGC volume, 1×10^5; maximum IT, 50 ms; loop count, 10; isolation window,

4.0 m/z; normalized collision energy, 20, 40, and 60 eV. Data are reported for the z = 1

acquisition for each indicated cyclic nucleotide. The chemical structures were drawn

using ChemDraw version 19.0.

Proteomics analysis

Proteomic analysis is performed by a commercial company vendor (Genecreate). The

protein samples were mixed with NuPAGE LDS sample buffer (Invitrogen) and

separated using NuPAGE 4 to 12%, Bis-Tris gel. The corresponding gel bands were

then cut and washed with 40% H_2O, 60% acetonitrile and 50 mM NH_4HCO_3 (pH 8.0),

and the protein bands to be identified were reduced and alkylated using TCEP and

chloroacetamide respectively. The gel pieces were digested with trypsin (20 ng/mL) in

50 mM NH_4HCO_3 (pH 8.0) at 37 ºC overnight. The digested peptides were extracted

with 40% H_2O, 60% acetonitrile and 1% formic acid, dried and reconstituted with 1%

formic acid for LC-MS/MS analysis. The tryptic peptides (100 ng) were analyzed on a

UltiMate 3000 RSLCnano coupled to an Q Exactive HF mass spectrometer (Thermo

Fisher Scientific). The peptides were firstly loaded onto a 75 μm×2 cm trap column and

separated on a 75 μm×25 cm Pepmap C18 analytical column (Thermo Fisher Scientific)

with a binary buffer system. Buffer A was 0.1% formic acid and 3% DMSO in 100%

H_2O and buffer B was 0.1% formic acid and 3% DMSO in 80% acetonitrile with 20%

H_2O. The Eclipse mass spectrometer was operated in data-dependent acquisition mode

with one full MS scan followed by MS/MS scans with higher-energy collision-induced

dissociation (HCD) fragmentation. The typical MS settings were spray voltage of 2.2

kV, heated capillary temperature of 320 ºC, HCD energy of 27. The LC-MS/MS RAW

data were processed with Maxquant (version 2.1.2.0) for protein identification using a

tailored database containing the sequences of E2 and cGAS and the *Escherichia coli*

strain B/BL21-DE3 proteomes database in Uniprot. The typical search parameters were

enzymatic specificity of Trypsin/P with maximum of two missed cleavages and variable

modifications of oxidation on methionine. Carbamidomethyl (Cys) was set as the fixed

modification. All results were screened at 1% protein and peptide level FDR. Only high

confidence identified peptides were chosen for downstream protein identification

analysis.