Structural determinants of relaxation dynamics in chemical reaction networks

Yusuke Himeoka[a,b], Julius B. Kirkegaard[b], Namiko Mitarai[b], and Sandeep Krishna[c]

[a]Universal Biology Institute, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku , Tokyo, Japan; [b]The Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, Copenhagen, 2100, Denmark; [c]Simons Centre for the Study of Living Machines, National Centre for Biological Sciences TIFR, GKVK Campus, Bellary Road, Bangalore 560065, India

Understanding the relationship between the structure of chemical reaction networks and their reaction dynamics is essential for the functioning of living organisms. However, while some network-structural features are known to relate to the steady-state characteristics of chemical reaction networks, mathematical frameworks describing the links between out-of-steady-state dynamics and network structure are still underdeveloped. Here, we examine a class of the chemical reaction networks consisting of the ligation and splitting reactions of the polymers. These networks satisfy mass-conservation and other key properties of chemical reaction networks. We simulated the dynamics of these chemical reactions, focusing on the relaxation dynamics after perturbation from a steady-state. We then find three types of relaxation dynamics: exponential, power-law, and plateau. We show that three features computed from a submatrix of the network's stoichiometric matrix are sufficient to determine the type of the dynamics, namely, (i) the rank, determining the existence of a steady-state; (ii) the left null-space, being related to conserved quantities in the dynamics; and (iii) the stoichiometric cone, dictating the range of achievable chemical concentrations. We further show that these three quantities also predict the type of relaxation dynamics of combinations of our minimum networks, larger networks with many redundant pathways, and a real example of a metabolic network. The unified method to predict the qualitative feature of the relaxation dynamics presented here will provide a basis for unveiling the design principles of the metabolic reaction networks.

chemical reaction| network | dynamics | metabolism

Origins of chemical reaction network theory can be found in the discovery of the law of mass action (1). Since then, mathematical laws governing chemical reactions have been uncovered, such as Le Chatelier principle (2), the Arrhenius equation (3), and the detailed-balance condition for the feasibility of chemical equilibrium (4).

For the development of quantitative theories of living entities, a deep understanding of the out-of-steady-state dynamics of chemical reactions is indispensable. Until roughly a decade ago, a large part of the experimental studies of microbial physiology focused on the steadily-growing cellular populations. In such situations, the assumption that the concentrations of the cellular metabolites are constant (at the steady-state) is reasonable, and static metabolic analysis such as Flux Balance Analysis (5–10) and the steady-state cell models (11, 12) have provided the mathematical description of stable cell growth.

However, recent experiments have been unveiling the dynamic nature of cellular physiology. Especially, the physiology of starved *Escherichia coli* cells are gathering much attention; The lag time — the duration that the cell restart growth after the substrate replenishment — was shown to depend on how long the cells are starved (13–17) and the death rates of the starved cells differ depending on the previous culture

conditions even though the starvation condition is identical (18). Such "memory" is a unique feature of out-of-steady-state systems. In addition, the time scales of the memories are much longer than that in the steadily-growing cells. It is therefore desirable to develop theoretical frameworks to understand how such slow relaxation dynamics and long-term memory emerge from the structure of chemical networks.

There are two broad types of approaches one might take to explore the link between the network structure and the relaxation characteristics of chemical networks. One is to examine random networks with structural characteristics (such as density of links and degree distribution) identical to natural chemical networks like metabolic networks (19–22). This approach enables us to explore a large class of network structures, but due to the simplification into a random network, it may miss crucial properties of real chemical networks. For example, the central principle of chemical reactions, the law of mass conservation, is often violated. Thus, any structure-dynamics connections uncovered by these approaches may not always apply to real chemical networks.

On the other hand, data-driven approaches such as reaction pathway prediction by using machine learning (23–25) generate realistic chemical reaction networks, which never violate mass conservation and other necessary properties. However, it is too computationally costly to explore a wide variety of possible chemical reaction networks. The generality of structure-dynamics connections, or lack of it, uncovered by

---

Significance Statement

The relationship between network structure and chemical reaction dynamics is of central interest in chemical reaction network theory, as it underlies chemical manufacturing, cellular metabolism, and bioengineering. The links between structure and steady-state properties have been extensively investigated. However, how far the network structure determines the out-of-steady-state, transient dynamics of chemical reactions is unexplored. Here we construct a chemical reaction network model that is simple but generate a wide variety of network instances. By exploring the networks' structural- and dynamical features, we found that three network-structural features are sufficient to predict the qualitative characteristics of the relaxation dynamics after the chemical concentrations are perturbed from their steady-state. Our findings provide a foundation for the structure-based prediction of chemical reaction dynamics.

Please provide details of author contributions here.

Please declare any competing interests here.

[1]A.O.(Author One) contributed equally to this work with A.T. (Author Two) (remove if not applicable).

[2]To whom correspondence should be addressed. E-mail: author.twoemail.com

these approaches may therefore be unclear.

Here we develop an alternative approach. We construct a class of the chemical reaction network consisting of chemical species that can be represented as monomer sequences (polymers), and chemical reactions that ligation (concatenation) and splitting of these polymers. The model can generate a wide variety of chemical reaction network structures. Yet, all the generated networks satisfy the mass conservation principle.

Microbial metabolic networks were one of our motivations for studying relaxation dynamics. Such metabolic networks, as well as chemical networks used in industrial manufacturing processes, have the specific function of converting certain input substrates into final product chemicals. For instance, in bacterial metabolism, carbon, nitrogen, phosphorous sources (e.g., glucose, potassium hydrogen phosphate, ammonium chloride) and other miscellaneous molecules are converted into biomass such as proteins, nucleic acids, and lipids. Therefore, we impose a further constraint on the networks we study - the ability to synthesize a given set of final products from a specified set of input substrates. The resultant chemical reaction networks we generate are, of course, a subset of all possible networks, but they form a well-defined subset so the generality of the structure-dynamics relationships we find is also well-defined. Our framework can also easily be extended to include other kinds of chemical species and reactions as desired.

We find that for these artificial reaction networks we construct, there is a single stable steady-state for the chemical concentrations. Computational simulations of the chemical reaction dynamics upon perturbation of these steady-states revealed four distinct types of relaxation dynamics. Interestingly, we found that the qualitative features of the relaxation dynamics are determined by only three properties of the stoichimetric matrix which encodes the chemical network structure, namely, (i) the rank, (ii) the dimension of the left null space, and (iii) the reachability of the final steady-state. Crudely, a non-zero rank gap indicates that no steady-state exists in the absence of any degradation of chemicals, and the left null space determines the existence of conserved quantities in the dynamics. The reachability of the final steady state appears to be a dynamical, and not structural, condition. However, we show that in fact it is determined entirely by the stoichiometry of the chemical reactions (more precisely, the 'stoichiometric cone', described in detail in later sections), a purely structural property. Finally, we tested the applicability of these three network-structural features to the prediction of relaxation dynamics of a real biological (metabolic) network. Our results and the general framework we have developed provides valuable tools for exploring the relationship between the network structure and chemical reaction dynamics, which may be applied both to complex industrial manufacturing processes as well as design of synthetic microorganisms designed to perform certain chemical tasks (26, 27).

Model

Constructing a class of artificial reaction networks. For the construction of the chemical reaction networks, we define a set of chemicals $\mathscr{C}$ and the set of the reactions $\mathscr{R}$ they participate in. In our artificial reaction network model, we consider each chemical species to be a one-dimensional chain of the monomers, such as AA or $AABA$ where A and B represent

different monomer species. Each monomer idealizes chemical elements such as hydrogen, carbon, nitrogen, and oxygen, or even larger chemical moeities. The chemical reactions possible are the splitting or ligation of the chemicals. For instance, there are three splitting reactions possible with $AABA$ as the substrate: $AABA \rightarrow A + ABA$, $AABA \rightarrow AA + BA$, and $AABA \rightarrow AAB + A$. In the present manuscript, we suppose that all reactions are reversible. That is, if there is a splitting reaction $AABA \rightarrow A + ABA$, the corresponding ligation reaction $A + ABA \rightarrow AABA$ exists.

With this set-up, the set of all possible chemicals $\mathscr{C}$ and the set of all possible reactions $\mathscr{R}$ are fully determined by setting the number of monomers M and the length of the longest polymer L. For instance, in the case of $M = 2$ and $L = 2$, $\mathscr{C}$ and $\mathscr{R}$ are given by

$$
\begin{aligned}
\mathscr{C} &= \{A, B, AA, AB, BB\}, \\
\mathscr{R} &= \{AA \rightleftharpoons 2A, AB \rightleftharpoons A + B, BB \rightleftharpoons 2B\}.
\end{aligned}
$$

To avoid redundancy, we omit the direction of the polymers, e.g., the polymer BA is identified as identical to AB. Given this, the number of chemicals ($|\mathscr{C}|$) and the number of reactions ($|\mathscr{R}|$) scale with the parameters M and L as

$$
|\mathscr{C}| \sim \sum_{l=1}^{L} M^l \sim M^L, \ |\mathscr{R}| \sim \sum_{l=1}^{L} (l-1)M^l \sim LM^L.
$$

We now construct individual chemical reaction networks by choosing appropriate subsets of $\mathscr{R}$. Importantly, as each reaction in $\mathscr{R}$ satisfies the mass-conservation principle, all the reaction networks made by combining the reactions from $\mathscr{R}$ satisfy this principle. The number of possible reaction networks formed by choosing a subset of $\mathscr{R}$, including disconnected ones, is roughly given by $2^{|\mathscr{R}|} \sim 2^{LM^L}$. This number grows astonishingly fast with M and L (256 for $M = 2$ and $L = 2$, but $\approx 10^{19}$ for $M = 2$ and $L = 4$). Thus, we focus only on the *minimum* networks which can produce a given set of N_p final product chemicals from a given set of N_s input substrates (in metabolic terms, those that can produce biomass from the available nutrients).

We define a minimum network to be the set of reactions $\mathscr{R}_0 \subset \mathscr{R}$ with the smallest $|\mathscr{R}_0|$ which can synthesize all the final products from the input substrates. In contrast a *minimal* network is one from which no reaction can be removed while still maintaining production of all final products from the input substrates. There are many minimal (but not minimum) networks where the number of reactions $|\mathscr{R}_0|$ is not the global minimum. Note also that we must choose the input substrates and final products such that the network consisting of all reactions in $\mathscr{R}_0$ satisfies the condition of being able to synthesize the final products from the input substrates, otherwise no subset will be able to do so. However, this 'fully connected network' is typically very far from being minimal, let alone minimum. Finding minimum reaction networks, where $|\mathscr{R}_0|$ is the global minimum, is computationally challenging. Here we exploit the fact that this computation can be represented in the form of the 3-satisfiability (3-SAT) problem. Being an NP-complete problem, no polynomial time algorithm exists (28). However, sophisticated 3-SAT solvers are available which work very well in practice — in particular for the present problem minimum networks (see Materials and Methods for more details).

In the following, we deal with the minimum networks generated from the fully-connected network with $M = 2$ and $L = 6$, with $N_s = 2$ and $N_p = 1$. The fully-connected network and examples of minimum networks are depicted in Fig.1 A and B, respectively.

Defining the chemical reaction dynamics of the constructed networks. There are multiple choices for implementing the kinetics of a given chemical reaction network. In the present manuscript, we utilize the simplest and widely used formulation — mass-action kinetics. Suppose that there is a following reaction with n_s and n_p reactants of the forward- and backward reaction, respectively.

$$\sum_{i=1}^{n_s-1} \sigma_{s(i)}^+ X_{s(i)} \rightleftharpoons \sum_{i=0}^{n_p-1} \sigma_{p(i)}^- X_{p(i)}.$$

By using mass-action kinetics, the rate of the reaction (also called the 'flux') is given by

$$J = v_+ \prod_{i=0}^{n_s-1} x_{s(i)}^{\sigma_{s(i)}^+} - v_- \prod_{i=0}^{n_p-1} x_{p(i)}^{\sigma_{p(i)}^-}, \qquad [1]$$

where the capital letter X_i indicates the ith chemical, and its concentration is denoted by the corresponding lower case letter. $s(i)$ and $p(i)$ represent the index of the ith reactant of the forward- and backward reactions, respectively. $\sigma_i^\pm$ specify the stoichiometric coefficients of the ith chemical in the forward- (σ_i^+) and backward (σ_i^-) reaction, respectively. $v_\pm$ indicates the rate of the forward (v_+) and the backward (v_-) reaction. When we summarize the information of the whole reactions by using the reaction stoichiometric matrix, the relationship between $\sigma_{ir}^\pm$ (now we added the index of the reaction r) and the stoichiometric matrix S is given by

$$\sigma_{ir}^+ = \begin{cases} -S_{ir} & (S_{ir} < 0) \\ 0 & (\text{otherwise}), \end{cases} \qquad [2]$$

$$\sigma_{ir}^- = \begin{cases} S_{ir} & (S_{ir} > 0) \\ 0 & (\text{otherwise}). \end{cases} \qquad [3]$$

Each (minimum) reaction network we construct can be represented by a corresponding reaction stoichiometry matrix S_0, which denotes the relationship of the interconversion among the chemicals by the splitting and ligation reactions. The exponents in the mass-action kinetics (Eq. (1)), $\sigma_{ir}^\pm$'s, are set by S_0 according to Eq. (2) and 3. This matrix however does not represent the processes of uptake of input substrates and consumption or secretion of the final products. Therefore, we add the uptake reaction $\tilde{X}_i^{(\text{sub})} \rightleftharpoons X_i^{(\text{sub})}$ for N_s substrates and the product synthesis reaction from N_p product precursors $\sum_{i=0}^{N_p-1} X_i^{(\text{prp})} \rightleftharpoons \tilde{X}^{(\text{pr})}$. Here, the symbols with tildes represent the chemicals in the external environment, and "sub", "prp", and "pr" are abbreviations of the (input) substrates, (final) product precursors, and the final products. The kinetics of each of these reactions is also implemented by the mass-action kinetics given by

$$J_i^{(\text{sub})} = u_+^{(i)} \tilde{x}_i^{(\text{sub})} - u_-^{(i)} x_i^{(\text{sub})}, \ (0 \le i < N_s) \qquad [4]$$

$$J^{(\text{pr})} = p_+ \prod_{i=0}^{N_p-1} x_i^{(\text{prp})} - p_- \tilde{x}^{(\text{pr})} \qquad [5]$$

We further suppose that all chemical species are spontaneously degraded or deactivated at a uniform rate ϕ which is sufficiently smaller than the rates of the other chemical reactions. Additionally, the external environment is large enough so the concentrations of chemicals there are set to be constant. Then, by summing up, the dynamics of the concentrations of chemical species follow the equation below

$$\dot{\boldsymbol{x}} = S_0 \boldsymbol{v}_0 + \boldsymbol{u} - \boldsymbol{p} - \phi \boldsymbol{x}, \qquad [6]$$

where $\dot{\boldsymbol{x}}$ represents the time-derivative of $\boldsymbol{x}$, $d\boldsymbol{x}/dt$. S_0 is the stoichiometry matrix of the network. $\boldsymbol{v}_0$ represents the reaction flux vector and each element has the form of Eq. (1). $\boldsymbol{u}$ and $\boldsymbol{p}$ denote the vectors representing the rate of substrate uptake and product synthesis, respectively. If the ith chemical X_i is one of the substrates (product precursors), the ith element of the vector $\boldsymbol{u}$ ($\boldsymbol{p}$) is given by the nonzero entity $J_i^{(\text{sub})}$ ($J^{(\text{pr})}$).

Once a reaction network and the substrates and products are specified, the differential equation (Eq. (6)) is uniquely given. In this sense, we call the Eq. (6) the accompanying ordinary differential equation (ODE) of the reaction network. For simplicity, we set all the rate constants except ϕ to unity. Also, we suppose that the synthesized product is harvested quickly, and thus, the external concentration of the product ($\tilde{x}^{(\text{pr})}$ in Eq. (5)) is set to zero.

Relaxation behaviours and network structures

In the following sections, we study the link between the network structure and the relaxation behaviour of the accompanying ODEs. First, we generate the reaction networks and check if there is a steady-state attractor. As far as we have confirmed, all the networks have a unique fixed-point attractor for each. For probing the relaxation characteristics, we applied perturbations to the attractor. Since the concentrations of the chemicals span a wide range at the attractors, we apply perturbations in a multiplicative manner. In addition, to normalize the strength of the perturbations, we generated the initial points on the surface of the $|\mathscr{C}_0|$-dimensional hypersphere, centred at the steady-state attractor, with a given diameter D. In concrete terms, a single initial point $\boldsymbol{x}_{\text{ini}}$ is given by $\ln \boldsymbol{x}_{\text{ini}} = \ln \boldsymbol{x}_{\text{att}} + D \cdot \boldsymbol{r}/\|\boldsymbol{r}\|$ with $\boldsymbol{x}_{\text{att}}$ is the steady-state attractor of the model, and $\boldsymbol{r}$ is a vector of the random numbers each lying in the interval [01], respectively (see Materials and Methods section for more details).

We simulated the accompanying ODE (Eq. (6)) of 2054 minimum networks with $N_{\text{ptb}}(= 256)$ initial points for each network. Visually, we found three typical types of relaxation: exponential, plateau, and power-law type. Examples of each are shown in Figs.1C-F. There appeared to be three groups in terms of classification of the networks; (i) the networks exhibiting only exponential relaxation regardless of the initial concentrations, (ii) the networks showing only the plateau type relaxation regardless of the initial concentrations, and (iii) the networks displaying either the plateau or the power-law type of relaxation depending on the initial concentrations. The plateau dynamics are observed commonly in the second and the third network groups. At this stage, we do not know if the plateau exhibited by the second and the third type of the networks are different in any sense, but for later arguments we refer to them as "Always Plateau" (independent of initial conditions) and "Conditional Plateau" (dependent on initial conditions) in Figs. 1D and E.

To make the classification more quantitative, we focused on two quantities: the relaxation time and the migration length. Here we define the relaxation time as the time point at which the Euclidean distance, in logarithmic scale, between the state and the steady-state attractor first becomes smaller than a certain threshold value ϵ [*], that is

$$T_{\mathrm{relax}} = \min\{t \mid \|\ln \boldsymbol{x}(t) - \ln \boldsymbol{x}_{att}\| < \epsilon\}. \qquad [7]$$

Next, to distinguish between the plateau and the power-law relaxation types, we quantified the total length of the trajectory in concentration space during a given time period. It is measured by the 'migration length' defined as

$$L = \int_a^b \left\| \frac{\mathrm{d}\ln \boldsymbol{x}}{\mathrm{d}t} \right\| dt, \qquad [8]$$

where we choose $a = 100$ and $b = 0.01/\phi$ in order to quantify the "motion" of the trajectory in the intermediate time scale (sufficiently larger than the time scale of the reaction rate constants and sufficiently smaller than that of the degradation). The migration length is computed in the logarithmic scale to take the effect of low-concentration chemicals into account. Additionally, we compute $L_{\max}$ given by Eq. (8) with 10^{10} (the end of the simulation) as the upper bound of the integral.

Our prior expectation, from visual examination of the trajectories (e.g., Figs.1C-F), was that since plateau type trajectories are almost static in the time range $t \in [100, 0.01/\phi]$, the normalized migration length $L/L_{\max}$ would have a relatively lower value, compared to that for power-law type dynamics. We also expected that exponential relaxation dynamics will result in lower T_{relax} values than the other two types.

Examples of the distribution of migration length of two sets of networks are presented in Fig. 1G. The upper panel is is the distribution over all initial conditions for one network that shows always-plateau dynamics, and the bottom panel is for one network exhibiting both conditional-plateau and power-law relaxation. The bottom distribution shows a clear double-peak due to bimodality of the relaxation dynamics, while the upper distribution is rather unimodal. We suppose that if networks exhibit a bimodal distribution due to the presence of both power-law relaxation in addition to conditional-plateau relaxation, the average value of the normalized migration length $L/L_{\max}$ (averaged over all initial conditions) would be larger than the corresponding value for always-plateau networks. Thus, we utilize $\langle L/L_{\max}\rangle$ as an indicator of the variety of the relaxation behaviour.

The average relaxation time and the average migration length are computed for each network, and we plotted the distribution over the networks. As shown in Fig.1H, the distribution has four distinct peaks, and we classified the networks from type-I to type-IV according to the nearest peak. While the separation of the type-II and type-IV peaks is less clear, the distribution-based classification of the networks matched very well to the visual grouping of the models: the type-I networks (small T_{relax} and large $L/L_{\max}$) exhibit only exponential relaxation; most of the type-II networks showed always-plateau relaxation; and, the type-III networks and most type-IV networks exhibited both the conditional-plateau and the power-law type relaxation depending on the initial concentrations.

Note that the final steady-state attractor is reached at the latest by $t \sim 1/\phi$, which is the timescale of the spontaneous degradation, and the qualitative differences between the different types of relaxation only appear in $t \ll 1/\phi$ region. These observations mean that the difference in the relaxation behaviour during $t \ll 1/\phi$ highlight structural features of the reaction networks and would not be changed by varying the value of $1/\phi$ as long as it is sufficiently large. What structural features of the networks result in the four different relaxation types observed in Fig.1H? We now show that the relaxation behaviour is indeed determined by three features of the stoichiometric matrix (which in turn encodes the structure of the network), namely, (I) the rank gap, (II) dimension of the left null-space, and (III) the stoichiometric cone.

I. The rank gap. The distinct relaxation behaviours are observed over timescales shorter than $1/\phi$, and thus, should stem from the dynamics (Eq. (6)) without the degradation term. The clear difference between the exponential relaxation and the others is the relaxation time. The exponential type trajectories already reach the attractor well before t exceeds $1/\phi$. We now show how measuring the ranks of the stoichiometric matrix and some enlargements can inform us about the reachability to the attractor in the absence of the degradation term.

First, we introduce additional notation. Let S_1 denote the enlarged matrix $(S_0|\boldsymbol{U}^{(0)}|\boldsymbol{U}^{(1)}|\cdots|\boldsymbol{U}^{(N_n-1)})$, where $\boldsymbol{U}^{(j)}$'s are vectors of the stoichiometry of the substrate uptake reactions, i.e., $U_i^{(j)}$ is unity if the ith chemical is the jth input substrate and zero otherwise. By vertically stacking $\boldsymbol{v}_0$ and $\boldsymbol{u}$ in Eq. (6) and denoting it as $\boldsymbol{v}_1$, we can rewrite Eq. (6) with S_1 instead of S_0. Then, removing the degradation term, we obtain

$$\dot{\boldsymbol{x}} = S_1\boldsymbol{v}_1 - \boldsymbol{p}. \qquad [9]$$

Note that Eq. (9) represents the dynamics of chemical concentration driven by the substrate uptake, internal reactions, and the product synthesis. This equation approximates the original equation Eq. (6) for $t \ll 1/\phi$.

For there to exist a steady-state of this equation, there must be a pair of vectors, $\boldsymbol{v}_1^*$ and $\boldsymbol{p}^*$, satisfying $S_1\boldsymbol{v}_1^* = \boldsymbol{p}^*$. Since we are not interested in the trivial solution $\boldsymbol{v}_1^* = \boldsymbol{0}$, the necessary and sufficient condition for the existence of a steady-state is that the rows of S_1 and $\boldsymbol{P}$ are linearly dependent, i.e, $\mathrm{rank}S_1 = \mathrm{rank}(S_1|\boldsymbol{P})$, where $\boldsymbol{P}$ is the stoichiometry of the final product synthesis (i.e., the ith element is unity if the ith chemical is one of the product precursors and zero for the rest).

We found that the the rank gap $\delta \equiv \mathrm{rank}(S_1|\boldsymbol{P}) - \mathrm{rank}S_1$ for all type-I networks type-I networks (showing only the exponential relaxation) is zero. Interestingly, all type-II networks also satisfied the $\delta = 0$ condition. On the other hand, all the type-III and type-IV networks have $\delta = 1$, i.e., in the absence of degradation no steady-state exists.

This implies that there are two types of plateau relaxation. The plateaux exhibited by type-II networks are the steady-states of the model equation without degradation (Eq. (6)). However, the plateaux exhibited by type-III and IV networks are not the steady-state of Eq. (6), but are extremely slow transient dynamics towards the final steady-state for the model *with* degradation. Indeed, we find that the type-II plateaux correspond to what we had earlier termed "always-plateau",

Fig. 1. (A) The fully-connected network for $M = 2$ and $L = 6$. (B) Two examples of the minimum network which synthesizes a single final product from two substrates. The substrates and product chemicals are indicated by the red arrows and the blue arrow, respectively. In A and B, the bipartite representation of the chemical reaction network is adopted. For example, the reaction $S_1 + S_2 \rightarrow P$ is depicted as follows (see inset of the panel B); first, the chemical nodes representing S_1 and S_2 (the gray disks) are wired to the reaction node (the pink diamond), and then, an edge connects the reaction node and the chemical node of the chemical P (the gray disk). Directions of the reactions are chosen to be consistent with the steady-state flux distribution. (C to F) Four typical relaxation dynamics emerged from the minimum networks: Exponential, Always-Plateau, Conditional-Plateau, and Power-law relaxation. The vertical and horizontal axes are on a logarithmic scale. (G) Example distributions of the migration length (see text for definition). The top panel is the distribution over all initial conditions for one network showing always-plateau type relaxation, and the bottom panel is from a network showing both the plateau and power-law type relaxations. (H) Classification of the minimum networks based on the average relaxation time (see text for definition) and the average migration length.

while the type-III and IV plateux corresond to what we called "conditional-plateau".

II. The left null-space. The rank gap does not distinguish between type-I and type-II networks (which both have $\delta = 0$). Fortunately, these networks exhibit a unique type of relaxation regardless of the initial concentrations (exponential or always-plateau). Thus, we can expect that the difference stems from the network structure.

Fig.2 shows two time courses of the same type-II network but starting from different initial concentrations. The plateau regions differ not only in the concentrations of the chemicals but also in the rank order of the chemical concentrations (the identical color is used for each chemical species in both figure panels).

Fig. 2. Two time courses emerged from the same minimum type-II network when started with different initial concentrations. Both the vertical and horizontal axes are plotted in the logarithmic scale.

Considering that each plateau corresponds to the steady-state of $\dot{\boldsymbol{x}} = S_1 \boldsymbol{v}_1 - \boldsymbol{p}$ and the model eventually reaches a unique final attractor, possible mechanisms to generate the variations are limited. We hypothesized that such type-II networks might be have combinations of chemical concentrations that, over timescales much shorter than $1/\phi$, are constrained to be almost-conserved. If different initial conditions have different values for such conserved quantities, the system might evolve towards a different steady-state of $\dot{\boldsymbol{x}} = S_1 \boldsymbol{v}_1 - \boldsymbol{p}$ for each of these. We therefore computed the dimension of the left null-space of the enlarged stoichiometry matrix, $I := \dim(\mathrm{coker}) S_2$, where $S_2 = (S_1 | \boldsymbol{P})$, because the left null-space of the reaction stoichiometry represents the conserved quantities of the reaction dynamics. We found that the left null-space of all the type-I models was $\{0\}$. In contrast, all the type-II models had non-zero I[†]. Further, we found that the type-III models had no conserved quantity ($I = 0$), while the type-IV networks had at least one ($I \geq 1$).

So far, we have fully characterized the type-I and type-II networks. The type-I networks have no rank gap, $\delta = \mathrm{rank} S_1 - \mathrm{rank} S_2 = 0$, and no conserved quantity $I = \dim(\mathrm{coker}) S_2 = 0$. Type-II networks also have no rank gap but have at least one conserved quantity, $I \geq 1$.

III. The stoichiometric cone. Now we move onto the networks with $\delta = 1$[‡]. Due to the non-zero rank gap, there is no steady-state until t becomes comparable with $1/\phi$, i.e, the spontaneous degradation starts affecting the dynamics of the concentration. Thus, in these type-III and type-IV networks, the relaxation dynamics over timescales smaller than $1/\phi$ is not to some meta-stable almost-steady-state, but a slow transient dynamics towards the final attractor with degradation. With power-law relaxation there is a slow but continuous approach towards the final attractor, while with the conditional-plateau dynamics the trajectory is practically static for a long time before suddenly "jumping" to the final attractor.

We therefore hypothesized that in the power-law case, the attractor is easy to reach from the initial point, but the degradation term $\phi \boldsymbol{x}$ is necessary for making the final concentrations a stable steady-state. By contrast, in the conditional-plateau case getting close to the attractor may itself be hindered in the absence of the degradation term. To check this, we introduce a subset of the phase space defined as

$$SC(\boldsymbol{x}_{\mathrm{ini}}) = \{\boldsymbol{x} \in \mathbb{R}_+^{|\mathscr{C}_0|} | \boldsymbol{x} = \boldsymbol{x}_{\mathrm{ini}} + S_2 \boldsymbol{v}_2, \boldsymbol{v}_2 \in \mathbb{R}^{|\mathscr{R}|_0}, J^{(\mathrm{pr})} \geq 0\}.$$

$SC(\boldsymbol{x}_{\mathrm{ini}})$ is the region in the phase space "reachable" from the initial point by controlling the chemical reaction fluxes in any way except synthesizing back the product precursors from the final product. Note that in the definition, no flux balance (steady-state condition) is required. Due to the non-negativity condition ($J^{(\boldsymbol{pr})} \geq 0$) and the non-zero rank gap δ, $SC(\boldsymbol{x}_{\mathrm{ini}})$ does not coincide with the whole space $\mathbb{R}_+^{|\mathscr{C}_0|}$. This $SC(\boldsymbol{x}_{\mathrm{ini}})$ is equivalent to the stoichiometric cone in the field of chemical reaction network theory (29), and thus, we term $SC(\boldsymbol{x})$ the stoichiometric cone corresponding to the initial condition $\boldsymbol{x}$ in the following arguments.

We can now mathematically formulate our hypothesis on the relationship between reachability and relaxation behaviours: for the initial points whose stoichiometric cone $SC(\boldsymbol{x}_{\mathrm{ini}})$ contains the final attractor, relaxation becomes power-law-like, whereas, if the stoichiometric cone of the initial point does not include the attractor, plateaux appear during the relaxation (for the algorithm of judging the inclusion relationship, see Methods).

For grouping the trajectories into the (conditional-) plateau relaxation and the power-law relaxation types, we used the migration length defined earlier (Eq. (8)). Recall that if the normalized migration length $L/L_{\max}$ has a small value, it indicates the relaxation dynamics is plateau type, and if it has a large value, the relaxation is power-law type.

To examine the relationship between the relaxation type and the position of the initial concentrations in the phase space, we computed the average of the normalized migration length $L/L_{\max}$ separately for the initial concentrations whose stoichiometric cone does and does not contain the final attractor[§]. Fig. 3 is the scatter plot of the two average values

---

[†]In the present minimum reaction models, there is a lower limit of I for the models with $\mathrm{rank} S_1 = \mathrm{rank} S_2$, given by $M - N_n$, where M is the number of monomer types and N_n is the number of input substrates. For more detail, see SI Text

[‡]Since the product formation reaction is represented as the single reaction (Eq. (5)), the rank gap between S_1 and S_2 is one at most.

[§]Since the type-IV networks have conserved quantities, the stoichiometric cone of an initial point

for all the minimum networks of type-III and IV that we constructed. The migration length averaged over the initial concentrations with $x_{att} \in SC(x_{ini})$, $\langle L/L_{max}\rangle_{in}$, and with $x_{att} \notin SC(x_{ini})$, $\langle L/L_{max}\rangle_{out}$ are plotted in the horizontal and vertical axis, respectively. In all networks, $\langle L/L_{max}\rangle_{in}$ is larger than $\langle L/L_{max}\rangle_{out}$. This result supports our intuition that some trajectories are confined in a region of the phase space and cannot get closer to the attractor, resulting in the conditional-plateau dynamics.

Fig. 3. The normalized migration length is averaged separately for the time courses with the initial points whose SC, defined in the text, contains (vertical axis) and does not contain (horizontal axis) the attractor. The $y = x$ line is plotted as a guide for the eye. The distributions of $\langle L/L_{max}\rangle_{in}$ and $\langle L/L_{max}\rangle_{out}$ are shown along the horizontal and the vertical axis, respectively.

To summarize: Networks that have zero rank gap and no conserved quantity (type-I) always exhibit exponential relaxation dynamics. Networks that have zero rank gap but some conserved quantities (type-II) always exhibit plateau relaxation. Type III and type IV networks both have a rank gap of 1, but the former have no conserved quantities, while the latter have at least one. Both these networks exhibit either power-law relaxation or plateaux, depending on the initial conditions. For both types of networks, the type of relaxation for a given initial condition is determined by whether the final attractor is reachable from that initial point. If it is, then we observe power-law relaxation, and if it is not, then we observe plateau relaxation. Thus, from the stoichiometric matrices associated with a given network, by computing three network-structural quantities, namely, (i) the rank gap δ, (ii) the dimension of the left null space $I = \dim(\mathrm{coker}) S_2$, and (iii) the stoichiometric cone, we can predict whether the relaxation will be exponential, power-law or plateau-type. A graphical summary of this result is presented in Fig. 4.

---

never contains the attractor unless the values of the conserved quantities at the initial point and the attractor are equal. This situation never happens by applying random perturbations to generate the initial points. Thus, we first search for the nearest point from the initial point, $\tilde{x}_{ini}$, that has the same values as the conserved quantities with the attractor (see Materials and Methods). Then, we checked if the attractor is in the stoichiometric cone of $\tilde{x}_{ini}$.

Fig. 4. The summary of the relaxation type and network-structural features. Network types are classified according to the rank gap δ and the dimension of the cokernel of the stoichiometry matrix S_2, I. In the networks with $\delta = 1$, conditional plateau or power-law dynamics emerges depending on whether the final attractor is in the stoichiometric cone of the initial point.

Applicability of the structure-dynamics relationship for non-minimal networks

So far, we have explored the relationship between the network structure and emerging relaxation dynamics. There, we focused on the networks capable of synthesizing the final product from given substrates with the fewest possible reactions. In this section we show that the three network-structural features predict the relaxation dynamics well in non-minimal networks also, and even in a real biological network.

Combining minimum networks. Here we combine two minimum networks which share the same substrates and the final products. Fig. 5 is one instance of such a the combination. The two networks, labelled G and H, are minimum networks that synthesize the same product from the same substrate. Both networks have a non-zero rank gap δ, and in addition, the dimension of the left-null space I is zero for network G but 1 for network H. The combined network is obtained as the union of the sets of chemical species and reactions of the two networks and, thus, has redundant reactions (Fig. 5A).

Interestingly, the combined network $G \cup H$ has zero rank gap ($\delta=0$) and no conserved quantity ($I = 0$). As a consequence, the relaxation dynamics of the combined network becomes a simple, exponential-type even though the original networks exhibit the power-law and the plateau relaxation, respectively (Fig. 5B-D).

We can also see that the effective structure of the network varies over time when we combine the two networks with different rate-constants. We construct a combined network which has rate constant unity for all reactions from network H, and rate constant $1/\tau \ll 1$ for those reactions only in G. This combined network has an effective structure with non-zero δ and I until the dynamics start to "feel" the additional part brought from G ($t \lesssim \tau$). Thus, the relaxation type becomes the power-law or the plateau type for $t \lesssim \tau$. Subsequently, once the dynamics feel the whole structure of the combined network, both δ and I are zero, and thus, the relaxation dynamics transition to the exponential-type and quickly relax to the steady-state (Fig. 6).

Fig. 5. A. The two networks G and H, sharing some chemical species, including the substrate and the product, are combined to form the union network $G \cup H$. Each network G and H has a non-zero rank gap δ, and network H has one conserved quantity. However, the combined network has zero rank gap and no conserved quantity. B and C. The time courses of the network G (B) and H (C) exhibit the power-law or the conditional-plateau type relaxation due to the non-zero rank gap. D. Since δ and I are both zero in the combined network $G \cup H$, the relaxation time courses is of the exponential type.

Large networks. Next, we ask if the three quantities, the rank gap, the left null-spaces, and the stoichiometric cone, also work as indicators of the relaxation dynamics of large networks. For this purpose, we generated a number of reaction networks consisting of the given numbers of chemicals and reactions utilizing Mixed Integer Linear Programming (for details, see Methods). In the following, we deal with $|\mathscr{C}_0| = |\mathscr{R}_0| = 64$ networks constructed from the fully-connected network with $M = 2$ and $L = 8$ model. The number of the input chemical N_s as 2 and the output chemical N_p as 1.

We classified the networks based on the rank gap (δ) and the dimension of the left null space (I) of the stoichiometric matrix. Examples of the dynamics of type-I and type-II networks are shown in the SI Text. In line with the analysis of the minimum networks, most of the type-I and type-II networks exhibited the exponential-, and plateau-type relaxation, respectively, regardless of the initial concentrations. However, when the concentrations of the chemicals at the attractor span a wide range, several reactions having chemicals with low concentrations are considerably slowed down, and the relaxation time course cannot be described simply by the words "exponential" or "plateau". The critical values of the concentration range are studied in the SI text section.

The relaxation dynamics get much more complicated for the networks with a non-zero rank gap $\delta > 0$. Three example time courses are shown in Fig. 7. It is reasonable to term the relaxation time courses plotted in the top and the middle panel as "power-law" and "plateau", while the one shown in the bottom panel is neither plateau nor power-law type. Certainly, the relaxation dynamics of such large networks with $\delta > 0$ can no longer be simply classified as "plateau" or "power-law". However, similar to the above analysis for minimum networks, we examine the relationship between the migration length and the confinement of the trajectory in the stoichiometric cone.

Fig. 8 is the scatter plot showing the relationship between $\langle L/L_{\max} \rangle_{\text{in}}$ and $\langle L/L_{\max} \rangle_{\text{out}}$ of each large network we constructed. In contrast to the minimum network case, the relationship $\langle L/L_{\max} \rangle_{\text{in}} > \langle L/L_{\max} \rangle_{\text{out}}$ is violated by a few large networks, but the broad trend remains the same – the trajectories confined in the stoichiometric cone tend to show lower values of normalized migration length.

A biological example

Lastly, we demonstrate that the features of the network structure play a role also in a real biological network, namely, the metabolic network model of glucose fermentation in *Lactococ-*

Fig. 6. Three time courses exemplifying the dynamics of the combined network $G \cup H$ with different rate-constants of the reactions (unity for reactions in H and $1/\tau$ for that only in G). The gray-dashed vertical lines indicates $t = \tau = 10^4$. The dynamics for $t \lesssim \tau$ shows the plateau- or the power-law relaxation behaviour, while after that, the state exponentially converges to the steady-state attractor, except the left bottom case.

Fig. 7. Three example time courses from three different large networks. The dynamics shown in the top and middle panels can be termed as the power-law and the plateau relaxation, respectively, while that in the bottom panel is more complicated.

cus lactis (Fig.9A). The model is adopted from the reference (30). The network takes up glucose (GLC) from the environment and ferments it. The final product of the fermentation — either lactate (LAC), ethanol (ETOH), acetate (AC), acetoin (ACET), or butanediol (BUT) — is released into the environment. We briefly note some details of the model: the reaction rates are given by the Michaelis-Menten type functions (31), and the spontaneous degradation term $(-\phi x)$ is not implemented in the original model. The reactions depicted in the network diagram are all the reactions in the model (for a full description of the model, see the original paper).

A significant feature of the model is the balance between NAD and NADH. Suppose that a single glucose molecule is converted into either lactose or ethanol, then the production and consumption of NAD and NADH balance. However, when a glucose molecule is converted to acetate and secreted, a single NAD molecule is consumed, and a single molecule of NADH is produced. Thus, if the pathway GLC → AC is utilized, the concentrations of NAD and NADH cannot balance without the NADH oxidase (NOX) reaction (NADH → NAD) drawn separately from the main network in Fig.9A. Indeed, the stoichiometric matrix of the fermentation model will have the rank gap $\delta = 1$ when NOX is removed from the model, while with this reaction included it has a zero rank gap.

Here, we expect to observe a qualitative difference in the relaxation dynamics that originate from the rank gap by comparing the relaxation dynamics of the model with- and without

Fig. 8. The normalized migration length is averaged separately for the time courses with the initial points whose SC contains (vertical axis) and does not contain (horizontal axis) the final attractor. The $y = x$ line is plotted as a guide for the eye. The distributions of $\langle L/L_{\max} \rangle_{\text{in}}$ and $\langle L/L_{\max} \rangle_{\text{out}}$ are shown along the horizontal and the vertical axis, respectively.

the NOX reaction. However, the model without the NOX reaction can never reach a steady-state attractor. Recall that the relaxation to the attractor in the polymer models is possible because of the spontaneous degradation. We cannot introduce it into this model because the total concentrations of the following chemical pairs are set to be constant: ATP and ADP, NAD and NADH, and Coenzyme-A (CoA, which does not appear in the network diagram) and Acetyl-CoA (AcCoA). Instead, we changed the rate constant of the NOX reaction, 100 times faster and slower than the value used in (30), which results in the effective structure of the network in a certain time range to be with- and without the NOX reaction. Here, the slow down of the NOX reaction corresponds to the decrease of oxigen in the medium. Also, we set the external acetate concentration to be 10^{-4} (mM), leaving the rest of the parameter values unchanged.

Figs. 9B and C show the examples of the relaxation time course with the fast (B) and slow (C) rate constant of the NOX reaction. As seen from the figures, the relaxations are exponential-like in the model with a fast NOX reaction, and are power-law or plateau-like in the model with the slow NOX reaction.

Discussion

In the present manuscript, we have studied a class of chemical reaction networks consisting of polymer chemicals that can split and ligate with each other. The networks satisfy the law of mass conservation and can generate a variety of mass-conserving sub-networks. The minimum networks capable of synthesizing a specified set of product precursors from a set of input substrates are computed within a reasonable time by utilizing the algorithm for solving the SAT problem.

We generated a large number of such minimum networks and simulated their chemical reaction dynamics to unveil the relationship between the network structure and the relaxation dynamics after perturbation of the steady-state. We found three types of relaxation dynamics: exponential, power-law and plateau type. We found that we could predict which relaxation behaviour will be exhibited from three structural features of the reaction networks, namely, (i) the rank gap, (ii) the dimension of the left null space, and (iii) the stoichiometric cone. We showed that these three features determine the relaxation behaviour not only in the minimum networks, but also in combinations of the minimum networks, in large networks with many redundant pathways, as well as in one example of a real metabolic network.

Our results therefore provide easily computable criteria for predicting whether a given chemical reaction network can lead to slow dynamics. By examining the rank gap and the left null-space, one can predict whether the concentration of the chemicals will quickly relax to a steady-state or not. A non-zero rank gap indicates that the reaction network cannot have a steady state. Thus, especially, we expect metabolic reaction networks of biological cells would have evolved to achieve a zero rank gap. However, as illustrated by the network combination example and the metabolic networks of *Lactococcus lactis*, the behaviours of the chemical reaction network reflect the effective network structure on a timescale of interest. Therefore, over certain timescales metabolic, and other real chemical networks, may effectively behave as if they have a non-zero rank gap, leading to non-exponential relaxation behaviours.

Finally, we would like to emphasise the flexibility of our theoretical framework. Even with only polymer chemical species, the size is easily controlled by manipulating the maximum length L and types of the monomers M. Further, the types of species and chemical reactions used to construct the networks can easily be expanded without affecting key chemical properties like mass conservation. For instance, one could include all species and reactions known to occur in microbes (32). The SAT formulation can be equally easily used in such a case also to construct minimum networks. We expect that our approach for constructing and analyzing such artificial reaction networks will provide a useful toolbox for exploring the extensive world of out-of-steady-state dynamics of chemical reaction networks.

Method

Generating artificial reaction networks. For the set of chemicals $\mathscr{C}$, we wish to identify the smallest set of reactions $\mathscr{R}_0 \subseteq \mathscr{R}$ such that growth factors can be produced from the available nutrients.

As input we have the set of available nutrients $N \subset \mathscr{C}$ and the set of growth factors $G \subset \mathscr{C}$ that must be produced. The main idea of our approach is to consider the Boolean expression

$$B = (|\mathscr{R}_0| \leq K) \wedge \bigwedge_{g \in G} \mathrm{CanMake}(g),$$

where $\mathrm{CanMake}(g)$ depends on the nutrient set N. We may then iteratively solve this SAT problem starting from $K = 1$ (to $|\mathscr{R}|$) and stop when we can satisfy the expression. This ensures that we find the global minimum solutions.

The Boolean *at most* expression, $|\mathscr{R}_0| \leq K$, can be expressed efficiently using auxiliary variables (33). What remains is the specification of the Boolean function CanMake. We can define this directly by the binary tree of reactions that leads from the nutrients to the growth factors. For instance if the required growth factor is AA and available nutrients are AB and AC, we could write

$$\mathrm{CanMake}(AA) = (R_{AB \rightleftharpoons A,B} \wedge R_{AA \rightleftharpoons A,A})$$
$$\vee (R_{AC \rightleftharpoons A,C} \wedge R_{AA \rightleftharpoons A,A}).$$

This fully defines a SAT-problem in the reaction R-variables. However, this approach makes exponentially long expressions for large sets of chemicals and reactions.

The reason for the long expressions is mainly repetitions: even in the tiny example above, the reaction variable $R_{AA \rightleftharpoons A,A}$ appeared in both sub-expressions. To remedy this, we use the concept of equisatisfiability to introduce auxiliary variables that describe if a specific chemical $c \in \mathscr{C}$ can be produced. Using these variables we can truncate the binary tree after one level. We thus intend to write the condition to make growth factors as

$$\bigwedge_{c \in G} c \wedge \bigwedge_{c \in \mathscr{C}} \mathrm{Relation}(c),$$

where the Boolean Relation function specifies the relationship between chemicals as specified by the set of reactions. For instance, for the above example:

$$\mathrm{Relation}(AA) = AA \odot \left[A \wedge R_{AA \rightleftharpoons A,A} \right], \qquad [10]$$

i.e. to make AA you need available A and the reaction $AA \rightleftharpoons A, A$. Here, $\odot$ denotes *exclusive nor* indicating for $X \odot Y$ that

Fig. 9. (A) The metabolic model of glucose fermentation in *Lactococcus lactis* adopted from (30). The red colored metabolites and the blue colored metabolites are the intracellular- and extracellular metabolites, respectively. (B and C) The representative time courses of the model with the fast NOX reaction (B) and the slow NOX reaction (C). The time courses shown in the neighboring panels start from the same initial concentrations.

either both X and Y are TRUE or both are FALSE. Likewise, for example

$$\text{Relation}(A) = A \odot \big[(AA \wedge R_{AA \rightleftharpoons A,A}) \vee (AB \wedge R_{AB \rightleftharpoons A,B})$$
$$\vee (AC \wedge R_{AC \rightleftharpoons A,C}) \big]$$
$$= A \odot \big[(AA \wedge R_{AA \rightleftharpoons A,A}) \vee R_{AB \rightleftharpoons A,B} \vee R_{AC \rightleftharpoons A,C} \big],$$

where we used the fact that AB and AC are available nutrients.

With this approach, however, we need to restrict the direction of the reactions: reactions should only be used in one direction as the network of reactions from nutrients to growth factors should be *directed* and *acyclic* (a DAG). To fix the directionality, we introduce Boolean direction variables $\{D_i\}$ with the convention that $D =$ TRUE indicates breakdown reactions and $D =$ FALSE indicates assembly. If we denote the reaction $AA \rightleftharpoons A, A$ by k, Eq. (10) thus becomes $\text{DRelation}(AA) = AA \odot [A \wedge R_k \wedge \neg D_k]$, whereas the DRelation for A would use D_k without the negation. When there is no reaction, we do not care about the direction of the (nonexistent) reaction and we thus include a term that enforce $D_k =$ TRUE if there is no reaction, i.e. $\neg R_k \Rightarrow D_k = R_k \vee D_k$.

The final reformulated expression is

$$B = (|\mathscr{R}_0| \leq K) \wedge \bigwedge_{c \in G} c \wedge \bigwedge_{c \in \mathscr{C}} \text{DRelation}(c) \wedge \bigwedge_{i \in \mathscr{R}} (R_i \vee D_i).$$

The only remaining complexity is the requirement that the solution of reactions is acyclic. We implement this requirement outside the language of Boolean expressions and simply discard such solutions. If only cyclic solutions are found at a certain K, these solutions are disallowed at larger K, thus avoiding a large number of degenerate cyclic solutions at large K.

All solutions to the SAT-problem are found using the GLUCOSAMINE solver via the BOOLEXPR interface.

Checking if the attractor is in the stoichiometric cone. For checking if the attractor is in the stoichiometric cone of the initial point, $SC(\boldsymbol{x}_{\text{ini}})$, we computed the L^∞ norm between the

attractor and the points in the stoichiometric cone and minimize the norm, by utilizing the following linear optimization problem.

$$\text{minimize} \quad \max_i \Big| x_{\text{att}}^{(i)} - (x_{\text{att}}^{(i)} + \sum_j (S_2)_{ij} v_2^{(j)}) \Big|, (0 \leq i < |\mathscr{C}_0|)$$

$$\text{subject to} \quad x_{\text{att}}^{(i)} + \sum_j (S_2)_{ij} v_2^{(j)} > 0, (0 \leq i < |\mathscr{C}_0|)$$

$$J^{(\boldsymbol{pr})} > 0$$

If the minimum distance is smaller than 10^{-12}, we regard that the attractorr is in the stoichiometric cone and *vice versa*.

Finding $\tilde{\boldsymbol{x}}_{\text{ini}}$. For obtaining $\tilde{\boldsymbol{x}}_{\text{ini}}$, the nearest point to the initial point with the same value of the conserved quantities $\boldsymbol{Q}$ with that of the attractor, we utilized the following quadratic optimization problem with $\boldsymbol{\delta x}$ as variables

$$\text{minimize} \quad \|\boldsymbol{\delta x}\|^2$$
$$\text{subject to} \quad \boldsymbol{Q}(\boldsymbol{x}_{\text{ini}} + \boldsymbol{\delta x}) = \boldsymbol{Q}(\boldsymbol{x}_{\text{att}})$$
$$\boldsymbol{x} \geq \boldsymbol{0}.$$

and obtained $\tilde{\boldsymbol{x}}_{\text{ini}}$ as $\tilde{\boldsymbol{x}}_{\text{ini}} = \boldsymbol{x}_{\text{ini}} + \boldsymbol{\delta x}$.

Generating networks with a specified size. Due to the computational cost, we adopt a mixed-integer linear (MIL) optimization approach for generating the large (non-minimal) networks, instead of the SAT approach. For obtain a network with $|\mathscr{C}_0|$ chemicals, $|\mathscr{R}_0|$ reactions, N_s substrates, and N_p

product precursors by solving the following MIL problem:

$$\text{minimize} \quad \mathbf{1}_{|\mathscr{C}|} \cdot \boldsymbol{\xi}$$

$$\text{subject to} \quad \tilde{S}(\boldsymbol{v}_s^{(+)} - \boldsymbol{v}_s^{(-)}) + U_{\max}\boldsymbol{n}_s + \boldsymbol{\xi} \geq \boldsymbol{g},$$

$$0 \leq \boldsymbol{v}_s^{(\pm)} \leq M\boldsymbol{\sigma},$$

$$2 - \sum_s \boldsymbol{n}_s - \boldsymbol{g} - \tilde{T}\boldsymbol{\sigma} \leq M(\mathbf{1}_{|\mathscr{C}|} - \boldsymbol{\theta})$$

$$3\boldsymbol{\sigma} \leq |\tilde{S}|\boldsymbol{\theta} \quad\quad [11]$$

$$\mathbf{1}_{|\mathscr{C}|} \cdot \boldsymbol{n}_s = 1$$

$$\mathbf{1}_{|\mathscr{C}|} \cdot \boldsymbol{n}_p = N_p$$

$$\mathbf{1}_{|\mathscr{C}|} \cdot \boldsymbol{\theta} = |\mathscr{C}_0|$$

$$\mathbf{1}_{|\mathscr{R}|} \cdot \boldsymbol{\sigma} = |\mathscr{R}_0|$$

where the continuous variables are $\boldsymbol{\xi} \geq 0$ and $\boldsymbol{v}_s^{(\pm)} \geq 0$, and the bool variables are $\boldsymbol{n}_s, \boldsymbol{n}_p, \boldsymbol{\theta}, \boldsymbol{\sigma}$. $\boldsymbol{\xi}$ and $\boldsymbol{v}_s^{(\pm)}$ are the slack variable and the forward- and backward reaction flux vectors under the environment s, respectively. $\boldsymbol{n}_s$ and $\boldsymbol{n}_p$ denotes the choice of the substrate and the product. If the ith element of the vector $\boldsymbol{n}_s$ ($\boldsymbol{n}_p$) is one, the ith chemical is the substrate supplied under the environment s (the product). $\boldsymbol{\theta}$ and $\boldsymbol{\sigma}$ denotes which chemicals ($\boldsymbol{\theta}$) and reactions ($\boldsymbol{\sigma}$) are used. If the entity is unity, it is used.

$\tilde{S}$ and $U_{\max}$ are the stoichiometric matrix of all reactions in $\mathscr{R}$, and the maximum substrate uptake rate. $\tilde{T}$ and $|\tilde{S}|$ are defined as $\tilde{T}_{ij} = |\text{sgn}(\tilde{S}_{ij})|$ and $|\tilde{S}|_{ij} = |\tilde{S}_{ij}|$. $\mathbf{1}_n$ is the abbreviated representation of n-dimensional vector with all the elements being unity.

Each constraint of Eq. (11) represents the following conditions. (First line) The total production rate of the ith chemical should be larger than p_i ($= 1$ if it is one of the products and $= 0$ otherwise). (Second line) The rth reaction flux can be turned-on if σ_r is one. (Third line) If θ_i can be unity only if two or more (one or more for the substrates and the products) reactions connected to the ith chemical are turned-on. (Forth line) To turn the r th chemical on, the substrates and the product of the reaction should be in the network. The fifth to the eighth line set the number of the substrates, products, total chemicals, and total reactions.

Eq. (11) is always feasible and if the optimal value of the objective is smaller than 10^{-12}, the original network $\{\mathscr{C}, \mathscr{R}\}$ has a subnetwork of the specified size. For generating multiple networks, we introduce an additional constraint

$$\sigma_i = 0, \ i \in K_{\text{off}}$$

where the index set K_{off} consists of randomly-chosen indices.

Acknowledgments

The authors thank Chikara Furusawa for fruitful discussion. This work is supported by the research grant (00028054) from VILLUM FONDEN and JSPS KAKENHI Grant Number JP 21K20626 and JP 22K15069. J.B.K. acknowledges funding from the Novo Nordisk Foundation, Grant Agreement NNF20OC0062047. N.M. acknowledges funding from Novo Nordisk foundation, Grant Agreement NNF21OC0068775. S.K. acknowledges funding from the Department of Atomic Energy, Government of India (Project Identification No. RTI 4006) and the Simons Foundation (Grant No. 287975).

1. P Érdi, J Tóth, *Mathematical models of chemical reactions: theory and applications of deterministic and stochastic models*. (Manchester University Press), (1989).
2. HB Callen, *Thermodynamics and an Introduction to Thermostatics*. (American Association of Physics Teachers), (1998).
3. KJ Laidler, J Keith, , et al., *Chemical kinetics*. (McGraw-Hill New York) Vol. 2, (1965).
4. NG Van Kampen, *Stochastic processes in physics and chemistry*. (Elsevier) Vol. 1, (1992).
5. A Varma, BO Palsson, Metabolic capabilities of escherichia coli: I. synthesis of biosynthetic precursors and cofactors. *J. theoretical biology* **165**, 477–502 (1993).
6. A Varma, BO Palsson, Metabolic capabilities of escherichia coli ii. optimal growth patterns. *J. Theor. Biol.* **165**, 503–522 (1993).
7. JS Edwards, BO Palsson, Robustness analysis of the escherichiacoli metabolic network. *Biotechnol. progress* **16**, 927–939 (2000).
8. JS Edwards, RU Ibarra, BO Palsson, In silico predictions of escherichia coli metabolic capabilities are consistent with experimental data. *Nat. biotechnology* **19**, 125–130 (2001).
9. MW Covert, CH Schilling, B Palsson, Regulation of gene expression in flux balance models of metabolism. *J. theoretical biology* **213**, 73–88 (2001).
10. DW Erickson, et al., A global resource allocation strategy governs growth transition kinetics of escherichia coli. *Nature* **551**, 119–123 (2017).
11. M Scott, CW Gunderson, EM Mateescu, Z Zhang, T Hwa, Interdependence of cell growth and gene expression: origins and consequences. *Science* **330**, 1099–1102 (2010).
12. S Klumpp, Z Zhang, T Hwa, Growth rate-dependent global effects on gene expression in bacteria. *Cell* **139**, 1366–1375 (2009).
13. Y Himeoka, K Kaneko, Theory for transitions between exponential and stationary phases: universal laws for lag time. *Phys. Rev. X* **7**, 021049 (2017).
14. Y Himeoka, B Gummesson, MA Sørensen, SL Svenningsen, N Mitarai, Distinct survival, growth lag, and rrna degradation kinetics during long-term starvation for carbon or phosphate. *mSphere* pp. e01006–21 (2022).
15. O Gefen, O Fridman, I Ronin, NQ Balaban, Direct observation of single stationary-phase bacteria reveals a surprisingly long period of constant protein production activity. *Proc. Natl. Acad. Sci.* **111**, 556–561 (2014).
16. Y Kaplan, et al., Observation of universal ageing dynamics in antibiotic persistence. *Nature* **600**, 290–294 (2021).
17. I Levin-Reisman, et al., Automated imaging with scanlag reveals previously undetectable bacterial growth phenotypes. *Nat. Methods* **7**, 737 (2010).
18. E Biselli, SJ Schink, U Gerland, Slower growth of escherichia coli leads to longer survival in carbon starvation due to a decrease in the maintenance rate. *Mol. Syst. Biol.* **16**, e9478 (2020).
19. K Kaneko, Recursiveness, switching, and fluctuations in a replicating catalytic network. *Phys. Rev. E* **68**, 031909 (2003).
20. D Segré, D Ben-Eli, D Lancet, Compositional genomes: prebiotic information transfer in mutually catalytic noncovalent assemblies. *Proc. Natl. Acad. Sci.* **97**, 4112–4117 (2000).
21. H Jeong, B Tombor, R Albert, ZN Oltvai, AL Barabási, The large-scale organization of metabolic networks. *Nature* **407**, 651–654 (2000).
22. E Ravasz, AL Barabási, Hierarchical organization in complex networks. *Phys. review E* **67**, 026112 (2003).
23. P Schwaller, et al., Molecular transformer: a model for uncertainty-calibrated chemical reaction prediction. *ACS central science* **5**, 1572–1583 (2019).
24. J Nam, J Kim, Linking the neural machine translation and the prediction of organic chemistry reactions. *arXiv preprint arXiv:1612.09529* (2016).
25. M Segler, M Preuß, MP Waller, Towards" alphachem": Chemical synthesis planning with tree search and deep neural network policies. *arXiv preprint arXiv:1702.00020* (2017).
26. J Zaldivar, J Nielsen, L Olsson, Fuel ethanol production from lignocellulose: a challenge for metabolic engineering and process integration. *Appl. microbiology biotechnology* **56**, 17–34 (2001).
27. B Hahn-Hägerdal, M Galbe, MF Gorwa-Grauslund, G Lidén, G Zacchi, Bio-ethanol–the fuel of tomorrow from the residues of today. *Trends biotechnology* **24**, 549–556 (2006).
28. C Moore, S Mertens, *The nature of computation*. (OUP Oxford), (2011).
29. M Feinberg, *Foundations of chemical reaction network theory*. (Springer), (2019).
30. MH Hoefnagel, et al., Metabolic engineering of lactic acid bacteria, the combined approach: kinetic modelling, metabolic control and experimental analysisthe genbank accession number for the sequence reported in this paper is ay046926. *Microbiology* **148**, 1003–1013 (2002).
31. A Cornish-Bowden, *Fundamentals of enzyme kinetics*. (John Wiley & Sons), (2013).
32. M Kanehisa, M Furumichi, Y Sato, M Ishiguro-Watanabe, M Tanabe, Kegg: integrating viruses and cellular organisms. *Nucleic acids research* **49**, D545–D551 (2021).
33. C Sinz, *International conference on principles and practice of constraint programming*. (Springer), pp. 827–831 (2005).