Chimeric 3D-gastruloids – a versatile tool for studies of mammalian peri-gastrulation development

Alexandra E. Wehmeyer[1], Katrin M. Schüle[1], Alexandra Conrad[1], Chiara M. Schröder[1,2,3,4], Simone Probst[1], and Sebastian J. Arnold[1,4]

[1]Institute of Experimental and Clinical Pharmacology and Toxicology, Faculty of Medicine, University of Freiburg, Albertstrasse 25, D-79104 Freiburg, Germany

[2]Spemann Graduate School of Biology and Medicine (SGBM), University of Freiburg, Albertstrasse 19a, D-79104 Freiburg, Germany

[3]Faculty of Biology, University of Freiburg, Schänzlestrasse 1, D-79104 Freiburg, Germany

[4]Signaling Research Centers BIOSS and CIBSS, University of Freiburg, Schänzlestrasse18, D-79104 Freiburg, Germany

For Correspondence: sebastian.arnold@pharmakol.uni-freiburg.de

ORCID Alexandra E. Wehmeyer 0000-0002-9981-8042
ORCID Katrin M. Schüle 0000-0001-6642-0030
ORCID Alexandra Conrad 0000-0001-7766-5033
ORCID Chiara M. Schröder 0000-0001-5345-3415
ORCID Simone Probst 0000-0002-0220-5400
ORCID Sebastian J. Arnold 0000-0002-2688-9210

Key words

Chimeric *3D*-Gastruloids, mESCs, *Eomes*, *Brachyury*, functional genetics, axis formation, cell lineage specification, Tbx transcription factors

Abstract

Stem cell-derived 3D-gastruloids show a remarkable capacity of self-organisation and recapitulate many aspects of gastrulation stage mammalian development. Gastruloids can be rapidly generated and offer several experimental advantages, such as scalability, observability, and accessibility for manipulation. Here, we present approaches to further expand the experimental potency of murine 3D-gastruloids by utilizing functional genetics in mouse embryonic stem cells (mESCs) to generate chimeric gastruloids. In chimeric gastruloids fluorescently labelled cells of different genotypes harbouring inducible gene-expression, or loss-of-function alleles, are combined with wildtype cells. We showcase this experimental approach in chimeric gastruloids of mESCs carrying homozygous deletions of the Tbx transcription factors *Brachyury,* or inducible expression of *Eomes.* Resulting chimeric gastruloids recapitulate reported *Eomes* and *Brachyury* functions, such as instructing cardiac fate and promoting posterior axial extension, respectively. Additionally, chimeric gastruloids revealed previously unrecognized phenotypes such as tissue sorting preference of *Brachyury*-deficient cells to endoderm, and cell non-autonomous effects of *Brachyury*-deficiency on *Wnt3a*-patterning along the embryonic axis, demonstrating some of the advantages of chimeric gastruloids as efficient tool for studies of mammalian gastrulation.

Introduction

The versatile experimental opportunities offered by functional genetics available in the mouse as main mammalian model system have greatly enhanced our understanding of embryonic development. However, studies of mammalian embryogenesis are hampered by the relative inaccessibility of the embryo due to intrauterine growth. *Ex vivo* embryo cultures partially overcome some of the experimental restrictions but rely on the isolation of often limiting numbers of embryos. More recently, novel approaches were developed to generate stem cell derived, 3D-embryoids reflecting different stages of mammalian development from the formation of the blastocyst (Kagawa et al., 2022; Li et al., 2019; Yu et al., 2021), to periimplantation embryos (Amadei et al., 2021; Harrison et al., 2017; Harrison et al., 2018), and postimplantation stages recapitulating early organogenesis (Beccari et al., 2018; Moris et al., 2020; Turner et al., 2017; van den Brink et al., 2020; van den Brink et al., 2014; Veenvliet et al., 2020). These embryoid models demonstrate the remarkable self-organisation and robustness of

embryonic programs that guide the processes of morphogenesis, growth, cell lineage
specification and differentiation in the embryo as well as by *in vitro* models.

Among widely used embryoid model system are 3D-gastruloids, offering a simple
experimental procedure to reproducibly generate embryoids that recapitulate
development stages from early gastrulation to somitogenesis and onset of
organogenesis (comparable to Embryonic days 6.5 (E6.5) – E9.0) reviewed in (van
den Brink and van Oudenaarden, 2021; Veenvliet et al., 2021). 3D-gastruloids are
formed by the aggregation of 100-300 mouse or human pluripotent embryonic stem
cells (m or hESCs) that are treated by a 24 h pulse of the GSK3b-inhibitor CHIR to
activate the canonical Wnt-cascade. This signalling stimulus induces an anterior-
posterior asymmetry in the aggregate (van den Brink et al., 2014), indicated by the
one-sided expression of the Tbx factor *Brachyury* that in the embryo marks the site of
primitive streak formation at gastrulation onset (Rivera-Perez and Magnuson, 2005;
reviewed in Arnold and Robertson, 2009). The *Brachyury*-expressing posterior pole of
3D-gastruloids elongates over several days thereby forming different tissues
resembling paraxial mesoderm, neural tube, and the primitive gut tube. 3D-gastruloids
show a remarkable self-organizing capacity as the different tissues are generated in
proper spatial organization and arranged according to the embryonic axes (Beccari et
al., 2018). 3D-gastruloids thus can be used as model systems for studies of various
gastrulation-associated morphogenetic processes, as exemplified for somitogenesis
(van den Brink et al., 2020; Veenvliet et al., 2020). However, some tissues of
gastrulation stage embryos are less represented in 3D-organoids, including anterior
mesoderm derivatives, such as cardiogenic mesoderm, and cranial structures such as
cranial neural tissues (van den Brink et al., 2020; Veenvliet et al., 2020). This
underrepresentation of anterior/cranial tissues most likely results from the initial
induction of ESC aggregates with CHIR which imposes a strong signal for tissue
identities of the posterior/caudal primitive streak (Dunty et al., 2008). However, this
and other aberrations of 3D-gastruloids compared to embryos can also be used to
investigate which additional regulatory requirements need to be met for proper
morphogenesis of tissues and organ *anlagen* to more closely recapitulate the embryo
(Veenvliet et al., 2021). Due to the scalability of this ESC- derived embryoid culture
system, multiple different environmental cues can be readily tested. It is thus expected
that further refinements of current protocols will lead to the generation of gastruloids

that more closely resemble the full spectrum of gastrulation and following stages of
embryogenesis.

Cell specification to mesoderm and definitive endoderm (DE) cell lineages is regulated
by two Tbx transcription factors *Eomesodermin* (*Eomes*) and *Brachyury (T)*. Functions
of both factors were previously extensively studied in mouse establishing crucial roles
of *Eomes* for cell lineage specification of definitive endoderm (DE) (Arnold et al., 2008;
Teo et al., 2011) and anterior mesoderm (Costello et al., 2011), and functions of
Brachyury for the generation of posterior mesoderm derivatives, and notochord
(Wymeersch et al., 2021). The compound genetic deletion of *Eomes* and *Brachyury*
completely abrogates specification of any ME during differentiation of pluripotent cells
(Tosic et al., 2019). While functions of these Tbx factors for specification of cell fate
are well described, their roles in tissue-wide morphogenetic processes are less
defined. For example, the genetic deletion of *Eomes* in the epiblast abrogates
formation of the mesoderm (and DE) cell layer, hindering studies of morphogenetic
functions of *Eomes*-regulated programs (Arnold et al., 2008). Similarly, studies of the
cell-intrinsic roles of *Brachyury* in posterior body axis extension are compromised by
cell non-autonomous functions of *Brachyury* in feed-forward regulatory loops to
maintain caudal Wnt-signals, that could independently act on cell specification and
morphogenetic programs (Arnold et al., 2000; Dunty et al., 2008; Martin and Kimelman,
2008; Turner et al., 2014; Yamaguchi et al., 1999). These constraints to distinguish
cell-autonomous effects of loss-of-gene-function from secondary effects, such as by
general disturbance of morphogenesis, can be experimentally addressed by the
generation of embryonic chimeras. Here, cell-autonomous gene functions can be
studied when limited numbers of genetically altered cells are placed in an otherwise
wildtype cellular environment. Chimeric approaches were previously reported for
studies of *Brachyury* functions, such as in zebrafish (Halpern et al., 1993) and mouse
(Wilson et al., 1993; Wilson et al., 1995; Wilson and Beddington, 1997), as well as for
Eomes functions (Arnold et al., 2008), indicating regulatory roles of both Tbx factors in
cell migration and morphogenesis. However, embryonic chimera analyses are highly
labour and resource intensive, especially in mouse, and thus are only infrequently
employed.

In this report, we demonstrate an approach to further increase the experimental
potency of murine 3D-gastruloids by expanding it to chimeric analyses. We use
genetically engineered, traceable mESCs to generate chimeric 3D-gastruloids that are

composed of cells with different genotypes. To showcase the advantages of this
approach and to test experimental feasibility we generated fluorescently labelled
mESC lines with homozygous loss-of-function, or inducible expression cassettes for
the two Tbx transcription factors *Brachyury* and *Eomes*. These are used to generate
chimeric gastruloids, e.g. by mixing defined ratios of inducible Tbx-expressing, or Tbx-
deficient and WT cells to generate chimeric situations with defined cell contribution that
are normally only difficult to achieve by conventional genetic tools or by cell injections
into embryos. In addition to demonstrating the efficiency and feasibility of this
experimental approach of chimeric gastruloids, this study also provides some
previously unrecognized insights into the morphogenetic functions of *Eomes* and
Brachyury.
145

Results and Discussion
To expand the experimental versatility of the 3D-gastruloid model system we combined
it with the use of genetically modified mESCs and fluorescent imaging. We generated
chimeric gastruloids composed of cells with different genetic backgrounds that can be
readily traced by different fluorescent membrane labels (Fig. 1A, B). WT mESCs are
permanently labelled by monoallelic knock-in of membrane GFP (mG) into the Rosa26
genomic locus (Fig. 1C), while membrane Tomato (mT) labelling was used for
genetically manipulated mESCs lines (Fig. 1C). Genetic alterations used in presented
experiments were the homozygous deletion of the Tbx transcription factor *Brachyury*
(*Bra$^{-/-}$*), and the ICE-mediated (induced cassette exchange) targeted integration of
Eomes 3' to a monoallelic Tet-responsive element (TRE) in A2lox (E14-based)
mESCs (Iacovino et al., 2014), resulting in TRE.*Eomes*GFP (short TRE.*Eo*). These
cell lines were first tested for their abilities to form regular gastruloids. This
demonstrated that fluorescent membrane labels don't have an impact on gastruloid
formation, while *Bra$^{-/-}$* and dox-induced TRE.*Eo* cells show a failure to form regularly
extended gastruloids at 120 h (Supplementary Fig. 1). To generate chimeric
gastruloids we followed two approaches (Fig. 1A, D, E), either by mixing of different
cells during the aggregation of mESCs at the beginning of gastruloid culture (Fig. 1E),
or by merging of two preformed mESCs aggregates composed of different cells 24 h
after the initial aggregation (Fig. 1 B, D). Such merged ESC aggregates rapidly fuse
and stably adhere after placing them together in 96-wells (Fig. 1D, Supplementary
Movie 1) or in hanging drops (not shown).

The mixing of cells at different ratios can be used to evaluate cell-autonomous vs. cell

non-autonomous gene functions, by testing how different levels of cell contribution

affects tissue behaviour, e.g. when using cells with loss-of-gene-function. Merging of

cell aggregates can be applied when the behaviour of coherent groups of cells is

analysed, such as in studies of inductive tissue interactions. A similar approach was

recently reported where a small aggregate of 50 cells was treated with BMP4 to induce

organizer-activities which was used instead of CHIR-treatment for the induction of

gastruloid-formation (Xu et al., 2021). Thus, chimeric gastruloids offer multiple

experimental opportunities to study gastrulation development (Fig. 1A) as exemplarily

demonstrated in following experiments.

First, we tested if chimeric gastruloids are a suitable model to study instructive gene

functions during lineage specification. Hence, we generated chimeric gastruloids by

merging preformed aggregates of WT mESCs (mG-label) and mESCs containing the

doxycycline (Dox) inducible *Eomes*.GFP expression cassette (TRE.*Eo*; mT-label) (Fig.

2A, B). *Eome*s is crucially required for lineage specification towards definitive

endoderm (Arnold et al., 2008; Teo et al., 2011) and anterior mesoderm, including the

cardiac lineage (Costello et al., 2011; Probst et al., 2021). We tested if forced *Eomes*

expression in a group of cells (TRE.*Eo*) within a gastruloid would be sufficient to induce

a coherent heart-like domain, which is only inconsistently forming in gastruloids

composed of WT mESCs (Rossi et al., 2021; van den Brink and van Oudenaarden,

2021). Indeed, Dox-induced *EomesGFP*-expressing cells form a domain of beating

cardiomyocytes (Fig. 2C, Supplementary movie 2) in chimeric gastruloids in 33.7% ($\pm$

3.0 SD) of cases, whereas similar beating areas are only observed in 1.5% ($\pm$ 2.5 SD)

in uninduced chimeric gastruloids (Fig. 2D) or using only A2lox WT mESCs (not

shown). To correlate induced *EomesGFP* expression with the cardiogenic domain we

performed *in situ* hybridization analyses for early cardiac markers *Mlc2a* and *Nkx2.5*,

which are expressed in mT-marked TRE.*Eo* cells that are forming a coherent domain

on one side of gastruloids (Fig. 2E, F). To analyse the position of TRE.*Eo* cell-derived

domains along the AP axis of merged chimeric gastruloids, we performed

immunostainings against BRACHYURY and CDX2, indicating the posterior pole of

extending gastruloids (Fig. 2G, H). The domains of TRE.*Eo* cells are predominantly

found on one side along the AP-axis of the gastruloids and oriented towards the

anterior pole (Fig. 2G, H). Of note, these coherent domains of cells are not found in

gastruloids when two preformed aggregates of WT cells are merged where cells

undergo increased mixing (Supplementary Fig. 2A). Interestingly, TRE.*Eo* cell

containing gastruloids consistently showed a reduction in axis extension, so that the

anterior-posterior (AP) axis of resulting gastruloids is less obvious in comparison to

only WT-derived gastruloids (Supplementary Fig. 2B, C). In short, these experiments

show that induced expression of *Eomes* suffices to cell-autonomously generate

coherent cardiogenic domains oriented to the anterior-lateral side of developing

gastruloids in a highly increased proportion of gastruloids compared to uninduced

gastruloids.

Next, we aimed for testing the feasibility of chimeric gastruloids for the assessment of

cell-autonomous vs. cell non-autonomous functions of *Brachyury* during posterior

elongation, as the shortening of the posterior body axis is the most prominent

phenotype of *Brachyury*-mutant embryos (Fig. 3A)(Wymeersch et al., 2021).

Gastruloids entirely generated from mT-labelled *Bra$^{-/-}$* mESCs present with a

phenotype of impaired elongation, reminiscent of the failure of tailbud elongation

observed in *Brachyury*-mutant embryos (Fig. 3B, E; Supplementary Figs. 1, 3B) (Inman

and Downs, 2006). Increasing the proportional contribution of WT cells (mG) to the

gastruloid by intermixing WT with *Bra$^{-/-}$* mESCs during the initial aggregation of 300

mESCs leads to the gradual extension and increase in overall tissue mass of the

posterior portion of the mixed gastruloids (Fig. 3B). We analysed BRACHYURY

presence in the WT cells in the posterior regions of chimeric gastruloids with high

(80%) and equal (50%) contribution of *Bra$^{-/-}$* cells at 24 h time intervals. The posterior

region of chimeric gastruloids with a contribution of *Bra$^{-/-}$* cells of 80% show less

confined and reduced a-BRACHYURY staining (Fig. 3C) compared to chimeric

gastruloids where *Bra$^{-/-}$* cell contribution is 50%. Here, *Brachyury* expression is found

robustly but with decreased intensity in the posterior pole when compared to WT

gastruloids (Fig. 3D, and Supplementary Fig. 3A).

The absence of *Brachyury* expression also from WT cells in gastruloids with high

contribution of *Bra$^{-/-}$* cells suggests cell non-autonomous effects of *Bra$^{-/-}$* cells. These

are most likely explained by disturbances of the previously described feed-forward

regulation of *Wnt3a* and *Brachyury* in the tail-bud region (Arnold et al., 2000; Dunty et

al., 2008; Martin and Kimelman, 2008; Turner et al., 2014; Yamaguchi et al., 1999). To

test this, we performed *in situ* hybridization to analyse expression of *Wnt3a* and the

putative *Brachyury* target gene and Wnt co-ligand *Rspo3 (*Koch *et al*., 2017*)* in WT

gastruloids, in gastruloids entirely composed of *Bra$^{-/-}$* cells, and in mixed gastruloids

with 80% and 50% *Bra^-/-* cell contribution at 72, 96 and 120 h (Fig. 3E). This analysis

demonstrates *Brachyury*-requirements for the sustained expression of *Wnt3a*, which

is absent from *Bra^-/-* gastruloids at 96 h and greatly reduced and mislocalised in mixed

gastruloids at 120 h. Interestingly *Wnt3a* expression, which normally is confined to the

posterior pole, is found at intermediate levels of mixed gastruloids, most likely resulting

from unequal cell distribution of WT and *Bra^-/-* cells, which accumulate at the posterior

of mixed *Bra^-/-* : WT gastruloids (see also Fig. 4 and discussion below). Analysis of

Rspo3 expression recapitulates the lack of *Brachyury* expression in *Bra^-/-* gastruloids

and the prematurely exhausted *Brachyury* expression in *Bra^-/-* : WT mixed gastruloids

(Fig. 3C-E).

Next, in addition to the tissue-wide, partially cell non-autonomous phenotype of

Brachyury-deficiency in gastruloids with a high contribution of *Bra^-/-* cells, we analysed

the cell-autonomous effects of *Brachyury*-deficiency in conditions with low contribution

of mutant cells (Fig. 4). We generated chimeric gastruloids by mixing cells in a 10:90

ratio of *Bra^-/-* : WT mESCs (Fig. 4A, B) and compared resulting gastruloids with controls

where 10% mT-labelled WT cells were used (Fig. 4C, D). Further controls were

included where cells were mixed at 50:50 ratios (Supplementary Fig. 4A, B). WT cells

randomly disperse, while *Bra^-/-* exhibit a tissue sorting behaviour (Fig. 4A-D,

Supplementary Fig. 4A, B). At 120 h of culture *Bra^-/-* cells are predominantly found

along the midline in the interior of the gastruloids, and in the posterior pole (Fig. 4B).

To determine tissue identity of these regions we used immunofluorescence (IF)

staining against FOXA2 and CDH1 (E-Cadherin), labelling DE (Viotti et al., 2014),

SOX2 labelling neural tube progenitors (Wood and Episkopou, 1999), and FOXC2 as

mesoderm marker, and combined IF-staining with the fluorescent labels of WT (mG)

and *Bra^-/-* cells (mT) (Fig. 4E-I). FOXA2 and CDH1 stainings indicate that *Bra^-/-* cells

are biased towards DE and favored to form the gut tube-like structure in the midline of

gastruloids (Fig. 4E, G, H). This finding not only demonstrates that *Brachyury* is

dispensable for DE lineage specification from pluripotent cells, but suggests that

Brachyury actually counteracts DE specification programs and thus may not represent

a suitable marker for early DE-forming cells as previously suggested (Kubo et al.,

2004). In addition to their contribution to endoderm structures of chimeric gastruloids,

Bra^-/- cells are predominantly found in the posterior region of chimeric gastruloids.

Here, *Bra^-/-* cells show SOX2 staining marking them as neural tube progenitors (Fig.

4F, G). Furthermore, *Bra^-/-* cells are rather excluded from the FOXC2-positive

mesoderm forming domain (Fig. 4I), in accordance with previous findings about NE-
repressive and mesoderm promoting functions of *Brachyury* (Koch et al., 2017; Tosic
et al., 2019).

In conclusion, this study illustrates the feasibility and experimental potency to use
functional genetics in mESCs to generate chimeric 3D-gastruloids. These allow for the
rapid and reproducible analysis of gain- and loss-of-gene functions, which may reveal
different phenotypic outcomes according to the degree of cellular contribution in
chimeric gastruloids. Close attention should be paid to control for the general variability
in mESC morphogenetic and differentiation behaviour, which might not strictly depend
on the genetic manipulation of each cell line. We did not experience these problems in
presented experiments, as they were all based on usage of mESCs lines originating
from the same parental clone (A2lox, Iacovino et al., 2014).

Here, we used the inducible expression of the Tbx transcription factor *Eomes* to
demonstrate how limitations of 3D gastruloids can be overcome by genetically
providing regulatory cues that are missing or underrepresented in CHIR-only treated
gastruloids, namely the induced formation of anterior mesoderm derivatives including
heart tissue. The analysis of *Brachyury*-deficient cells in chimeric gastruloids
remarkably reflects aspects previous studies of embryonic chimeras (Wilson et al.,
1993; Wilson et al., 1995; Wilson and Beddington, 1997). Here, $Bra^{-/-}$ cells
progressively accumulate in the tailbud region, most likely as result of *Brachyury*
functions in regulation of mesoderm cell migration leading to passive replacement of
mutant cells into the most posterior embryonic regions (Wilson et al., 1995; Wilson and
Beddington, 1997). It thus will be interesting to follow morphogenetic cell movements
of $Bra^{-/-}$ cells in chimeric gastruloids. In addition, the analysis of *Brachyury* deficient
cells in chimeras with low cell contribution showed a cell-autonomous bias towards the
DE, a cell lineage that strictly depends on *Eomes* functions (Arnold et al., 2008; Teo et
al., 2011). Since *Brachyury* and *Eomes* are co-expressed in cells of the early primitive
streak in early gastrulating embryos (Tosic et al., 2019), this poses the interesting
question about reciprocal regulatory interactions between these two related Tbx
factors.

In summary, this study demonstrates that chimeric gastruloids represent a powerful
experimental tool for the analysis of gastrulation stage embryogenesis. Gastruloids
already exhibit a high degree of experimental accessibility, observability and
scalability. The additional implementation of chimeric gastruloids allows to analyse

gene-functions at cellular level without affecting the general gastruloid architecture and might thus enhance our understanding of cell-signal interactions during gastrulation morphogenesis.

Materials and Methods

Cell lines

A2lox mESCs (Iacovino et al., 2014) were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 15 % fetal bovine serum (FBS, Gibco), 2 mM L-glutamine, 1X non-essential amino acids (NEAA), 1 mM sodium-pyruvate, 1X penicillin/streptomycin (all from Gibco), 100 µM β-mercaptoethanol (Sigma), Leukemia inhibitory factor (ESGRO LIF, Merck Millipore, 1000 U/ml), and 2i: CHIR99021 (Axon Medchem, 3 µM) and PD0325901 (Axon Medchem, 1 µM) on 0.1% gelatine-coated dishes. The medium was changed daily and mESCs were passaged every other day. The generation of homozygous $Bra^{-/-}$ mESCs and of A2lox mESCs harbouring monoallelic dox- inducible expression cassette for $Eomes$.GFP were described previously (Tosic et al., 2019). A2lox cells with membrane-tagged fluorescent labels (membrane-Tomato, mT, and membrane-GFP, mG) were generated by targeted integration of a mT/mG targeting vector (Muzumdar et al., 2007), Addgene plasmid #17787) into the Rosa26 locus. 1×10^6 A2lox mESCs (WT and $Bra^{-/-}$) were transfected with 2.5 µg of linearized vector using the Nucleofector ESC kit (Lonza) and G418 selected (350 µg/ml) on a monolayer of MitoC (Sigma)-mitotically inactivated STO feeder cells. mT-expressing ESC clones were picked on day 9 of selection. To convert the expression of the membrane-Tomato (mT) to membrane-GFP (mG) in WT A2lox mESCs, cells were treated for 24 h with 5 µg/ml Doxycyclin (Sigma, D9891) for induced expression of the Cre-recombinase from the Dox-inducible locus of A2lox WT cells to excise the loxP-flanked mT expression cassette and bring the mG expression cassette under the transcriptional control of the Rosa26 gene locus. After Cre-excision mG-expressing WT A2lox mESCs underwent one round of clonal selection by minimal dilution of 500 cells onto a 10 cm cell culture dish.

Generation of chimeric gastruloids

Gastruloids were generated using published protocols (van den Brink et al., 2020) with some modifications as outlined below. Gastruloid formation was performed in ESGRO Complete Basal Medium (Merck Millipore) in the absence of Matrigel. To generate

chimeric gastruloids using different mESCs by merging of preformed aggregates 150

cells of each ESC line were aggregated in 40 µl of ESGRO basal medium in 96-well

format (Greiner ultra-low attachment plates, No. 650970) for 24 h before merging by

combining two aggregates into the same well. At 48 h following first aggregate

formation, fused gastruloids were induced by administration of 3 µM CHIR and if

indicated with doxycycline (1 µg/µl, Sigma) for 24 h. In the course of gastruloid culture,

the medium was changed daily at 72, 96, 120 and 144 h. For the generation of

gastruloids by mixing of different ESC lines, aggregates were formed from a total of

300 mESCs at various ratios between the two ESC lines, and further gastruloid

formation followed by previously used protocols (van den Brink et al., 2020).

Whole mount *in situ* hybridization

Whole-mount *in situ* hybridization was performed according to standard protocols

using standard probes for *Mlc2a, Nkx2.5, Wnt3a*, and *Rspo3* (Details can be requested

from the authors). In brief, gastruloids were fixed in 4% PFA/PBS o/n at 4°C,

dehydrated via a methanol series and stored in methanol at -20°C. After rehydration

gastruloids were bleached in 6 % H_2O_2 for 5 min, digested by 1.6 µg/ml ProteinaseK

in PBT for 2 min, and postfixed in 4% PFA/0,2% glutaraldehyde for 20 min before

prehybridization for 2 h and hybridization o/n according to standard protocols. DIG-

labelled RNA probe was detected using anti-Digoxigenin-AP Fab fragments (Roche)

in 1% sheep serum, 2% BBR in MAB (0.1 M Maleic acid, 0.3 M NaCl, NaOH, 1%

Tween-20 in H_2O, pH 7.5) and incubation at 4°C o/n. Antibody was washed out by

extensive washes in MAB (>24 h, RT), and color reaction performed in BM purple

staining solution (Roche) for 2-6 h at RT.

Whole mount immunofluorescence staining

Gastruloids were fixed in 4% PFA /PBS for 1 h at 4°C, permeabilized (0.3% Triton X-

100/ PBT, 30 min) and blocked in 1% BSA / PBT for 1h at RT. Primary antibody

incubation was performed at 4°C o/n in 1% BSA / PBT, gastruloids washed 4x in PBT

before secondary fluorescence-conjugated antibody incubation for 3 h followed by

DAPI staining for 30 min at RT. Primary antibodies used were BRACHYURY (R&D

Systems; AF2085), CDX2 (Biogenex; MU392A) FOXA2 (Cell Signaling; 8186S), E-

Cadherin (BD Transduction Laboratories; 610182), SOX2 (R&D Systems; AF2018),

FOXC2 (R&D Systems; AF6989) at suggested dilutions. Secondary anti-goat, anti-

rabbit, anti-mouse and anti-sheep Alexa Fluor 647-conjugated antibodies (Thermo Fisher), and anti-goat CF405M (Biotium).

Histological sections of gastruloids

Fixed gastruloids were processed through 15% and 30% sucrose/PBS and incubated for at least 1h in embedding medium (15% sucrose, 7,5% gelatin in PBS) before cryo-embedding. Embedded gastruloids were cut with into 8 µm sections, mounted with ProLong Diamond Antifade Mountant (Life Technologies, P36970) and imaged as described below.

Imaging

Images were acquired on a Leica DMi8 Thunder Imager System or a Leica M165FC Stereo microscope. Images were processed in the Leica LASX software and Affinity Photo. During time lapse imaging gastruloids were maintained under constant conditions at 37°C, 5% CO_2.

Data acquisition and statistics

The percentage of beating gastruloids (Fig. 2D) was determined by manually counting during life imaging in three individual experiments. Following numbers of gastruloids were evaluated: 23 (+DOX) and 21 (-DOX) for n1; 61(+DOX) and 57 (-DOX) for n2, and 91(+DOX) and 90 (-DOX) for n3.

The length of the anterior-posterior axis of gastruloids (Fig. 2J) was measured using the LASX software (Leica) for n=34 (WTmG-WTmT) and n=36 (WTmG-TRE.*Eo*mT). Graphs (Scatter plot; Mean with SD) and column statistics were acquired with the GraphPad Prism software (Version 5.04).

Acknowledgements

We thank T. Bass for excellent technical assistance and Michael Kyba for the A2lox ESC line.

Funding

This work was supported by the German Research Foundation (DFG) through the Heisenberg Program (AR 732/3-1), project grant (AR 732/2-1,) project B07 of SFB 1140 (project ID 246781735), project A03 of SFB 850 (project ID 89986987), project

P7 of SFB 1453 (project ID 431984000), and Germany's Excellence Strategy (CIBSS – EXC-2189 – Project ID 390939984) to S.J.A., and by the MOTI-VATE program of the Freiburg Medical Faculty supported by the Else-Kröner-Fresenius-Stiftung to A.C.

Competing interests

The authors declare no competing interests.

Author contributions

A.E.W., K.M.S., A.C., C.M.S. generated different ESC lines and performed experiments. A.E.W, S.P. and S.J.A. planned and analysed experiments. A.E.W. and S.J.A. prepared figures, wrote and edited the manuscript with input from all authors. S.J.A. conceived the study.

References

Amadei, G., Lau, K. Y. C., De Jonghe, J., Gantner, C. W., Sozen, B., Chan, C., Zhu, M., Kyprianou, C., Hollfelder, F. and Zernicka-Goetz, M. (2021). Inducible Stem-Cell-Derived Embryos Capture Mouse Morphogenetic Events In Vitro. *Dev Cell* **56**, 366-382 e369.

Arnold, S. J., Hofmann, U. K., Bikoff, E. K. and Robertson, E. J. (2008). Pivotal roles for eomesodermin during axis formation, epithelium-to-mesenchyme transition and endoderm specification in the mouse. *Development* **135**, 501-511.

Arnold, S. J. and Robertson, E. J. (2009). Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. *Nat Rev Mol Cell Biol* **10**, 91-103.

Arnold, S. J., Stappert, J., Bauer, A., Kispert, A., Herrmann, B. G. and Kemler, R. (2000). Brachyury is a target gene of the Wnt/beta-catenin signaling pathway. *Mech Dev* **91**, 249-258.

Beccari, L., Moris, N., Girgin, M., Turner, D. A., Baillie-Johnson, P., Cossy, A. C., Lutolf, M. P., Duboule, D. and Arias, A. M. (2018). Multi-axial self-organization properties of mouse embryonic stem cells into gastruloids. *Nature* **562**, 272-276.

Costello, I., Pimeisl, I. M., Drager, S., Bikoff, E. K., Robertson, E. J. and Arnold, S. J. (2011). The T-box transcription factor Eomesodermin acts upstream of Mesp1 to specify cardiac mesoderm during mouse gastrulation. *Nat Cell Biol* **13**, 1084-1091.

Dunty, W. C., Jr., Biris, K. K., Chalamalasetty, R. B., Taketo, M. M., Lewandoski, M. and Yamaguchi, T. P. (2008). Wnt3a/beta-catenin signaling controls posterior body development by coordinating mesoderm formation and segmentation. *Development* **135**, 85-94.

Halpern, M. E., Ho, R. K., Walker, C. and Kimmel, C. B. (1993). Induction of muscle pioneers
and floor plate is distinguished by the zebrafish no tail mutation. *Cell* **75**, 99-111.

Harrison, S. E., Sozen, B., Christodoulou, N., Kyprianou, C. and Zernicka-Goetz, M.
(2017). Assembly of embryonic and extraembryonic stem cells to mimic embryogenesis
in vitro. *Science* **356**.

Harrison, S. E., Sozen, B. and Zernicka-Goetz, M. (2018). In vitro generation of mouse
polarized embryo-like structures from embryonic and trophoblast stem cells. *Nat Protoc*
13, 1586-1602.

Iacovino, M., Roth, M. E. and Kyba, M. (2014). Rapid genetic modification of mouse
embryonic stem cells by Inducible Cassette Exchange recombination. *Methods Mol Biol*
1101, 339-351.

Inman, K. E. and Downs, K. M. (2006). Brachyury is required for elongation and
vasculogenesis in the murine allantois. *Development* **133**, 2947-2959.

Kagawa, H., Javali, A., Khoei, H. H., Sommer, T. M., Sestini, G., Novatchkova, M., Scholte
Op Reimer, Y., Castel, G., Bruneau, A., Maenhoudt, N., et al. (2022). Human blastoids
model blastocyst development and implantation. *Nature* **601**, 600-605.

Koch, F., Scholze, M., Wittler, L., Schifferl, D., Sudheer, S., Grote, P., Timmermann, B.,
Macura, K. and Herrmann, B. G. (2017). Antagonistic Activities of Sox2 and Brachyury
Control the Fate Choice of Neuro-Mesodermal Progenitors. *Dev Cell* **42**, 514-526 e517.

Kubo, A., Shinozaki, K., Shannon, J. M., Kouskoff, V., Kennedy, M., Woo, S., Fehling, H.
J. and Keller, G. (2004). Development of definitive endoderm from embryonic stem cells
in culture. *Development* **131**, 1651-1662.

Li, R., Zhong, C., Yu, Y., Liu, H., Sakurai, M., Yu, L., Min, Z., Shi, L., Wei, Y., Takahashi,
Y., et al. (2019). Generation of Blastocyst-like Structures from Mouse Embryonic and
Adult Cell Cultures. *Cell* **179**, 687-702 e618.

Martin, B. L. and Kimelman, D. (2008). Regulation of canonical Wnt signaling by Brachyury
is essential for posterior mesoderm formation. *Dev Cell* **15**, 121-133.

Moris, N., Anlas, K., van den Brink, S. C., Alemany, A., Schroder, J., Ghimire, S., Balayo,
T., van Oudenaarden, A. and Martinez Arias, A. (2020). An in vitro model of early
anteroposterior organization during human development. *Nature* **582**, 410-415.

Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. and Luo, L. (2007). A global double-
fluorescent Cre reporter mouse. *Genesis* **45**, 593-605.

Probst, S., Sagar, Tosic, J., Schwan, C., Grun, D. and Arnold, S. J. (2021). Spatiotemporal
sequence of mesoderm and endoderm lineage segregation during mouse gastrulation.
Development **148**.

Rivera-Perez, J. A. and Magnuson, T. (2005). Primitive streak formation in mice is preceded
by localized activation of Brachyury and Wnt3. *Dev Biol* **288**, 363-371.

Rossi, G., Broguiere, N., Miyamoto, M., Boni, A., Guiet, R., Girgin, M., Kelly, R. G., Kwon,
C. and Lutolf, M. P. (2021). Capturing Cardiogenesis in Gastruloids. *Cell Stem Cell* **28**,
230-240 e236.

Teo, A. K., Arnold, S. J., Trotter, M. W., Brown, S., Ang, L. T., Chng, Z., Robertson, E. J.,
Dunn, N. R. and Vallier, L. (2011). Pluripotency factors regulate definitive endoderm
specification through eomesodermin. *Genes Dev* **25**, 238-250.

Tosic, J., Kim, G. J., Pavlovic, M., Schroder, C. M., Mersiowsky, S. L., Barg, M., Hofherr,
A., Probst, S., Kottgen, M., Hein, L., et al. (2019). Eomes and Brachyury control
pluripotency exit and germ-layer segregation by changing the chromatin state. *Nat Cell*
Biol **21**, 1518-1531.

Turner, D. A., Girgin, M., Alonso-Crisostomo, L., Trivedi, V., Baillie-Johnson, P.,
Glodowski, C. R., Hayward, P. C., Collignon, J., Gustavsen, C., Serup, P., et al.
(2017). Anteroposterior polarity and elongation in the absence of extra-embryonic tissues
and of spatially localised signalling in gastruloids: mammalian embryonic organoids.
Development **144**, 3894-3906.

Turner, D. A., Rue, P., Mackenzie, J. P., Davies, E. and Martinez Arias, A. (2014).
Brachyury cooperates with Wnt/beta-catenin signalling to elicit primitive-streak-like
behaviour in differentiating mouse embryonic stem cells. *BMC Biol* **12**, 63.

van den Brink, S. C., Alemany, A., van Batenburg, V., Moris, N., Blotenburg, M., Vivie, J.,
Baillie-Johnson, P., Nichols, J., Sonnen, K. F., Martinez Arias, A., et al. (2020). Single-
cell and spatial transcriptomics reveal somitogenesis in gastruloids. *Nature* **582**, 405-409.

van den Brink, S. C., Baillie-Johnson, P., Balayo, T., Hadjantonakis, A. K., Nowotschin,
S., Turner, D. A. and Martinez Arias, A. (2014). Symmetry breaking, germ layer
specification and axial organisation in aggregates of mouse embryonic stem cells.
Development **141**, 4231-4242.

van den Brink, S. C. and van Oudenaarden, A. (2021). 3D gastruloids: a novel frontier in
stem cell-based in vitro modeling of mammalian gastrulation. *Trends Cell Biol* **31**, 747-
759.

Veenvliet, J. V., Bolondi, A., Kretzmer, H., Haut, L., Scholze-Wittler, M., Schifferl, D.,
Koch, F., Guignard, L., Kumar, A. S., Pustet, M., et al. (2020). Mouse embryonic stem
cells self-organize into trunk-like structures with neural tube and somites. *Science* **370**.

Veenvliet, J. V., Lenne, P. F., Turner, D. A., Nachman, I. and Trivedi, V. (2021). Sculpting
with stem cells: how models of embryo development take shape. *Development* **148**.

Viotti, M., Nowotschin, S. and Hadjantonakis, A. K. (2014). SOX17 links gut endoderm
morphogenesis and germ layer segregation. *Nat Cell Biol* **16**, 1146-1156.

Wilson, V. and Beddington, R. (1997). Expression of T protein in the primitive streak is necessary and sufficient for posterior mesoderm movement and somite differentiation. *Dev Biol* **192**, 45-58.

Wilson, V., Manson, L., Skarnes, W. C. and Beddington, R. S. (1995). The T gene is necessary for normal mesodermal morphogenetic cell movements during gastrulation. *Development* **121**, 877-886.

Wilson, V., Rashbass, P. and Beddington, R. S. (1993). Chimeric analysis of T (Brachyury) gene function. *Development* **117**, 1321-1331.

Wood, H. B. and Episkopou, V. (1999). Comparative expression of the mouse Sox1, Sox2 and Sox3 genes from pre-gastrulation to early somite stages. *Mech Dev* **86**, 197-201.

Wymeersch, F. J., Wilson, V. and Tsakiridis, A. (2021). Understanding axial progenitor biology in vivo and in vitro. *Development* **148**.

Xu, P. F., Borges, R. M., Fillatre, J., de Oliveira-Melo, M., Cheng, T., Thisse, B. and Thisse, C. (2021). Construction of a mammalian embryo model from stem cells organized by a morphogen signalling centre. *Nat Commun* **12**, 3277.

Yamaguchi, T. P., Takada, S., Yoshikawa, Y., Wu, N. and McMahon, A. P. (1999). T (Brachyury) is a direct target of Wnt3a during paraxial mesoderm specification. *Genes Dev* **13**, 3185-3190.

Yu, L., Wei, Y., Duan, J., Schmitz, D. A., Sakurai, M., Wang, L., Wang, K., Zhao, S., Hon, G. C. and Wu, J. (2021). Blastocyst-like structures generated from human pluripotent stem cells. *Nature* **591**, 620-626.

FIGURE LEGENDS

Fig. 1. Experimental approaches for the generation of chimeric gastruloids

(A) Schematic of two alternative experimental approaches to generate chimeric gastruloids from fluorescently labelled embryonic stem cells (mESCs). Wildtype (WT) mESCs are marked by membrane-GFP (mG) and combined with membrane-Tomato (mT) labelled, genetically modified mESCs. The genetic modifications of mESCs comprise homozygous gene deletions (*GeneX*) or inducible gene-expression by the targeting of cDNAs into a fully controllable pre-engineered locus containing a Tetracycline responsive element (TRE) for dox-dependent induction of gene expression (*GeneY*). Chimeric gastruloids are generated by either mixing of mESCs at the beginning of gastruloid culture, or by merging preformed mESCs aggregates preceeding the induction of gastruloids by CHIR. Chimeric gastruloids can be used as versatile novel model system for various types of studies and embryonic research questions as indicated exemplary.

(B) Schematic of the protocol to form chimeric gastruloids indicating timepoints for either mixing of cells, or merging of aggregates to create chimeric gastruloids.

(C) Membrane-GFP (mG) labelling of WT mESCs and membrane-Tomato (mT) labelling for genetically modified mESCs allows for distinguishing different cell types in chimeric gastruloids. Scale bars 10 μm.

(D) Timelapse-imaging of the merging process of pre-formed ESC aggregates at indicated time points. 150 mG and mT mESCs were aggregated for 24 h before merging them by placing them together in 96-well plates. After 1 hr two aggregates spontaneously aggregated and formed stable contacts. Scale bars 100 μm. Also see **Supplementary Movie 1** for a 12 h time-lapse movie.

(E) Examples of mixed ESC aggregates at 24 h after aggregation of 300 mESCs by mixing mG WT mESCs and mT *Bra*$^{-/-}$ mESCs at indicated ratios. Scale bars 100 μm.

Fig. 2. Instructive functions of *Eomes* for cardiac lineage specification in merged chimeric gastruloids.

(A) Schematic illustrating the generation and culture of chimeric gastruloids by merging preformed aggregates of mG-labelled WT mESCs and mT-labelled mESCs harbouring *EomesGFP* in the Doxycycline (Dox)-inducible gene locus (TRE.*EomesGFP, short* TRE.*Eo*).

(B) Fluorescent microscopy of TRE.*Eo* mESCs showing nuclear expression of doxycycline-induced (+DOX) EOMES.GFP after 24 h of administration that is absent in -DOX conditions. Scale bars 10 μm.

(C) Brightfield (left) and fluorescent (right) images of a chimeric gastruloid at 168 h following induced *Eomes* expression showing the region of beating cardiomyocytes within the gastruloid that is mostly derived of mT-labelled TRE.*Eo* cells. The dashed line indicates the domain of beating cells. See also **Supplementary Movie 2** for the corresponding time-lapse movie.

(D) Statistics comparing the percentage of beating gastruloids at 168 h in gastruloids with forced *Eomes* expression +DOX (33.7% $\pm$ 3.0) and uninduced controls -DOX (1.5% $\pm$ 2.5) in n=3 independent experiments that are colour-encoded in orange, blue, and purple. Error bars indicate SD.

(E, F) Fluorescent microscopy (left) and whole-mount *in situ* hybridization (right) of the same chimeric gastruloids following induced *Eomes*-expression (+DOX) showing

instructive functions of forced *Eomes*-expression in TRE.*Eo* cells for the induction of
cardiac progenitors, marked by expression of (**E**) *Mlc2a* (n=12), and (**F**) *Nkx2.5* (n=10).
(G, H) Fluorescent microscopy (left) and immunofluorescence staining (right) showing
positioning of TRE.*Eo* cells at the opposite pole of posterior **(G)** BRACHYURY (a-BRA)
(n=24) and **(H)** CDX2 (n=10) expression. Scale bars 100 µm in C, E-H.

Fig. 3. *Brachyury* functions during axis-elongation in chimeric gastruloids
generated by cell mixing with high *Bra*$^{-/-}$ cell contribution.
(A) Schematic illustrating the generation of chimeric gastruloids by mixing of mG-
labelled WT mESCs and mT-labelled Brachyury-deficient (*Bra*$^{-/-}$) mESCs.
(B) Chimeric gastruloids at 120 h generated by mixing of *Bra*$^{-/-}$ (mT) and WT (mG)
mESCs at indicated ratios of cell numbers. Gastruloids with a contribution of *Bra*$^{-/-}$ cells
above 80 % show reduced axial elongation. Gastruloids entirely generated from *Bra*$^{-/-}$
cells fail to axially extend beyond an oval shape. At WT (mG) cell contribution > 50 %
axial elongation is similar to WT gastruloids but occasionally shows a thinning of the
posterior portion. Across all experiments *Bra*$^{-/-}$ (mT) cells preferentially contribute to the
posterior portion of mixed gastruloids. Anterior is oriented to the top and posterior to
the bottom in all images.
(C, D) Immunofluorescence staining against BRACHYURY (a-BRA) in gastruloids
generated by cell mixing of *Bra*$^{-/-}$ and WT cells in ratios of 80:20 (**C**) and 50:50 (**D**) at
indicated timepoints, showing less confined and reduced BRACHYURY expression in
80:20 chimeras at 96 and 120 h in comparison to posteriorly localized, robust
BRACHYURY expression in 50:50 chimeras. Images are representatives of n=14 (72
h); n=9 (96 h); n=14 (120 h) in (**C**), and n=16 (72 h); n=25 (96 h), n=18 (120 h)
gastruloids in (**D**).
(E) Whole-mount *in situ* hybridization for *Wnt3a* and *Rspo3* of WT, Bra$^{-/-}$-null and
chimeric gastruloids of indicated cell ratios at 72 h, 96 h, and 120 h showing
mislocalised and reduced expression of *Wnt3a* and *Rspo3* in Bra-deficient and
chimeric WT: Bra$^{-/-}$ mixed gastruloids. The frequencies of representative staining
pattern per numbers of examined gastruloid are indicated below each image. Scale
bars 100 µm in B-D.

Fig. 4. Brachyury functions for cell lineage specification and tissue sorting in
chimeric gastruloids with low *Bra*$^{-/-}$ cell contribution.

(A-D) Chimeric gastruloids generated by mixing of 10% $Bra^{-/-}$ (mT) and 90 % WT (mG)

ESCs at **(A)** 72 h and 96 h**,** and **(B)** three representative gastruloids at 120 h showing

the spectrum of distribution of $Bra^{-/-}$ cells between the posterior pole and the midline of

chimeric gastruloids.

(**C, D**) Control chimeric gastruloids generated by mixing of 10% WT (mT) and 90% WT

(mG) ESCs at **(C)** 72 h, 96 h, and **(D)** three replicates at 120 h don't show distinct

patterns of cell distribution.

(E-G) Immunofluorescence staining against (**E**) FOXA2 (n=6) and (**F**) SOX2 (n=8)

individually, and **(G)** double IF staining showing the tissue contribution of $Bra^{-/-}$ cells to

FOXA2+ endoderm and SOX2+ posterior cells (n=14).

(**H**) $Bra^{-/-}$ cells contribute to the forming primitive gut tubes (arrows) and are positive for

the epithelial marker CDH1 (E-Cadherin) (n=3).

(I) $Bra^{-/-}$ cells are mostly excluded from a mesodermal domain that is positive for

FOXC2 (n=7). Scale bars 100 µm.

A

WT ESCs

Genetically modified ESCs

⇨ *GeneX*

⇨ TRE *GeneY*

Merging
preformed aggregates

Mixing
cells

Chimeric Gastruloids

Advantages:

→ Versatile genetics
   by gain-and loss-of-function
→ direct observability
→ scalability
→ complex genetic settings
→ cell-autonomous vs.
   non-autonomous functions

B

Basal Medium	+CHIR/+DOX	Basal Medium

0 h   24 h   48 h   72 h   96 h   120 h

⇧ Mixing   ⇧ Merging

C

WT | Genetically modified

mG | mT

D

Merging aggregates   24 h

+ 0 h   + 1 h   + 2 h   + 4 h   + 8 h

E

Mixing cells   24 h

90:10   70:30   10:90

Fig. 1

A

0 h — 24 h — 48 h — 72 h — 96 h — 120 h — 144 h — 168 h

| Basal Medium | +CHIR +DOX | Basal Medium |

Merging

WT

TRE.*Eo*

B

TRE.*Eo* EOMES.GFP

+ DOX (24 h) — — - DOX

C

TRE.*Eo*
WT

+DOX — 168 h

D

% of beating gastruloids

+DOX — -DOX

E

TRE.*Eo*
WT

+DOX — *Mlc2a* — 168 h

F

TRE.*Eo*
WT

+DOX — *Nkx2.5* — 168 h

G

TRE.*Eo*
WT

+DOX — DAPI α-BRA — 120h

H

TRE.*Eo*
WT

+DOX — DAPI α-CDX2 — 120h

Fig. 2

Fig. 3

Fig. 4