A whole blood approach improves speed and accuracy when measuring mitochondrial

respiration in intact avian hematocytes

Authors: Andreas Nord[1], Imen Chamkha[2], Eskil Elmér[2]

[1] Lund University, Department of Biology, Section for Evolutionary Ecology, Sölvegatan 37,

SE-223 62 Lund, Sweden

[2] Lund University, Department of Clinical Sciences, Mitochondrial Medicine, Sölvegatan 17,

SE-221 84, Lund, Sweden

* Author for correspondence: AN, e-mail: andreas.nord@biol.lu.semi

ORCID numbers:

AN: 0000-0001-6170-689X

IM: 0000-0002-6919-585X

EE: 0000-0001-7901-1826

AUTHOR CONTRIBUTIONS:

AN and EE conceived the idea and designed the experimental protocol together with IC. AN

and IC performed the laboratory work. AN analysed the data and produced the graphical

material. All authors interpreted the results. AN wrote the first draft of paper, which was

edited by EE and IC. All authors approved submission and agreed to be accountable for all

contents.

ABSTRACT

Understanding mitochondrial biology and pathology is key to understanding the evolution of

animal form and function. However, mitochondrial measurement often involves invasive, or

even terminal, sampling, which can be difficult to reconcile in wild models or in longitudinal

studies. Non-mammal vertebrates contain mitochondria in their red blood cells, which can be

exploited for minimally invasive mitochondrial measurement. Several recent bird studies have

measured mitochondrial function using isolated blood cells. Isolation adds time in the

laboratory and might be associated with physiological complications. Inference may also be

constrained on biological grounds by lack of tissue context. We developed and validated a

protocol to measure mitochondrial respiration in bird whole blood. Endogenous respiration

was comparable between isolated blood cells and whole blood. However, oxidative

respiration was higher in whole blood, and whole blood mitochondria were better coupled and

had higher maximum working capacity. Whole blood measurement was also more

reproducible than measurement on isolated cells for all traits considered. Measurements were

feasible over a 10-fold range of sample volumes, though both small and large volumes were

associated with changes to respiratory traits. The protocol was compatible with long-term

storage: after 24 h at 5 °C without agitation all respiration traits but maximum working

capacity remained unchanged, the latter decreasing by 14%. Our study suggests that whole

blood measurement provides faster, more reproducible, and more biologically (tissue context)

and physiologically (mitochondrial integrity) relevant assessment of mitochondrial

respiration. We recommend future studies to take a whole blood approach unless specific

circumstances require the use of isolated blood cells.

Keywords: blood, blood cells, cell respiration, energy, erythrocyte, metabolism, mitochondria

INTRODUCTION

The ability to convert energy contained in food into a currency that can be used to sustain bodily housekeeping functions and key fitness-related traits such as growth, reproduction, movement, and survival, is a fundamental requirement for life. This occurs by digestion of food items into macromolecules that are subsequently used to create energy for working cells in the form of adenosine triphosphate (ATP). More than 90% of ATP is produced during cellular respiration in the mitochondria (Nicholls & Ferguson 2013). It is therefore not surprising that mitochondrial respiration and/or volume increase when organismal demand for energy increase (e.g. Liknes & Swanson 2011; Zheng *et al.* 2014; Dawson *et al.* 2016). Mitochondrial ATP is created by transport of electrons through a series of protein complexes (the electron transport system, ETS) whilst protons are being actively pumped across the inner mitochondrial membrane to establish electrochemical potential. The energy required for oxidative phosphorylation (OXPHOS) is harnessed when protons backflow into the mitochondrial matrix at complex V, the ATP-synthase. However, some protons forego OXPHOS by leaking back into the mitochondrial matrix spontaneously or under the influence of uncoupling proteins (Nicholls & Ferguson 2013). Proton leak thus uncouples mitochondrial respiration from ATP production (LEAK) but generates heat and reduces the rate at which potentially biodegrading reactive oxygen species (ROS) are formed (Brand *et al.* 2004; Speakman *et al.* 2004). It follows that there is intricate regulation of the balance between LEAK and OXPHOS, such that mitochondria can be made more effective for a given substrate oxidation rate when energy is in short supply (Trzcionka *et al.* 2008; Bourguignon *et al.* 2017; Roussel *et al.* 2020), or more thermogenic to aid endothermic heat production (Walter & Seebacher 2009).

Studies of mitochondrial function can provide proximate and ultimate depth in many

fundamental areas of research within ecology and evolution (see Koch *et al.* 2021 for a recent

review). However, such studies are rare in comparison to the rich body of literature on

organismal-level adaptation, function, and performance, particularly in the wild. This

probably reflects logistic and ethical considerations, because mitochondrial measurement

often requires terminal sampling for organ harvest or anaesthesia to collect biopsies, together

with speedy handling of samples to ensure organelle integrity; all of which are easier to

reconcile with laboratory models. However, birds and some other non-mammal vertebrates

retain the nucleus and functional organelles throughout erythroblast maturation and so contain

mitochondria in their erythrocytes (red blood cells) (Moritz, Lim & Wintour 1997; Moras,

Lefevre & Ostuni 2017). Studies on birds have found that this makes it possible to assess

mitochondrial function even from small (25-50 µl) blood samples (Stier *et al.* 2013; Stier *et*

al. 2017; Stier *et al.* 2019) that can be safely collected in organisms < 5 g (Kramer & Harris

2010). This is not possible when measurements are performed using other mitochondria-rich

hematocytes, as is necessary in mammals (e.g. Sjövall *et al.* 2013a; 2013b). While much is

still unknown about the function and regulation of blood cell mitochondria relative those in

other tissues, several recent bird studies indicate that blood cell respiration varies predictably

in line with organismal-level energy demand over long (Stier *et al.* 2019; Casagrande *et al.*

2020) and short (Garcia Diaz et al. 2022) time frames, and in the contexts of seasonal

adaptation (Nord *et al.* 2021) and metabolic senescence (Dawson & Salmón 2020). Moreover,

blood cell mitochondrial function seems amenable to epigenetic programming during

embryonic development (Udino *et al.* 2021; Stier, Monaghan & Metcalfe 2022), much like

what is a well-known fact within human medicine (Gyllenhammer *et al.* 2020). Studies have

even found that blood cell respiration can potentially be used to predict variation in whole-

organism metabolic rate (Malkoc, Casagrande & Hau 2021). Thus, there is ample evidence

that mitochondrial measurement in red blood cells can provide minimally invasive insights

into several key areas within the broader fields of ecology, ecological physiology, and

evolutionary biology.

To the best of our knowledge, all studies on avian blood cell mitochondria have been

performed using isolated blood cells, prepared using standard laboratory techniques including

pipetting, vortexing and centrifugation. These procedures could impact the status of the cells

going into the experiment (e.g. Wiegmann *et al.* 2017). If standard handling impacts cell

health, mitochondrial function in isolated hematocytes may not be as representative of the *in*

vivo state, or could introduce experimental variation, but this notion remains untested.

Furthermore, while erythrocytes will be responsible for most of the respiration in the avian

circulation (c.f. Scanes 2022), it can be difficult to reliably separate cell types by

centrifugation in the small samples that are typically collected from wild models. Thus, the

exact composition of a cell isolate may, or may not, correspond to that in whole blood.

Moreover, on conceptual grounds it can be argued that isolate-based measurements lack some

biological value   since the removal of plasma before measurement effectively also removes

the tissue-context of the blood. On this background, we asked whether mitochondrial

measurement on whole blood samples were comparable to measurements   on isolated

hematocytes from the same blood volume. If cell isolation impacts mitochondrial integrity or

function of the electron transport chain, we predicted that isolate measurements would be

associated with leakier mitochondria and/or lower overall respiration compared to whole

blood samples. To explore the utility of whole blood measurement, we proceeded by testing

the resilience of the assay to variation in sample volume and long-term storage. Finally, we

undertook initial validation of a protocol for permeabilising whole blood to permit studies of

mitochondrial function with unlimited substrate supply.

MATERIALS AND METHODS

Animals and housing

The study was performed using captive zebra finches (*Taeniopygia guttata* Vieillot) (*n* = 23;

12 females and 11 males). All birds were in their second calendar year, or older, and body

masses ranged 12.1 to 19.4 g. The birds originated from a captive population kept in outdoor

flight aviaries at the Lund University field station Stensoffa (WGS84 DD: N55.69534,

E13.44739) and were brought to light- and temperature controlled indoor facilities at Lund

University 5-7 d before the experiment started. Birds were kept in cages measuring 120×80

$\times 100$ cm (length $\times$ depth $\times$ height) at a density of ≤ 6 birds to a cage. Room temperature was

maintained at 19-20°C and photoperiod was 12:12 LD. Food (mixed millet seeds), cuttlebone,

and water was provided *ad libitum.*

Anaesthesia and blood sampling

We sterilised the ventral skin area covering the distal part of the sternum using 70% ethanol

and immediately anaesthetised the birds by intraperitoneal injection of pentobarbital sodium

(30-40 µg per g body mass). When the bird was reactionless to an external stimulus (toe

pinching), which occurred within 6-10 min after the injection, we collected a maximal blood

sample (250-650 µl) from the jugular vein and then euthanised the bird by cervical

dislocation. Samples were stored at 10-15°C in 2 mL K2-EDTA (ethylenediaminetetraacetic

acid) tubes (BD Vacutainer®, Franklin Lakes, NJ, USA) without agitation until analysed 15 to

45 min later.

Mitochondrial respiration measurements and experiments

Mitochondrial respiration was measured at bird body temperature (41°C) (Prinzinger,

Pressmar & Schleucher 1991) using Oxygraph O2k high-resolution respirometers (Oroboros

Instruments, Innsbruck, Austria). Respiration was inferred from the decline in O_2

concentration in a 2 ml suspension of sample and respirometry medium (MiR05: 0.5 mM of

EGTA, 3 mM of $MgCl_2$, 60 mM of K-lactobionate, 20 mM of taurine, 10 mM of KH_2PO_4, 20

mM of HEPES, 110 mM of sucrose, and 1 g l^{-1} free fatty acid bovine serum albumin, pH 7.1).

All measurements were performed on intact cells. We allowed 10-15 min for O_2 consumption

to stabilise after adding sample to the chambers, and then recorded baseline phosphorylating

respiration on endogenous substrates for 2-4 min ("ROUTINE"). We then added 1 μg ml^{-1}

oligomycin to inhibit ATP synthase, thus preventing oxidative phosphorylation. The

remaining respiration in this stage is used to offset the leak of protons that occur across the

inner mitochondrial membrane ("LEAK"). Thus, the part of respiration devoted to oxidative

phosphorylation ("OXPHOS") can be derived as the difference between the ROUTINE and

LEAK states. This was followed by titrating the mitochondrial uncoupler carbonyl cyanide-p-

trifluoro-methoxyphenyl-hydrazone (FCCP) in 0.5 μl, 1 mM, aliquots until maximum

respiration was reached (final concentration: 1-2 μM). At maximal non-inhibiting

concentration, FCCP abolishes the proton gradient across the inner mitochondrial membrane,

forcing the electron transport system to work at its maximum capacity ("ETS") to restore it.

After FCCP titration, we inhibited mitochondrial complex I, using 2 μM rotenone, and then

added 1 μg ml^{-1} of the complex III inhibitor antimycin A to stop electron transport. Any

respiration remaining after addition of antimycin A is considered of non-mitochondrial origin.

There was never a meaningful further reduction in respiration when antimycin A was added

onto rotenone (data not shown). Hence, we adjusted all respiratory states for non-

mitochondrial respiration by subtracting whichever was the lowest of respiration on rotenone

and rotenone+antimycin A, under the assumption that any absolute differences between the

two were due to random noise.

In the first experiment, we compared mitochondrial respiration traits measured in whole blood

and isolated blood cells. Data were collected from 7 birds (3 males, 4 females) and all

samples were run in simultaneous duplicates within 45 min of collection. We manually mixed

the blood by gently tilting the sample tubes for 4-5 min. Then, we collected two 100 µl whole

blood samples to use either for immediate respiration measurement, or for isolating the

hematocytes. We isolated the blood cells following Stier et al. (2017) with slight

modifications according to Dawson & Salmón (2020) and Nord et al. (2021). Thus, the

sample was first centrifuged at 3000 g for 10 min at room temperature to separate plasma

from the blood cells. The pellet was then re-suspended in 500 µl cold MiR05, centrifuged at

1000 g for 5 min and the supernatant was discarded. The remaining pellet, containing all cells

from the 100 µl blood sample, was immediately re-suspended in 750 µl MiR05 pre-

equilibrated at 41 °C and added to 1.35 ml MiR05 contained in the respiration chamber (final

volume 2 ml) for use in the assay. The sample volume used here is the same as in the original

protocol for mitochondrial measurement in bird blood cells (Stier *et al.* 2017) and is within

range of that used in other studies (e.g. Dawson & Salmón 2020; Nord *et al.* 2021).

In the second experiment, we asked how blood sample volume affected respiration traits,

attempting to refine the protocol by defining the lowest possible volume yielding

representative data. Blood samples were drawn and managed as above ($n = 7$; 3 females, 4

males). Mitochondrial respiration traits were then assayed simultaneously in 10, 25, 50 and

100 µl whole blood samples. The volume of MiR05 in the chambers was adjusted accordingly

to maintain a total volume of 2 mL

In the third experiment, we investigated the effect of storage time on whole blood respiration,

using data from 6 birds (3 females, 3 males). Within 30 min of blood sampling (henceforth,

time = '0 h'), we followed the protocol above to measure mitochondrial respiration in 50 µl

blood samples in 1.95 mL MiR05. The tube with remaining blood was then stored without

agitation at 5.2 ± 0.7 °C (mean $\pm$ s.d.; range 3.9 to 6.5 °C) and another 50 µl aliquot from the

same sample was measured again (after 5 min of mixing) 24 h later (henceforth, time = '24

h'). Samples from each of two birds were also opportunistically measured at 72 h and 96 h

after collection.

Data analyses

We calculated three flux control ratios (FCR) to address integrity of mitochondrial respiration

following the method and terminology proposed by Gnaiger (2020): 1) E-R control efficiency

(1 - ROUTINE / ETS), which is a measure of the proportion of maximum working capacity

remaining during endogenous respiration; 2) R-L control efficiency ((ROUTINE - LEAK) /

ROUTINE), which is the proportion of endogenous respiration channelled towards ATP

production via oxidative phosphorylation; 3) E-L coupling efficiency (1 - LEAK / ETS),

which is indicative of how 'tightly' electron transport is coupled to ATP production under a

stimulated cellular state.

Statistical analyses were performed using R 4.1.2 for Windows (R Core Team 2021). To

compare the utility of blood cells and whole blood (i.e., Experiment 1), we fitted trait- and

FCR-specific linear mixed effect models (lmer in the lme4 package) (Bates *et al.* 2015) with

sample type as a factor and bird ID as a random intercept to account for the dependence of

observations between duplicates and samples. Repeatability of traits (*R*) for each sample type

was expressed as the intraclass correlation coefficient with 95% confidence intervals based on

1000 bootstrap iterations using the rptR package (Stoffel, Nakagawa & Schielzeth 2017). We

used lmer models to address the effect of sample volume on respiration rates and FCRs (i.e.,

Experiment 2) using the same general model structure. Predicted values and their standard

errors were calculated using the emmeans package (Lenth 2019) and used the pairs function

in this package to perform post hoc tests for Experiment 2 data. The effects of storage time

(i.e., Experiment 3) were assessed using paired t-tests between the 0 h and 24 h samples (t.test

function in R base).

RESULTS

A representative trace of oxygen consumption in a whole blood experiment is provided in the

Electronic Supplementary Materials (Fig. S1 in ESM1).

Effects of sample type – blood cells vs. whole blood

ROUTINE respiration did not differ between isolated blood cell and whole blood samples

(Table 1, Fig. 1A). However, OXPHOS and ETS were significantly higher when measured on

whole blood samples compared to on blood cell isolates (1.2 and 1.3-fold, respectively) (Figs

1B-C, Table 1). In contrast, LEAK was 30% lower in the whole blood samples (Fig. 1D,

Table 1). Consequently, the ER control efficiency (which measures proportionally how much

respiration can be increased from ROUTINE) was significantly higher in the whole blood

samples (0.511, compared to 0.365 in isolated cells; Fig. 2A, Table 1). Whole blood samples

also had significantly higher phosphorylating efficiency and higher coupling efficiency during

maximal FCCP-stimulated respiration (i.e., R-L control and E-L coupling efficiencies were

higher) (Fig. 2B-C, Table 1).

Respiration traits measured on both whole blood and isolated blood cells were significantly

and highly reproducible (blood cells: R_{min} = 0.790, R_{maz} = 0.968; whole blood: R_{min} = 0.849,

R_{max} = 0.985). However, whole blood samples were more reproducible, and the calculated

repeatabilities had markedly narrower confidence intervals, for all respiration traits and FCRs

investigated (Table 1).

Effects of blood volume

There were non-linear effects on blood sample volume on respiration traits and FCRs, with

the lowest and highest sample volumes showing the most pronounced deviations.

Accordingly, ROUTINE was significantly higher (by a factor 1.5 to 1.6) for all volumes 10-

50 µl compared to the 100 µl sample (Fig. 3A, Table 2). OXPHOS was the highest, and

similar for 25 and 50 µl, and the lowest for 10 and 100 µl (which did not differ) (Fig. 3A,

Table 2). OXPHOS measured using 50 µl was 1.3-fold higher than the measurement using 10

µl, though this difference was not statistically significant. ETS was the highest on 50 µl

samples (up to 1.4-fold), and significantly different from all volumes but 25 µl (despite a 1.2-

fold difference; Table 2). ETS was the lowest on 100 µl samples, though this volume differed

statistically only from 50 µl (Table 2). LEAK was significantly higher on 10 µl compared to

all other volumes, and the lowest on 100 µl (though this volume was not statistically different

from 50 µl) (Table 2).

FCRs, which provide rate-independent insight into mitochondrial function, were also volume-

specific. Specifically, E-R control efficiency was similar in 50 and 100 µl and significantly

higher (reflecting more surplus capacity) compared to the two other volumes (Fig. 3B). R-L

control efficiency (i.e., the efficiency with which O_2 consumption is channelled towards

oxidative phosphorylation) was the lowest in 10 µl (reflecting higher LEAK) compared to the

other volumes (Table 2, Fig. 3B). E-L coupling efficiency was also the lowest in 10 µl

followed by 25 µl, but higher in both 50 and 100 µl (which were not statistically different)

(Table 2, Fig. 3B).

Effects of storage

Endogenous (ROUTINE), phosphorylating (OXPHOS) and leak (LEAK) respiration did not

change significantly with 24 h storage at 5°C (all $P > 0.2$) (Fig. 4A, Table 3). However

maximal respiration capacity (ETS) was 16% lower at 24 h compared to at 0 h. As a result, E-

R control efficiency and E-L coupling efficiency was significantly lower at 24 h (Fig. 4B,

Table 3). However, there was no change in R-L control efficiency, in keeping with stable

ROUTINE and LEAK over 24 h storage. By 72 h, ROUTINE, OXPHOS and ETS had

decreased somewhat from the 24 h level and LEAK had increased, with corresponding

changes to FCRs (Fig. S2 in ESM2). These changes became more exaggerated with 96 h

storage (Fig. S2).

DISCUSSION

In this study, we asked whether measurement on whole blood could provide a feasible and

potentially more biologically contextualised approach to studying mitochondrial respiration in

intact avian blood cells. While endogenous respiration did not differ between whole blood

samples and isolated cells produced from the same blood volume, we found that both

phosphorylating (i.e., OXPHOS; Fig. 1B) and maximal (i.e., ETS; Fig. 1C) respiration was

significantly higher in whole blood, and that mitochondria in isolated blood cells displayed

significantly more LEAK (i.e., they were more uncoupled) than whole blood (Fig. 1D). This

translated to significant differences in flux control ratios (FCRs), which indicated that the

isolated blood cells had lower surplus respiratory capacity (Fig. 2A), lower phosphorylating

efficiency (Fig. 2B) and reduced coupling efficiency when stimulated to work at maximum

(Fig. 2C). Moreover, while our repeatability estimates for isolated erythrocytes were

consistently high and mostly within the range for the same traits reported elsewhere (Stier et

al. 2017), the whole blood samples were consistently more reproducible (all $R > 0.93$) and

less variable for all traits considered (Table 1). Thus, in our study whole blood measurement

was more precise, and biological inference differed in several key traits depending on whether

measurements were performed on the intact tissue or on a derived sample. We believe that

this could be caused by physical damage or reduced integrity of mitochondria in the isolated

blood cells caused by more handling, because lower ETS, higher LEAK and associated

changes to FCRs in repeated samples of the same subject are all indicative of declining

mitochondrial health/viability (e.g. Sjövall *et al.* 2013a) (see also Fig. 4). Studies on

mammalian erythrocytes show centrifugation alone can cause time-dependent structural and

physiological damage to red blood cells (such as increased free and corpuscular haemoglobin

and increased ATP release) at spinning speeds considerably lower than those used here ($<$

1500 *g*) (Kiss *et al.* 2016; Urbina *et al.* 2016; Wiegmann *et al.* 2017; Mancuso, Jayaraman &

Ristenpart 2018). We are not suggesting that our whole blood samples were in a 'pristine'

condition, because drawing blood into a syringe, and subsequent pipetting of a blood sample,

may also impact the physiological and physical state of the red blood cells (Wiegmann *et al.*

2017). However, our study indicates that whole blood measurement may salvage more of the

physiological integrity of the avian blood cell.

We found non-linearity of volume-adjusted respiration rates, manifested as inhibited or

enhanced respiration on the highest and lowest blood volumes (Fig. 3A). On average, optimal

trait combinations for coupled and uncoupled respiration were achieved using 50 µl samples,

but these runs were most often comparable to those on 25 µl. FCR for phosphorylating and

coupling efficiencies were similar in the range of 25 to 100 µl, whereas 10 µl samples were

less well coupled. Thus, working in a range of 25 to 50 µl whole blood provided

representative results for respiration traits, whereas a sample volume range of 25 to 100 µl

was acceptable if FCRs only were considered. Our findings contrast previous studies on pig

liver homogenates where complete linearity of phosphorylating respiration was demonstrated

across a 3-fold range in sample volume (Kuznetsov *et al.* 2002). However, studies on isolated

human platelets show reduced uncoupled respiration on small sample volumes (Sjövall *et al.*

2013a). This could be caused e.g., by reduced accuracy on low sample concentrations acting

in combination with enhanced sensitivity/exposure to inhibitors. Measurements on pig

skeletal muscle mitochondria also suggest suppressed respiration when samples were

prepared from smaller biopsies (Isner-Horobeti *et al.* 2014), though this probably reflects

heterogeneity in sample composition that was not accounted for by weight alone. It is less

clear why we observed inhibition of respiration (but not FCRs) on large blood sample

volumes. We do not believe that this was caused by the presence of plasma in the sample for

at least two reasons: 1) both intact and open blood cells can be measured using plasma as

medium with inhibition of respiration (Sjövall *et al.* 2010; Grip *et al.* 2016); 2) ROUTINE

was identical in whole blood and isolated blood cells (Fig. 1A), which is not expected if some

whole blood constituents act suppressively. The explanation for these observations should be

addressed in future work, including measurement of mitochondrial content in differently sized

samples, and using both intact and open-cell protocols. Meanwhile, we recommend

experimenters to use relatively constant sample volumes to the furthest extent possible.

The whole blood samples were robust to cold storage. After 24 h at 5°C without agitation, all

respiration traits but ETS (which decreased by 14%) remained unaltered. This corroborates

previous work on both human and avian blood cells (Sjövall *et al.* 2013a; Stier *et al.* 2017)

and pig liver homogenates (Kuznetsov *et al.* 2002). However, neither Sjövall et al. (2013) or

Stier et al. (2017) found any reduction in ETS over 24 h. These authors measured the cells 3-5

h after blood collection, whereas our first measurement was performed within 45 min of

bleeding the birds. It is tempting to speculate that most changes to ETS occur within the first

hour of collection. Sjövall (2013) found that respiration of permeabilised human platelets

remained largely unaltered up to 48 h when the blood was stored at room temperature. This

does not seem to be the case in birds: while ROUTINE, OXPHOS and LEAK remained

relatively stable when the blood was stored 2-2.5 h at room temperature under constant

motion, ETS dropped by 30 to 90% over the same period (data not shown).

CONCLUSIONS

We found that whole blood measurement provides more rapid, more biologically

contextualised, and more precise and reproducible assessment of mitochondrial respiration in

intact avian hematocytes compared to measurement on isolated cells. We also give data-based

recommendations on sample volume ranges, show that 24 h cold-storage in the blood

collection tubes is possible at acceptable changes to respiration traits and that whole blood can

be permeabilised. Changes to respiration parameters of the intact cells were indicative of

handling-induced damage. However, it is possible that less extensive handling protocols (see

e.g. Stier *et al.* 2019) will provide more comparable results. Even so, our study suggests that

centrifugation and removal of plasma are unnecessary complications that add processing time

and reduce biological context. Thus, we advocate use of whole blood whenever possible.

When measurements on isolated cells are necessary, such as when low sample volume is not

sufficient for complementary analyses of plasma metabolites, care should be taken to keep

mechanical stress to a minimum. We also recommend use of relatively constant blood

volumes to minimize variation within an experiment, and suggest samples are run on the day

of collection.

ACKNOWLEDGEMENTS

The authors thank Elsie Ye Xiong and Michael Tobler for providing zebra finches for use in

the study. Camilla Björklöv and Agnieszka Czopek excellently assisted with care for the

experimental animals. Comments from Elisa Thoral improved a previous version of the

manuscript.

FUNDING

This study was supported by the Royal Physiographic Society / The Birgit and Hellmuth

Hertz Foundation (grant no. 2017-39034) and the Swedish Research Council (grant no. 2020-

04686) (to AN).

ETHICS

All procedures were approved by the Malmö/Lund Animal Ethics Committee, acting under

authority of the Swedish Board of Agriculture (permit no. 9246-19, 16977-19).

DATA AVAILABILITY

Data will be deposited in Figshare upon acceptance of the manuscript.

CONFLICT OF INTEREST

The authors declare that they have no competing or commercial interests.

REFERENCES

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software,* **67,** 1-48.

Bourguignon, A., Rameau, A., Toullec, G., Romestaing, C. & Roussel, D. (2017) Increased

mitochondrial energy efficiency in skeletal muscle after long-term fasting: its

relevance to animal performance. *Journal of Experimental Biology,* **220,** 2445-2451.

Brand, M.D., Buckingham, J.A., Esteves, T.C., Green, K., Lambert, A.J., Miwa, S., Murphy,

M.P., Pakay, J.L., Talbot, D.A. & Echtay, K.S. (2004) Mitochondrial superoxide and

aging: uncoupling-protein activity and superoxide production. *Biochemical Society*

*Symposium***,** 203-213.

Casagrande, S., Stier, A., Monaghan, P., Loveland, J.L., Boner, W., Lupi, S., Trevisi, R. &

Hau, M. (2020) Increased glucocorticoid concentrations in early life cause

mitochondrial inefficiency and short telomeres. *Journal of Experimental Biology,* **223,**

jeb222513.

Dawson, N.J., Ivy, C.M., Alza, L., Cheek, R., York, J.M., Chua, B., Milsom, W.K.,

McCracken, K.G. & Scott, G.R. (2016) Mitochondrial physiology in the skeletal and

cardiac muscles is altered in torrent ducks, *Merganetta armata*, from high altitudes in

the Andes. *Journal of Experimental Biology,* **219,** 3719-3728.

Dawson, N.J. & Salmón, P. (2020) Age-related increase in mitochondrial quantity may

mitigate a decline in mitochondrial quality in red blood cells from zebra finches

(*Taeniopygia guttata*). *Experimental Gerontology,* **133,** 110883.

Gnaiger, E. (2020) Mitochondrial pathways and respiratory control: An Introduction to

OXPHOS Analysis. 5th ed.

Grip, J., Jakobsson, T., Tardif, N. & Rooyackers, O. (2016) The effect of plasma from septic

ICU patients on healthy rat muscle mitochondria. *Intensive Care Medicine*

Experimental, **4,** 20.

Gyllenhammer, L.E., Entringer, S., Buss, C. & Wadhwa, P.D. (2020) Developmental

programming of mitochondrial biology: a conceptual framework and review.

Proceedings of the Royal Society B, **287,** 20192713.

Isner-Horobeti, M.E., Charton, A., Daussin, F., Geny, B., Dufour, S.P. & Richard, R. (2014)

Microbiopsies versus Bergström needle for skeletal muscle sampling: impact on

maximal mitochondrial respiration rate. *European Journal of Applied Physiology,* **114,**

885-889.

Kiss, F., Toth, E., Miszti-Blasius, K. & Nemeth, N. (2016) The effect of centrifugation at

various g force levels on rheological properties of rat, dog, pig and human red blood

cells. *Clinical Hemorheology and Microcirculation,* **62,** 215-227.

Koch, R.E., Buchanan, K.L., Casagrande, S., Crino, O., Dowling, D.K., Hill, G.E., Hood,

W.R., McKenzie, M., Mariette, M.M., Noble, D.W.A., Pavlova, A., Seebacher, F.,

Sunnucks, P., Udino, E., White, C.R., Salin, K. & Stier, A. (2021) Integrating

Mitochondrial Aerobic Metabolism into Ecology and Evolution. *Trends in Ecology &*

Evolution, **36,** 321-332.

Kramer, M.H. & Harris, D.J. (2010) Avian Blood Collection. *Journal of Exotic Pet Medicine,*

19, 82-86.

Kuznetsov, A.V., Strobl, D., Ruttmann, E., Königsrainer, A., Margreiter, R. & Gnaiger, E.

(2002) Evaluation of mitochondrial respiratory function in small biopsies of liver.

Analytical Biochemistry, **305,** 186-194.

Lenth, R.V. (2019) emmeans: Estimated marginal means, aka least-squares means. R package

version 1.3.3. https://CRAN.R-project.org/package=emmeans.

Liknes, E.T. & Swanson, D.L. (2011) Phenotypic flexibility in passerine birds: Seasonal

variation of aerobic enzyme activities in skeletal muscle. *Journal of Thermal Biology,*

36, 430-436.

Malkoc, K., Casagrande, S. & Hau, M. (2021) Inferring Whole-Organism Metabolic Rate

452   From Red Blood Cells in Birds. *Frontiers in Physiology,* **12,** 691633-691633.

453   Mancuso, J.E., Jayaraman, A. & Ristenpart, W.D. (2018) Centrifugation-induced release of

454   ATP from red blood cells. *PLoS One,* **13,** e0203270.

455   Moras, M., Lefevre, S.D. & Ostuni, M.A. (2017) From Erythroblasts to Mature Red Blood

456   Cells: Organelle Clearance in Mammals. *Frontiers in Physiology,* **8,** 201701076.

457   Moritz, K.M., Lim, G.B. & Wintour, E.M. (1997) Developmental regulation of erythropoietin

458   and erythropoiesis. *American Journal of Physiology,* **273,** R1829-R1844.

459   Nicholls, D.G. & Ferguson, S.J. (2013) *Bioenergetics,* 4th ed. Academic Press, Cambridge,

460   MA, USA.

461   Nord, A., Metcalfe, N.B., Page, J.L., Huxtable, A., McCafferty, D.J. & Dawson, N.J. (2021)

462   Avian red blood cell mitochondria produce more heat in winter than in autumn. *The*

463   *FASEB Journal,* **35,** e21490.

464   Prinzinger, R., Pressmar, A. & Schleucher, E. (1991) Body temperature in birds. *Comparative*

465   *Biochemistry and Physiology A,* **99,** 499-506.

466   R Core Team (2021) R: A language and environment for statistical computing. R Foundation

467   for Statistical Computing, Vienna, Austria.

468   Roussel, D., Marmillot, V., Monternier, P.-A., Bourguignon, A., Toullec, G., Romestaing, C.

469   & Duchamp, C. (2020) Skeletal muscle metabolism in sea-acclimatized king penguins.

470   II. Improved efficiency of mitochondrial bioenergetics. *Journal of Experimental*

471   *Biology,* **223,** jeb233684.

472   Scanes, C.G. (2022) Blood. *Sturkie's Avian Physiology* (ed. C.G. Scanes, Dridi, S). Academic

473   Press, Cambridge, Massachusetts.

474   Sjövall, F., Ehinger, J.K.H., Marelsson, S.E., Morota, S., Åsander Frostner, E., Uchino, H.,

475   Lundgren, J., Arnbjörnsson, E., Hansson, M.J., Fellman, V. & Elmér, E. (2013a)

Mitochondrial respiration in human viable platelets—Methodology and influence of gender, age and storage. *Mitochondrion,* **13,** 7-14.

Sjövall, F., Morota, S., Hansson, M.J., Friberg, H., Gnaiger, E. & Elmér, E. (2010) Temporal increase of platelet mitochondrial respiration is negatively associated with clinical outcome in patients with sepsis. *Critical Care,* **14,** R214.

Sjövall, F., Morota, S., Persson, J., Hansson, M.J. & Elmér, E. (2013b) Patients with sepsis exhibit increased mitochondrial respiratory capacity in peripheral blood immune cells. *Critical Care,* **17,** R152.

Speakman, J.R., Talbot, D.A., Selman, C., Snart, S., McLaren, J.S., Redman, P., Krol, E., Jackson, D.M., Johnson, M.S. & Brand, M.D. (2004) Uncoupled and surviving: individual mice with high metabolism have greater mitochondrial uncoupling and live longer. *Aging Cell,* **3,** 87-95.

Stier, A., Bize, P., Hsu, B.-Y. & Ruuskanen, S. (2019) Plastic but repeatable: rapid adjustments of mitochondrial function and density during reproduction in a wild bird species. *Biology Letters,* **15,** 20190536.

Stier, A., Bize, P., Schull, Q., Zoll, J., Singh, F., Geny, B., Gros, F., Royer, C., Massemin, S. & Criscuolo, F. (2013) Avian erythrocytes have functional mitochondria, opening novel perspectives for birds as animal models in the study of ageing. *Frontiers in Zoology,* **10,** 33.

Stier, A., Monaghan, P. & Metcalfe, N.B. (2022) Experimental demonstration of prenatal programming of mitochondrial aerobic metabolism lasting until adulthood. *Proceedings of the Royal Society B,* **289,** 20212679.

Stier, A., Romestaing, C., Schull, Q., Lefol, E., Robin, J.-P., Roussel, D. & Bize, P. (2017) How to measure mitochondrial function in birds using red blood cells: a case study in

the king penguin and perspectives in ecology and evolution. *Methods in Ecology and Evolution,* **8,** 1172-1182.

Stoffel, M.A., Nakagawa, S. & Schielzeth, H. (2017) rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution,* **8,** 1639-1644.

Trzcionka, M., Withers, K.W., Klingenspor, M. & Jastroch, M. (2008) The effects of fasting and cold exposure on metabolic rate and mitochondrial proton leak in liver and skeletal muscle of an amphibian, the cane toad *Bufo marinus. Journal of Experimental Biology,* **211,** 1911-1918.

Udino, E., George, J.M., McKenzie, M., Pessato, A., Crino, L., Buchanan, K.L. & Mariette, M.M. (2021) Prenatal acoustic programming of mitochondrial function for high temperatures in an arid-adapted bird. *Proceedings of the Royal Society B,* **288,** 20211893.

Urbina, A., Godoy-Silva, R., Hoyos, M. & Camacho, M. (2016) Acute hydrodynamic damage induced by SPLITT fractionation and centrifugation in red blood cells. *Journal of Chromatography B,* **1020,** 53-61.

Walter, I. & Seebacher, F. (2009) Endothermy in birds: underlying molecular mechanisms. *Journal of Experimental Biology,* **212,** 2328-2336.

Wiegmann, L., de Zélicourt, D.A., Speer, O., Muller, A., Goede, J.S., Seifert, B. & Kurtcuoglu, V. (2017) Influence of standard laboratory procedures on measures of erythrocyte damage. *Frontiers in Physiology,* **8**.

Zheng, W.H., Li, M., Liu, J.-S., Shao, S.-L. & Xu, X.-J. (2014) Seasonal variation of metabolic thermogenesis in Eurasian tree Sparrows (*Passer montanus*) over a latitudinal gradient. *Physiological and Biochemical Zoology,* **87,** 704-718.

 TABLES

526 **Table 1. Comparison of mitochondrial respiration metrics in whole blood and in isolated**

527 **blood cells from the same blood volume.** The table shows model estimates, test statistics,

528 degrees of freedom and resultant *P*-values, when assaying mitochondrial respiration in zebra

529 finches using either 100 µl whole blood or all cells from 100 µl blood. All samples were run

530 in duplicate. *R* refers to intraclass repeatability of these samples and is presented with 95%

531 confidence bands inferred from 1000 bootstrap iterations. Significant (i.e., *P* < 0.05) effects

532 appear in bold font. Abbreviations: CI = Confidence Interval; d.f. = Degrees of Freedom; *LRT*

533 = Likelihood Ratio Test statistic; *R* = Intraclass Repeatability; s.e.m. = Standard Error of

534 Mean.

Model and parameter	Estimate (s.e.m.)	d.f.	*LRT*	*P*	Fold difference	*R* (95% CI)
ROUTINE (pmol $O_2 \times s^{-1} \times$ µl blood^{-1}) Sample		1	0.4	0.5328	0.98	
Blood cells	0.397 (0.035)					0.911 (0.606-0.982)
Whole blood	0.391 (0.035)					0.985 (0.921-0.997)
OXPHOS (pmol $O_2 \times s^{-1} \times$ µl blood^{-1}) Sample		1	8.0	**0.0046**	1.17	
Blood cells	0.225 (0.023)					0.808 (0.269-0.957)
Whole blood	0.264 (0.023)					0.948 (0.741-0.989)
ETS (pmol $O_2 \times s^{-1} \times$ µl blood^{-1}) Sample		1	19.1	**< 0.0001**	1.26	
Blood cells	0.648 (0.058)					0.886 (0.507-0.977)
Whole blood	0.815 (0.058)					0.965 (0.775-0.993)
LEAK (pmol $O_2 \times s^{-1} \times$ µl blood^{-1}) Sample		1	12.7	**< 0.0001**	0.74	
Blood cells	0.172 (0.019)					0.908 (0.548-0.981)
Whole blood	0.128 (0.019)					0.967 (0.827-0.992)
E-R control efficiency Sample		1	19.0	**< 0.0001**	0.77	
Blood cells	0.365 (0.049)					0.968 (0.806-0.994)
Whole blood	0.511 (0.049)					0.982 (0.893-0.996)
R-L control efficiency Sample		1	12.2	**0.0004**	0.71	
Blood cells	0.559 (0.034)					0.849 (0.325-0.972)
Whole blood	0.685 (0.034)					0.933 (0.691-0.988)
E-L coupling efficiency Sample		1	26.4	**< 0.0001**	0.55	
Blood cells	0.726 (0.023)					0.790 (0.202-0.950)
Whole blood	0.848					0.949 (0.711-0.990)

	(0.023)					

Model and parameter	Estimate (s.e.m.)	d.f.	*LRT*	*P*
ROUTINE (pmol $O_2 \times s^{-1} \times µl\ blood^{-1}$) Blood volume		1	27.5	**< 0.0001**
10 µl [A]	0.669 (0.037)			
25 µl [A]	0.656 (0.034)			
50 µl [A]	0.607 (0.034)			
100 µl [B]	0.411 (0.034)			
OXPHOS (pmol $O_2 \times s^{-1} \times µl\ blood^{-1}$) Blood volume		1	22.2	**< 0.0001**
10 µl [AB]	0.328 (0.029)			
25 µl [C]*	0.437 (0.026)			
50 µl [BC]	0.416 (0.026)			
100 µl [A]	0.259 (0.026)			
ETS (pmol $O_2 \times s^{-1} \times µl\ blood^{-1}$) Blood volume		1	13.9	**0.0031**
10 µl [A]	0.784 (0.060)			
25 µl [AB]	0.835 (0.055)			
50 µl [B]	1.005 (0.055)			
100 µl [A]	0.715 (0.055)			
LEAK (pmol $O_2 \times s^{-1} \times µl\ blood^{-1}$) Blood volume		1	39.9	**< 0.0001**
10 µl [A]	0.345 (0.018)			
25 µl [B]	0.219 (0.017)			
50 µl [BC]	0.191 (0.017)			
100 µl [C]	0.152 (0.017)			
E-R control efficiency Blood volume		1	29.6	**< 0.0001**
10 µl [A]	0.159 (0.051)			
25 µl [A]	0.191 (0.049)			
50 µl [B]	0.389 (0.049)			
100 µl [B]	0.421 (0.049)			

		1	28.5	**< 0.0001**

R-L control efficiency
Blood volume
    10 µl [A]
 	
    25 µl [B]
 	
    50 µl [B]
 	
    100 µl [B]
	
	0.486
	(0.025)
	0.664
	(0.022)
	0.685
	(0.022)
	0.625
	(0.022)
		1	28.5	**< 0.0001**

E-L coupling efficiency
Blood volume
    10 µl [A]
 	
    25 µl [B]
 	
    50 µl [C]
 	
    100 µl [C]
	
	0.566
	(0.023)
	0.726
	(0.022)
	0.803
	(0.022)
	0.785
	(0.022)
		1	49.7	**< 0.0001**

Table 3. Effects of storage time on mitochondrial respiration metrics in zebra finch whole blood. The table shows mean (± standard error; s.e.m.) trait values at 0 h and after 24 h storage at 5°C, as well as test statistics, degrees of freedom and corresponding P-values for paired t-tests of the effects of storage time on respiration traits. Significant (i.e., $P < 0.05$) effects appear in bold font. Abbreviations: d.f. = Degrees of Freedom; s.e.m. = Standard Error of Mean; t = t-test statistic.

Model and parameter	Mean (s.e.m.)	d.f.	t	P
ROUTINE (pmol $O_2 \times s^{-1} \times \mu l$ blood^{-1}) Sample 　0 h 　24 h	 0.673 (0.047) 0.731 (0.047)	5	-1.40	0.2214
OXPHOS (pmol $O_2 \times s^{-1} \times \mu l$ blood^{-1}) Sample 　0 h 　24 h	 0.498 (0.046) 0.542 (0.035)	5	-1.47	0.2015
ETS (pmol $O_2 \times s^{-1} \times \mu l$ blood^{-1}) Sample 　0 h 　24 h	 1.169 (0.088) 1.011 (0.111)	5	3.23	**0.0233**
LEAK (pmol $O_2 \times s^{-1} \times \mu l$ blood^{-1}) Sample 　0 h 　24 h	 0.175 (0.008) 0.188 (0.019)	5	-0.94	0.3905
E-R control efficiency Sample 　0 h 　24 h	 0.404 (0.066) 0.255 (0.049)	5	4.16	**0.0089**
R-L control efficiency Sample 　0 h 　24 h	 0.736 (0.019) 0.744 (0.016)	5	-0.74	0.4904
E-L coupling efficiency Sample 　0 h 　24 h	 0.845 (0.017) 0.809 (0.016)	5	3.29	**0.0217**

Figure 1. Difference in mitochondrial respiration traits as assayed in either isolated blood cells or whole blood from zebra finches. A) baseline ("ROUTINE") oxygen consumption in the endogenous cellular state; B) oxygen consumption devoted to oxidative phosphorylation alone, where ATP is produced ("OXPHOS); C) maximum working capacity of the electron transport system when uncoupled from ATP production ("ETS"); D) the part of ROUTINE used to offset the leak of protons across the inner mitochondrial membrane ("LEAK). Coloured plotting symbols show raw data means ± 1 standard error. Gray points and lines show individual responses. Significances are reported in Table 1.

Figure 2. Flux Control Ratios for mitochondrial respiration traits measured in isolated

blood cells or whole blood. A) The proportion of maximum working capacity (ETS)

remaining during endogenous respiration (ROUTINE); B) Efficiency of ATP-producing

respiration (OXPHOS) under endogenous cellular conditions; C) Coupling efficiency of

electron transport in a FCCP-stimulated cellular state. Coloured plotting symbols show raw

data means ± 1 standard error. Gray points and lines show individual responses. Details on

calculation of FCRs are provided in the main text, and significances appear in Table 1.

Figure 3. Effects of sample volume when assaying mitochondrial function in zebra finch whole blood. The panels show mitochondrial respiration traits (A) and Flux Control Ratios (FCR) (B) in relation to blood sample volume. Derivation of traits and FCRs are detailed in the main text and in the legend to Fig. 2. Coloured plotting symbols show raw data means ± 1 standard error, and coloured lines show the averaged response. Gray lines show individual responses. Significances are reported in Table 2.

Figure 4. Effects of storage time when assaying mitochondrial function in zebra finch

whole blood. The panels show mitochondrial respiration traits (A) and Flux Control Ratios

(FCR) (B) in relation to blood sample volume. All individuals were measured at 0 h and after

storage without agitation for 24 h at 5°C. Derivation of traits and FCRs are detailed in the

main text and in the legend to Fig. 2. Coloured plotting symbols show raw data means ± 1

standard error, and coloured lines show the averaged response. Gray lines show individual

responses. Significances are reported in Table 3.