Plasticity of mitochondrial function safeguards phosphorylating respiration during *in vitro*

simulation of rest-phase hypothermia

Authors: Carmen C. García-Díaz[1], Imen Chamkha[2], Eskil Elmér[2], Andreas Nord[1,*]

[1] Lund University, Department of Biology, Section for Evolutionary Ecology, Sölvegatan 37,

SE-223 62 Lund, Sweden

[2] Lund University, Department of Clinical Sciences, Mitochondrial Medicine, Sölvegatan 17,

SE-221 84, Lund, Sweden

* Author for correspondence: AN, e-mail: andreas.nord@biol.lu.se

ORCID numbers:

CCGD: 0000-0002-7827-8104

IM: 0000-0002-6919-585X

EE: 0000-0001-7901-1826

AN: 0000-0001-6170-689X

AUTHOR CONTRIBUTIONS: AN conceived the idea and designed the protocol with input

from EE. AN performed the fieldwork, and CCGD, IM and AN performed the laboratory

work. All authors interpreted the results. CCGD and AN analysed the data and wrote the first

draft. CCGD and AN produced the graphic material. All authors critically revised and edited

the manuscript. AN procured funding.

ABSTRACT

Many animals downregulate body temperature to save energy when resting (rest-phase

hypothermia). Small birds that winter at high latitude have comparatively limited capacity for

hypothermia and so pay large energy costs for thermoregulation during cold nights. Available

evidence suggests this process is fuelled by adenosine triphosphate (ATP)-dependent

mechanisms. Most ATP is produced by oxidative phosphorylation in the mitochondria, but

mitochondrial respiration can be lower during hypothermia because of the temperature-

dependence of biological processes. This can create conflict between increased organismal

ATP demand and a lower mitochondrial capacity to provide it. We studied this in blood cell

mitochondria of wild great tits (*Parus major*) by simulating rest-phase hypothermia *via* a 6°C

reduction in assay temperature *in vitro*. The birds had spent the night preceding the

experiment in thermoneutrality or in temperatures representing mild or very cold winter

nights. Night temperature did not affect mitochondrial respiration. Across treatments,

endogenous respiration was 14% lower in hypothermia. This did not reflect general thermal

suppression because phosphorylating respiration was unaffected by thermal state. Instead,

hypothermia was associated with a threefold reduction of leak respiration, from 17% in

normothermia to 4% in hypothermia. Thus, coupling of total respiration to ATP production

was 96% in hypothermia, compared to 83% in normothermia. Our study shows that thermal

insensitivity of phosphorylation combined with short-term plasticity of leak respiration may

safeguard ATP production when endogenous respiration is suppressed. This casts new light

on the process by which small birds endure harsh winter cold and warrants future tests across

tissues *in vivo*.

Keywords: blood cells, body temperature, cell respiration, great tit, mitochondria,

thermoregulation

INTRODUCTION

Endothermic animals, such as mammals and birds, use considerable amounts of energy to regulate a high and relatively stable body temperature across a wide range of air temperatures. The associated energy cost of thermoregulation can sometimes become prohibitively high, for example when food is scarce or when weather conditions deteriorate. Many small to medium-sized endotherms use rest-phase hypothermia, a regulated reduction of resting body temperature below normal levels, to mitigate such energetic challenges. This may reduce metabolic demands by 50% or more [1]. It follows that most animals using rest-phase hypothermia have high metabolic intensity, lose heat quickly and rely on high-energy or ephemeral food sources [2]. This fits the description of "the little bird in winter" [3], i.e., small (< 20 g) songbirds that are year-round residents at high latitudes where they routinely meet environmental temperatures 50-60 °C below body temperature during the cold season. Yet, most birds in northern forests, such as tits and chickadees, typically show daily variation in body temperature within 5 °C [e.g. 4, 5, 6] and generally do not exceed 7-8 °C even in severe cold [7-9, but see 10 for an exception]. Theoretical models predict that even such moderate body temperature reduction can carry survival benefits [11, 12]. However, the modest extent of hypothermia means that the little bird in winter can never forego a significant increase in energetically costly heat production to counter winter cold, even when maximally hypothermic [e.g. 13].

Most of the energy used by the working body is in the form of adenosine triphosphate (ATP) that is produced mainly by oxidative phosphorylation in the mitochondria [14]. Since more energy is needed to stay warm in winter, it is not surprising that many endotherms upregulate mitochondrial content and/or respiration rate in key metabolic tissues when cold exposed, such as in the wintertime [15-18, e.g. 19]. However, little is known about mitochondrial

adjustment in response to shorter-term variation in air temperature, for example during a

sudden drop in night-time temperature coincident with clearing of an overcast sky. Moreover,

a potential problem for a hypothermic animal is that the activity of the respiratory chain could

be slower in a colder body [e.g. 20], causing a conflict between increased organismal demand

for fuel but reduced mitochondrial capacity to provide it. However, not all the proton motive

force is used to produce ATP since some protons backflow into the mitochondrial matrix

bypassing ATP synthase. This is mediated by three principal mechanisms. Physiological

properties maintain a basal level of proton conductance of the mitochondrial membrane that

can be further up- or downregulated under the influence of uncoupling proteins [21]. A

functionally similar phenomenon occurs when protons spontaneously "slip" back through the

proton pumps of the electron transport chain, though the contribution of slippage to total

uncoupling is probably low [reviewed by 21]. Compensation for these processes make up

what is known as leak respiration. This constitutes some 20-25% of resting metabolic rate in

endotherms [22, 23], meaning there is scope to adjust the coupling of electron transport to

ATP production according to the animal's need. For example, increased mitochondrial

uncoupling in brown adipose tissue (BAT) is the basis for non-shivering heat production in

mammals [14]. In contrast, birds lack BAT and with few exceptions are thought to produce

heat using ATP-dependent shivering in skeletal muscles [24, 25; but see 26-27 for

exceptions]. Thus, in birds it can be speculated that acute cold exposure could be associated

with improved coupling, to make ATP production more effective for a given respiration rate,

which has been suggested for other energy demanding processes such as fasting [e.g. 28-31].

This could secure adequate fuel delivery even when cell respiration is reduced, such as when

heat production must remain high, but rest-phase hypothermia causes the mitochondria to

work at slower pace. There is support for this compensation hypothesis from studies on

ectotherms [32-34; but see 35, 36 for exceptions]. Similar studies on endothermic mammals

suggest that while phosphorylating respiration is often suppressed in hypometabolic animals,

non-phosphorylating (i.e., leak) respiration is sometimes reduced, sometimes unaltered, and

sometimes increased [reviewed by 37].

Previous endotherm studies have largely been performed in mammals during hibernation or in

deep torpor; thermal states characterised by substantial metabolic downregulation [1] beyond

what is feasible in most birds [but see 38]. Thus, previous data are not easily extrapolated to

the problems faced by the little bird in winter that must increase energy expenditure sharply to

counter cold temperature even when hypothermic and must do so using ATP-dependent

thermogenesis. Hence, we studied mitochondrial responses to both cold environments and

rest-phase hypothermia in the great tit (*Parus major* L.) – a small (16-20 g) songbird that

winters at high latitude and show pronounced diel variation in body temperature [4, 39]. We

exploited the fact that birds contain functional mitochondria in their red blood cells. Previous

studies have found that avian blood cell mitochondria respond in the short-term to variation in

energetic demands [40-42], over longer time periods in line with age-related metabolic

senescence [43] and according to environmental context [18]. There is also evidence

suggesting blood cell respiration is correlated to that in in key tissues for heat production

(skeletal muscle) and energy management (heart, kidney, brain) in mammals and birds [44-

46], all of which show pronounced seasonal acclimatisation to winter conditions in small,

northern birds [47]. This emphasises the need for further studies exploring the utility of blood

cell respirometry as a minimally invasive method for assessing mitochondrial responses to

environmental stressors. On this basis, we, firstly, tested if the immediate thermal

environment of the bird impacted mitochondrial function by manipulating the night preceding

measurements to be representative of mild or very cold winter temperature, or

thermoneutrality. Then, we performed an *in vitro* simulation of rest-phase hypothermia, to test

if lower body temperature caused a reduction in mitochondrial respiration and if this was

associated with any changes in the efficiency of ATP-production. We expected that low night-

time temperature would be associated with increased mitochondrial respiration rate to

facilitate ATP-dependent heat production. In contrast, we predicted that lower assay

temperature would cause thermal suppression of respiration traits. However, if better coupled

mitochondria serve as an adaptation to safeguard ATP production when cold ambient

temperature increases organismal-level demand for ATP, or when lower body temperature

reduces overall energy expenditure of the cell, we expected tighter coupling of electron

transport to ATP production both in response to lower air temperature at night and during *in*

vitro hypothermia.

MATERIALS AND METHODS

The experimental design and timeline are outlined in Fig. 1.

Study site and animals

Great tits were caught at two sites in the vicinity of Lund in southernmost Sweden – Räften (n

= 37) and Vomb (n = 21) – between January and February 2021. Räften (55° 43'N, 13° 17'E")

is an oak (*Quercus robur*) dominated 32 ha woodlot surrounded by arable fields. The Vomb

site (55° 39'N, 13° 33'E) is part of a pine (*Pinus sylvestris*) plantation with a rich deciduous

understory composed mainly of sallow (*Salix* spp.), birch (*Betula pendula*) and rowan (*Sorbus*

aucuparia). Birds in Räften were caught in mist nets close to feeders baited with sunflower

seeds that were erected 1.5 m above ground in January 2021. The birds in Vomb were caught

shortly before sunrise when roosting in nest boxes. Four to eight birds were caught each day

of the study and were brought to Lund University within 2 h of capture, where they were

alternately allocated to temperature treatment groups in batches of four. We housed the birds

in cages measuring $77 \times 44 \times 77$ cm (length $\times$ width $\times$ height) at 5 °C under simulated natural

photoperiod and were provided sunflower seeds, peanuts, and water *ad libitum*. One-two days

after capture the birds were implanted with a temperature-sensitive PIT tag (BioTherm13,

Biomark, Boise, ID, USA) into the intraperitoneal cavity under local anaesthesia (5%

lidocaine) as part of a different study. Eight to 34 h later, the birds were put in a 1 L

hermetically sealed glass container ventilated with dry air at 400-500 ml / min (recorded using

a FB-8 mass flow meter, Sable Systems, Las Vegas, NV, USA) for measurement of resting

metabolic rate (RMR; as oxygen consumption recorded in carbon dioxide-free air using a FC-

10 oxygen analyser, Sable Systems) and body temperature (recorded using a custom-built

multiplexed antennae system from BioMark) in a climate chamber (Weiss Umwelttechnik

C180 [Reiskirchen, Germany]) at thermoneutrality (25 °C; n = 20) [13], during a simulated

mild winter night (5 °C: n = 19) and during a simulated very cold winter night (-15 °C; n =

19) for 13-15 h. Four birds were measured each night. Mean RMR was 0.36 ± 0.01 W (mean

$\pm$ s.e.m.), 0.51 ± 0.01 W and 0.68 ± 0.02 W in +25, +5 and -15 °C, respectively. Mean nightly

body temperature was 37.9 ± 0.3 °C (range of means: 35.2 to 40.1 °C; minimum: 33.8 °C),

39.3 ± 0.1°C (range of means: 37.6 to 40.2 °C; minimum: 37.0 °C), and 39.2 ± 0.1 °C (range

of means: 38.0 to 40.3 °C; minimum: 37.6 °C), in the cold, mild, and thermoneutral night

conditions.

Air temperature in Lund (10-40 km from the study sites) ranged -14 °C to 7 °C during the

study period and photoperiod ranged 8:20 h to 10:20 h.

Blood sampling and mitochondrial measurement in whole blood

In the morning after the metabolic measurements, a 100-150 µl blood sample ≤ 10% of

estimated total blood volume was obtained from the jugular vein using a heparinized syringe

within 5 min of removing the bird from the metabolic chamber. The sample was stored at 10-

12 °C in 2 mL EDTA (ethylenediaminetetraacetic acid) tubes (BD Vacutainer®, Franklin

Lakes, NJ, USA) without agitation until analysed 0.5-2.5 h later. All birds were released at the

site of capture within 2 h of blood sampling.

We measured mitochondrial respiration in whole blood using an Oxygraph O2k high

resolution respirometer (Oroboros Instruments, Innsbruck, Austria), following the protocol

described in Nord, Chamkha & Elmér [48]. To test how rest-phase hypothermia affected

mitochondrial function, we ran duplicate samples from the same individual simultaneously in

each of two thermal states: a representative normothermic daytime bird body temperature (41

°C; henceforth "normothermia") and a hypothermic night-time body temperature in the lower

range of the measured natural variation (35 °C; henceforth "hypothermia") [49] using two

respirometers. These were calibrated at assay temperature (i.e., 35 °C or 41 °C) daily, and we

changed instrument/assay temperature combinations between days to avoid confounding

experimental results by any variation pertaining to the specific instruments. The main purpose

of our study was to demonstrate the presence of a putative mitochondrial response to acute

body temperature change. To achieve this goal, we purposely selected a low but biologically

relevant hypothermic assay temperature, under the assumption that any phenotypic response

would be linear over the range of *in vitro* temperatures considered.

The blood samples were allowed 5-10 min on a rotating mixer at room temperature before the

start of the assay. Then, we added 50 µL sample to 1.95 mL MiR05 respiration medium (0.5

mM EGTA, 3 mM MgCl$_2$, 60 mM K-lactobionate, 20 mM taurine, 10 mM KH$_2$PO$_4$, 20 mM

Hepes, 110 mM sucrose, free fatty acid bovine serum albumin (1 g L^{-1}), pH 7.1) that was

contained within the respirometer chambers and so was pre-equilibrated at assay temperature.

The final volume contained in the respirometer chamber was, thus, 2 mL.

Mitochondrial respiration rate was measured as the rate of decline in O$_2$ concentration in the

chamber. Once the O$_2$ consumption had stabilized upon closing the chamber, baseline O$_2$

consumption on endogenous substrates ('ROUTINE') was recorded for 2-3 min. Then, we

assessed mitochondrial function by sequential addition of three mitochondrial agonists and

antagonists. Firstly, we added the ATP-synthase inhibitor oligomycin at a final concentration

of 5 µmol L^{-1}, which blocks the process of oxidative phosphorylation ('OXPHOS'). Thus, the

remaining O$_2$ after oligomycin addition is attributed to proton leak ('LEAK'). Next, we added

the mitochondrial uncoupler FCCP (carbonyl cyanide-p-trifluoro-methoxyphenyl-hydrazone),

which abolishes the proton gradient across the inner mitochondrial membrane, causing the

electron transport system to work at maximum capacity to restore the gradient ('ETS').

Because an excessive concentration of FCCP inhibits mitochondrial respiration and can lead

to acidification of the cell cytoplasm [14], we first added 1.5 µL of 1 mmol L^{-1} FCCP and

then titrated FCCP in 0.5 µL increments until maximum O$_2$ consumption was reached. The

final concentration ranged 1-1.75 µmol L^{-1}. Finally, we added 5 µmol L^{-1} of antimycin A,

which blocks Complex III of the electron transport chain. Therefore, the remaining O$_2$

consumption after antimycin A addition is due to non-mitochondrial oxygen utilisation.

Data handling and statistical analyses

We excluded five of the 57 duplicate samples measured at 41 °C since they were not inhibited

by the addition of oligomycin. Thus, the final data set consisted of 57 observations at 35 °C

and 52 observations at 41 °C. We then calculated mitochondrial respiration states and flux

control ratios (FCR; i.e., ratios between the respiratory states) following Gnaiger (2020) [50].

We considered three FCRs: 1) E-R control efficiency (1 - ROUTINE / ETS), which measures

how much excess capacity there is in the electron transport system until maximum respiration

is reached); 2) R-L control efficiency ((ROUTINE - LEAK) / ROUTINE), which measures

the proportion of endogenous respiration that is channeled towards oxidative phosphorylation;

3) E-L coupling effiency (1 - LEAK / ETS) which is indicative of mitochondrial efficiency in

the context of coupling between electron transport and ATP production in a stimulated

cellular state. Derivation and definition of all respiration traits and FCRs are detailed in Table

1.

All statistical analyses were performed using R version 4.1.2. To test how a sudden change in

night temperature affected mitochondrial respiration the morning after, we used linear models

(lm in R base) to test how each respiration state and FCR assayed at normothermic

temperature (i.e., 41°C) was affected by preceding night temperature. We used linear mixed

models (lmer in the lme4 package) [51] to test how respiration traits and FCRs were affected

by thermal state (i.e., hypo- or normothermia) using bird ID as a random factor to account for

repeated measurements. To test if the phenotypic response to a change in thermal state

differed depending on the depth of hypothermia during the preceding night, we included mean

nightly body temperature and body temperature × thermal state in these models. Significances

were assessed using F-tests with degrees of freedom based on the Kenward-Roger

approximation, implemented using the KRmodcomp function in the pbkrtest package [52].

The significance of the random term (i.e., bird ID) in final models was assessed using log

likelihood tests using the drop1 function in base R. The interaction was removed when non-

significant ($P > 0.05$), but all main effects were retained. Predicted values and standard errors

(s.e.m.) for the lmer models were calculated using the emmeans function in the emmeans

package [53]. For the tests of night temperature effects, estimates and their s.e.m. were taken

from the linear model output.

RESULTS

A representative normothermic experiment is presented in Fig. S1. All details of the statistical

tests pertaining to effects of night temperature are presented in Table S1, and those pertaining

to the effects of thermal state and body temperature are presented in Table S2.

Effects of preceding night temperature

All but one of the mitochondrial respiration metrics were unaffected by temperature of the

preceding night (all $P > 0.16$; Table S1). There was a tendency for R-L control efficiency,

which measures how much of the ROUTINE respiration that is channelled towards ATP

production, to vary with night temperature ($P = 0.063$). Specifically, 86% of endogenous

respiration was used to drive oxidative phosphorylation after a night in -15°C, which was

higher than after a night at +5°C (81%) but similar to a night in +25°C (83%) (Table S1).

Effects of thermal state and nightly body temperature

There were profound changes when assay temperature changed from a normothermic to a

hypothermic thermal state. ROUTINE respiration was 14 % higher in normothermia

(0.567 ± 0.019 pmol O_2 s^{-1} µl^{-1}) than in hypothermia (0.499 ± 0.019 pmol O_2 s^{-1} µl^{-1}) ($P <$

0.0001) (Fig. 2A), i.e., endogenous respiration was lower during *in vitro* hypothermia.

However, neither OXPHOS nor ETS were affected by thermal state ($P = 0.7$ and 0.6,

respectively) (Figs. 2B-C). Instead, the drop in ROUTINE was mediated by a pronounced

reduction in LEAK respiration ($P < 0.0001$), which was more than three times higher during

normothermia (0.096 ± 0.004 pmol O_2 s^{-1} μl^{-1}) compared to hypothermia (0.023 ± 0.004 pmol

O_2 s^{-1} μl^{-1}) (Fig. 2D).

E-R control efficiency was lower during hypothermia ($P < 0.0001$), meaning a hypothermic

bird had more mitochondrial reserve capacity compared to a normothermic bird (0.505 ± 0.012

and 0.434 ± 0.012, of 1, respectively) (Fig. 3A). The efficiency of oxidative phosphorylation

was higher in the hypothermic state (i.e., R-L control efficiency increased) such that nearly all

(96%) of endogenous respiration was used to produce ATP in hypothermia, compared to 83%

in normothermia (Fig. 3B). In other words, LEAK accounted for nearly 17% of endogenous

respiration during normothermia, but in hypothermia LEAK contributed only 4% to

ROUTINE. E-L coupling efficiency (i.e., coupling of the ETS when maximal respiration was

stimulated using FCCP) increased from 91% (0.907 ± 0.004) in normothermia to 98%

(0.978 ± 0.004) in hypothermia (Fig. 3C). Thus, the electron transport chain was tighter in the

hypothermic state.

Most mitochondrial respiration traits were unaffected by nightly mean body temperature, both

as main effects and in interaction with thermal state (Table S2; Fig. 4). However, E-R control

efficiency increased with increasing body temperature (by 0.041 ± 0.010 per °C) across thermal

states (Fig .4), suggesting that birds that were more hypothermic during the preceding night

had mitochondria that respired at a greater proportion of maximum. There was also a thermal-

state-dependent relationship between R-L control and body temperature, whereby oxidative

efficiency decreased by 0.012 ± 0.008 per °C in the normothermic state but was unaffected by

body temperature in the hypothermic state. However, when tested in assay temperature-

specific models, the linear regression between R-L control and mean night-time body

temperature was not significant (hypothermia: $P = 0.39$; normothermia: $P = 0.13$).

DISCUSSION

We tested whether plasticity of mitochondrial function could compensate for any reduced cellular respiration rate during rest-phase hypothermia in great tit blood cells. As predicted, endogenous respiration (ROUTINE) was lower in the hypothermic thermal state (Fig. 2A). Surprisingly, this was not a reflection of a general downregulation of cellular respiration during hypothermia, because there was no temperature-dependence of phosphorylating respiration (OXPHOS; in which ATP is produced) (Fig. 2B). Instead, lower ROUTINE was primarily caused by a threefold reduction in LEAK (Fig. 2D). Lack of temperature effects on OXPHOS contrasts previous studies on mammals, where a lowering of assay temperature by 10-12°C to simulate a deep hypothermic state was associated with a 50-70% decrease of both phosphorylating and non-phosphorylating respiration [54-55]. It has been proposed that, in hibernating mammals, active suppression of phosphorylating respiration only becomes noticeable at body temperatures < 30°C [37], which could explain part of the thermal insensitivity of OXPHOS recorded here for a 6°C temperature reduction. If this applies also to birds, a logical conclusion is that passive thermal effects do not influence phosphorylating respiration strongly over the range of temperatures in our study. By contrast, leak respiration showed thermal sensitivity at, or beyond, that recorded in other studies [54-55], but our experimental design does not permit analysis of the contribution of passive thermal and actively regulated effects to this response. Mechanisms aside, the combined actions of thermal insensitivity of OXPHOS and pronounced thermal sensitivity of LEAK meant that the hypothermic thermal state was associated with significant increases in both phosphorylating efficiency (i.e., R-L control efficiency was higher; Fig. 3B) and coupling efficiency of electron transport when the cells were stimulated to work at maximal rate (i.e., E-L coupling efficiency was higher; Fig. 3C), broadly in keeping with results from mammal studies that

manipulated assay temperature [54-55]. In fact, nearly all respiration (96%) was directed

towards ATP production during hypothermia, and the electron transport system was almost

perfectly coupled (0.98%) in this thermal state. It is also interesting to note that we found no

effect of simulated hypothermia on maximally stimulated mitochondrial respiration (i.e., ETS;

Fig. 2C), suggesting lack of noticeable thermal sensitivity of the electron transport system. As

a result, hypothermic cells had higher reserve capacity (Fig. 3A), which could have functional

significance in face of a challenge to energy metabolism. Collectively, our study indicates

that plasticity of mitochondrial leak respiration combined with thermal insensitivity of

phosphorylating and maximal respiration may allow the little bird in winter to secure

sufficient ATP production to meet thermoregulatory demands during rest-phase hypothermia.

This notion now needs confirmation in other tissues and by *in vivo* studies.

Reduced LEAK and improved phosphorylating and coupling efficiency during hypothermia is

in keeping with the hypothesis of biochemical compensation for thermal suppression of cell

respiration at low body temperature. However, tighter coupling can potentially result in

increased production of reactive oxygen species (ROS) as a by-product of electron transport

[21, 56], provided that the resultant increase in protonmotive force is not dissipated through

ATP synthase. Unless quenched, ROS cause oxidative stress with resultant macromolecular

damage [e.g. 57], which is related to many medical disorders [58]. It is interesting to speculate

that a hypothermic bird might have to accept an oxidative cost for the improved oxidative and

coupling efficiencies. In this context, our study could provide proximate insights into why

oxidative stress has been found to be higher in birds that are more hypothermic at night [59].

However, the role of mitochondria as sources of cellular ROS has been contended [60-62] and

empirical evidence suggests that the relationship between body temperature, ATP production

and ROS generation is not straightforward [31, 63, 64]. In line with this, even though

(pharmacological) uncoupling protects birds from cold-induced oxidative stress [65], the

evidence for clear fitness effects thereof is equivocal [66, 67].

We found no strong effects of temperature of the preceding night on mitochondrial respiration

traits and flux control ratios, though phosphorylating capacity was the highest after the coldest

night-time temperature (Table S1). This contrasts what has been recorded as part of seasonal

acclimatisation of blood cell respiration in the same model system [18] and in response to

cold exposure in other tissue types [15-17]. Responses in other studies were likely driven by

seasonal upregulation of mitochondrial content (e.g. [18]) and so is possible that the period of

exposure (one night) and acute nature of the air temperature manipulation was too short or too

stochastic to serve as a reliable environmental cue prompting a change to cellular respiratory

capacity. It is also possible that seasonal changes in mitochondrial content or respiration rate

in the studies above were not causally related to cold temperature. In line with this, cold

acclimation did not impact the respiration of liver mitochondria in the lesser hedgehog tenrec

(*Echinops telfairi*) [54]. Even so, the difference in short- and long-term responses of

mitochondrial respiration coincident with colder environmental temperature suggests it would

be interesting to study the temporal resolution of mitochondrial thermal acclimatisation in

some detail. For example, it is possible that, in the short term, mitochondrial function

responds primarily to a change of tissue temperature in endothermic animals. This could be

beneficial because endotherm body temperature is determined by the balance between heat

production and heat loss rates and so may be partly or fully independent of air temperature.

On average, there were few effects of night-time body temperature on mitochondrial

respiration traits the day after. Similar results were obtained for liver mitochondria in the

golden spiny mouse (*Acomys russatus*) measured at a constant assay temperature [68]. Thus,

for many of the mitochondrial traits measured here, between-individual variation (which

explained 30-70% of total variance in the statistical models; Table S2) did not constrain

thermal plasticity. One notable exception was the mitochondrial reserve capacity (i.e., E-R

control ratio), which was higher in the hypothermic state (Fig. 3A) but showed a negative

relationship with overnight body temperature (Fig. 4). Thus, *in vivo*, more hypothermic birds

had mitochondria with *lower* surplus capacity. This invites the speculation that the scope for

plastic upregulation of mitochondrial work rate, e.g. in response to sudden environmental

change, may be limited by individual differences in the underlying mitochondrial phenotype.

It remains to be tested if such constraints on plasticity hinders adaptation to the environment

over short- and long temporal scales. Finally, it is interesting to note that the reduction in

surplus mitochondrial capacity with decreasing body temperature is well in line with the

observation that more hypothermic birds are typically of lower phenotypic quality [e.g., 69-

70]. This raises the question of whether rest-phase hypothermia in the little bird in winter is an

adaptive response to save energy or if it instead reflects a phenotypic expression enforced by

the inability to stay warm.

This study provides new perspectives to our understanding of how cellular and whole animal

metabolism interact when the little bird in winter adapts to low temperatures. We conclude

that *in vitro* hypothermia was associated with an immediate reduction in proton leak and

resultant increased in coupling and oxidative efficiency, which can explain why blood cell

mitochondria could maintain phosphorylating respiration despite acute temperature reduction.

Nor did low assay temperature suppress maximum working capacity of the electron transport

system, leading to increased mitochondrial respiratory scope during hypothermia (i.e., E-R

control efficiency was higher, Fig. 3A). However, it is possible that mitochondrial plasticity

during rest-phase hypothermia may come at the cost of increased oxidative stress [cf. 59].

Future work should explore this hypothesis by direct measurement of ATP and ROS

production, and membrane potential, to conclude on the consequences of changes to the

efficiency of phosphorylating respiration for redox balance in normo- and hypothermic birds.

Several endotherm studies show that mitochondrial function in blood cells correlates to that in

tissues and organs with clearly defined roles in heat production and seasonal acclimatisation

(see above), suggesting data presented here may be representative of cellular responses to

hypothermia at organismal levels. More broadly, our work aligns with the body of evidence

highlighting the utility of blood cell respirometry to gain minimally invasive insight into

functional ecology, evolutionary physiology, and gerontology in birds [40, 41, 43,71]. Even

so, it is important that future studies formally address the uniformity of mitochondrial

plasticity across tissues and organs. Likewise, future investigations should aim to increasingly

integrate tissue-based studies with an *in vivo* approach to understand mitochondrial regulation

during hypothermia at the organismal level. More broadly, it would be relevant to increase

taxonomic and phenotypic coverage of these studies, spanning cold-hardy, non-hypothermic,

birds [e.g. 72] to those that routinely use shallow [e.g. 73] or deep torpor [74-76].

ACKNOWLEDGEMENTS

The authors thank Stefan Nord for managing the Räften feeding stations by which birds for

the study were caught. SN also assisted with fieldwork and provided miscellaneous technical

support during the study. Two anonymous reviewers provided critical comments that

improved the first draft.

FUNDING

This study was supported by the Birgit and Hellmuth Hertz Foundation (grant no. 2017-
39034), The Royal Physiographic Society in Lund (grant no. 2019-41011), the Swedish
Research Council (grant no. 2020-04686) and Stiftelsen Lunds Djurskyddsfond (all to AN).

ETHICS

All procedures were approved by the Malmö/Lund Animal Ethics Committee, acting under

jurisdiction of the Swedish Board of Agriculture (permit no. 9246-19). Bird ringing was

permitted by the Swedish Bird Ringing Centre at Natural History Museum of Sweden (license

number 723, to AN).

DATA AVAILABILITY

Data are deposited in Figshare: 10.6084/m9.figshare.19902277

CONFLICT OF INTEREST

The authors declare that they have no competing or financial interests.

REFERENCES

1. Ruf T., Geiser F. 2015 Daily torpor and hibernation in birds and mammals. *Biol Rev*

90(3), 891-926. (doi:10.1111/brv.12137).

2. Geiser F. 2021 *Ecological physiology of daily torpor and hibernation*. Cham,

Switzerland, Springer Nature Switzerland AG; 317 p.

3. Brodin A. 2007 Theoretical models of adaptive energy management in small wintering

birds. *Phil Trans R Soc B* **362**(1486), 1857-1871.

4. Nilsson J.F., Nilsson J.-Å., Broggi J., Watson H. 2020 Predictability of food supply

modulates nocturnal hypothermia in a small passerine. *Biol Lett* **16**(6), 20200133.

(doi:doi:10.1098/rsbl.2020.0133).

5. Nord A., Nilsson J.F., Nilsson J.-Å. 2011 Nocturnal body temperature in wintering blue

tits is affected by roost-site temperature and body reserves. *Oecologia* **167**(1), 21-25.

6. Nord A., Nilsson J.F., Sandell M.I., Nilsson J.-Å. 2009 Patterns and dynamics of rest-

phase hypothermia in wild and captive blue tits during winter. *J Comp Physiol B* **179**(6),

737-745. (doi:10.1007/s00360-009-0357-1 [doi]).

7. Reinertsen R.E., Haftorn S. 1984 The effect of short-time fasting on metabolism and

nocturnal hypothermia in the willow tit *Parus montanus*. *J Comp Physiol* **154**(1), 23-28.

8. Reinertsen R.E., Haftorn S. 1983 Nocturnal hypothermia and metabolism in the willow tit

Parus montanus at 63 degrees N. *J Comp Physiol* **151**(2), 109-118.

9. Haftorn S. 1972 Hypothermia of tits in the Arctic winter. *Orn Scand* **3**(2), 153-166.

10. Cooper S.J., Gessaman J.A. 2005 Nocturnal hypothermia in seasonally acclimatized

Mountain Chickadees and Juniper Titmice. *Condor* **107**(1), 151-155.

11. Brodin A., Nilsson J.-Å., Nord A. 2017 Adaptive temperature regulation in the little bird

in winter: predictions from a stochastic dynamic programming model. *Oecologia* **185**(1),

43-54. (doi:10.1007/s00442-017-3923-3).

12. Welton N.J., Houston A.I., Ekman J., McNamara J.M. 2002 A dynamic model of

hypothermia as an adaptive response by small birds to winter conditions. *Acta

Biotheoretica* **50**(1), 39-56.

13. Andreasson F., Nord A., Nilsson J.-Å. 2020 Age differences in night-time metabolic rate

and body temperature in a small passerine. *J Comp Physiol B* **190**(3), 349-359.

(doi:10.1007/s00360-020-01266-5).

472 14. Nicholls D.G., Ferguson S.J. 2013 *Bioenergetics*. 4th ed. Cambridge, MA, USA,

473   Academic Press.

474 15. Liknes E.T., Swanson D.L. 2011 Phenotypic flexibility in passerine birds: Seasonal

475   variation of aerobic enzyme activities in skeletal muscle. *J Therm Biol* **36**(7), 430-436.

476   (doi:http://dx.doi.org/10.1016/j.jtherbio.2011.07.011).

477 16. Wang Y., Shan S., Zhang H., Dong B., Zheng W., Liu J. 2019 Physiological and

478   biochemical thermoregulatory responses in male Chinese hwameis to seasonal

479   acclimatization: Phenotypic flexibility in a small passerine. *Zool Stud* **58**, e6-e6.

480   (doi:10.6620/ZS.2019.58-06).

481 17. Zheng W.H., Li M., Liu J.-S., Shao S.-L., Xu X.-J. 2014 Seasonal variation of metabolic

482   thermogenesis in Eurasian tree Sparrows (*Passer montanus*) over a latitudinal gradient.

483   *Physiol Biochem Zool* **87**(5), 704-718. (doi:doi:10.1086/676832).

484 18. Nord A., Metcalfe N.B., Page J.L., Huxtable A., McCafferty D.J., Dawson N.J. 2021

485   Avian red blood cell mitochondria produce more heat in winter than in autumn. *FASEB J*

486   **35**(5), e21490. (doi:https://doi.org/10.1096/fj.202100107R).

487 19. Schaeffer P.J., Villarin J.J., Lindstedt S.L. 2003 Chronic Cold Exposure Increases

488   Skeletal Muscle Oxidative Structure and Function in Monodelphis domestica, a

489   Marsupial Lacking Brown Adipose Tissue. *Physiol Biochem Zool* **76**(6), 877-887.

490   (doi:10.1086/378916).

491 20. Wright K.P., Jr., Hull J.T., Czeisler C.A. 2002 Relationship between alertness,

492   performance, and body temperature in humans. *Am J Physiol* **283**(6), R1370-1377.

493   (doi:10.1152/ajpregu.00205.2002).

494 21. Divakaruni A.S., Brand M.D. 2011 The Regulation and Physiology of Mitochondrial

495   Proton Leak. *Physiology* **26**(3), 192-205. (doi:10.1152/physiol.00046.2010).

22. Porter R.K., Brand M.D. 1995 Causes of differences in respiration rate of hepatocytes
from mammals of different body mass. *Am J Physiol* **269**(5), R1213-1224.
(doi:10.1152/ajpregu.1995.269.5.R1213).

23. Rolfe D.F., Brand M.D. 1996 Contribution of mitochondrial proton leak to skeletal
muscle respiration and to standard metabolic rate. *Am J Physiol* **271**(4), C1380-1389.
(doi:10.1152/ajpcell.1996.271.4.C1380).

24. Stager, M., Cheviron, Z.A. 2020 Is there a role for sarcolipin in avian facultative
thermogenesis in extreme cold? *Biol Lett* **16**, 20200078.

25. Hohtola, E. 2004 Shivering thermogenesis in birds and mammals. In Barnes, B.M.,
Carey, H.V. (eds) Life in the Cold – Mechanisms, adaptation and application, pp. 241-
252. Fairbanks: University of Alaska.

26. Teulier, L., Rouanet, J.L., Letexier, D., Romestaing, C., Belouze, M., Rey, B., Duchamp,
C., Roussel, D. 2010 Cold-acclimation-induced non-shivering thermogenesis in birds is
associated with upregulation of avian UCP but not with innate uncoupling or altered ATP
efficiency. *J Exp Biol* **14**, 2476-2482.

27. Nowack, J., Giroud, S., Arnold, W., Ruf, T. 2017 Muscle non-shivering thermogenesis
and its role in the evolution of endothermy. *Front Physiol* **8**, 889.

28. Roussel D., Marmillot V., Monternier P.-A., Bourguignon A., Toullec G., Romestaing C.,
Duchamp C. 2020 Skeletal muscle metabolism in sea-acclimatized king penguins. II.
Improved efficiency of mitochondrial bioenergetics. *J Exp Biol* **223**(21), jeb233684.
(doi:10.1242/jeb.233684).

29. Bourguignon A., Rameau A., Toullec G., Romestaing C., Roussel D. 2017 Increased
mitochondrial energy efficiency in skeletal muscle after long-term fasting: its relevance
to animal performance. *J Exp Biol* **220**(13), 2445-2451. (doi:10.1242/jeb.159087).

30. Monternier P.A., Marmillot V., Rouanet J.L., Roussel D. 2014 Mitochondrial phenotypic

flexibility enhances energy savings during winter fast in king penguin chicks. *J Exp Biol*

217(15), 2691-2697. (doi:10.1242/jeb.104505).

31. Boël M., Romestaing C., Duchamp C., Veyrunes F., Renaud S., Roussel D., Voituron Y.

2020 Improved mitochondrial coupling as a response to high mass-specific metabolic rate

in extremely small mammals. *J Exp Biol* **223**(5), jeb215558. (doi:10.1242/jeb.215558).

32. Glanville E.J., Seebacher F. 2006 Compensation for environmental change by

complementary shifts of thermal sensitivity and thermoregulatory behaviour in an

ectotherm. *J Exp Biol* **209**(24), 4869-4877. (doi:10.1242/jeb.02585).

33. Seebacher F. 2005 A review of thermoregulation and physiological performance in

reptiles: what is the role of phenotypic flexibility? *J Comp Physiol B* **175**(7), 453-461.

(doi:10.1007/s00360-005-0010-6).

34. Bouchard P., Guderley H. 2003 Time course of the response of mitochondria from

oxidative muscle during thermal acclimation of rainbow trout, *Oncorhynchus mykiss*. *J*

Exp Biol **206**(Pt 19), 3455-3465. (doi:10.1242/jeb.00578).

35. St-Pierre J., Brand M.D., Boutilier R.G. 2000 The effect of metabolic depression on

proton leak rate in mitochondria from hibernating frogs. *J Exp Biol* **203**(9), 1469-1476.

36. Bishop T., St-Pierre J., Brand M.D. 2002 Primary causes of decreased mitochondrial

oxygen consumption during metabolic depression in snail cells. *Am J Physiol* **282**(2),

R372-R382. (doi:10.1152/ajpregu.00401.2001).

37. Staples J.F., Brown J.C.L. 2008 Mitochondrial metabolism in hibernation and daily

torpor: a review. *J Comp Physiol B* **178**(7), 811-827.

38. McKechnie A.E., Lovegrove B.G. 2002 Avian facultative hypothermic responses: A

review. *Condor* **104**(4), 705-724.

39. Nord A., Chiriac S., Hasselquist D., Nilsson J.-Å. 2013 Endotoxin injection attenuates
rest-phase hypothermia in wintering great tits through the onset of fever. *Funct Ecol*
27(1), 236-244. (doi:10.1111/1365-2435.12003).

40. Casagrande S., Stier A., Monaghan P., Loveland J.L., Boner W., Lupi S., Trevisi R., Hau
M. 2020 Increased glucocorticoid concentrations in early life cause mitochondrial
inefficiency and short telomeres. *J Exp Biol* **223**(15), jeb222513.
(doi:10.1242/jeb.222513).

41. Stier A., Bize P., Hsu B.-Y., Ruuskanen S. 2019 Plastic but repeatable: rapid adjustments
of mitochondrial function and density during reproduction in a wild bird species. *Biol*
Lett **15**(11), 20190536. (doi:doi:10.1098/rsbl.2019.0536).

42. Udino E., George J.M., McKenzie M., Pessato A., Crino L., Buchanan K.L., Mariette
M.M. 2021 Prenatal acoustic programming of mitochondrial function for high
temperatures in an arid-adapted bird. *Proc R Soc Lond B* **288**, 20211893.
(doi:doi:10.1098/rspb.2021.1893).

43. Dawson N.J., Salmón P. 2020 Age-related increase in mitochondrial quantity may
mitigate a decline in mitochondrial quality in red blood cells from zebra finches
(*Taeniopygia guttata*). *Exp Gerontol* **133**, 110883.
(doi:https://doi.org/10.1016/j.exger.2020.110883).

44. Stier A., Romestaing C., Schull Q., Lefol E., Robin J.-P., Roussel D., Bize P. 2017 How
to measure mitochondrial function in birds using red blood cells: a case study in the king
penguin and perspectives in ecology and evolution. *Meth Ecol Evol* **8**(10), 1172-1182.
(doi:10.1111/2041-210x.12724).

45. Stier A., Monaghan P., Metcalfe N.B. 2022 Experimental demonstration of prenatal
programming of mitochondrial aerobic metabolism lasting until adulthood. *Proc R Soc*
Lond B **289**(1970), 20212679. (doi:doi:10.1098/rspb.2021.2679).

46. Karamercan M.A., Weiss S.L., Villarroel J.P., Guan Y., Werlin E., Figueredo R., Becker

L.B., Sims C. 2013 Can peripheral blood mononuclear cells be used as a proxy for

mitochondrial dysfunction in vital organs during hemorrhagic shock and resuscitation?

Shock **40**(6), 476-484. (doi:10.1097/shk.0000000000000026).

47. Petit M., Lewden A., Vézina F. 2014 How does flexibility in body composition relate to

seasonal changes in metabolic performance in a small passerine wintering at northern

latitude? *Physiol Biochem Zool***87**(4), 539-549. (doi:10.1086/676669).

48. Nord A., Chamkha I., Elmér E. 2022 A whole blood approach improves speed and

accuracy when measuring mitochondrial respiration in avian hematocytes. *bioRxiv*,

2022.2005.2025.493398. (doi:10.1101/2022.05.25.493398).

49. Prinzinger R., Pressmar A., Schleucher E. 1991 Body temperature in birds. *Comp*

Biochem Physiol A **99**(4), 499-506.

50. Gnaiger E. 2020 Mitochondrial pathways and respiratory control: An Introduction to

OXPHOS Analysis. 5th ed.

51. Bates D., Maechler M., Bolker B., Walker S. 2015 Fitting linear mixed-effects models

using lme4. *J Stat Soft* **67**, 1-48.

52. Halekoh, U., Højsgaard, S. 2014 A Kenward-Roger approximation and parametric

bootstrap methods for tests in linear mixed models – The R package pbkrtest. *J Stat Soft*

59(9), 1-32.

53. Lenth R.V. 2019 emmeans: Estimated marginal means, aka least-squares means. R

package version 1.3.3. https://CRAN.R-project.org/package=emmeans.

54. Polymeropoulos, E.T., Oelkrug, R., Jastroch, M. 2017 Mitochondrial proton leak

compensates for reduced oxidative power during frequent hypothermic events in a

protoendothermic mammal, *Echinops telfairi*. *Front Physiol* **8**, 909.

55. Dufour, S., Rousse, N., Canioni, P, Diolez, P. 1996 Top-down control analysis of temperature effect on oxidative phosphorylation. *Biochem J* **314**, 743-751.

56. Demine S., Renard P., Arnould T. 2019 Mitochondrial uncoupling: A key controller of biological processes in physiology and diseases. *Cells* **8**(8), 795. (doi:10.3390/cells8080795).

57. Brand M.D., Buckingham J.A., Esteves T.C., Green K., Lambert A.J., Miwa S., Murphy M.P., Pakay J.L., Talbot D.A., Echtay K.S. 2004 Mitochondrial superoxide and aging: uncoupling-protein activity and superoxide production. *Biochem Soc Symp* (71), 203-213. (doi:10.1042/bss0710203).

58. Grasemann H., Holguin F. 2021 Oxidative stress and obesity-related asthma. *Paed Resp Rev* **37**, 18-21. (doi:10.1016/j.prrv.2020.05.004).

59. Zagkle E., Grosiak M., Bauchinger U., Sadowska E.T. 2020 Rest-phase hypothermia reveals a link between aging and oxidative stress: A novel hypothesis. *Front Physiol* **11**, 575060. (doi:10.3389/fphys.2020.575060).

60. Zhang Y., Wong H.S. 2021 Are mitochondria the main contributor of reactive oxygen species in cells? *J Exp Biol* **224**(5), eb221606. (doi:10.1242/jeb.221606).

61. Mookerjee S.A., Divakaruni A.S., Jastroch M., Brand M.D. 2010 Mitochondrial uncoupling and lifespan. *Mech Ageing Devel* **131**(7-8), 463-472. (doi:10.1016/j.mad.2010.03.010).

62. Brown G.C., Borutaite V. 2012 There is no evidence that mitochondria are the main source of reactive oxygen species in mammalian cells. *Mitochondrion* **12**(1), 1-4. (doi:https://doi.org/10.1016/j.mito.2011.02.001).

63. Roussel D., Voituron Y. 2020 Mitochondrial costs of being hot: Effects of acute thermal change on liver bioenergetics in toads (*Bufo bufo*). *Front Physiol* **11**, 153. (doi:10.3389/fphys.2020.00153).

64. Boël M., Veyrunes F., Durieux A.-C., Freyssenet D., Voituron Y., Roussel D. 2022 Does

high mitochondrial efficiency carry an oxidative cost? The case of the African pygmy

mouse (*Mus mattheyi*). *Comp Biochem Physiol A* **264**, 111111.

(doi:https://doi.org/10.1016/j.cbpa.2021.111111).

65. Stier A., Massemin S., Criscuolo F. 2014 Chronic mitochondrial uncoupling treatment

prevents acute cold-induced oxidative stress in birds. *J Comp Physiol B* **184**(8), 1021-

1029. (doi:10.1007/s00360-014-0856-6).

66. Stier A., Bize P., Roussel D., Schull Q., Massemin S., Criscuolo F. 2014 Mitochondrial

uncoupling as a regulator of life-history trajectories in birds: an experimental study in the

zebra finch. *J Exp Biol* **217**(19), 3579-3589. (doi:10.1242/jeb.103945).

67. Stier A., Bize P., Massemin S., Criscuolo F. 2021 Long-term intake of the illegal diet pill

DNP reduces lifespan in a captive bird model. *Comp Biochem Physiol C* **242**, 108944.

(doi:https://doi.org/10.1016/j.cbpc.2020.108944).

68. Grimpo, K., Kutschke, M., Kastl, A., Meyer, C.W., Heldmaier, G., Exner, C., Jastroch,

M. 2014 Metabolic depression during warm torpor in the Golden spiny mouse (*Acomys*

russatus) does not affect mitochondrial respiration and hydrogen peroxide release. *Comp*

Biochem Physiol A **167**, 7-14.

69. Nord, A., Nilsson, J.F., Sandell, M.I., Nilsson, J.-Å. 2009 Patterns and dynamics of rest-

phase hypothermia in wild and captive blue tits during winter. *J Comp Physiol B* **179**(6),

737-745.

70. Nord, A., Nilsson, J.F., Nilsson, J.-Å. 2011 Nocturnal body temperature in wintering blue

tits is affected by roost-site temperature and body reserves. *Oecologia* **167**(1), 21-25.

71. Brodin, A., Nilsson, J.-Å., Nord, A. 2017 Adaptive temperature regulation in the little

bird in winter: predictions from a stochastic dynamic programming model. *Oecologia*

185(1), 43-54.

72. Stier A., Bize P., Schull Q., Zoll J., Singh F., Geny B., Gros F., Royer C., Massemin S.,

Criscuolo F. 2013 Avian erythrocytes have functional mitochondria, opening novel

perspectives for birds as animal models in the study of ageing. *Frontiers in Zoology*

10(1), 33. (doi:10.1186/1742-9994-10-33).

73. Nord A., Folkow L.P. 2018 Seasonal variation in the thermal responses to changing

environmental temperature in the world's northernmost land bird. *J Exp Biol* **221**(1),

jeb171124. (doi:10.1242/jeb.171124).

74. Romano A.B., Hunt A., Welbergen J.A., Turbill C. 2019 Nocturnal torpor by superb

fairy-wrens: a key mechanism for reducing winter daily energy expenditure. *Biol Lett*

15(6), 20190211. (doi:doi:10.1098/rsbl.2019.0211).

75. Hiebert S.M. 1990 Energy costs and temporal organization of torpor in the rufous

hummingbird (*Selasphorus rufus*). *Physiol Zool* **63**, 1082-1097.

76. Brigham R.M. 1992 Daily torpor in a free-ranging goatsucker, the common poorwill

(*Phalaenoptilus nuttallii*). *Physiol Zool* **65**(2), 457-472. (doi:10.2307/30158263).

TABLES

Table 1. Calculation method and definition of mitochondrial respiration states and flux

control ratios. All traits were derived based on changes in O_2 consumption rate in response to

the addition of mitochondrial agonists and antagonists during the assay as described in the

main text.

Parameter	Calculation method	Definition
ROUTINE	$R_{Baseline} - R_{Antimycin\ A}$	Baseline O_2 consumption during oxidative phosphorylation under endogenous cellular conditions.
OXPHOS	$R_{Baseline} - R_{Oligomycin}$	O_2 consumption used to drive oxidative phosphorylation where ATP is produced.
ETS	$R_{FCCP} - R_{Antimycin\ A}$	O_2 consumption when the electron transport system is operating at its maximum capacity.
LEAK	$R_{Oligomycin} - R_{Antimycin\ A}$	O_2 consumption needed to compensate for proton leakiness of the mitochondria.
E-R control efficiency	1 - (ROUTINE / ETS)	Proportion of ETS maximum working capacity remaining under endogenous cellular conditions.
R-L control efficiency	(ROUTINE - LEAK) / ROUTINE	Proportion of endogenous respiration used for ATP production via oxidative phosphorylation.
E-L coupling efficiency	1 - (LEAK / ETS)	Tightness of electron transport during ATP production in a stimulated cellular state.

Figure 1. Overview of the experimental design and timeline of events when studying the effect of thermal state (normothermia or hypothermia) on functional properties of mitochondria in avian blood cells. Studies were performed using wild great tits (*Parus major*) that wintered outside the city of Lund in southernmost Sweden.

Figure 2. Mitochondrial respiration traits in whole blood of winter-adapted, wild, great tits (*Parus major*) at a representative daytime normothermic body temperature (41 °C) and a representative night-time hypothermic body temperature (35 °C). The same individuals were measured in both thermal states. Solid plotting symbols with error bars represent model-predicted means ± s.e.m. Gradient lines show individual responses. Sample sizes and statistics are reported in Table S2. (A) baseline ("ROUTINE") oxygen consumption during oxidative phosphorylation. (B) oxygen consumption devoted to ATP production alone ("OXPHOS") (C) maximum working capacity of the electron transport system ("ETS"). (D) the part of ROUTINE used to offset proton leak across the inner mitochondrial membrane ("LEAK").

Figure 3. Control ratios for mitochondrial respiration in great tit whole blood in a

normothermic (41 ºC) and a hypothermic (35 ºC) thermal state. (A) E-R control efficiency

indicates how intensively the mitochondria were respiring under endogenous conditions (i.e.,

ROUTINE) relative to their maximum working capacity (i.e., ETS). For example, a value of

0.4 indicates that mitochondrial respiration can increase by 40% before reaching maximum.

(B) The fraction of endogenous respiration that is used for oxidative phosphorylation where

ATP is formed. (C) Coupling efficiency of electron transport when the ETS is stimulated to

work at maximum capacity, where a value of 0 indicates a fully uncoupled system and a

value of 1 a fully coupled system (i.e., where all respiration is towards phosphorylation).

Samples were collected from wild birds and the same individual was measured in both

thermal states. Solid plotting symbols with error bars represent model-predicted means ±

s.e.m. Gradient lines show individual responses. Sample sizes and statistics are reported in

Table S2.

Figure 4. The relationship between mitochondrial respiration traits measured in great tit

whole blood and mean body temperature of the individual during the preceding night. Panels

A-C show these relationships for ROUTINE, ETS, and LEAK respiration, neither of which

were statistically significant as denoted by dotted lines. Panel D) shows the significant

relationship with E-R control efficiency, an index of mitochondrial reserve capacity, denoted

by the solid lines. Data were collected from wild great tits and respiration was assayed at

each of a normothermic (41°C; red triangles) and a hypothermic (35°C; blue circles) thermal

state *in vitro*. Sample sizes and statistics are reported in Table S2.