Plasticity of mitochondrial function restores fuel production during rest-phase hypothermia in
a winter bird

Authors: Carmen C. García-Díaz[1], Imen Chamkha[2], Eskil Elmér[2], Andreas Nord[1,*]

[1] Lund University, Department of Biology, Section for Evolutionary Ecology, Sölvegatan 37,
SE-223 62 Lund, Sweden
[2] Lund University, Department of Clinical Sciences, Mitochondrial Medicine, Sölvegatan 17,
SE-221 84, Lund, Sweden

* Author for correspondence: AN, e-mail: andreas.nord@biol.lu.se

ORCID numbers:
CCGD: 0000-0002-7827-8104
IM: 0000-0002-6919-585X
EE: 0000-0001-7901-1826
AN: 0000-0001-6170-689X

AUTHOR CONTRIBUTIONS: AN conceived the idea and designed the protocol with input
from EE. AN performed the fieldwork, and CCGD, IM and AN performed the laboratory
work. All authors interpreted the results. CCGD and AN analysed the data and wrote the first
draft. CCGD produced the graphic material. All authors critically revised and edited the
manuscript. AN procured funding.

ABSTRACT

Many small animals routinely regulate resting body temperature several degrees below active-

phase levels (rest-phase hypothermia), which reduces heat transfer rate and tissue oxygen

demand and so may confer energy savings. Small birds that winter at high latitude show

limited capacity for rest-phase hypothermia, meaning they cannot avoid upregulating heat

production when facing winter cold. Substrates for upholding body temperature are provided

by mitochondria in the form of adenosine triphosphate (ATP), but mitochondrial respiration is

probably lower during hypothermia because of the temperature-dependence of biological

processes (i.e., Q_{10} effect). Thus, there could be a conflict between increased organismal fuel

demand and a lower mitochondrial capacity to provide it during rest-phase hypothermia. We

studied this enigma by assessing mitochondrial function in the blood cells of wintering great

tits (*Parus major*) that had spent the preceding night in warm, mild, or cold temperatures, at

each of a hypothermic and a normothermic thermal state *in vitro*. Hypothermia reduced

mitochondrial respiration by 13 %. However, this did not affect respiration allocated to

oxidative phosphorylation, because hypothermia was also associated with a reduction in non-

phosphorylating respiration, from 17 % in normothermia to 4 % in hypothermia. The

proportion of non-phosphorylating respiration also decreased in response to colder night

temperatures Thus, the coupling efficiency between electron transport and ATP production in

maximally stimulated cells increased from 91% in normothermia to 98% in hypothermia. Our

study shows that mitochondrial function can be highly plastic on short temporal scales. Thus,

functional changes in the electron transport system might safeguard ATP production at lower

tissue temperatures and when organismal demand for fuel increases in cold winter

temperatures.

Keywords: blood cells, body temperature, cell respiration, great tit, mitochondria,

thermoregulation, winter

INTRODUCTION

Mammals and birds rely mostly on endogenous heat production to thermoregulate. An endothermic lifestyle carries many benefits, such as the possibility of sustaining high work rate (largely) independently of variation in air temperature. However, it can be cumbersome to amass the resources needed to support the high metabolic throughput machinery that is required for this to happen. This is the case during high latitude winters, when cold temperature increases the energy cost of staying warm at the same time as hours of daylight for foraging, and food availability itself, reach their annual lows. Animals that remain resident in such biomes throughout the year adapt accordingly by a combination of behavioural plasticity and seasonal morphological and physiological adjustments, rendering winter phenotypes better equipped to produce heat and withstand heat loss see [1,2 for reviews].

Many small to medium-sized endotherms save energy by rest-phase hypothermia, a regulated reduction in resting body temperature body temperature below the euthermic level during the resting phase [3]. This typically occurs in response to various energetic challenges, such as food scarcity or inclement weather, and may reduce metabolic demands by more than 50% [4]. It follows that most animals using rest-phase hypothermia have high metabolic intensity, resultant high heat loss rate, and utilises high-energy or ephemeral food sources [3]. This fits the description of "the little bird in winter" sensu [5], i.e., small (< 20 g) songbirds that are year-round residents at high latitudes where they routinely meet environmental temperatures 50-60 °C below body temperature. Yet, most winter birds in northern forests, such as tits and chickadees, rarely show daily variation in body temperature > 5 °C e.g. [6-8] and generally do not exceed 7-8 °C even in severe cold [9-11] (but see [12]). Theoretical models predict even such moderate body temperature reduction can carry survival benefits [13,14]. However, the modest extent of hypothermia means that the little bird in winter can never forego a significant

increase in energetically costly heat production to counter winter cold, even when maximally

hypothermic (e.g. [15]).

Most of the energy used by the working body is in the form of adenosine triphosphate (ATP)

that is produced at the expense of oxygen in the mitochondria. This occurs by dissipation of

the energy contained in a proton gradient across the inner mitochondrial membrane that is

accumulated by the mitochondria's electron transport system (ETS) [16]. Since more energy is

needed to stay warm in winter, it is not surprising that many endotherms upregulate

mitochondrial content and/or respiration rate in key metabolic tissues when cold exposed [17-21].

However, a potential problem for a hypothermic animal is that the activity of the respiratory

chain could be slower in a colder body (e.g.[22]), resulting in a conflict between increased

organismal demand for fuel but reduced mitochondrial capacity to provide it. However, not all

the proton motive force generated by the ETS is used to produce substrate, because under the

influence of uncoupling proteins and due to spontaneous "slip", protons also leak back into

the mitochondrial matrix without generating ATP in the process [23]. This process, known as

mitochondrial proton leak, constitutes some 20-25% of resting metabolic rate in endotherms

[24,25]. Thus, there could be scope to adjust the coupling of ETS to ATP production according to

the animal's need. For example, increased mitochondrial coupling during energy limitation

makes ATP production more effective for a given respiration rate [26-29]. It could, thus, be

speculated that improved coupling could secure adequate ATP production even when the

overall energy expenditure of the cell is reduced, such as when heat production must remain

high, but substrate delivery is reduced because rest-phase hypothermia causes the

mitochondria to work at a slower pace. There is support for this compensation hypothesis

from studies on ectotherms [30-32] but see [33,34]. Similar studies on endotherms suggests that

while phosphorylating respiration is often actively suppressed in hypometabolic animals, non-

phosphorylating (i.e., leak) respiration is sometimes suppressed, sometimes unaltered, and

sometimes increased reviewed by [35].

Previous endotherm studies have largely been performed in hibernators or in animals capable

of deep torpor; thermal states characterised by substantial metabolic downregulation [4]. In such

models, it is likely that the combination of direct thermal suppression and active

downregulation of phosphorylating respiration allows for sufficient energy conservation

without a concomitant need to tighten coupling. This is not easily extrapolated to the

problems faced by the little bird in winter, which must increase energy expenditure sharply to

counter cold temperature even when hypothermic. Hence, we studied mitochondrial responses

to *in vitro* simulation of rest-phase hypothermia in the great tit (*Parus major* L.) – a small (16-

20 g) songbird that winters at high latitude and show pronounced diel variation in body

temperature [6,36]. The birds spent the night preceding these measurements in conditions

representative of mild or very cold winter temperature, or in thermoneutrality. This allowed us

to investigate how short-term changes to the thermal environment affected the mitochondrial

response to reduced body temperature. We exploited the fact that birds contain functional

mitochondria in their red blood cells, which makes it possible to gain non-terminal insights

into mitochondrial function. Previous studies have found that bird blood cell mitochondria

respond in the short-term to variation in energetic demands [37-39], over longer time periods in

line with age-related metabolic senescence [40] and according to environmental context [20]. If

better coupled mitochondria serve as an adaptation to secure a high level of cellular energy

production even when the overall energy expenditure of the cell is reduced, we expected that

coupling should increase in response to low night-time air temperature and during

hypothermia *in vitro*. If, on the other hand, mitochondrial uncoupling can reduce the need for

shivering thermogenesis (the requires ATP) or reduce the amount of ROS produced during in

shivering [41,42], hypothermia and low air temperature could also be associated with increased

proton leak.

MATERIALS AND METHODS

The experimental design and timeline are outlined in Fig. 1.

Study site and animals

Great tits were caught at two sites in the vicinity of Lund in southernmost Sweden – Räften (n

= 37) and Vomb (n = 21) – between January and February 2021. Räften (55° 43'N, 13° 17'E")

is an oak (*Quercus robur*) dominated 32 ha woodlot surrounded by arable fields. The Vomb

site (55° 39'N, 13° 33'E) is part of a pine (*Pinus sylvestris*) plantation with a rich deciduous

understory composed mainly of sallow (*Salix* spp.), birch (*Betula pendula*) and rowan (*Sorbus*

aucuparia). Birds in Räften were caught in mist nets close to feeders baited with sunflower

seeds that were erected 1.5 m above ground in January 2021. The birds in Vomb were caught

shortly before sunrise when roosting in nest boxes. All birds were brought to Lund University

within 2 h of capture, where they were housed in cages measuring $77 \times 44 \times 77$ cm (length $\times$

width $\times$ height) at 5 °C under simulated natural photoperiod and were provided sunflower

seeds, peanuts, and water *ad libitum.* One-two days after capture the birds were implanted

with a temperature-sensitive PIT tag (BioTherm13, Biomark, Boise, ID, USA) into the

intraperitoneal cavity under local anaesthesia (5% lidocaine) as part of a different study. Eight

to 34 h later, the birds were put in a 1 L hermetically sealed glass container ventilated with

dry air at 400-500 ml / min for measurement of metabolic rate and body temperature in a

climate chamber (Weiss Umwelttechnik C180, Reiskirchen, Germany) at thermoneutrality (25

°C; n = 20; Andreasson et al., 2020), during a simulated mild winter night (5 °C: n = 19) and

during a simulated very cold winter night (-15 °C; n = 19) for 13-15 h, again for use in the

other study.

Ambient temperature in Lund (10-40 km from the study sites) ranged -14 °C to 7 °C during

the study period and photoperiod ranged 8:20 h to 10:20 h.

Blood sampling and mitochondrial measurement in whole blood

In the morning after the metabolic measurements, a blood sample ≤ 10% of estimated total

blood volume was obtained from the jugular vein using a heparinized syringe within 5 min of

removing the bird from the metabolic chamber. The sample was stored at 10-12 °C in 2 mL

EDTA (ethylenediaminetetraacetic acid) tubes (BD Vacutainer®, Franklin Lakes, NJ, USA)

without agitation until analysed 0.5-2.5 h later.

We measured mitochondrial respiration in whole blood using an Oxygraph O2k high

resolution respirometer (Oroboros Instruments, Innsbruck, Austria), following the protocol

described in Nord et al. (2022). To test how rest-phase hypothermia affected mitochondrial

function, we ran duplicate samples from the same individual simultaneously in each of two

thermal states: a representative normothermic daytime bird body temperature (41 °C;

henceforth "normothermia") and a representative hypothermic night-time body temperature

(35 °C; henceforth "hypothermia") [43] using two respirometers. These were calibrated at assay

temperature (i.e., 35 °C or 41 °C) daily, and we changed instrument/assay temperature

combinations between days.

The blood samples were allowed 5-10 min on a rotating mixer in room temperature before the

start of the assay. Then, we added 50 µL sample to 1.95 mL MiR05 respiration medium (0.5

mM EGTA, 3 mM $MgCl_2$, 60 mM K-lactobionate, 20 mM taurine, 10 mM KH_2PO_4, 20 mM

Hepes, 110 mM sucrose, free fatty acid bovine serum albumin (1 g L^{-1}), pH 7.1) that was

contained within the respirometer chambers and so was pre-equilibrated at assay temperature.

The final volume contained in the respirometer chamber was, thus, 2 mL.

Mitochondrial respiration rate was measured as the rate of decline in O_2 concentration in the

chamber. Once the O_2 consumption had stabilized upon closing the chamber, recorded

baseline O_2 consumption on endogenous substrates ('ROUTINE') for 2-3 min. Then, we

assessed mitochondrial function by sequential addition of three mitochondrial agonists and

antagonists. Firstly, we added the ATP-synthase inhibitor oligomycin at a final concentration

of 5 µmol L^{-1}, which blocks the process of oxidative phosphorylation ('OXPHOS'). Thus, the

remaining O_2 after oligomycin addition is attributed to proton leak ('LEAK'). Next, we added

the mitochondrial uncoupler FCCP (carbonyl cyanide-p-trifluoro-methoxyphenyl-hydrazone),

which abolishes the proton gradient across the inner mitochondrial membrane, causing the

electron transport system to work at maximum capacity to restore the gradient ('ETS').

Because an excessive concentration of FCCP inhibits mitochondrial respiration and can lead

to acidification of the cell cytoplasm [16], we first added 1.5 µL of 1 mmol L^{-1} FCCP and then

titrated FCCP in 0.5 µL increments until maximum O_2 consumption was reached. The final

concentration ranged 1-1.75 µmol L^{-1}. Finally, we added 5 µmol L^{-1} of antimycin A, which

blocks Complex III in the electron transport chain. Therefore, the remaining O_2 consumption

after antimycin A addition is due to non-mitochondrial respiration.

Statistical analyses

We excluded five of the 57 duplicate samples measured at 41 °C since they were not inhibited

by the addition of oligomycin. Thus, the final data set consisted of 57 observations at 35 °C

and 52 observations at 41 °C. We then calculated mitochondrial respiration states, flux control

ratios (FCR) and coupling efficiency according to Table 1.

All statistical analyses were performed using R version 4.0.4 (R Core Team, 2020). We used

linear mixed models (lmer in the lme4 package) [44] to test how each respiration state and flow

control ratio was affected by assay temperature, night temperature and assay temperature ×

night temperature. Bird ID nested in night temperature was used as a random factor to account

for repeated measurements. Final models were derived using backward elimination of non-

significant terms based on likelihood ratio tests [45]. Predicted values and their standard errors

(s.e.m.) were calculated using the emmeans functions in the emmeans package [46]. Post hoc

tests were performed using the pairs function in emmeans.

RESULTS

A representative normothermic experiment is presented in Fig. 2. All details on the statistical

tests are presented in Table 2.

The effect of thermal state on ROUTINE was similar in all night temperatures (i.e., assay

temperature × night temperature: $P = 0.568$) and ROUTINE was not affected by night

temperature ($P = 0.406$). However, ROUTINE was 13 % higher in normothermia

(0.572±0.020 pmol O_2 s^{-1} μl^{-1}) than at hypothermia (0.551±0.020 pmol O_2 s^{-1} μl^{-1}) ($P < 0.001$)

(Fig. 3A), i.e., endogenous respiration was slower when the blood was colder. OXPHOS was

not influenced by the interaction between thermal state and night temperature ($P = 0.895$) and

was also not affected by thermal state ($P = 0.450$) or night temperature ($P = 0.218$) alone (Fig.

3B). This was also true for ETS (thermal state × night temperature: $P = 0.214$; thermal state:

P = 0.480; night temperature: *P* = 0.308) (Fig. 3C). However, LEAK was markedly affected

by assay temperature ($P < 0.0001$), being more than three times higher during normothermic

(0.096±0.004 pmol O_2 s^{-1} μl^{-1}) compared to hypothermic (0.023±0.004 pmol O_2 s^{-1} μl^{-1})

assays (Fig. 3D).

The mitochondria respired at a higher proportion of their maximum capacity when

normothermic (0.570±0.013, of 1) than when hypothermic (0.496±0.012) (i.e.,

$FCR_{ROUTINE/ETS}$ was lower during hypothermia). In other words, hypothermic cells had more

surplus respiratory capacity compared to normothermic cells (Fig. 4A). The transition from a

normothermic to a hypothermic state also affected the contribution of LEAK to endogenous

respiration (i.e., there was a significant change in $FCR_{LEAK/ROUTINE}$; Fig. 4B). Specifically,

during normothermia LEAK accounted for nearly 17% of endogenous respiration

(0.166±0.007), but in hypothermia LEAK contributed only 4% to ROUTINE (0.042±0.007)

(Fig. 4). Coupling efficiency when the cells were induced to work harder (i.e., when maximal

respiration was stimulated using FCCP) increased from 91% (0.906±0.004) in normothermia

to 98% (0.979±0.004) in hypothermia (Fig. 4C). Thus, the electron transport chain was tighter

when in a hypothermic state. When thermal states were accounted for, there was also a

tendency for a change to $FCR_{LEAK/ROUTINE}$ in response to night temperature ($P = 0.059$),

whereby LEAK contributed less to ROUTINE after a night in -15°C than after a night at -5

and 25°C, though differences in $FCR_{LEAK/ROUTINE}$ were small between the coldest and

warmest night temperatures (Table 2). Fig. 4

DISCUSSION

We tested whether mitochondrial respiration in great tit blood cells varied with thermal state

(i.e., normothermia or hypothermia) and air temperature. Endogenous respiration

('ROUTINE') was lower during *in vitro* hypothermia (Fig. 3A). However, the part of

respiration devoted to oxidative phosphorylation (OXPHOS; in which ATP is produced)

remained constant during the transition from normo- to hypothermia (Fig. 3B). At the same

time, non-phosphorylating (i.e., LEAK) respiration decreased threefold (Fig. 3D). As a result,

LEAK constituted significantly less of endogenous respiration rate in the hypothermic state

(i.e., $FCR_{LEAK/ROUTINE}$ was lower; Fig. 4B) and coupling efficiency when the cells were

stimulated to work hard also increased during hypothermia Fig. 4C). Non-uniformity of

changes to respiration traits indicates that the mitochondrial response to cooling were actively

regulated and not merely a consequence of thermal suppression of the ETS (cf. [47,48]). Hence,

our study indicates that flexibility in mitochondrial function may allow the little bird in winter

to secure sufficient substrate delivery to meet thermoregulatory demands despite a general

depression of cellular respiration during hypothermia. This notion should be explored by

direct measurement of ATP production and efficiency of phosphorylating respiration in

normo- and hypothermic birds. Future studies should also address if the response recorded in

blood cells mirrors that in other tissues with more clearly defined roles in energy metabolism

and thermogenesis. For example, hibernating or torpid mammals show downregulation of

phosphorylating and non-phosphorylating respiration in some tissues (e.g., the liver) [49,50] but

not in others (e.g., skeletal muscles) (see [35,50] for reviews). In some cases, metabolically

suppressed states may even be associated with reduced coupling of the ETS, possibly to aid

local thermogenesis in the central nervous system [51].

Reduced LEAK and improved coupling efficiency during hypothermia is in keeping with the

hypothesis of biochemical compensation for thermal suppression of cell respiration at lower

body temperature. However, tighter coupling can potentially result in increased production of

reactive oxygen species (ROS) as a by-product of electron transport [23,52], provided that the

resultant increase in protonmotive force is not dissipated through ATP synthase. Unless

quenched, ROS cause oxidative stress with resultant macromolecular damage e.g. [53], which is

related to many medical disorders [54]. This has given rise to the "uncoupling to survive"

hypothesis, in which mitochondrial ROS are implicated in the aging process [55,56]. It is

interesting to speculate that a hypothermic bird might have to accept an oxidative cost to

uphold ATP production. This could explain why the suppressing effect of low night

temperature on LEAK was only evident at the normothermic assay temperature (Fig. 3D) but

stabilised at a constant low level in the hypothermic state. In this context, our study could

provide proximate insights into why oxidative stress has been found to be higher in birds that

are more hypothermic at night [57]. However, the role of mitochondria as sources of cellular

ROS has been contended [58-60] and empirical evidence suggests that the relationship between

body temperature, ATP production and ROS generation is not straightforward [29,61,62]. In line

with this, even though (pharmacological) uncoupling protects birds from cold-induced

oxidative stress [41], the evidence for clear effects thereof on fitness in this taxon are equivocal

[63,64]. It is advisable that future studies measure both ATP- and ROS production, and

membrane potential, to determine if a hypothermic bird pays an oxidative cost.

We found evidence of improved coupling of the ETS after a 12-14 h acute exposure to lower

night-time temperature (Fig. 4A). This contrasts previous studies on birds and mammals,

where cold exposure or cold acclimatisation was associated with increased non-

phosphorylating respiration or increased expression of uncoupling proteins [65-69]. These

responses have been studied most frequently in skeletal muscle, and have been implicated as a

transition towards increased reliance on non-shivering thermogenesis, though its occurrence

and significance in the skeletal muscles of birds remains dubious [70] but see e.g. [68,71]. Similar

increases in LEAK respiration were also found in the blood cells of three temperate species of

tits during the transition from mild autumn to cold winter temperature [20]. It is interesting to

note that the studies cited above have measured changes to mitochondrial respiration after

longer-term exposure to cold temperature whereas we measured the response to an immediate

temperature change. It is possible that the short-term response to colder environmental

temperature is a tightening of the ETS to secure substrate delivery, whereas the longer-term

response may reflect a balance between the competing demands of producing ATP and

keeping ROS levels minimal that could be facilitated by a simultaneous increase of

mitochondrial content (e.g.[20]).

This study adds new perspectives to our understanding of how cellular- and whole animal

metabolism interact when the little bird in winter adapts to low temperatures. We conclude

that, in the short term, blood cell mitochondria maintain phosphorylating respiration by

reducing proton leak and increasing coupling efficiency when body or air temperature

decreases. It is possible, therefore, that rest-phase hypothermia may come at the cost of

increased oxidative stress (cf. [57]). It is interesting to note that this acute response to reduced

air temperature is opposite to what has been recorded as part of seasonal acclimatisation in the

same model system [20] and in response to cold exposure in tissues with more clearly defined

metabolic or thermogenic roles in other models (see references above). These striking

difference in short- and long-term responses to colder temperature demands a need to study

the temporal resolution of blood cell responses in some detail. Even so, this work adds to the

body of evidence highlighting the utility of studying blood cells to gain minimally invasive

insight when integrating mitochondrial function into the study of functional ecology,

evolutionary physiology, and gerontology in birds [37,38,40,72,73]. Future studies should address

how changes to phosphorylating and non-phosphorylating respiration during hypothermia in

this model correlates with simultaneous changes to ATP production and ROS generation and

how responses in blood cells correlate with those in other tissues. More broadly, it would be

relevant to increase taxonomic and phenotypic coverage of these studies, spanning cold-

hardy, non-hypothermic, birds (e.g. [74]) to those that routinely use shallow (e.g. [75]) or deep

torpor ([76-78]).

ACKNOWLEDGEMENTS

The authors thank Stefan Nord for managing the Räften feeding stations by which birds for

the study were caught. SN also assisted with fieldwork and provided miscellaneous technical

support during the study.

FUNDING

This study was supported by the Birgit and Hellmuth Hertz Foundation (grant no. 2017-

39034), The Royal Physiographic Society in Lund (grant no. 2019-41011), the Swedish

Research Council (grant no. 2020-04686) and Stiftelsen Lunds Djurskyddsfond (all to AN).

ETHICS

All procedures were approved by the Malmö/Lund Animal Ethics Committee, acting under

jurisdiction of the Swedish Board of Agriculture (permit no. 9246-19). Bird ringing was

permitted by the Swedish Bird Ringing Centre at Natural History Museum of Sweden (license

number 723, to AN).

DATA AVAILABILITY

Data will be deposited in Figshare upon acceptance of the manuscript.

CONFLICT OF INTEREST

The authors declare that they have no competing or financial interests.

REFERENCES

1. Blix AS. Adaptations to polar life in mammals and birds. *J Exp Biol.* 2016;219(8):1093-1105.

2. Swanson DL, Vézina F. Environmental, ecological and mechanistic drivers of avian seasonal metabolic flexibility in response to cold winters. *J Ornithol.* 2015;156(1):377-388.

3. Geiser F. *Ecological physiology of daily torpor and hibernation.* Cham, Switzerland: Springer Nature Switzerland AG; 2021.

4. Ruf T, Geiser F. Daily torpor and hibernation in birds and mammals. *Biol Rev.* 2015;90(3):891-926.

5. Brodin A. Theoretical models of adaptive energy management in small wintering birds. *Philosophical Transactions of the Royal Society B.* 2007;362(1486):1857-1871.

6. Nilsson JF, Nilsson J-Å, Broggi J, Watson H. Predictability of food supply modulates nocturnal hypothermia in a small passerine. *Biol Lett.* 2020;16(6):20200133.

7. Nord A, Nilsson JF, Nilsson J-Å. Nocturnal body temperature in wintering blue tits is affected by roost-site temperature and body reserves. *Oecologia.* 2011;167(1):21-25.

8. Nord A, Nilsson JF, Sandell MI, Nilsson J-Å. Patterns and dynamics of rest-phase hypothermia in wild and captive blue tits during winter. *J Comp Physiol B.* 2009;179(6):737-745.

9. Reinertsen RE, Haftorn S. The effect of short-time fasting on metabolism and nocturnal hypothermia in the willow tit *Parus montanus*. *J Comp Physiol.* 1984;154(1):23-28.

10. Reinertsen RE, Haftorn S. Nocturnal hypothermia and metabolism in the willow tit *Parus montanus* at 63 degrees N. *J Comp Physiol.* 1983;151(2):109-118.

11. Haftorn S. Hypothermia of tits in the Arctic winter. *Orn Scand.* 1972;3(2):153-166.

381   12.   Cooper SJ, Gessaman JA. Nocturnal hypothermia in seasonally acclimatized Mountain
382   Chickadees and Juniper Titmice. *Condor.* 2005;107(1):151-155.

383   13.   Brodin A, Nilsson J-Å, Nord A. Adaptive temperature regulation in the little bird in
384   winter: predictions from a stochastic dynamic programming model. *Oecologia.*
385   2017;185(1):43-54.

386   14.   Welton NJ, Houston AI, Ekman J, McNamara JM. A dynamic model of hypothermia
387   as an adaptive response by small birds to winter conditions. *Acta Biotheor.*
388   2002;50(1):39-56.

389   15.   Andreasson F, Nord A, Nilsson J-Å. Age differences in night-time metabolic rate and
390   body temperature in a small passerine. *J Comp Physiol B.* 2020;190(3):349-359.

391   16.   Nicholls DG, Ferguson SJ. *Bioenergetics.* 4th ed. Cambridge, MA, USA: Academic
392   Press; 2013.

393   17.   Liknes ET, Swanson DL. Phenotypic flexibility in passerine birds: Seasonal variation
394   of aerobic enzyme activities in skeletal muscle. *J Therm Biol.* 2011;36(7):430-436.

395   18.   Wang Y, Shan S, Zhang H, Dong B, Zheng W, Liu J. Physiological and biochemical
396   thermoregulatory responses in male Chinese hwameis to seasonal acclimatization:
397   Phenotypic flexibility in a small passerine. *Zool Stud.* 2019;58:e6-e6.

398   19.   Zheng WH, Li M, Liu J-S, Shao S-L, Xu X-J. Seasonal variation of metabolic
399   thermogenesis in Eurasian tree sparrows (*Passer montanus*) over a latitudinal gradient.
400   *Physiol Biochem Zool.* 2014;87(5):704-718.

401   20.   Nord A, Metcalfe NB, Page JL, Huxtable A, McCafferty DJ, Dawson NJ. Avian red
402   blood cell mitochondria produce more heat in winter than in autumn. *FASEB J.*
403   2021;35(5):e21490.

21. Schaeffer PJ, Villarin JJ, Lindstedt SL. Chronic cold exposure increases skeletal
muscle oxidative structure and function in *Monodelphis domestica*, a marsupial
lacking brown adipose tissue. *Physiol Biochem Zool.* 2003;76(6):877-887.

22. Wright KP, Jr., Hull JT, Czeisler CA. Relationship between alertness, performance,
and body temperature in humans. *Am J Physiol.* 2002;283(6):R1370-1377.

23. Divakaruni AS, Brand MD. The Regulation and Physiology of Mitochondrial Proton
Leak. *Physiology.* 2011;26(3):192-205.

24. Porter RK, Brand MD. Causes of differences in respiration rate of hepatocytes from
mammals of different body mass. *Am J Physiol.* 1995;269(5 Pt 2):R1213-1224.

25. Rolfe DF, Brand MD. Contribution of mitochondrial proton leak to skeletal muscle
respiration and to standard metabolic rate. *Am J Physiol.* 1996;271(4 Pt 1):C1380-
1389.

26. Roussel D, Marmillot V, Monternier P-A, et al. Skeletal muscle metabolism in sea-
acclimatized king penguins. II. Improved efficiency of mitochondrial bioenergetics. *J
Exp Biol.* 2020;223(21):jeb233684.

27. Bourguignon A, Rameau A, Toullec G, Romestaing C, Roussel D. Increased
mitochondrial energy efficiency in skeletal muscle after long-term fasting: its
relevance to animal performance. *J Exp Biol.* 2017;220(13):2445-2451.

28. Monternier PA, Marmillot V, Rouanet JL, Roussel D. Mitochondrial phenotypic
flexibility enhances energy savings during winter fast in king penguin chicks. *J Exp
Biol.* 2014;217(15):2691-2697.

29. Boël M, Romestaing C, Duchamp C, et al. Improved mitochondrial coupling as a
response to high mass-specific metabolic rate in extremely small mammals. *J Exp
Biol.* 2020;223(5):jeb215558.

30. Glanville EJ, Seebacher F. Compensation for environmental change by

complementary shifts of thermal sensitivity and thermoregulatory behaviour in an

ectotherm. *J Exp Biol.* 2006;209(24):4869-4877.

31. Seebacher F. A review of thermoregulation and physiological performance in reptiles:

what is the role of phenotypic flexibility? *J Comp Physiol B.* 2005;175(7):453-461.

32. Bouchard P, Guderley H. Time course of the response of mitochondria from oxidative

muscle during thermal acclimation of rainbow trout, Oncorhynchus mykiss. *J Exp*

Biol. 2003;206(19):3455-3465.

33. St-Pierre J, Brand MD, Boutilier RG. The effect of metabolic depression on proton

leak rate in mitochondria from hibernating frogs. *J Exp Biol.* 2000;203(9):1469-1476.

34. Bishop T, St-Pierre J, Brand MD. Primary causes of decreased mitochondrial oxygen

consumption during metabolic depression in snail cells. *Am J Physiol.*

2002;282(2):R372-R382.

35. Staples JF, Brown JCL. Mitochondrial metabolism in hibernation and daily torpor: a

review. *J Comp Physiol B.* 2008;178(7):811-827.

36. Nord A, Chiriac S, Hasselquist D, Nilsson J-Å. Endotoxin injection attenuates rest-

phase hypothermia in wintering great tits through the onset of fever. *Funct Ecol.*

2013;27(1):236-244.

37. Casagrande S, Stier A, Monaghan P, et al. Increased glucocorticoid concentrations in

early life cause mitochondrial inefficiency and short telomeres. *J Exp Biol.*

2020;223(15):jeb222513.

38. Stier A, Bize P, Hsu B-Y, Ruuskanen S. Plastic but repeatable: rapid adjustments of

mitochondrial function and density during reproduction in a wild bird species. *Biol*

Lett. 2019;15(11):20190536.

39.   Udino E, George JM, McKenzie M, et al. Prenatal acoustic programming of
453   mitochondrial function for high temperatures in an arid-adapted bird. *Proc R Soc Lond*
454   *B.* 2021;288(1964):20211893.

455   40.   Dawson NJ, Salmón P. Age-related increase in mitochondrial quantity may mitigate a
456   decline in mitochondrial quality in red blood cells from zebra finches (*Taeniopygia*
457   *guttata*). *Exp Gerontol.* 2020;133:110883.

458   41.   Stier A, Massemin S, Criscuolo F. Chronic mitochondrial uncoupling treatment
459   prevents acute cold-induced oxidative stress in birds. *J Comp Physiol B.*
460   2014;184(8):1021-1029.

461   42.   Stier A, Bize P, Habold C, Bouillaud F, Massemin S, Criscuolo F. Mitochondrial
462   uncoupling prevents cold-induced oxidative stress: a case study using UCP1 knockout
463   mice. *J Exp Biol.* 2014;217(4):624-630.

464   43.   Prinzinger R, Pressmar A, Schleucher E. Body temperature in birds. *Comp Biochem*
465   *Physiol A.* 1991;99(4):499-506.

466   44.   Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using
467   lme4. *J Stat Soft.* 2015;67:1-48.

468   45.   Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. *Mixed effects models and*
469   *extensions in ecology with R.* New York: Springer; 2009.

470   46.   Lenth RV. emmeans: Estimated marginal means, aka least-squares means. R package
471   version 1.3.3. https://CRAN.R-project.org/package=emmeans. 2019.

472   47.   Frye LD, Edidin M. The rapid intermixing of cell surface antigens after formation of
473   mouse-muman heterokaryons. *J Cell Sci.* 1970;7(2):319-335.

474   48.   Blicher A, Wodzinska K, Fidorra M, Winterhalter M, Heimburg T. The temperature
475   dependence of lipid membrane permeability, its quantized nature, and the influence of
476   anesthetics. *Biophys J.* 2009;96(11):4581-4591.

477   49.   Brown JCL, Gerson AR, Staples JF. Mitochondrial metabolism during daily torpor in

478   the dwarf Siberian hamster: role of active regulated changes and passive thermal

479   effects. *Am J Physiol.* 2007;293(5):R1833-R1845.

480   50.   Barger JL, Brand MD, Barnes BM, Boyer BB. Tissue-specific depression of

481   mitochondrial proton leak and substrate oxidation in hibernating arctic ground

482   squirrels. *Am J Physiol.* 2003;284(5):R1306-R1313.

483   51.   Laursen WJ, Mastrotto M, Pesta D, et al. Neuronal UCP1 expression suggests a

484   mechanism for local thermogenesis during hibernation. *PNAS.* 2015;112(5):1607-

485   1612.

486   52.   Demine S, Renard P, Arnould T. Mitochondrial uncoupling: A key controller of

487   biological processes in physiology and diseases. *Cells.* 2019;8(8):795.

488   53.   Brand MD, Buckingham JA, Esteves TC, et al. Mitochondrial superoxide and aging:

489   uncoupling-protein activity and superoxide production. *Biochem Soc Symp.*

490   2004(71):203-213.

491   54.   Grasemann H, Holguin F. Oxidative stress and obesity-related asthma. *Paed Resp Rev.*

492   2021;37:18-21.

493   55.   Brand MD. Uncoupling to survive? The role of mitochondrial inefficiency in ageing.

494   *Exp Gerontol.* 2000;35(6-7):811-820.

495   56.   Speakman JR, Talbot DA, Selman C, et al. Uncoupled and surviving: individual mice

496   with high metabolism have greater mitochondrial uncoupling and live longer. *Aging*

497   *Cell.* 2004;3(3):87-95.

498   57.   Zagkle E, Grosiak M, Bauchinger U, Sadowska ET. Rest-phase hypothermia reveals a

499   link between aging and oxidative stress: A novel hypothesis. *Front Physiol.* 2020;11:

500   575060.

58.   Zhang Y, Wong HS. Are mitochondria the main contributor of reactive oxygen
species in cells? *J Exp Biol.* 2021;224(5): jeb221606.

59.   Mookerjee SA, Divakaruni AS, Jastroch M, Brand MD. Mitochondrial uncoupling and
lifespan. *Mech Age Develop.* 2010;131(7-8):463-472.

60.   Brown GC, Borutaite V. There is no evidence that mitochondria are the main source
of reactive oxygen species in mammalian cells. *Mitochondrion.* 2012;12(1):1-4.

61.   Roussel D, Voituron Y. Mitochondrial Costs of Being Hot: Effects of acute thermal
change on liver bioenergetics in toads (*Bufo bufo*). *Front Physiol.* 2020;11: 153.

62.   Boël M, Veyrunes F, Durieux A-C, Freyssenet D, Voituron Y, Roussel D. Does high
mitochondrial efficiency carry an oxidative cost? The case of the African pygmy
mouse (*Mus mattheyi*). *Comp Biochem Physiol A.* 2022;264:111111.

63.   Stier A, Bize P, Roussel D, Schull Q, Massemin S, Criscuolo F. Mitochondrial
uncoupling as a regulator of life-history trajectories in birds: an experimental study in
the zebra finch. *J Exp Biol.* 2014;217(19):3579-3589.

64.   Stier A, Bize P, Massemin S, Criscuolo F. Long-term intake of the illegal diet pill
DNP reduces lifespan in a captive bird model. *Comp Biochem Physiol C.*
2021;242:108944.

65.   Raimbault S, Dridi S, Denjean F, et al. An uncoupling protein homologue putatively
involved in facultative muscle thermogenesis in birds. *Biochem J.* 2001;353(3):441-
444.

66.   Toyomizu M, Ueda M, Sato S, Seki Y, Sato K, Akiba Y. Cold-induced mitochondrial
uncoupling and expression of chicken UCP and ANT mRNA in chicken skeletal
muscle. *FEBS Lett.* 2002;529(2-3):313-318.

67.   Teulier L, Rouanet J-L, Rey B, Roussel D. Ontogeny of non-shivering thermogenesis
525   in Muscovy ducklings (*Cairina moschata*). *Comp Biochem Physiol A.*
526   2014;175(0):82-89.

527   68.   Teulier L, Rouanet JL, Letexier D, et al. Cold-acclimation-induced non-shivering
528   thermogenesis in birds is associated with upregulation of avian UCP but not with
529   innate uncoupling or altered ATP efficiency. *J Exp Biol.* 2010;213(14):2476-2482.

530   69.   Mahalingam S, Cheviron ZA, Storz JF, McClelland GB, Scott GR. Chronic cold
531   exposure induces mitochondrial plasticity in deer mice native to high altitudes. *J
532   Physiol.* 2020;598(23):5411-5426.

533   70.   Stager M, Cheviron ZA. Is there a role for sarcolipin in avian facultative
534   thermogenesis in extreme cold? *Biol Lett.* 2020;16(6):20200078.

535   71.   Nowack J, Giroud S, Arnold W, Ruf T. Muscle Non-shivering thermogenesis and its
536   role in the evolution of endothermy. *Front Physiol.* 2017;8: 889.

537   72.   Ton R, Stier A, Cooper CE, Griffith SC. Effects of heat waves during post-natal
538   development on mitochondrial and whole body physiology: An experimental study in
539   zebra finches. *Front Physiol.* 2021;12: 554.

540   73.   Stier A, Bize P, Schull Q, et al. Avian erythrocytes have functional mitochondria,
541   opening novel perspectives for birds as animal models in the study of ageing. *Front
542   Zool.* 2013;10(1):33.

543   74.   Nord A, Folkow LP. Seasonal variation in the thermal responses to changing
544   environmental temperature in the world's northernmost land bird. *J Exp Biol.*
545   2018;221(1):jeb171124.

546   75.   Romano AB, Hunt A, Welbergen JA, Turbill C. Nocturnal torpor by superb fairy-
547   wrens: a key mechanism for reducing winter daily energy expenditure. *Biol Lett.*
548   2019;15(6):20190211.

76.   Hiebert SM. Energy costs and temporal organization of torpor in the rufous
hummingbird (*Selasphorus rufus*). *Physiol Zool.* 1990;63:1082-1097.

77.   McKechnie AE, Ashdown RAM, Christian MB, Brigham RM. Torpor in an African
caprimulgid, the freckled nightjar *Caprimulgus tristigma*. *J Avian Biol.*
2007;38(3):261-266.

78.   Brigham RM. Daily torpor in a free-ranging goatsucker, the common poorwill
(*Phalaenoptilus nuttallii*). *Physiol Zool.* 1992;65(2):457-472.

TABLES

Table 1. Calculation method and definition of mitochondrial respiration states. Mitochondrial

parameters were measured in terms of changes in O_2 concentration after the addition of

agonists and antagonists of mitochondrial respiration.

Parameter	Calculation method	Definition
ROUTINE	$R_{Baseline} - R_{AntimycinA}$	Baseline O_2 consumption during oxidative phosphorylation under endogenous cellular conditions.
OXPHOS	$R_{Baseline} - R_{Oligomycin}$	O_2 consumption used to drive oxidative phosphorylation where ATP is produced.
ETS	$R_{FCCP} - R_{AntimycinA}$	O_2 consumption when the electron transport system is operating at its maximum capacity.
LEAK	$R_{Oligomycin} - R_{AntimycinA}$	O_2 consumption needed to compensate for proton leakiness of the mitochondria.
$FCR_{ROUTINE/ETS}$	ROUTINE / ETS	Proportion of ETS maximum working capacity used under endogenous cellular conditions.
$FCR_{LEAK/ROUTINE}$	LEAK / ROUTINE	The proportion of endogenous respiration used to compensate for proton leak under non-stimulated cellular conditions.
Coupling Efficiency	1 – (LEAK / ETS)	Coupling efficiency of ATP production in a stimulated cellular state.

Table 2. Results from final models comparing mitochondrial respiration parameters in great

tit whole blood in each of two assay temperatures. Means are presented ± s.e.m. Values for

significant effects (i.e., $P \leq 0.05$) are presented in bold font, and values for effects where 0.05

$< P < 0.10$ are given in italics. Note that test statistics and parameter estimates for main

effects are not provided when interactions are retained in the final model. $*T_{\text{night}[-15°C]}$ vs.

$T_{\text{night}[5°C]}$: $P = 0.058$.

Model and Variable	Estimate (± s.e.m.)	$n_{35°C}$ / $n_{41°C}$	*LRT*	*P*
ROUTINE (pmol O$_2$ s^{-1} μl^{-1})		57 / 52		
Night Temp × Assay Temp			1.13	0.568
Night Temp (-15°C, 5°C or 25°C)			1.80	0.406
Thermal State			17.77	**< 0.001**
Hypothermia (35°C)	0.505±0.020			
Normothermia (41°C)	0.572±0.020			
OXPHOS (pmol O$_2$ s^{-1} μl^{-1})				
Night Temp × Assay Temp		57 / 52	0.22	0.895
Thermal State			0.57	0.450
Night Temp (-15°C, 5°C or 25°C)			3.05	0.218
ETS (pmol O$_2$ s^{-1} μl^{-1})				
Night Temp × Assay Temp		57 / 52	3.08	0.214
Thermal State			0.50	0.480
Night Temp (-15°C, 5°C or 25°C)			2.35	0.308
LEAK (pmol O$_2$ s^{-1} μl^{-1})				
Night Temp × Assay Temp			3.56	0.168
Night Temp (-15°C, 5°C or 25°C)		57/52	2.08	0.354
Thermal State			77.57	**< 0.001**
Hypothermia (35°C)	0.023±0.006			
Normothermia (41°C)	0.096±0.006			
FCR ROUTINE / ETS				
Night Temp × Assay Temp			0.43	0.807
Night Temp (-15°C, 5°C or 25°C)			3.56	0.169
Thermal State		57/52	37.72	**< 0.001**
Hypothermia (35°C)	0.496±0.012			
Normothermia (41°C)	0.570±0.012			
FCR LEAK / ROUTINE				
Night Temp × Assay Temp			1.81	0.404
Night Temp (-15°C, 5°C or 25°C)			5.65	*0.059*
Night Temp = -15°C [A]*	0.088±0.010	18 / 17		
Night Temp = 5°C [A]	0.122±0.010	19 / 17		
Night Temp = 25°C [A]	0.101±0.010	20 / 18		
Thermal State			95.51	**< 0.001**
Hypothermia (35°C)	0.042±0.007			
Normothermia (41°C)	0.166±0.007			
Coupling Efficiency		57 / 52		
Night Temp × Assay Temp			0.90	0.636
Night Temp (-15°C, 5°C or 25°C)			5.15	*0.076*
Thermal State			92.57	**< 0.001**
Hypothermia (35°C)	0.979±0.004			
Normothermia (41°C)	0.906±0.004			

FIGURES

Figure 1. Overview of the experimental design and timeline of events when studying the

effect of thermal state (normothermia or hypothermia) on functional properties of

mitochondria in avian blood cells. Studies were performed using wild great tits (*Parus major*)

that wintered outside the city of Lund in southernmost Sweden.

Figure 2. Representative trace of mitochondrial respiration in whole blood of a winter-
adapted great tit (*Parus major*, individual 2KV07141) when measured at a normothermic
daytime bird body temperature (41 °C). The blue line shows oxygen (O$_2$) concentration in the
test chamber, and the red line shows O$_2$ consumption by the mitochondria on endogenous
substrates (ROUTINE), after the addition of ATP synthase inhibitor oligomycin (Omy)
(OXPHOS), when maximally stimulated by the mitochondrial uncoupler FCCP (carbonyl
cyanide-p-trifluoro-methoxyphenyl-hydrazone) (ETS), and after inhibition by the cytochrome
c reductase inhibitor antimycin A (Ama). LEAK refers to the part of respiration devoted to
compensating for protons escaping over the inner mitochondrial membrane. "Non-mito"
refers to non-mitochondrial respiration, i.e., oxygen consumption by the sample (such as
binding of O$_2$ by haemoglobin) that is not due to mitochondrial use. Respiration states are
defined in Table 1 and were superimposed on the raw data graph for clarity.

Figure 3. Mitochondrial respiration in whole blood of winter-adapted, wild, great tits (*Parus*
major) at a representative daytime normothermic body temperature (41 °C) and a
representative night-time hypothermic body temperature (35 °C). The same individuals were
measured at both thermal states. Solid plotting symbols with error bars represent raw data
means ± s.e.m. Gradient lines show individual responses. Sample sizes and statistics are
reported in Table 2. (A) baseline ("ROUTINE") oxygen consumption during oxidative
phosphorylation. (B) oxygen consumption devoted to ATP production alone ("OXPHOS")
(C) maximum working capacity of the electron transport system when uncoupled from ATP
production ("ETS"). (D) the part of ROUTINE used to offset proton leak across the inner
mitochondrial membrane ("LEAK") in different assay temperatures after previous exposure to
thermoneutral, mild, or very cold nights.

Figure 4. Flux control ratios (FCR) for mitochondrial respiration in great tit whole blood at

a normothermic (41 ℃) and a hypothermic (35 ℃) thermal state. (A) An index of how

intensively the mitochondria were respiring under endogenous conditions (ROUTINE)

relative to their maximum working capacity (i.e., ETS). (B) The fraction of endogenous

(ROUTINE) respiration that is used to compensate for LEAK respiration. (C) Coupling

efficiency of electron transport to ATP production under maximally stimulated working

capacity, i.e., the fraction of maximal respiration that is directed towards oxidative

phosphorylation. Samples were collected from wild birds and the same individual was

measured in both thermal states. Solid plotting symbols with error bars represent raw data

means ± s.e.m. Gradient lines show individual responses. Sample sizes and statistics are

reported in Table 2.