A comparison of software for analysis of rare and common short tandem repeat (STR)

variation using human genome sequences from clinical and population-based samples

John W. Oketch (1), Louise V. Wain (2,3), Edward J. Hollox (1*)

1.   Department of Genetics and Genome Biology, University of Leicester, Leicester UK

2.   Department of Population Health Sciences, University of Leicester, Leicester UK

3.   National Institute for Health Research, Leicester Respiratory Biomedical

Research Centre, Glenfield Hospital, Leicester UK

* Corresponding author:

Dr E Hollox,

Department of Genetics and Genome Biology,

University of Leicester,

Leicester

UK

ejh33@le.ac.uk

Keywords

Short tandem repeat, STR, Genome-In-A-Bottle, 1000 Genomes Project, sequencing, genotyping,

forensic identification, CODIS

Abstract

Short tandem repeat (STR) variation is an often overlooked source of variation between genomes.

STRs comprise about 3% of the human genome and are highly polymorphic. Some cause

Mendelian disease, and others affect gene expression. Their contribution to common disease is

not well-understood, but recent software tools designed to genotype STRs using short read

sequencing data are beginning to address this. Here, we compare software that genotypes

common STRs and rarer STR expansions genome-wide, with the aim of applying them to

population-scale genomes. By using the Genome-In-A-Bottle (GIAB) consortium and 1000

Genomes Project sequencing data, we compare performance in terms of sequence length, depth,

computing resources needed, genotyping accuracy and number of STRs genotyped. To ensure

broad applicability of our findings, we also measure genotyping performance against a set of

genomes from clinical samples with known STR expansions, and a set of STRs commonly used

for forensic identification. We find that HipSTR, ExpansionHunter and GangSTR perform well in

genotyping common STRs, including the CODIS 13 core STRs used for forensic analysis.

GangSTR and ExpansionHunter outperform HipSTR for genotyping call rate and memory usage.

ExpansionHunter denovo (EHdn), STRling and GangSTR outperformed STRetch for detecting

expanded STRs, and EHdn and STRling used considerably less processor time compared to

GangSTR. Analysis on shared genomic sequence data provided by the GIAB consortium allows

future performance comparisons of new software approaches on a common set of data, facilitating

comparisons and allowing researchers to choose the best software that fulfils their needs.

Introduction

Over the past two decades, most research on the contribution of genetic variation in the human

genome to disease has focused on single nucleotide variation. Short tandem repeat (STR)

variation has been generally overlooked, as it is not readily assayed by chip-based hybridisation

approaches. STRs are short DNA sequence motifs, typically 2-6bp in size, that are repeated

multiple times in tandem. They are distributed throughout the human genome, comprising about

3% of the genome sequence, and have a mutation rate generally much higher than single

nucleotide variants (around 2×10^{-3} per locus per generation for STRs compared to 10^{-8} for single

nucleotide variants) (Brinkmann et al., 1998; Burgarella and Navascués, 2011; Willems et al.,

2014; Gymrek, 2017). They are frequently polymorphic with multiple alleles in a population, as

expected by neutral population genetics theory for loci with high mutation rates.

Particular STRs cause a variety of severe rare monogenic diseases that are inherited in a

Mendelian manner (Hannan, 2018). These include triplet-repeat expansion diseases, such as

Huntington's disease and myotonic dystrophy, caused by expansion of 3 bp repeat STRs in the

coding region of *HTT*, and *DMPK* genes respectively. Some disorders also occur with other repeat

motifs in both coding and in non-coding regions (Hannan, 2018; Paulson, 2018; Lieberman et al.,

2019; van Kuilenburg et al., 2019). Emerging evidence shows that STRs modulate disease risk in

several complex diseases such as autism and neurodegenerative disorders (Hannan, 2020; Trost

et al., 2020).

STRs can affect the function of genes in three main ways, either by directly altering the coding

region of genes, by disrupting introns, or by affecting expression levels as part of enhancer

elements. For example, in addition to STRs encoding coding CAG-repeats leading to

polyglutamine tracts in genes such as *HTT*, polymorphic STRs with GCA, GCC, GCG or GCT

motifs also encode polyalanine tracts of varying length in a variety of genes (Lavoie et al., 2003).

Other STRs can disrupt introns, such as the expansion of an intronic ATTCT-repeat in

spinocerebellar ataxia type 10 (Matsuura et al, 2000). It is well-established that STR variation is

associated with expression levels of genes both at a single gene level and at a genome-wide

level (Contente et al., 2002; Gymrek et al., 2016; Fotsing et al., 2019). It is unclear the extent to

which STRs are responsible for genomewide association signals detected using SNPs. Linkage

disequilibrium between STRs and SNPs varies, but is expected to be lower than between two

SNPs at the same recombination distance because the high mutation rate of STRs rapidly breaks

down linkage disequilibrium over time. This is shown in attempts to impute STR length genotypes

from SNP genotypes, as biallelic STRs can generally be reliably imputed but multi-allelic STRs

are much less reliably imputed (Saini et al., 2018).

Given the potential importance of STRs in human disease, genome-wide studies examining STR

variation are clearly warranted. The first generation of STR-calling software, including LobSTR

(Gymrek et al., 2012), RepeatSeq (Highnam et al., 2013) and HipSTR (Willems et al., 2017),

aimed to genotype STRs using sequencing reads that completely span the STR. This led to the

limitation that STRs longer than the sequencing read length could not be genotyped. The

second generation of STR-calling software uses information from the distance between the

paired sequencing reads, in addition to direct sequence information across the repeat, to

genotype the STR. This attempts to capture information on STRs that are longer than the

sequencing read length and is particularly effective at identifying large expansions at an STR

(Bahlo et al., 2018).

This study aimed to compare software, suitable for biobank-scale data analysis, that genotypes

common STRs and identifies rarer STR expansions genome-wide, with a view to identifying an

approach to investigate the role of genomic STR variants in polygenic human diseases. We

selected software tools that had at least one of the following characteristics: not limited by read

length, able to estimate STR allele length, or had not been compared previously. Three recent

pieces of software are focused on profiling large repeat expansions: Expansion Hunter De Novo

(EHdn) (Dolzhenko et al., 2020), STRetch (Dashnow et al., 2018) and STRling (Dashnow et al.,

2022). They all use sequence reads that span STRs and mate pair distance information to

identify large expansions, however, unlike STRetch, EHdn and STRling do not require a

predefined catalogue of STRs to genotype an STR locus. Therefore EHdn and STRling can also

identify novel STR loci not assembled in the reference genome. STRetch, STRling and EHdn

can all detect and genotype large repeat expansions in a sample that are outliers from the

population distribution of allele lengths. Some of the key differences between these tools are

summarised in Table 1. Here we assess the ability of STRetch, STRling and EHdn to identity

rarer STR expansions by running them on 116 PCR free WGS containing clinically-validated

trinucleotide repeat expansions, normal repeat sizes and or pre-mutations (Dolzhenko et al.,

2017).

We also compare three software tools that focus on genome-wide genotyping of both short and

expanded (longer than sequencing read length) repeat arrays given a reference catalogue of

STR genomic coordinates: ExpansionHunter,GangSTR and HipSTR. ExpansionHunter and

GangSTR use sequence mate-pair distance information together with STR-spanning sequence

reads. Although HipSTR does not use mate-pair distance information and therefore limited by

sequence read length, a previous study showed that it was shown to outperform its counterparts

(Willems et al., 2017), and therefore was included in this study. These three tools report diploid

allele lengths. In addition, HipSTR genotypes the allele sequence. HipSTR was compared to

both GangSTR and ExpansionHunter to genotyping common STR genome wide. We use

different sequence coverages, read lengths and genomes from different populations to estimate

practical parameters with broad applicability, such as run time, memory usage and proportion of

STRs called genome wide. We use both Mendelian inheritance in parent-offspring trios and

comparison with capillary electrophoresis-based genotypes of forensically-relevant STRs to

estimate the genotyping error rate of different approaches. Both GangSTR and

ExpansionHunter can also identify large STR expansions and were compared against EHdn,

STRling and STRetch to assess their ability to call rarer large STR expansions.

Table 1 in methods summarises difference/similarities of the tools analysed in this study. These

tools have been partially compared in literature by the authors but there is still lack of

comprehensive comparison of the STR calling tools. Two additional tools that call STR

expansions but are excluded from this study are exSTRa and Tredparse. exSTRa does not

estimate the allele length and Tredparse cannot estimate repeat lengths longer than sequencing

mate-pair length (Bahlo et al., 2018, Mousavi et al., 2019)

The genome data analysed here includes those from the Genome in a Bottle consortium and the

1000 Genomes project, which are collections of publicly-available samples. Our aim in this study is

to benchmark different STR-calling software tools primarily using these reference samples. In

particular, the Genome in a Bottle consortium provides key reference standard samples analysed

by many different methods, so as new technical approaches become available we encourage

others to use these same reference samples to facilitate fair comparison of software tools between

studies (Zook et al., 2016).

Methods

Samples and datasets

An overview of the approaches used in this paper is shown in Figure 1, and the different features of

the software tools are shown in Table 1.

Table 1: Software for genotyping tandem repeats compared

Software	Repeat unit size range (bp)	STR catalogue required?	Genomewide?	Approach	Estimates repeat length and/or sequence	Reference
HipSTR v.0.6.2	1-6	Yes	Yes; only < read length	Individual calling	Both	Willems et al., 2017
GangSTR v.2.5.4	1-20 ;20+	Yes	Yes	Individual calling	Only length	Mousavi et al., 2019
ExpansionHunter v.5.0.0	1-6 ;6+	Yes	Yes	Individual calling	Only length	Dolzhenko et., 2017
STRetch	1-6	Yes	yes, only expansions	Case/control - outlier	Only length	Dashnow et al., 2018
STRling	1-6	No	yes, only expansions	Individual calling & Case/control - outlier	Both	Dashnow et al., 2022
ExpansionHunter Denovo	2-20; 20+	No	yes, only expansions	Case/control -outlier	Both	Dolzhenko et al., 2020

The six tools were benchmarked by analysing three different datasets. Firstly, ten gold-standard

high coverage human genome sequences from the Genome in a Bottle consortium

(https://github.com/genome-in-a-bottle/giab_data_indexes) consisting of two child-parent trios.

Secondly, four child-parent trios randomly selected from the 1000 Genomes project

(https://www.internationalgenome.org/data) and thirdly, 116 samples with known trinucleotide

repeat mutations obtained from the Coriell Institute (EGA accession number

EGAD00001003562). The accession numbers for samples obtained from the Genome in a

Bottle (GIAB) and the 1000 Genomes Project are listed in Table 2.

The 116 whole genomes including previously validated disease-causing STR expansions obtained

(Dolzhenko et al., 2017) have been validated with one the following repeat expansions: Fragile X

Syndrome (FMR1), Huntington disease (HTT), Friedreich's ataxia (FXN), Myotonic Dystrophy

(DM1), Spinocerebellar Ataxia 1/3 (ATXN1/3), Spinal and Bulbar Muscular Atrophy (SBMA). The

samples had been sequenced at 2x150 bp reads on Illumina HiSeqX and repeat expansions

previously detected using ExpansionHunter and standard PCR techniques (Dolzhenko et., al

2017). All samples had been Illumina sequenced using PCR-free methods. The sequencing

methods for these data sets have been described in Dolzhenko et al., 2017,

https://doi.org/10.1038/sdata.2016.25 and Byrska-Bishop et al., 2022.

Bam file preparation

The seven GIAB bam files were down-sampled from their original coverage of either 300x or

100x to a final coverage of ~30x. In summary, a total of 5/7 GIAB samples had an initial

genome coverage of ~300x and 2/7 had a genome coverage of ~100x. The 300x genomes

were down-sampled to 100x and all samples were down-sampled further to ~30x genome

coverage using Picard v2.6.0 software. The final 30x coverage was chosen to assess the

applicability of STR genotyping tools on large cohorts of genomes sequenced at 30x genome

coverage, which is a standard coverage adopted by the 1000 Genomes project, for example.

The bam files were refined to remove sequence duplicates and the quality of the bam files

assessed using Qualimap v2.2.1 (Okonechnikov et., al 2016). This data set was collated with

twelve 30x genomes from the 1000G Project (1 KGP). Overall, we generated 3 datasets: 300x

(n = 5), 100x (n = 7) and 30x (n =19) genomes, including 6 child-parent trios, with three

individuals sequenced using both 150bp and 250bp paired-end reads (Table 2). These

sequences had been aligned to the GRCh38 human genome assembly (Zook et al., 2016). All

the bam files were sorted and indexed using samtools v1.9 (Li et., 2009) before calling for STR

genotypes.

Table 2 - Samples used in this study

Sample ID	Population	Trio	Collection	Average fold sequence coverage	Paired-end read length (bp)	Accession number
HG002	Ashkanazi	Son	GIAB	30, 100, 300	150, 250	NA24385
HG003	Ashkanazi	Father	GIAB	30, 100, 300	150, 250	NA24149
HG004	Ashkanazi	Mother	GIAB	30, 100, 300	150, 250	NA24143
HG005	Chinese	Son	GIAB	30, 100, 300	250	NA24631
HG006	Chinese	Father	GIAB	30, 100	150	NA24694
HG007	Chinese	Mother	GIAB	30, 100	150	NA24691
NA12878	CEU	-	GIAB	30, 100, 300	150	NA12878
HG00403	CHS	Father	1000G	30	150	HG00403
HG00404	CHS	Mother	1000G	30	150	HG00404
HG00405	CHS	Son	1000G	30	150	HG00405
NA18485	YRI	Son	1000G	30	150	NA18485
NA18487	YRI	Father	1000G	30	150	NA18487
NA18489	YRI	Mother	1000G	30	150	NA18489
HG01500	IBS	Father	1000G	30	150	HG01500
HG01501	IBS	Mother	1000G	30	150	HG01501
HG01502	IBS	Son	1000G	30	150	HG01502
NA06984	CEU	Father	1000G	30	150	NA06984
NA06989	CEU	Mother	1000G	30	150	NA06989
NA12329	CEU	Daughter	1000G	30	150	NA12329

GIAB=Genome in a Bottle

1000G=1000 Genomes Project

Computing memory and run time evaluation.

To evaluate compute resource usage, we recorded the time and memory taken to process each

bam file from five samples sequenced at 30x coverage, using a single core of an Intel Xeon

Skylake CPU running at 2.6GHz clock speed with 80Gb RAM available. GangSTR, HipSTR,

STRetch and ExpansionHunter were run on a custom STR catalogue containing 811899 STRs of 2-

6 bp repeat units. This catalogue was built from GangSTR's catalogue (hg38_v13.bed) available at

https://github.com/gymreklab/GangSTR, limited to 2-6 bp repeat units. Because ExpansionHunter is

sensitive to the presence of unambiguous 'NNs' in a reference genome around the STR, some of

the loci were dropped leaving a total of 790661 loci. This custom catalogue used is available at :

https://doi.org/10.25392/leicester.data.22041020. GangSTR, HipSTR, and ExpansionHunter were

run at default parameters using this custom catalogue. STRetch was run using whole genome

sequencing pipeline starting from mapped bam files (Dashnow et al., 2018). STRling and

ExpansionHunter Denovo (EHdn) do not require a reference catalogue. STRling was run using

single sample pipeline (Dashnow et al., 2022). Both EHdn and STRetch require a control set of

samples. To explore STR expansion profiles in each sample, for EHdn, the sample to be examined

was treated as a case sample and the rest of the samples as controls and performed outlier

analysis. For STRetch, we build a control set from the remaining subset of the genomes analysed

in this study using STRetch pipeline (Dashnow et al., 2018).

Comparison of software for genotyping STRs

HipSTR, GangSTR and ExpansionHunter performance was compared genome-wide by assessing:

(a) the proportion of STRs genotyped (b) accuracy of the calls made by analysing Mendelian

inheritance patterns in 6 child-parent trios and sample call-concordance compared across varying

sequence depths and sequence read lengths. First, each tool was run using its own STR catalogue

published and tested on the respective tool. These catalogues are of different sizes , with GangSTR

listing 832,380 STR loci, HipSTR listing 1,638,945 loci and ExpansionHunter listing 174,293 loci.

These catalogues are available at: https://github.com/HipSTR-Tool/HipSTR,

https://github.com/gymreklab/GangSTR and https://github.com/Illumina/RepeatCatalogs. HipSTR

and ExpansionHunter catalogues consist of STRs of 2-6 bp units while GangSTR consist of 2-20bp

repeat units. GangSTR was run using default parameters. HipSTR was run using some non-default

parameters : --min-reads 10 and --def-stutter-model. The parameters were used to set the

minimum total reads required to genotype a locus from 100 and allow a default stutter model,

recommended for running few samples (Willems et al., 2017). ExpansionHunter was run at default

parameters, see codes used in data availability section. For comparisons between the three

software tools, they were all run using the custom catalogue of 790661 STRs described above. The

raw variant calling files (VCF) were filtered using dumpSTR tool to remove calls with read coverage

below 10 and those with abnormally high coverage by setting the max call depth for all tools to

1000 (Mousavi et al., 2021). Additional parameters were used for GangSTR: --filter- spanbound-

only and --filter-badCI, to filter out calls where only spanning or bounding reads were found, or

calls with the maximum likelihood genotype estimates outside of the 95% bootstrap confidence

interval as recommended by the GangSTR authors (Mousavi et al 2019). The performance of the

three methods was further assed by comparing the genotype calls to CE data (published in Kim et

al., 2019 for sample NA12878) across 13 forensic STRs, known as core Combined DNA Index

System (CODIS), supplementary file 1. This panel was further extended to include 9 additional

forensic STRs and assessed the accuracy of the 3 software by assessing the mendelian

inheritance patterns in the 5 parent-child trios.

Comparison of software for detecting and genotyping STR expansions

To assess the sensitivity of GangSTR, EHdn, STRling and STRetch at detecting STR

expansions, the four tools were run on samples with known clinical STR expansions that had

been analysed by ExpansionHunter in a previous publication (Dolzhenko, et. al 2017). EHdn was

run in a case- control mode as opposed to outlier analysis (Dolzhenko et al 2020). For each

clinical STR expansion locus, the samples with those STR expansions were treated as cases,

and samples with different STR expansions or normal allele sizes used as controls. For STRetch,

samples were used as a control to each other, and the depth normalised read counts compared

to identify significantly expanded loci in each sample relative to the rest. Both STRetch and

GangSTR were run using STR catalogues containing disease loci

(https://stripy.org/expansionhunter-catalog-creator). GangSTR was run with the same

parameters described above. However, for Fragile X Syndrome and Spinal and Bulbar Muscular

Atrophy, GangSTR was run by specifying the ploidy of the X chromosome for male and female.

STRling was run using joint calling pipeline (Dashnow et al., 2022). Our results were compared to

the previously reported validated calls for these samples (Dolzhenko et., al 2017).

Data availability

Code for analyses, and the full set of genotype calls at the clinical and forensic loci are available

at https://doi.org/10.25392/leicester.data.22041020.

Genotype call vcf files for GangSTR and HipSTR and ExpansionHunter are available for the

Genome In a bottle samples are at https://ftp-trace.ncbi.nlm.nih.gov/ReferenceSamples/giab/data/

and https://doi.org/10.25392/leicester.data.22041020

Genotype call vcf files for GangSTR, ExpansionHunter and HipSTR are available for the 1000

Genomes samples used are at https://doi.org/10.25392/leicester.data.22041020

Figure 1 - Flowchart of analyses and samples used in this study

a)   Genome in a bottle (GIAB) samples

b)   1000 Genomes project samples

c)   Known clinical samples available from Coriell

..

Results

Computing resource usage

For large scale studies involving thousands of samples, the relative processing time of different

software tools can become an important aspect of software choice. We first compared the

software tools that use a STR catalogue (Table 3). GangSTR, HipSTR, ExpansionHunter and

STRetch were run on the same STR catalogue containing 790661 loci. GangSTR took an average

of 10 hours per diploid genome across five samples, STRetch took about 7 hours and HipSTR

took about 3 hours. ExpansionHunter default mode (seeking mode) needed more than 7 days and

was not run to completion. For the software tools that do not require a software catalogue, EHdn

and STRling took around 30 minutes per diploid genome.

STRetch and ExpansionHunter are the only software tools tested with a multithreading option. For

STRetch increasing the core number from 1 to 28 reduces the time from 7 hours to about 1.5

hours per diploid genome. Using ExpansionHunter's streaming mode (that is recommended for

large genomic catalogs) with at least 16 threads noticeably improved its performance, reducing its

run time to an average of 1.5 hours per diploid genome (Table 3). ExpansionHunter was notable

for its high memory usage – typically around 70Gb in contrast to the other 5 software tools using

either ~17 Gb or less than 1GB (Table 3).

Table 3. Comparison of processing times and memory usage of STR-calling software

	CPU Time (hours:minutes:seconds)							RAM in Gb						
Sample	GangSTR	HipSTR	STRetch	EH (default mode)	EH (streaming mode)	EHdn	STRling	GangSTR	HipSTR	STRetch	EH (default mode)	EH (streaming mode)	EHdn	STRling
NA18485	10:33:01	3:58:53	06:39:38	> 168 h	01:33:22	00:30:50	00:24:33	0.04	0.50	17.63	n/a	69.80	0.57	1.16
NA18487	10:27:03	3:00:32	07:10:47	> 168 h	01:28:28	00:30:09	00:24:03	0.03	0.49	17.60	n/a	72.56	0.48	0.87
NA06984	10:13:42	2:47:50	06:21:20	> 168 h	01:23:40	00:30:33	00:24:19	0.04	0.40	17.42	n/a	70.11	0.58	1.16
NA06989	8:41:14	2:34:05	05:57:05	> 168 h	01:17:33	00:27:24	00:22:11	0.04	0.42	17.51	n/a	69.82	0.57	0.82
NA12329	10:24:04	2:50:49	07:15:59	> 168 h	01:19:26	00:29:49	00:23:42	0.05	0.34	17.40	n/a	72.57	0.51	0.76
Average	10:03:48	03:02:25	06:40:57	n/a	01:24:29	00:29:45	00:23:45	0.04	0.43	17.51	n/a	70.97	0.54	0.96

Number of common STR loci genotyped and genotype call concordance of common STRs.

We compared GangSTR, ExpansionHunter and HipSTR, which are the three software tools that

aim to genotype all STRs across the genome. They all rely on a catalogue of STR loci provided by

the user, and the number of the loci in that catalogue provide the upper limit on the number of

genotyped STRs. Therefore, our results are presented as a percentage of STRs called using that

particular catalogue. We first used the different catalogues provided with each software tool, and

found several patterns common to both GangSTR, ExpansionHunter and HipSTR (Supplementary

table 1). Firstly, all software tools call a high proportion of the STRs in their respective catalogues

across all samples, with GangSTR and ExpansionHunter calling a higher proportion than HipSTR,

although the different sizes of their catalogues must be taken into account. Secondly, more calls

are made from 100x coverage data than 30x. The increase is small across most samples but is

large (over 10 percentage points) for HG005 for HipSTR and GangSTR, where both tools call fewer

than 90% of STRs in the 30x coverage data. Taken together, 30x is not only sufficient but appears

optimal for STR calling, with little benefit in increasing sequencing depth. For read length, 150bp

paired-end is optimal and increased read length seems slightly detrimental, at least for HipSTR,

which made fewer calls across the four samples that were sequenced at 2x250bp.

To ensure a fair comparison across tools, we repeated our analysis using the same STR catalogue

containing 790661 loci (Table 4). For all software tools and samples, 90% of STRs in this catalogue

were called, with most calling >99%. We compared the actual genotype calls made on the STRs

called by three tools by plotting each allele call for each on a scatterplot, normalised against the

allele in the reference genome (Figure 2). This shows that over 85% of the calls were identical

between the three methods, and of those calls, at least 89.0% were identical with the reference

genome. There was no strong bias in under- or over-calling repeat number made by HipSTR,

ExpansionHunter or GangSTR. There was high concordance of the genotype calls made at

2x150bp compared to 2x250bp for each tool (Supplementary table 1).

Table 4 – Percentage of loci called in a shared STR catalogue by GangSTR and HipSTR and ExpansionHunter

Sample	Read length	GangSTR					HipSTR					ExpansionHunter				
		Called at 30x (%)	Called at 100x (%)	Calls in both (%)	Identical by allele length (%)	Identical by allele sequence (%)	Called at 30x (%)	Called at 100x (%)	Calls in both (%)	Identical by allele length (%)	Identical by allele sequence (%)	Called at 30x (%)	Called at 100x (%)	Calls in both (%)	Identical by allele length (%)	Identical by allele sequence (%)
HG002	2x150	99.7	99.7	99.6	99.4	NA	96.2	96.0	95.5	99.8	99.9	99.8	99.9	99.8	98.9	NA
HG003	2x150	99.7	99.7	99.6	99.4	NA	96.1	96.0	95.5	99.8	99.9	99.8	99.9	99.8	99.8	NA
HG004	2x150	99.2	99.2	99.1	99.4	NA	95.8	95.5	95.2	99.8	99.9	99.3	99.5	99.3	98.9	NA
HG005	2x250	98.5	99.6	98.4	99.6	NA	93.9	94.3	92.9	99.7	99.1	99.8	99.9	99.8	98.4	NA
HG006	2x150	99.4	99.5	99.3	99.3	NA	96.0	95.9	95.0	99.8	99.9	99.7	99.9	99.7	98.7	NA
HG007	2x150	99.0	99.1	98.9	99.3	NA	95.7	95.4	95.0	99.8	99.8	99.3	99.5	99.3	98.7	NA
NA12878	2x150	99.1	99.1	98.9	99.3	NA	95.7	95.4	95.0	99.8	99.8	99.3	99.5	99.3	98.8	NA
HG002	2x250	96.1	NA	NA	NA	NA	94.1	NA	NA	NA	NA	99.3	NA	NA	NA	NA
HG003	2x250	95.8	NA	NA	NA	NA	93.9	NA	NA	NA	NA	99.8	NA	NA	NA	NA
HG004	2x250	96.8	NA	NA	NA	NA	93.9	NA	NA	NA	NA	99.4	NA	NA	NA	NA

NA: not analysed

Figure 2 – Comparison of STR calls from ExpansionHunter, HipSTR and GangSTR

A). Comparison of the HG002 genotypes of the STRs called by both GangSTR and HipSTR. The

x-axis represents the call made by GangSTR in comparison to the reference sequence (0=same

as reference sequence, -100 represents 100 fewer repeat units than the reference sequence.

The y-axis represents the call made by HipSTR. B) Comparison of the genotypes of the STRs

called by either GangSTR or HipSTR compared to ExpansionHunter (x-axis). The dotted red

lines shows correlation between the calls compared.

Accuracy of common STR calls assessed using Mendelian inheritance

Accuracy of the genotypes made by GangSTR, ExpansionHunter and HipSTR can be measured

using the expectation of Mendelian inheritance of alleles, whereby we would expect to see one

allele of a child's genotype in their mother and one allele in their father. Any allele in a child not

observed in one of the parents is due either to a genotyping error in the child or the parent, or a

de novo mutation in the child. Assuming a rate of de novo mutations at STR loci as 10^{-4} per

generation, (Willems et al., 2016), the contribution to observed inheritance errors of de novo

mutation is minimal and therefore differences in inconsistencies can be attributed to differences

in STR genotyping accuracy.

Analysis of Mendelian inconsistencies using bcftools software across the five mother-father-child

trios with 150bp sequence reads shows very little difference in genotyping accuracy across the

three methods. GangSTR and HipSTR showed the highest genotyping accuracy (table 5).

Comparison between the HG002,HG003,HG004 trio sequenced at 100x coverage and 30x

coverage shows that increased sequencing coverage does not improve genotyping accuracy.

Table 5 – Mendelian inheritance of STR alleles

Parent-offspring Trio	Coverage	GangSTR Mendelian consistent (%)	HipSTR Mendelian consistent (%)	ExpansionHunter Mendelian consistent (%)
HG002,HG003,HG004	100x	99.5	99.6	99.2
HG002,HG003,HG004	30x	99.7	99.5	98.7
NA18485,NA18489,NA18487	30x	99.6	99.4	98.5
NA06984,NA06989,NA12329	30x	99.8	99.9	98.9
HG00403,HG00404,HG00405	30x	99.9	99.9	99.0
HG01500,HG01501,HG01502	30x	99.7	99.6	98.7

In order to investigate the possible reasons for erroneous genotype calls, we stratified the

genotypes that showed Mendelian inconsistencies by repeat unit size normalised against the

repeat-unit counts in the reference list (Figure 3). It is clear that for the three tools;

ExpansionHunter, HipSTR and GangSTR, there is overrepresentation of 2 bp repeat unit STRs

in the incorrect genotype calls. Therefore, for these methods 2 bp repeat STR genotypes have

the highest error rate, over twice the error rate of other STRs (supplementary figure 1).

Genotyping of known forensic STR loci

Thirteen STRs, known as core Combined DNA Index System (CODIS) STRs, are long STRs that

show extensive variation between individuals, and are used across many different forensic STR

panels for identification of individuals from genomic DNA (Butler, 2006; Brearley et al., 2020).

Analysis of the performance of GangSTR, ExpansionHunter and HipSTR in calling these STRs in

comparison to capillary electrophoresis (CE) data is useful both for practical forensic analysis, but

also as a measure of the error rate of the software, if we accept that capillary electrophoresis can

be regarded as a gold standard. Matched capillary electrophoresis data for the 13 core STRs,

generated using Promega Powerplex Fusion 24 assay, has been published (Kim et al., 2019),

and our calls for NA12878 were compared against these data (table 6). ExpansionHunter calls

matched CE data at 10/13 loci, called one allele incorrectly at the THO1 locus and two alleles at

both FGA and D21S11 loci. GangSTR genotypes matched CE data for 11/13 loci, with one locus

(D21S11) not called and one (TH01) called incorrectly. HipSTR showed the same results as

GangSTR, except that, in addition, it incorrectly called one allele at D13S317.The three tools were

also consistent across 19/22 forensic STRs analysed for Mendelian inheritance errors across the

six trios (Supplementary file 1).

Detection of STR expansions at known clinical loci

To assess the reliability in calling expanded STRs in a clinical situation, we compared STRetch,

EHdn and GangSTR performance in calling expanded STRs in samples with known STR

expansions (Table 7). All four methods showed high sensitivity, with EHdn (95% sensitivity),

GangSTR (89% sensitivity) and STRling (94% sensitivity) outperforming STRetch (68%

sensitivity). Although STRetch, STRling and GangSTR were able to flag repeat expansions in

most of the genes analysed, the three tools underestimated the repeat lengths in comparison to

Southern blot results at *DMPK*, *FMR1* and *FXN* loci (Dolzhenko et al., 2017). These loci were

characterised by longer and or double expansions in both alleles ranging between 50 to 1000

repeat units in the samples that were screened (supplementary file 1). Both STRetch and STRling

performed poorly at the *FMR1* locus (Table 7). Of the samples identified as expanded (p<=0.05;

adjusted for multiple testing) at the *FMR1* locus by STRetch, all had reported repeat lengths

below the expected *FMR1* premutation ranges (Hannan et al., 2018). For STRling, half the

samples identified had reported repeat lengths below the expected *FMR1* premutation ranges. To

assess specificity, 21 samples with known non-pathogenic allele length at *FMR1* were analysed,

with STRling called two of these as expanded and GangSTR identifying one sample with an

expanded allele length (supplementary file 2). This is a small dataset to robustly test false positive

rate, but suggests overall high specificity for the three software tools used.

Figure 3 – Repeat unit size distribution of Mendelian inconsistencies

a) ExpansionHunter, GangSTR and HipSTR calls that are Mendelian inconsistent as a proportion

of motif counts in the catalogue, b) all STRs in the catalogue used by the three software tools.

417
418
419

Table 6 - Comparison of STR genotype calls at core forensic loci for NA12878

Locus	Chrom	Start	End	Motif	GangSTR 100x	GangSTR 30x	HipSTR 100x	HipSTR 30x	ExpansionHunter 100x	ExpansionHunter 30x	CE
CSF1PO	5	150076324	150076375	ATCT	10,11	10,11	10,11	10,11	10,11	10,11	10,11
D5S818	5	123775556	123775599	ATCT	12,12	12,12	12,12	12,12	12,12	12,12	12,12
D7S820	7	84160226	84160277	TATC	8,10	8,10	8,10	8,10	8,10	8,10	8,10
D13S317	13	82148025	82148068	TATC	11,12	11,12	11.3,12.3	11.3,12.3	11,12	11,12	11,12
D16S539	16	86352702	86352745	GATA	10,11	10,11	10,11	10,11	10,12	10,11	10,11
D21S11	21	19181973	19182099	TCTA	-	-	-	-	18,34	34,34	30,30
TH01	11	2171088	2171115	AATG	7,8	7,7	7,9.8	7,9.8	7,10	7,10	7,9.3
TPOX	2	1489653	1489684	AATG	8,8	8,8	8,8	8,8	8,8	8,8	8,8
vWA	12	5983977	5984044	AGAT	15,17	15,15	15,17	15,17	15,17	15,17	15,17
D3S1358	3	45540739	45540802	TCTA	16,17	16,17	16,17	16,17	16,17	16,17	16,17
D8S1179	8	124894865	124894916	TATC	12,12	12,12	12,12	12,12	12,13	12,13	12,12
D18S51	18	63281667	63281738	AGAA	16,17	16,17	16,17	16,17	16,17	16,17	16,17
FGA	4	154587736	154587823	GGAA	22,24	22,24	22,24	22,24	23,25	23,25	22,24

N/B: F=failed QC; CE=capillary electrophoresis genotype

Table 7 - Detection of STR expansions at known clinical loci

Disease	Gene	Total analysed	Identified using EHdn	Identified using STRetch	Identified using GangSTR	Identified using STRling
Spinal-bulbar muscular atrophy	*AR*	1	0	0	0	0
Myotonic dystrophy type 1	*DMPK*	16	16	16	16	16
Fragile X	*FMR1*	34	34	19	32	33
Friedreich's ataxia	*FXN*	25	25	11	19	24
Huntington's disease	*HTT*	13	13	13	13	12
Dentatorubral-pallidoluysian atrophy	*ATN1*	2	2	2	1	2
Spinocerebellar ataxia type 1	*ATXN1*	3	0	3	3	1
Spinocerebellar ataxia type 3	*ATXN3*	1	0	1	1	1
Total		95	90 (95%)	65 (68%)	85 (89%)	89 (94%)
Unaffected		21	0	0	1	2

Discussion

Genomewide analysis of STRs using short read sequencing has lagged behind studies of other

variations, including single nucleotide variation and, to a certain extent, structural variation. It is

likely that STRs underly some undiscovered genetic associations with complex disease (Hannan

2010). Several software tools have been developed with the aim of accurately calling STR

genotypes genome-wide, but with different ultimate aims. Some have been developed to detect

and genotype large repeat expansions that are outliers from the population distribution of allele

lengths for that particular STR. Development of these was motivated by well- established repeat

expansions causing a variety of Mendelian diseases, and more recent discoveries showing that

large repeat expansions can underlie a large amount of complex disease, such as ALS

(DeJesus-Hernandez et al., 2011; Renton et al., 2011). Other software tools aim to genotype

STRs genome-wide irrespective of their alleles, using a catalogue of known STRs.

We selected the most recent software tools most appropriate to our needs in detecting both STR

expansions and genome-wide STR genotypes in a population-based sample. A previous study

describes detection of six disease-causing STR expansions from whole exome sequences using

several STR-calling software tools, including GangSTR and STRetch but not EHdn and STRling

(Rajan-Babu et al., 2021). ExpansionHunter (Dolzhenko et al., 2019), STRetch and exSTRa

(Tankard et al., 2018) were used to detect clinically-related STR expansions in those 6 genes with

high specificity. Other reports include those focused on forensic STR detection (Valle-Silva et al.,

2022), or are focused on genotyping a particular, or small number, of STR loci (Budiš et al., 2019).

Our aim was to assess the feasibility of applying these software tools, both in terms of quality of

the resulting data and practicality of using the software, for large cohorts of genomes. We

assessed this in a variety of ways. Processor and memory usage, in conjunction with processing

time, was measured so that we could assess the feasibility of scaling up analysis to thousands of

genomes given current computing resources. For the software using a catalogue to genotype

known STRs (GangSTR, HipSTR and Expansionhunter), we examined the effect of increasing

sequencing depth on the proportion of STRs genotyped, and accuracy. We then examined the

concordance between GangSTR, ExpansionHunter and HipSTR using common STR catalogues,

and the relative number of Mendelian inconsistencies in trios to assess STR genotyping accuracy.

We found sequencing depth to have no effect on the number or quality of STR genotypes called

by HipSTR, but higher sequencing depth had a modest positive effect on GangSTR STR genotype

calling. Quality of STR calls made by Expansionhunter, GangSTR and HipSTR was very similar

but slightly higher between GangSTR and HipSTR when given the same STR catalogue. Both

GangSTR and HipSTR used less memory than Expansionhunter, but GangSTR took about 3x

more CPU time. Both GangSTR and HipSTR performed very well, but not perfectly, in genotyping

STR loci used for forensics.

The software tools STRetch, STRling, ExpansionHunter, EHdn and GangSTR can all call

expanded repeats, with only EHdn and STRling not requiring a predefined catalogue of STR loci.

Because expanded repeats are expected to be rare, and underlie several clinical conditions, we

used a previous dataset to test the ability of STRetch, STRling, EHdn and GangSTR to genotype

known STR expansions. EHdn, STRling and GangSTR show higher sensitivity than STRetch, at

least under the conditions tested. EHdn and STRling used the least resource, analysing a genome

in about half an hour using about half a Gb of memory under our conditions.

For case-control studies where thousands of genomes have been sequenced at ~30x coverage

for both large expansions and known STRs genomewide, we have decided to use GangSTR for

known STRs and EHdn for large expansions. Although STRling used similar time and memory as

EHdn when run in a single sample mode, its resource usage can increase linearly in joint calling

mode which is appropriate for case-control outlier analysis (Dashnow *et al*, 2022). EHdn and

STRling are the only STR genotypers that can genotype expansions without a prior defined

catalogue, broadening their scope to identify previously unknown expansions. GangSTR and

HipSTR are similar, with the increased computing resources needed by GangSTR offset in our

view by its ability to detect larger STR expansions longer than the read length, to support and

extend EHdn calls. ExpansionHunter default mode was unusable at genotyping a catalog of

790661 STR loci as it needed more than 7 days but efficient in streaming mode albeit higher

memory requirements about 70 Gb per genome using 16 cores.

We note that, although the clinical repeat expansions and the forensic STRs have been validated

by orthogonal data and methods, this is not the case with the other STR genotypes. For STRs

that are associated with disease in subsequent analyses, it will be important to validate those

particular STRs on a subset of samples using alternative methods, such as capillary

electrophoresis. STR genotype calling from high throughput sequencing remains an area of active

development, and we hope that further progress can be made in reliably calling STRs from both

short read and long read sequencing data.

Our study had some limitations inherent to different tools. First, we did not analyse the 1bp repeat

unit motifs. Both GangSTR and EHdn have been optimised to genotype 2-20bp repeat motifs, of

which EHdn only genotypes expanded repeats. 1bp repeat motifs present a challenge to many

sequencing technologies due to PCR stutter noise and may introduce artefacts in sequence,

however, a detailed assessment of accuracy of 1bp repeat motif STR genotyping in PCR-free

sequencing is needed. Secondly, apart from EHdn and STRling, the tools require defined STR

coordinates built from reference genomes. Therefore, STRs not assembled in the reference

genome, including the highly complex regions of the genome that are often hard to sequence and

assemble, cannot be genotyped. However, with improvements in PCR free protocols and long

read sequencing, combined with new genome assemblies, the ability to genotype these regions

will improve.

Acknowledgements

JWO is funded by a Wellcome Trust PhD studentship as part of the Wellcome Trust Genetic

Epidemiology and Public Health Genomics Doctoral Training Programme by grant number

218505/Z/19/Z . LWV holds a GSK/Asthma+Lung UK Chair in Respiratory Research (C17-1). The

research was partially supported by the National Institute for Health Research (NIHR) Leicester

Biomedical Research Centre; the views expressed are those of the author(s) and not necessarily

those of the National Health Service (NHS), the NIHR or the Department of Health. This research

used the ALICE High Performance Computing Facility at the University of Leicester.

Ethics statement

All sequencing data used in this study are from previous studies, from fully consented individuals.

Sample collections are from the 1000 Genomes project

(https://www.internationalgenome.org/sample_collection_principles/) , Personal Genome Project

Canada (https://personalgenomes.ca/) , or NIGMS Human Genetic Cell Repository at the Coriell

Institute for Medical Research.

Authorship statement

All authors conceived and designed the study. JWO sourced data and carried out the analysis.

EJH and JWO wrote the paper, with input from all authors.

Conflict of Interest statement

The authors declare no conflict of interest.

References

Bahlo M, Bennett MF, Degorski P, Tankard RM, Delatycki MB, Lockhart PJ. 2018. Recent

advances in the detection of repeat expansions with short-read next-generation sequencing.

F1000Research 7:736

Brearley EJ, Singh P, Bhatti JS, Mastana S. 2020. Genetic variation and differentiation among a

native British and five migrant South Asian populations of the East Midlands (UK) based on

CODIS forensic STR loci. Ann Hum Biol 47:572–583.

Brinkmann B, Klintschar M, Neuhuber F, Hühne J, Rolf B. 1998. Mutation rate in human

microsatellites: influence of the structure and length of the tandem repeat. Am J Hum Genet

62:1408–1415.

Budiš J, Kucharík M, Ďuriš F, Gazdarica J, Zrubcová M, Ficek A, Szemes T, Brejová B,

Radvanszky J. 2019. Dante: genotyping of known complex and expanded short tandem

repeats. Bioinformatics 35:1310–1317.

Burgarella C, Navascués M. 2011. Mutation rate estimates for 110 Y-chromosome STRs combining

population and father–son pair data. Eur J Hum Genet 19:70–75.

Butler JM. 2006. Genetics and genomics of core short tandem repeat loci used in human

identity testing. J Forensic Sci 51:253–265.

Byrska-Bishop M, Evani US, Zhao X, Basile AO, Abel HJ, Regier AA, Corvelo A, Clarke

WE, Musunuri R, Nagulapalli K, Fairley S, Runnels A, Winterkorn L, Lowy E; Human

Genome Structural Variation Consortium; Flicek P, Germer S, Brand H, Hall IM, Talkowski

ME, Narzisi G, Zody MC. 2022. High-coverage whole-genome sequencing of the

expanded 1000 Genomes Project cohort including 602 trios. Cell. 185(18):3426-3440.e19.

Contente A, Dittmer A, Koch MC, Roth J, Dobbelstein M. 2002. A polymorphic microsatellite that

mediates induction of PIG3 by p53. Nat Genet 30:315–320.

Dashnow H, Lek M, Phipson B, Halman A, Sadedin S, Lonsdale A, Davis M, Lamont P, Clayton

JS, Laing NG, MacArthur DG, Oshlack A. 2018. STRetch: detecting and discovering pathogenic

short tandem repeat expansions. Genome Biol 19:121.

DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, Nicholson

AM, Finch NA, Flynn H, Adamson J, Kouri N, Wojtas A, et al. 2011. Expanded GGGGCC

hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and

ALS. Neuron 72:245–256.

Dolzhenko E, Bennett MF, Richmond PA, Trost B, Chen S, Vugt JJFA van, Nguyen C, Narzisi G,

Gainullin VG, Gross AM, Lajoie BR, Taft RJ, et al. 2020. ExpansionHunter Denovo: a

computational method for locating known and novel repeat expansions in short-read sequencing

data. Genome Biol 21:102.

Dolzhenko E, Deshpande V, Schlesinger F, Krusche P, Petrovski R, Chen S, Emig-Agius D,

Gross A, Narzisi G, Bowman B, Scheffler K, Vugt JJFA van, et al. 2019. ExpansionHunter: a
sequence-graph-based tool to analyze variation in short tandem repeat regions. Bioinformatics
35:4754–4756.

Dolzhenko E, Vugt JJFA van, Shaw RJ, Bekritsky MA, Blitterswijk M van, Narzisi G, Ajay SS,
Rajan V, Lajoie BR, Johnson NH, Kingsbury Z, Humphray SJ, et al. 2017. Detection of long
repeat expansions from PCR-free whole-genome sequence data. Genome Res 27:1895–1903.

Fotsing SF, Margoliash J, Wang C, Saini S, Yanicky R, Shleizer-Burko S, Goren A, Gymrek M.
2019. The impact of short tandem repeat variation on gene expression. Nat Genet 51:1652– 1659.

Gymrek M. 2017. A genomic view of short tandem repeats. Curr Opin Genet Dev 44:9–16.

Gymrek M, Golan D, Rosset S, Erlich Y. 2012. lobSTR: A short tandem repeat profiler for
personal genomes. Genome Res 22:1154–1162.

Gymrek M, Willems T, Guilmatre A, Zeng H, Markus B, Georgiev S, Daly MJ, Price AL, Pritchard
JK, Sharp AJ, Erlich Y. 2016. Abundant contribution of short tandem repeats to gene expression
variation in humans. Nat Genet 48:22–29.

Hannan AJ. 2010. Tandem repeat polymorphisms: modulators of disease susceptibility and
candidates for 'missing heritability'. Trends Genet.26:59-65.

Hannan AJ. 2018. Tandem repeats mediating genetic plasticity in health and disease. Nat Rev
Genet 19:286–298.

Hannan AJ. 2020. Expanding genes, repeating themes and therapeutic schemes: The
neurobiology of tandem repeat disorders. Neurobiol Dis 144:105053.

Highnam G, Franck C, Martin A, Stephens C, Puthige A, Mittelman D. 2013. Accurate human
microsatellite genotypes from high-throughput resequencing data using informed error profiles.
Nucleic Acids Res 41:e32.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37:907–915.

Kuilenburg ABP van, Tarailo-Graovac M, Richmond PA, Drögemöller BI, Pouladi MA, Leen R,
Brand-Arzamendi K, Dobritzsch D, Dolzhenko E, Eberle MA, Hayward B, Jones MJ, et al. 2019.
Glutaminase Deficiency Caused by Short Tandem Repeat Expansion in GLS. N Engl J Med
380:1433–1441.

Lavoie H, Debeane F, Trinh Q-D, Turcotte J-F, Corbeil-Girard L-P, Dicaire M-J, Saint-Denis A,
Pagé M, Rouleau GA, Brais B. 2003. Polymorphism, shared functions and convergent evolution
of genes with sequences coding for polyalanine domains. Hum Mol Genet 12:2967–2979.

Lieberman AP, Shakkottai VG, Albin RL. 2019. Polyglutamine Repeats in Neurodegenerative
Diseases. Annu Rev Pathol 14:1–27.

Matsuura T, Yamagata T, Burgess DL, Rasmussen A, Grewal RP, Watase K, Khajavi M,
McCall AE, Davis CF, Zu L, Achari M, Pulst SM, et al. 2000. Large expansion of the ATTCT

pentanucleotide repeat in spinocerebellar ataxia type 10. Nat Genet. 26:191-4.

Mousavi N, Shleizer-Burko S, Yanicky R, Gymrek M. 2019. Profiling the genome-wide landscape

of tandem repeat expansions. Nucleic Acids Res 47:e90.

Paulson H. 2018. Chapter 9 - Repeat expansion diseases. In: Geschwind DH, Paulson HL,

Klein C, editors. Handbook of Clinical Neurology, Elsevier, p 105–123.

Rajan-Babu I-S, Peng JJ, Chiu R, Birch P, Couse M, Guimond C, Lehman A, Mwenifumbo J,

Karnebeek C van, Friedman J, Adam S, Souich CD, et al. 2021. Genome-wide sequencing as a

first-tier screening test for short tandem repeat expansions. Genome Med 13:126.

Renton AE, Majounie E, Waite A, Simón-Sánchez J, Rollinson S, Gibbs JR, Schymick JC,

Laaksovirta H, Swieten JC van, Myllykangas L, Kalimo H, Paetau A, et al. 2011. A

hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-

FTD. Neuron 72:257–268.

Saini S, Mitra I, Mousavi N et al. 2018. A reference haplotype panel for genome-wide imputation

of short tandem repeats. Nat Commun 9, 4397.

Tankard RM, Bennett MF, Degorski P, Delatycki MB, Lockhart PJ, Bahlo M. 2018. Detecting

Expansions of Tandem Repeats in Cohorts Sequenced with Short-Read Sequencing Data. Am J

Hum Genet 103:858–873.

Trost B, Engchuan W, Nguyen CM, Thiruvahindrapuram B, Dolzhenko E, Backstrom I, Mirceta

M, Mojarad BA, Yin Y, Dov A, Chandrakumar I, Prasolava T, et al. 2020. Genome-wide

detection of tandem DNA repeats that are expanded in autism. Nature 586:80–86.

Valle-Silva G, Frontanilla TS, Ayala J, Donadi EA, Simões AL, Castelli EC, Mendes-Junior CT.

2022. Analysis and comparison of the STR genotypes called with HipSTR, STRait Razor and

toaSTR by using next generation sequencing data in a Brazilian population sample. Forensic Sci

Int Genet 58:102676.

Willems T, Gymrek M, Highnam G, 1000 Genomes Project Consortium, Mittelman D, Erlich Y.

2014. The landscape of human STR variation. Genome Res 24:1894–1904.

Willems T, Gymrek M, Poznik GD, Tyler-Smith C, 1000 Genomes Project Chromosome Y

Group, Erlich Y. 2016. Population-Scale Sequencing Data Enable Precise Estimates of Y-STR

Mutation Rates. Am J Hum Genet 98:919–933.

Willems T, Zielinski D, Yuan J, Gordon A, Gymrek M, Erlich Y. 2017. Genome-wide profiling of

heritable and de novo STR variations. Nat Methods 14:590–592.

Zook JM, Catoe D, McDaniel J, Vang L, Spies N, Sidow A, Weng Z, Liu Y, Mason CE,

Alexander N, Henaff E, McIntyre ABR, et al. 2016. Extensive sequencing of seven human

genomes to characterize benchmark reference materials. Sci Data 3:160025.

644