MHC class I Ligands of Rhesus Macaque Killer-Cell Immunoglobulin-Like Receptors

Jennifer L. Anderson[*], Kjell Sandstrom[*], Willow R. Smith[*], Molly Wetzel[*], Vadim A. Klenchin[*] and David T. Evans[*,†]

[*]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison, Madison, WI

[†]Wisconsin National Primate Research Center, University of Wisconsin-Madison, Madison, WI

Corresponding author:

David T. Evans

dtevans2@wisc.edu

Phone: (608) 265-1485

Fax: (608) 265-8084

Running Title: Rhesus Macaque KIR Ligands

Abstract

Definition of MHC class I ligands of rhesus macaque KIRs is fundamental to NK cell biology in

this species as an animal model for infectious diseases, reproductive biology, and transplantation.

To provide a more complete foundation for studying NK cell responses, rhesus macaque KIRs

representing common allotypes of lineage II *KIR* genes were tested for interactions with MHC

class I molecules representing diverse *Mamu-A, -B, -E, -F, -I* and *-AG* alleles. KIR-MHC class I

interactions were identified by co-incubating reporter cell lines bearing chimeric KIR-CD3ζ

receptors with target cells expressing individual MHC class I molecules and were corroborated

by staining with KIR IgG-Fc fusion proteins. Ligands for 11 KIRs of previously unknown

specificity were identified that fell into two general categories: interactions with multiple Mamu-

Bw4 molecules or with Mamu-A-related molecules, including several allotypes of Mamu-AG

and the hybrid Mamu-B*045:03 molecule. Although both groups include inhibitory and

activating receptors, the majority of KIRs found to interact with Mamu-Bw4 are inhibitory,

whereas most of the KIRs that interact with Mamu-AG are activating. We also identified Mamu-

A1*012:01 as a ligand for KIR3DLw03*002, which belongs to a phylogenetically distinct group

of macaque KIRs with a three amino acid deletion in D0 that is also present in human

KIR3DL1/S1 and KIR3DL2. This study more than doubles the number of rhesus macaque KIRs

with defined MHC class I ligands and identifies novel interactions with Mamu-AG, -B*045, and

-A1*012. These findings support overlapping, but nonredundant, patterns of ligand recognition

that reflects extensive functional diversification of these receptors.

Introduction

Natural killer cell responses in humans and other primate species are regulated in part by interactions between two highly polymorphic sets of molecules; the killer cell Ig-like receptors (KIRs) expressed on the surface of NK cells and their major histocompatibility complex (MHC) class I ligands on target cells (1, 2). As indicated by their nomenclature, KIRs typically have two or three extracellular Ig-like domains (2D or 3D) and either long (L) or short (S) cytoplasmic tails. KIRs with long cytoplasmic tails contain a pair of ITIMs that transduce inhibitory signals, whereas those with short cytoplasmic tails associate with adaptor molecules such as DAP12 or FcεRIγ to transduce activating signals (2, 3). Inhibitory KIRs normally suppress NK cell responses through interactions with MHC class I ligands on the surface of heathy cells. However, if these interactions are somehow perturbed, for instance as a result of MHC class I downregulation in virus-infected cells or deletion of MHC class I genes in tumor cells, the loss of these inhibitory signals can trigger an NK cell response (4-9). Although the molecular mechanisms are not as well defined, activating KIRs can also interact with MHC class I molecules on unhealthy cells to directly trigger an NK cell response (2, 10-12).

Polymorphisms in the *KIR* and *HLA class I* genes have been associated with differences in the course of infection for a number of human viral pathogens and with the outcome of cancer immunotherapy (12-19). *KIR* and *HLA class I* polymorphisms have also been linked to complications during pregnancy and to the success of hematopoietic stem cell and solid organ transplantation (20-23). However, studies to address the immunological mechanisms underlying these observations have been hampered by the lack of a suitable animal model. Mice and other rodents do not have KIRs, but instead express C type lectin-like molecules on their NK cells as polymorphic receptors for MHC class I ligands (2, 24). Moreover, due to the rapid pace of KIR

and MHC class I evolution (25-27), it is not possible to predict KIR ligands in non-human

primates based on sequence similarity with human KIRs.

In contrast to humans, which have three classical *HLA class I* genes (*HLA-A, -B* and *-C*),

Old World monkeys express an expanded array of polymorphic MHC class I genes related to

HLA-A and *-B*. However, these species do not have an *HLA-C* ortholog, since *MHC-C* arose in

hominids as a duplication of an *MHC-B* gene after their divergence from Old World monkeys

(28, 29). The *MHC class I* haplotypes of Old World monkeys are polygenic, and rhesus

macaques typically express two or three <u>*Macaca mulatta (Mamu)-A*</u> genes and 4-11 *Mamu-B*

genes (30-32). Depending on their transcriptional abundance, the *MHC class I* genes of

macaques can be further classified as "major" or "minor" loci (32). Products of the major genes

exhibit the greatest polymorphism and present peptides to virus-specific $CD8^+$ T cells (33),

suggesting functional similarities to HLA-A and -B in humans.

Four non-classical *MHC class I* genes have also been identified in Old World monkeys.

Orthologs of *HLA-E* and *-F* (*Mamu-E* and *-F* in rhesus macaques) are well conserved and serve

similar functions as their human counterparts (34-37). Macaques and other Old World monkeys

also have an ortholog of *HLA-G*; however the *MHC-G* genes of these species have accumulated

multiple stop-codons and no longer encode functional proteins (38, 39). In the absence of a

functional *MHC-G* gene, Old World monkeys have evolved duplicated *MHC-A* genes, termed

MHC-AG (*Mamu-AG* in rhesus macaques), that appear to serve a similar function (40-43). Like

HLA-G, Mamu-AG has limited polymorphism, a truncated cytoplasmic tail, and is only

expressed on placental trophoblasts (40, 41, 44). Mamu-AG is therefore thought to contribute to

pregnancy success through interactions with receptors on NK cells and myeloid cells that

promote placental vascularization, facilitate fetal tolerance, and protect the maternal-fetal

interface from invading microorganisms (44-46). Rhesus macaques also express Mamu-I, which

is the product of a fixed duplication of an *MHC-B* gene with limited sequence variation (47).

In concert with the tandem duplication of their *Mamu-A* and *-B* genes, macaques have

evolved an expanded complement of lineage II *KIRs* that encode 3D KIRs for Mamu-A and -B

ligands (26, 48-52). Whereas humans only have two lineage II *KIR* genes that encode

KIR3DL1/S1 and KIR3DL2 as receptors for HLA-Bw4 and –A3/A11, respectively, rhesus

macaques have at least 19 lineage II *KIRs* that encode inhibitory and activating receptors for

Mamu-A and -B molecules (50, 52-56). Furthermore, in contrast to humans, which have multiple

lineage III *KIR* genes encoding KIR2DL/S receptors for HLA-C1 and -C2 ligands (1, 2),

macaques only have two lineage III *KIR* genes, one is a pseudogene (KIR3DP1) and the other

(KIR1D) expresses a single domain KIR of uncertain function (52). Thus, while human NK cell

regulation predominately depends on KIR2DL/S-HLA-C interactions, macaque NK cells are

dependent on KIR3DL/S interactions with Mamu-A and -B ligands (52).

The rhesus macaque has become an increasingly valuable animal model for infectious

diseases (57-68), reproductive biology (44, 69), and transplantation (70, 71). Although MHC

class I ligands have been identified for a few macaque KIRs (72-76), ligands for most of these

receptors remain undefined. We therefore performed a systematic survey of MHC class I

interactions for common lineage II KIR allotypes representing 16 predicted genes of Indian-

origin rhesus macaques. By measuring the responses of KIR-CD3ζ-transduced reporter cell lines

to target cells expressing individual MHC class I molecules and confirming interactions by

staining with KIR IgG-Fc domain (KIR-Fc) fusion proteins, we identified MHC class I ligands

for 11 rhesus macaque KIRs of previously undefined specificity. This represents the most

comprehensive analysis of ligand recognition by macaque KIRs to date and reveals novel

interactions as well as extensive functional diversification of these receptors.

Materials and Methods

Jurkat NFAT luciferase reporter cells expressing chimeric KIR-CD3ζ receptors

cDNA sequences encoding the leader peptide of Mamu-KIR3DL05*008 followed by a Flag-tag

(DYKDDDDK) and the D0, D1, D2 and stem domains of rhesus macaque KIRs were

synthesized (Integrated DNA Technologies) and cloned into pQCXIP in-frame with sequences

encoding the transmembrane and cytoplasmic domains of human CD3ζ. Accuracy of all plasmid

constructs was verified by Sanger sequencing of the entire ORF. Retroviral vectors were

packaged into VSV G–pseudotyped murine leukemia virus (MLV) particles by co-transfecting

GP2-293 cells (Clontech Laboratories) with pQCXIP constructs and pVSV-G (Clontech

Laboratories). Jurkat NFAT luciferase (JNL) cells (Signosis) were transduced via spinoculation

for one hour with filtered supernatant collected from transfected GP2-293 cells and 2 days later

were placed under selection in RPMI medium supplemented with 10% FBS, L-glutamine,

penicillin, and streptomycin (R10 medium) plus 0.1 mg/ml hygromycin (Invitrogen) to maintain

the luciferase reporter gene. Selection for the KIR-CD3ζ receptor was performed by increasing

the puromycin (Invitrogen) concentration to 1.0 µg/ml over 3-4 weeks and then maintaining at

0.5 µg/ml. The GenBank accession numbers for rhesus macaque KIR and human CD3ζ

sequences are as follows: *Mamu-KIR3DL02*004:01* (LT963635.1), *-KIR3DLw03*002*

(EU419055.1), *-KIR3DL04*001:02* (GU299490.1), *-KIR3DL07*004* (EU419060.1), -

*KIR3DL10*002:01* (GU112259.1), *-KIR3DL11*001* (GU112271.1), *-KIR3DLw34*002*

(previously *-KIR3DLw03*003*; EU419031.1), *-KIR3DS01*003* (GU564161.1), -

*KIR3DS02*004:02* (GU014296.1), *-KIR3DS03*003* (EU702454.1), *-KIR3DS04*003:01*

(EU419029.1), *-KIR3DS05*002:01* (GU112262.1), *-KIR3DS06*002:02* (GU112298.1), -

KIR3DSw07*001 (GU112272.1), *-KIR3DSw08*001* (AY505479.1), *-KIR3DSw09*003*

(MF164924.1) and *CD3ζ* (J04132.1).

Rhesus macaque MHC class I-expressing 721.221 cells

Codon-optimized cDNA sequences encoding rhesus macaque MHC class I heavy chain

constructs were cloned into pQCXIP or pQCXIN and verified by Sanger sequencing of the entire

ORF. Retroviral vectors were packaged into VSV-G-pseudotyped MLV particles as described

above. 721.221 cells were transduced via spinoculation for one hour with filtered supernatant

collected from transfected GP2-293 cells and placed under selection in R10 medium containing

0.4 mg/ml puromycin or 0.5 mg/ml G418, depending on the vector, to select for MHC class I

expression. Selection for the MHC class I receptor was performed for pQCXIN constructs by

increasing the G418 (Enzo Life Sciences) concentration to 1.0 mg/ml (based on active units) and

then maintaining at 0.5 mg/ml and for pQCXIP constructs by increasing the puromycin

(Invitrogen) concentration to 1.0 µg/ml and then maintaining at 0.4 µg/ml. The GenBank

accession numbers for rhesus macaque MHC class I sequences are as follows: *Mamu-*

A1*001:01:01:01 (LT908117), *-A1*002:01:01:01* (LN899621), *-A1*004:01:01:01*

(FN396407.1), *-A1*008:01:01:01* (LN899628), *-A1*011:01:01:01* (AJ542579), *-*

A1*012:01:01:01 (AF157398.1), *-A2*05:02:01:01* (LN899639), *-A2*05:04:01:01*

(LT908121.1), *-A3*13:02:01:01* (LN899654.1), *-A3*13:03:01:01* (AF157401), *-*

A4*14:03:01:01 (GU080236.1), *-A6*01:03:02:01* (EF602318.1), *-B*001:01:02:01*

(LM608018), *-B*002:01:01:01* (LN851852), *-B*005:01:01:01* (LM608023), *-B*007:01:01:01*

(U41829.1), *-B*008:01:01:01* (U41830), *-B*015:01* (AM902541.1), *-B*017:01:01:01*

(AF199358), *-B*022:01:01:01* (LN899675), *-B*036:01:01:01* (AJ556886.1), *-B*041:01:01:01*

(LN899682), *-B*043:01:01:01* (AJ556893), *-B*045:03:01:01* (LM608047), *-B*056:01:01:01*

(GQ902079.1), *-B*065:01:01:01* (AJ620416.1), *-I*01:01:01* (LT908864), *-E*02:01:02*

(previously E*02:01; NM_001114966.1) *-F*01:01* (LT899414), *-AG1*05:01* (FJ409466), -

*AG2*01:01* (previously AG*01:01; U84783.1), *AG2*01:02:01:01* (LR989653)-

*AG3*02:01:01:01* (U84785.1), *-AG3*03:01:01:01* (U84787.1), *-AG4*08:01:01:01* (LR989670),

*-AG4*09:02:02:02* (LR990451), *-AG5*07:06:01:01* (LR989633), *-AG5*07:10:01:02*

(LR989642), and *-AG6*04:01:01:01* (LR989697). Only partial-length cDNA sequence was

available for *-AG1*05:01*. For the construct expressing this allele, sequences encoding the α3-,

transmembrane and cytoplasmic domains were inferred from the *Mamu-AG* consensus sequence.

KIR-Fc fusion proteins

All KIR-Fc fusion constructs were cloned into an in-house made expression construct (pCIW)

that contains the following regulatory elements: CMV promoter/enhancer, SV40 intron,

woodchuck hepatitis virus posttranscriptional regulatory element and SV40 polyadenylation

signal. The constructs encoded an artificial "secrecon" leader peptide (77) followed by KIR

ectodomain fused to the Fc domain of murine IgG2A. Following cloning of the initial

KIR3DL01 construct using a synthetic gBlock DNA fragment (IDT DNA), all other KIR

fragments were cloned in place of the KIR3DL01 using the "QuickChange cloning" protocol as

described by Klenchin et al (78). All fusion proteins were expressed in Expi-293F cells in 250-

280 ml culture using the ExpiFectamine 293 Transfection Kit (Life Technologies) according to

the manufacturer's recommendations. Five-six days after transfection, the supernatant was

collected, filtered through a 0.45 µM PES filter, and loaded on a rProtein A GraviTrap column

(GE Healthcare). After thorough washing of the column with PBS (at least 30 ml), KIR-Fc

fusion proteins were eluted with 3 ml Gentle Ag/Ab Buffer (Pierce) and buffer exchanged into

50 mM sodium citrate, (pH 6.5) using three cycles of concentration/dilution in Amicon Ultra-15

50K centrifugal Filter unit (Millipore). The final concentrations of the purified KIR-Fc fusion

proteins were calculated from absorbance at 280 nm using theoretical extinction coefficients

calculated by ProtParam (https://web.expasy.org/protparam/). After addition of 3 mM sodium

azide, the purified protein preparations were clarified by centrifugation at 30,000 g for 30 min at

20 °C and the supernatants were snap frozen in liquid nitrogen as 50 µl aliquots and stored at -80

°C.

Flow cytometry

KIR-CD3ζ expression on the surface of JNL cells and MHC class I expression on 721.221 cells

was confirmed by staining with Near-IR fluorescent dye (Invitrogen) followed by 30 min

staining at room temperature with PE-conjugated anti-Flag (clone REA216, Miltenyi Biotec) or

pan-HLA class I-specific antibodies (clone W6/32; Life Technologies). The cells were then

washed twice in PBS plus 1% FBS (FACS buffer) and fixed in 2% formaldehyde in PBS. To

assess KIR binding, MHC class I-transduced 721.221 cells were stained for 30 minutes at room

temperature with Near-IR fluorescent dye, washed twice with FACS buffer and stained for 30

minutes at room temperature with KIR-Fc fusions proteins (10-20 µg/ml). The cells were then

washed twice with FACS buffer, stained for 45 minutes with Alexa Flour 647-conjugated goat

anti-mouse IgG (RRID AB_2338925, Jackson ImmunoResearch), washed twice, and stained for

30 minutes at room temperature with the PE-conjugated pan-MHC class I-specific antibody

W6/32. The cells were washed again and fixed in 2% formaldehyde. All flow cytometry data

were collected using a FACSymphony™ A3 Cell Analyzer (BD Biosciences) and analyzed using

FlowJo version 10.8.0. After gating on viable cells, KIR-CD3ζ (Flag-tag) and MHC class I

staining was compared to parental JNL and 721.221 cells, respectively. KIR-MHC class I

binding was similarly assessed by comparing KIR-Fc versus W6/32 staining on the surface of

MHC class I-transduced 721.221 cells to their levels of non-specific staining on parental 721.221

cells.

KIR-CD3ζ JNL reporter assays

KIR-CD3ζ JNL cells (100,000) and MHC class I-transduced 721.221 cells (100,000) were co-

cultured overnight at 37 °C and 5% CO_2 in R10 medium without antibiotics. KIR-CD3ζ JNL

cells were also incubated with parental 721.221 cells as negative control and with anti-Flag-tag

(5 µg/ml) (clone 5A8E5, GenScript) and goat anti-mouse (10 µg/ml) (Poly4053, Biolegend)

antibodies as a positive "X-link" control. Each combination was tested in triplicate wells (100

µl/well). After 12-18 hours, BriteLite Plus luciferase substrate (PerkinElmer) was added to each

well (100 µl/well) and the relative light units (RLU) of luciferase activity was measured using a

VICTOR X4 multiplate reader (PerkinElmer).

Alignments and Phylogenetic trees

Amino acid sequences of KIR and MHC class I molecules were aligned using Geneious Prime

2021.2.2 (Biomatters Ltd). Phylogenetic trees were constructed using raw difference neighbor-

joining analysis of nucleic acid sequence for the α1-α3 domains of Mamu-MHC and the D0-D2

domains of human and rhesus macaque KIRs was performed using MAFFT (version 7, publicly

available at https://mafft.cbrc.jp/alignment/server/) (79, 80). All analyses utilized G-INS-I

iterative refinement with 1000 bootstrap replicates. Trees were viewed using Phylo.io version

1.0.0 (81). For the analysis of KIR sequences, human lineage II KIRs were included as an

outgroup (KIR3DL1*001, AY760023.1 and KIR3DL2*001, MN196233.1).

Statistical analysis

The luciferase induction by KIR-CD3ζ JNL cells (RLU data) was analyzed by one-way ANOVA

for multiple comparisons of mean responses to each of the MHC class I-expressing 721.221 cells

with mean responses to the parental 721.221 cells. Statistical analyses were performed using

GraphPad Prism for Mac OS version 10.8.0.

Results

Extensive sequencing and segregation analyses of *KIR* transcripts support the existence of at

least 22 *KIR* genes in Indian-origin rhesus macaques (53-56). Although our group and others

have identified ligands for a few rhesus macaque KIRs (72-76), ligands for most of the KIR in

this species remain undefined. To provide a more complete foundation for studying NK cell

biology in this species, we investigated the MHC class I interactions of 16 KIRs representing

common alleles of distinct lineage II genes in animals of Indian descent (Fig. 1A). Jurkat cells

harboring an NFAT-inducible luciferase reporter gene (JNL cells) were transduced with

retroviral vectors expressing chimeric KIR-CD3ζ receptors consisting of the extracellular

domains of macaque KIRs (D0, D1, D2 and stem domains) fused to the transmembrane and

cytoplasmic domains of human CD3ζ. To confirm surface expression, a Flag-tag was appended

to the N-terminus of the D0 domain of each KIR-CD3ζ receptor (Fig. 1B). 721.221 cells were

similarly transduced with vectors expressing rhesus macaque MHC class I molecules and surface

expression was confirmed with a pan-MHC class I antibody (Supplemental Figs 1-3). After co-

incubating the KIR-CD3ζ JNL cells overnight with MHC class I-transduced 721.221 cells,

ligand recognition was detected by the MHC class I-dependent upregulation of luciferase. KIR-

MHC class I interactions were also corroborated by staining MHC class I-expressing 721.221

cells with KIR-Fc fusion proteins consisting of the extracellular domains of macaque KIRs fused

to the Fc domain of murine IgG2A.

Mamu-Bw4 recognition by multiple rhesus macaque KIRs

Four of the rhesus macaque KIRs recognized Mamu-Bw4 ligands. KIR3DL04*001- and

KIR3DL10*002-CD3ζ JNL cells responded to 721.221 cells expressing Mamu-B*007:01, -

B*022:01, -B*041:01, -B*043:01 and -B*065:01 (Fig. 2A, Supplemental Fig. 4A). Similarly,

KIR3DL11*001-CD3ζ JNL cells responded to Mamu-B*007:01, -B*041:01 and -B*065:01, and

KIR3DS04*003-CD3ζ JNL cells responded to Mamu-B*22:01, -B*041:01 and -B*065:01 (Fig.

2A, Supplemental Fig. 4A). These interactions were corroborated by KIR-Fc staining of 721.221

cells expressing these molecules (Fig. 2B, Supplemental Fig. 4B).

Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01 and -B*065:01 share a Bw4 motif

at residues 77-83 (Supplemental Fig. 2), which was previously shown to contribute to

recognition by KIR3DL01 as well as three other rhesus macaque KIRs (73, 75). The

identification of Mamu-Bw4 ligands for KIR3DL04*001, KIR3DL10*002, KIR3DL11*001 and

KIR3DS04*003 doubles the number of rhesus macaque KIRs known to interact with this group

of molecules and reveals an extraordinary expansion of the lineage II KIRs in macaques with this

specificity (Table I). Differences in the particular Mamu-Bw4 molecules recognized by these

KIRs and the relative strength of these interactions suggests extensive functional diversification

of these receptors.

Multiple KIRs recognize Mamu-AG molecules

Six rhesus macaque KIRs were found to interact with Mamu-AG, a non-classical MHC class I

molecule expressed at the maternal-fetal interface of the placenta (40-44). JNL cells expressing

KIR-CD3ζ chimeras for KIR3DL07*004, KIR3DS01*003, KIR3DS02*004, KIR3DS06*002,

KIR3DSw07*001 and KIR3DSw09*003 all responded strongly to 721.221 cells expressing

Mamu-AG2*01:01 and -AG6*04:01 (Fig. 3A, Supplemental Fig. 5A). In addition,

KIR3DS02*004-CD3ζ JNL cells also engaged 721.221 cells expressing Mamu-AG2*01:02 (Fig.

3A). The binding of these KIRs to Mamu-AG2*01:01 and -AG6*04:01 was corroborated by

KIR-Fc staining of Mamu-AG2-expressing 721.221 cells (Fig. 3B, Supplemental Fig. 5B). KIR-

Fc staining also revealed detectable interactions with Mamu-AG1*05:01 for KIR3DS02*004,

KIR3DS06*002 and KIR3DSw09*003, and Mamu-AG2*01:02 for KIR3DL07*004 and

KIR3DSw09*003 that were not observed in cellular assays with KIR-CD3ζ JNL cells (Fig. 3B,

Supplemental Fig. 5B). In this case, the detection of binding interactions with Mamu-AG1*05:01

and -AG2*01:02 that were not detectable with KIR-CD3ζ JNL cells may reflect greater

sensitivity of KIR-Fc staining or possibly interference of the Flag-tag appended to the N-

terminus of the KIR-CD3ζ receptor in the reporter cell assay. These findings identify multiple

Mamu-AG ligands for the products of six different KIR genes, five of which encode activating

receptors. This brings the total number of rhesus macaque KIRs known to interact with Mamu-

AG to seven (Table I).

*Recognition of Mamu-B*045:03 by KIR3DL07*004, KIR3DS02*004 and KIR3DSw09*003*

In addition to Mamu-AG, JNL cells expressing KIR3DL07*004-CD3ζ also responded to Mamu-

B*045:03 (Fig. 3A, Supplemental Fig. 6A). KIR-Fc staining corroborated this interaction and

additionally also revealed low but detectable interactions for Mamu-B*045:03 with

KIR3DS02*004 and KIR3DSw09*003 (Fig. 3B, Supplemental Fig. 6B). To better understand

how these KIR, which predominately recognize Mamu-AG ligands, can also interact with

B*045:03, we compared the amino acid sequences for Mamu-B*045:03 with other Mamu-A, -B

and -AG molecules. Although Mamu-B*045:03 resembles a typical Mamu-B molecule in its α2

domain, we noted a region of sequence similarity with other Mamu-A molecules in the α1-

domain (Fig. 4A). These observations were also supported by phylogenetic analysis. Whereas

the full-length nucleotide sequence for Mamu-B*045:03 clusters with Mamu-B molecules (Fig.

4B), the α1-domain is more similar to Mamu-A molecules (Fig. 4C). A similar pattern of

clustering was observed for Mamu-B*036, which is a product of recombination between *MHC-A*

and *-B* genes and was previously identified as a ligand for KIR3DL05 (75). Mamu-B*045:03

recognition by KIR3DL07*004, KIR3DS02*004 and KIR3DSw09*003 may therefore be

explained by its origins as the product of recombination between ancestral *MHC-A* and *-B* genes.

*Recognition of Mamu A1*012:01 by KIR3DLw03*002*

KIR3DLw03*002-CD3ζ JNL cells robustly responded to 721.221 cells expressing Mamu-

A*012:01 (Fig. 5A, Supplemental Fig. 7A), but not to other MHC class I molecules tested in this

study. The binding of KIR3DLw03*002 to Mamu-A*012:01 was also corroborated by KIR-Fc

staining (Fig. 5B, Supplemental Fig. 7B). KIR3DLw03*002 is one of four KIRs that display an

absence of 3 amino acids at positions 37-39 of the D0 domain compared to the other rhesus

macaque KIRs tested in this study (Fig. 1A).

Phylogenetic relationship of rhesus macaque KIRs is concordant with patterns of ligand

recognition

To investigate the relationship between the extracellular domains of rhesus macaque KIRs and

their patterns of ligand recognition, we constructed a phylogenic tree of the D0-D2 domains of

rhesus macaque KIRs using human KIR3DL1 and KIR3DL2 as outgroups. The clustering of

macaque KIRs generally reflected their ligand recognition. Six of the eight KIRs found to

interact with Mamu-Bw4 ligands clustered together on a branch that includes KIR3DSw39*002,

KIR3DL06*001, KIR3DS04*003, KIR3DL11*001, KIR3DL04*001 and KIR3DL08*002 (Fig.

6). Likewise, six of the seven KIRs found to interact with Mamu-AG ligands clustered together

on a separate branch that includes KIR3DS06*002, KIR3DS01*003, KIR3DL07*004,

KIR3DSw09*003, KIR3DS02*004 and KIR3DL05*008 (Fig. 6). KIR3DL10*002,

KIR3DSw07*001 and KIR3DLw03*002 were exceptions. Although KIR3DL10*002 and

KIR3DSw07*001 interact with Mamu-Bw4 and -AG ligands, respectively, these receptors

grouped together on the same branch (Fig. 6). In contrast, KIR3DLw03*002, which interacts

with Mamu-A1*012:01, was found on a distinct branch with three other KIRs for which ligands

have not yet been identified (KIR3DS05*002, KIR3DLw34*002 and KIR3DL02*004) (Fig. 6).

This group of KIRs is separated from the others by strong bootstrap support and is distinguished

by the absence of three amino acids (residues 37-39) in the D0 domain (Fig. 1A) that are also

missing in human KIR3DL1/S1 and KIR3DL2.

Discussion

Establishing the MHC class I ligands of rhesus macaques KIRs is fundamental to NK cell

biology in this species as a preclinical model for infectious diseases, transplantation, and

reproductive biology. Because of the rapid pace of KIR evolution in primates, it is not possible to

predict the ligands for macaque KIRs on the basis of sequence similarity with their human

counterparts. The *KIR* genes of macaques are evolving at a particularly rapid pace characterized

by extensive gene duplication and recombination, generating allelic variation and haplotype

diversity that greatly exceeds the genetic diversity of human *KIRs* (55, 56, 82). Although the

number of alleles and predicted *KIR* genes varies among geographically distinct populations

(83), extensive sequencing and segregation analyses strongly support the existence of at least 22

distinct *KIR* genes in rhesus macaques of Indian origin (55, 56, 82). Of these, 19 are predicted to

encode lineage II KIRs for Mamu-A and -B ligands. To provide a foundation for studying NK

cell responses in macaques, we selected common allotypes representing 16 lineage II KIRs for

analysis of their MHC class I interactions. By screening 721.221 cells expressing MHC class I

molecules corresponding to 39 *Mamu-A, -B, -E, -F, -I* and *-AG* alleles in assays with KIR-

CD3ζ JNL reporter cell lines and by staining these MHC class I-expressing 721.221 cells with

KIR-Fc fusion proteins, we identified ligands for 11 previously uncharacterized KIRs.

These analyses revealed two general patterns of ligand recognition; interactions with

Mamu-Bw4 molecules and interactions with Mamu-A-related molecules, particularly Mamu-

AG. Of the 16 KIRs tested in this study, four (KIR3DL04*001, KIR3DL10*002,

KIR3DL11*001 and KIR3DS04*003) were found to interact with molecules that contain a Bw4

motif at residues 77-83. This demonstrates a remarkable expansion of Bw4 specificity in the

rhesus macaque. In contrast to humans, which only have a single *KIR3DL1/S1* gene that encodes

receptors for HLA-Bw4, rhesus macaques have at least eight lineage II *KIRs* that encode

receptors for Bw4 molecules (Table I). Six of the KIRs in this study (KIR3DL07*004,

KIR3DS01*003, KIR3DS02*004, KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003)

were found to interact with Mamu-AG2*01:01 and AG6*04:01, either alone or in combination

with Mamu-AG1*05:01, -AG2*01:02, and -B*45:03. Together with earlier studies defining

similar ligands for KIR3DL05 (72, 74, 76), at least eight rhesus macaque *KIRs* are known to

encode receptors for Mamu-A or -AG ligands (Table I). We also identified Mamu-A1*012:01 as

a ligand for KIR3DLw03*002, which belongs to a distinct group of receptors with a

characteristic deletion of three amino acids in the D0 domain.

All the KIRs that interact with Bw4 ligands recognize Mamu-B*041:01 and -B*065:01

with varying efficiency, suggesting that these molecules may provide a common reference for

this group of receptors (Table I). However, interactions with other Mamu-Bw4 molecules were

more variable. Several KIRs, including KIR3DL11*001 and KIR3DS04*003, lack detectable

responses to Mamu-B*007:01, -B*022:01 or -B*043:01. Differences in the magnitude of KIR-

CD3ζ JNL cell responses and KIR-Fc staining also suggest variation in the strength of these

interactions. Although most of the KIRs responding to Mamu-Bw4 ligands are inhibitory,

interactions with a couple of activating KIRs (KIR3DS04*003 and KIR3DSw39*002) were also

observed (Table I). Thus, differences in the preferential recognition of individual Mamu-Bw4

molecules, variation in the strength of these interactions, and the identification of inhibitory and

activating KIRs that respond to this group of molecules, indicates extensive functional

diversification of this group of receptors.

Rhesus macaque MHC class I haplotypes typically express two or three *Mamu-A* genes

and 4-11 *Mamu-B* genes that can be classified as major or minor loci based on their

transcriptional abundance (30-32). The Mamu-Bw4 molecules identified as KIR ligands are

products of transcriptionally abundant genes (32). However, phylogenetic and segregation

analyses suggest that they are distinct from the *Mamu-B* genes that encode molecules known to

present viral peptides to CD8$^+$ T cells (30-33, 84). This is further supported by unique

polymorphisms in the cytoplasmic domains of Mamu-B*007:01, -B*022:01, -B*041:01, -

B*043:01 and -B*065:01 that are not present in the cytoplasmic tails of molecules such as

Mamu-B*008 and -B*017 that are known to present SIV epitopes for recognition by CD8+ T

cells (84, 85). We previously demonstrated that these cytoplasmic domain polymorphisms

prevent MHC class I downmodulation by SIV Nef (86). Selective downmodulation of HLA-A

and -B (but not HLA-C) by HIV-1 Nef reduces the susceptibility of virus-infected cells to CD8+

T cells (6, 7, 87). Although the Vpu proteins of certain HIV-1 isolates may downmodulate HLA-

C at later stages of infection (88, 89), HLA-C is initially retained on the cell surface, presumably

to maintain interactions with inhibitory KIR (6). In a similar manner, SIV Nef downmodulates

Mamu-A and -B molecules known to present viral peptides to CD8+ T cells, but not the products

of *Mamu-Bw4* genes that serve as ligands for multiple KIRs (75, 86). Hence, parallels in both

immunogenetics and virology suggest functional specialization, whereby the products of certain

Mamu-B genes play a dominant role in restricting CD8+ T cell responses while others serve as

KIR ligands to modulate NK cell responses.

Another group of rhesus macaque KIRs interacts with Mamu-A-related ligands, including

certain Mamu-AG allotypes and in some cases Mamu-B*045:03. KIR3DL07*004,

KIR3DS01*003, KIR3DS02*004, KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003 all

recognize Mamu-AG2*01:01 and -AG6*04:01 (Table I). Additional interactions with Mamu-

AG1*05:01 were detectable for KIR3DS02*004, KIR3DS06*002 and KIR3DSw09*003 and

with Mamu-AG2*01:02 for KIR3DL07*004, KIR3DS02*004, and KIR3DSw09*003 (Table I).

Mamu-B*045:03 was also identified a ligand for KIR3DL07*004, KIR3DS02*004 and

KIR3DSw09*003 (Table I). Although Mamu-B*045:03 is a product of a *Mamu-B* allele, a

region of its α1-domain appears to be derived from an *MHC-A* gene. Thus, the hybrid origins of

Mamu-B*045:03 probably account for its recognition by these KIRs.

In contrast to KIRs with specificity for Mamu-Bw4, most receptors for Mamu-AG are

activating. Five of the seven rhesus macaque KIRs that interact with Mamu-AG2*01:01 and -

AG6*04:01 are products of *KIR3DS* genes (Table I). In humans, the ligation of activating lineage

III KIRs (KIR2DS) and KIR2DL4 on maternal NK cells of the placental decidua by HLA-C and

soluble HLA-G stimulates the release of pro-inflammatory and pro-angiogenic factors that

promote placental vascularization (20, 21, 45). Since macaques do not express orthologs of

HLA-C or -G (28, 29, 38), it is conceivable that KIR3DS interactions with Mamu-AG play a

similar role in placental development. Unlike human *KIRs*, which can be broadly categorized

into inhibitory or activating haplotypes based on fixed differences in the number of activating

genes, *KIR* haplotypes in macaques are much more variable (82). There are typically 4-17

expressed *KIR* genes per haplotype in the rhesus macaque, including at least one activating gene

(56, 82). The expansion of lineage II *KIR* that encode receptors for Mamu-AG may therefore

ensure that every animal has at least one activating KIR capable of engaging this non-classical

molecule in the placenta.

The identification of Mamu-A1*012:01 as a ligand for KIR3DLw03*002 provides at

least one MHC class I ligand for a phylogenetically distinct group of receptors with a

characteristic deletion of residues 37-39 in D0. Additional comparison of the D0 domains of

KIRs expressed in other primate species, revealed that these residues are also absent from the

lineage II KIRs of apes and humans. This suggests that hominid lineage II KIRs evolved from a

KIR gene with a corresponding 9 base pair deletion in exon 3 that was present in the common

ancestor of apes and Old World monkeys. Further inspection indicates that this deletion results in

polymorphic differences in potential N-linked glycosylation (PNG) sites that are close to a

predicted MHC class I contact site (90). Whereas most rhesus macaque KIRs have two tandem

PNG sites ($\underline{N}_{36}$FT$\underline{N}_{39}$FT), KIR3DL02, KIR3DLw03, KIR3DLw34 and KIR3DS05 only contain

a single PNG site in this region ($\underline{N}_{36}$FT) (Fig. 1). Furthermore, because of an additional

threonine-to-methionine substitution at residue 38 (N_{36}FM), human lineage II KIRs (KIR3DL1

and KIR3DL2) lack a PNG site in this region. Hence, these species-specific differences in

glycosylation may influence MHC class I interactions with lineage II KIRs.

The present study more than doubles the number of rhesus macaque KIRs with defined

MHC class I ligands. At least one MHC class I ligand has now been identified for KIRs

representing 16 different lineage II *KIR* genes in animals of Indian origin. These findings reveal

a remarkable expansion of lineage II KIRs with broad specificity for Mamu-Bw4 or -AG ligands

and novel receptor interactions with Mamu-AG, -B*045 and -A1*012 allotypes. The recognition

of overlapping, but non-redundant sets of ligands by both inhibitory and activating KIRs

indicates extensive diversification of these receptors in concert with an expanded complement of

Mamu-A and *-B* genes. These observations advance our basic understanding of KIR and MHC

class I co-evolution in primates and provide a valuable foundation for studying NK cell

responses in the rhesus macaque as a pre-clinical model.

References

1.   Parham, P. 2005. MHC class I molecules and KIRs in human history, health and survival.

Nat. Rev. Immunol. 5: 201-214.

2.   Lanier, L. L. 2005. NK cell recognition. *Annu. Rev. Immunol.* 23: 225-274.

3.   Hasan, M. Z., and L. Walter. 2021. Rhesus Macaque Activating Killer Immunoglobulin-

Like Receptors Associate With Fc Receptor Gamma (FCER1G) and Not With DAP12

Adaptor Proteins Resulting in Stabilized Expression and Enabling Signal Transduction.

Front. Immunol. 12: 678964.

4.   Huard, B., and K. Fruh. 2000. A role for MHC class I down-regulation in NK cell lysis of

herpes virus-infected cells. *Eur. J. Immunol.* 30: 509-515.

5.   Falk, C. S., M. Mach, D. J. Schendel, E. H. Weiss, I. Hilgert, and G. Hahn. 2002. NK cell

activity during human cytomegalovirus infection is dominated by US2-11-mediated HLA

class I down-regulation. *J. Immunol.* 169: 3257-3266.

6.   Cohen, G. B., R. T. Gandhi, D. M. Davis, O. Mandelboim, B. K. Chen, J. L. Strominger,

and D. Baltimore. 1999. The selective downregulation of class I major histocompatibility

complex proteins by HIV-1 protects HIV-infected cells from NK cells. *Immunity* 10: 661-

671.

7.   Schwartz, O., V. Marechal, S. LeGall, F. Lemonnier, and J.-M. Heard. 1996. Endocytosis

of major histocompatibility complex class I molecules is induced by the HIV-1 Nef

protein. *Nat. Med.* 2: 338-342.

8.   Porgador, A., O. Mandelboim, N. P. Restifo, and J. L. Strominger. 1997. Natural killer

cell lines kill autologous beta2-microglobulin-deficient melanoma cells: implications for

cancer immunotherapy. *Proc. Natl. Acad. Sci. U. S. A.* 94: 13140-13145.

9.   Alter, G., M. P. Martin, N. Teigen, W. H. Carr, T. J. Suscovich, A. Schneidewind, H.

Streeck, M. Waring, A. Meier, C. Brander, J. D. Lifson, T. M. Allen, M. Carrington, and

M. Altfeld. 2007. Differential natural killer cell-mediated inhibition of HIV-1 replication

based on distinct KIR/HLA subtypes. *J. Exp. Med.* 204: 3027-3036.

10. O'Connor, G. M., J. P. Vivian, E. Gostick, P. Pymm, B. A. Lafont, D. A. Price, J.

Rossjohn, A. G. Brooks, and D. W. McVicar. 2015. Peptide-dependent recognition of

HLA-B*57:01 by KIR3DS1. *J. Virol.* 89: 5213-5221.

11. Garcia-Beltran, W. F., A. Holzemer, G. Martrus, A. W. Chung, Y. Pacheco, C. R.

Simoneau, M. Rucevic, P. A. Lamothe-Molina, T. Pertel, T. E. Kim, H. Dugan, G. Alter,

J. Dechanet-Merville, S. Jost, M. Carrington, and M. Altfeld. 2016. Open conformers of

HLA-F are high-affinity ligands of the activating NK-cell receptor KIR3DS1. *Nat.*

Immunol. 17: 1067-1074.

12. Martin, M. P., X. Gao, J-H. Lee, G. W. Nelson, R. Detels, J. J. Goedert, S. Buchbinder,

K. Hoots, D. Vlahov, J. Trowsdale, M. Wilson, S. J. O'Brien, and M. Carrington. 2002.

Epistatic interaction between KIR3DS1 and HLA-B delays the progression to AIDS. *Nat.*

Genet. 31: 429-434.

13. Martin, M. P., Y. Qi, X. Gao, E. Yamada, J. N. Martin, F. Pereyra, S. Colombo, E. E.

Brown, W. L. Shupert, J. Phair, J. J. Goedert, S. Buchbinder, G. D. Kirk, A. Telenti, M.

Connors, S. J. O'Brien, B. D. Walker, P. Parham, S. G. Deeks, D. W. McVicar, and M.

Carrington. 2007. Innate partnership of HLA-B and KIR3DL1 subtypes against HIV-1.

Nat. Genet. 39: 733-740.

14. Khakoo, S. I., C. L. Thio, M. P. Martin, C. R. Brooks, X. Gao, J. Astemborski, J. Cheng,

J. J. Godert, D. Vlahov, M. Hilgartner, S. Cox, A.-M. Little, G. J. Alexander, M. E.

Cramp, S. J. O'Brien, W. M. C. Rosenberg, D. L. Thomas, and M. Carrington. 2004. HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. *Science* 305: 872-874.

15. Carrington, M., S. Wang, M. P. Martin, X. Gao, M. Schiffman, J. Cheng, R. Herrero, A. C. Rodriguez, R. Kurman, R. Mortel, P. Schwartz, A. Glass, and A. Hildesheim. 2005. Hierarchy of resistance to cervical neoplasia mediated by combinations of killer immunoglobulin-like receptor and human leukocyte antigen loci. *J. Exp. Med.* 201: 1069-1075.

16. Chen, C., M. Busson, V. Rocha, M.-L. Appert, V. Lepage, N. Dulphy, P. Haas, G. Socie, A. Toubert, D. Charron, and P. Loiseau. 2006. Activating KIR genes are associated with CMV reactivation and survival after non-T-cell depleted HLA-identical sibling bone marrow transplantation for malignant disorders. *Bone Marrow Transplant.* 38: 437-444.

17. Cooley, S., E. Trachtenberg, T. L. Bergemann, K. Saeteurn, J. Klein, C. T. Le, S. G. Marsh, L. A. Guethlein, P. Parham, J. S. Miller, and D. J. Weisdorf. 2009. Donors with group B KIR haplotypes improve relapse-free survival after unrelated hematopoietic cell transplantation for acute myelogenous leukemia. *Blood* 113: 726-732.

18. Cooley, S., D. J. Weisdorf, L. A. Guethlein, J. P. Klein, T. Wang, C. T. Le, S. G. Marsh, D. Geraghty, S. Spellman, M. D. Haagenson, M. Ladner, E. Trachtenberg, P. Parham, and J. S. Miller. 2010. Donor selection for natural killer cell receptor genes leads to superior survival after unrelated transplantation for acute myelogenous leukemia. *Blood* 116: 2411-2419.

19. Delgado, D. C., J. A. Hank, J. Kolesar, D. Lorentzen, J. Gan, S. Seo, K. Kim, S. Shusterman, S. D. Gillies, R. A. Reisfeld, R. Yang, B. Gadbaw, K. B. DeSantes, W. B.

London, R. C. Seeger, J. M. Maris, and P. M. Sondel. 2010. Genotypes of NK cell KIR

receptors, their ligands, and Fcgamma receptors in the response of neuroblastoma

patients to Hu14.18-IL2 immunotherapy. *Cancer Res.* 70: 9554-9561.

20.   Hiby, S. E., J. J. Walker, M. O'Shaughnessy K, C. W. Redman, M. Carrington, J.

Trowsdale, and A. Moffett. 2004. Combinations of maternal KIR and fetal HLA-C genes

influence the risk of preeclampsia and reproductive success. *J. Exp. Med.* 200: 957-965.

21.   Hiby, S. E., R. Apps, A. M. Sharkey, L. E. Farrell, L. Gardner, A. Mulder, F. H. Claas, J.

J. Walker, C. W. Redman, L. Morgan, C. Tower, L. Regan, G. E. Moore, M. Carrington,

and A. Moffett. 2010. Maternal activating KIRs protect against human reproductive

failure mediated by fetal HLA-C2. *J. Clin. Invest.* 120: 4102-4110.

22.   Orrantia, A., I. Terren, G. Astarloa-Pando, C. Gonzalez, A. Uranga, J. J. Mateos-Mazon,

J. C. Garcia-Ruiz, M. Rinon, M. Rey, S. Perez-Fernandez, O. Zenarruzabeitia, and F.

Borrego. 2021. NK Cell Reconstitution After Autologous Hematopoietic Stem Cell

Transplantation: Association Between NK Cell Maturation Stage and Outcome in

Multiple Myeloma. *Front. Immunol.* 12: 748207.

23.   Littera, R., G. Piredda, D. Argiolas, S. Lai, E. Congeddu, P. Ragatzu, M. Melis, E. Carta,

M. B. Michittu, D. Valentini, L. Cappai, R. Porcella, F. Alba, M. Serra, V. Loi, R. Maddi,

S. Orru, G. La Nasa, G. Caocci, R. Cusano, M. Arras, M. Frongia, A. Pani, and C.

Carcassi. 2017. KIR and their HLA Class I ligands: Two more pieces towards completing

the puzzle of chronic rejection and graft loss in kidney transplantation. *PLoS One* 12:

e0180831.

24. Yokoyama, W. M., P. J. Kehn, D. I. Cohen, and E. M. Shevach. 1990. Chromosomal

location of the Ly-49 (A1, YE1/48) multigene family, genetic association with the NK1.1

antigen. *J. Immunol.* 145: 2353-2358.

25. Khakoo, S. I., R. Rajalingam, B. P. Shum, K. Weidenbach, L. Flodin, D. G. Muir, F.

Canavez, S. L. Cooper, N. M. Valiante, L. L. Lanier, and P. Parham. 2000. Rapid

evolution of NK cell receptor systems demonstrated by comparison of chimpanzee and

humans. *Immunity* 12: 687-698.

26. Sambrook, J. G., A. Bashirova, S. Palmer, S. Sims, J. Trowsdale, L. Abi-Rached, P.

Parham, M. Carrington, and S. Beck. 2005. Single haplotype analysis demonstrates rapid

evolution of the killer immunoglobulin-like receptor (KIR) loci in primates. *Genome Res.*

15: 25-35.

27. Parham, P., L. Abi-Rached, L. Matevosyan, A. K. Moesta, P. J. Norman, A. M. O.

Aguilar, and L. A. Guethlein. 2010. Primate-specific regulation of natural killer cells. *J.*

Med. Primatol. 39: 194-212.

28. Boyson, J. E., C. Schufflebotham, L. F. Cadavid, J. A. Urvater, L. A. Knapp, A. L.

Huges, and D. I. Watkins. 1996. The MHC class I genes of the rhesus monkey: Different

evolutionary histories of the MHC class I and II genes in primates. *J. Immunol.* 156:

4656-4665.

29. Adams, E. J., and P. Parham. 2001. Species-specific evolution of MHC class I genes in

the higher primates. *Immunol. Rev.* 183: 41-64.

30. Otting, N., C. M. C. Heijmans, R. C. Noort, N. G. d. Groot, G. G. M. Doxiadis, J. J.

vanRood, D. I. Watkins, and R. E. Bontrop. 2005. Unparalleled complexity of the MHC

class I region in rhesus macaques. *Proc. Natl. Acad. Sci. U. S. A.* 102: 1626-1631.

31.   Daza-Vamenta, R., G. Glusman, L. Rowen, B. Guthrie, and D. E. Geraghty. 2004.

Genetic divergence of the rhesus macaque major histocompatibility complex. *Genome*

Res. 14: 1501-1515.

32.   Wiseman, R. W., J. A. Karl, P. S. Bohn, F. A. Nimityongskul, G. J. Starrett, and D. H.

O'Connor. 2013. Haplessly hoping: macaque major histocompatibility complex made

easy. *ILAR journal / National Research Council, Institute of Laboratory Animal*

Resources 54: 196-210.

33.   Evans, D. T., P. Jing, T. M. Allen, D. H. O'Connor, H. Horton, J. E. Venham, M.

Piekarczyk, J. Dzuris, M. Dykhuzen, J. Mitchen, R. A. Rudersdorf, C. D. Pauza, A. Sette,

R. E. Bontrop, R. DeMars, and D. I. Watkins. 2000. Definition of five new simian

immunodeficiency virus cytotoxic T-lymphocyte epitopes and their restricting major

histocompatibility complex class I molecules: evidence for an influence on disease

progression. *J. Virol.* 74: 7400-7410.

34.   Boyson, J. E., S. N. McAdam, A. Gallimore, T. G. Golos, X. Liu, F. M. Gotch, A. L.

Hughes, and D. I. Watkins. 1995. The MHC E locus in macaques is polymorphic and is

conserved between macaques and humans. *Immunogenetics* 41: 59-68.

35.   Knapp, L., L. F. Cadavid, and D. I. Watkins. 1998. The MHC-E locus is the most well

conserved of all known primate class I histocompatibility genes. *J. Immunol.* 160: 189-

196.

36.   Godfrey, D. I., J. Le Nours, D. M. Andrews, A. P. Uldrich, and J. Rossjohn. 2018.

Unconventional T cell targets for cancer immunotherapy. *Immunity* 48: 453-473.

37. Walter, L., and B. Petersen. 2017. Diversification of both KIR and NKG2 natural killer

cell receptor genes in macaques - implications for highly complex MHC-dependent

regulation of natural killer cells. *Immunology* 150: 139-145.

38. Boyson, J. E., K. K. Iwanaga, T. G. Golos, and D. I. Watkins. 1996. Identifiction of the

rhesus monkey HLA-G ortholog. Mamu-G is a pseudogene. *J. Immunol.* 157: 5428-5437.

39. Castro, M. J., P. Morales, V. Fernandez-Soria, B. Suarez, M. J. Recio, M. Alvarez, M.

Martin-Villa, and A. Arnaiz-Villena. 1996. Allelic diversity at the primate Mhc-G locus:

exon 3 bears stop codons in all Cercopithecinae sequences. *Immunogenetics* 43: 327-336.

40. Boyson, J. E., K. K. Iwanaga, T. G. Golos, and D. I. Watkins. 1997. Identification of a

novel MHC class I gene, Mamu-AG, expressed in the placenta of a primate with an

inactivated G locus. *J. Immunol.* 159: 3311-3321.

41. Boyson, J. E., K. K. Iwanaga, J. A. Urvater, A. L. Hughes, T. G. Golos, and D. I.

Watkins. 1999. Evolution of a new nonclassical MHC class I locus in two Old World

primate species. *Immunogenetics* 49: 86-98.

42. Langat, D. K., P. J. Morales, A. T. Fazleabas, J. M. Mwenda, and J. S. Hunt. 2002.

Baboon placentas express soluble and membrane-bound Paan-AG proteins encoded by

alternatively spliced transcripts of the class Ib major histocompatibility complex gene,

Paan-AG. *Immunogenetics* 54: 164-173.

43. Bondarenko, G. I., S. V. Dambaeva, R. L. Grendell, A. L. Hughes, M. Durning, M. A.

Garthwaite, and T. G. Golos. 2009. Characterization of cynomolgus and vervet monkey

placental MHC class I expression: diversity of the nonhuman primate AG locus.

Immunogenetics 61: 431-442.

44.   Slukvin, II, D. P. Lunn, D. I. Watkins, and T. G. Golos. 2000. Placental expression of the

588   nonclassical MHC class I molecule Mamu-AG at implantation in the rhesus monkey.

589   *Proc. Natl. Acad. Sci. U. S. A.* 97: 9104-9109.

590   45.   Rajagopalan, S. 2014. HLA-G-mediated NK cell senescence promotes vascular

591   remodeling: implications for reproduction. *Cell. Mol. Immunol.* 11: 460-466.

592   46.   Arnaiz-Villena, A., I. Juarez, F. Suarez-Trujillo, A. Lopez-Nares, C. Vaquero, J. Palacio-

593   Gruber, and J. M. Martin-Villa. 2020. HLA-G: Function, polymorphisms and pathology.

594   *Int. J. Immunogenet*. 48: 172-192.

595   47.   Urvater, J. A., N. Otting, J. H. Loehrke, R. Rudersdorf, I. I. Slukvin, M. S. Piekarczyk, T.

596   G. Golos, A. L. Hughes, R. E. Bonrop, and D. I. Watkins. 2000. Mamu-I: A novel

597   primate MHC class I B-related locus with unusually low variability. *J. Immunol.* 164:

598   1386-1398.

599   48.   Hershberger, K. L., R. Shyam, A. Miura, and N. L. Letvin. 2001. Diversity of the killer

600   cell Ig-like receptors of rhesus monkeys. *J. Immunol.* 166: 4380-4390.

601   49.   Bimber, B. N., A. J. Moreland, R. W. Wiseman, A. L. Hughes, and D. H. O'Connor.

602   2008. Complete characterization of killer Ig-like receptor (KIR) haplotypes in Mauritian

603   cynomolgus macaques: Novel insights into nonhuman primate KIR gene content and

604   organization. *J. Immunol.* 181: 6301-6308.

605   50.   Kruse, P. H., C. Rosner, and L. Walter. 2010. Characterization of rhesus macaque KIR

606   genotypes and haplotypes. *Immunogenetics* 62: 281-293.

607   51.   Blokhuis, J. H., M. K. v. d. Wiel, G. G. Doxiadis, and R. E. Bontrop. 2010. The mosaic

608   of KIR haplotypes in rhesus macaques. *Immunogenetics* 62: 295-306.

609   52.   Bimber, B. N., and D. T. Evans. 2015. The killer-cell immunoglobulin-like receptors of

610   macaques. *Immunol. Rev.* 267: 246-258.

611   53.   Blokhuis, J. H., M. K. van der Wiel, G. G. Doxiadis, and R. E. Bontrop. 2011. The

612   extreme plasticity of killer cell Ig-like receptor (KIR) haplotypes differentiates rhesus

613   macaques from humans. *Eur. J. Immunol.* 41: 2719-2728.

614   54.   Moreland, A. J., L. A. Guethlein, R. K. Reeves, K. W. Broman, R. P. Johnson, P.

615   Parham, D. H. O'Connor, and B. N. Bimber. 2011. Characterization of killer

616   immunoglobulin-like receptor genetics and comprehensive genotyping by

617   pyrosequencing in rhesus macaques. *BMC genomics* 12: 295.

618   55.   Bruijnesteijn, J., M. K. H. van der Wiel, W. T. N. Swelsen, N. Otting, A. J. M. de Vos-

619   Rouweler, D. Elferink, G. G. Doxiadis, F. H. J. Claas, N. M. Lardy, N. G. de Groot, and

620   R. E. Bontrop. 2018. Human and rhesus macaque KIR haplotypes defined by their

621   transcriptomes. *J. Immunol.* 200: 1692-1701.

622   56.   Bruijnesteijn, J., N. de Groot, M. K. H. van der Wiel, N. Otting, A. J. M. de Vos-

623   Rouweler, N. G. de Groot, and R. E. Bontrop. 2020. Unparalleled Rapid Evolution of

624   KIR Genes in Rhesus and Cynomolgus Macaque Populations. *J. Immunol.* 204: 1770-

625   1786.

626   57.   Hatziioannou, T., and D. T. Evans. 2012. Animal models for HIV/AIDS research. *Nat.*

627   *Rev. Microbiol.* 10: 852-867.

628   58.   Barouch, D. H., J. B. Whitney, B. Moldt, F. Klein, T. Y. Oliveira, J. Liu, K. E.

629   Stephenson, H. W. Chang, K. Shekhar, S. Gupta, J. P. Nkolola, M. S. Seaman, K. M.

630   Smith, E. N. Borducchi, C. Cabral, J. Y. Smith, S. Blackmore, S. Sanisetty, J. R. Perry,

631   M. Beck, M. G. Lewis, W. Rinaldi, A. K. Chakraborty, P. Poignard, M. C. Nussenzweig,

632  and D. R. Burton. 2013. Therapeutic efficacy of potent neutralizing HIV-1-specific

633  monoclonal antibodies in SHIV-infected rhesus monkeys. *Nature* 503: 224-228.

634 59. Esquivel, R. N., A. Patel, S. B. Kudchodkar, D. H. Park, K. Stettler, M. Beltramello, J.

635  W. Allen, J. Mendoza, S. Ramos, H. Choi, P. Borole, K. Asija, M. Bah, S. Shaheen, J.

636  Chen, J. Yan, A. C. Durham, T. R. F. Smith, K. Broderick, G. Guibinga, K. Muthumani,

637  D. Corti, L. Humeau, and D. B. Weiner. 2019. In Vivo Delivery of a DNA-Encoded

638  Monoclonal Antibody Protects Non-human Primates against Zika Virus. *Mol. Ther.* 27:

639  974-985.

640 60. Hessell, A. J., L. Hangartner, M. Hunter, C. E. Havenith, F. J. Beurskens, J. M. Bakker,

641  C. M. Lanigan, G. Landucci, D. N. Forthal, P. W. Parren, P. A. Marx, and D. R. Burton.

642  2007. Fc receptor but not complement binding is important in antibody protection against

643  HIV. *Nature* 449: 101-104.

644 61. Hessell, A. J., P. Poignard, M. Hunter, L. Hangartner, D. M. Tehrani, W. K. Bleeker, P.

645  W. Parren, P. A. Marx, and D. R. Burton. 2009. Effective, low-titer antibody protection

646  against low-dose repeated mucosal SHIV challenge in macaques. *Nat. Med.* 15: 951-954.

647 62. Magnani, D. M., T. F. Rogers, N. J. Maness, N. D. Grubaugh, N. Beutler, V. K. Bailey,

648  L. Gonzalez-Nieto, M. J. Gutman, N. Pedreño-Lopez, J. M. Kwal, M. J. Ricciardi, T. A.

649  Myers, J. G. Julander, R. P. Bohm, M. H. Gilbert, F. Schiro, P. P. Aye, R. V. Blair, M. A.

650  Martins, K. P. Falkenstein, A. Kaur, C. L. Curry, E. G. Kallas, R. C. Desrosiers, P. J.

651  Goldschmidt-Clermont, S. S. Whitehead, K. G. Andersen, M. C. Bonaldo, A. A. Lackner,

652  A. T. Panganiban, D. R. Burton, and D. I. Watkins. 2018. Fetal demise and failed

653  antibody therapy during Zika virus infection of pregnant macaques. *Nat. Commun.* 9:

654  1624.

655   63.   Martinez-Navio, J. M., S. P. Fuchs, S. N. Pantry, W. A. Lauer, N. N. Duggan, B. F.

656   Keele, E. G. Rakasz, G. Gao, J. D. Lifson, and R. C. Desrosiers. 2019. Adeno-Associated

657   Virus Delivery of Anti-HIV Monoclonal Antibodies Can Drive Long-Term Virologic

658   Suppression. *Immunity* 50: 567-575.e565.

659   64.   Pauthner, M. G., J. P. Nkolola, C. Havenar-Daughton, B. Murrell, S. M. Reiss, R.

660   Bastidas, J. Prévost, R. Nedellec, B. von Bredow, P. Abbink, C. A. Cottrell, D. W. Kulp,

661   T. Tokatlian, B. Nogal, M. Bianchi, H. Li, J. H. Lee, S. T. Butera, D. T. Evans, L.

662   Hangartner, A. Finzi, I. A. Wilson, R. T. Wyatt, D. J. Irvine, W. R. Schief, A. B. Ward,

663   R. W. Sanders, S. Crotty, G. M. Shaw, D. H. Barouch, and D. R. Burton. 2019. Vaccine-

664   Induced Protection from Homologous Tier 2 SHIV Challenge in Nonhuman Primates

665   Depends on Serum-Neutralizing Antibody Titers. *Immunity* 50: 241-252.e246.

666   65.   Shingai, M., O. K. Donau, R. J. Plishka, A. Buckler-White, J. R. Mascola, G. J. Nabel,

667   M. C. Nason, D. Montefiori, B. Moldt, P. Poignard, R. Diskin, P. J. Bjorkman, M. A.

668   Eckhaus, F. Klein, H. Mouquet, J. C. Cetrulo Lorenzi, A. Gazumyan, D. R. Burton, M. C.

669   Nussenzweig, M. A. Martin, and Y. Nishimura. 2014. Passive transfer of modest titers of

670   potent and broadly neutralizing anti-HIV monoclonal antibodies block SHIV infection in

671   macaques. *J. Exp. Med.* 211: 2061-2074.

672   66.   Welles, H. C., M. F. Jennewein, R. D. Mason, S. Narpala, L. Wang, C. Cheng, Y. Zhang,

673   J. P. Todd, J. D. Lifson, A. B. Balazs, G. Alter, A. B. McDermott, J. R. Mascola, and M.

674   Roederer. 2018. Vectored delivery of anti-SIV envelope targeting mAb via AAV8

675   protects rhesus macaques from repeated limiting dose intrarectal swarm SIVsmE660

676   challenge. *PLoS Pathog.* 14: e1007395.

677   67.   Lu, S., Y. Zhao, W. Yu, Y. Yang, J. Gao, J. Wang, D. Kuang, M. Yang, J. Yang, C. Ma,

678   J. Xu, X. Qian, H. Li, S. Zhao, J. Li, H. Wang, H. Long, J. Zhou, F. Luo, K. Ding, D.

679   Wu, Y. Zhang, Y. Dong, Y. Liu, Y. Zheng, X. Lin, L. Jiao, H. Zheng, Q. Dai, Q. Sun, Y.

680   Hu, C. Ke, H. Liu, and X. Peng. 2020. Comparison of SARS-CoV-2 infections among 3

681   species of non-human primates. *bioRxiv*: https://doi.org/10.1101/2020.04.08.031807.

682   68.   Roark, H. K., J. A. Jenks, S. R. Permar, and M. R. Schleiss. 2020. Animal Models of

683   Congenital Cytomegalovirus Transmission: Implications for Vaccine Development. *J.*

684   *Infect. Dis.* 221: S60-S73.

685   69.   Bondarenko, G. I., D. W. Burleigh, M. Durning, E. E. Breburda, R. L. Grendell, and T.

686   G. Golos. 2007. Passive immunization against the MHC class I molecule Mamu-AG

687   disrupts rhesus placental development and endometrial responses. *J. Immunol.* 179: 8042-

688   8050.

689   70.   Torrealba, J. R., M. Katayama, J. H. Fechner, Jr., E. Jankowska-Gan, S. Kusaka, Q. Xu,

690   J. M. Schultz, T. D. Oberley, H. Hu, M. M. Hamawy, M. Jonker, J. Wubben, G.

691   Doxiadis, R. Bontrop, W. J. Burlingham, and S. J. Knechtle. 2004. Metastable tolerance

692   to rhesus monkey renal transplants is correlated with allograft TGF-beta 1+CD4+ T

693   regulatory cell infiltrates. *J. Immunol.* 172: 5753-5764.

694   71.   Forrest, L., J. Fechner, J. Post, N. Van Asselt, K. Kvasnica, L. D. Haynes, J. Coonen, K.

695   Brunner, W. J. Haynes, C. Little, W. J. Burlingham, P. Hematti, S. Strober, and D. B.

696   Kaufman. 2021. Tomotherapy Applied Total Lymphoid Irradiation and Allogeneic

697   Hematopoietic Cell Transplantation Generates Mixed Chimerism in the Rhesus Macaque

698   Model. *Radiat. Res.* 196: 623-632.

72.   Colantonio, A. D., B. N. Bimber, W. J. Neidermyer, R. K. Reeves, G. Alter, M. Altfeld,

R. P. Johnson, M. Carrington, D. H. O'Connor, and D. T. Evans. 2011. KIR

polymorphisms modulate peptide-dependent binding to an MHC class I ligand with a

Bw6 motif. *PLoS Pathog.* 7: e1001316.

73.   Schafer, J. L., A. D. Colantonio, W. J. Neidermyer, D. M. Dudley, M. Connole, D. H.

O'Connor, and D. T. Evans. 2014. KIR3DL01 recognition of Bw4 ligands in the rhesus

macaque: maintenance of Bw4 specificity since the divergence of apes and Old World

monkeys. *J. Immunol.* 192: 1907-1917.

74.   Rosner, C., P. H. Kruse, M. Hermes, N. Otto, and L. Walter. 2011. Rhesus macaque

inhibitory and activating KIR3D interact with Mamu-A-encoded ligands. *J. Immunol.*

186: 2156-2163.

75.   Banerjee, P., M. Ries, S. K. Janaka, A. G. Grandea, 3rd, R. Wiseman, D. H. O'Connor, T.

G. Golos, and D. T. Evans. 2018. Diversification of Bw4 Specificity and Recognition of a

Nonclassical MHC Class I Molecule Implicated in Maternal-Fetal Tolerance by Killer

Cell Ig-like Receptors of the Rhesus Macaque. *J. Immunol.* 201: 2776-2786.

76.   Nicholas, R. E., K. Sandstrom, J. L. Anderson, W. R. Smith, M. Wetzel, P. Banerjee, S.

K. Janaka, and D. T. Evans. 2022. KIR3DL05 and KIR3DS02 recognition of a

nonclassical MHC class I molecule in the rhesus macaque implicated in pregnancy

success. *Front. Immunol.* 13: 841136.

77.   Barash, S., W. Wang, and Y. Shi. 2002. Human secretory signal peptide description by

hidden Markov model and generation of strong artificial signal peptide for secreted

protein expression. *Biochem. Biophys. Res. Commun.* 294: 835-842.

78. Klenchin, V. A., J. J. Frye, M. H. Jones, M. Winey, and I. Rayment. 2011. Structure-

Function Analysis of the C-terminal Domain of CNM67, a Core Component of the

Saccharomyces cerevisiae Spindle Pole Body. *J. Biol. Chem.* 286: 18240-18250.

79. Nakamura, T., K. D. Yamada, K. Tomii, and K. Katoh. 2018. Parallelization of MAFFT

for large-scale multiple sequence alignments. *Bioinformatics* 34: 2490-2492.

80. Katoh, K., J. Rozewicki, and K. D. Yamada. 2019. MAFFT online service: multiple

sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* 20:

1160-1166.

81. Robinson, O., D. Dylus, and C. Dessimoz. 2016. Phylo.io: Interactive Viewing and

Comparison of Large Phylogenetic Trees on the Web. *Mol. Biol. Evol.* 33: 2163-2166.

82. Bruijnesteijn, J., N. de Groot, A. J. M. de Vos-Rouweler, N. G. de Groot, and R. E.

Bontrop. 2022. Comparative genetics of KIR haplotype diversity in humans and rhesus

macaques: the balancing act. *Immunogenetics*. 74: 313-326.

83. Bruijnesteijn, J., N. G. de Groot, N. Otting, G. Maccari, L. A. Guethlein, J. Robinson, S.

G. E. Marsh, L. Walter, D. H. O'Connor, J. A. Hammond, P. Parham, and R. E. Bontrop.

2020. Nomenclature report for killer-cell immunoglobulin-like receptors (KIR) in

macaque species: new genes/alleles, renaming recombinant entities and IPD-NHKIR

updates. *Immunogenetics* 72: 37-47.

84. Budde, M. L., J. J. Lhost, B. J. Burwitz, E. A. Becker, C. M. Burns, S. L. O'Connor, J. A.

Karl, R. W. Wiseman, B. N. Bimber, G. L. Zhang, W. Hildebrand, V. Brusic, and D. H.

O'Connor. 2011. Transcriptionally abundant major histocompatibility complex class I

alleles are fundamental to nonhuman primate simian immunodeficiency virus-specific

CD8+ T cell responses. *J. Virol.* 85: 3250-3261.

85.   Evans, D. T., J. Peicheng, T. M. Allen, D. H. OConnor, H. Horton, J. E. Venham, M.

Piekarczyk, J. Dzuris, M. Dykhuzen, J. Mitchen, R. A. Rudersdorf, C. D. Pauza, A. Sette,

R. E. Bontrop, R. DeMars, and D. Watkins. 2000. Defintion of Five New Simian

Immunodeficiency Virus Cytoxic T-Lymphocyte Epitores and Their Resrticting Major

Histocompatibility Comple Class I Molecules: Evidence for an Influence on Disease

Progression. *J. Virol.* 74: 7400-7410.

86.   DeGottardi, M. Q., A. Specht, B. Metcalf, A. Kaur, F. Kirchhoff, and D. T. Evans. 2008.

Selective downregulation of rhesus macaque and sooty mangabey major

histocompatibility complex class I molecules by nef alleles of simian immunodeficiency

virus and human immunodeficiency virus type 2. *J. Virol.* 82: 3139-3146.

87.   Collins, K. L., B. K. Chen, S. A. Kalamis, B. D. Walker, and D. Baltimore. 1998. HIV-1

nef protein protects infected cells against killing by cytotoxic T lymphocytes. *Nature*

391: 397-401.

88.   Apps, R., G. Q. Del Prete, P. Chatterjee, A. Lara, Z. L. Brumme, M. A. Brockman, S.

Neil, S. Pickering, D. K. Schneider, A. Piechocka-Trocha, B. D. Walker, R. Thomas, G.

M. Shaw, B. H. Hahn, B. F. Keele, J. D. Lifson, and M. Carrington. 2016. HIV-1 Vpu

Mediates HLA-C Downregulation. *Cell Host Microbe* 19: 686-695.

89.   Bachtel, N. D., G. Umviligihozo, S. Pickering, T. M. Mota, H. Liang, G. Q. Del Prete, P.

Chatterjee, G. Q. Lee, R. Thomas, M. A. Brockman, S. Neil, M. Carrington, B. Bwana,

D. R. Bangsberg, J. N. Martin, E. G. Kallas, C. S. Donini, N. B. Cerqueira, U. T.

O'Doherty, B. H. Hahn, R. B. Jones, Z. L. Brumme, D. F. Nixon, and R. Apps. 2018.

HLA-C downregulation by HIV-1 adapts to host HLA genotype. *PLoS Pathog.* 14:

e1007257.

90.   Vivian, J. P., R. C. Duncan, R. Berry, G. M. O'Connor, H. H. Reid, T. Beddoe, S. Gras,

P. M. Saunders, M. A. Olshina, J. M. Widjaja, C. M. Harpur, J. Lin, S. M. Maloveste, D.

A. Price, B. A. Lafont, D. W. McVicar, C. S. Clements, A. G. Brooks, and J. Rossjohn.

2011. Killer cell immunoglobulin-like receptor 3DL1-mediated recognition of human

leukocyte antigen B. *Nature* 479: 401-405.

Footnotes

This work was supported by Public Health Service grants AI095098, AI155163, AI161816,

AI121135, AI148379 and AI098485 to DTE. Additional support was provided by PHS grant

OD011106 to the WNPRC. DTE is an Elizabeth Glaser Scientist of the Elizabeth Glaser

Pediatric AIDS Foundation and The UW Medical Foundation Professor of Pathology and

Laboratory Medicine.

Abbreviations used in this article: KIR, killer-cell Ig-like receptor; MHC, major

histocompatibility complex; *Mamu, Macaca mulatta*; JNL, Jurkat NFAT luciferase

 Table

 Table I. MHC class I ligands of rhesus macaque lineage II KIRs

KIR Gene[a]	Allotype[b]	MHC class I ligands[c]	Group[d]	Ref[e]
KIR3DL01	KIR3DL01*012	Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01, -B*065:01	Bw4	(73)
KIR3DL02	**KIR3DL02*004**	none detected[f]		This study
KIR3DLw03	**KIR3DLw03*002**	**Mamu-A1*012:01**	A	This study
KIR3DL04	**KIR3DL04*001**	**Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01, -B*065:01**	Bw4	This study
KIR3DL05	KIR3DL05*008	Mamu-A1*002:01, -A3*13:03, -B*036:01, -AG2*01:01, -AG1*05:01	AG/A-related	(72, 74, 75)
KIR3DL06	KIR3DL06*001	Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01, -B*065:01	Bw4	(75)
KIR3DL07	**KIR3DL07*004**	**Mamu-AG2*01:01, -AG2*01:02, -AG6*04:01, -B*045:03**	AG/A-related	This study
KIR3DL08	KIR3DL08*002	Mamu-B*002:01, -B*007:01, -B*022:01, -B*041:01, -B*065:01	Bw4	(75)
KIR3DL10	**KIR3DL10*002**	**Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01, -B*065:01**	Bw4	This study
KIR3DL11	**KIR3DL11*001**	**Mamu-B*007:01, -B*041:01, -B*065:01**	Bw4	This study
KIR3DLw34	**KIR3DLw34*002**	none detected[f]		This study
KIR3DS01	**KIR3DS01*003**	**Mamu-AG2*01:01, -AG6*04:01**	AG	This study
KIR3DS02	**KIR3DS02*004**	**Mamu-AG2*01:01, -AG2*01:02,, -AG6*04:01, -AG1*05:01, -B*045:03**	AG/A-related	(76), This study
KIR3DS03	**KIR3DS03*003**	none detected[f]		This study
KIR3DS04	**KIR3DS04*003**	**Mamu-B*022:01, -B*041:01, -B*065:01**	Bw4	This study
KIR3DS05	**KIR3DS05*002**	none detected[f]		This study
KIR3DS06	**KIR3DS06*002**	**Mamu-AG2*01:01, -AG1*05:01, -AG6*04:01**	AG	This study
KIR3DSw07	**KIR3DSw07*001**	**Mamu-AG2*01:01, -AG6*04:01**	AG	This study
KIR3DSw08	**KIR3DSw08*001**	none detected[f]		This study
KIR3DSw09	**KIR3DSw09*003**	**Mamu-AG2*01:01, -AG2*01:02, -AG1*05:01, -B*045:03**	AG/A-related	This study
KIR3DSw39	KIR3DSw39*002	Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01, -B*065:01	Bw4	(75)

 [a]Rhesus macaque lineage II KIR genes.
 [b]Representative KIR allotypes. KIR allotypes tested in this study are indicated in bold.
 [c]Novel MHC class I ligands defined in this study are indicated in bold.
 [d]Group of MHC class I ligands.
 [e]References for previously reported KIR ligands.
 [f]Supplemental Figure 8

	Figure Legends

FIGURE 1. Rhesus macaque KIRs and KIR-CD3ζ expression on JNL cells. (**A**) Alignment of

amino acid sequences for the D0, D1 and D2 domains of rhesus macaque KIRs. Shaded

sequences correspond to predicted MHC class I contact sites based on the crystal structure of

HLA-B*57 in complex with human KIR3DL1 (90). Positions of identity are indicated by periods

and amino acid differences are identified by their single-letter code. (**B**) KIR-CD3ζ-transduced

JNL cells were stained with anti-Flag antibody and Near-IR LIVE/DEAD stain. After excluding

dead cells, the fluorescence intensity of KIR-CD3ζ (Flag-tag) staining on KIR-CD3ζ-transduced

JNL cells (open) was compared to background staining on parental JNL cells (shaded).

FIGURE 2. KIR3D04*001, KIR3DL10*002, KIR3DL11*001 and KIR3DS04*003 interact with

multiple Mamu-Bw4 ligands. (**A**) KIR3DL04*001-, KIR3DL10*002-, KIR3DL11*001- and

KIR3DS04*003-CD3ζ JNL cells were incubated with 721.221 cells expressing the indicated

Mamu-B molecules. The bar graphs represent the mean and standard deviation (error bars) of

luciferase activity (RLU) from triplicate wells of KIR-CD3ζ JNL cells incubated with Mamu-B +

721.221 cells (blue), parental 721.221 cells (red) and anti-Flag plus anti-mouse antibodies (X-

link, gray). Asterisks denote significant differences by one-way ANOVA with Dunnett test (**

p=0.0002, and *** p < 0.0001). These results are representative of at least three independent

experiments. (**B**) Parental and Mamu-B+721.221 cells were stained with Near-IR LIVE/DEAD

dye, KIR3DL04*001-, KIR3DL10*002-, KIR3DL11*001- and KIR3DS04*003-Fc followed by

goat anti-mouse IgG and MHC class I-specific antibody (W6/32).

FIGURE 3. KIR3DL07*004, KIR3DS01*003, KIR3DS02*004, KIR3DS06*002,

KIR3DSw07*001 and KIR3DSw09*003 interact with Mamu-AG and -B*045:03. (**A**)

KIR3DL07*004-, KIR3DS01*003-, KIR3DS02*004-, KIR3DS06*002-, KIR3DSw07*001- and

KIR3DSw09*003-CD3ζ JNL cells were incubated with 721.221 cells expressing the indicated

Mamu-AG or -B molecules. The bar graphs represent the mean and standard deviation (error

bars) of luciferase activity (RLU) from triplicate wells of KIR-CD3ζ JNL cells incubated with

Mamu-AG+721.221 cells (purple), Mamu-B+721.221 (blue), parental 721.221 cells (red) and

anti-Flag plus anti-mouse antibodies (X-link, gray). Asterisks denote significant differences by

one-way ANOVA with Dunnett test (** $p < 0.005$, and *** $p < 0.0001$). These results are

representative of at least three independent experiments. (**B**) Parental, Mamu-AG and Mamu-

B+721.221 cells were stained with Near-IR LIVE/DEAD dye, KIR3DL07*004-,

KIR3DS01*003-, KIR3DS02*004-, KIR3DS06*002-, KIR3DSw07*001- and

KIR3DSw09*003-Fc followed by goat anti-mouse IgG and MHC class I-specific antibody

(W6/32).

FIGURE 4. Mamu-B*045:03 has an α1-domain similar to Mamu-A molecules. (**A**) Amino acid

sequence alignment for the α1- and α2-domains of rhesus macaque MHC class I molecules.

Residues 77–83 corresponding to the Bw4/Bw6 motif are underlined and predicted KIR contact

sites based on the crystal structure of HLA-B*57 in complex with human KIR3DL1 are shaded

(90). Positions of identity are indicated by periods and amino acid differences are identified by

their single-letter code. (**B & C**) Phylogenetic trees of nucleotide sequence coding for the α1-α3

domains (**B**) or the α1 domain (**C**) of selected Mamu-A, -B and -AG molecules. Neighbor-

joining analysis was performed using MAFFT software (version 7, publicly available at

https://mafft.cbrc.jp/alignment/server/). Bootstrap support of greater than 50% is indicated.

FIGURE 5. KIR3DLw03*002 interacts with Mamu-A*012:01. (**A**) KIR3DLw03*002-CD3ζ

JNL cells were incubated with 721.221 cells expressing the indicated rhesus macaque MHC class

I molecules. The bar graphs represent the mean and standard deviation (error bars) of luciferase

activity (RLU) from triplicate wells of KIR-CD3ζ JNL cells incubated with Mamu-

A*012:01+721.221 (green), nonclassical (purple), parental 721.221 cells (red) and anti-Flag plus

anti-mouse antibodies (X-link, gray). Asterisks denote significant differences by one-way

ANOVA with Dunnett test (*** $p < 0.0001$). These results are representative of at least three

independent experiments. (**B**) Parental and Mamu-A1*012:01+721.221 cells were stained with

Near-IR LIVE/DEAD dye, and KIR3DLw03*002-Fc followed by goat anti-mouse IgG and

MHC class I-specific antibody (W6/32).

FIGURE 6. Phylogenetic clustering of rhesus macaque KIRs reflect their patterns of ligand

recognition. Neighbor-joining analysis of nucleotide sequences coding for the D0-D2 domains of

human and rhesus macaque KIRs was performed using MAFFT software. Bootstrap support of

greater than 50% is indicated. Human KIR3DL1 and KIR3DL02 were used included as an

outgroup. Rhesus macaque KIRs are color coded according to their MHC class I ligands:

unknown (gray), Mamu-Bw4 (green), Mamu-AG (dark blue), and Mamu-A1*012:01 (purple).

Figure 1

A

D0 Domain

```
                    10        20        30        40        50        60        70        80        90        100
Consensus     HTGGQDKTFLSARPSAVVPQGGHVTLRCXYRRGLNNFTNFTLYKDDRSHVPIFHSRIFQESFLMGPVTPAHAGTYRCRGSYPHSPTEWSALSDPLAIRVT
KIR3DL02*004  Y.....N....W...L.........H..G.F.---..........................................L.....TP.N......
KIR3DLw03*002 Y.....N....W...P.........H..G.F.---..........V...............................TP.N......
KIR3DL04*001  ...........L.............Y...H..................PG...HQ..........D...........Q...........M..
KIR3DL07*004  .........F...............Y..D..............................................................M..
KIR3DL10*002  Y.........VQ...L........Y...F...............N....Q..........D...........................M..
KIR3DL11*001  ......S...W...P.........Q.H.....F...............PG...HQ..................................M..
KIR3DLw34*002 Y.....N......W...P.......H..G.F.---.........I..............................................
KIR3DS01*003  ....................FQ.Y.H...............Q.................................................
KIR3DS02*004  .........F...............Y..D..............................................................M..
KIR3DS03*003  ......I...VQ...L....M....H....Y...............N..........D...........................M..
KIR3DS04*003  ......N..........Q.H.....F...............V.............................................M..
KIR3DS05*002  .....NA....W...P.........H..G.F.---....................................................P.N......
KIR3DS06*002  ...A...N..F.W...P.....L....Y.H.........................................................M..
KIR3DSw07*001 ................L.......Y..D..........................V..N................................
KIR3DSw08*001 .......S.........L.......H...................V...........................................H....
KIR3DSw09*002 ...D......F................Y..D..........................................................
```

D1 Domain

```
                    110       120       130       140       150       160       170       180       190       200
Consensus     GVHRKPSLLALPGPLVKSGETVTLQCSSDTVFXHFFLHSEVTFEXPLHLVGELHGGGSQANYSINSTTSDLAGTYRCYGSVTHSPYVLSAPSDPLDIVIT
KIR3DL02*004  ...K.....................E....R...L.E................MM......F.....F..................
KIR3DLw03*002 ...K..............I......E....R...L.E...........S.......MMYT.....F..................
KIR3DL04*001  ..................I.G.........EL...........................................
KIR3DL07*004  ......................N...K...................................D....
KIR3DL10*002  ..................I.E.........K....................EE.F..............N....
KIR3DL11*001  ...............G.........K.....................E.F..............N....
KIR3DLw34*002 ...K.....................E....R...L.E...........K.................F..........
KIR3DS01*003  ...K...............G.........K.....................K.............
KIR3DS02*004  ...................G.........K......R........KM......................H......
KIR3DS03*003  ...................G.........K.........K...E.................
KIR3DS04*003  ..................I...I.G.........EL...................................T....
KIR3DS05*002  ...K.....................E....R...L.E...........F...F.....F..................
KIR3DS06*002  ...............L.........E.......N..K...........K.E......................R....
KIR3DSw07*001 .................D.......G.........K.................K....E..........
KIR3DSw08*001 ..................I.G.........N.EL...................................T....
KIR3DSw09*002 ..................I.......E.......N..K.........................
```

D2 Domain

```
                    210       220       230       240       250       260       270       280       290
Consensus     GKYEKPSLSAQPGPTVQAGENVTLSCSSXNSFDMYHLSREGEARELSLSAVPSVNGTFQADFPLGPATHGGTYRCFGSFRTAPYKWSDPSDPLPVSVT
KIR3DL02*004  .................I.......RC................................................Q.......
KIR3DLw03*002 .L......................RC............R....G...................H......
KIR3DL04*001  ...................Q....................S...............Q.......H....
KIR3DL07*004  ...................Q.......................................
KIR3DL10*002  ...................Q....S...V...R....G.........I.......IT.....NS...L....
KIR3DL11*001  ..............I.......RC............................
KIR3DLw34*002 .L......................RC............R....G...............N.....H.....
KIR3DS01*003  .L...........D....QI..............................S...Q.......HI...
KIR3DS02*004  .L......................RR................R....G.........H.T......H......
KIR3DS03*003  ...................Q............G....Q..............
KIR3DS04*003  .................RC...........I.......................T...............K
KIR3DS05*002  .L......................RR....................-............TS....H....
KIR3DS06*002  .L...................Q............A..R....G.........H.T......H....
KIR3DSw07*001 ...................Q...........G.....Q..R..................Q.......
KIR3DSw08*001 ...................Q...........G.....Q.......N.................
KIR3DSw09*002 .L......................RC................R....G....................H....
```

B

KIR3DL02*004 KIR3DLw03*002 KIR3DL04*001 KIR3DL07*004 KIR3DL10*002 KIR3DL11*001 KIR3DLw34*002 KIR3DS01*003

KIR3DS02*004 KIR3DS03*003 KIR3DS04*003 KIR3DS05*002 KIR3DS06*002 KIR3DSw08*001 KIR3DSw07*001 KIR3DSw09*003

Percent Max

KIR-CD3ζ →

Figure 2

A

B

Figure 3

A

B

Figure 4

A

Figure 5

A

KIR3DLw03*002

B

Figure 6