Broad Recognition of Mamu-Bw4 and -A-related MHC class I Ligands by Rhesus Macaque Killer-Cell Immunoglobulin-Like Receptors

Jennifer L. Anderson[*], Kjell Sandstrom[*], Rachel E. Nicholas[*], Willow R. Smith[*], Molly Wetzel[*], Vadim A. Klenchin[*], and David T. Evans[*,†]

[*]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison, Madison, WI

[†]Wisconsin National Primate Research Center, University of Wisconsin-Madison, Madison, WI

This work was supported by Public Health Service Grants AI095098, AI155163, AI161816, AI121135, AI148379 and AI098485 (to DTE) and OD011106 (to the WNPRC). DTE is an Elizabeth Glaser Scientist of the Elizabeth Glaser Pediatric AIDS Foundation and The UW Medical Foundation Professor of Pathology and Laboratory Medicine.

Corresponding author:
David T. Evans
dtevans2@wisc.edu
Phone: (608) 265-1485
Fax: (608) 265-8084

Running Title: Rhesus Macaque KIR Ligands

Abstract

Definition of the MHC class I ligands of rhesus macaque KIRs is fundamental to NK cell

biology in this species as an animal model for infectious diseases, reproductive biology and

transplantation. To provide a more complete foundation for studying NK cell responses in this

species, rhesus macaque KIRs representing common allotypes of lineage II *KIR* genes were

tested for interactions with MHC class I molecules representing diverse *Mamu-A, -B, -E, -F, -I*

and *-AG* alleles. KIR-MHC class I interactions were identified by co-incubating reporter cell

lines bearing chimeric KIR-CD3ζ receptors with target cells expressing individual MHC class I

molecules and were corroborated by staining with KIR-Ig Fc domain fusion proteins. Ligands for

10 KIRs of previously unknown specificity were identified that fell into two general categories;

interactions with multiple Mamu-Bw4 molecules (KIR3DL04*001, KIR3DL10*002,

KIR3DL11*001 and KIR3DS04*003) or with Mamu-A-related molecules (KIR3DL07*004,

KIR3DS01*003, KIR3DS02*004, KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003).

Among the latter group, three KIRs (KIR3DL07*004, KIR3DS02*004 and KIR3DSw09*003)

also interacted with Mamu-B*045:03, a hybrid molecule with Mamu-A α1-domain sequences.

Both groups include inhibitory and activating receptors that vary in the specificity and strength

of their MHC class I interactions. However, whereas the majority of KIRs that recognize Mamu-

Bw4 ligands are inhibitory, most of the KIRs that recognize Mamu-A-related ligands are

activating. These findings more than double the number of rhesus macaque KIRs with defined

MHC class I ligands and reveal an overlapping, but nonredundant, pattern of ligand recognition

that supports extensive functional diversification of these receptors.

Introduction

Natural killer cell responses in humans and other primate species are regulated in part by interactions between two highly polymorphic sets of molecules; the killer cell Ig-like receptors (KIRs) expressed on the surface of NK cells and their MHC class I ligands on target cells (1, 2). As indicated by their nomenclature, KIRs typically have two or three extracellular Ig-like domains (2D or 3D) and either long (L) or short (S) cytoplasmic tails. KIRs with long cytoplasmic tails contain a pair of ITIMs that transduce inhibitory signals, whereas those with short cytoplasmic tails associate with adaptor molecules such as DAP12 or FcεRIγ to transduce activating signals (2, 3). Inhibitory KIRs normally suppress NK cell responses through interactions with MHC class I ligands on the surface of heathy cells. However, if these interactions are somehow perturbed, for instance as a result of MHC class I downregulation in virus-infected cells or deletion of MHC class I genes in tumor cells, the loss of inhibitory signals can trigger an NK cell response (4-8). Activating KIRs can also interact with MHC class I molecules on unhealthy cells to directly trigger an NK cell response, although their molecular mechanisms are not as well defined (2, 9-11).

Polymorphisms in the *KIR* and *HLA class I* genes have been associated with differences in the course of infection for a number of human viral pathogens and with the outcome of cancer immunotherapy (12-19). *KIR* and *HLA class I* polymorphisms have also been linked to complications during pregnancy and to the success of hematopoietic stem cell and solid organ transplantation (20-23). However, studies to address the immunological mechanisms underlying these observations have been hampered by the lack of a suitable animal model. Mice and other rodents do not have KIRs, but instead express C type lectin-like molecules on their NK cells as polymorphic receptors for MHC class I ligands (2, 24). Moreover, as a consequence of the rapid

pace of KIR and MHC class I evolution (25-27), it is not possible to predict KIR ligands in Old

World monkeys on the basis of sequence similarity with human KIRs.

In contrast to humans, which have three classical *HLA class I* genes (*HLA-A, -B* and *-C*),

Old World monkeys express an expanded array of polymorphic MHC class I genes related to

HLA-A and *-B*. However, these species do not have an *HLA-C* ortholog, since *MHC-C* arose in

hominids as a duplication of an *MHC-B* gene after their divergence from Old World monkeys

(28, 29). The *MHC class I* haplotypes of Old World monkeys are polygenic, and in the rhesus

macaque typically express two or three *Macaca mulatta (Mamu)-A* genes and four to eleven

Mamu-B genes (30-32). Depending on their transcriptional abundance, the *MHC class I* genes of

macaques can be further classified as "major" or "minor" loci (32). Products of the major genes

exhibit the greatest polymorphism and present peptides to virus-specific $CD8^+$ T cells (33),

suggesting functional similarities to HLA-A and -B in humans.

Four non-classical *MHC class I* genes have also been identified in Old World monkeys.

Orthologs of *HLA-E* and *-F* (*Mamu-E* and *-F* in rhesus macaques) are well conserved and serve

similar functions as their human counterparts (34-37). Macaques and other Old World monkeys

also have an ortholog of *HLA-G*; however the *MHC-G* genes of these species have accumulated

numerous stop-codons and no longer encode functional proteins (38, 39). In lieu of a functional

MHC-G locus, Old World monkeys have evolved duplicated *MHC-A* genes, designated *MHC-*

AG, that appear to serve a similar function (40-43). Like HLA-G, MHC-AG has limited

polymorphism, a truncated cytoplasmic tail, and is only expressed on placental trophoblasts that

form a barrier between the mother and developing fetus (40, 41, 44). Mamu-AG is therefore

thought to contribute to pregnancy success in rhesus macaques through interactions with

receptors on NK cells and myeloid cells that stimulate placental vascularization, facilitate fetal

tolerance, and protect the maternal-fetal interface from invading microorganisms (44-46). Rhesus

macaques also express Mamu-I, which is the product of a fixed duplication of an *MHC-B* gene

with limited sequence variation (47).

In concert with the expansion of their *Mamu-A* and *-B* genes, macaques have evolved an

expanded complement of lineage II *KIRs* (26, 48-52). Whereas humans only have two lineage II

KIR genes that encode KIR3DL1/S1 and KIR3DL2 as receptors for HLA-Bw4 and –A3/A11,

respectively, rhesus macaques have at least 19 lineage II *KIRs* that encode inhibitory and

activating receptors for Mamu-A and -B molecules (50, 52-56). Furthermore, in contrast to

humans, which have multiple lineage III *KIR* genes encoding KIR2DL/S receptors for HLA-C1

and -C2 ligands (1, 2), macaques only have one lineage III *KIR* that expresses a single domain

KIR (KIR1D) of uncertain function (52). Thus, while human NK cells are predominantly

regulated by KIR2DL/S-HLA-C interactions, macaque NK cells are dependent on KIR3DL/S

interactions with Mamu-A and -B ligands (52).

The rhesus macaque has become an increasingly valuable animal model for infectious

diseases (57-68), reproductive biology (44, 69), and transplantation (70, 71). Although MHC

class I ligands have been identified for a few macaque KIRs (72-75), ligands for the majority of

these receptors remain undefined. We therefore performed a systematic survey of MHC class I

interactions for common KIR allotypes representing 14 predicted lineage II *KIR* genes in Indian-

origin rhesus macaques. By measuring the responses of KIR-CD3ζ-transduced reporter cell lines

to target cells expressing individual MHC class I molecules and confirming interactions by

staining with KIR-Ig Fc domain (KIR-Fc) fusion proteins, we identified MHC class I ligands for

10 rhesus macaque KIRs of previously undefined specificity. This represents the most

comprehensive survey of ligand recognition by macaque KIRs to date and reveals extensive

diversification of the lineage II KIRs that interact with Mamu-Bw4 and -A-related ligands.

Materials and Methods

Jurkat NFAT luciferase reporter cells expressing chimeric KIR-CD3ζ receptors

Complementary DNA sequences encoding the leader peptide of Mamu-KIR3DL05*008

followed by a Flag®-tag (DYKDDDDK) and the D0, D1, D2 and stem domains of rhesus

macaque KIRs were synthesized (Integrated DNA Technologies) and cloned into pQCXIP in-

frame with sequences encoding the transmembrane and cytoplasmic domains human CD3ζ.

Retroviral vectors were packaged into VSV G–pseudotyped murine leukemia virus (MLV)

particles by co-transfecting GP2-293 cells (Clontech Laboratories) with pQCXIP constructs and

pVSV-G (Clontech Laboratories). Jurkat NFAT luciferase (JNL) cells (Signosis) were

transduced via spinoculation for one hour with filtered supernatant collected from transfected

GP2-293 cells and 2 d later were placed under selection in RPMI medium supplemented with

10% FBS, L-glutamine, penicillin and streptomycin (R10 medium) plus 0.1 mg/ml hygromycin

(Invitrogen) to maintain the luciferase reporter gene and 0.5 µg/ml puromycin (Invitrogen) to

select for the KIR -CD3ζ receptor. The puromycin concentration in the medium was gradually

increased to 1.0 µg/ml over 3-4 weeks to eliminate untransduced cells and then maintained at a

concentration of 0.5 µg/ml. The GenBank accession numbers for rhesus macaque KIR and

human CD3ζ sequences are as follows: *Mamu-KIR3DL02*004:01* (LT963635.1), -

*KIR3DL04*001:02* (GU299490.1), -*KIR3DL07*004* (EU419060.1), -*KIR3DL10*002:01*

(GU112259.1), -*KIR3DL11*001* (GU112271.1), -*KIR3DLw34*002* (previously -

*KIR3DLw03*003*; EU419031.1), -*KIR3DS01*003* (GU564161.1), -*KIR3DS02*004:02*

(GU014296.1), -*KIR3DS03*003* (EU702454.1), -*KIR3DS04*003:01* (EU419029.1), -

*KIR3DS05*002:01* (GU112262.1), -*KIR3DS06*002:02* (GU112298.1), -*KIR3DSw07*001*

(GU112272.1), -*KIR3DSw09*003* (MF164924.1) and *CD3ζ* (J04132.1).

Rhesus macaque MHC class I-expressing 721.221 cells

Codon-optimized cDNA sequences encoding rhesus macaque MHC class I molecules were

cloned into pQCXIP or pQCXIN. Retroviral vectors were packaged into VSV-G-pseudotyped

MLV particles as described above. 721.221 cells were transduced via spinoculation for one hour

with filtered supernatant collected from transfected GP2-293 cells and placed under selection in

R10 medium containing 0.4 mg/ml puromycin or 0.5 mg/ml G418, depending on the vector, to

select for MHC class I expression. The concentrations of puromycin and G418 in the medium

were gradually increased to 1.0 µg/ml and 1.0 mg/ml, respectively, over 3-4 weeks to eliminate

untransduced cells and then maintained at a concentration of 0.4 µg/ml and 0.5 mg/ml. The

GenBank accession numbers for rhesus macaque MHC class I sequences are as follows: *Mamu-*

*A1*001:01:01:01* (LT908117), *-A1*002:01:01:01* (LN899621), *-A1*004:01:01:01*

(FN396407.1), *-A1*008:01:01:01* (LN899628), *-A1*011:01:01:01* (AJ542579), *-*

*A1*012:01:01:01* (AF157398.1), *-A2*05:02:01:01* (LN899639), *-A2*05:04:01:01*

(LT908121.1), *-A3*13:02:01:01* (LN899654.1), *-A3*13:03:01:01* (AF157401), *-*

*A4*14:03:01:01* (GU080236.1), *-A6*01:03:02:01* (EF602318.1), *-B*001:01:02:01*

(LM608018), *-B*002:01:01:01* (LN851852), *-B*005:01:01:01* (LM608023), *-B*007:01:01:01*

(U41829.1), *-B*008:01:01:01* (U41830), *-B*015:01* (AM902541.1), *-B*017:01:01:01*

(AF199358), *-B*022:01:01:01* (LN899675), *-B*036:01:01:01* (AJ556886.1), *-B*041:01:01:01*

(LN899682), *-B*043:01:01:01* (AJ556893), *-B*045:03:01:01* (LM608047), *-B*056:01:01:01*

(GQ902079.1), *-B*065:01:01:01* (AJ620416.1), *-I*01:01:01* (LT908864), *-E*02:01:02*

(previously E*02:01; NM_001114966.1) *-F*01:01* (LT899414), *-AG2*01:01* (previously

AG*01:01; U84783.1), *-AG3*02:01:01:01* (U84785.1), *-AG3*02:02:01* (U84786.1), *-*

*AG3*03:01:01:01* (U84787.1), *-AG3*03:02* (U84789.1), *-AG3*03:03* (KF855159.1), *-*

AG3*03:04 (KF855160.1), and *-AG1*05:01* (FJ409466). Only partial-length cDNA sequences

were available for *Mamu-AG3*03:03*, *-AG3*03:04*, and *-AG1*05:01*. For constructs expressing

these alleles, sequences encoding the α3-, transmembrane and cytoplasmic domains were

inferred from the *Mamu-AG* consensus sequence.

KIR-Fc fusion proteins

Complementary DNA sequences encoding the artificial secrecon leader peptide

(MWWRLWWLLLLLLLLLWPMVWA), the D0, D1, D2 and stem domains of macaque KIRs,

and the Fc domain of murine IgG2A were synthesized (Integrated DNA Technologies) and

cloned downstream of the CMV promoter in pCIW using unique *Hind* III and *Not* I sites.

Subsequent *KIR* sequences were introduced by primer overlap extension. These sequences were

PCR amplified to create a product with arms complementary to the insertion site of the vector (5'

secrecon, 3' msFc), and then employed as a megaprimer to amplify the entire plasmid with Pfu

Ultra II Fusion HS (Agilent). The template vector was eliminated by digestion with Fermentas'

FastDigest Dpn I for one hour at 37 °C. After sequence confirmation, plasmid DNA maxipreps

were prepared for each construct. Expi-293T cells (7.5×10^8) were transfected with the pCIW-

KIR-Fc plasmid DNA (250 µg) using the ExpiFectamine 293 Transfection Kit (Life

Technologies) according to the manufacturer's recommendations. Five days after transfection,

the supernatant was collected, filtered through a 0.45 µM PES filter (Fisher Scientific), and

loaded on a Protein A column (GE Healthcare). After washing the columns three times with

PBS, KIR-Fc fusion proteins were eluted in gentle Ag/Ab buffer (Thermo Scientific) and buffer

exchanged with 50 mM Sodium citrate using an Amicon Ultra-15 Ultracel 50K Centrifugal Filter

unit (Millipore). The final concentrations of the purified KIR-Fc fusion proteins were calculated

by dividing the UV absorbance at 280 nm by the theoretical extinction coefficient obtained from

ExPASy ProParam (https://web.expasy.org/protparam/). After adding sodium azide (3 mM), the

KIR-Fc fusion stocks were snap frozen in liquid nitrogen and stored at -80 °C.

Flow cytometry

KIR-CD3ζ expression on the surface of JNL cells and MHC class I expression on 721.221 cells

was confirmed by staining with PE-conjugated anti-Flag® (Miltenyi Biotec) and pan-HLA class

I-specific antibodies (clone W6/32; Life Technologies), respectively. These cells were then

stained with by Near-IR fluorescent dye (Invitrogen), washed twice in PBS plus 1% FBS (FACS

buffer), and fixed in 2% paraformaldehyde PBS. To assess KIR binding, MHC class I-transduced

721.221 cells were stained for 30 minutes at room temperature with Near-IR fluorescent dye,

washed twice with FACS buffer and stained for 30 minutes at room temperature with KIR-Fc

fusions proteins (10-20 µg). The cells were then washed twice with FACS buffer, stained for 45

minutes with Alex 647-conjugated goat anti-mouse IgG (Jackson ImmunoResearch), washed

twice more, and stained for 30 minutes at room temperature with the PE-conjugated pan-MHC

class I-specific antibody W6/32. The cells were then washed again and fixed in 2%

paraformaldehyde PBS. Data was collected using a FACSymphony™ A3 Cell Analyzer (BD

Biosciences) and analyzed using FlowJo version 10.8.0 software (Becton Dickinson). After

gating on viable cells, KIR-CD3ζ (Flag-tag) and MHC class I staining was compared to parental

JNL and 721.221 cells, respectively. KIR-MHC class I binding was similarly assessed by

comparing KIR-Fc versus W6/32 staining on the surface of MHC class I-transduced 721.221

cells to their levels of non-specific staining on parental 721.221 cells.

KIR-CD3ζ⁺ JNL reporter assays

KIR-CD3ζ JNL cells (1x10□) and MHC class I-transduced 721.221 cells (1x10□) were co-

cultured overnight at 37°C and 5% CO□ in R10 medium without antibiotics. KIR-CD3ζ□ JNL

cells were also incubated with parental 721.221 cells as negative control and with anti-Flag-tag

(5 µg/ml) (GenScript) and goat anti-mouse (10 µg/ml) (Poly4053, Biolegend) antibodies as a

positive control. Each combination was tested in triplicate wells (100 µl/well). After 12-18

hours, BriteLite Plus luciferase substrate (PerkinElmer) was added to each well (100 µl/well) and

the relative light units (RLU) of luciferase activity was measured using a VICTOR X4 multiplate

reader (PerkinElmer).

Alignments and Phylogenetic trees

The predicted amino acid sequences of KIR and MHC class I molecules were aligned using

Geneious Prime® 2021.2.2 (Biomatters Ltd) and the text format was exported to Microsoft Word

for figure construction. Neighbor-joining analysis of CDS nucleotide sequences for the α1-α3

domains of Mamu-A and -B and the D0-D2 domains of KIR molecules was performed using

MAFFT (version 7, publicly available at https://mafft.cbrc.jp/alignment/server/) (76, 77). For the

analysis of KIR sequences, human lineage II KIRs were included as an outgroup

(KIR3DL1*0010101, NP_037421.2 and KIR3DL02*0010101, NP_006728.2). All analyses

utilized G-INS-I iterative refinement with 1000 bootstrap replicates. Trees were viewed using

Phylo.io (1.0.0) (78).

Statistical analysis

The luciferase induction by KIR-CD3ζ☐ JNL cells (RLU data) was analyzed by one-way

ANOVA followed by Dunnett's test for multiple comparisons of mean responses to each of the

MHC class I-expressing 721.221 cells with mean responses to the parental 721.221 cells as a

negative control. Statistical analyses were performed using GraphPad Prism for Mac OS version

9.2.0.

Results

Extensive sequencing and segregation analyses of *KIR* transcripts support the existence of at

least twenty-two *KIR* genes in Indian-origin rhesus macaques (53-56). Although our group and

others have identified ligands for a few rhesus macaque KIRs (72-75), ligands for most of the

receptors in this species remain undefined. To provide a more complete foundation for studying

NK cell biology in the rhesus macaque, we investigated the MHC class I interactions of 14 KIRs

representing common alleles of lineage II genes in animals of Indian descent for which ligands

have not been previously reported (Fig. 1A). Jurkat cells harboring an NFAT-inducible luciferase

reporter gene (JNL cells) were transduced with retroviral vectors expressing KIR-CD3ζ chimeric

receptors consisting of the extracellular domains of macaque KIRs (D0, D1, D2 and stem

domains) fused to the transmembrane and cytoplasmic domains of human CD3ζ. To confirm

surface expression, a Flag-tag was appended to the N-terminus of the D0 domain of each

receptor (Fig. 1B). After an overnight incubation with HLA class I-negative 721.221 cells

transduced with individual rhesus macaque MHC class I molecules (Supplemental Figs. 1-3),

ligand recognition was detected by the MHC class I-dependent upregulation of luciferase by

KIR-CD3ζ JNL cells (Supplemental Figs. 4-5). KIR-MHC class I interactions were corroborated

by staining MHC class I-expressing 721.221 cells with KIR-Fc fusion proteins consisting of the

extracellular domains of macaque KIRs fused to the Fc domain of murine IgG2A (Supplemental

Figs. 6-8).

Mamu-Bw4 recognition by multiple rhesus macaque KIRs

KIR3DL04*001- and KIR3DL10*002-CD3ζ JNL cells responded strongly to 721.221 cells

expressing Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01 and -B*065:01 (Fig. 2A). The

magnitude of these responses was generally reflected by the relative intensity of

KIR3DL04*001- and KIR3DL10*002-Fc staining on the surface of Mamu-B*007:01+, -

B*022:01+, -B*041:01+, -B*043:01+ and -B*065:01+ 721.221 cells (Fig. 2B). Similar

responses were also observed for two other KIRs. KIR3DL11*001-CD3ζ JNL cells responded to

Mamu-B*007:01, -B*041:01 and -B*065:01, and KIR3DS04*003-CD3ζ JNL cells responded to

Mamu-B*22:01, -B*041:01 and -B*065:01 (Fig. 2A). These interactions were further

corroborated by KIR-Fc staining of 721.221 cells expressing these molecules (Fig. 2B). KIR-Fc

staining affords a comparison of the relative strength of MHC class I interactions across different

KIRs that is not possible using independently derived JNL cell lines that may vary in KIR-CD3ζ

expression and luciferase induction (Fig. 2B). In the case of KIR3DS04*003, a lower binding

affinity for fewer ligands may reflect a role for MHC class I-bound peptide or other cellular

factors in reaching a binding threshold necessary to trigger an NK cell response through this

activating receptor.

Mamu-B*007:01, -B*022:01, -B*041:01, -B*043:01 and -B*065:01 share a Bw4 motif

at residues 77-83 (Supplemental Fig. 2), which was previously shown to contribute to

recognition by KIR3DL01 as well as three other rhesus macaque KIRs (73, 75). The

identification of Mamu-Bw4 ligands for KIR3DL04*001, KIR3DL10*002, KIR3DL11*001 and

KIR3DS04*003 doubles the number of rhesus macaque KIRs known to interact with this group

of molecules and reveals an extraordinary expansion of the lineage II KIRs in macaques with

Bw4 specificity (Table I). Differences in the particular Mamu-Bw4 molecules recognized by

these KIRs and the relative strength of these interactions suggests extensive functional

diversification of these receptors.

Multiple KIRs recognize Mamu-AG molecules

Six rhesus macaque KIRs were found to interact with Mamu-AG2*01:01, a non-classical MHC

class I molecule expressed by fetal trophoblasts at the maternal-fetal interface of the placenta

(40). JNL cells expressing KIR-CD3ζ chimeras for KIR3DL07*004, KIR3DS01*003,

KIR3DS02*004, KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003 all responded

strongly to 721.221 cells expressing Mamu-AG2*01:01 but not other Mamu-AG allotypes (Fig.

3A). The binding of these KIRs to Mamu-AG2*01:01 was further corroborated by KIR-Fc

staining of Mamu-AG2*01:01+721.221 cells (Fig. 3B). KIR-Fc staining also revealed low but

detectable interactions with Mamu-AG1*05:01 for KIR3DS02*004, KIR3DS06*002 and

KIR3DSw09*003 that were not observed in cellular assays with KIR-CD3ζ JNL cells (Fig. 3B).

In this case, the detection of weak binding interactions with Mamu-AG*01*05:01 that were not

detectable with KIR-CD3ζ JNL cells may reflect somewhat greater sensitivity of KIR-Fc

staining. Together with the previous identification of Mamu-AG ligands for KIR3DL05 and

KIR3DS02*004 (75, 79), these findings bring the number of rhesus macaque KIRs known to

interact with Mamu-AG to seven (Table I).

*Mamu-B*045:03 recognition by KIR3DL07*004, KIR3DS02*004 and KIR3DSw09*003*

In addition to Mamu-AG, JNL cells expressing KIR3DL07*004-, KIR3DS02*004- and

KIR3DSw09*003-CD3ζ also responded to Mamu-B*045:03 (Fig. 4A). The binding of these

KIRs to Mamu-B*045:03 was also corroborated by KIR-Fc staining (Fig. 4B). Mamu-B*045:03

has not been previously identified as a KIR ligand, and in contrast to Mamu-AG2*01:01, has a

Bw6 motif (NLRTLRG) at α1-domain residues 77-83 (Supplemental Fig. 3). To better

understand this unusual pattern of responses, we compared the predicted amino acid sequences

for Mamu-B*045:03 and -AG2*01:01. Although Mamu-B*045:03 looks like a typical Mamu-B

molecule in its α2-cytoplasmic domains, we noted a patch of sequence similarity with other

Mamu-A molecules in residues 13-44 of the α1-domain (Fig. 5A). This included a region of

identity at positions 16-19 previously defined as a KIR contact site based on the crystal structure

of HLA-B*57 in complex with human KIR3DL1*001 (80). These observations were supported

by phylogenetic comparisons. Whereas the full-length amino acid sequence of Mamu-B*045:03

clusters with other Mamu-B molecules (Fig. 5B), the α1-domain of Mamu-B*045:03 is more

similar to Mamu-A and -AG molecules (Fig. 5C). A similar pattern of clustering is exhibited by

Mamu-B*036, which is hybrid molecule with MHC-A extracellular domains and an MHC-B

cytoplasmic tail (Fig. 5C). The recognition of Mamu-B*045:03 by KIR3DL07*004,

KIR3DS02*004 and KIR3DSw09*003 may therefore be explained by the origins of this

molecule as the product of an ancestral recombination between *MHC-A* and *-B* genes.

Phylogenetic analysis of the D0-D2 of rhesus macaque KIRs reflects ligand recognition

To investigate the relationship between the extracellular domains of rhesus macaque KIRs and

their patterns of ligand recognition, we conducted phylogenic analysis of the D0-D2 domains of

rhesus KIRs using the human KIR3DL1 and KIR3DL2 as an outgroup. Seven of the eight KIRs

that interact with Mamu-Bw4 ligands cluster together on one branch and five of the six KIRs that

interact with Mamu-AG and -A-related ligands cluster on separate branch (Fig. 6). Furthermore,

within the broader group of receptors that respond to Mamu-AG, the three KIRs that recognize

Mamu-B*045:03 all cluster together. However, KIR3DSw07 was an exception. Although

KIR3DSw07 recognizes Mamu-AG, its D0-D2 domain sequences group with the Bw4-specific

KIRs (Fig. 6). This analysis also revealed that three of the four KIRs for which we were unable

to define a ligand (KIR3DL02*004, KIR3DLw34*002 and KIR3DS05*002) form a distinct

phylogenic group (Fig. 6). A common feature of these KIRs is the conspicuous absence of D0

domain residues 34-36 (Fig. 1A), which could explain the inability to detect interactions with

any of the MHC class I molecules in our panel. In the case of KIR3DL01*012 and

KIR3DL05*008, while these receptors interact respectively with Mamu-Bw4 and -A-related

ligands, they exhibit an earlier branching pattern distinct from other macaque KIRs (Fig. 6). It is

therefore conceivable that *KIR3DL01* and *KIR3DL05* represent more ancestral lineage II genes in

macaques that gave rise to the duplicated genes encoding diverse receptors for Mamu-Bw4 and -

A-related ligands.

Discussion

316
Definition of the MHC class I ligands for rhesus macaques KIRs is fundamental to NK cell

biology in this species as a nonhuman model for infectious diseases, transplantation and

reproductive biology. As consequence of the rapid pace of KIR evolution in primates, it is not

possible to predict the ligands for macaque KIRs on the basis of sequence similarity with their

human counterparts. The *KIR* genes of macaques are evolving at a particularly rapid pace

characterized by extensive gene duplication and recombination that has generated allelic

variation and haplotype diversity that exceeds the diversity of human *KIRs* (55, 56, 81).

Although the number of alleles and predicted *KIR* genes varies among geographically distinct

populations of macaques, extensive sequencing and segregation analyses support the existence of

at least 22 distinct *KIR* genes in rhesus macaques of Indian origin (55, 56, 81). Of these, 19 are

predicted to encode lineage II KIRs for Mamu-A and -B ligands. To provide a foundation for

studying NK cell responses in rhesus macaques and for basic comparative immunogenetics in

primates, we selected common allotypes representing 14 lineage II KIRs for analysis of their

MHC class I interactions. By screening 721.221 cells expressing MHC class I molecules

corresponding to 37 *Mamu-A, -B, -E, -F, -I* and *-AG* alleles in assays with KIR-CD3ζ JNL

reporter cell lines and by staining with KIR-Fc fusion proteins, we identified ligands for 10

previously uncharacterized KIRs.

These analyses revealed two general patterns of ligand recognition; interactions with

Mamu-B molecules that contain a Bw4 motif at residues 77-83 and interactions with one or more

Mamu-A-related molecules. Of the 14 KIRs tested in this study, four (KIR3DL04*001,

KIR3DL10*002, KIR3DL11*001 and KIR3DS04*003) were found to interact with multiple

Mamu-Bw4 ligands. Together with our previous studies identifying Mamu-Bw4 ligands for

KIR3DL01*012, KIR3DL06*001, KIR3DL08*002 and KIR3DSw39*002 (73, 75), these

findings reveal a remarkable expansion of Bw4 specificity in the rhesus macaque. In contrast to

humans, which only have a single *KIR3DL1/S1* gene that encodes receptors for HLA-Bw4,

rhesus macaques have at least eight lineage II *KIRs* that encode receptors for Mamu-Bw4 (Table

I). KIRs that interact with Mamu-A- or -A-like ligands have also been greatly expanded in rhesus

macaques. Six of the KIRs in this study (KIR3DL07*004, KIR3DS01*003, KIR3DS02*004,

KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003) were found to interact with Mamu-

AG2*01:01 alone or in combination with Mamu-B*45:03, a hybrid molecule that contains

MHC-A α1-domain sequences. Together with earlier studies defining similar ligands for

KIR3DL05 (72, 74, 82), at least seven rhesus macaque *KIRs* are known to encode receptors for

Mamu-A or -A-related ligands (Table I).

All of the KIRs that interact with Bw4 ligands recognize Mamu-B*041:01 and -B*065:01

with varying efficiency, suggesting that these molecules may provide a common reference for

this specificity (Table I). However, interactions with other Mamu-Bw4 molecules were more

variable. Several KIRs, including KIR3DL11*001 and KIR3DS04*003, lack detectable

responses to Mamu-B*007:01, -B*022:01 or -B*043:01 (Table I). Differences in the magnitude

of KIR-CD3ζ JNL cell responses and KIR-Fc staining reveal additional variation in the strength

of these interactions. Although most of the KIRs responding to Mamu-Bw4 ligands are

inhibitory, interactions with a couple of activating KIRs (KIR3DS04*003 and KIR3DSw39*002)

were identified (Table I). Thus, differences in the preferential recognition of individual Mamu-

Bw4 molecules, variation in the strength of these interactions, and the identification of inhibitory

and activating KIRs that respond to this group of molecules, suggests extensive functional

diversification of this group of receptors.

Rhesus macaque MHC class I haplotypes typically express two or three *Mamu-A* genes

and four to eleven *Mamu-B* genes that can be classified as major or minor loci on the basis of

their transcriptional abundance (30-32). The Mamu-Bw4 molecules identified as KIR ligands are

products of transcriptionally abundant major genes (32). However, phylogenetic and segregation

analyses suggest that they are distinct from the *Mamu-B* genes that encode molecules known to

present viral peptides to $CD8^+$ T cells (30-33, 83). This is further supported by unique

polymorphisms in the cytoplasmic domains of Mamu-B*007:01, -B*022:01, -B*041:01, -

B*043:01 and -B*065:01 that are not present in the cytoplasmic tails of molecules such as

Mamu-B*008 and -B*017 that present SIV epitopes for recognition by CD8+ T cells (84).

We previously demonstrated that these cytoplasmic domain polymorphisms prevent

MHC class I downmodulation by SIV Nef similar to polymorphisms in the cytoplasmic tail of

HLA-C that prevent downmodulation by HIV-1 Nef (6, 7, 84, 85). Selective downmodulation of

HLA-A and -B (but not HLA-C) by HIV-1 Nef reduces the susceptibility of virus-infected cells

to CD8+ T cells (7, 85). While the Vpu proteins of certain HIV-1 isolates may downmodulate

HLA-C at later stages of infection (86, 87), HLA-C is initially retained on the cell surface to

inhibit NK cells (6). In a similar manner, SIV Nef downmodulates Mamu-A and -B molecules

known to present viral peptides to CD8+ T cells, but not the products of duplicated *Mamu-B*

genes that serve as ligands for multiple KIRs (75, 84). Hence, parallels in both immunogenetics

and virology suggest functional specialization similar to *HLA-B* and *-C* in humans, where the

products of certain *Mamu-B* genes have a dominant role in restricting CD8+ T cell responses

while others predominantly serve as KIR ligands to regulate NK cells.

Another major group of rhesus macaque KIRs appears to interact exclusively with

Mamu-A or A-related molecules. KIR3DL07*004, KIR3DS01*003, KIR3DS02*004,

KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003 all recognize Mamu-AG2*01:01

(Table I). Additional interactions with Mamu-AG1*05:01 were detectable for KIR3DS02*004,

KIR3DS06*002 and KIR3DSw09*003, and with Mamu-B*045:03 for KIR3DL07*004,

KIR3DS02*004 and KIR3DSw09*003 (Table I). Although Mamu-B*045:03 is a Mamu-B

molecule, its α1-domain appears to be derived from an *MHC-A* gene (Fig. 5). Thus, the hybrid

origins of this molecule as a result of ancestral recombination between *MHC-A* and *-B* genes

probably accounts for its interactions with these KIRs. Mamu-B*045:03 recognition may in turn

be dependent on sequences shared by these KIRs, since KIR3DL07*004, KIR3DS02*004 and

KIR3DSw09*003 have nearly identical D0 domains (Fig. 1). As with Mamu-Bw4 interactions,

differences in the preferential recognition of individual Mamu-A-related ligands, the strength of

these interactions, and the identification of both inhibitory and activating receptors that recognize

these molecules indicates extensive functional diversification of this group of KIRs.

In contrast to the KIRs found to interact with Mamu-Bw4 ligands, the majority of the

receptors for Mamu-A and -A-like ligands are activating. Five of the seven rhesus macaque KIRs

that interact with Mamu-AG2*01:01 are products of *KIR3DS* genes (Table I). In humans, the

ligation of activating lineage III KIRs (KIR2DS) and KIR2DL4 on maternal NK cells of the

placental decidua by HLA-C and soluble HLA-G stimulates the release of pro-inflammatory and

pro-angiogenic factors that promote placental vascularization to increase blood supply to the

developing fetus (20, 21, 45). Since macaques do not express orthologs of HLA-C or -G (28, 29,

38), it is tempting to speculate that Mamu-AG interactions with activating lineage II KIRs may

play a similar role in placental development. In contrast to human *KIR* haplotypes, which can be

broadly categorized into groups A and B that correspond to inhibitory and activating profiles

based largely on fixed differences in the content of activating genes, *KIR* haplotypes in macaques

are much more variable (81). *KIR* haplotypes in rhesus macaques usually contain 4-17 expressed

genes, and although the number of activating genes varies, most include at least one activating

KIR (56, 81). Thus, multiple *KIR3DS* genes that encode receptors for Mamu-AG increases the

likelihood that a given animal will express at least one activating KIR capable of engaging this

molecule in the placenta. It is therefore conceivable that in the absence of functional orthologs of

HLA-C or -G, macaques may depend on KIR3DS interactions with Mamu-AG to promote

placental vascularization.

The present study more than doubles the number of rhesus macaque KIRs with defined

MHC class I ligands. At least one MHC class I ligand has now been identified for KIRs

representing 15 different lineage II *KIR* genes in animals of Indian origin (Table I). These

findings reveal a remarkable expansion of two groups of lineage II KIRs with broad specificity

for either Mamu-Bw4 or -A-related molecules. The recognition of overlapping, but non-

redundant, sets of ligands by both inhibiting and activating KIRs indicates extensive

diversification of these receptors in concert with products an expanded complement of *Mamu-A*

and *-B* genes. These observations provide a valuable foundation for studying NK cell responses

in the rhesus macaque as a pre-clinical model for infectious diseases, transplantation and

reproductive biology in addition to advancing our basic understanding of KIR and MHC class I

co-evolution in primates.

References

1. Parham, P. 2005. MHC class I molecules and KIRs in human history, health and survival. *Nat. Rev. Immunol.* 5: 201-214.

2. Lanier, L. L. 2005. NK cell recognition. *Annu. Rev. Immunol.* 23: 225-274.

3. Hasan, M. Z., and L. Walter. 2021. Rhesus Macaque Activating Killer Immunoglobulin-Like Receptors Associate With Fc Receptor Gamma (FCER1G) and Not With DAP12 Adaptor Proteins Resulting in Stabilized Expression and Enabling Signal Transduction. *Front Immunol* 12: 678964.

4. Huard, B., and K. Fruh. 2000. A role for MHC class I down-regulation in NK cell lysis of herpes virus-infected cells. *Eur. J. Immunol.* 30: 509-515.

5. Falk, C. S., M. Mach, D. J. Schendel, E. H. Weiss, I. Hilgert, and G. Hahn. 2002. NK cell activity during human cytomegalovirus infection is dominated by US2-11-mediated HLA class I down-regulation. *J. Immunol.* 169: 3257-3266.

6. Cohen, G. B., R. T. Gandhi, D. M. Davis, O. Mandelboim, B. K. Chen, J. L. Strominger, and D. Baltimore. 1999. The selective downregulation of class I major histocompatibility complex proteins by HIV-1 protects HIV-infected cells from NK cells. *Immunity* 10: 661-671.

7. Schwartz, O., V. Marechal, S. LeGall, F. Lemonnier, and J.-M. Heard. 1996. Endocytosis of major histocompatibility complex class I molecules is induced by the HIV-1 Nef protein. *Nat. Med.* 2: 338-342.

8. Porgador, A., O. Mandelboim, N. P. Restifo, and J. L. Strominger. 1997. Natural killer cell lines kill autologous beta2-microglobulin-deficient melanoma cells: implications for cancer immunotherapy. *Proc. Natl. Acad. Sci. U. S. A.* 94: 13140-13145.

449   9.   Alter, G., M. P. Martin, N. Teigen, W. H. Carr, T. J. Suscovich, A. Schneidewind, H.

450   Streeck, M. Waring, A. Meier, C. Brander, J. D. Lifson, T. M. Allen, M. Carrington, and

451   M. Altfeld. 2007. Differential natural killer cell-mediated inhibition of HIV-1 replication

452   based on distinct KIR/HLA subtypes. *J. Exp. Med.* 204: 3027-3036.

453   10.   O'Connor, G. M., J. P. Vivian, E. Gostick, P. Pymm, B. A. Lafont, D. A. Price, J.

454   Rossjohn, A. G. Brooks, and D. W. McVicar. 2015. Peptide-dependent recognition of

455   HLA-B*57:01 by KIR3DS1. *J. Virol.* 89: 5213-5221.

456   11.   Garcia-Beltran, W. F., A. Holzemer, G. Martrus, A. W. Chung, Y. Pacheco, C. R.

457   Simoneau, M. Rucevic, P. A. Lamothe-Molina, T. Pertel, T. E. Kim, H. Dugan, G. Alter,

458   J. Dechanet-Merville, S. Jost, M. Carrington, and M. Altfeld. 2016. Open conformers of

459   HLA-F are high-affinity ligands of the activating NK-cell receptor KIR3DS1. *Nat.*

460   *Immunol.* 17: 1067-1074.

461   12.   Martin, M. P., X. Gao, J.-H. Lee, G. W. Nelson, R. Detels, J. J. Goedert, S. Buchbinder,

462   K. Hoots, D. Vlahov, J. Trowsdale, M. Wilson, S. J. O'Brien, and M. Carrington. 2002.

463   Epistatic interaction between KIR3DS1 and HLA-B delays the progression to AIDS. *Nat.*

464   *Genet.* 31: 429-434.

465   13.   Martin, M. P., Y. Qi, X. Gao, E. Yamada, J. N. Martin, F. Pereyra, S. Colombo, E. E.

466   Brown, W. L. Shupert, J. Phair, J. J. Goedert, S. Buchbinder, G. D. Kirk, A. Telenti, M.

467   Connors, S. J. O'Brien, B. D. Walker, P. Parham, S. G. Deeks, D. W. McVicar, and M.

468   Carrington. 2007. Innate partnership of HLA-B and KIR3DL1 subtypes against HIV-1.

469   *Nat. Genet.* 39: 733-740.

470   14.   Khakoo, S. I., C. L. Thio, M. P. Martin, C. R. Brooks, X. Gao, J. Astemborski, J. Cheng,

471   J. J. Godert, D. Vlahov, M. Hilgartner, S. Cox, A.-M. Little, G. J. Alexander, M. E.

Cramp, S. J. O'Brien, W. M. C. Rosenberg, D. L. Thomas, and M. Carrington. 2004. HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. *Science* 305: 872-874.

15. Carrington, M., S. Wang, M. P. Martin, X. Gao, M. Schiffman, J. Cheng, R. Herrero, A. C. Rodriguez, R. Kurman, R. Mortel, P. Schwartz, A. Glass, and A. Hildesheim. 2005. Hierarchy of resistance to cervical neoplasia mediated by combinations of killer immunoglobulin-like receptor and human leukocyte antigen loci. *J. Exp. Med.* 201: 1069-1075.

16. Chen, C., M. Busson, V. Rocha, M.-L. Appert, V. Lepage, N. Dulphy, P. Haas, G. Socie, A. Toubert, D. Charron, and P. Loiseau. 2006. Activating KIR genes are associated with CMV reactivation and survival after non-T-cell depleted HLA-identical sibling bone marrow transplantation for malignant disorders. *Bone Marrow Transplant.* 38: 437-444.

17. Cooley, S., E. Trachtenberg, T. L. Bergemann, K. Saeteurn, J. Klein, C. T. Le, S. G. Marsh, L. A. Guethlein, P. Parham, J. S. Miller, and D. J. Weisdorf. 2009. Donors with group B KIR haplotypes improve relapse-free survival after unrelated hematopoietic cell transplantation for acute myelogenous leukemia. *Blood* 113: 726-732.

18. Cooley, S., D. J. Weisdorf, L. A. Guethlein, J. P. Klein, T. Wang, C. T. Le, S. G. Marsh, D. Geraghty, S. Spellman, M. D. Haagenson, M. Ladner, E. Trachtenberg, P. Parham, and J. S. Miller. 2010. Donor selection for natural killer cell receptor genes leads to superior survival after unrelated transplantation for acute myelogenous leukemia. *Blood* 116: 2411-2419.

19. Delgado, D. C., J. A. Hank, J. Kolesar, D. Lorentzen, J. Gan, S. Seo, K. Kim, S. Shusterman, S. D. Gillies, R. A. Reisfeld, R. Yang, B. Gadbaw, K. B. DeSantes, W. B.

London, R. C. Seeger, J. M. Maris, and P. M. Sondel. 2010. Genotypes of NK cell KIR

receptors, their ligands, and Fcgamma receptors in the response of neuroblastoma

patients to Hu14.18-IL2 immunotherapy. *Cancer Res.* 70: 9554-9561.

20. Hiby, S. E., J. J. Walker, M. O'Shaughnessy K, C. W. Redman, M. Carrington, J.

Trowsdale, and A. Moffett. 2004. Combinations of maternal KIR and fetal HLA-C genes

influence the risk of preeclampsia and reproductive success. *J. Exp. Med.* 200: 957-965.

21. Hiby, S. E., R. Apps, A. M. Sharkey, L. E. Farrell, L. Gardner, A. Mulder, F. H. Claas, J.

J. Walker, C. W. Redman, L. Morgan, C. Tower, L. Regan, G. E. Moore, M. Carrington,

and A. Moffett. 2010. Maternal activating KIRs protect against human reproductive

failure mediated by fetal HLA-C2. *J. Clin. Invest.* 120: 4102-4110.

22. Orrantia, A., I. Terren, G. Astarloa-Pando, C. Gonzalez, A. Uranga, J. J. Mateos-Mazon,

J. C. Garcia-Ruiz, M. Rinon, M. Rey, S. Perez-Fernandez, O. Zenarruzabeitia, and F.

Borrego. 2021. NK Cell Reconstitution After Autologous Hematopoietic Stem Cell

Transplantation: Association Between NK Cell Maturation Stage and Outcome in

Multiple Myeloma. *Frontiers in immunology* 12: 748207.

23. Littera, R., G. Piredda, D. Argiolas, S. Lai, E. Congeddu, P. Ragatzu, M. Melis, E. Carta,

M. B. Michittu, D. Valentini, L. Cappai, R. Porcella, F. Alba, M. Serra, V. Loi, R. Maddi,

S. Orru, G. La Nasa, G. Caocci, R. Cusano, M. Arras, M. Frongia, A. Pani, and C.

Carcassi. 2017. KIR and their HLA Class I ligands: Two more pieces towards completing

the puzzle of chronic rejection and graft loss in kidney transplantation. *PLoS One* 12:

e0180831.

24.   Yokoyama, W. M., P. J. Kehn, D. I. Cohen, and E. M. Shevach. 1990. Chromosomal

517   location of the Ly-49 (A1, YE1/48) multigene family, genetic association with the NK1.1

518   antigen. *J. Immunol.* 145: 2353-2358.

519   25.   Khakoo, S. I., R. Rajalingam, B. P. Shum, K. Weidenbach, L. Flodin, D. G. Muir, F.

520   Canavez, S. L. Cooper, N. M. Valiante, L. L. Lanier, and P. Parham. 2000. Rapid

521   evolution of NK cell receptor systems demonstrated by comparison of chimpanzee and

522   humans. *Immunity* 12: 687-698.

523   26.   Sambrook, J. G., A. Bashirova, S. Palmer, S. Sims, J. Trowsdale, L. Abi-Rached, P.

524   Parham, M. Carrington, and S. Beck. 2005. Single haplotype analysis demonstrates rapid

525   evolution of the killer immunoglobulin-like receptor (KIR) loci in primates. *Genome Res.*

526   15: 25-35.

527   27.   Parham, P., L. Abi-Rached, L. Matevosyan, A. K. Moesta, P. J. Norman, A. M. O.

528   Aguilar, and L. A. Guethlein. 2010. Primate-specific regulation of natural killer cells. *J.*

529   *Med. Primatol.* 39: 194-212.

530   28.   Boyson, J. E., C. Schufflebotham, L. F. Cadavid, J. A. Urvater, L. A. Knapp, A. L.

531   Huges, and D. I. Watkins. 1996. The MHC class I genes of the rhesus monkey: Different

532   evolutionary histories of the MHC class I and II genes in primates. *J. Immunol.* 156:

533   4656-4665.

534   29.   Adams, E. J., and P. Parham. 2001. Species-specific evolution of MHC class I genes in

535   the higher primates. *Immunol. Rev.* 183: 41-64.

536   30.   Otting, N., C. M. C. Heijmans, R. C. Noort, N. G. d. Groot, G. G. M. Doxiadis, J. J.

537   vanRood, D. I. Watkins, and R. E. Bontrop. 2005. Unparalleled complexity of the MHC

538   class I region in rhesus macaques. *Proc. Natl. Acad. Sci. U. S. A.* 102: 1626-1631.

31.   Daza-Vamenta, R., G. Glusman, L. Rowen, B. Guthrie, and D. E. Geraghty. 2004.

Genetic divergence of the rhesus macaque major histocompatibility complex. *Genome*

Res. 14: 1501-1515.

32.   Wiseman, R. W., J. A. Karl, P. S. Bohn, F. A. Nimityongskul, G. J. Starrett, and D. H.

O'Connor. 2013. Haplessly hoping: macaque major histocompatibility complex made

easy. *ILAR journal / National Research Council, Institute of Laboratory Animal*

Resources 54: 196-210.

33.   Evans, D. T., P. Jing, T. M. Allen, D. H. O'Connor, H. Horton, J. E. Venham, M.

Piekarczyk, J. Dzuris, M. Dykhuzen, J. Mitchen, R. A. Rudersdorf, C. D. Pauza, A. Sette,

R. E. Bontrop, R. DeMars, and D. I. Watkins. 2000. Definition of five new simian

immunodeficiency virus cytotoxic T-lymphocyte epitopes and their restricting major

histocompatibility complex class I molecules: evidence for an influence on disease

progression. *J. Virol.* 74: 7400-7410.

34.   Boyson, J. E., S. N. McAdam, A. Gallimore, T. G. Golos, X. Liu, F. M. Gotch, A. L.

Hughes, and D. I. Watkins. 1995. The MHC E locus in macaques is polymorphic and is

conserved between macaques and humans. *Immunogenetics* 41: 59-68.

35.   Knapp, L., L. F. Cadavid, and D. I. Watkins. 1998. The MHC-E locus is the most well

conserved of all known primate class I histocompatibility genes. *J. Immunol.* 160: 189-

196.

36.   Godfrey, D. I., J. Le Nours, D. M. Andrews, A. P. Uldrich, and J. Rossjohn. 2018.

Unconventional T cell targets for cancer immunotherapy. *Immunity* 48: 453-473.

37. Walter, L., and B. Petersen. 2017. Diversification of both KIR and NKG2 natural killer

cell receptor genes in macaques - implications for highly complex MHC-dependent

regulation of natural killer cells. *Immunology* 150: 139-145.

38. Boyson, J. E., K. K. Iwanaga, T. G. Golos, and D. I. Watkins. 1996. Identifiction of the

rhesus monkey HLA-G ortholog. Mamu-G is a pseudogene. *J. Immunol.* 157: 5428-5437.

39. Castro, M. J., P. Morales, V. Fernandez-Soria, B. Suarez, M. J. Recio, M. Alvarez, M.

Martin-Villa, and A. Arnaiz-Villena. 1996. Allelic diversity at the primate Mhc-G locus:

exon 3 bears stop codons in all Cercopithecinae sequences. *Immunogenetics* 43: 327-336.

40. Boyson, J. E., K. K. Iwanaga, T. G. Golos, and D. I. Watkins. 1997. Identification of a

novel MHC class I gene, Mamu-AG, expressed in the placenta of a primate with an

inactivated G locus. *J. Immunol.* 159: 3311-3321.

41. Boyson, J. E., K. K. Iwanaga, J. A. Urvater, A. L. Hughes, T. G. Golos, and D. I.

Watkins. 1999. Evolution of a new nonclassical MHC class I locus in two Old World

primate species. *Immunogenetics* 49: 86-98.

42. Langat, D. K., P. J. Morales, A. T. Fazleabas, J. M. Mwenda, and J. S. Hunt. 2002.

Baboon placentas express soluble and membrane-bound Paan-AG proteins encoded by

alternatively spliced transcripts of the class Ib major histocompatibility complex gene,

Paan-AG. *Immunogenetics* 54: 164-173.

43. Bondarenko, G. I., S. V. Dambaeva, R. L. Grendell, A. L. Hughes, M. Durning, M. A.

Garthwaite, and T. G. Golos. 2009. Characterization of cynomolgus and vervet monkey

placental MHC class I expression: diversity of the nonhuman primate AG locus.

Immunogenetics 61: 431-442.

44. Slukvin, II, D. P. Lunn, D. I. Watkins, and T. G. Golos. 2000. Placental expression of the

nonclassical MHC class I molecule Mamu-AG at implantation in the rhesus monkey.

Proc. Natl. Acad. Sci. U. S. A. 97: 9104-9109.

45. Rajagopalan, S. 2014. HLA-G-mediated NK cell senescence promotes vascular

remodeling: implications for reproduction. *Cell Mol Immunol* 11: 460-466.

46. Arnaiz-Villena, A., I. Juarez, F. Suarez-Trujillo, A. Lopez-Nares, C. Vaquero, J. Palacio-

Gruber, and J. M. Martin-Villa. 2020. HLA-G: Function, polymorphisms and pathology.

Int J Immunogenet.

47. Urvater, J. A., N. Otting, J. H. Loehrke, R. Rudersdorf, I. I. Slukvin, M. S. Piekarczyk, T.

G. Golos, A. L. Hughes, R. E. Bonrop, and D. I. Watkins. 2000. Mamu-I: A novel

primate MHC class I B-related locus with unusually low variability. *J. Immunol.* 164:

1386-1398.

48. Hershberger, K. L., R. Shyam, A. Miura, and N. L. Letvin. 2001. Diversity of the killer

cell Ig-like receptors of rhesus monkeys. *J. Immunol.* 166: 4380-4390.

49. Bimber, B. N., A. J. Moreland, R. W. Wiseman, A. L. Hughes, and D. H. O'Connor.

2008. Complete characterization of killer Ig-like receptor (KIR) haplotypes in Mauritian

cynomolgus macaques: Novel insights into nonhuman primate KIR gene content and

organization. *J. Immunol.* 181: 6301-6308.

50. Kruse, P. H., C. Rosner, and L. Walter. 2010. Characterization of rhesus macaque KIR

genotypes and haplotypes. *Immunogenetics* 62: 281-293.

51. Blokhuis, J. H., M. K. v. d. Wiel, G. G. Doxiadis, and R. E. Bontrop. 2010. The mosaic

of KIR haplotypes in rhesus macaques. *Immunogenetics* 62: 295-306.

52.   Bimber, B. N., and D. T. Evans. 2015. The killer-cell immunoglobulin-like receptors of

605   macaques. *Immunol. Rev.* 267: 246-258.

606   53.   Blokhuis, J. H., M. K. van der Wiel, G. G. Doxiadis, and R. E. Bontrop. 2011. The

607   extreme plasticity of killer cell Ig-like receptor (KIR) haplotypes differentiates rhesus

608   macaques from humans. *Eur. J. Immunol.* 41: 2719-2728.

609   54.   Moreland, A. J., L. A. Guethlein, R. K. Reeves, K. W. Broman, R. P. Johnson, P.

610   Parham, D. H. O'Connor, and B. N. Bimber. 2011. Characterization of killer

611   immunoglobulin-like receptor genetics and comprehensive genotyping by

612   pyrosequencing in rhesus macaques. *BMC genomics* 12: 295.

613   55.   Bruijnesteijn, J., M. K. H. van der Wiel, W. T. N. Swelsen, N. Otting, A. J. M. de Vos-

614   Rouweler, D. Elferink, G. G. Doxiadis, F. H. J. Claas, N. M. Lardy, N. G. de Groot, and

615   R. E. Bontrop. 2018. Human and rhesus macaque KIR haplotypes defined by their

616   transcriptomes. *J. Immunol.* 200: 1692-1701.

617   56.   Bruijnesteijn, J., N. de Groot, M. K. H. van der Wiel, N. Otting, A. J. M. de Vos-

618   Rouweler, N. G. de Groot, and R. E. Bontrop. 2020. Unparalleled Rapid Evolution of

619   KIR Genes in Rhesus and Cynomolgus Macaque Populations. *J. Immunol.* 204: 1770-

620   1786.

621   57.   Hatziioannou, T., and D. T. Evans. 2012. Animal models for HIV/AIDS research. *Nature*

622   *Reviews Microbiology* 10: 852-867.

623   58.   Barouch, D. H., J. B. Whitney, B. Moldt, F. Klein, T. Y. Oliveira, J. Liu, K. E.

624   Stephenson, H. W. Chang, K. Shekhar, S. Gupta, J. P. Nkolola, M. S. Seaman, K. M.

625   Smith, E. N. Borducchi, C. Cabral, J. Y. Smith, S. Blackmore, S. Sanisetty, J. R. Perry,

626   M. Beck, M. G. Lewis, W. Rinaldi, A. K. Chakraborty, P. Poignard, M. C. Nussenzweig,

627   and D. R. Burton. 2013. Therapeutic efficacy of potent neutralizing HIV-1-specific

628   monoclonal antibodies in SHIV-infected rhesus monkeys. *Nature* 503: 224-228.

629 59. Esquivel, R. N., A. Patel, S. B. Kudchodkar, D. H. Park, K. Stettler, M. Beltramello, J.

630   W. Allen, J. Mendoza, S. Ramos, H. Choi, P. Borole, K. Asija, M. Bah, S. Shaheen, J.

631   Chen, J. Yan, A. C. Durham, T. R. F. Smith, K. Broderick, G. Guibinga, K. Muthumani,

632   D. Corti, L. Humeau, and D. B. Weiner. 2019. In Vivo Delivery of a DNA-Encoded

633   Monoclonal Antibody Protects Non-human Primates against Zika Virus. *Mol Ther* 27:

634   974-985.

635 60. Hessell, A. J., L. Hangartner, M. Hunter, C. E. Havenith, F. J. Beurskens, J. M. Bakker,

636   C. M. Lanigan, G. Landucci, D. N. Forthal, P. W. Parren, P. A. Marx, and D. R. Burton.

637   2007. Fc receptor but not complement binding is important in antibody protection against

638   HIV. *Nature* 449: 101-104.

639 61. Hessell, A. J., P. Poignard, M. Hunter, L. Hangartner, D. M. Tehrani, W. K. Bleeker, P.

640   W. Parren, P. A. Marx, and D. R. Burton. 2009. Effective, low-titer antibody protection

641   against low-dose repeated mucosal SHIV challenge in macaques. *Nat Med* 15: 951-954.

642 62. Magnani, D. M., T. F. Rogers, N. J. Maness, N. D. Grubaugh, N. Beutler, V. K. Bailey,

643   L. Gonzalez-Nieto, M. J. Gutman, N. Pedreño-Lopez, J. M. Kwal, M. J. Ricciardi, T. A.

644   Myers, J. G. Julander, R. P. Bohm, M. H. Gilbert, F. Schiro, P. P. Aye, R. V. Blair, M. A.

645   Martins, K. P. Falkenstein, A. Kaur, C. L. Curry, E. G. Kallas, R. C. Desrosiers, P. J.

646   Goldschmidt-Clermont, S. S. Whitehead, K. G. Andersen, M. C. Bonaldo, A. A. Lackner,

647   A. T. Panganiban, D. R. Burton, and D. I. Watkins. 2018. Fetal demise and failed

648   antibody therapy during Zika virus infection of pregnant macaques. *Nat Commun* 9:

649   1624.

63.   Martinez-Navio, J. M., S. P. Fuchs, S. N. Pantry, W. A. Lauer, N. N. Duggan, B. F.

Keele, E. G. Rakasz, G. Gao, J. D. Lifson, and R. C. Desrosiers. 2019. Adeno-Associated

Virus Delivery of Anti-HIV Monoclonal Antibodies Can Drive Long-Term Virologic

Suppression. *Immunity* 50: 567-575.e565.

64.   Pauthner, M. G., J. P. Nkolola, C. Havenar-Daughton, B. Murrell, S. M. Reiss, R.

Bastidas, J. Prévost, R. Nedellec, B. von Bredow, P. Abbink, C. A. Cottrell, D. W. Kulp,

T. Tokatlian, B. Nogal, M. Bianchi, H. Li, J. H. Lee, S. T. Butera, D. T. Evans, L.

Hangartner, A. Finzi, I. A. Wilson, R. T. Wyatt, D. J. Irvine, W. R. Schief, A. B. Ward,

R. W. Sanders, S. Crotty, G. M. Shaw, D. H. Barouch, and D. R. Burton. 2019. Vaccine-

Induced Protection from Homologous Tier 2 SHIV Challenge in Nonhuman Primates

Depends on Serum-Neutralizing Antibody Titers. *Immunity* 50: 241-252.e246.

65.   Shingai, M., O. K. Donau, R. J. Plishka, A. Buckler-White, J. R. Mascola, G. J. Nabel,

M. C. Nason, D. Montefiori, B. Moldt, P. Poignard, R. Diskin, P. J. Bjorkman, M. A.

Eckhaus, F. Klein, H. Mouquet, J. C. Cetrulo Lorenzi, A. Gazumyan, D. R. Burton, M. C.

Nussenzweig, M. A. Martin, and Y. Nishimura. 2014. Passive transfer of modest titers of

potent and broadly neutralizing anti-HIV monoclonal antibodies block SHIV infection in

macaques. *J. Exp. Med.* 211: 2061-2074.

66.   Welles, H. C., M. F. Jennewein, R. D. Mason, S. Narpala, L. Wang, C. Cheng, Y. Zhang,

J. P. Todd, J. D. Lifson, A. B. Balazs, G. Alter, A. B. McDermott, J. R. Mascola, and M.

Roederer. 2018. Vectored delivery of anti-SIV envelope targeting mAb via AAV8

protects rhesus macaques from repeated limiting dose intrarectal swarm SIVsmE660

challenge. *PLoS Pathog.* 14: e1007395.

67. Lu, S., Y. Zhao, W. Yu, Y. Yang, J. Gao, J. Wang, D. Kuang, M. Yang, J. Yang, C. Ma,

J. Xu, X. Qian, H. Li, S. Zhao, J. Li, H. Wang, H. Long, J. Zhou, F. Luo, K. Ding, D.

Wu, Y. Zhang, Y. Dong, Y. Liu, Y. Zheng, X. Lin, L. Jiao, H. Zheng, Q. Dai, Q. Sun, Y.

Hu, C. Ke, H. Liu, and X. Peng. 2020. Comparison of SARS-CoV-2 infections among 3

species of non-human primates. *bioRxiv*: 2020.2004.2008.031807.

68. Roark, H. K., J. A. Jenks, S. R. Permar, and M. R. Schleiss. 2020. Animal Models of

Congenital Cytomegalovirus Transmission: Implications for Vaccine Development. *J.*

Infect. Dis. 221: S60-S73.

69. Bondarenko, G. I., D. W. Burleigh, M. Durning, E. E. Breburda, R. L. Grendell, and T.

G. Golos. 2007. Passive immunization against the MHC class I molecule Mamu-AG

disrupts rhesus placental development and endometrial responses. *J. Immunol.* 179: 8042-

8050.

70. Torrealba, J. R., M. Katayama, J. H. Fechner, Jr., E. Jankowska-Gan, S. Kusaka, Q. Xu,

J. M. Schultz, T. D. Oberley, H. Hu, M. M. Hamawy, M. Jonker, J. Wubben, G.

Doxiadis, R. Bontrop, W. J. Burlingham, and S. J. Knechtle. 2004. Metastable tolerance

to rhesus monkey renal transplants is correlated with allograft TGF-beta 1+CD4+ T

regulatory cell infiltrates. *J. Immunol.* 172: 5753-5764.

71. Forrest, L., J. Fechner, J. Post, N. Van Asselt, K. Kvasnica, L. D. Haynes, J. Coonen, K.

Brunner, W. J. Haynes, C. Little, W. J. Burlingham, P. Hematti, S. Strober, and D. B.

Kaufman. 2021. Tomotherapy Applied Total Lymphoid Irradiation and Allogeneic

Hematopoietic Cell Transplantation Generates Mixed Chimerism in the Rhesus Macaque

Model. *Radiat. Res.*

72.   Colantonio, A. D., B. N. Bimber, W. J. Neidermyer, R. K. Reeves, G. Alter, M. Altfeld,

R. P. Johnson, M. Carrington, D. H. O'Connor, and D. T. Evans. 2011. KIR

polymorphisms modulate peptide-dependent binding to an MHC class I ligand with a

Bw6 motif. *PLoS Pathog.* 7: e1001316.

73.   Schafer, J. L., A. D. Colantonio, W. J. Neidermyer, D. M. Dudley, M. Connole, D. H.

O'Connor, and D. T. Evans. 2014. KIR3DL01 recognition of Bw4 ligands in the rhesus

macaque: maintenance of Bw4 specificity since the divergence of apes and Old World

monkeys. *J. Immunol.* 192: 1907-1917.

74.   Rosner, C., P. H. Kruse, M. Hermes, N. Otto, and L. Walter. 2011. Rhesus macaque

inhibitory and activating KIR3D interact with Mamu-A-encoded ligands. *J. Immunol.*

186: 2156-2163.

75.   Banerjee, P., M. Ries, S. K. Janaka, A. G. Grandea, 3rd, R. Wiseman, D. H. O'Connor, T.

G. Golos, and D. T. Evans. 2018. Diversification of Bw4 Specificity and Recognition of a

Nonclassical MHC Class I Molecule Implicated in Maternal-Fetal Tolerance by Killer

Cell Ig-like Receptors of the Rhesus Macaque. *J. Immunol.* 201: 2776-2786.

76.   Nakamura, T., K. D. Yamada, K. Tomii, and K. Katoh. 2018. Parallelization of MAFFT

for large-scale multiple sequence alignments. *Bioinformatics* 34: 2490-2492.

77.   Katoh, K., J. Rozewicki, and K. D. Yamada. 2019. MAFFT online service: multiple

sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* 20:

1160-1166.

78.   Robinson, O., D. Dylus, and C. Dessimoz. 2016. Phylo.io: Interactive Viewing and

Comparison of Large Phylogenetic Trees on the Web. *Mol Biol Evol* 33: 2163-2166.

79. Nicholas, R. E., K. Sandstrom, J. L. Anderson, W. R. Smith, M. Wetzel, P. Banerjee, S.

K. Janaka, and D. T. Evans. 2022. KIR3DL05 and KIR3DS02 recognition of a

nonclassical MHC class I molecule in the rhesus macaque implicated in pregnancy

success. *Frontiers in Immunolgy* 13: in press.

80. Vivian, J. P., R. C. Duncan, R. Berry, G. M. O'Connor, H. H. Reid, T. Beddoe, S. Gras,

P. M. Saunders, M. A. Olshina, J. M. Widjaja, C. M. Harpur, J. Lin, S. M. Maloveste, D.

A. Price, B. A. Lafont, D. W. McVicar, C. S. Clements, A. G. Brooks, and J. Rossjohn.

2011. Killer cell immunoglobulin-like receptor 3DL1-mediated recognition of human

leukocyte antigen B. *Nature* 479: 401-405.

81. Bruijnesteijn, J., N. de Groot, A. J. M. de Vos-Rouweler, N. G. de Groot, and R. E.

Bontrop. 2022. Comparative genetics of KIR haplotype diversity in humans and rhesus

macaques: the balancing act. *Immunogenetics*.

82. Nicholas, R. E., K. Sandstrom, J. L. Anderson, W. R. Smith, M. Wetzel, P. Banerjee, S.

K. Janaka, and D. T. Evans. 2022. KIR3DL05 and KIR3DS02 Recognition of a

Nonclassical MHC Class I Molecule in the Rhesus Macaque Implicated in Pregnancy

Success. *Frontiers in immunology* 13: 841136.

83. Budde, M. L., J. J. Lhost, B. J. Burwitz, E. A. Becker, C. M. Burns, S. L. O'Connor, J. A.

Karl, R. W. Wiseman, B. N. Bimber, G. L. Zhang, W. Hildebrand, V. Brusic, and D. H.

O'Connor. 2011. Transcriptionally abundant major histocompatibility complex class I

alleles are fundamental to nonhuman primate simian immunodeficiency virus-specific

CD8+ T cell responses. *J. Virol.* 85: 3250-3261.

84. DeGottardi, M. Q., A. Specht, B. Metcalf, A. Kaur, F. Kirchhoff, and D. T. Evans. 2008.

Selective downregulation of rhesus macaque and sooty mangabey major

histocompatibility complex class I molecules by nef alleles of simian immunodeficiency

virus and human immunodeficiency virus type 2. *J. Virol.* 82: 3139-3146.

85. Collins, K. L., B. K. Chen, S. A. Kalamis, B. D. Walker, and D. Baltimore. 1998. HIV-1

nef protein protects infected cells against killing by cytotoxic T lymphocytes. *Nature*

391: 397-401.

86. Apps, R., G. Q. Del Prete, P. Chatterjee, A. Lara, Z. L. Brumme, M. A. Brockman, S.

Neil, S. Pickering, D. K. Schneider, A. Piechocka-Trocha, B. D. Walker, R. Thomas, G.

M. Shaw, B. H. Hahn, B. F. Keele, J. D. Lifson, and M. Carrington. 2016. HIV-1 Vpu

Mediates HLA-C Downregulation. *Cell Host Microbe* 19: 686-695.

87. Bachtel, N. D., G. Umviligihozo, S. Pickering, T. M. Mota, H. Liang, G. Q. Del Prete, P.

Chatterjee, G. Q. Lee, R. Thomas, M. A. Brockman, S. Neil, M. Carrington, B. Bwana,

D. R. Bangsberg, J. N. Martin, E. G. Kallas, C. S. Donini, N. B. Cerqueira, U. T.

O'Doherty, B. H. Hahn, R. B. Jones, Z. L. Brumme, D. F. Nixon, and R. Apps. 2018.

HLA-C downregulation by HIV-1 adapts to host HLA genotype. *PLoS Pathog.* 14:

e1007257.

88. Bruijnesteijn, J., N. G. de Groot, N. Otting, G. Maccari, L. A. Guethlein, J. Robinson, S.

G. E. Marsh, L. Walter, D. H. O'Connor, J. A. Hammond, P. Parham, and R. E. Bontrop.

2020. Nomenclature report for killer-cell immunoglobulin-like receptors (KIR) in

macaque species: new genes/alleles, renaming recombinant entities and IPD-NHKIR

updates. *Immunogenetics* 72: 37-47.

Footnotes

This work was supported by Public Health Service grants AI095098, AI155163, AI161816, AI121135, AI148379 and AI098485 to DTE. Additional support was provided by PHS grant OD011106 to the WNPRC. DTE is an Elizabeth Glaser Scientist of the Elizabeth Glaser Pediatric AIDS Foundation and The UW Medical Foundation Professor of Pathology and Laboratory Medicine.

Abbreviations used in this article: KIR, killer-cell Ig-like receptor; *Mamu, Macaca mulatta*; JNL, Jurkat NFAT luciferase

771 **Table**

772 Table I. MHC class I ligands of rhesus macaque lineage II KIRs

KIR Gene[a]	Allotype[b]	MHC class I ligands[c]	Group[d]	Refs[e]
KIR3DL01	KIR3DL01*012	Mamu-B*007, -B*022, -B*041, -B*043, -B*065	Bw4	[73]
KIR3DL02	**KIR3DL02*004**	**none detected**		
KIR3DL04	**KIR3DL04*001**	**Mamu-B*007, -B*022, -B*041, -B*043, -B*065**	**Bw4**	
KIR3DL05	KIR3DL05*008	Mamu-A1*002, -A3*13, -B*036, -AG2*01, -AG1*05	A/A-related	[72, 74, 75]
KIR3DL06	KIR3DL06*001	Mamu-B*007, -B*022, -B*041, -B*043, -B*065	Bw4	[75]
KIR3DL07	**KIR3DL07*004**	**Mamu-AG2*01, -B*045**	**A-related**	
KIR3DL08	KIR3DL08*002	Mamu-B*002, -B*007, -B*022, -B*041, -B*065	Bw4	[75]
KIR3DL10	**KIR3DL10*002**	**Mamu-B*007, -B*022, -B*041, -B*043, -B*065**	**Bw4**	
KIR3DL11	**KIR3DL11*001**	**Mamu-B*007, -B*041, -B*065**	**Bw4**	
KIR3DLw34	**KIR3DLw34*002**[f]	**none detected**		
KIR3DS01	**KIR3DS01*003**	**Mamu-AG2*01**	**A-related**	
KIR3DS02	**KIR3DS02*004**	Mamu-AG2*01**, -AG1*05, -B*045**	**A-related**	[82]
KIR3DS03	**KIR3DS03*003**	**none detected**		
KIR3DS04	**KIR3DS04*003**	**Mamu-B*022:01, -B*041, -B*065**	**Bw4**	
KIR3DS05	**KIR3DS05*002**	**none detected**		
KIR3DS06	**KIR3DS06*002**	**Mamu-AG2*01, -AG1*05**	**A-related**	
KIR3DSw07	**KIR3DSw07*001**	**Mamu-AG2*01**	**A-related**	
KIR3DSw09	**KIR3DSw09*003**	**Mamu-AG2*01, -AG1*05, -B*045**	**A-related**	
KIR3DSw39	KIR3DSw39*002[g]	Mamu-B*007, -B*022, -B*041, -B*043, -B*065	Bw4	[75]

[a]Rhesus macaque lineage II KIR genes.

[b]KIR allotypes tested in this study are indicated in bold.

[c]Novel MHC class I ligands defined in this study are indicated in bold.

[d]Group of MHC class I ligands.

[e]References for previously reported KIR ligands.

[f]Previously KIR3DLw03*003 (88).

[g]Previously KIR3DSw08*006 (88).

 Figure Legends

FIGURE 1. Rhesus macaque KIRs and KIR-CD3ζ expression on JNL cells. (**A**) An alignment

showing the predicted amino acid sequences for the D0, D1 and D2 domains of representative

allotypes of rhesus macaque lineage II KIRs. Shaded sequences correspond to predicted MHC

class I contact sites (80). Positions of identity are indicated by periods and amino acid

differences are identified by their single-letter code. (**B**) Parental and KIR-CD3ζ-transduced JNL

cells were stained with a PE-conjugated anti-Flag antibody and Near-IR LIVE/DEAD stain.

After excluding dead cells, the fluorescence intensity of KIR-CD3ζ (Flag-tag) staining on KIR-

CD3ζ-transduced JNL cells (open) was compared to background staining on parental JNL cells

(shaded). Flow cytometry data was analyzed using FlowJo 10.6 for Mac OSX.

FIGURE 2. KIR3D04*001, KIR3DL10*002, KIR3DL11*001 and KIR3DS04*003 interact with

multiple Mamu-Bw4 ligands. (**A**) JNL cells expressing the extracellular domains of

KIR3DL04*001, KIR3DL10*002, KIR3DL11*001 and KIR3DS04*003 fused to the

transmembrane and cytoplasmic domains of human CD3ζ were incubated overnight at a 1:1 E:T

ratio with 721.221 cells expressing the indicated Mamu-B molecules. The bar graphs represent

the mean and standard deviation (error bars) of luciferase activity (RLU) from triplicate wells of

KIR-CD3ζ JNL cells incubated with Mamu-B+721.221 cells (blue), parental 721.221 cells (red)

and anti-Flag plus anti-mouse antibodies (gray). Asterisks denote significant differences by one-

way ANOVA with Dunnett test (**** $p < 0.0001$). These results are representative of at least

three independent experiments. (**B**) Parental and Mamu-B+ 721.221 cells were stained with

Near-IR LIVE/DEAD dye, KIR-Fc fusion proteins consisting of the extracellular domains

KIR3DL04*001, KIR3DL10*002, KIR3DL11*001 and KIR3DS04*003 fused to the Fc domain

of murine IgG2A followed by AF647-conjugated goat anti-mouse IgG and PE-conjugated MHC

class I-specific antibody (W6/32). Data was collected using a FACSymphony™ A3 Cell

Analyzer and analyzed using FlowJo 10.8.0 for Mac OSX.

FIGURE 3. KIR3DL07*004, KIR3DS01*003, KIR3DS02*004, KIR3DS06*002,

KIR3DSw07*001 and KIR3DSw09*003 interact with Mamu-AG ligands. (**A**) JNL cells

expressing the extracellular domains of KIR3DL07*004, KIR3DS01*003, KIR3DS02*004,

KIR3DS06*002, KIR3DSw07*001 and KIR3DSw09*003 fused to the transmembrane and

cytoplasmic domains of human CD3ζ were incubated overnight at a 1:1 E:T ratio with 721.221

cells expressing the indicated Mamu-AG molecules. The bar graphs represent the mean and

standard deviation (error bars) of luciferase activity (RLU) from triplicate wells of KIR-CD3ζ

JNL cells incubated with Mamu-AG+ 721.221 cells (purple), parental 721.221 cells (red) and

anti-Flag plus anti-mouse antibodies (gray). Asterisks denote significant differences by one-way

ANOVA with Dunnett test (**** $p < 0.0001$). These results are representative of at least three

independent experiments. (**B**) Parental and Mamu-AG+721.221 cells were stained with Near-IR

LIVE/DEAD dye, KIR-Fc fusion proteins consisting of the extracellular domains

KIR3DL07*004, KIR3DS01*003, KIR3DS02*004, KIR3DS06*002, KIR3DSw07*001 and

KIR3DSw09*003 fused to the Fc domain of murine IgG2A followed by AF647-conjugated goat

anti-mouse IgG and PE-conjugated MHC class I-specific antibody (W6/32). Data was collected

using a FACSymphony™ A3 Cell Analyzer and analyzed using FlowJo 10.8.0 for Mac OSX.

FIGURE 4. KIR3DL07*004, KIR3DS02*004, and KIR3DSw09*003 interact with Mamu-

B*045:03. (**A**) JNL cells expressing the extracellular domains of KIR3DL07*004,

KIR3DS02*004, and KIR3DSw09*003 fused to the transmembrane and cytoplasmic domains of

human CD3ζ were incubated overnight at a 1:1 E:T ratio with 721.221 cells expressing the

indicated Mamu-B molecules. The bar graphs represent the mean and standard deviation (error

bars) of luciferase activity (RLU) from triplicate wells of KIR-CD3ζ JNL cells incubated with

Mamu-B+721.221 cells (blue), parental 721.221 cells (red) and anti-Flag plus anti-mouse

antibodies (gray). Asterisks denote significant differences by one-way ANOVA with Dunnett

test (**** $p < 0.0001$). These results are representative of at least three independent experiments.

(**B**) Parental and Mamu-B*045:03+721.221 cells were stained with Near-IR LIVE/DEAD dye,

KIR-Fc fusion proteins consisting of the extracellular domains KIR3DL07*004,

KIR3DS02*004, and KIR3DSw09*003 fused to the Fc domain of murine IgG2A followed by

AF647-conjugated goat anti-mouse IgG and PE-conjugated MHC class I-specific antibody

(W6/32). Data was collected using a FACSymphony™ A3 Cell Analyzer and analyzed using

FlowJo 10.8.0 for Mac OSX.

FIGURE 5. Mamu-B*045:03 has an α1-domain similar to Mamu-A molecules. (**A**) Amino acid

sequence alignment of rhesus macaque MHC class I molecules. α1 domain residues 77-83

corresponding to Bw4 or Bw6 motifs are underlined and shaded sequences correspond to

predicted KIR contact sites (80). Positions of identity are indicated by periods and amino acid

differences are identified by their single-letter code. Phylogenetic trees of coding nucleotide

sequence for the α1-α3 domains of Mamu-A and -B (**B**) and the α1 domains of Mamu-A and -B

molecules (**C**). Neighbor-joining analysis was performed using MAFFT (version 7, publicly

available at https://mafft.cbrc.jp/alignment/server/). Bootstrap support of greater than 50% is

indicated.

FIGURE 6. Phylogeny of the D0-D2 domains of rhesus macaque KIRs reflects ligand

recognition. Neighbor-joining analysis of predicted amino acid sequences of the D0-D2 domains

of human and rhesus macaque KIRs was performed using MAFFT. Bootstrap support of greater

than 50% is indicated. Human KIR3DL1*0010101 and KIR3DL02*0010101 were used as an

outgroup. Rhesus macaque KIRs are color coded according to their MHC class I ligands: Mamu-

A-related (dark blue), Mamu-AG2*01:01 (blue), Mamu-B*045:03 (light blue), Mamu-Bw4

(green), and no ligands found (gray).

A

D0 Domain

```
                    10        20        30        40        50        60        70        80        90        100
Consensus   HTGGQDKTFLSARPSAVVPQGGHVTLRCYYRRGLNNFTNFTLYKDDRSHVPIFHSRIFQESFLMGPVTPAHAGTYRCRGSYPHSPTEWSALSDPLAIXVT
KIR3DL02*004  Y....N...W..L........H..G.---.N...........................................L....TP.N....R..
KIR3DL04*001  ................L........H..........PG..HQ...........D.........Q.........E....M..
KIR3DL07*004  .......F.............D...........................................R..
KIR3DL10*002  Y......VQ..L........F..........N...Q.........D.........M..
KIR3DL11*001  .....S..W..P.........Q.H....F.........PG..HQ...........M..
KIR3DLw34*002 Y....N...W..P.........H..G.---.N.......I...............R..
KIR3DS01*003  .................FQ..H................Q.........R..
KIR3DS02*004  ......F.............D..................................D..
KIR3DS03*003  .....I..VQ..L.......M..H....Y..........N.........D.........M..
KIR3DS04*003  ...........Q.H....F...........V...A.........M..
KIR3DS05*002  ...NA...W..P.........H..G.---.N..................P.N....R..
KIR3DS06*002  ..A..N..F.W..P.....L.....H.........................M..
KIR3DSw07*001 .......L.............D..............V..N...........R..
KIR3DSw09*003 ...D...F.............D................R..
```

D1 Domain

```
                    110       120       130       140       150       160       170       180       190       200
Consensus   GVHRKPSLLALPGPLVKSGETVTLQCSSDTVFXHFFLHSEVTFEKPLHLVGELHGGGSQANYSINSTTSDLAGTYRCYGSVTHSPYVLSAPSDPLDIVIT
KIR3DL02*004  ...K.................E....R...L.E.........................MM...F....F..
KIR3DL04*001  .............I..G..........EL...........................
KIR3DL07*004  ...............E.......N.....................D..
KIR3DL10*002  .............I..G..........EE.F...........N..
KIR3DL11*001  ...............G..........EL.............E.F...........N..
KIR3DLw34*002 ...K..............E....R...L.E..................K.......F..
KIR3DS01*003  ...K..............G..............................
KIR3DS02*004  ...............................R.......KM....H..
KIR3DS03*003  ...............G....................K.....E..
KIR3DS04*003  ...........I...I..G..........EL...........T..
KIR3DS05*002  ...K..............E....R...L.E..................K.......F...F....F..
KIR3DS06*002  ...............L................N...........K..E..........R..
KIR3DSw07*001 ...........D........G.....................K...E..
KIR3DSw09*003 ...............I.......E..........N..
```

D2 Domain

```
                    210       220       230       240       250       260       270       280       290       298
Consensus   GKYEKPSLSAQPGPTVQAGENVTLSCSSXNSFDMYHLSREGEARELSLSAVPSVNGTFQADFPLGPATHGGTYRCFGSFRTAPYKWSDPSDPLPVSVT
KIR3DL02*004  ...............I........RC...............................Q..
KIR3DL04*001  ........................Q................S..............Q.....H..
KIR3DL07*004  ........................Q..........................
KIR3DL10*002  ........................Q......S...V...R.....G..............I.....IT....NS....L..
KIR3DL11*001  ........................I........RC..............................
KIR3DLw34*002 .L......................RC..........R..........G.....N..H..
KIR3DS01*003  .L..............D.......QI...............S...Q.....HI..
KIR3DS02*004  .L......................RR..........R..........G.....H.T....H..
KIR3DS03*003  ........................Q..........G.....Q.........................
KIR3DS04*003  ........................RC..........I.................T.....K..
KIR3DS05*002  .L......................RR..........-.......................TS...H..
KIR3DS06*002  .L......................Q.........A..R.....G.....H.T....H..
KIR3DSw07*001 ........................Q..........G.....Q..R..................Q..
KIR3DSw09*003 ........................RC..........R..........G.....H..
```

B