**Title:** Change point estimation by the mouse medial frontal cortex during probabilistic reward

learning

**Authors:** Huriye Atilgan[1], Cayla E. Murphy[1], Hongli Wang[2], Heather K. Ortega[2], Lucas Pinto[3,4],

Alex C. Kwan[1,2,5]*

[1]Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut, 06511,

USA

[2]Interdepartmental Neuroscience Program, Yale University School of Medicine, New Haven,

Connecticut, 06511, USA

[3]Princeton Neuroscience Institute, Princeton University, Princeton, NJ 08544, USA

[4]Department of Neuroscience, Northwestern University Feinberg School of Medicine, Chicago, IL

60611, USA

[5]Department of Neuroscience, Yale University School of Medicine, New Haven, Connecticut,

06511, USA

*Correspondence: alex.kwan@yale.edu

**Summary**

There are often sudden changes in the state of environment. For a decision maker, accurate

prediction and detection of change points are crucial for optimizing performance. Still unclear,

however, is whether rodents are simply reactive to reinforcements, or if they can be proactive to

estimate future change points during value-based decision making. In this study, we characterize

head-fixed mice performing a two-armed bandit task with probabilistic reward reversals. Choice

behavior deviates from classic reinforcement learning, but instead suggests a strategy involving

belief updating, consistent with the anticipation of change points to exploit the task structure.

Excitotoxic lesion and optogenetic inactivation implicate the anterior cingulate and premotor regions

of medial frontal cortex. Specifically, over-estimation of hazard rate arises from imbalance across

frontal hemispheres during the time window before the choice is made. Collectively, the results

demonstrate that mice can capitalize on their knowledge of task regularities, and this estimation of

future changes in the environment may be a main computational function of the rodent dorsal

medial frontal cortex.

## Introduction

In life, we experience twists and turns – discrete events that abruptly alter the state of environment. In some cases, the change is a one-time occurrence that is impossible to predict. We must then adjust by assessing the new situation following the change. However, in other cases, the changes may occur repeatedly with certain tendencies. For example, a favorite chef in a restaurant may have a recurring schedule where she cooks throughout the year, except in the summer for 3 – 5 weeks when she would take a vacation and lets a substitute take over. As a patron, it would be advantageous to learn this pattern, anticipate the impending switches, and maximize the chance of receiving a delicious outcome. While it is evident that humans can estimate change points and leverage the information in decision-making, whether animals such as mice have this ability and the neural substrates supporting the computations remain unclear.

A classic paradigm to study decision-making in response to repeated changes is the two-armed bandit task. Each trial, the animal has two options, and each option is associated probabilistically with a reward. After a certain number of trials, the reward probabilities are switched among the options. The two-armed bandit task is widely used because it can be tested in different species including humans (Tsuchida, Doll, and Fellows 2010; Evers et al. 2005; O'Doherty et al. 2001), monkeys (Costa et al. 2015; Donahue and Lee 2015; Clarke, Robbins, and Roberts 2008), rats (Groman et al. 2019; Hamid et al. 2016; Sul et al. 2010; Bari et al. 2010; Ito and Doya 2009; Samejima et al. 2005), and mice (Grossman, Bari, and Cohen 2022; Hattori et al. 2019; Tai et al. 2012). Moreover, the paradigm has translational significance because it can reveal defects from pharmacological interventions or in animal models for psychiatric disorders (Costa et al. 2015; Groman et al. 2018; Liao and Kwan 2021).

Most analyses of rodents performing two-armed bandit and related decision-making tasks have relied on simple reinforcement learning schemes such as Q-learning algorithms (Wang et al. 2022; Bari et al. 2019; Hattori et al. 2019; Groman et al. 2019; Ito and Doya 2015; Sul et al. 2010), but see (Ito and Doya 2015). Q-learning algorithms assume that animals learn from experience, and therefore choice behavior adapts only after a change point has occurred. By contrast, recent studies in monkeys and humans have challenged this assumption. Namely, primates can exploit predictable structure in a task and adjust for an impending change point (Bartolo and Averbeck 2020; Jang et al. 2019; Costa et al. 2015). Indeed, under a few select situations, rodents also seem to make inferences about hidden states (Liu, Xin, and Xu 2021; Vertechi et al. 2020; Starkweather,

Gershman, and Uchida 2018). These recent results hint at the possibility that mice may leverage

their knowledge of task structure for probabilistic reward learning.

To test the possibility that rodents may estimate change points during probabilistic reward learning,

we trained head-fixed mice on a two-armed bandit task. By analyzing a sizable data set totaling

1,007 sessions involving 15,352 reversals, we demonstrate that mice are sensitive to impending

switches in reward probabilities, because they alter their choices prior to the actual change points.

We show that the animals' choice behavior can be modelled effectively with a Bayesian framework

involving belief updating and choice kernels. Furthermore, we performed unilateral and bilateral

excitotoxic lesions and optogenetic inactivation to demonstrate mechanistically how the anterior

cingulate and premotor regions of medial frontal cortex may be involved in the computation.

Together, the results indicate that mice can take advantage of the task structure to solve a classic

probabilistic reward learning task and implicate the dorsal medial frontal cortex as a locus in the

accurate estimation of future changes in the state of environment.

**Results**

**Mice use their knowledge of the task structure during a two-armed bandit task**

We trained head-fixed C57BL/6J mice on a two-armed bandit task involving probabilistic reward

reversals. On each trial, the mouse could choose left or right by a directional tongue lick. The two

options were associated with different reward probabilities, e.g., "70:10" for 70% and 10% chance

to receive water from the left and right spouts respectively (**Figure 1A – B**). The reward

probabilities would flip when the animal reaches the switching condition, which is a performance-

dependent number of trials to fulfill a criterion ($L_{Criterion}$, 10 trials choosing the better option) followed

by a performance-independent random number of trials ($L_{Random}$, drawn from a geometric

distribution with $p = 0.0909$ and truncated at 30). In an example session shown in **Figure 1C**, the

animal performed more than 500 trials, including 15 reversals of 70:10 and 10:70 blocks. To

visualize how animals adjust to the sudden changes in reward probabilities, we aligned the trials by

the time of block switches. As expected, mice primarily chose the better option pre-switch, and then

quickly adapted their preferred action post-switch (**Figure 1D**, n = 31 mice, 617 sessions, 9,163

blocks).

**Figure 1: Mice were sensitive to block length and leverage this information during the two-armed bandit task**

**(A)** The mouse makes a left or right choice via tongue lick after the go cue. Depending on the reward probabilities, the choice might lead to water. **(B)** Trials were organized into blocks, each with distinct reward probabilities: "70:10" (70% chance to receive water for left choice; 10% for right) or "10:70" (10% for left; 70% for right). The block switches after the animal choose the high-reward-probability side ten times ($L_{Criterion}$) plus an additional random number of trials ($L_{Random}$, drawn from exponential distribution, up to 30 trials). **(C)** Performance of a mouse in one example session. The top row shows reward probabilities for left and right options. The bottom row shows the animal's choices and the outcomes. **(D)** Choice behavior around block switches. Thin line, mean values for individual animal. Thick line, mean values and SEM for all animals. **(E)** Histogram of $L_{Random}$. For all blocks with $L_{Criterian} \leq 20$. Colors indicate the 4 ranges of $L_{Random}$ for subsequent analyses. **(F)** Choice behavior around block switches, plotted separately for the 4 ranges of $L_{Random}$. Mean values and SEM for all animals. **(G)** The probability of choosing the better option on the trial immediately preceding the switch, as a function of $L_{Random}$ for the block preceding the switch. Mean values and SEM for all animals. (H) The number of trials to reach midpoint (when animal is equally likely to choose either option) as a function of $L_{Random}$ for the block preceding the switch. Mean values and SEM for all animals. n = 31 mice, 617 sessions.

An important parameter in our task is $L_{Random}$, which dictates the frequency of reversals. Although the animals cannot know the exact value of $L_{Random}$ before each switch because it is drawn randomly, it is possible for the mice to learn the statistical distribution of $L_{Random}$. This knowledge may then be used to infer that the more trials that an animal stays at the better option, the more likely that a block switch might have already occurred. To determine if mice were making use of such knowledge of the task structure, we analyzed the subset of 7,396 blocks in which animals were performing at expert level ($L_{Criterion} \leq 20$). **Figure 1E** shows the histogram of $L_{Random}$ values for

these trial blocks, exhibiting the geometric distribution as the task was designed. We found that if

$L_{Random}$ was large in the preceding block, the animals tended to choose the better option less

frequently prior to the block switch, and subsequently adapted faster after the block switch (**Figure**

**1F**). These results suggest that the mice may anticipate an impending change point and adjust their

behavior prior to the block switch.

To quantify the observations, we computed *P (better option) $_{pre\text{-}switch}$*, the probability of selecting the

better option in the trial immediately before the block switch, and *trials to reach midpoint*, the

number of trials from switch for *P (better option)* to reach 0.5. These analyses confirmed the

influence of $L_{Random}$ on choice patterns around a block switch (main effect of $L_{Random}$: F (30, 5544) =

6.0743, *P* < 0.001, one-way ANOVA; **Figure 1G**) and their speed to adjust after a block switch

(main effect of $L_{Random}$: F (30, 711) = 2.1316, *P* < 0.001, one-way ANOVA; **Figure 1H**). We reiterate

that the animal could not predict the value of $L_{Random}$, which was drawn randomly for each block.

However, the tendencies around a block switch are consistent with learning the task structure,

namely aspects of the statistical distribution of $L_{Random}$, presumably through repeated training over

dozens of sessions on the two-armed bandit task. As the animal dwells in a block selecting the

same better option for many trials, it becomes more probable that a change point in reward

probabilities has occurred and therefore the animals should explore the alternate option more.

Overall, these results demonstrate that mice were sensitive to the block length – a key feature of

the task structure – and could leverage this information during the two-armed bandit task.

**Effects of unilateral lesion of ACAd/MOs on choice behavior around switches**

Prior studies implicated the anterior cingulate cortex in behavioral flexibility in the face of variability

in the environment (Behrens et al. 2007; Soltani and Izquierdo 2019). The related region in the

mouse is the dorsal aspect of the medial frontal cortex, encompassing the anterior cingulate

(ACAd) and medial secondary motor (MOs) areas (Barthas and Kwan 2017; Laubach et al. 2018).

To determine the role of ACAd/MOs, we trained mice until they reached expert performance, and

then performed unilateral excitotoxic lesion by injecting ibotenic acid into the ACAd/MOs region in

the left or right hemisphere (n = 5 and 4 mice respectively, 200 pre-lesion and 142 post-lesion

sessions in total; **Figure 2A**). For clarity, we will collapse the two groups and refer to trial blocks as

'lesion', if the lesioned side was the better option, or 'contra' if the side contralateral to the lesion

was the better option (**Figure 2B**). Post hoc histology with cresyl violet staining confirmed the loss

of cell bodies at the targeted ACAd/MOs location (**Figure 2C – D**). After the lesion, animals

performed a similar number of trials and block switches (**Supplementary Figure 2.1**) and had no

motor deficit in licking (**Supplementary Figure 2.2)**. Post-lesion mice exhibited block length-

dependent choice patterns (**Figure 2E**). However, with the unilateral loss of ACAd/MOs, for the

switch from lesion block to contra block, this tendency to choose the worse option pre-switch

exacerbated with increasing $L_{Random}$ (left panel, **Figure 2E**). Summary of the data reaffirmed that

animals with unilateral ACAd/MOs lesion were selecting the worse option at the expense of

exploiting the better option pre-switch, specifically when $L_{Random}$ was large in the preceding block

(main effect of lesion: $P < 0.001$, main effect of $L_{Random}$: $P < 0.001$, lesion * $L_{Random}$ interaction: $P =$

0.044, three-way ANOVA; **Figure 2F, Supplementary Table 2.1**). Although they appeared to adapt

faster after the switch relative to control animals (main effect of lesion: $P < 0.001$, main effect of

side: $P = 0.003$, lesion * side interaction: $P = 0.038$, $L_{Random}$ * side interaction: $P = 0.004$, three-way

ANOVA; **Figure 2G**), overall the performance suffered after the lesion (main effect of lesion: $P =$

0.027, main effect of side: $P = 0.045$, main effect of $L_{Random}$: $P < 0.001$, lesion * $L_{Random}$ interaction: $P$

$= 0.027$, three-way ANOVA; **Figure 2H**). The data therefore show that unilateral lesion of the

ACAd/MOs impairs the proper estimate and use of task structure knowledge during probabilistic

reward learning.

**Figure 2: Unilateral lesion of ACAd/MOs altered block-length-dependent choice behavior and impaired overall**

**performance**

**(A)** Schematic representation of the unilateral excitotoxic lesion via injection of ibotenic acid. **(B)** Lesion blocks refers to

blocks in which the lesioned side is the better option. Contra blocks refer to blocks in which the lesioned side is

contralateral to the better option. **(C, D)** Post hoc histology with cresyl violet staining to confirm the loss of neurons in

ACAd/MOs. **(E)** Choice behavior around block switches, plotted separately for the 4 ranges of $L_{Random}$. Black, pre-lesion.

Green, post-lesion. Left, switches from lesion block to contra block. Right, switches from contra block to lesion block.

Mean values and SEM for all animals. **(F)** The probability of choosing the better option on the trial immediately preceding

the switch, as a function of $L_{Random}$ for the block preceding the switch. Black, pre-lesion. Green, post-lesion. Mean values

and SEM for all animals. **(G)** Similar to (F) for number of trials to reach midpoint (when animal is equally likely to choose

either option). **(H)** Similar to (F) for hit rate (probability for animal to choose the better option). For (F) – (H), significant

main effects and interactions from three-way ANOVA were indicated ($P < 0.05$). n = 9 mice, 200 pre-lesion sessions and

142 post-lesion sessions.

**A hybrid model of belief and choice kernels to explain the animals' behavior**

To gain insight into the empirical findings, we fitted different computational models to the data. We

were specifically drawn to two emerging ideas in the field of decision-making. First, the concept of

belief enables an agent to apply their knowledge of the task structure (Jang et al. 2019). Namely,

the two-armed bandit task in this study can be deconstructed as an environment with 2 states

('70:10' and '10:70'), each with their optimal action (choose left and right, respectively). In this

scheme, each trial, the agent maintains a belief, which is a mixture of probabilities that the task is

currently at each of the states, and acts accordingly. After the outcome is presented, the belief is

updated based on the outcome and the agent's estimate of the likelihood of a change point, i.e., the

reversal of reward probabilities, which is known as the hazard rate $H$. Second, choice kernels can

be used to capture an agent's tendency to repeat the previous actions (Wilson and Collins 2019).

The choice kernels are updated based on the prior action, scaled by a learning rate $\alpha_K$. Our belief-

CK model contains components for belief and choice kernels, and integrates their outputs for action

selection based on a softmax function with separate inverse temperature parameters for belief, $\beta$,

and choice kernel, $\beta_K$ (**Figure 3A, Supplementary Figure 3.1 - 3.4**). Fit to an example session of

animal data suggests that this 4-parameter belief-CK model can recapitulate the choice behavior of

the mouse in the two-armed bandit task (**Figure 3B – E**).

**Figure 3: A hybrid model of beliefs and choice kernels to explain the behavior**

**(A)** The schematic representation of the belief with choice kernel model (belief-CK). The model has four parameters: $H$

(hazard rate), $\beta$ (inverse temperature for belief), $\alpha_K$ (learning rate for choice kernel) and $\beta_K$ (inverse temperature for

choice kernel). **(B – E)** An example session along with the fits from the belief-CK model, including reward probabilities for

left and right options (B) the running-average of probability of choosing right for the animal (black) and model (purple) (C),

the belief that the left option is associated with reward probability of 10% (pL10, blue) or 70% (pL70, red) (D), and the

choice kernels for left (blue) and right options (red) (E). **(F)** Model comparison between the belief-CK model and 7 other

models. Lower log BIC values indicate a better fit. **(G)** The tally of the best-fitting model for each animal. **(H)** The

probability of choosing the better option on the trial immediately preceding the switch, as a function of $L_{Random}$ for the

block preceding the switch. Black, mice. Purple, simulated performance using the belief-CK model with best-fitting

parameters. Mean values and SEM for all animals. **(I)** Similar to (H) for number of trials to reach midpoint (when animal is

equally likely to choose either option). **(J)** Similar to (H) for the tendency to win-stay on the 5 trials preceding the switch.

**(K)** Similar to (H) for the tendency to lose-switch on the 5 trials preceding the switch. n = 31 mice, 617 sessions.

We compared the belief-CK model against 7 other computational models (see Methods). We

started with the win-stay, lose-switch (WSLS) and 3 classic reinforcement learning algorithms

including Q-learning (Q-RPE), Q-learning with forgetting (F-Q-RPE), and Q-learning with differential

forgetting (DF-Q-RPE) (Ito and Doya 2015, 2015). We then examined effects of adding choice

kernels, by testing DF-Q-RPE with choice kernels (DF-Q-RPE-CK), because DF-Q-RPE was the

best fit in the initial set of 4 algorithms, and F-Q-RPE with choice kernels (F-Q-RPE-CK), because

this model has the same number of free parameters as the belief-CK model. Finally, we also tested

the belief model alone without choice kernels. Model comparison based on Bayesian information

criterion (BIC) revealed that inclusion of choice kernels improved the fits significantly. Moreover, the

belief-CK model had the lowest BIC values (**Figure 3F**; belief-CK versus F-Q-RPE-CK: $t_{60}$= 2.562,

$P$ = 0.013, paired t-test; belief-CK versus DF-Q-RPE-CK: $t_{60}$= 2.313, $P$ = 0.024), and was the best

fit for 30 out of 31 animals in this study (**Figure 3G**). For each session, we can simulate the belief-

CK model using the best-fitting parameters and compare the tendencies of the simulated and

experimental data. This exercise shows that the belief-CK model can capture the $L_{Random}$-dependent

choice behavior in the experimental data (**Figure 3H – K**), which is not possible with the classic

reinforcement learning algorithm DF-Q-RPE (**Supplementary Figure 3.5**). These analyses

demonstrate that simple models of reward-based learning such as Q-learning algorithms cannot

fully account for the observed choice behavior. Instead, the results support our intuition that mice

were estimating change points, which is formalized as the hazard rate $H$ in the belief-CK model.

**Unilateral ACAd/MOs lesion led to side-specific increase in hazard rate for change points**

Next, we applied the computational model to quantify the effect of unilateral ACAd/MOs lesion. To

account for the possibility of side-specific alterations, we modified the 4-parameter belief-CK model

to include 6 parameters to include differential learning for the sides ipsilateral and contralateral to

lesion (see Methods; $H_{lesion}, H_{contra}, \alpha_{K\ lesion}, \alpha_{K\ contra}, \beta,$ and $\beta_K$). After fitting the expanded

model to animal data, we compared pre- versus post-lesion performance in two ways. First, on a

per-animal basis, sessions before or after the lesion were concatenated for fitting to yield one set of

pre-lesion parameters and one set of post-lesion parameters for each animal. Second, on a per-

session basis, each session was analyzed separately and the fitted parameters were summarized.

These analyses revealed a side-specific increase in hazard rate after unilateral ACAd/MOs lesion.

The exaggerated hazard rate for the side contralateral to lesion $H_{contra}$ was detected on a per-

animal basis (**Figure 4A**; pre- vs. post-lesion, $H_{contra}$: $P$ = 0.004; post-lesion, $H_{contra}$ vs. $H_{lesion}$: $P$

= 0.810, Wilcoxon signed-rank test), and on a per-session basis (**Figure 4B**; pre- vs. post-lesion,

$H_{contra}$: $P$ = 0.001; post-lesion, $H_{contra}$ vs. $H_{lesion}$: $P$ < 0.001, Wilcoxon rank sum test). Unilateral

ACAd/MOs lesion also led to an increase in choice perseveration for both sides, reflected as higher

choice-kernel learning rates (**Figure 4C – D**; per-animal, pre- vs. post-lesion, $\alpha_{K\ lesion}$: $P$ = 0.012;

$\alpha_{K\ contra}$: $P$ = 0.004; per-session, pre- vs. post-lesion, $\alpha_{K\ lesion}$: $P$ < 0.001; $\alpha_{K\ contra}$: $P$ < 0.001,

Wilcoxon ranked sum test). Action selection depends on the inverse temperature sum, $\beta + \beta_K$,

reflecting the exploration-exploitation balance, and inverse temperature ratio,——, reflecting the

relative reliance on belief over choice kernels. There was no detected difference in inverse

temperature sum between pre- and post-lesion animals (**Figure 4E – F**, per animal, $P = 0.567$; per

session $P = 0.858$, Wilcoxon signed-rank test). By contrast, the inverse temperature ratio was

heightened after the lesion (**Figure 4G – H**, per animal $P = 0.038$; per session $P = 0.001$, Wilcoxon

signed-rank test). Collectively, these analyses show that the consequences of unilateral ACAd/MOs

lesion are a contralateral side-specific increase in hazard rate, and broad increases in choice

perseveration and reliance on belief for action selection.

268
**Figure 4: Effects of unilateral lesion of ACAd/MOs is consistent with a side-specific increase in hazard rate**

**(A)** The hazard rates, before and after lesion, extracted by fitting the belief-with-choice-kernel model on a per-animal

basis. Square, hazard rate for side ipsilateral to lesion. Cross, hazard rate for side contralateral to lesion. Inset, violin plot

of the same data. **(B)** The hazard rates, before and after lesion, on a per-session basis. Mean and SEM. **(C – D)** Similar

to (A – B) for learning rate for choice kernel. **(E)** The inverse temperature sum, before and after lesion, on a per-animal

basis. **(F)** The inverse temperature sum, before and after lesion, on a per-session basis. **(G - H)** Similar to (E – F) for

inverse temperature ratio. *, $P < 0.05$. n.s., not significant. n = 9 mice, 190 pre-lesion sessions and 140 post-lesion

sessions.

**Accurate change point estimation depends on the balance between the left and right hemispheres**

The results so far from unilateral lesions suggest two potential mechanisms for change point estimation. The first possibility is that the computation of change point estimation is lateralized, such that the left hemisphere is involved in estimation for the right side, and vice versa. If this is the case, for a bilateral lesion, we would expect aberrant increases of hazard rates for both sides. The second possibility is that the estimation of change points involves inter-hemispheric coordination, which was perturbed by disruption of one hemisphere. If true, the lack of medial frontal cortex on both sides could nullify their respective maladaptive influences on behavior, and we may observe no or milder deficit after a bilateral lesion of ACAd/MOs. To distinguish between these two possibilities, we injected ibotenic acid bilaterally to the left and right ACAd/MOs regions in expert mice. Animals with bilateral lesions performed fewer trials per session, and accordingly fewer block switches (**Figure 5A – B, Supplementary Figure 5.1;** $P < 0.001$, Wilcoxon signed-rank test), but had no motor deficit (**Supplementary Figure 5.2**). Surprisingly, and in line with inter-hemispheric coordination, there was no detectable change in the $L_{Random}$-dependent choice behavior (**Figure 5C – D**, **Supplementary table 5.1**), and no significant changes in the latent decision parameters including hazard rates (**Figure 5E – H**). Comparison to sham animals in which saline was injected unilaterally (**Figure 5I-P, Supplementary table 5.2**) highlights again that the only effect of bilateral lesion was diminished motivation to perform the task. The reduced motivation to perform trials is consistent with a prior work from the lab (Siniscalchi et al. 2016). More importantly, these results argue against change point estimation as a lateralized computation in ACAd/MOs, but rather point to unbalance between the hemispheres as the reason for behavioral deficits.

300
**Figure 5: Effects of bilateral and sham lesions of ACAd/MOs**

**(A)** The number of trials performed in each session, before and after bilateral lesion, on a per-session basis. Mean and

SEM. **(B)** Similar to (A) for the number of block switches in each session. **(C)** The probability of choosing the better option

on the trial immediately preceding the switch, as a function of $L_{Random}$ for the block preceding the switch, before and after

bilateral lesion, on a per-session basis. Mean and SEM. Significant main effects and interactions from three-way ANOVA

were indicated ($P < 0.05$). **(D)** Similar to (C) for number of trials to reach midpoint (when animal is equally likely to choose

either option). **(E)** The hazard rates, before and after bilateral lesion, extracted by fitting the belief-CK model on a per-

session basis. Mean and SEM. **(F)** Similar to (E) for learning rate for choice kernel. **(G)** Similar to (E) for inverse

temperature sum. **(H)** Similar to (E) for inverse temperature ratio. **(I – P)** Similar to (A – H) for sham controls with

unilateral saline injection. n.s., not significant. For bilateral lesion, n = 4 mice, 105 pre-lesion sessions and 61 post-lesion

sessions. For saline control, n = 4 mice, 117 pre-lesion sessions and 53 post-lesion sessions.

**Medial frontal cortex impacts the decisions during action selection**

Permanent lesions are advantageous because, compared to transient inactivation, they avoid

confounds from potential acute off-target effects (Otchy et al. 2015). However, it is unclear which of

the lesion-induced behavioral changes are due directly to ACAd/MOs disruption, and which other

consequences are compensatory adjustments. For this reason, we performed transient inactivation

experiments using optogenetics. Mice were implanted with a clear-skull cap that has ~50% optical

transmission (**Supplementary Figure 6.1A**). For photostimulation, we used a laser-steering system

(Pinto et al. 2019), in which the excitation beam from a 473 nm laser was steered by a set of mirror

galvanometers to specific locations with high spatial and temporal resolutions (**Figure 6A**). We

calibrated the linearity of the steered coordinates as a function of galvanometer voltages and the

spatial profile of the laser beam (**Supplementary Figure 6.1B – C**). We demonstrated that the

system can effectively manipulate neural activity by showing elevated c-fos immunohistostaining

after unilateral photostimulation of ACAd/MOs in *CaMKIIa$^{Cre}$;Ai32* animals (**Supplementary Figure**

**6.1D – F**), and inducing biased tongue licks during the two-armed bandit task by inhibiting the

anterolateral motor cortex in *Pvalb$^{Cre}$;Ai32* animals **(Supplementary Figure 6.2)**

**Figure 6: Optogenetic inactivation in pre-choice, but not post-choice, period reproduced the deficit in change-point estimation**

**(A)** The schematic representation of experimental setup. **(B)** CCD image of a mouse with a cleared skull cap. The two blue crosses indicate the locations of the photostimulation, i.e. left and right ACAd/MOs. **(C - D)** The trial and block structures, and the timing of the photostimulation. **(E)** Choice behavior around block switches, plotted separately for the 4 ranges of $L_{Random}$. Black, control trials. Green, pre-choice inactivation trials. Left, switches from ipsi block (photostimulated side was the high-reward-probability option). Right, switches from contra block ((photostimulated side was the low-reward-probability option). Mean values and SEM for all animals. **(F)** Similar to (E) for post-choice inactivation. **(G)** The hazard rates extracted by fitting a modified belief-CK model, for pre- and post-choice inactivation, on a per-animal basis. **(H)** Similar to (G) for learning rate for choice kernel. **(I)** Similar to (G) for inverse temperature sum. **(J)** Similar to (G) for inverse temperature ratio. n = 6 animals.

To suppress excitatory activity in ACAd/MOs during block switches, we used *Pvalb^Cre^;Ai32* animals in which the channelrhodopsin ChR2 was selectively expressed in parvalbumin-expressing (PV) GABAergic interneurons (n = 6). Targeted photostimulation would activate PV interneurons in the left or right ACAd/MOs (**Figure 6B**), which would in turn silence local excitatory spiking activity

(Guo et al. 2014; Li et al. 2019). These transient inactivations were applied either at the time of

action selection ("pre-choice", from cue onset to lick response) or after the outcome ("post-choice",

from lick response for 2 s; **Figure 6C**), and on every trials across two consecutive blocks such that

activity was suppressed before and after certain block switches (**Figure 6D**). **Figures 6E–F** show

the effects of pre- and post-choice inactivation on behavior around block switches, separating the

data based on whether the high-reward-probability side is ipsilateral or contralateral to the

perturbation. Quantification of the behavior showed that pre-choice inactivation, but not post-choice

inactivation, significantly affected the change point estimation (**Supplementary Figure 6.3,** main

effect of block length: F(3, 1999) = 49.778, $P < 0.001$; main effect of stimulation type: F (1, 1999) =

7.974, $P = 0.004$; interaction between block length and stimulation: F(3, 1999) = 2.674, $P = 0.046$;

interaction between block length, stimulation and stimulation type: F(3, 1999) = 2.625, $P$ = 0.049).

Fitting to the belief-CK model, expanded to account for the optogenetic stimulation (see Methods;

$H_{control}$, $H_{ipsi\ stim}$, $H_{contra\ stim}$, $\alpha_{K\ control}$, $\alpha_{K\ ipsi\ stim}$, $\alpha_{K\ contra\ stim}$, $\beta_{control}$, $\beta_{stim}$, $\beta_{K\ control}$ and

$\beta_{K\ stim}$), highlights the strongest effect is an acute change to hazard rate contralateral to the

transient inactivation, although there were variations across individual animals and effect was not

statistically significant (**Figure 6G**; pre-choice inactivation, $H_{contra\ stim}$ vs. $H_{ipsi\ stim}$: $P = 0.156$;

post-choice inactivation, $H_{contra\ stim}$ vs. $H_{ipsi\ stim}$: $P = 0.687$; pre-choice inactivation, $H_{contra\ stim}$ vs.

$H_{control}$: $P = 0.094$; post-choice inactivation, $H_{contra\ stim}$ vs. $H_{control}$: $P = 0.562$, Wilcoxon signed-

rank test). There were no detectable effects of transient inactivation on $\alpha_K$, $\frac{\beta}{\beta+\beta_K}$, and $\beta + \beta_K$

(**Figure 6H – J**). Together with the results from unilateral lesions, we interpret these findings from

acute inactivation to indicate that ACAd/MOs is involved specifically in the change point estimation

process, which occurs during the pre-choice period.

**Discussion**

This study provides evidence that mice anticipate impending change points by altering their choices

prior to switches in a classic probabilistic reward learning task. Computational analyses indicate

that the animals' choice behavior is consistent with a model of belief updating and choice

perseveration. Causal perturbation experiments emphasize the role of the ACAd/MOs region of the

medial frontal cortex. Crucially, as discussed below, the collective results from the range of

manipulations employed – unilateral and bilateral lesions, as well as pre- and post-choice

optogenetic inactivation – provide important insights that can constrain the potential neural

mechanisms underlying change-point estimation during decision-making.

Although many studies of decision-making in rodents relied on analyses involving Q-learning
algorithms (Wang et al. 2022; Bari et al. 2019; Hattori et al. 2019; Groman et al. 2019; Ito and Doya
2015; Sul et al. 2010), there are other reports suggesting deviations from simple reinforcement
learning. For instance, a pioneering study demonstrated that rats are exceedingly sensitive to
changes in reward rates, approximating an ideal observer (Gallistel et al. 2001). Moreover, several
studies using a variety of timing, operant conditioning, decision, and sensory categorization tasks
have found neural and behavioral data consistent with the use of belief in rodents (Karlsson, Tervo,
and Karpova 2012; Li and Dudman 2013; Liu et al. 2021; Starkweather et al. 2017; Vertechi et al.
2020). However, these prior studies are different because in some cases, the use of belief did not
necessarily confer better performance over other strategies for the task (Starkweather et al. 2017).
In other cases, the task employs a series of deterministic outcomes (Vertechi et al. 2020), which
strongly favors switching behaviors. Our study therefore extends these past results by showing
decisions consistent with belief updating in one of the most popular value-based decision-making
tasks used for human and animal studies (Sutton and Barto 2018).
393
To quantify the animals' behavior, we proposed a model involving belief updating with a fixed
hazard rate, which was motivated by a prior study (Jang et al. 2019) and adapted to fit our task
design. In this model, the agent understands that each action is associated with one of two reward
probabilities. Given information from its choice and reward history as well as knowledge of the
probability of a switch in reward probabilities, the agent infers the likelihood of the current states
associated with the actions. This is in sharp contrast to the Q-learning algorithms, where the agent
is implicitly ignorant of the task structure, and simply updates action values based on the last trial's
action and outcome. For belief updating, one limitation for our model is that the hazard rate is a
constant value within a session. This assumption seems reasonable because mice were trained on
the task extensively and probably accrue knowledge of the hazard rate based on experience of
multiple switches across many sessions. That said, in principle, it is possible for an agent to infer
the hazard rate as the task proceeds (Wilson, Nassar, and Gold 2010), which could be a
refinement for future analyses.
407
To determine the role of the medial frontal cortex, we started with excitotoxic lesions because the
approach has numerous merits (Vaidya et al. 2019). Specifically, lesions can mitigate diaschisis,
where the sudden loss of cortical excitations can be a confounding factor (Otchy et al. 2015).
Moreover, we employed unilateral lesions, such that we can compare effects between sides
ipsilateral and contralateral to the lesion in the same animal, serving as a rigorous internal control.

Results from these experiments demonstrated lateralized deficits from lesions of the medial frontal

cortex. This finding may be surprising, because although sensory and motor functions are typically

expected to be lateralized, it is less obvious that cognitive function may also be side-dependent.

However, we note that a few prior studies have also found side-specific decision-making deficits

from unilateral manipulations, such as effects of dorsal striatum on action values (Tai et al. 2012)

and effects of MOs on lapses in a multisensory task (Pisupati et al. 2021). The exact reason for

side-specific effects is unclear, although one possibility is that when decisions are intimately tied to

responses associated with lateralized motor actions, then there is embodiment and motor and

premotor cortical regions become involved in the neural computation (Bennur and Gold 2011).

The various deficits arising from lesion and optogenetic manipulations are useful for thinking about

the potential mechanisms for how the medial frontal cortex contributes to belief updating and more

specifically change-point estimation. An inaccurate estimate, which would reflect as altered hazard

rate in our computational model, can occur for several reasons: (1) error in estimating the value of

hazard rate, (2) error when using the hazard rate to update belief, and (3) error when using the

prior choice and reward to update belief. Reason (3) was not explicitly tested in our model fits but

could manifest as an apparent change in hazard rate. Among these possibilities, the first option

seems unlikely. In our task, an accurate value for hazard rate cannot be determined quickly but

must be calibrated by experiencing many switches across multiple sessions. This is difficult to

reconcile with the immediate deficit observed with pre-choice optogenetic inactivation. The third

option is also unlikely. Previous studies have shown that choice- and outcome-related signals arise

in the medial frontal cortex shortly after the outcome (Siniscalchi, Wang, and Kwan 2019; Sul et al.

2011), whereas optogenetic inactivation during this post-choice period was ineffective. Therefore, it

may be the case that the medial frontal cortex is involved in incorporating the likelihood of an

impending change point for estimating the current task state.

Furthermore, rather than computing using a probability such as the hazard rate, the animal may

instead approximate the process by employing simpler heuristics to predict the impending

occurrence of a change point. One intuitive heuristic, consistent with the reason for lateralized

deficits, is that the animals may rely on the recent choice history of the number of better options

chosen, which would indicate a higher likelihood of an impending switch. Here, the lack of effect

from bilateral lesions can shed light on the form of the heuristic. For example, one heuristic that can

work is a ratio of the number of recent left choices divided by the number of recent right choices,

and if the unilateral lesion effectively adds a multiplier to the side's choice history, then a bilateral

perturbation would lead to a null effect. Heuristics based on choice history are plausible because
the medial frontal cortex has long-lasting, persistent representation of past choices (Bari et al.
2019; Hattori et al. 2019; Siniscalchi et al. 2016; Sul et al. 2011). One caveat for this line of logic is
that it is based on a specific belief updating model. However, we discuss the implications to
illustrate how the results can inform the underlying neural basis.
452
To sum, the two-armed bandit task has gained widespread use in neuroscience and artificial
intelligence research because of its simplicity, translational significance, and amenability to
computational modeling. Our results show that mice may perform the task by not only updating
based on choices and outcomes, but also leverage knowledge of the environment to estimate
change points. The diminished or exaggerated use of this prior knowledge represents suboptimal
decision-making, which may underlie pathological behaviors in neuropsychiatric disorders that
involve dysfunctions of the medial frontal cortex.
460
461
462  **METHODS**
463  **Lead Contact**
464  Further information and requests for resources and reagents should be directed to and will be
465  fulfilled by the Lead Contact, Alex Kwan (alex.kwan@yale.edu)
466
467  **Materials Availability**
468  All published reagents and mouse lines will be shared upon request within the limits of the
469  respective material transfer agreements. Detailed plans including parts list for constructing the
470  behavioral training apparatus is available at https://github.com/Kwan-Lab/behavioral-rigs.
471
472  **Data and Code Availability**
473  Data and analysis software for this paper will be available at Github (https://github.com/Kwan-Lab).
474
**EXPERIMENTAL MODEL AND SUBJECT DETAILS**
**Mouse lines**
In this study, we used a total of 30 adult male mice (**Table 1**; 2 - 8 months old), including 24
C57BL/6J wild-type mice (#000664, Jackson Laboratory) for the lesion experiments, and 6
$Pvalb^{Cre}$;$ROSA^{CAG-ChR2-EYFP}$(Ai32) mice for the photostimulation experiments. The $Pvalb^{Cre}$;$ROSA^{CAG-}$
$^{ChR2-EYFP}$(Ai32) mice were generated by crossing the $Pvalb^{Cre}$ (B6.129P2-$Pvalb^{tm1(cre)Arbr}$/J; #017320,

Jackson laboratory) and *ROSA*$^{CAG-ChR2-EYFP}$*(Ai32)* (B6.Cg-*Gt(ROSA)26Sor*$^{tm32\ (CAG-}$

$^{COP4*H134R/EYFP)Hze}$/J; #024109, Jackson laboratory) strains. Mice were housed in groups of 2 – 5 per

cage in a 12h:12h light:dark cycle with ad libitum access to food. All of the experiments were

completed during the light cycle. Experimental procedures were approved by the Yale University

Institutional Animal Care and Use Committee.

**METHOD DETAILS**

**Surgery for lesion experiments**

All of the mice in the lesion study underwent two surgeries. In the first surgery, a stainless steel

headplate was attached to the skull to facilitate behavioral training. After collecting baseline

behavioral data, a second surgery consisting of either an excitotoxic or sham lesion was performed.

Before each surgery, the animal was treated pre-operatively with carprofen (5 mg/kg, i.p.; 024751,

Butler Animal Health) and dexamethasone (3 mg/kg, i.p.; Dexaject SP, #002459, Henry Schein

Animal Health). At the start of each surgery, anesthesia was induced with 2% isoflurane in oxygen,

and the animal was placed on a water-circulating heating pad (TP-700, Gaymar Stryker). The head

was secured in a stereotaxic frame with ear bars (David Kopf Instruments). Following induction,

isoflurane concentration was lowered to 1 – 1.5% based on the animal's weight and breathing

pattern.

For the first surgery, the scalp was shaved using scissors and cleaned with povidone-iodine

(Betadine, Perdue Products L.P.). A narrow portion of the scalp was removed along the midline

from the interaural line to a line visualized just posterior to the eyes. The scalp was retracted to

expose the dorsal aspect of the skull and washed thoroughly with artificial cerebrospinal fluid

(ACSF; in mM: 5 KCl, 5 HEPES, 135 NaCl, 1 MgCl2, and 1.8 CaCl2; pH 7.3). A scalpel and a

ballpoint pen were used to scratch and paint marks onto the skull at the secondary motor and

anterior cingulate cortices (MOs/ACAd; +1.5 mm AP, +0.3 mm ML from bregma), to be used as a

landmark for the second surgery. A custom-made stainless-steel head plate (eMachineShop) was

then bonded to the skull with cyanoacrylate glue (Loctite 454, Henkel) and transparent dental

acrylic (C&B Metabond, Parkell Inc.), with care taken to cover any remaining exposed skull. The

post-operative care was provided immediately, and for three consecutive days following surgery,

consisting of carprofen (5 mg/kg, i.p.) for analgesia and preservative-free 0.9% NaCl (0.5 mL, i.p.)

for fluid support. The animal had at least one week for post-operative recovery prior to the onset of

behavioral training.

For the second surgery, a 1-mm-diameter circular craniotomy was made over the marked spot using a high-speed rotary drill (K.1070, Foredom). A total of ~300 nL of ibotenic acid (5 mg/mL in saline; 505024, Abcam) was injected into two locations (+1.5 mm and +1.7mm AP, +0.3 mm ML from bregma; 0.4 mm DV) through a glass micropipette attached to a microinjection unit (Nanoject II, Drummond). More specifically, each location would receive 15 pulses of 9.6 nL of the prepared solution. To minimize backflow of the injected solution, there was a 1 min gap between each pulse, and the micropipette was left in place for 20 min after the last pulse. Sham animals underwent the same surgical procedure, but saline was delivered instead of ibotenic acid. The exposed skull was covered with dental cement. The animal had two weeks of post-operative recovery prior to resuming behavioral testing.

**Surgery for photostimulation experiments**

All of the mice in the photostimulation study underwent one surgery. The animal was anesthetized in the same way as described above. Procedures to prepare the skull were nearly identical to those described in (Pinto et al. 2019). Briefly, the scalp covering the dorsal skull surface was excised and the periosteum over the skull was removed using a micro-curette (VWR Buck Micro Curette, 10806-346). The skull was washed thoroughly with ACSF. A custom stainless-steel head-plate (eMachineShop) was affixed at points above the cerebellum and olfactory bulbs with cyanoacrylate glue (Loctite 454, Henkel) and transparent dental acrylic (C&B Metabond, Parkell Inc.). The exposed skull was covered with a thin layer of cyanoacrylate glue (Apollo 2000, Cyberbond) and transparent dental acrylic, then polished with an acrylic polishing kit (0321, PearsonDental), and finally covered with transparent nail polish (72180, Electron Microscopy Services). The animal had at least one week of post-operative recovery prior to the onset of behavioral training.

**Behavioral training apparatus**

The apparatus for training head-fixed mice was adapted from (Siniscalchi et al. 2016). Detailed plans including parts list for constructing the behavioral training apparatus is available at https://github.com/Kwan-Lab/behavioral-rigs. Briefly, the behavioral box was constructed using a closed compartment of an audio-visual cart (4731T74, McMaster-Carr) that was soundproofed with acoustic foam (5692T49, McMaster-Carr). The mouse was placed in an acrylic tube (8486K433, McMaster-Carr), which allowed for postural adjustments but restricted large body movements. Two metal screws were used to attach the head plate of the mouse onto a custom stainless-steel mount (eMachineShop). The lickometer was based on a 3D-printed part that held two lick ports

constructed from 20-gauge needles, and was placed in front of the mouse such that the lick ports
are on the left and right of the animal's mouth. The position of the lick ports relative to the mouse
could induce considerable side bias and influence response time. To mitigate variations across
sessions, the lickometer was attached to an XYZ translation stage (MT3, Thorlabs) for precise
positioning, and the same set of coordinates were used for the same mouse between sessions.
Water was supplied to the lick ports via Tygon tubing (EW- 95666-01, Cole-Parmer). A touch
detector circuit was used for detecting tongue licks onto each lick port. Water was delivered at the
lick ports by gravity feed and controlled by solenoid valves (EV-2-24; Clippard or MB202-V-A-3-0-L-
204, Gems Sensor Solenoid). The water amount is controlled by the duration of a TTL pulse, and
we calibrated the solenoid to deliver ~2 µL per pulse. All of the electrical circuits for water delivery
and lick detection were connected to a desktop computer via a data acquisition board (USB-201,
Measurement Computing). A pair of speakers (S120, Logitech) were positioned in front of the
animal for auditory stimuli (calibrated to 80 dB). The tasks were programmed in scripts using the
Presentation software (Neurobehavioral Systems), which controlled the entire behavioral apparatus
including stimulus presentation. A table lamp (LT-T6, Aukey) was placed in each box, behind the
mouse, to provide dim ambient light in the box. A camera (SV-USBFHD01M-BFV, Svpro) was used
to optimize the lick port position at the beginning of each session and monitor the animal's behavior
throughout the session.
567
568    **Two-armed bandit task**
569    Mice were fluid-restricted. On training days, animals received all of their water intake from
570    behavioral training that occurred 1 session per day, 5 days per week. On non-training days and
571    days when weight measurements fell below 85% of their pre-restriction weight, water was provided
572    ad libitum in the home cage for 5 minutes.
573
574    Prior to any behavioral training, the animal was handled and habituated to head fixation for
575    increasing durations over three days. Water was manually provided via the lick ports to familiarize
576    mice with receiving fluid from the lickometer. After 1 – 2 days of habituation, the animal underwent
577    two phases of shaping.
578
In the first phase, the animal was trained to alternate between the two lick ports to receive water
rewards. More specifically, on each trial, there would be an auditory cue (duration = 0.2 s, tone with
5 kHz carrier frequency). The onset of the auditory cue is the start of a 5-s long response window,
during which the first lick detected is the animal's response. The playback of the auditory cue was

terminated early if the response was recorded before the entire stimulus was played. The animal

was required to alternate between left and right responses to earn water rewards: if the last

rewarded response was left, then the mouse must make a right response to receive water, and vice

versa. The inter-trial interval had a fixed duration, such that the auditory cue for the next trial would

occur 3.1 s after the animal's response. A session would end when the animal did not lick during

the response window ('miss') for 20 consecutive trials. When the animal could attain at least ~60

rewards in a session, the shaping would proceed to the second phase.

In the second phase, the animal still had to alternate, but was trained to the trial structure including

withholding licks between trials. The second phase was similar to the first phase, with two

exceptions. First, the onset of the auditory cue is the start of a 2-s long response window, during

which the first lick detected is the animal's response. Second, the addition of a no-lick period

between trials. The no-lick period began 3 s after the animal's response. Initially, the duration of the

no-lick period was drawn from a truncated exponential distribution ($\lambda$ = 0.33333, minimum = 1,

maximum = 5). If any lick was detected during the no-lick period, then another duration drawn from

the same truncated exponential distribution would be added onto the end of the first no-lick period.

The addition could repeat for up to 5 times if the animal could not withhold licking. Therefore, the

possible duration for the entire no-lick period ranged between 1 and 25 s, and was dependent on

whether the animal could withhold licking. Subsequently, the auditory cue for the next trial would

occur 0.1 s after the end of the no-lick period. When the animal could receive rewards in at least

~40% of all trials, it would be advanced to the two-armed bandit task. Typically, the animal would

proceed through each shaping phase in 3 or fewer sessions.

In the two-armed bandit task, the auditory stimulus, response timing, and inter-trial interval

including no-lick period were exactly the same as the second shaping phase. However, the

outcome of each trial was probabilistically determined. In a 10:70 block of trials, the left lick port had

a 10% chance of delivering water if chosen and the right lick port had a 70% chance of delivering

water if chosen. By contrast, in a 70:10 block of trials, the reward probabilities associated with the

left and right ports were reversed. Hence, the better option is right in a 10:70 block, but left in a

70:10 block. At the start of each session, the block type (10:70 or 70:10) was randomly chosen.

The block type would switch when the mouse fulfilled the switching condition: perform trials

($L_{Criterion}$) until the animal accumulated 10 choices selecting the side with high reward probability,

and then perform additional trials ($L_{Random}$) that were drawn from a truncated geometric distribution

($p$ = 0.0909, no minimum = 0, maximum = 30). Notably, $L_{Criterion}$ depended on the animal's

performance, whereas $L_{Random}$ was random and independent of performance. The block type would
continue to switch, as long as the animal was fulfilling the switching condition of each block. In the
lesion experiments, they would be tested on the two-armed bandit task in daily sessions until at
least 150 switches were collected for each of the pre- and post-lesion conditions.
621
**Photostimulation**
The photostimulation rig allowed for rapid adjustment of the position of the laser. The rig was
constructed based on the design in (Pinto et al. 2019). Briefly, a 473 nm laser beam (Obis LX 473
nm, 75 mW; 1193830, Coherent) was steered by a set of XY galvo mirrors (6210H, Cambridge
Technologies) mounted in a ThorLabs 60 mm cage system. The laser was sent through a F-theta
scan lens (f = 160 mm; FTH160-1064-M39, ThorLabs) and directed onto the animal's head. A
monochromatic camera (Grasshopper3; GS3-U3-23S6M-C, Point Grey) equipped with a telecentric
lens (TEC-55, Computar) was used to visualize the cortical surface and to calibrate the position of
the laser beam relative to bregma. The laser, mirrors, and camera were controlled via a data
acquisition board (PCIe-6343, National Instruments) by custom software written in MATLAB on a
desktop computer. The laser was calibrated to yield a time-averaged power of 1.5 mW at the
sample. Light transmission through the clear-skull cap (dental cement and skull) was measured by
placing the cap at the sample plane, and positioning a laser power meter underneath the cap.
635
Animals underwent the same shaping phases and task training. For the photostimulation
experiments, the animal was tested on the two-armed bandit task in a behavioral setup within the
photostimulation rig. Temporally, the photostimulation could occur either before or after the animal's
response. For pre-choice photostimulation, the laser was turned on at the onset of the auditory cue
and turned off immediately when a response was detected. For post-choice photostimulation, the
laser was turned on immediately when a response was detected and turned off 2 s later. Spatially,
the photostimulation was targeted to one of two possible locations: left MOs/ACAd (+1.5 mm AP, -
0.3 mm ML from bregma) or right MOs/ACAd (+1.5 mm AP, +0.3 mm ML from bregma).
644
At the start of each session, the timing of the photostimulation (pre-choice or post-choice) was
randomly chosen and stayed the same for the entire session. The initial 3 – 5 blocks were always
control blocks, i.e., no photostimulation. The rationale was to make sure the animal was performing
the task well that day before any perturbation. Subsequently, the next 2 blocks would be
photostimulation blocks targeting the same spatial location, followed by 2 control blocks, followed
by 2 photostimulation blocks targeting the same spatial location, and so on. For the

photostimulation blocks, the spatial location was randomly selected to be left or right MOs/ACAd

each time. In other words, in the same session, the animal may receive perturbation of both left and

right MOs/ACAd, albeit in different trial blocks.

To prevent the animal from using stray laser light to distinguish photostimulation from control

blocks, we implemented a masking stimulus by shining a blue LED at the eyes. The masking

stimulus had the same onset timing and duration as the photostimulation used for the session, and

was applied for every trial in both control and photostimulation blocks.

**Histology**

To determine the extent of the lesions, following behavioral experiments, the mouse was deeply

anaesthetized with an overdose of isoflurane and transcardially perfused with chilled formaldehyde

solution (4%, in phosphate-buffered saline (PBS)) at a rate of 5 mL/min. The brain was quickly

removed, stored overnight in the formaldehyde solution at 4 °C, and then switched to PBS for long-

term storage. Coronal sections with a thickness of 100 µm were cut using a vibratome (VT1000 S,

Leica).

For cresyl violet staining, cresyl violet (1 g/L; 10510-54-0, Sigma Aldrich) was added to filtered $H_2O$

and stirred overnight. The next day, glacial acetic acid (2.5 mL/L; 64-19-7, Sigma Aldrich) was

added to the solution. The tissue sections were washed with filtered $H_2O$ before mounting on glass

slides and stained with a pre-warmed (50°C) cresyl violet solution. The sections were dehydrated

with ascending grades of alcohol (95% for 10 minutes, 100% twice for 10 minutes each), cleared

with xylene (twice for 5 minutes each), and mounted with DPX mounting medium (06522,

MilliporeSigma).

For NeuN staining, tissue sections were washed three times with PBS and then incubated with a

blocking solution (5% normal goat serum, 0.3% Triton X-100, in PBS) for 1 hour at room

temperature. Subsequently, sections were incubated with rabbit monoclonal primary antibody

against NeuN (1:500 dilution; ab177487, Abcam Inc,) overnight at 4 °C on the shaker. After

washing three times with PBS, tissue sections were incubated with goat anti-rabbit secondary

antibody with conjugated Alexa 488 (1:500 dilution; ab150077, Abcam Inc,) for 2.5 hours at room

temperature. After washing with PBS, nuclear staining was performed by incubating with a 4',6-

diamidino-2-phenylindole (DAPI) staining solution (ab228549, Abcam Inc.) for 10 minutes. Finally,

sections were washed three times with PBS and then with filtered $H_2O$, before mounting on slides

with DPX mounting medium (06522, MilliporeSigma). A motorized upright fluorescence microscope

(Olympus BX61, Olympus) was used to image the sections.

**QUANTIFICATION AND STATISTICAL ANALYSIS**

**Analysis of behavioral data**

Timestamps of the behavioral events, including cue onsets, outcome onsets, licks, and reward

probabilities were logged to a text file by the NBS Presentation software. The text files were parsed

and analyzed using scripts written in MATLAB (MathWorks, Inc.). For all of the analyses, we

excluded the session if the animal had fewer than 4 block switches. We analyzed all of the trials up

to the last switch, and ignored the trials in the last incomplete block where by definition had many

miss trials.

When analyzing the consequences of unilateral lesions, for simplicity, we used the term _lesion_

_blocks_ and _contra blocks_. This is because unilateral lesions were randomly assigned to the left or

right hemisphere for each animal. Lesion blocks refer to those blocks where the lesioned side is the

same as the better option. In other words, if the animal had a unilateral lesion on the right

hemisphere, then the lesion blocks correspond to 10:70 blocks. If the animal had a unilateral lesion

on the left hemisphere, then the lesion blocks correspond to the 70:10 blocks. The remainder was

referred to as the contra blocks.

**Analysis of behavioral data – effects of block length**

For analyses involving block lengths, we used the subset of data in which $L_{Criterion} \leq 20$ trials for the

pre-switch block, in order to restrict the analyses to situations where the performance was similar.

The probability of choosing the better option pre-switch, $P$ (_better option_) $_{pre-switch}$, was determined

for each animal by examining the last five trials before each block switch, dividing the number of

times in which the animal chose the initial better option (i.e., the side with 70% reward probability

before switch) by the number of switches. _Hit rate_ was the proportion of trials in which the animal

selected the better option. The win-stay probability, $P$ (_stay | win_), was the fraction of trials in which

animals repeated a choice after a rewarded trial. The lose-switch probability, $P$ (_switch | lose_), was

the fraction of trials in which an animal switched its choice after an unrewarded trial. For all of these

performance metrics, we computed the metric on a session-by-session basis, then averaged

across sessions to obtain per-animal value. In the lesion data, $P$ (_better option_) $_{pre-switch}$, $P$ (_stay |_

_win_) and $P$ (_switch | lose_) was calculated in the last five trials before each block switch

*Trials to reach midpoint* was determined by first calculating the fraction of trials for choosing the

initial better option around the block switch for the animal, and then identifying the trial from the

switch where the fraction of trials choosing the initial better option was closest to 0.5. To compute

the trials to reach the midpoint metric, we would first concatenate data across sessions including

inserting 20 NaN in the gaps, then compute the metric to obtain the per-animal value. We did this

because the per-session values for trials to reach midpoint can be quite variable.

**Analysis of behavioral data – reinforcement learning models**

The response-by-response behavior of the animal was fitted with eight models: (1) win-stay lose-

switch (WSLS); (2) Q-learning (Q-RPE); (3) Q-learning with forgetting (F-Q-RPE); (4) Q-learning

with differential forgetting (DF-Q-RPE); (5) F-Q-RPE with choice kernel (F-Q-RPE-CK), which

captured the tendency to repeat the same option; (6) DF-Q-RPE with choice kernel (DF-Q-RPE-

CK); (7) belief model that uses the prior knowledge of a change point in reward probabilities to

make a decision (7) belief model with choice kernel (belief-CK). We will describe these models in

detail in the following paragraphs.

For *win-stay lose-switch (WSLS)*, if the last trial was rewarded, the agent would repeat to choose

the same option with probability $p$. Else, if the last trial was unrewarded, the agent would switch to

choose the other option with probability $p$. This model has 1 free parameter: $p$.

For the three *simple Q-learning models (Q-RPE, F-Q-RPE, DF-Q-RPE)*, the updating rules are as

follows. On trial $n$, for a choice $c_n$ that leads to an outcome $r_n$, the reward prediction error $\delta_n$ is:

$$\delta_n = r_n - Q_n^i \qquad\qquad\qquad (1)$$

where $Q_n^i$ is the action value associated with the chosen action $i$. In our task, there are two options,

so $i \in \{L, R\}$. For the outcome, $r_n$ = 1 for reward, 0 for no reward. The action value for each action

is then updated accordingly:

$$Q_{n+1}^i = \begin{cases} Q_n^i + \alpha\,\delta_n & \text{if } c_n = i \\ (1-\lambda)Q_n^i & \text{if } c_n \neq i \end{cases} \qquad\qquad (2)$$

where $\alpha$ is the learning rate, $\lambda$ are the forgetting terms for the unchosen action. Then on the next

trial, the probability of choosing each action was determined by a softmax rule:

$$P(c_{n+1} = i) = \frac{exp\ (\beta Q_{n+1}^i)}{\sum_i exp\ (\beta Q_{n+1}^i)} \tag{3}$$

where $\beta$ is the inverse temperature parameter.

The model as stated with 3 free parameters — $\alpha$, $\lambda$, and $\beta$ — is referred to as Q-learning with

differential forgetting (DF-Q-RPE). A special case of this model is when $\lambda = \alpha$, which is referred to

as Q-learning with forgetting (F-Q-RPE). Another special case is when $\lambda = 0$, which is referred to

as Q-learning (Q-RPE).

For two Q-learning models with choice kernel (*F-Q-RPE-CK, DF-Q-RPE-CK)*, choice kernel was

implemented to capture the tendency of choosing the previous choice. We adapted the formulation

from (Wilson and Collins 2019). The choice kernel $K_t^i$ on trial $t$ associated with action $i$ is updated

in a manner analogous to the action values:

$$K_{n+1}^i = \begin{cases} K_n^i + \alpha_K(1 - K_n^i) & \text{if } c_n = i \\ (1 - \alpha_K)K_n^i & \text{if } c_n \neq i \end{cases} \tag{4}$$

769    where $\alpha_K$ is the choice-kernel learning rate. For action selection with both action values and choice

770    kernels, the probability of choosing each action was determined by a softmax rule:

$$P(c_{n+1} = i) = \frac{exp\ (\beta Q_{n+1}^i + \beta_K K_{n+1}^i)}{\sum_i exp\ (\beta Q_{n+1}^i + \beta_K K_{n+1}^i)} \tag{5}$$

774    where $\beta$ and $\beta_K$ are the inverse temperature parameters for the action values and choice kernels

775    respectively. Note that the term within the numerator on the right-hand side can be re-arranged:

$$\beta Q_{n+1}^i + \beta_K K_{n+1}^i = \left(\frac{\beta}{\beta + \beta_K} Q_{n+1}^i + \frac{\beta_K}{\beta + \beta_K} K_{n+1}^i\right) \tag{6}$$

779    Where $(\beta + \beta_K)$ is the effective inverse temperature parameter reflecting the exploration-

780    exploitation balance, and $\frac{\beta}{\beta + \beta_K}$ is a ratio indicating the relative reliance on expected reward rather

781    than perseveration in action selection.

**Analysis of behavioral data – belief models**

In the <u>belief model</u>, the agent knows two aspects about the task structure. First, the left option can have reward probabilities of either 10% or 70%. It follows that the right option would have the other reward probability. These are the two possible hidden states of the environment. Second, the reward probabilities will reverse with a certain frequency characterized by a hazard rate, $H$. In each trial, the animal has a belief, which consists of the likelihood that the left option has a reward probability of 10%, $\rho_{L10}$, and the likelihood that the left option has a reward probability of 70%, $\rho_{L70}$. The constraints are that $\rho_{L10} + \rho_{L70} = 1$, $\rho_{R10} = 1 - \rho_{L10}$ and $\rho_{R70} = 1 - \rho_{L10}$. At the start of a session, we set the prior as a uniform distribution, so the $\rho_{L10} = \rho_{L70} = \rho_{prior} = 0.5$. At the end of each trial, the belief is updated. The possibility of a reward probability switch is considered:

$$\rho_{L10} = \rho_{L10} * (1 - H) + \rho_{prior} * H \qquad (7)$$

Similarly, $p_{L70}$ is updated and then $p_{L10}$ and $p_{L70}$ are normalized to sum to 1. Next, inference is made based on the outcome following Bayes' rule, which states that P (belief | observation) = P(belief) * P (observation | belief):

$$\rho_{L10} = \rho_{L10} * 0.1 \qquad \text{if left and rewarded} \quad (8)$$
$$\rho_{L70} = \rho_{L70} * 0.7 \qquad \text{if left and rewarded}$$

$$\rho_{L10} = \rho_{L10} * (1 - 0.1) \qquad \text{if left and unrewarded}$$
$$\rho_{L70} = \rho_{L70} * (1 - 0.7) \qquad \text{if left and unrewarded}$$

or if the animal chooses right, then $\rho_{R10}$ and $\rho_{R70}$ would be updated instead. Again, the probabilities for the belief are normalized to sum to 1. With the updated belief, the expected rewards for the left and right options can be calculated directly, for example:

$$\mu_L = \rho_{L10} * 0.1 + \rho_{L70} * 0.7 \qquad (9)$$

Action selection then proceeds using the same softmax equation as Equations 3 or 5, with the expected rewards replacing the action value terms, for the belief and belief with choice kernels models respectively. The belief model has two free parameters, $H$ and $\beta$. The belief model with choice kernel model has four free parameters $H$, $\beta$, $\alpha_K$, and $\beta_K$.

**Parameter fitting and model evaluation**

For each animal, trials across sessions were concatenated. The values for the free parameters

were determined by fitting each model to the concatenated data using the Bayesian adaptive direct

search (BADS) algorithm with default settings (Acerbi and Ma 2017). The initial values for $\alpha$, $\lambda$, $H$,

$\alpha_K$, $\beta$ and $\beta_K$ were set to 0.3, 0.3, 0.1, 0.2, 5 and 5 respectively. The lower bound of parameters

were set to 0 and the upper bound was set to 100 for inverse temperatures and 1 for the rest of the

parameters. To evaluate the models, we calculated the Bayes information criterion (BIC).

825
$$BIC = N_m \log T + 2LL \qquad\qquad (10)$$

where $N_m$ is the number of parameters in model m. T is the number of trials used to estimate the

parameters and LL is the negative log-likelihood value at the best fitting parameter settings. The

model that best fits the data should have the smallest BIC score as the positive effect of the

number of parameters, $N_m$, has an explicit penalty for free parameters.

The parameters used for the simulated data for the belief model with choice kernel was the best-

fitting parameters of one animal ($H$ = 0.320, $\beta$ = 1.387, $\alpha_K$=0.468, $\beta_K$ = 2.543 with 300,000 trials,

approximately 9000 switches as in the experiment data). The belief model with choice kernel was

used to analyze the latent variables for lesion and photostimulation data.

To fit the lesion data, we modified the belief-CK model. Different parameters for hazard rate and

choice kernel learning rate were used depending on if the animal's choice in the current trial is

ipsilateral or contralateral to the lesion side. This yields an expanded model with 6 parameters:

$H_{lesion}$, $H_{contra}$, $\alpha_{K\ lesion}$, $\alpha_{K\ control}$, $\beta$, and $\beta_K$. To fit the data on a per-animal basis, trials across

sessions before lesion were concatenated, and trials across sessions after lesion were

concatenated. To fit the data on a per-session basis, we estimated the parameters for each

session.

To fit the optogenetics data, we modified the belief-CK model. Different parameters for hazard rate

and choice kernel learning rate were used depending on if the animal's choice in the current trial is

ipsilateral or contralateral to the photostimulated side, or if the animal's choice occurred in a control

trial with no photostimulation. Different parameters for inverse temperatures were used depending

on if the current trial was photostimulation or control. This yields an expanded model with 10

parameters: $H_{ipsi\ stim}$, $H_{contra\ stim}$, $H_{control}$, $\alpha_{K\ ipsi\ stim}$, $\alpha_{K\ contra\ stim}$, $\alpha_{K\ control}$, $\beta_{control}$, $\beta_{stim}$,

$\beta_{K\ control}$, and $\beta_{K\ stim}$. Per each animal fittings, trials across sessions for pre-choice stimulation and

post-choice stimulation were concatenated. Per session fittings was not used for this dataset as

total number of trials within a session did not give a reliable estimate for the ten parameters fittings.

**Statistical Analyses**

All statistical analyses were completed using MATLAB (version 2019b, MathWorks). Three-way

ANOVA was used to examine the effect of the lesion on behavioral performance. For datasets with

a matched number of data points, Wilcoxon signed-rank test was used; otherwise, Wilcoxon ranked

sum test was used. Unless otherwise specified, we used an alpha level of 0.05 for all statistical

tests.

**Acknowledgments**

We thank Neil Savalia and John-Anthony Fraga for assistance with behavioral training, Michael

Siniscalchi for use of help with the behavioral apparatus, Farhan Ali for advice on histology,

Stephan Thiberge and David Tank for sharing design of photostimulation rig, and Alireza Soltani for

comments on an earlier version of the analysis. This work was supported by NIH/NIMH grants

R01MH112750 (A.C.K.), R21MH118596 (A.C.K.), F32NS101871 (L.P.), K99MH120047 (L.P.),

China Scholarship Council-Yale World Scholars Fellowship (H.W.), Gruber Science Fellowship

(H.K.O.), NIH training grant T32NS007224 (H.K.O.), and Kavli Institute for Neuroscience

Postdoctoral Fellowship (H.A.).

**Author Contributions**

H.A. and A.C.K. conceived the project. H.A. performed all surgeries. H.A. and C.E.M performed

behavioral training including lesion and optogenetic inactivation experiments, and histology. H.K.O.

performed the ALM inactivation experiment. H.W., H.A., and A.C.K. put together the optogenetic

activation rig, based on L.P.'s plans. H.A. and A.C.K. analyzed the data and wrote the paper with

input from all other authors.

**Declaration of Interests**

The authors declare no competing financial interests.

 **Supplementary Information**

| | df | P(better option) pre-switch | | Trials to reach midpoint | | Hit rates | |
|---|---|---|---|---|---|---|---|
| | | F | p | F | p | F | p |
| *Lesion* | *1* | **32.71** | **.000** | **33.36** | **.000** | **4.87** | **.027** |
| *Side* | *1* | 0.63 | .857 | **8.68** | **.003** | **4.02** | **.045** |
| $L_{Random}$ | *3* | **57.67** | **.000** | 0.52 | .666 | **21.46** | **.000** |
| *Lesion*Side* | *1* | *1.42* | .233 | **4.30** | **.038** | 0.86 | .355 |
| *Lesion*$L_{Random}$* | *3* | **2.71** | **.044** | 0.57 | .635 | **3.06** | **.027** |
| $L_{Random}$*Side* | *3* | *0.18* | .910 | **4.30** | **.004** | 0.49 | .688 |
| *Lesion*$L_{Random}$*Side* | *3* | *2.40* | .066 | 1.60 | .187 | 1.07 | .361 |

884　**Supplementary Table 2.1**: The results of three-way between-subjects ANOVA with factors of

885　lesion (pre- and post-lesion), side (lesion blocks and Contra blocks), and $L_{Random}$ (4 $L_{Random}$ ranges)

886　for P(better option)$_{pre-switch}$, trials to reach midpoint and hit rates. $p < 0.05$ in bold. All dependent

887　variables calculated for each block across sessions. (Error = 3285; 2399; 3285; 3285; 2816;3187

888　for P (better option) $_{pre-switch}$, trials to reach midpoint and hit rates respectively

**Supplementary Figure 2.1: No decrease in overall performance after unilateral lesion of ACAd/MOs**

The total number of trials and block switches per session before (pre) and after (post) the unilateral lesion.

896 **Supplementary Figure 2.2: No motor deficits after unilateral lesion of ACAd/MOs**

897 Mean left and right lick density for each possible combination for choice (left or right) and outcome

898 (reward or no reward). No significant difference was detected between pre- and post-unilateral

899 lesion.

# Hazard rate simulation

n = 300,000 trials; $\alpha_K$ = 0.4675; $\beta$= 2.3872; $\beta_K$ = 2.5426;
All parameters fixed except H:

**Supplementary Figure 3.1: Belief-CK model: effect of varying the hazard rate**

The belief-CK model was used to simulate an agent's choice behavior in the two-armed bandit task

with probabilistic reward reversal. Parameters were selected based on the best fitting values from

an animal. Each column shows the results using a different hazard rate (   = 0.01, 0.25, 0.5, 0.75,

1) while all other parameters were kept constant (n = 300,000 trials,   = 1.387,   =0.468,   =

2.543). Top row shows the mean fraction of trials choosing the better and worse options for 4

different $L_{Random}$ ranges for 10 trials before and after the block switch. Middle row shows the P

908  (better option) $_{\text{pre-switch}}$ as a function of $L_{\text{Random}}$. Mean and SEM. Bottom row shows the mean number

909  of trials to reach midpoint as a function of $L_{\text{Random}}$.

**Choice kernel learning rate simulation**

n = 300,000 trials; H = 0.3201; β= 2.3872; $β_K$ = 2.5426;

All parameters fixed except $α_K$:

912 **Supplementary Figure 3.2: Belief-CK model: effect of varying choice kernel learning rate.**

913 Similar to Supplementary Figure 3.1, with different choice kernel learning rates ( = 0.01, 0.25,

914 0.5, 0.75, 1) while all other parameters were kept constant (n = 300,000 trials, = 0.320, =

915 1.387, = 2.543).

**Beta sum simulation**

n = 300,000 trials; H = 0.3201; $\alpha_K$ = 0.4675;
All parameters fixed except $\beta$ = $\beta_K$ = ($\beta$ + $\beta_K$)/2

**Supplementary Figure 3.3: Belief-CK model: effect of varying beta sum**.

Similar to Supplementary Figure 3.1, with different beta sum (          = 0, 1, 3, 5, 10) while all other

parameters were kept constant (n = 300,000 trials,      = 0.320,      = 0.468).      was set to equal to    .

**Supplementary Figure 3.4: Belief-CK model: effect of varying beta ratio.**

Similar to Supplementary Figure 3.1, with different beta ratios (0.01, 0.25, 0.5, 0.75, 1) while all
other parameters were kept constant (n = 300,000 trials,  = 0.320,  =0.468,  = 2.543).  was
fixed and  was calculated based on the beta ratio values.

**Supplementary Figure 3.5: DF-Q-RPE algorithm cannot reproduce the $L_{Random}$-dependent**

**trends in the experimental data.**

**(A)** The probability of choosing the better option on the trial immediately preceding the switch, as a

function of $L_{Random}$ for the block preceding the switch. Black, mice. Purple, simulated performance

using the DF-Q-RPE model with best-fitting parameters. Mean values and SEM for all animals. **(B)**

Similar to (A) for number of trials to reach midpoint (when animal is equally likely to choose either

option). **(C)** Similar to (A) for the tendency to win-stay on the 5 trials preceding the switch. **(D)**

Similar to (A) for the tendency to lose-switch on the 5 trials preceding the switch. Mean and SEM. n

= 31 mice, 617 sessions.

**Supplementary Figure 5.1: Fewer trials but similar performance after bilateral lesion of ACAd/MOs**

The total number of left- and right-responding trials, reward rates, and hit rates before (pre) and after (post) the lesion.

**Supplementary Figure 5.2: No motor deficits after bilateral lesion of ACAd/MOs**

Mean left and right lick density for each possible combination for choice (left or right) and outcome

(reward or no reward). No significant difference was detected between pre- and post-bilateral

lesion.

|  | df | P(better option) pre-switch | | Trials to reach midpoint | | Hit rates | |
|---|---|---|---|---|---|---|---|
|  |  | F | p | F | p | F | p |
| *Lesion* | *1* | 1.968 | *.161* | 0.001 | .974 | 1.0814 | 0.2986 |
| *L<sub>Random</sub>* | *3* | 20.179 | *.000* | 1.525 | .206 | 1.8643 | 0.1337 |
| *Lesion*L<sub>Random</sub>* | *3* | *1.285* | *.278* | 0.202 | .895 | 0.4492 | 0.7179 |

950 **Supplementary Table 5.1**: The results of two-way between-subjects ANOVA for bilaterally injected

951 animals with factors of lesion (pre- and post-lesion and $L_{Random}$ (4 $L_{Random}$ ranges) for P(better

952 option) $_{pre-switch}$, trials to reach midpoint and hit rates. $p < 0.05$ in bold. All dependent variables

953 calculated for each block across sessions. (Error = 1356;911;1356; for P (better option) $_{pre-switch}$,

954 trials to reach midpoint and hit rates respectively

|  | df | P(better option) pre-switch | | Trials to reach midpoint | | Hit rates | |
|---|---|---|---|---|---|---|---|
|  |  | F | p | F | p | F | p |
| *Lesion* | *1* | 0.0167 | *.897* | 9.7782 | *.002* | 0.0325 | 0.857 |
| *L<sub>Random</sub>* | *3* | 20.430 | *.000* | 1.0813 | *.356* | 9.624 | 0.000 |
| *Lesion*L<sub>Random</sub>* | *3* | *0.082* | *0.969* | *0.0594* | *.981* | 0.190 | 0.903 |

**Supplementary Table 5.2**: The results of two-way between-subjects ANOVA for saline injected

animals with factors of lesion (pre- and post-lesion and $L_{Random}$ (4 $L_{Random}$ ranges) for P (better

option) $_{pre-switch}$, trials to reach midpoint, hit rates, P(lose | switch), P(win | stay). $p < 0.05$ in bold. All

dependent variables calculated for each block across sessions. (Error = 1871;1217; 1871 for P

(better option) $_{pre-switch}$, trials to reach midpoint and hit rates respectively.

**Supplementary Figure 6.1: Validation of the laser steering system for optogenetic**

**manipulation: characterization and c-Fos staining**

**(A)** Optical transmission of the clear skull cap preparation was measured by illuminating with a

laser and recording intensity using a power meter. Mean and SEM. n = 5. **(B)** Linearity of the

galvanometers in the x and y directions. **(C)** Beam profile was measured at the sample plane by

inserting and moving a razor blade across the plane using a micromanipulator. **(D - F)** In

*CaMKIIa$^{Cre}$;Ai32* animals, cortical excitatory neurons express ChR2. After unilateral

photostimulation of the left ACAd/MOs region (40 Hz, 1.5 mW, 1 min on then 1 min off repeatedly

for 20 min), immunohistostaining with a c-Fos antibody showed elevated signals.

**Supplementary Figure 6.2: Inactivating left and right ALM during two-armed bandit task**

**(A)** In *Pvalb<sup>Cre</sup>;Ai32* animals, parvalbumin-expressing neurons including fast-spiking interneurons in

the neocortex express ChR2. Photostimulation of a brain region drives spiking in the interneurons,

which in turn suppresses excitatory activity. Lick raster recorded in an example session, in which

trials were sorted based on the photostimulation (None: no stimulation; ALM-L: left anterior lateral

motor cortex, AP=2.5 mm, ML=-1.5 mm; ALM-R: right anterior lateral motor cortex, AP=2.5 mm,

ML=1.5 mm; V1-L: left primary visual cortex, AP=-2.7 mm, ML=-2.5 mm; V1-R: right primary visual

cortex, AP=-2.7 mm, ML=2.5 mm). **(B)** The number of trials of each type per session. **(C)** Percent

of trials resulted in a miss, as a function of trial type. **(D)** Percent of trials resulted in a left response,

as a function of trial type. **(E)** Percent of trials resulted in a right response, as a function of trial type.

These results show that transient inactivation of ALM increased ipsilateral responses at the

expense of contralateral responses. 9 sessions from 3 animals.

**Supplementary Figure 6.3: Pre-choice inactivation, but not post-choice inactivation, significantly affected the change point estimation. (A)** Schematic of stimulation scheme. **(B)** The probability of choosing the better option on the trial immediately preceding the switch, as a function of $L_{Random}$ for the block preceding the switch, for pre-choice inactivation. Mean values and SEM for all animals (n = 6 mice). **(C)** Similar to (B) for post-choice inactivation.

**Reference**

Acerbi, Luigi, and Wei Ji Ma. 2017. "Practical Bayesian Optimization for Model Fitting with Bayesian
Adaptive Direct Search." *ArXiv Preprint ArXiv:1705.04405*.

Bari, Andrea, David E. Theobald, Daniele Caprioli, Adam C. Mar, Alex Aidoo-Micah, Jeffrey W.
Dalley, and Trevor W. Robbins. 2010. "Serotonin Modulates Sensitivity to Reward and
Negative Feedback in a Probabilistic Reversal Learning Task in Rats."
*Neuropsychopharmacology* 35(6):1290–1301. doi: 10.1038/npp.2009.233.

Bari, Bilal A., Cooper D. Grossman, Emily E. Lubin, Adithya E. Rajagopalan, Jianna I. Cressy, and
Jeremiah Y. Cohen. 2019. "Stable Representations of Decision Variables for Flexible
Behavior." *Neuron* 103(5):922-933.e7. doi: 10.1016/j.neuron.2019.06.001.

Barthas, Florent, and Alex C. Kwan. 2017. "Secondary Motor Cortex: Where 'Sensory'Meets
'Motor'in the Rodent Frontal Cortex." *Trends in Neurosciences* 40(3):181–93.

Bartolo, Ramon, and Bruno B. Averbeck. 2020. "Prefrontal Cortex Predicts State Switches during
Reversal Learning." *Neuron* 106(6):1044-1054.e4. doi: 10.1016/j.neuron.2020.03.024.

Behrens, Timothy E. J., Mark W. Woolrich, Mark E. Walton, and Matthew F. S. Rushworth. 2007.
"Learning the Value of Information in an Uncertain World." *Nature Neuroscience*
10(9):1214–21. doi: 10.1038/nn1954.

Bennur, Sharath, and Joshua I. Gold. 2011. "Distinct Representations of a Perceptual Decision and
the Associated Oculomotor Plan in the Monkey Lateral Intraparietal Area." *Journal of
Neuroscience* 31(3):913–21.

Clarke, Hannah F., Trevor W. Robbins, and Angela C. Roberts. 2008. "Lesions of the Medial
Striatum in Monkeys Produce Perseverative Impairments during Reversal Learning Similar
to Those Produced by Lesions of the Orbitofrontal Cortex." *Journal of Neuroscience*
28(43):10972–82.

Costa, Vincent D., Valery L. Tran, Janita Turchi, and Bruno B. Averbeck. 2015. "Reversal Learning
and Dopamine: A Bayesian Perspective." *The Journal of Neuroscience* 35(6):2407. doi:
10.1523/JNEUROSCI.1989-14.2015.

Donahue, Christopher H., and Daeyeol Lee. 2015. "Dynamic Routing of Task-Relevant Signals for
Decision Making in Dorsolateral Prefrontal Cortex." *Nature Neuroscience* 18(2):295–301.
doi: 10.1038/nn.3918.

Evers, Elizabeth A. T., Roshan Cools, Luke Clark, Frederik M. van der Veen, Jelle Jolles, Barbara
J. Sahakian, and Trevor W. Robbins. 2005. "Serotonergic Modulation of Prefrontal Cortex
during Negative Feedback in Probabilistic Reversal Learning." *Neuropsychopharmacology*
30(6):1138–47. doi: 10.1038/sj.npp.1300663.

Gallistel, C. R., Terence A. Mark, Adam Philip King, and P. E. Latham. 2001. "The Rat
Approximates an Ideal Detector of Changes in Rates of Reward: Implications for the Law of
Effect." *Journal of Experimental Psychology: Animal Behavior Processes* 27(4):354.

Groman, Stephanie M., Colby Keistler, Alex J. Keip, Emma Hammarlund, Ralph J. DiLeone,
Christopher Pittenger, Daeyeol Lee, and Jane R. Taylor. 2019. "Orbitofrontal Circuits

Control Multiple Reinforcement-Learning Processes." *Neuron* 103(4):734-746.e3. doi: 10.1016/j.neuron.2019.05.042.

Groman, Stephanie M., Katherine M. Rich, Nathaniel J. Smith, Daeyeol Lee, and Jane R. Taylor. 2018. "Chronic Exposure to Methamphetamine Disrupts Reinforcement-Based Decision Making in Rats." *Neuropsychopharmacology* 43(4):770–80.

Grossman, Cooper D., Bilal A. Bari, and Jeremiah Y. Cohen. 2022. "Serotonin Neurons Modulate Learning Rate through Uncertainty." *Current Biology* 32(3):586-599. e7.

Guo, Zengcai V., Nuo Li, Daniel Huber, Eran Ophir, Diego Gutnisky, Jonathan T. Ting, Guoping Feng, and Karel Svoboda. 2014. "Flow of Cortical Activity Underlying a Tactile Decision in Mice." *Neuron* 81(1):179–94.

Hamid, Arif A., Jeffrey R. Pettibone, Omar S. Mabrouk, Vaughn L. Hetrick, Robert Schmidt, Caitlin M. Vander Weele, Robert T. Kennedy, Brandon J. Aragona, and Joshua D. Berke. 2016. "Mesolimbic Dopamine Signals the Value of Work." *Nature Neuroscience* 19(1):117–26. doi: 10.1038/nn.4173.

Hattori, Ryoma, Bethanny Danskin, Zeljana Babic, Nicole Mlynaryk, and Takaki Komiyama. 2019. "Area-Specificity and Plasticity of History-Dependent Value Coding During Learning." *Cell* 177(7):1858-1872.e15. doi: 10.1016/j.cell.2019.04.027.

Ito, Makoto, and Kenji Doya. 2009. "Validation of Decision-Making Models and Analysis of Decision Variables in the Rat Basal Ganglia." *Journal of Neuroscience* 29(31):9861–74.

Ito, Makoto, and Kenji Doya. 2015. "Distinct Neural Representation in the Dorsolateral, Dorsomedial, and Ventral Parts of the Striatum during Fixed- and Free-Choice Tasks." *The Journal of Neuroscience* 35(8):3499. doi: 10.1523/JNEUROSCI.1962-14.2015.

Ito, Makoto, and Kenji Doya. 2015. "Parallel Representation of Value-Based and Finite State-Based Strategies in the Ventral and Dorsal Striatum." *PLoS Computational Biology* 11(11):e1004540.

Jang, Anthony I., Matthew R. Nassar, Daniel G. Dillon, and Michael J. Frank. 2019. "Positive Reward Prediction Errors during Decision-Making Strengthen Memory Encoding." *Nature Human Behaviour* 3(7):719–32. doi: 10.1038/s41562-019-0597-3.

Karlsson, Mattias P., Dougal GR Tervo, and Alla Y. Karpova. 2012. "Network Resets in Medial Prefrontal Cortex Mark the Onset of Behavioral Uncertainty." *Science* 338(6103):135–39.

Laubach, Mark, Linda M. Amarante, Kyra Swanson, and Samantha R. White. 2018. "What, If Anything, Is Rodent Prefrontal Cortex?" *Eneuro* 5(5):ENEURO.0315-18.2018. doi: 10.1523/ENEURO.0315-18.2018.

Li, Nuo, Susu Chen, Zengcai V. Guo, Han Chen, Yan Huo, Hidehiko K. Inagaki, Guang Chen, Courtney Davis, David Hansel, and Caiying Guo. 2019. "Spatiotemporal Constraints on Optogenetic Inactivation in Cortical Circuits." *Elife* 8:e48622.

Li, Yi, and Joshua Tate Dudman. 2013. "Mice Infer Probabilistic Models for Timing." *Proceedings of the National Academy of Sciences* 110(42):17154–59.

Liao, Clara, and Alex C. Kwan. 2021. "Applying Reinforcement Learning to Rodent Stress Research." *Chronic Stress* 5:2470547020984732.

Liu, Yanhe, Yu Xin, and Ning-long Xu. 2021. "A Cortical Circuit Mechanism for Structural Knowledge-Based Flexible Sensorimotor Decision-Making." *Neuron*.

O'Doherty, J., M. L. Kringelbach, E. T. Rolls, J. Hornak, and C. Andrews. 2001. "Abstract Reward and Punishment Representations in the Human Orbitofrontal Cortex." *Nature Neuroscience* 4(1):95–102. doi: 10.1038/82959.

Otchy, Timothy M., Steffen BE Wolff, Juliana Y. Rhee, Cengiz Pehlevan, Risa Kawai, Alexandre Kempf, Sharon MH Gobes, and Bence P. Ölveczky. 2015. "Acute Off-Target Effects of Neural Circuit Manipulations." *Nature* 528(7582):358–63.

Pinto, Lucas, Kanaka Rajan, Brian DePasquale, Stephan Y. Thiberge, David W. Tank, and Carlos D. Brody. 2019. "Task-Dependent Changes in the Large-Scale Dynamics and Necessity of Cortical Regions." *Neuron* 104(4):810–24.

Pisupati, Sashank, Lital Chartarifsky-Lynn, Anup Khanal, and Anne K. Churchland. 2021. "Lapses in Perceptual Decisions Reflect Exploration." *Elife* 10:e55490.

Samejima, Kazuyuki, Yasumasa Ueda, Kenji Doya, and Minoru Kimura. 2005. "Representation of Action-Specific Reward Values in the Striatum." *Science* 310(5752):1337. doi: 10.1126/science.1115270.

Siniscalchi, Michael J., Victoria Phoumthipphavong, Farhan Ali, Marc Lozano, and Alex C. Kwan. 2016. "Fast and Slow Transitions in Frontal Ensemble Activity during Flexible Sensorimotor Behavior." *Nature Neuroscience* 19(9):1234–42.

Siniscalchi, Michael J., Hongli Wang, and Alex C. Kwan. 2019. "Enhanced Population Coding for Rewarded Choices in the Medial Frontal Cortex of the Mouse." *Cerebral Cortex* 29(10):4090–4106.

Soltani, Alireza, and Alicia Izquierdo. 2019. "Adaptive Learning under Expected and Unexpected Uncertainty." *Nature Reviews Neuroscience* 20(10):635–44. doi: 10.1038/s41583-019-0180-y.

Starkweather, Clara Kwon, Benedicte M. Babayan, Naoshige Uchida, and Samuel J. Gershman. 2017. "Dopamine Reward Prediction Errors Reflect Hidden-State Inference across Time." *Nature Neuroscience* 20(4):581–89.

Starkweather, Clara Kwon, Samuel J. Gershman, and Naoshige Uchida. 2018. "The Medial Prefrontal Cortex Shapes Dopamine Reward Prediction Errors under State Uncertainty." *Neuron* 98(3):616-629. e6.

Sul, Jung Hoon, Suhyun Jo, Daeyeol Lee, and Min Whan Jung. 2011. "Role of Rodent Secondary Motor Cortex in Value-Based Action Selection." *Nature Neuroscience* 14(9):1202–8.

Sul, Jung Hoon, Hoseok Kim, Namjung Huh, Daeyeol Lee, and Min Whan Jung. 2010. "Distinct Roles of Rodent Orbitofrontal and Medial Prefrontal Cortex in Decision Making." *Neuron* 66(3):449–60. doi: 10.1016/j.neuron.2010.03.033.

Sutton, Richard S., and Andrew G. Barto. 2018. *Reinforcement Learning: An Introduction*. MIT
press.

Tai, Lung-Hao, A. Moses Lee, Nora Benavidez, Antonello Bonci, and Linda Wilbrecht. 2012.
"Transient Stimulation of Distinct Subpopulations of Striatal Neurons Mimics Changes in
Action Value." *Nature Neuroscience* 15(9):1281–89.

Tsuchida, Ami, Bradley B. Doll, and Lesley K. Fellows. 2010. "Beyond Reversal: A Critical Role for
Human Orbitofrontal Cortex in Flexible Learning from Probabilistic Feedback." *The Journal
of Neuroscience* 30(50):16868. doi: 10.1523/JNEUROSCI.1958-10.2010.

Vaidya, Avinash R., Maia S. Pujara, Michael Petrides, Elisabeth A. Murray, and Lesley K. Fellows.
2019. "Lesion Studies in Contemporary Neuroscience." *Trends in Cognitive Sciences*
23(8):653–71.

Vertechi, Pietro, Eran Lottem, Dario Sarra, Beatriz Godinho, Isaac Treves, Tiago Quendera,
Matthijs Nicolai Oude Lohuis, and Zachary F. Mainen. 2020. "Inference-Based Decisions in
a Hidden State Foraging Task: Differential Contributions of Prefrontal Cortical Areas."
*Neuron* 106(1):166-176. e6.

Wang, Hongli, Heather K. Ortega, Huriye Atilgan, Cayla E. Murphy, and Alex C. Kwan. 2022. "Pupil
Correlates of Decision Variables in Mice Playing a Competitive Mixed-Strategy Game."
*Eneuro* 9(2).

Wilson, Robert C., and Anne GE Collins. 2019. "Ten Simple Rules for the Computational Modeling
of Behavioral Data." *Elife* 8:e49547.

Wilson, Robert C., Matthew R. Nassar, and Joshua I. Gold. 2010. "Bayesian Online Learning of the
Hazard Rate in Change-Point Problems." *Neural Computation* 22(9):2452–76.

1132