Spatial and lineage dependent processes underpin floristic assembly in the megadiverse Eastern South American mountains

Yago Barros-Souza [1,2] and Leonardo M. Borges [1]

Departamento de Botânica, Universidade de São Carlos, Brazil [1]
Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Brazil [2]
 Corresponding author: souzayagob@gmail.com

Acknowledgments

We thank Fernando Silveira, Rosa Scherson, Suzette Flantua, Rafael Izbicki, Renske Onstein, Kaori Nagata, two anonymous reviewers and the editors for their criticisms and suggestions. YB was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. No permits were needed to carry this work.

Abstract

Aim The astonishing diversity of ancient mountains was likely shaped by multiple evolutionary processes. However, there is an ongoing debate on what were the main processes driving the assembly of *campos rupestres*, the mega-diverse flora of Eastern South American mountains. Although the ancient nature of these mountains suggests their flora should be composed by relatively older lineages, they harbour a number of recently diverged clades. To better understand the evolution of ancient mountains' floras, we tested if the *campos rupestres* are mainly composed by relatively old or recent communities and if angiosperm diversity is geographically structured using analyses of diversity and endemism.

Location Eastern South America.

Time period Oligocene/Miocene to the present.

Major taxa studied Flowering plants.

Methods We used analyses of diversity and endemism for 10% of the *campos rupestres flora*. We obtained distribution data from online databases, and phylogenetic hypotheses from the literature. With these datasets, we estimated alpha and beta metrics of taxonomic and phylogenetic diversity, and conducted categorical analyses of neo- and paleo-endemism.

Results Phylogenetic overdispersion predominates in the *campos rupestres*. However, this general pattern is permeated by both lineage- and site-specific phylogenetic clustering, suggesting that recent diversification events depend on particular regional conditions and on the overall maintenance of old lineages. Although endemism patterns vary among different *campos rupestres* sites, paleo-endemism is widespread and particularly prominent where phylogenetic

overdispersion is evident. Moreover, phylogenetic composition indicates variable past spatial
connections across different sites, taxonomic composition is highly geographically structured and
seems to be influenced by the vegetation surrounding the *campos rupestres* and/or by abiotic
conditions.

Main conclusions Our results reinforce the idiosyncratic nature of diversification patterns in ancient
mountains and suggest that old, climatically buffered, infertile montane ecosystems not only
include both relatively old and recent lineages, but that recent diversification is lineage and
spatially dependent.

Keywords: CANAPE; Jaccard; OCBIL; Pleistocene; species-pump; UniFrac.

1 Introduction

Despite covering a strikingly small area of the globe, mountains house most of terrestrial biodiversity (Myers, Mittermeier, Mittermeier, Da Fonseca & Kent, 2000; Rafiqpoor, Kier, & Kreft, 2005; Kreft & Jetz, 2007; Rahbek et al., 2019). Their island-like nature and heterogeneous landscape promote diversification at multiple scales (Hughes & Eastwood, 2006; Drummond, Eastwood, Miotto & Hughes, 2012; Hughes & Atchison, 2015; Cortés, Garzón, Valencia & Madriñán, 2018; Nürk, Atchison, & Hughes, 2019; García-Rodríguez et al., 2021). However, the diversity of different mountains may have distinct origins. For example, new environments and geographic barriers that emerged during the Andes uplift led to fast and recent species diversification (Hughes & Eastwood, 2006; Särkinen, Pennington, Lavin, Simon & Hughes, 2012; Hughes, Pennington, & Antonelli, 2013). On the other hand, the long-term climatic and topographic stability of ancient mountains may have favoured the persistence of old lineages (Berry & Riina, 2005; Warren & Hawkins, 2006; Barbosa, Fernandes, & Sanchez-Azofeifa, 2015). Although these and other examples suggest that the evolution of montane diversity was intricate (e.g. Hughes & Atchison, 2015; Badgley et al., 2017; Antonelli et al., 2018), there is an ongoing debate over the main processes behind the origin of a hyper-diverse montane vegetation in Eastern South America (Silveira et al., 2016), the *campos rupestres*.

The *campos rupestres* are one of the richest and most endemic floras in the tropics (BFG, 2015). With more than 5000 plant species, it is similar to or almost twice as diverse as some biodiversity hotspots, such as Japan (5600 species), the Horn of Africa (5000 species), and New Zealand (2300 species) (Mittermeier, Turner, Larsen, Brooks & Gascon, 2011; Silveira et al., 2016). This floristic diversity is also spatially limited to an area smaller than Ireland or less than 1% of the Brazilian terrestrial territory (Silveira et al., 2016). *Campos rupestres* occur in old, climatically buffered, infertile landscapes (OCBIL; Silveira et al., 2016), over topographically heterogeneous and ancient quartzite, sandstone and ironstone formations ranging from 900 to over 2,000 m a.s.l. (Pedreira & De Waele, 2008; Fernandes, 2016). The vegetation mosaic of these areas, which include grasslands and rock-dwelling plant communities intermingled with gallery forest patches (for a detailed review, see Fernandes, 2016), occurs mostly in the Espinhaço Range, a mountain chain distributed along the Brazilian states of Minas Gerais and Bahia (Giulietti, Pirani, & Harley, 1997). Other disjunct sites, set apart by lowland vegetation are at the Brazilian Central Plateau (Goiás state and Distrito Federal) and at *Serra da Canastra* (western Minas Gerais state; Fig. 1; Silveira et al., 2016). The *campos rupestres* discontinuity is also reflected in species distribution, many of which are narrow and/or occur as small, disjunct populations (Giulietti et al., 1997; Echternacht, Trovó, & Takeo Sano, 2010; Echternacht, Trovó, Oliveira & Pirani, 2011; Lousada, Lovato, & Borba, 2013; Trovó, De Andrade, Sano, Ribeiro & Van den Berg, 2013).

Figure. 1: Main distribution of the *campos rupestres* within the Brazilian territory and across the Distrito Federal (DF) and the states of Bahia (BA), Minas Gerais (MG) and Goiás (GO). This vegetation complex is concentrated in the Espinhaço Range, but also occurs in the Brazilian Central Plateau and the *Serra da Canastra*. Here we considered the *campos rupestres* as divided into four different sections: the Southern and Northern Espinhaço, the Brazilian Central Plateau and the *Serra da Canastra*. Adapted from Silveira et al. (2016).

The distribution and diversity of *campos rupestres* plants have often been viewed as the result of a
species-pump mechanism, which promotes recurrent bursts of diversification similar to the ones
seen in the Andes (Harley, 1988; Giulietti et al., 1997; Flantua, O'dea, Onstein, Giraldo &
Hooghiemstra, 2019). According to this hypothesis, recurrent climatic variations during the
Tertiary and Quaternary (Adams, Maslin, & Thomas, 1999; Zachos, Dickens, & Zeebe, 2008) led to
population expansion-contraction cycles and intermittent gene flow. In this context, isolation –
sometimes together with hybridisation (e.g. Antonelli, Verola, Parisod & Gustafsson, 2010) – would
promote local diversification (Giulietti et al., 1997; Vasconcelos et al., 2020). Indeed, shifts in the
range of *campos rupestres* populations might have been common during the Pleistocene (Collevatti,
Rabelo, & Vieira, 2009; Bonatelli et al., 2014; Barbosa, Fernandes & Sanchez-Azofeifa, 2015; Barres,
Batalha-Filho, Schnadelbach & Roque, 2019), a period of high diversification (Souza et al., 2013;
Trovó et al., 2013; Ribeiro, Rapini, Damascena & Van den Berg, 2014; Rando et al., 2016;
Vasconcelos et al., 2020). Therefore, if these processes were as common as expected by the species-
pump hypothesis, recently diverged lineages (i.e. neo-endemics) and signatures of recent
diversification (e.g. phylogenetic clustering) should dominate *campos rupestres* communities.

However, similarly to other OCBILs, such as the Cape and southwestern Australia (Cowling et al.,
2014), old communities should prevail in the *campos rupestres* due to the long-term climatic and
topographical stability of these ancient mountains (Hopper, 2009). Although this prediction has
been recently contested (Cortez et al., 2020), several evidences indicate that the *campos*
rupestres may be one of the oldest modern day open vegetation in Tropical America, and include
lineages diversified as early as the Oligocene/Miocene (Rapini, Berg, & Liede-Schumann, 2007;
Antonelli et al., 2010; Gustafsson, Verola, & Antonelli, 2010; Hughes, Pennington & Antonelli, 2013;
Zappi, Moro, Meagher & Nic Lughadha, 2017; Alcantara, Ree, & Mello-Silva, 2018). Thus, *campos*
rupestres hyper-diversity would be, at least in part, due to the accumulation of old lineages.

Whichever processes were in action during formation of the flora of ancient mountains, their
nature and/or intensity likely varied across space (Croizat, 1962; Nürk et al., 2020). However, little
is known about how the *campos rupestres* high levels of topographical, climatic and edaphic
heterogeneity (Giulietti et al., 1997; Schaefer, Cândido, Corrêa, Nunes & Arruda, 2016; Neves et al.,
2018) shaped the distribution of old and/or recent lineages (but see Bitencourt & Rapini, 2013).
Hence, to better understand the *campos rupestres*' evolutionary history, it is necessary to evaluate
how diversity is distributed across space.

Spatial analyses of phylogenetic diversity and endemism are reliable approaches to address this
issue (e.g. Forest et al., 2007; Mishler et al., 2014; Heenan, Millar, Smissen, McGlone & Wilton, 2017;
Scherson et al., 2017). Phylogenetic diversity (Faith, 1992) informs the proportion of recently and
deeply diverged lineages in local communities (Webb, Ackerly, McPeek & Donoghue, 2002; Divya,
Ramesh & Karanth, 2021), while the distinction between centres of neo- and paleo-endemism
shows how these different types of lineages are distributed in space (Mishler et al., 2014). These
patterns can be complemented by analyses of beta and phylobeta diversity (e.g. Jaccard, 1901;
Lozupone & Knight, 2005), which point to past and present connections between areas, as well as
to historical processes structuring communities (Graham & Fine, 2008).

Here we aimed to identify which evolutionary processes may have shaped the flora of an ancient
Tropical mountain and how they are distributed across space. To do so, we assessed fine spatial
patterns of phylogenetic diversity and endemism for over 10% of the *campos rupestres* flora. If
predictions for the persistence of old lineages in ancient landscapes are met, patterns of
phylogenetic overdispersion and paleo-endemism should prevail in the *campos rupestres*. On the
other hand, widespread phylogenetic clustering and neo-endemism would indicate communities to
be composed by recently diversified lineages, reinforcing the role of a species-pump mechanism in

the diversification of altitudinal floras (e.g. Flantua et al., 2019; Vasconcelos et al., 2020). Moreover,
a geographical organisation of the taxonomic and phylogenetic composition of the *campos*
rupestres would indicate the biological connection of these mountains to be under topographical,
climatic and edaphic constraints.

2 Methods

To answer our questions, we used 572 species from nine angiosperm groups for which well-
sampled phylogenetic hypotheses are available: *Mimosa* L. (Leguminosae), *Diplusodon* Pohl
(Lythraceae), *Comanthera* L.B. Sm. (Eriocaulaceae), Velloziaceae J. Agardh, Lychnophorinae Benth.
(Asteraceae), *Paepalanthus* Mart. (Eriocaulaceae), *Calliandra* Benth. (Leguminosae), *Minaria* T.U.P.
Konno (Apocynaceae) and Marcetieae M.J.R. Rocha, P.J.F. Guim. & Michelang. (Melastomataceae).
Although this strategy could bias analyses towards the evolutionary history of sampled lineages,
these groups are well-represented in the *campos rupestres*, their life forms range from herbs to
trees and they comprise both monocots and eudicots. They are also mostly restricted to grasslands
and rocky outcrops, commonly not occurring in the gallery forest patches dispersed across
the *campos rupestres* area. Therefore, they are an adequate proxy for general *campos*
rupestres diversity patterns.

Our analyses were divided into two parts: (1) data compilation and cleaning, and (2) spatial
analyses. Below we present a general workflow for each step, all of which were conducted in the R
statistical computing environment (R Core Team, 2013). For details about each analysis, see the
Supporting Information (Appendix 1).

2.1 Data compilation and cleaning

We retrieved occurrence data for our model groups from two complementary databases:
GBIF (GBIF.org, 2020) and speciesLink (CRIA, 2020). Data extracted from these databases were
concatenated into a single data set to facilitate posterior treatment. To reduce data dimensionality,
we removed all variables but the ones relevant for data cleaning and spatial analyses.

To clean our data, we first pruned duplicated records and removed records without identification
at species level, without names of determiners or without vouchers. Then we followed the methods
proposed by Magdalena et al. (2018), which consist of: (1) defining a coordinate reference system;
(2) standardising names of geographic units; (3) selecting and evaluating records with geographic
coordinates; (4) identifying records without geographic coordinates; and (5) inferring geographic
coordinates based on municipalities or field notes. To avoid bias due to species misidentification,
we also updated the taxonomy, corrected typos in species names and limited the data set to records
identified by taxonomic specialists (authors of monographs, floras or taxonomic revisions on the
family of the focal group).

Finally, we removed all records whose coordinates fell outside the *campos rupestres*'
distribution (Silveira et al., 2016).

2.2 Spatial analyses

To understand the taxonomic and phylogenetic composition and structure of plant communities in
the *campos rupestres*, we calculated the following metrics: species richness and phylogenetic

diversity (PD; Faith, 1992), Jaccard distance (Jaccard, 1901) and UniFrac (Lozupone & Knight,
2005). We also conducted categorical analyses of neo- and paleo-endemism (CANAPE; Mishler et
al., 2014) to assess the distribution of neo- and paleo-endemic lineages.

Since part of our analyses rely on phylogenetic trees, we obtained the most recent phylogenetic
hypothesis for each group (Echternacht et al., 2014; Loeuille, Semir, Lohmann & Pirani, 2015;
Alcantara et al., 2018; Inglis & Cavalcanti, 2018; Andrino et al., 2020; Vasconcelos et al., 2020), all of
which include at least 50% of their diversity. Trees and distribution matrices were both pruned to
include only species present in both. Also, we scaled every tree by total tree length, making it
possible to compare the results for different groups and with future studies. All analyses were
conducted at species level.

201   2.2.1 Spatial grids

Although it is hard to define the best resolution of the spatial grids needed for diversity
analyses (Blackburn & Gaston, 2002), their configuration affects the results of biogeographical
studies (Willis & Whittaker, 2002; Rahbek, 2005). Thus, we based grid cells' size on redundancy
values, which take sampling effort in consideration (Scherson et al., 2017). To do so, we first
removed grid cells that had only one sample per species. Then we calculated redundancy for all
cells at different sizes and assessed the relationship between grid cells' size and redundancy (i.e.,
the median of all redundancy values at a given cell's size). To investigate if the diversity of each cell
could be biased by unequal sampling effort, we estimated asymptotic species richness (Chao et al.,
2014), a measure of "true" diversity, for each taxonomic group and for all of them combined. The
high correlation between observed and estimated diversity ($R^2 \geq 0.82$ for two groups, $R^2 \geq 0.93$ for
the other groups, $R^2 = 0.99$ for all groups combined, and p-values < 0.05 for all groups individually
and combined) indicates our dataset is evenly sampled and can be used in subsequent analyses
without any corrections (e.g. interpolation or extrapolation (Chao et al., 2014). See the supporting
information (Appendix 1) for details and results of these analyses.

216   2.2.2 Alpha diversity

We calculated two metrics of alpha diversity: richness and phylogenetic diversity (PD). For PD, we
used the 'picante' package (Kembel et al., 2010) to sum the branch lengths connecting a given set of
species to the root of a phylogenetic tree (Faith, 1992). We also tested the correlation between
richness and PD with Spearman's correlation tests (Spearman, 1904).

Because PD is often correlated with richness (Faith, 1992; Polasky, Csuti, Vossler & Meyers, 2001;
Rodrigues & Gaston, 2002; Tôrres & Diniz-Filho, 2004; Forest et al., 2007; Scherson et al., 2017),
decoupling these two metrics provides a powerful tool to assess community phylogenetic structure
and, thus, offers valuable insights into macroecological and biogeographical processes (Webb et al.,
2002). We did this by using the residuals of linear regression models between phylogenetic
diversity and richness. Residual values above or below zero indicate PD higher or lower than
expected based on richness, respectively (Forest et al., 2007; Brown et al., 2020; Colville et al.,
2020).

Finally, to gain a broader perspective on the spatial distribution of alpha diversity, we calculated
total richness and total PD for all groups occurring in each grid cell. Then, we calculated the
residuals of each metric for all groups combined (Brown et al., 2020).

232   2.2.3 Beta diversity

To assess communities' structure both in the present as in the past (Wiens & Donoghue, 2004;
Emerson & Gillespie, 2008; Graham & Fine, 2008), we conducted UPGMA analyses for each group
using dissimilarity matrices based on Jaccard distance (beta diversity; Jaccard, 1901) and UniFrac
(phylobeta diversity; Lozupone & Knight, 2005). As with alpha diversity metrics, we also performed
an UPGMA analysis based on Jaccard distance for all groups combined simply by merging their
distribution matrices. This approach was not possible for UniFrac, as it relies on pairwise
calculations based on unique phylogenetic trees.

Before generating the consensus dendrograms (1000 replicates; consensus criterion of 0.5) for
each group, we removed singletons (species whose occurrence is restricted to a single cell) in order
to avoid noise. Then, to better visualise spatial structure, we subdivided each dendrogram using
fusion level values (Borcard, Gillet, & Legendre, 2018), while attempting to maintain a consistent
number of clusters between all groups. All beta diversity analyses were conducted with
'recluster' (Dapporto et al., 2013).

246 2.2.4 CANAPE

The categorical analyses of neo- and paleo-endemism were conducted following the methods
developed by Mishler et al. (2014), which involve prior calculation of phylogenetic endemism
(PE; Rosauer, Laffan, Crisp, Donnellan & Cook, 2009) and relative phylogenetic endemism
(RPE; Mishler et al., 2014). After calculating PE and RPE with the 'PDcalc' package (D. Nipperess;
under development), we performed all CANAPE's statistical tests with a custom code. CANAPE
includes four endemism categories: neo-endemism (rare, younger lineages), paleo-endemism (rare,
older lineages), mixed endemism (a mix of neo- and paleo-endemics), and super-endemism (a
strong signal of mixed endemism). Please bear in mind that the distinction between paleo- and neo-
endemism is not absolute, but relative to the age of sampled lineages. In our analyses, we found
some grid cells that could not be classified in any of CANAPE's categories, as they do not meet the
criteria for all its statistical tests. Thus, we defined their endemism as "uncertain". See Supporting
Information for details about this analysis (Appendix 1).

Because we did not include occurrence data from areas surrounding the *campos rupestres* adjacent
areas, our endemism analyses could be biased. However, CANAPE relies on the range of
phylogenetic branches, not that of terminals. Also, most *campos rupestres* species are restricted to
small areas within it (e.g. Rapini, 2010; Echternacht et al., 2011; Bitencourt & Rapini, 2013;
Trovó et al., 2013). Thus, we believe our analyses were robust enough to assess the distribution of
different types of endemism within the *campos rupestres*' range. Nevertheless, including the
surrounding vegetation in future studies will certainly improve knowledge on these patterns.

267 3 Results

268 3.1 Redundancy

269 Redundancy values for all groups steadily increase together with grid cells' size until reaching a
270 plateau around 0.9 (Fig. S2.1). However, a value of 0.9 for redundancy defines cells' sizes too large
271 to recover variations in spatial patterns. Hence, aiming for a balance between resolution and
272 redundancy, we used cells of 0.6 per 0.6 decimal degrees (roughly 66 km per 66 km), whose
273 redundancy varies between 0.6 and 0.8.

3.2 Richness and phylogenetic diversity

Richness in the *campos rupestres* is concentrated in the Southern Espinhaço, where some cells include more than 200 species, and in the Brazilian Central Plateau (Figs. 2a and S2.2). The richness peak in the national capital could be a bias due to automatic georreferencing using countries' capitals (Zizka et al., 2019) or higher collecting effort in this area, which harbours several herbaria and research institutions. However, as some of our model groups are very diverse in this area (*Mimosa* and *Diplusodon*; Barneby, 1991; Cavalcanti, 1995; Cavalcanti, 2007) and our dataset is relatively evenly sampled, we decided to keep records for this particular cell to avoid underestimation of *campos rupestres* diversity in the Brazilian Central Plateau.

Figure 2: Combined results of (a) species richness and (b) phylogenetic diversity for all groups. Phylogenetic trees were scaled by total tree length so phylogenetic diversity results could be combined. Transparent grid cells indicate absent data

As expected (Faith, 1992; Polasky et al., 2001; Rodrigues & Gaston, 2002; Tôrres & Diniz-Filho, 2004; Forest et al., 2007; Scherson et al., 2017), PD follows richness distribution (Fig. S2.3), and highest PD values occur in the Southern Espinhaço, followed by the Brazilian Central Plateau and the Northern Espinhaço (Figs. 2b and S2.4). Although we noticed some individual groups may dominate and skew the residuals of the regression between total richness and total PD (Fig. S2.5), the combined residuals of all groups present a more accurate general view of the phylogenetic structure of the *campos rupestres* communities.

The PD of a large proportion of *campos rupestres* communities follow what is expected by richness (50% of residual values range from -0.2 to 0.2), particularly in the Brazilian Central Plateau. Nonetheless, the residual values show that communities in which PD is higher than expected by richness (phylogenetic overdispersion; positive residuals) are more common than communities in which PD is lower than the expected (phylogenetic clustering; negative residuals), more so in the Espinhaço Range and *Serra da Canastra* (Fig. 3). Although a single Southern Espinhaço site has an extremely low residual value, phylogenetic overdispersion is usually stronger than phylogenetic clustering (Fig. 3). The highest levels of phylogenetic overdispersion occur in the Northern Espinhaço, particularly when individual groups are considered.

302
Figure 3: Spatial distribution of residuals from linear regressions between species richness and phylogenetic diversity
across the *campos rupestres* for all groups combined (a). Positive and negative residuals respectively indicate
phylogenetic overdispersion and clustering (i.e. communities in which phylogenetic diversity is higher or lower than
expected based on richness). Transparent grid cells indicate absent data). The violin boxplot (b) depict the distribution of
residuals along the total range.

Residual values for each focal group (Fig. S2.6) agree with a general pattern of phylogenetic
overdispersion, but also show variations in the occurrence of phylogenetic clustering. For example,
phylogenetic clustered communities not clearly seen in the joint analysis occur in the Brazilian
Central Plateau (*Mimosa* and *Diplusodon*), in the Southern Espinhaço (*Comanthera* and *Minaria*), in
the Northern Espinhaço (Marcetieae and *Calliandra*), and across the whole Espinhaço Range
(*Paepalanthus*).

3.3 CANAPE

Although the distribution of endemism categories varies among individual groups, CANAPE results
indicate that paleo-endemic lineages predominate in the Northern Espinhaço, while other *campos*
rupestres areas include a mix of paleo- and neo-endemism (Fig. 4; see also Fig. S2.7 for detailed
information on each group). *Mimosa*, Velloziaceae and Lychnophorinae are paleoendemic to the
Northern Espinhaço, which also includes a case of mixed endemism for *Comanthera*. All endemism
categories occur in the Southern Espinhaço: on its northern part, paleo-, super and mixed
endemism are seen for *Paepalanthus*, *Comanthera* and Velloziaceae, whereas super, mixed, paleo-
and neo-endemism of *Paepalanthus*, *Comanthera*, *Calliandra*, and *Minaria* all occur on its northern
part. Finally, mixed endemism for *Diplusodon* and Velloziacae defines the *Serra da Canastra*, while
neo-, paleo- and super endemism characterise the Brazilian Central Plateau
(*Mimosa*, *Diplusodon*, *Comanthera* and Marcetieae).

Figure 4: CANAPE results of all groups combined. Neo-endemic cells contain a predominance of short, rare phylogenetic branches; paleo-endemic cells contain a predominance of long, rare branches; and cells with mixed endemism contain a mixture of both. Mixed endemism can be further categorised as super endemism in a cell in which it is statistically significant at a lower confidence interval ($\alpha \leq 0.001$). Combining the results for all groups results in some cells harbouring multiple endemism types. Transparent grid cells indicate absent data.

It is important to note that we were not able to categorise a few grid cells for *Comanthera*. This is likely due to low statistical power: *Comanthera* is a small genus (Echternacht et al., 2014), and the number of species assessed in this work was further reduced so that the phylogenetic tree could match the distribution matrix (see Methods). Thus, low sampling may have negatively affected the permutations performed during CANAPE analyses.

3.4 Jaccard distance and UniFrac

UPGMA analyses based on Jaccard distance (Fig. 5) showed that plant communities are geographically structured. Although the limits of these geographical patterns might have changed over time, the Northern Espinhaço flora stands in contrast with the area encompassing the Brazilian Central Plateau, the *Serra da Canastra* and the Southern Espinhaço Range (hereafter referred to as the Central Plateau-Southern Espinhaço arch).

343
Figure 5: Hierarchical clustering of *campos rupestres* communities based on their taxonomic composition. Colours and
cells in the map correspond to colours and terminals in the dendrogram, respectively. The closer the terminals are, more
taxonomic identity the corresponding cells share. Transparent grid cells indicate absent data. See supporting information
for bootstrap values.

The remarkable contrast between the taxonomic composition of the Northern Espinhaço and that
of other *campos rupestres* is evident for all groups combined and also for each taxon individually,
attesting to the consistency of this pattern (Figure 5). The floristic cohesiveness between the
Brazilian Central Plateau, the *Serra da Canastra* and the Southern Espinhaço is also very strong.
Even in the few cases in which the Brazilian Central Plateau and/or the Southern Espinhaço appear
as taxonomically unique (e.g. *Diplusodon* and *Paepalanthus*), these regions are nested within a
larger Central Plateau-Southern Espinhaço group.

A divide between the Central Plateau-Southern Espinhaço and the Northern Espinhaço is also seen
in the phylogenetic composition of *campos rupestres* communities, particularly
in *Mimosa*, *Diplusodon*, Velloziaceae, *Paepalanthus* and Marcetieae, but with fuzzier geographic
limits (Fig. 6). For example, the results for *Mimosa* extend the limits of a Central Plateau-Southern
Espinhaço's cluster further north, while a Northern Espinhaço's cluster includes close areas in the
Southern Espinhaço.

Figure 6: Hierarchical clustering of *campos rupestres* communities based on their phylogenetic composition. Colours and
cells in the map correspond to colours and terminals in the dendrogram, respectively. The closer the terminals are, more
phylogenetic identity the corresponding cells share. Transparent grid cells indicate absent data. See supporting
information for bootstrap values.

There are exceptions to this general geographic structure. For example, in *Calliandra* the Brazilian
Central Plateau appears as the most evolutionary contrasting region, while the Southern and
Northern Espinhaço form a single, cohesive region. For *Minaria*, the Espinhaço Range and the *Serra*
da Canastra form a cluster that contrasts to the Brazilian Central Plateau, even though both regions
are nested into a widespread cluster. Moreover, the central Southern Espinhaço frequently appears
as an evolutionary unique area, as seen for *Mimosa*, *Diplusodon*, Lychnophorinae, and *Minaria*.

4 Discussion

Aiming to understand the evolutionary history of ancient tropical mountains, here we investigated
spatial patterns of richness, phylogenetic diversity and endemism for the *campos rupestres* mega-
diverse flora. We also assessed phylogenetic and taxonomic differences between communities using
beta and phylobeta diversity metrics. Although particular results reinforce that the *campos*
rupestres are a complex assemblage of different evolutionary histories, these are not evenly
distributed across space. Below we discuss the relationship between these spatial patterns and
what processes might have shaped them.

4.1 Spatial distribution of recent and old lineages

Richness and phylogenetic diversity are not evenly distributed in the *campos rupestres*, but
concentrated in the Brazilian Central Plateau and the Southern Espinhaço. Decoupling these two
metrics from each other reveals different aspects of community composition. The combined
residuals of all groups indicate that phylogenetic overdispersion is more common than
phylogenetic clustering, even though PD follows what is expected by richness in a large number of
sites (ca. 50% of grid cells). The particular predominance of expected PD values in the Brazilian
Central Plateau, however, must be seen with caution, as there may be a bias against groups whose
communities are phylogenetically clustered in this region (e.g. *Mimosa* and *Diplusodon*; Fig. S2.5).
Regarding endemism patterns, paleo-endemism predominates in the Northern Espinhaço, while the
Southern Espinhaço, *Serra da Canastra* and Brazilian Central Plateau floras (Central Plateau-
Southern Espinhaço arch) include a mixture of paleo- and neo-endemic lineages.

Widespread phylogenetic overdispersion indicates the predominance of relatively older (deeply
diverged) lineages (Lososová et al., 2015; Colville et al., 2020; Divya et al., 2021). CANAPE
reinforces this by showing that endemics commonly belong to old lineages (paleo-endemism). This
pattern is especially prominent in the Northern Espinhaço, where both paleo-endemism and
phylogenetic overdispersion predominate. Contrasting with the Northern Espinhaço, the Central
Plateau-Southern Espinhaço arch includes more areas of phylogenetic clustering, with a single
Southern Espinhaço site showing the lowest residual value (close to -1), and a mix of paleo- and
neo-endemism or even well-marked neo-endemism. This pattern is clearer in analyses of individual
taxa, especially in the Brazilian Central Plateau, where neo-endemism
for *Mimosa* and *Diplusodon* overlaps with phylogenetically clustered communities. These results
show that the Central Plateau-Southern Espinhaço arch flora includes patches of relatively young
(short branches) and closely related lineages (Webb et al., 2002; Divya et al., 2021), as previously
indicated by phylogenetic and diversification analyses (Trovó et al., 2013; Bonatelli et al., 2014;
Rando et al., 2016; Vasconcelos et al., 2020).

Taken together, these patterns indicate that recent diversification is spatially restricted, lineage-
dependent and embedded in a widespread and relatively older flora (Figures 3 and 4). As
expected (Hopper, Silveira, & Fiedler, 2016; Silveira et al., 2016; Hopper, Lambers, Silveira &
Fiedler, 2021), long-term climatic and topographical stability (Pedreira & De Waele, 2008;
Silveira et al., 2016) may have favoured the persistence of old lineages, promoting an accumulation
of species over time similar to that seen in the Cape and southwestern Australia (Cowling et al.,
2014). Such a reservoir of older lineages was likely the raw material for bursts of diversification,
which could have been driven by the reassembly of ancient alleles (Richards et al., 2021), for
example. Moreover, high heterogeneous topography, specific edaphic conditions, and particular
ecophysiological requirements (Giulietti et al., 1997; Fernandes, 2016; Oliveira et al., 2016, Schaefer
et al., 2016) likely constrained plant diversification across the *campos rupestres*. Thus, although
probably present (e.g. (Antonelli et al., 2010; Bonatelli et al., 2014; Barbosa et al., 2015; Barres et

al., 2019), climatic driven species-pumps apparently were not the main process shaping floristic
diversity in the *campos rupestres* and other ancient mountains (e.g. Onstein & Linder, 2016).

A matrix of phylogenetic overdispersed communities could have been assembled by the free flow of
multiple unrelated lineages across the vegetation surrounding the *campos rupestres*, as predicted
by a model invoking changes in niche breadth as the precursors of lineage range expansion and
diversification (Rapini, Bitencourt, Luebert & Cardoso, 2020). However, processes other than the
balance between adaptive radiations and extinctions after drift invoked by this model could have
shaped *campos rupestres* diversity. Hybridisation, for example, although relatively rare in
OCBILs (Hopper, 2018), is widespread in angiosperms (Rieseberg & Willis, 2007), not uncommon
in the *campos rupestres* (e.g., Borba & Semir, 1998; Ribeiro et al., 2008; Yamagishi-Costa & Forni-
Martins, 2009), and often linked to macroevolutionary diversification events (Rieseberg & Willis,
2007). Also, species already well-established in the *campos rupestres* were likely less prone to
extinction than species of similar ecology generated during bursts of recent
diversification (McPeek, 2007). The role of such ecologically similar and transient species in
the *campos rupestres* flora is reinforced by the occurrence of phylogenetic clustering in species-
rich sites (McPeek, 2007), shown here for individual taxa, as well as by the overall high proportion
of short branches seen for the Central Plateau–Southern Espinhaço arch (Cortez et al., 2020).
Repetition of these localised birth and death events over time could produce the spatter of recent
diversification within a matrix of relatively older lineages we observed here. Why local
diversification is more common in the Central Plateau–Southern Espinhaço arch than in the
Northern Espinhaço can be better discussed in the context of beta and phylobeta diversity analyses.

4.2 Present and past geographical structure of plant communities

As expected for island-like systems (e.g. Garcia-Verdugo, Forrest, Balaguer, Fay & Vargas, 2010;
Kubota, Hirao, Fujii & Murakami, 2011; Carstensen et al., 2012), beta diversity analyses show a
geographic structuring of the *campos rupestres* flora. Although the Southern Espinhaço and the
Brazilian Central Plateau differ in species composition, together with the *Serra da Canastra* they
form a cohesive group (the Central Plateau-Southern Espinhaço arch), which stands in taxonomic
and phylogenetic contrast to the Northern Espinhaço. The limits between these two major areas,
however, apparently varied across time.

A floristic dissimilarity between the Northern Espinhaço and the Southern Espinhaço has been
previously stressed (Harley, 1988; Rapini, 2010; Bitencourt & Rapini, 2013; Trovó et al., 2013;
Neves et al., 2018) and attributed to the role of the lowland between the two portions of the
Espinhaço Range as a barrier to migration (Harley, 1988; Bitencourt & Rapini, 2013). However,
even though *campos rupestres* species are known for their limited dispersal potential (Giulietti et
al., 1997; Conceição et al., 2016; Arruda et al., 2020), the connections between distant areas
belonging to the Central Plateau-Southern Espinhaço arch questions the role, or at least the
strength, of such geographical barriers. Indeed, changes in community composition across
the *campos rupestres* may be better predicted by environmental conditions (Neves et al., 2018):
while the Northern Espinhaço is relatively less fragmented and located in the driest extreme of the
Brazilian precipitation gradient, the Central Plateau-Southern Espinhaço arch includes highly
fragmented sites and lies in a relatively moister environment (Bitencourt & Rapini, 2013; Neves et
al., 2018). At the same time, the propensity of closer sites to share lineages indicates that
geographic proximity may have also influenced the assembly of *campos rupestres* communities.
Although the role of distance has been dismissed for the *campos rupestres* woody flora (Neves et

al., 2018), our analyses including both woody and herbaceous taxa indicate it should be further
tested.

Differences in the type of vegetation surrounding the *campos rupestres*, which have a remarkable
influence on its woody flora (Neves et al., 2018; Assunção-Silva & Assis, 2021), probably also
shaped the distribution of species composition. While the Central Plateau-Southern Espinhaço arch
is chiefly embedded in a savanna-like vegetation (the Brazilian *cerrado*), the Northern Espinhaço is
predominantly surrounded by a massive seasonally dry tropical forest, the *caatinga* (Conceição et
al., 2016; Neves et al., 2018). One example of the role of the surrounding vegetation comes
from *Calliandra*, in which a lineage from the *caatinga* radiated into a clade mostly endemic to the
Northern Espinhaço (Souza et al., 2013). Similarly, radiations in the Cape usually occurred after
transitions from forest to fynbos (Onstein, Carter, Xing & Linder, 2014), reinforcing the idea that
adjacent environments might have played a major role in the assembly of mega-diverse flora in
ancient mountains. Different processes could explain how lineages move from one vegetation to the
another, including exaptations (*sensu* Gould & Vrba, 1982), the previously mentioned changes in
niche breadth (Rapini et al., 2020), or sharing of similar environmental filters. Although these
processes are not mutually exclusive, if the *campos rupestres* distribution area was not affected by
Pleistocene climatic fluctuations (Rapini et al., 2020), lineage migration between disjunct sites must
have occurred through the vegetation surrounding the *campos rupestres*. Thus, a scenario in which
the *cerrado* chiefly exchanged lineages with the Central Plateau-Southern Espinhaço arch, while the
Northern Espinhaço interacted mainly with the *caatinga* flora helps to explain the division of
the *campos rupestres* into these two distinct floristic groups.

The putative influence of the surrounding vegetation may also explain phylogenetic diversity and
endemism patterns, particularly the predominance of neo-endemism and phylogenetic clustering in
the Central Plateau-Southern Espinhaço Arch. Because the *caatinga* is also an ancient and
climatically stable habitat (Werneck, Costa, Colli, Prado & Sites Jr, 2011; Queiroz, Cardoso,
Fernandes & Moro, 2017), a higher proportion of paleo-endemic and deeply diverged lineages are
expected to colonise the Northern Espinhaço. On the other hand, the effects of Pleistocene climatic
fluctuations over the more recently assembled *cerrado* (Behling, 2002; Safford, 2007; Simon et al.,
2009; Werneck, Nogueira, Colli, Sites Jr & Costa, 2012) could have promoted recent lineage- and
site-specific diversification in the Central Plateau-Southern Espinhaço arch, especially if
morphological innovations matched climatic shifts followed by the emergence of ecological
opportunities (Onstein et al., 2014). Nonetheless, it is important to note that the Northern
Espinhaço and the Central Plateau-Southern Espinhaço arch are not completely evolutionary
independent, as shown by the variations in their limits seen for different lineages and across
relative time (Fig. 6). This indicates that, even though some lineages were unable to disperse
(possibly due to intrinsic limitations, as suggested by a few evolutionary unique areas; Fig. 6),
disjunct sites were probably connected multiple times and by different lineages in the past.
Whether expansion of the *campos rupestres* area (e.g. Antonelli et al., 2010), niche shifts to the
surrounding vegetation (Rapini et al., 2020), or other processes, such as phenotypic plasticity,
promoted lineage dispersion is still debatable. Regardless of these uncertainties, it is clear that
studies incorporating multiple aspects, such as abiotic, biotic, and clade-specific features of
the *campos rupestres* and their surrounding floras may help to elucidate how historical
connections between disjunct ancient mountains were possible.

506 5 Conclusion

Competing evolutionary processes have been evoked to explain the origins and diversity of both
young and ancient mountains across the world (Hopper, 2009; Onstein et al., 2014; Flantua et al.,

2019; Rapini et al., 2020; Vasconcelos et al., 2020), including long-term lineage
persistence (Hopper, 2009), changes in niche breadth followed by range expansion (Rapini et al.,
2020), and local diversification driven by climatic shifts (Vasconcelos et al., 2020) among others.
However, even though evolution is strongly dependent on space (Croizat, 1962; Nürk et al., 2020),
species and lineage distribution are often neglected when assessing the role of these processes in
the assembly of mega-diverse floras, such as those of Eastern South American mountains,
the *campos rupestres*. Here we aimed to assess the distribution of multiple evolutionary processes
across the landscape of one of the richest floras in the world by combining spatial and phylogenetic
data.

We show that the *campos rupestres* are split into two large areas of cohesive floristic diversity: the
Central Plateau-Southern Espinhaço arch and the Northern Espinhaço. Although likely variable in
the past, the contrast between these regions is reinforced by patterns of phylogenetic diversity and
endemism. While phylogenetic overdispersion and paleo-endemism are dominant in the Northern
Espinhaço, phylogenetic clustering and neo-endemism are mostly restricted to the Central Plateau-
Southern Espinhaço arch, with very few exceptions. This is not only in agreement with patterns of
floristic composition, but also with the distribution of environmental conditions and surrounding
vegetation (Neves et al., 2018), which might have played an important role in the assembly of old,
climatically buffered, infertile vegetations (Hopper, 2009; Onstein et al., 2014; Hopper et al., 2016).

Overall, this species-rich, highly endemic vegetation is mainly composed by relatively older
lineages, spattered with patches of recently diversified lineages. Therefore, our spatial approach
reconciles the apparent conflicting evidences for recent and fast diversification and the persistence
of old lineages in ancient, climatic buffered, infertile landscapes, such as the Greater Cape Floristic
Region (Linder, 2008; Cowling et al., 2014) and the *campos rupestres* (Simon et al., 2009;
Bitencourt & Rapini, 2013; Hughes et al., 2013; Souza et al., 2013; Trovó et al., 2013; Bonatelli et al.,
2014; Silveira et al., 2016). Nonetheless, our work shows that recent diversification events in long-
term stable mountains can be both lineage- and site-dependent and may rely on the overall
maintenance of relatively older lineages. These findings emphasise the idiosyncratic nature of
diversification events and highlight the importance of considering both the spatial component and
intrinsic lineage aspects when investigating diversity-related questions, as space clearly affected
the balance between different evolutionary processes behind the astonishing diversity of ancient
mountains.

540 Data availability statement

541 The data that support the findings of this study are openly available at
542 https://zenodo.org/record/7226249 (DOI: 10.5281/zenodo.7226249).
543

544 Conflict of interest statement

545 All authors declare that they have no conflicts of interest.

547 References

Adams, J., Maslin, M., & Thomas, E. (1999). Sudden climate transitions during the Quaternary. *Progress in*
physical geography, 23(1), 1–36.

Alcantara, S., Ree, R. H., & Mello-Silva, R. (2018). Accelerated diversification and functional trait evolution
in Velloziaceae reveal new insights into the origins of the *campos rupestres*' exceptional floristic richness.
Annals of botany, *122*(1), 165–180.

Andrino, C. O., Sano, P. T., Inglis, P. W., Hensold, N., Costa, F. N., & Simon, M. F. (2020). Phylogenetics
of *Paepalanthus* (Eriocaulaceae), a diverse Neotropical monocot lineage. *Botanical Journal of the Linnean*
Society.

Antonelli, A., Kissling, W. D., Flantua, S. G., Bermúdez, M. A., Mulch, A., Muellner-Riehl, A. N., Kreft, H., et al.
(2018). Geological and climatic influences on mountain biodiversity. *Nature Geoscience*, *11*(10), 718–725.

Antonelli, A., Verola, C. F., Parisod, C., & Gustafsson, A. L. S. (2010). Climate cooling promoted the expansion
and radiation of a threatened group of South American orchids (Epidendroideae: laeliinae). *Biological Journal*
of the Linnean Society, *100*(3), 597–607.

Arruda, A. J., Junqueira, P. A., Rodrigues, H. T., Yvanez, F., Poschlod, P., Silveira, F. A., & Buisson, E.
(2020). Limited seed dispersability in a megadiverse OCBIL grassland. *Biological Journal of the Linnean*
Society.

Assunção-Silva, C. C., & Assis, L. C. S. (2021). Areas of endemism of Lauraceae: new insights on the
biogeographic regionalization of the Espinhaço Range, Brazil. *Cladistics*. Wiley Online Library.
doi:10.1111/cla.12481

Badgley, C., Smiley, T. M., Terry, R., Davis, E. B., DeSantis, L. R., Fox, D. L., Hopkins, S. S., et al. (2017).
Biodiversity and topographic complexity: Modern and geohistorical perspectives. *Trends in Ecology &*
Evolution, *32*(3), 211–226.

Barbosa, N. P. U., Fernandes, G. W., & Sanchez-Azofeifa, A. (2015). A relict species restricted to a quartzitic
mountain in tropical America: An example of microrefugium? *Acta Botanica Brasilica*, *29*(3), 299–309.

Barneby, R. (1991). Sensitivae censitae: A description of the genus *mimosa* Linnaeus (Mimosaceae) in the
New World. *Memoirs of The New York Botanical Garden*, *65*.

Barres, L., Batalha-Filho, H., Schnadelbach, A. S., & Roque, N. (2019). Pleistocene climatic changes drove
dispersal and isolation of *Richterago discoidea* (Asteraceae), an endemic plant of campos rupestres in the
central and eastern Brazilian sky islands. *Botanical Journal of the Linnean Society*, *189*(2), 132–152.

Behling, H. (2002). South and southeast Brazilian grasslands during Late Quaternary times: a synthesis.
Palaeogeography, Palaeoclimatology, Palaeoecology, *177*(1-2), 19–27.

Berry, P. E., & Riina, R. (2005). Insights into the diversity of the Pantepui flora and the biogeographic
complexity of the Guayana Shield. *Biologiske Skrifter*, *55*, 145–167.

BFG. (2015). Growing knowledge: An overview of seed plant diversity in Brazil. *Rodriguésia*, *66*(4), 1085–
1113.

Bitencourt, C., & Rapini, A. (2013). Centres of endemism in the Espinhaço Range: Identifying cradles and
museums of Asclepiadoideae (Apocynaceae). *Systematics and Biodiversity*, *11*(4), 525–536.

Blackburn, T. M., & Gaston, K. J. (2002). Scale in macroecology. *Global Ecology and Biogeography*, *11*(3), 185–
189.

Bonatelli, I. A., Perez, M. F., Peterson, A. T., Taylor, N. P., Zappi, D. C., Machado, M. C., Koch, I., et al. (2014).
Interglacial microrefugia and diversification of a cactus species complex: Phylogeography and
palaeodistributional reconstructions for *pilosocereus aurisetus* and allies. *Molecular Ecology*, *23*(12), 3044–
3063.

Borba, E. L., & Semir, J. (1998). Bulbophyllum xcipoense (orchidaceae), a new natural hybrid from the
brazilian" campos rupestres": Description and biology. *Lindleyana*, *13*, 113–120.

Borcard, D., Gillet, F., & Legendre, P. (2018). *Numerical ecology with R.* Springer.

Brown, J. L., Paz, A., Reginato, M., Renata, C. A., Assis, C., Lyra, M., Caddah, M. K., et al. (2020). Seeing the forest
through many trees: Multi-taxon patterns of phylogenetic diversity in the Atlantic Forest hotspot. *Diversity*
and Distributions, *26*(9), 1160–1176.

Carstensen, D. W., Dalsgaard, B., Svenning, J.-C., Rahbek, C., Fjeldså, J., Sutherland, W. J., & Olesen, J. M.
(2012). Biogeographical modules and island roles: a comparison of Wallacea and the West Indies. *Journal of*
biogeography, *39*(4), 739–749.

Cavalcanti, Taciana B. (1995). *Revisão de* Diplusodon *Pohl (Lythraceae)* (PhD thesis). University of São Paulo.

Cavalcanti, Taciana Barbosa. (2007). Diversidade e distribuição em *Diplusodon* Pohl (Lythraceae).
Fontqueria, *55*(49), 397–404.

Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., & Ellison, A. M. (2014). Rarefaction and
extrapolation with Hill numbers: a framework for sampling and estimation in species diversity
studies. *Ecological monographs*, *84*(1), 45-67.

Collevatti, R. G., Rabelo, S. G., & Vieira, R. F. (2009). Phylogeography and disjunct distribution
in *Lychnophora ericoides* (Asteraceae), an endangered cerrado shrub species. *Annals of Botany*, *104*(4), 655–
664.

Colville, J. F., Beale, C. M., Forest, F., Altwegg, R., Huntley, B., & Cowling, R. M. (2020). Plant richness, turnover,
and evolutionary diversity track gradients of stability and ecological opportunity in a megadiversity center.
Proceedings of the National Academy of Sciences, *117*(33), 20027–20037.

Conceição, A. A., Rapini, A., Carmo, F. F. do, Brito, J. C., Silva, G. A., Neves, S. P., & Jacobi, C. M. (2016).
Rupestrian grassland vegetation, diversity, and origin. *Ecology and conservation of mountaintop grasslands*
in brazil (pp. 105–127). Springer.

Cortés, A. J., Garzón, L. N., Valencia, J. B., & Madriñán, S. (2018). On the causes of rapid diversification in the
páramos: isolation by ecology and genomic divergence in Espeletia. *Frontiers in plant science*, *9*, 1700.

Cortez, M. B. S., Folk, R. A., Grady, C. J., Spoelhof, J. P., Smith, S. A., Soltis, D. E., & Soltis, P. S. (2020). Is the age of
plant communities predicted by the age, stability and soil composition of the underlying landscapes? An
investigation of OCBILs. *Biological Journal of the Linnean Society*.

Cowling, R. M., Potts, A. J., Bradshaw, P. L., Colville, J., Arianoutsou, M., Ferrier, S., Forest, F., et al.
(2014). Variation in plant diversity in mediterranean-climate ecosystems: The role of climatic and
topographical stability. *Journal of Biogeography*, *42*(3), 552–564.

CRIA. (2020). Rede *species*Link. Published on the Internet <<http://http://www.splink.org.br/>>.

Croizat, L. (1962). Space, time, form: The biological synthesis. *Published by the author*.

Dapporto, L., Ramazzotti, M., Fattorini, S., Talavera, G., Vila, R., & Dennis, R. L. (2013). Recluster: An unbiased
clustering procedure for beta-diversity turnover. *Ecography*, *36*(10), 1070–1075.

Divya, B., Ramesh, B. & Karanth, K. P. (2021). Contrasting patterns of phylogenetic diversity across climatic
zones of western ghats: A biodiversity hotspot in peninsular india. *Journal of Systematics and Evolution*,
59(2), 240.

Drummond, C. S., Eastwood, R. J., Miotto, S. T., & Hughes, C. E. (2012). Multiple continental radiations and
correlates of diversification in Lupinus (Leguminosae): testing for key innovation with incomplete taxon
sampling. *Systematic biology*, *61*(3), 443–460.

Echternacht, L., Sano, P. T., Bonillo, C., Cruaud, C., Couloux, A., & Dubuisson, J.-Y. (2014). Phylogeny and
taxonomy of *Syngonanthus* and *Comanthera* (Eriocaulaceae): Evidence from expanded sampling. *Taxon*,
63(1), 47–63.

Echternacht, L., Trovó, M., Oliveira, C. T., & Pirani, J. R. (2011). Areas of endemism in the Espinhaço
Range in Minas Gerais, Brazil. *Flora-Morphology, Distribution, Functional Ecology of Plants*, *206*(9), 782–791.

Echternacht, L., Trovó, M., & Takeo Sano, P. (2010). Rediscoveries in Eriocaulaceae: seven narrowly
distributed taxa from the Espinhaço Range in Minas Gerais, Brazil. *Feddes Repertorium*, *121*(3-4), 117–126.

Emerson, B. C., & Gillespie, R. G. (2008). Phylogenetic analysis of community assembly and structure over
space and time. *Trends in ecology & evolution*, *23*(11), 619–630.

Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biological conservation*, *61*(1), 1–10.

Fernandes, G. W. (2016). The megadiverse rupestrian grassland. *Ecology and conservation of mountaintop*
grasslands in brazil (pp. 3–14). Springer.

Flantua, S. G., O'dea, A., Onstein, R. E., Giraldo, C., & Hooghiemstra, H. (2019). The flickering connectivity
system of the north Andean páramos. *Journal of Biogeography*, *46*(8), 1808–1825.

Forest, F., Grenyer, R., Rouget, M., Davies, T. J., Cowling, R. M., Faith, D. P., Balmford, A., et al. (2007).
Preserving the evolutionary potential of floras in biodiversity hotspots. *Nature*, *445*(7129), 757.

Garcia-Verdugo, C., Forrest, A. D., Balaguer, L., Fay, M. F., & Vargas, P. (2010). Parallel evolution of insular *Olea*
europaea subspecies based on geographical structuring of plastid DNA variation and phenotypic similarity in
leaf traits. *Botanical Journal of the Linnean Society*, *162*(1), 54–63.

García-Rodríguez, A., Martínez, P. A., Oliveira, B. F., Velasco, J. A., Pyron, R. A., & Costa, G. C. (2021). Amphibian
speciation rates support a general role of mountains as biodiversity pumps. *The American Naturalist*, *198*(3),
E68–E79.

GBIF.org. (2020). GBIF Home Page. Published on the Internet <<https://www.gbif.org>>.

Giulietti, A. M., Pirani, J. R., & Harley, R. M. (1997). Espinhaço Range region - eastern Brazil. *Centres of plant*
diversity: a guide and strategy for their conservation.

Gould, S. J., & Vrba, E. S. (1982). Exaptation—a missing term in the science of form. *Paleobiology*, *8*(1), 4–15.

Graham, C. H., & Fine, P. V. (2008). Phylogenetic beta diversity: Linking ecological and evolutionary processes
across space in time. *Ecology letters*, *11*(12), 1265–1277.

Gustafsson, A. L. S., Verola, C. F., & Antonelli, A. (2010). Reassessing the temporal evolution of orchids with
new fossils and a Bayesian relaxed clock, with implications for the diversification of the rare South American
genus *Hoffmannseggella* (Orchidaceae: Epidendroideae). *BMC Evolutionary Biology*, *10*(1), 1–13.

Harley, R. (1988). Evolution and distribution of *eriope* (Labiatae) and its relatives in Brazil. *Proceedings of a*
workshop on neotropical distributions patterns. Rio de janeiro, academia brasileira de ciências (pp. 71–120).

Heenan, P. B., Millar, T. R., Smissen, R. D., McGlone, M. S., & Wilton, A. D. (2017). Phylogenetic measures of
neo-and palaeo-endemism in the indigenous vascular flora of the New Zealand archipelago. *Australian*
Systematic Botany, *30*(2), 124–133.

Hopper, S. (2018). Natural hybridization in the context of OCBIL theory. *South African Journal of Botany*, *118*,
284–289.

Hopper, S. D. (2009). OCBIL theory: Towards an integrated understanding of the evolution, ecology and
conservation of biodiversity on old, climatically buffered, infertile landscapes. *Plant and Soil*, *322*(1-2), 49–86.

Hopper, S. D., Silveira, F. A., & Fiedler, P. L. (2016). Biodiversity hotspots and Ocbil theory. *Plant and Soil*,
403(1-2), 167–216.

Hopper, S. D., Lambers, H., Silveira, F. A., & Fiedler, P. L. (2021). OCBIL theory examined: reassessing
evolution, ecology and conservation in the world's ancient, climatically buffered and infertile
landscapes. *Biological Journal of the Linnean Society*, *133*(2), 266-296.

Hughes, C. E., & Atchison, G. W. (2015). The ubiquity of alpine plant radiations: from the Andes to the
Hengduan Mountains. *New Phytologist*, *207*(2), 275–282. doi:10.1111/nph.13230

Hughes, C. E., Pennington, R. T., & Antonelli, A. (2013). Neotropical plant evolution: Assembling the big
picture. *Botanical Journal of the Linnean Society*, *171*(1), 1–18.

Hughes, C., & Eastwood, R. (2006). Island radiation on a continental scale: Exceptional rates of plant
diversification after uplift of the Andes. *Proceedings of the National Academy of Sciences*, *103*(27), 10334–
10339.

Inglis, P. W., & Cavalcanti, T. B. (2018). A molecular phylogeny of the genus *Diplusodon* (Lythraceae),
endemic to the campos rupestres and cerrados of South America. *Taxon*, *67*(1), 66–82.

Jaccard, P. (1901). Distribution de la flore alpine dans le bassin des Dranses et dans quelques régions
voisines. *Bull Soc Vaudoise Sci Nat*, *37*, 241–272.

Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D., Blomberg, S. P., et al.
(2010). Picante: R tools for integrating phylogenies and ecology. *Bioinformatics*, *26*(11), 1463–1464.

Kreft, H., & Jetz, W. (2007). Global patterns and determinants of vascular plant diversity. *Proceedings of the
National Academy of Sciences*, *104*(14), 5925–5930.

Kubota, Y., Hirao, T., Fujii, S., & Murakami, M. (2011). Phylogenetic beta diversity reveals historical effects in
the assemblage of the tree floras of the Ryukyu Archipelago. *Journal of biogeography*, *38*(5), 1006–1008.

Linder, H. P. (2008). Plant species radiations: Where, when, why? *Philosophical Transactions of the Royal
Society B: Biological Sciences*, *363*(1506), 3097–3105.

Loeuille, B., Semir, J., Lohmann, L. G., & Pirani, J. R. (2015). A phylogenetic analysis of Lychnophorinae
(Asteraceae: Vernonieae) based on molecular and morphological data. *Systematic Botany*, *40*(1), 299–315.

Lososová, Z., Šmarda, P., Chytrỳ, M., Purschke, O., Pyšek, P., Sádlo, J., Tichỳ, L., et al. (2015). Phylogenetic
structure of plant species pools reflects habitat age on the geological time scale. *Journal of Vegetation Science*,
26(6), 1080–1089.

Lousada, J. M., Lovato, M. B., & Borba, E. L. (2013). High genetic divergence and low genetic variability in
disjunct populations of the endemic *Vellozia compacta* (Velloziaceae) occurring in two edaphic environments
of Brazilian campos rupestres. *Brazilian Journal of Botany*, *36*(1), 45–53.

Lozupone, C., & Knight, R. (2005). UniFrac: A new phylogenetic method for comparing microbial
communities. *Appl. Environ. Microbiol.*, *71*(12), 8228–8235.

Magdalena, U. R., Silva, L. A. E., Lima, R. O., Bellon, E., Ribeiro, R., Oliveira, F. A., Siqueira, M. F., et al. (2018). A
new methodology for the retrieval and evaluation of geographic coordinates within databases of scientific
plant collections. *Applied Geography*, *96*, 11–15.

McPeek, M. A. (2007). The macroevolutionary consequences of ecological differences among species.
Palaeontology, *50*(1), 111–129.

Mishler, B. D., Knerr, N., González-Orozco, C. E., Thornhill, A. H., Laffan, S. W., & Miller, J. T. (2014).
Phylogenetic measures of biodiversity and neo- and paleo-endemism in Australian *Acacia*. *Nature*
Communications, *5*, 4473.

Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global biodiversity
conservation: The critical role of hotspots. *Biodiversity hotspots* (pp. 3–22). Springer.

Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A., & Kent, J. (2000). Biodiversity hotspots for
conservation priorities. *Nature*, *403*(6772), 853–858.

Neves, D. M., Dexter, K. G., Pennington, R. T., Bueno, M. L., Miranda, P. L. de, & Oliveira-Filho, A. T. (2018). Lack
of floristic identity in *campos rupestres*—A hyperdiverse mosaic of rocky montane savannas in South
America. *Flora*, *238*, 24–31.

Nürk, N. M., Atchison, G. W., & Hughes, C. E. (2019). Island woodiness underpins accelerated disparification in
plant radiations. *New Phytologist*, *224*(1), 518–531.

Nürk, N. M., Linder, H. P., Onstein, R. E., Larcombe, M. J., Hughes, C. E., Piñeiro Fernández, L., Schlüter, P. M., et
al. (2020). Diversification in evolutionary arenas—assessment and synthesis. *Ecology and Evolution*, *10*(12),
6163–6182.

Oliveira, R. S., Abrahão, A., Pereira, C., Teodoro, G. S., Brum, M., Alcantara, S., & Lambers, H. (2016).
Ecophysiology of *campos rupestres* plants. *Ecology and conservation of mountaintop grasslands in brazil* (pp.
227–272). Springer.

Onstein, R. E., Carter, R. J., Xing, Y., & Linder, H. P. (2014). Diversification rate shifts in the cape floristic region:
The right traits in the right place at the right time. *Perspectives in Plant Ecology, Evolution and Systematics*,
16(6), 331–340.

Onstein, R. E., & Linder, H. P. (2016). Beyond climate: Convergence in fast evolving sclerophylls in cape and
australian rhamnaceae predates the mediterranean climate. *Journal of Ecology*, *104*(3), 665–677.

Pedreira, A., & De Waele, B. (2008). Contemporaneous evolution of the Palaeoproterozoic–Mesoproterozoic
sedimentary basins of the São Francisco–Congo Craton. *Geological Society, London, Special Publications*,
294(1), 33–48.

Polasky, S., Csuti, B., Vossler, C. A., & Meyers, S. M. (2001). A comparison of taxonomic distinctness versus
richness as criteria for setting conservation priorities for North American birds. *Biological Conservation*,
97(1), 99–105.

Queiroz, L. P. de, Cardoso, D., Fernandes, M. F., & Moro, M. F. (2017). Diversity and evolution of flowering
plants of the Caatinga Domain. *Caatinga* (pp. 23–63). Springer.

R Core Team. (2013). R: A language and environment for statistical computing.

Rafiqpoor, D., Kier, G., & Kreft, H. (2005). Global centers of vascular plant diversity. *Nova Acta Leopoldina NF*,
92(342), 61–83.

Rahbek, C. (2005). The role of spatial scale and the perception of large-scale species-richness patterns.
Ecology letters, *8*(2), 224–239.

Rahbek, C., Borregaard, M. K., Colwell, R. K., Dalsgaard, B., Holt, B. G., Morueta-Holme, N., Nogues-Bravo, D., et
al. (2019). Humboldt's enigma: What causes global patterns of mountain biodiversity? *Science*, *365*(6458),
1108–1113.

Rando, J. G., Zuntini, A. R., Conceição, A. S., Berg, C. van den, Pirani, J. R., & Queiroz, L. P. de. (2016). Phylogeny
of *Chamaecrista* ser. *Coriaceae* (Leguminosae) unveils a lineage recently diversified in Brazilian campo
rupestre vegetation. *International Journal of Plant Sciences*, *177*(1), 3–17.

Rapini, A. (2010). Revisitando as Asclepiadoideae (Apocynaceae) da Cadeia do Espinhaço. *Boletim de
Botânica*, *28*(2), 97–123.

Rapini, A., Berg, C. van den, & Liede-Schumann, S. (2007). Diversification of Asclepiadoideae (Apocynaceae) in
the New World. *Annals of the Missouri Botanical Garden*, *94*(2), 407–422.

Rapini, A., Bitencourt, C., Luebert, F., & Cardoso, D. (2020). An escape-to-radiate model for explaining the high
plant diversity and endemism in campos rupestres. *Biological Journal of the Linnean Society*.

Ribeiro, P. Luz, Borba, E. L., Camargo Smidt, E. de, Lambert, S. M., Schnadelbach, A. S., & Van den Berg, C.
(2008). Genetic and morphological variation in the *Bulbophyllum exaltatum* (Orchidaceae) complex
occurring in the Brazilian "campos rupestres": implications for taxonomy and biogeography. *Plant
Systematics and Evolution*, *270*(1), 109–137.

Ribeiro, P. L., Rapini, A., Damascena, L. S., & Van den Berg, C. (2014). Plant diversification in the Espinhaço
Range: insights from the biogeography of *Minaria* (Apocynaceae). *Taxon*, *63*(6), 1253-1264.

Richards, E. J., McGirr, J. A., Wang, J. R., John, M. E. S., Poelstra, J. W., Solano, M. J., O'Connell, D. C., et al. (2021).
A vertebrate adaptive radiation is assembled from an ancient and disjunct spatiotemporal landscape.
Proceedings of the National Academy of Sciences, *118*(20).

Rieseberg, L. H., & Willis, J. H. (2007). Plant speciation. *Science*, *317*(5840), 910–914.

Rodrigues, A. S., & Gaston, K. J. (2002). Maximising phylogenetic diversity in the selection of networks of
conservation areas. *Biological Conservation*, *105*(1), 103–111.

Rosauer, D., Laffan, S. W., Crisp, M. D., Donnellan, S. C., & Cook, L. G. (2009). Phylogenetic endemism: A new
approach for identifying geographic concentrations of evolutionary history. *Molecular Ecology*, *18*(19),
4061–4072.

Safford, H. D. (2007). Brazilian Páramos IV. Phytogeography of the campos de altitude. *Journal of
Biogeography*, *34*(10), 1701–1722.

Särkinen, T., Pennington, R. T., Lavin, M., Simon, M. F., & Hughes, C. E. (2012). Evolutionary islands in the
Andes: Persistence and isolation explain high endemism in andean dry tropical forests. *Journal of
Biogeography*, *39*(5), 884–900. doi:10.1111/j.1365-2699.2011.02644.x

Schaefer, Carlos E., Cândido, H. G., Corrêa, G. R., Nunes, J. A., & Arruda, D. M. (2016). Soils associated with
rupestrian grasslands. *Ecology and conservation of mountaintop grasslands in brazil* (pp. 55–69). Springer.

Scherson, R. A., Thornhill, A. H., Urbina-Casanova, R., Freyman, W. A., Pliscoff, P. A., & Mishler, B. D. (2017).
Spatial phylogenetics of the vascular flora of Chile. *Molecular phylogenetics and evolution*, *112*, 88–95.

Silveira, F. A., Negreiros, D., Barbosa, N. P., Buisson, E., Carmo, F. F., Carstensen, D. W., Conceição, A. A., et al.
(2016). Ecology and evolution of plant diversity in the endangered *campo rupestre*: A neglected conservation
priority. *Plant and soil*, *403*(1-2), 129–152.

Simon, M. F., Grether, R., Queiroz, L. P. de, Skema, C., Pennington, R. T., & Hughes, C. E. (2009). Recent
assembly of the cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire.
Proceedings of the National Academy of Sciences, *106*(48), 20359–20364.

Souza, É. R. de, Lewis, G. P., Forest, F., Schnadelbach, A. S., Berg, C. van den, & Queiroz, L. P. de. (2013).
Phylogeny of *calliandra* (Leguminosae: Mimosoideae) based on nuclear and plastid molecular markers.
Taxon, *62*(6), 1200–1219.

Spearman, C. (1904). The proof and measurement of association between two things. *American journal of*
Psychology, *15*(1), 72–101.

Tôrres, N. M., & Diniz-Filho, J. A. F. (2004). Phylogenetic autocorrelation and evolutionary diversity
of Carnivora (Mammalia) in conservation units of the New World. *Genetics and Molecular Biology*, *27*(4),
511–516.

Trovó, M., De Andrade, M. J. G., Sano, P. T., Ribeiro, P. L., & Van den Berg, C. (2013). Molecular phylogenetics
and biogeography of neotropical Paepalanthoideae with emphasis on Brazilian *Paepalanthus* (Eriocaulaceae).
Botanical Journal of the Linnean Society, *171*(1), 225–243.

Vasconcelos, T. N., Alcantara, S., Andrino, C. O., Forest, F., Reginato, M., Simon, M. F., & Pirani, J. R. (2020). Fast
diversification through a mosaic of evolutionary histories characterizes the endemic flora of ancient
neotropical mountains. *Proceedings of the Royal Society B*, *287*(1923), 20192933.

Warren, B. H., & Hawkins, J. A. (2006). The distribution of species diversity across a flora's component
lineages: Dating the cape's "relicts." *Proceedings of the Royal Society B: Biological Sciences*, *273*(1598), 2149–
2158.

Webb, C. O., Ackerly, D. D., McPeek, M. A., & Donoghue, M. J. (2002). Phylogenies and community ecology.
Annual review of ecology and systematics, *33*(1), 475–505.

Werneck, F. P., Costa, G. C., Colli, G. R., Prado, D. E., & Sites Jr, J. W. (2011). Revisiting the historical distribution
of Seasonally Dry Tropical Forests: new insights based on palaeodistribution modelling and palynological
evidence. *Global Ecology and Biogeography*, *20*(2), 272–288.

Werneck, F. P., Nogueira, C., Colli, G. R., Sites Jr, J. W., & Costa, G. C. (2012). Climatic stability in the brazilian
cerrado: Implications for biogeographical connections of south american savannas, species richness and
conservation in a biodiversity hotspot. *Journal of Biogeography*, *39*(9), 1695–1706.

Wiens, J. J., & Donoghue, M. J. (2004). Historical biogeography, ecology and species richness. *Trends in*
ecology & evolution, *19*(12), 639–644.

Willis, K. J., & Whittaker, R. J. (2002). Species diversity–scale matters. *Science*, *295*(5558), 1245–1248.

Yamagishi-Costa, J., & Forni-Martins, E. R. (2009). Hybridization and polyploidy: cytogenetic indications for
Hoffmannseggella (Orchidaceae) species evolution. *International Journal of Botany*.

Zachos, J. C., Dickens, G. R., & Zeebe, R. E. (2008). An early Cenozoic perspective on greenhouse warming and
carbon-cycle dynamics. *Nature*, *451*(7176), 279-283.

Zappi, D. C., Moro, M. F., Meagher, T. R., & Nic Lughadha, E. (2017). Plant biodiversity drivers in
Brazilian *campos rupestres*: insights from phylogenetic structure. *Frontiers in plant science*, *8*, 2141.

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., et al. (2019).
CoordinateCleaner: Standardized cleaning of occurrence records from biological collection databases.
Methods in Ecology and Evolution, *10*(5), 744–751.

Biosketch

Yago Barros-Souza is interested in the biogeography of neotropical vegetations and the role of morphological traits in plant evolution. This work was carried out as his master's thesis at the Universidade de São Paulo, Brazil. **Leonardo M. Borges** studies the taxonomy and evolution of *Mimosa* and related genera in the legume family. He is also interested in the theory and practice of ontology, homology, and morphological disparity of plants. Except for data collection and analyses, led by YB, both authors contributed equally to this work.

Editor: Carina Hoorn

richness

phylogenetic diversity

Mimosa

Diplusodon

Comanthera

Velloziaceae

Lychnophorinae

Paepalanthus

Calliandra

Minaria

Marcetieae

All groups

Campos rupestres at *Serra da Canastra*