Title:

Activation of ChvG-ChvI regulon by cell wall stress confers resistance to β-lactam

antibiotics and initiates surface spreading in *Agrobacterium tumefaciens*

Authors:

Michelle A. Williams*$, Jacob M. Bouchier*, Amara K. Mason, and Pamela J.B. Brown#

Affiliations:

Division of Biological Sciences, University of Missouri-Columbia, Columbia, Missouri,

65211

Running title: Cell wall stress activates the ChvG-ChvI pathway in *A. tumefaciens*

*denotes equal contributions

#**Corresponding author:**

brownpb@missouri.edu

(573) 884-0384

Present address:

$Boulevard Brewery, 2501 Southwest Blvd, Kansas City, MO 64108

ABSTRACT

A core component of nearly all bacteria, the cell wall is an ideal target for broad

spectrum antibiotics. Many bacteria have evolved strategies to sense and respond to

antibiotics targeting cell wall synthesis, especially in the soil where antibiotic-producing

bacteria compete with one another. Here we show that cell wall stress caused by both

chemical and genetic inhibition of the essential, bifunctional penicillin-binding protein

PBP1a prevents microcolony formation and activates the canonical host-invasion two-

component system ChvG-ChvI in *Agrobacterium tumefaciens*. Using RNA-seq, we

show that depletion of PBP1a for 6 hours results in a downregulation in transcription of

flagellum-dependent motility genes and an upregulation in transcription of type VI

secretion and succinoglycan biosynthesis genes, a hallmark of the ChvG-ChvI regulon.

Depletion of PBP1a for 16 hours, results in differential expression of many additional

genes and may promote a general stress response. Remarkably, the overproduction of

succinoglycan causes cell spreading and deletion of the succinoglycan biosynthesis

gene *exoA* restores microcolony formation. Treatment with cefsulodin phenocopies

depletion of PBP1a and we correspondingly find that *chvG* and *chvI* mutants are

hypersensitive to cefsulodin. This hypersensitivity only occurs in response to treatment

β-lactam antibiotics and moenomycin, suggesting that the ChvG-ChvI pathway may play

a key role in resistance to antibiotics targeting cell wall synthesis. Finally, we provide

evidence that ChvG-ChvI likely has a conserved role in conferring resistance to cell wall

stress within the Alphaproteobacteria.

AUTHOR SUMMARY

Soil dwelling bacteria reside in changing environments requiring them to frequently adapt to stressful conditions to ensure survival. The bacterial envelope provides structural integrity and protection against osmotic stress and turgor pressure imposed by the environment. While the mechanisms of cell membrane and cell wall biogenesis have been extensively studied, our understanding of how diverse microbes respond to cell envelope and cell wall stress to increase their fitness remains limited. In this work, we identify ChvG-ChvI regulon as an envelope stress response system that confers protection under cell wall stress conditions in the bacterial plant pathogen *Agrobacterium tumefaciens*. This is a new function for the well-characterized ChvG-ChvI pathway which is also acid induced and promotes plant host invasion. Our results suggest that the ChvG-ChvI pathway has a broadly conserved role in protecting Alphaproteobacterial cells from extracellular stress and a more specific role in response to acid stress and promoting plant-microbe interactions.

INTRODUCTION

The soil environment is constantly in flux and can undergo rapid changes in hydration, nutrient availability, temperature, acidity levels and many other abiotic and biotic factors [1]. To survive in these conditions, soil-dwelling bacteria must be able to monitor and respond to the changes around them. One of the main mechanisms bacteria employ to monitor changes in their environment is coupling environmental stimuli to transcriptional regulation using two-component systems (TCS) [2]. In turn transcriptional changes can modify bacterial behavior. In the plant-pathogen *Agrobacterium tumefaciens*, two TCS sense the presence of a potential host and initiate transcription programs that transition the bacterium into a virulent state [3].

The VirA-VirG histidine kinase/response-regulator pair is a characteristic of

Agrobacteria and responds to plant phenolic compounds such as acetosyringone.

Activation induces expression of the *vir* regulon, which encodes genes that are required

for pathogenicity and plant transformation [4]. The ChvG-ChvI TCS is more broadly

conserved across many Alphaproteobacteria but has been best characterized among

the plant symbionts of Rhizobiales such as *Sinorhizobium meliloti* [5,6]. Activation of

ChvG-ChvI is required for the transition from a free-living bacterium to a host-associated

lifestyle [7].

In *S. meliloti* and *A. tumefaciens* ChvG-ChvI is regulated by the periplasmic

protein ExoR. Under neutral conditions, ExoR binds to and represses ChvG; however,

when cells are exposed to acidic conditions, ExoR is proteolyzed, which allows for

activation of the ChvG-ChvI TCS [8,9]. ChvI induces transcriptional changes in many

genes across several major pathways. For example, ChvI upregulates transcription of

mirA, encoding a repressor of the motility response regulator Rem and ultimately

resulting in suppression of genes for motility and chemotaxis [10]. ChvI also upregulates

genes for exopolysaccharide production and, in *A. tumefaciens,* induction of the Type VI

Secretion System (T6SS) [11].

Conservation of the ChvG-ChvI TCS is taxonomically constrained to several

orders of Alphaproteobacteria, many of whom have free-living lifestyles that are never

host-associated [6]. This begs the question: why is the ChvG-ChvI pathway conserved

in so many non-host-associated bacteria? Recent interest in the ChvG-ChvI pathway of

Caulobacter crescentus (ChvGI) provides a glimpse at the function of the pathway in the

context of a bacterium with a drastically different ecological niche to that of *A.*

tumefaciens or *S. meliloti*. ChvGI of *C. crescentus* senses and responds to osmotic

stress and mutants of ChvGI are sensitive to several cell-wall targeting antibiotics

[12,13]. It remains unclear if this function is solely a characteristic of *C. crescentus*

ChvGI or if it is conserved across ChvG-ChvI orthologs.

Although the cell wall is an essential feature of bacteria that protects them from

environmental stressors, relatively little is known about how bacteria sense and respond

to changes in the composition of their cell wall. Peptidoglycan (PG) is a heteroglycan

decorated with cross-linked peptide stems and is the primary component of bacterial

cell walls. During elongation in *A. tumefaciens*, nascent PG insertion is constrained to

the pole. Polar growth is a characteristic of Rhizobiales and does not require the

canonical MreB-RodA-PBP2 elongation complex. Indeed, all members of Rhizobiales

have lost this complex entirely [14,15].

We showed that PBP1a is essential in *A. tumefaciens* and is the primary driver of

polar growth. Depletion of PBP1a eliminates nascent PG insertion at the growth pole,

leading to shorter cells that have compositional changes in PG [16]. In addition to its

role in polar PG insertion, here we observe that during PBP1a depletion cells spread

apart rather than form microcolonies. To better understand this phenotype, we used

RNA-seq to obtain transcriptional profiles of cells depleted of PBP1a after 6 hours,

corresponding to the onset of the spreading phenotype, and after 16 hours.

Transcriptomic changes closely mimic the transcriptome changes seen when ChvG-

ChvI is activated in *A. tumefaciens*, including downregulation of genes for motility and

chemotaxis and upregulation of genes for exopolysaccharide biosynthesis and T6SS.

Here we experimentally validate the RNA-seq results, confirming the impacts of PBP1a

depletion on the physiology and behavior of *A. tumefaciens*.

RESULTS

PBP1a depletion prevents proper microcolony formation

Here, we grew PBP1a depleted cells on agarose pads and saw that these cells

exhibit surface spreading rather than forming closely packed microcolonies (Fig 1A).

Additionally, when centrifuging cultures of PBP1a-depleted cells, we observed that the

cells did not pellet (Fig S1). Considering the possibility that depletion of PBP1a

somehow signals for these phenomena, we decided to look at RgsM, another enzyme

required for polar elongation. Previous work points to RgsM activity being required for

incorporation of nascent PG by PBP1a [17]. However, depletion of RgSM did not cause

surface spreading (Fig 1A) indicating that an imbalance of PG hydrolysis and synthesis

triggers spreading and the inability pellet in *A. tumefaciens*. Deletions of genes

encoding other high molecular weight PBPs and *mtgA*, a PG transglycosylase did not

induce spreading (Fig 1B).

Timelapse microscopy revealed that after ~6 hours of PBP1a depletion cells

spread apart, though the movement of cell appears to be confined within a relatively

small region of the agarose pad (Fig 1C, Movie S1). Since spreading is confined and

occurs over the course of many hours, we suspected that this phenomenon was not

simply caused by the activation of swimming motility.

PBP1a depletion induces global transcriptome changes

To understand the spreading phenotype caused by PBP1a depletion, we

compared the transcriptomes of cells at the onset or late stage of the surface spreading

phenotype. Cells were grown with or without the inducer Isopropyl β-D-1-

thiogalactopyranoside (IPTG) for *mrcA,* encoding PBP1a, expression for 6 or 16 hours

(Fig. 2A). As a baseline, we compared transcriptional profiles of WT in the presence and

absence of IPTG to the PBP1a depletion strain in the presence of IPTG. The addition of

IPTG did not alter gene expression profiles of WT cells, and only minor differences were

apparent between the PBP1a depletion strain background and WT when both strains

are grown in the presence of IPTG (Fig S2). We next compared differences in the

PBP1a replete strain to the PBP1a depleted strain at either 6- or 16-hours post

depletion (Fig 2A). Using a p-value < 0.05 and log2 fold-change (L2FC) > 2.0, we

identified 91 and 306 genes that were differentially expressed in the + PBP1a strain

compared to the 6- or 16-hour depletion, corresponding to 2% and 6% of the total

genes, respectively.

Overall, we observed large-scale changes in a diverse and widespread range of

genes that are regulated in response to PBP1a depletion. Initially, the response to

PBP1a depletion is primarily mediated by chromosomally encoded rather than plasmid

encoded genes. *A. tumefaciens* has a circular chromosome, which houses roughly half

(51.7%) of the protein-coding genes, a linear chromosome (34.7%) and two mega

plasmids, the At plasmid (10%) and Ti plasmid (3.6%). Most of the genes differentially

expressed at both time points during PBP1a depletion were encoded on the linear and

circular chromosomes (Fig. 2A). Most of the differentially abundant genes from the 6-

hour timepoint were also present in the 16-hour timepoint. For several of these genes,

the magnitude of differential transcript abundance remained relatively constant. For

example, the response regulator ChvI, had an increased relative abundance early in

response to PBP1a depletion that remained constant in the 16-hour timepoint. In

contrast, several genes displayed a continuous increase or decrease in transcript

abundance between the 6- and 16-hour timepoints, including genes that encode

proteins necessary for assembly of flagella and type 6 secretion system machinery.

Finally, several genes were only differentially abundant at the 16-hour timepoint,

including many genes encoding proteins important for cell envelope homeostasis such

as the Tol-Pal system [18], and >30 ABC transporters.

To further categorize the diverse set of differentially abundant genes we

identified Clusters of Orthologous Groups (COGs) in the 16-hour timepoint and

classified them based on functional categories represented by a single letter code

(Figure 2B) [19,20]. The most affected COG category was motility (N). Decreased

abundance of mRNAs containing genes which encode structural flagella proteins further

supports the hypothesis that the spreading phenotype is independent of flagella-based

motility. The COG category with the largest proportion of increased differentially

abundant genes was cell wall, membrane, and envelope biogenesis (M). Notably, no

significant changes in the transcripts of other penicillin-binding proteins or

glycosyltransferases were observed in response to loss of PBP1a (Table S1). However,

significant changes in transcripts encoding cell wall remodeling enzymes such as LD-

transpeptidases, endopeptidases, and soluble lytic transglycosylases were detected

(Table S1). Atu0844, an LD-transpeptidase, was strongly downregulated suggesting it

may play an important role in polar growth alongside PBP1a. Additionally, one putative

β-lactamase gene, Atu0933, was strongly upregulated, which may provide a strategy to

protect against cell wall damage. In addition, many of the genes found in this COG

category encode cell envelope homeostasis and osmotic stress response proteins,

including the Tol-Pal system, several outer membrane proteins (i.e. AopB), and

periplasmic sensors (i.e. CreD).

At the 16-hour timepoint, the largest changes to cellular metabolism occurred in

the inorganic ion transport and metabolism (P) and amino acid metabolism and

transport (E) COG categories, suggesting a shift in nutrient uptake and metabolism, as

expected during a general stress response. Interestingly, the six most downregulated

genes in the 16-hour timepoint, with the exception of *mrcA*, encoding PBP1a, were *yciC*

(Atu3181), *zinT* (Atu1049), *troC* (Atu3180), *troB* (Atu3179), *troA* (Atu3179), and

Atu3184, all of which are major components of cytoplasmic zinc uptake in

Agrobacterium tumefaciens (Fig 2B, bottom) [21].

We also observed large increases and decreases in the transcript abundance of

signal transduction genes. Transcription of *exoR* (Atu1715), *chvG* (Atu0033), and *chvI*

(Atu0034) were upregulated at both the 6-hour (L2FC = 0.995, 1.42, and 2.18) and 16-

hour (L2FC = 2.20, 1.41, and 2.27) depletions of PBP1a. Transcription of genes

encoding three other two-component systems were also strongly upregulated (Fig S3,

Table S1).

Transcriptome changes during PBP1a depletion mimic activation of the ChvG-

ChvI two-component system.

Transcription of *virG*, encoding a TCS response regulator, was also strongly

upregulated in both the 6-hour (L2FC = 2.61) and 16-hour (L2FC = 4.02) timepoints (Fig

S3, Table S1). Transcription of *virG* has been reported to be upregulated under both

host-invasion and virulence-inducing conditions [11]. Because *virG* was also

upregulated during depletion of PBP1a, we reasoned that PBP1a depletion may be

mimicking one of these two conditions. Using comparative transcriptomics, we

compared the 150 most differentially expressed genes (DEGs) against published

datasets that simulate host-invasion conditions (Δ*exoR* & pH 5.5) and virulence-

inducing conditions (acetosyringone treatment & growth on AB media) [22][23]. We

found that L2FC values of the 6-hour PBP1a depletion RNA-seq dataset correlated with

the two host-invasion conditions and not with the virulence-inducing datasets, as

indicated by the spearmen rho correlation coefficient for each comparison (Δ*exoR*, rho =

0.875; pH 5.5, rho = 0.766) (Fig 3 and S4). Rho values near 1 indicate similar DEGs

between each dataset. Rho values near 0 would indicate no similar DEGs between

each dataset (Fig 3A). Interestingly, each of these two datasets have been implicated in

activation of the ChvG-ChvI pathway [9,11]. Correlation with each strongly implicates

ChvG-ChvI activation in our RNA-seq, suggesting that depletion of PBP1a may provide

a signal leading to similar changes described to occur during the transition to a host-

invasion lifestyle. While this trend was maintained in the 16-hour timepoint, we observed

additional genes that were differentially expressed under depletion of PBP1a, but not in

the Δ*exoR* and pH 5.5 datasets (Fig 3, Fig S4B). Indeed, the rho values for the 16-hour

depletion of PBP1a compared to the host-invasion datasets (Δ*exoR*, rho = 0.529; pH

5.5, rho = 0.739 were lower than the 6-hour comparisons. Additionally, we found 217

more genes with L2FC > 2.0 in the 16-hour depletion than in the 6-hour depletion (Fig

3B). Together, these findings suggest that longer depletions of PBP1a may result in the

activation of additional regulons beyond ChvG-ChvI.

Overall, a large number and variety of genes are regulated in response to

depletion of PBP1a. Although many of these changes in gene expression have been

reported previously in response to low pH or deletion of the ChvG-ChvI negative

regulator ExoR, these changes have never been associated with loss of a cell wall

synthase in *A. tumefaciens*. These observations indicate that there are additional

mechanisms that can activate the ChvG-ChvI TCS.

Succinoglycan overproduction is required for cell spreading

Previous work has clearly associated activation of ChvG-ChvI to a specific

transcriptomic pattern involving downregulation of flagellar motility genes and

upregulation of T6SS and succinoglycan biosynthesis genes [9,11,24]. Indeed, this

same pattern was observed in the 6- and 16-hour PBP1a depletion datasets (Fig 4A).

To confirm that the spreading phenotype is unrelated to flagella-dependent motility, we

made an in-frame deletion of *rem*, which encodes a transcriptional regulator of genes

encoding structural flagella proteins [25,26], in the PBP1a depletion strain. Deletion of

rem prevents swimming in *A. tumefaciens* and does not impact microcolony formation

on agarose pads (Fig 4B). Upon depletion of PBP1a, *rem* mutants continued to spread,

suggesting that the cause of this phenotype is independent of flagella-mediated

swimming motility (Fig 4B).

All genes in the *imp* and *hcp* operons, which are located on the linear

chromosome and encode the structural and toxin proteins of T6SS respectively, are

upregulated at both timepoints (Fig 4A). In *A. tumefaciens*, activation of the T6SS

results in the production of a contractile nanomachine which delivers effector proteins to

antagonize and compete with other bacteria [27]. Among agrobacteria, T6SS is

activated by different signals, is important during different stages of the lifecycle, and

may be used to acquire nutrients [28]. To determine if the T6SS contributes to the

spreading response observed during PBP1a depletion we deleted the intergenic gap

between the *hcp* and *imp* operons (ΔT6SSpro). This deletion prevented expression of

proteins from both the *hcp* and *imp* operons (Fig S5) [29]. Cell spreading persisted in

ΔT6SSpro during depletion of PBP1a suggesting that the activation of T6SS is not

responsible for this behavior.

Another possibility is that spreading might be caused by sliding motility, where

secretion of a surfactant gives the cells a slippery surface to "slide" across. Notably, *S.*

meliloti has been reported to undergo entropy-driven surface spreading during the

overproduction of succinoglycan [30]. Succinoglycan is a β-1,4-linked sugar made of

glucose and galactose, and is the most abundant exopolysaccharide produced by *A.*

tumefaciens and related bacteria [31]. Genes associated with the biosynthesis and

secretion of succinoglycan were strongly upregulated in both timepoints. To test if

entropy-driven surface spreading is causing PBP1a-depleted *A. tumefaciens* cells to

spread, we used a microscopy-based assay to observe succinoglycan production in *A.*

tumefaciens. Cells were spotted on agarose pads containing calcofluor white and grown

overnight, then imaged using the DAPI filter to detect succinoglycan production (Fig

4C). Wild-type *A. tumefaciens* and the PBP1a replete strains secrete some

succinoglycan that enriched near the center of the microcolony (Fig 4C). In comparison,

depletion of PBP1a triggers secretion of a large quantity of succinoglycan that defines

the boundary of where the cells spread. An in-frame deletion of *exoA*, which encodes a

glycosyltransferase required for succinoglycan production in *A. tumefaciens* [32],

prevents succinoglycan production (Fig 4D). During PBP1a depletion, microcolony

formation is restored in the Δ*exoA* mutant (Fig 4D). Together, these data illustrate that

succinoglycan is indeed overproduced and contributes to the surface spreading of the

PBP1a depletion.

Deletion of *chvG* or *chvI* results in hypersensitivity to β-lactam antibiotics

Since activation of succinoglycan production is known to be part of the ChvG-

ChvI regulon, we next wanted to test if the PBP1a depletion is activating SGN

production through the ChvG-ChvI signaling pathway. We made an in-frame deletion of

chvI in the PBP1a depletion background and found *chvI* mutants repleted with PBP1a

appear morphologically wild-type when grown in minimal media (Fig S6A). However,

PBP1a-depleted Δ*chvI* cells were extremely sick. Previously we reported that the

PBP1a depletion produces viable daughter cells for up to 5-6 generations [16], however,

the PBP1a-depleted Δ*chvI* strain was incapable of a single division event. Instead, the

cells exhibited growth arrest and eventually cell lysis shortly after depletion initiation (Fig

S6A, Movie S2).

To further assess the enhanced sensitivity of Δ*chvI* to PBP1a depletion, we

identified an antibiotic that specifically targets PBP1a enzymatic activity. Treatment with

cefsulodin at a concentration of 20 µg/mL resulted in a spreading phenotype similar to

the PBP1a depletion (Fig 5A). Additionally, treatment with cefsulodin results in short,

round cells, further phenocopying the PBP1a depletion (Fig 5B). Together, these data

suggest that cefsulodin targets the transpeptidase activity of PBP1a in *Agrobacterium*

tumefaciens, similar to reported cefsulodin specificity in *E. coli* [33]. Next, we observed

relative cefsulodin sensitivities in WT, Δ*chvG*, Δ*chvI*, and Δ*exoR* strains. Remarkably,

we found that while growth of wild-type and Δ*exoR* cells appeared relatively unaffected

by treatment with 10µg/mL of cefsulodin, Δ*chvG* and Δ*chvI* cells were hypersensitive

(Fig 4C). Next, we targeted the glycosyltransferase activity of PBP1a through treatment

with moenomycin [34], and found that moenomycin-treated cells also spread on

agarose pads (Fig S6B). These findings suggest that the ChvG-ChvI TCS is essential

for growth when the activity of the major PG synthase is inhibited either chemically or

genetically.

We broadened our investigation by testing Δ*chvG* and Δ*chvI* against ten

additional antibiotics. four that block protein synthesis (chloramphenicol, tetracycline,

erythromycin, and gentamicin); one that blocks DNA replication (nalidixic acid); one that

blocks transcription (rifampin); and four other cell wall synthesis inhibiting β-lactam

antibiotics (meropenem, cefoxitin, ampicillin, and ticarcillin). To measure changes in

sensitivity to each antibiotic compared to wildtype, Δ*chvG* and Δ*chvI* were spread on

ATGN minimal media and disks containing each antibiotic were applied. Diameters of

the zones of inhibition (ZOI) were measured and the difference in ZOIs for each mutant

strain compared to wildtype are shown (Fig 5D). Of the antibiotics tested, Δ*chvG* and

Δ*chvI* showed increased sensitivity only to β-lactam antibiotics, suggesting specificity of

the ChvG-ChvI pathway in conferring resistance to this antibiotic class.

ChvG and ChvI are conserved in Alphaproteobacteria but the presence of ExoR is

more constrained

The absence of PBP1a activity at the growth pole during elongation activates

ChvG-ChvI, the canonical host-invasion pathway of *Agrobacterium tumefaciens*. The

ChvG-ChvI pathway is most well known to be activated by environmental changes

associated with conditions favorable for plant association, yet this TCS is retained in

many non-plant-associated Alphaproteobacteria (Fig 6A). Remarkably, while ChvG-

ChvI is conserved in a large proportion of Alphaproteobacteria, ExoR is not (Fig 6A).

Predicted structures of the sensor domains of ChvG in bacteria with ChvG-ChvI

orthologs show two structural loops (L1 and L2; Fig 6A, Fig S7). While L2 is conserved

across the orthologous structures, L1 is expanded solely in the Rhizobiales (Fig S8).

This expansion coincides with the retention of ExoR, making it a compelling target for

ExoR-ChvG association studies (Fig 6A). The more broadly conserved L2, on the other

hand, may enable a conserved, ExoR-independent mechanism for activation of ChvG-

ChvI.

We hypothesized that the ChvG-ChvI pathway may confer resistance to cell wall

stress in other host-associated Rhizobiales. Indeed, depletion of PBP1a in

Sinorhizobium meliloti causes cells to spread suggest activation of ChvG-ChvI (ExoS-

ChvI) pathway (Fig 6B). Like *A. tumefaciens*, this response is specific to depletion of

PBP1a and is not triggered by deletion of the 5 other high molecular weight PBPs.

These results suggest that cell wall stress may be a well conserved trigger for activation

of ChvG-ChvI pathways in the Rhizobiales. Notably, *C. crescentus* does not spread

upon treatment with 80 μg/mL of cefsulodin (Fig S6C), despite recent findings that

ChvGI likely confers resistance at this concentration [12]. While *S. meliloti* and *A.*

tumefaciens encode succinoglycan biosynthesis operons, *C. crescentus* does not.

These findings support the hypothesis that the ChvG-ChvI response to cell wall stress is

conserved across Alphaproteobacterial species that have orthologs of ChvG-ChvI.

DISCUSSION

Why is ChvG-ChvI TCS activated upon inhibition of polar PG synthesis in *A.*

tumefaciens? PBP1a depletion results in a compositional shift in cell wall muropeptide

composition and cell wall crosslinking [16]. Here we find that PBP1a depletion causes

upregulation in transcription of LD-transpeptidases, endopeptidases, and soluble lytic

transglycosylases, indicating cells are attempting to compensate for compromised cells

walls. Remarkably, transcripts from these same genes are upregulated in the Δ*exoR*

and pH 5.5 datasets, suggesting that remodeling of the cell wall is a part of the ChvG-

ChvI regulon. While the role of cell wall remodeling during host-invasion is unclear it is

possible that these modifications may be protective for the bacterium when host

associated. The upregulation in transcription of genes encoding AopB and the Tol/Pal

system may indicate that PBP1a-depleted cells are succumbing to osmotic pressure, a

possible explanation for the increase in cell width in PBP1a-depleted and cefsulodin-

treated cells (Fig 5B). Notably, this concept aligns well with recent findings in

Caulobacter crescentus where it was shown that ChvGI activation was required for

osmoregulation [35]. Together, these findings indicate a conserved role within the

Alphaproteobacteria for the ChvG-ChvI TCS in sensing and responding to envelope

stress. Possible signals may include the accumulation of cytoplasmic peptidoglycan

precursors such as lipid II, increased levels of naked glycan strands in the cell wall, the

production of peptidoglycan hydrolysis products, or more conventional stress responses

due to osmotic sensitivity [36]. Future investigations will be necessary to determine if

ChvG-ChvI proteins themselves directly sense and respond to cell wall stress or if other

regulators which, presumably interact with the ChvG-ChvI are involved in this response.

If there is a conserved role in sensing cell wall stress, why would this pathway be

required for host invasion within *Agrobacterium tumefaciens*? One explanation could be

that during plant colonization, *A. tumefaciens* decreases cell wall biogenesis to form

cells which are relatively persistent in order to evade host recognition and survive the

harsh *in plantae* environment [37]. Alternatively, perhaps the integrity and composition

of the cell wall is routinely monitored and used as a signal for the activation of pathways

associated with lifestyle choice. Decreased activity of PBP1a may mimic one or more of

the conditions *A. tumefaciens* and *S. meliloti* encounter during host-invasion, leading to

the activation of ChvG-ChvI. Alternatively, perhaps the absence of PBP1a activity leads

to a destabilization of the polar growth complex leading to decreased cell envelope

integrity. It will be of interest to determine if other components of the polar growth

complex such as GPR [38], RgsP [39], or PopZ [40,41] have increased sensitivity to β-

lactam antibiotics and induce surface spreading. This possibility is in agreement with the

observation in *C. crescentus* that resistance to cell wall targeting antibiotics is

dependent on factors such as TipN that maintain the integrity of the cell envelope [42].

The overproduction of succinoglycan provides cells with passive protection

against several stresses *A. tumefaciens* may encounter during host invasion including

detergents, salt, acidity, heat, antimicrobial peptides, and reactive oxygen species [33-

36]. Production of succinoglycan may also help protect against cell-wall-synthesis

targeting antibiotics produced by competing bacteria and fungi in the soil. However,

succinoglycan production is taxonomically constrained within plant-host-associated

bacteria, indicating that there are other conserved mechanisms regulated by ChvG-ChvI

in resistance to these antibiotics. The surface spreading phenomenon that we

connected to overproduction of succinoglycan and that has been previously described

in *S. meliloti* [30], may be involved in cell dispersal during host invasion.

The increased sensitivity to mecillinam, vancomycin, cefsulodin, and

moenomycin in ChvGI mutants suggests that this pathway may confer resistance to

antibiotics inhibiting cell wall synthesis [35]. Our observation that the depletion of *A.*

tumefaciens RgsM mutant does not induce surface spreading further hints that

increased cell wall hydrolysis may be responsible for activation of the ChvG-ChvI

pathway. While our findings suggest a conserved signal in ChvG-ChvI activation, further

studies will need to be conducted to identify the signal(s), determine if they are species-

specific, and explore the conditions which lead to signal production.

Longer depletions of PBP1a may induce a more general stress response. While

previous reports propose *A. tumefaciens* to have stress-specific responses instead of a

single general response [43] , the similarities between the transcriptional changes at the

16-hour time point and the general stress responses of other bacteria are striking [44].

For example, the general stress response of *Escherichia coli*, regulated by the ECF

family sigma factor σ^E, upregulates transcription of genes encoding ABC transporters

associated with uptake of amino acids, peptides, phosphate sources, and polyamines

[45]. Indeed, the 16-hour time point shows differential expression of over 30 ABC

transporters. However, previous studies found no common response to stresses

including osmotic stress, heat shock, and low pH in *A. tumefaciens* [43]. While there are

several ECF family sigma factors encoded in the *A. tumefaciens* genome, it is unclear

what, if any, role they may be playing in the transcriptional changes exhibited under

long depletions of PBP1a.

Together our findings support a model that ChvG-ChvI can be acid induced by

dissociation and proteolysis of ExoR in a Rhizobiales specific response (Fig 6C, left)

and through a more broadly conserved cell envelope stress response (Fig 6C, middle).

The role of ExoR regulation is well established for the acid-induction of ChvG-ChvI but

the lack of ExoR conservation suggests that the cell envelope stress response may be

ExoR-independent. Finally, longer periods of PBP1a depletion results in a more general

stress response, affecting transcription of genes associated with nutrient uptake and

metabolism (Fig 6C, right). Overall, the data presented here are in agreement with

recent works in *Caulobacter* [35,42] which suggest that ChvGI activation is important in

oligotrophic free-living bacteria as a cell envelope or osmotic stress response.

Remarkably, it seems that the ChvG-ChvI pathway has a dual purpose in protecting the

bacterium and invading its host in *A. tumefaciens*, and other host-associated

Rhizobiales.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. A list of all bacterial strains and

plasmids used in this study is provided in Table S2. *Agrobacterium tumefaciens* C58

and derived strains were grown in ATGN minimal media [46] without exogenous iron at

28°C with shaking. When appropriate, kanamycin (KAN) was used at the working

concentration of 300 µg/ml. When indicated, isopropyl β-D-1-thio-galactopyranoside

(IPTG) was used as an inducer at a concentration of 1 mM, and cumate. *Sinorhizobium*

meliloti stains were grown in TY medium [47] at 28°C. When appropriate, KAN was

used at the working concentration of 200 µg/ml, gentamycin (GM) was used at 20

µg/ml, and IPTG was used at a concentration of 500 µg/ml. *C. crescentus* strains were

grown in PYE medium [48] at 28°C. *E. coli* DH5α and S17-1 λ pir were grown in Luria-

Bertani medium at 37°C and when appropriate 50 µg/ml or 30 µg/ml of KM were added,

respectively.

Construction of deletion/depletion plasmids and strains. A list of all primers and

synthetic DNAs used in this study is provided in Table S3. Vectors for gene deletion by

allelic exchange were constructed using recommended methods for *A. tumefaciens*

[49]. Briefly, 500-bp fragments upstream and 500 bp downstream of the target gene

were amplified using primer pairs P1/P2 and P3/4 respectively. Amplicons were spliced

together by SOEing using primer pair P1/P4. The amplicon was digested and ligated

into pNTPS139. The deletion plasmids were introduced into *A. tumefaciens* by mating

using an *E. coli* S17 conjugation strain to create KM resistant, sucrose sensitive primary

integrants. Primary integrants were grown overnight in media with no selection.

Secondary recombinants were screened by patching for sucrose resistance and KM

sensitivity. Colony PCR with primers P5/P6 for the respective gene target was used to

confirm deletion. PCR products from P5/P6 primer sets were sequenced to further

confirm deletions.

Phase and fluorescence microscopy. A small volume (~1 µl) of cells in exponential

phase (OD600 = 0.2 - 0.4) was applied to a 1% ATGN agarose pad as described

previously [50]. DIC, Phase contrast and epifluorescence microscopy were performed

with an inverted Nikon Eclipse TiE and a QImaging Rolera em-c2 123 1K EMCCD

camera with Nikon Elements Imaging Software. For time-lapse microscopy, images

were collected every ten minutes, unless otherwise stated. For calcofluor agar pad

assays, calcofluor was added to agarose pads at a concentration of 25 µg/mL and

exposed to DAPI filter for 50ms. When appropriate agar pads were supplemented with

1mM IPTG. For quantitative image analysis, live cells were imaged using phase-

contrast microscopy, and cell length and width distributions of the indicated number of

cells per strain were determined as measured using MicrobeJ software [51]. T-tests

were performed using the compare_means() function of the ggpubr R library.

RNA isolation, sequencing and analysis. Four cultures each of WT, WT + IPTG and

12 cultures of + PBP1a depletion cells were grown overnight in 2 ml of ATGN minimal

media at 28°C with shaking; the + PBP1a strains and WT + IPTG strains were

supplemented with 1mM IPTG. Cells were then pelleted by centrifugation at 7000 x g for

5 minutes. Cell pellets were washed three times with ATGN by centrifugation and

resuspension to remove IPTG. After the final wash the cell pellets from WT, WT + IPTG,

and four of the 12 + PBP1a strains were resuspended to an OD600 of 0.05 in 6mL

ATGN, or ATGN with 1mM IPTG. The other eight + PBP1a strains were resuspended to

an OD600 of 0.05 in 6 ml ATGN without IPTG. This resulted in 4 replicate cultures each

of WT, WT +IPTG, + PBP1a, - PBP1a_6hr and - PBP1a_16hr. Growth of the cultures

was monitored and supplemented with fresh medium as needed so that the OD600

never went over 0.3. RNA was isolated from the -PBP1a_6hr strains after 6 hours of

growth, and RNA was isolated from all other strains after 16 hours of growth. To

prepare samples, a culture volume equivalent to 6 ml at an optical density at 600 nm

(OD600) of 0.2-0.3 was pelleted by centrifugation at 7000 x g for 5 minutes and pellets

were resuspended in 1mL of ATGN media and incubated with 2 mL of RNAProtect

reagent (QIAgen) for 15 min at room temperature. Cells were lysed with 10 mg

lysozyme, and RNA was extracted using the QIAgen RNEasy kit.

DNA libraries for sequencing were constructed following the manufacturer's protocol

with reagents supplied in Illumina's TruSeq mRNA stranded sample preparation kit. The

sample concentration was determined by Qubit flourometer (Invitrogen) using the Qubit

HS RNA assay kit, and the RNA integrity was checked using the Fragment Analyzer

automated electrophoresis system. Briefly, the poly-A containing mRNA is purified from

total RNA, RNA is fragmented, double-stranded cDNA is generated from fragmented

RNA, and the index containing adapters are ligated to the ends. The amplified cDNA

constructs were purified by addition of Axyprep Mag PCR Clean-up beads. The final

construct of each purified library was evaluated using the Fragment Analyzer automated

electrophoresis system, quantified with the Qubit flourometer using the Qubit HS dsDNA

assay kit, and diluted according to Illumina's standard sequencing protocol for

sequencing on the NextSeq 500.

For all samples, when adapter sequence was detected, it was removed using cutadapt

(0.16) [52]. All samples were purged of reads that mapped to transcripts for rRNA genes

using bowtie2 (2.3.4.3) [53]. The reads were then mapped to the *A. fabrum* str. C58

genome using STAR (version 2.5.4b) [54], which also produces the number of read

counts per gene. The index files used by STAR were derived from the files

*Agrobacterium*_fabrum_str_c58.ASM9202v1.dna.toplevel.fa and

*Agrobacterium*_fabrum_str_c58.ASM9202v1.40.gtf, both of which are part of Ensembl

release 40 (http://bacteria.ensembl.org/index.html). Pairwise comparisons were

performed to test for differential expression of genes using the Bioconductor package

DESeq2 [55]. Gene annotations were collected from the annotations included with the

file of cDNAs also at Ensembl

*Agrobacterium*_fabrum_str_c58.ASM9202v1.cdna.all.fa.gz.

COG functional annotation. Amino acid sequences for all proteins in *A. tumefaciens*

were downloaded in a single FASTA file from GenBank and uploaded to EGGNOG-

MAPPER [56,57]. COG terms were outputted, and Python code was written to pull out

transcripts from 16-hour depletion of PBP1a with L2FC > 2.0 or < -2.0. Some transcripts

had multiple COG annotations and were therefore replicated for visualization according

to the number of annotations it had.

Comparative transcriptomics. Transcripts and L2FC values from each dataset were opened in Python code written to screen for and exclude any genes that were not present in both datasets. Statistics and visualization was done in R. Spearman correlation statistical test was run on the L2FC of the 150 most differentially expressed genes in the PBP1a depletion and their corresponding L2FC values in the comparison dataset.

Cell viability assays. For cell viability spot assays, cultures were grown overnight and diluted to an OD_{600} = 0.05 and serially diluted in ATGN and spotted on ATGN agar plates containing antibiotics as indicated. Four microliters of each dilution was spotted and plates were incubated at 28°C for 48 h before imaging.

Disk diffusion assays. Wild-type, Δ*chvG*, and Δ*chvI* cells were overnight and then knocked down to an OD_{600} of 1.0. Cells were then lawned on ATGN minimal media. Sterile paper disks either soaked in concentrations of each antibiotic or not (blank controls) were applied to the plate. Each plate was grown for ~48 hours at 28°C before being imaged. Zone of inhibition diameters were measured from each image using ImageJ software.

Phylogenetics and structure prediction. A seed of 22 amino acid sequences containing the annotated ChvG sensor domain (PF13755) were initially downloaded from Pfam [58]. Each was blasted against its corresponding proteome to retrieve the full protein sequence [59]. Additional sequences of relevant bacteria such as *S. meliloti,*

Brucella melitensis, and *C. crescentus,* were added by blasting the amino acid

sequence from *A. tumefaciens* ChvG (Atu0033) against each organism's proteome. All

sequences were aligned using MUSCLE and trimmed in Jalview according to Uniprot

predicted periplasmic region of Atu0033 [60–62]. A maximum parsimony phylogenetic

tree of these sequences was generated using MEGA-X [63].

Each trimmed sequence underwent one-to-one threading in Phyre2 with the complete

structure of Atu0033 predicted by AlphaFold as a template [64,65]. Local alignment and

a secondary structure weight of 0.1 was used. Structural analysis and structure

alignment was done in ChimeraX [66].

The amino acid sequence of *A. tumefaciens* ExoR (Atu1715) was blasted against each

organism's proteome and max score values of top hits were recorded. Max score values

under 50 were deemed too different and were therefore not considered an ExoR

ortholog. Additionally, sequences of each top hit were blasted against the proteome of

A. tumefaciens. If the top hit was not ExoR, it was also not considered an ExoR ortholog

in this analysis.

DATA AVAILABILITY

Raw RNA-seq read files (.fastq) and complete list of differentially expressed genes for each

comparison are publicly available through the NCBI Gene Expression Omnibus (GEO) under

the accession GSE173921.

ACKNOWLEDGEMENTS

We thank Elizaveta Krol and Anke Becker for providing *Sinorhizobium meliloti* strains and

micrographs and Erh-Min Lai for providing antibodies used in this work. We thank Amelia

Randich, Alex Quintero, Regis Hallez and members of the Brown lab for critical feedback on this

manuscript. This work was supported by the National Science Foundation, IOS1557806,

to PJBB. MAW and JMB were supported by the Life Sciences Fellowship at the

University of Missouri and AKM was supported as an IMSD/MARC fellow with support

from the National Institutes of Health, T34 GM136493.

The molecular graphics and analyses were performed with UCSF ChimeraX, developed

by the Resource for Biocomputing, Visualization, and Informatics at the University of

California, San Francisco, with support from National Institutes of Health R01-

GM129325 and the Office of Cyber Infrastructure and Computational Biology, National

Institute of Allergy and Infectious Diseases.

REFERENCES

1.   Venturi V, Keel C. Signaling in the Rhizobiales. Trends Plant Sci. 2016;21: 187–
198.
2.   Capra EJ, Laub MT. Evolution of two-component signal transduction systems.
Annu Rev Microbiol. 2012;66: 325–347.
3.   Yuan Z-C, Haudecoeur E, Faure D, Kerr KF, Nester EW. Comparative
transcriptome analysis of *Agrobacterium tumefaciens* in response to plant signal
salicylic acid, indole-3-acetic acid and γ-amino butyric acid reveals signalling
cross-talk and *Agrobacterium*-plant co-evolution. Cell Microbiol. 2008;10: 2339–
2354.
4.   Cho H, Winans SC. VirA and VirG activate the Ti plasmid *repABC* operon,
elevating plasmid copy number in response to wound-released chemical signals.
Proc Natl Acad Sci U S A. 2005;102: 14843–14848.
5.   Xue S, Biondi EG. Coordination of symbiosis and cell cycle functions in
Sinorhizobium meliloti. Biochim Biophys acta Gene Regul Mech. 2019;1862: 691–
696.
6.   Heavner ME, Qiu W-G, Cheng H-P. Phylogenetic co-occurrence of ExoR, ExoS,
and ChvI, components of the RSI bacterial invasion switch, suggests a key

adaptive mechanism regulating the transition between free-living and host-invading phases in Rhizobiales. Mergaert P, editor. PLoS One. 2015;10: e0135655.

7. Wells DH, Chen EJ, Fisher RF, Long SR. ExoR is genetically coupled to the ExoS-ChvI two-component system and located in the periplasm of *Sinorhizobium meliloti*. Mol Microbiol. 2007;64: 647–664.

8. Lu HY, Luo L, Yang MH, Chenga HP. *Sinorhizobium meliloti* ExoR is the target of periplasmic proteolysis. J Bacteriol. 2012;194: 4029–4040.

9. Yuan ZC, Liu P, Saenkham P, Kerr K, Nester EW. Transcriptome profiling and functional analysis of *Agrobacterium tumefaciens* reveals a general conserved response to acidic conditions (pH 5.5) and a complex acid-mediated signaling involved in *Agrobacterium*-plant interactions. J Bacteriol. 2008;190: 494–507.

10. Alakavuklar MA, Heckel BC, Stoner AM, Stembel JA, Fuqua C. Motility control through an anti-activation mechanism in *Agrobacterium tumefaciens*. Mol Microbiol. 2021;116: 1281–1297.

11. Heckel BC, Tomlinson AD, Morton ER, Choi JH, Fuqua C. *Agrobacterium tumefaciens* ExoR controls acid response genes and impacts exopolysaccharide synthesis, horizontal gene transfer, and virulence gene expression. J Bacteriol. 2014;196: 3221–3233.

12. Quintero-Yanes A, Mayard A, Hallez R. The two-component system ChvGI maintains cell envelope homeostasis in *Caulobacter crescentus*. bioRxiv. 2022; 2022.01.18.476748.

13. Vallet SU, Hansen LH, Bistrup FC, Laursen SA, Chapalay JB, Chambon M, et al. Loss of bacterial cell pole stabilization in *Caulobacter crescentus* sensitizes to outer membrane stress and peptidoglycan-directed antibiotics. MBio. 2020;11.

14. Brown PJB, De Pedro MA, Kysela DT, Van Der Henst C, Kim J, De Bolle X, et al. Polar growth in the Alphaproteobacterial order Rhizobiales. PNAS. 2012;109: 1697-1701.

15. Margolin W. Sculpting the Bacterial Cell. Curr Biol. 2009;19: R812.

16. Williams MA, Aliashkevich A, Krol E, Kuru E, Bouchier JM, Rittichier J, et al. Unipolar Peptidoglycan Synthesis in the Rhizobiales Requires an Essential Class A Penicillin-Binding Protein. 2021;12.

17. Figueroa-Cuilan WM, Randich AM, Dunn CM, Santiago-Collazo G, Yowell A, Brown PJB. Diversification of LytM in and of *Agrobacterium tumefaciens*. Front Microbiol. 2021;12: 2341.

18. Krol E, Yau HCL, Lechner M, Schäper S, Bange G, Vollmer W, et al. Tol-pal system and proteins interact to promote unipolar growth and cell division in *Sinorhizobium meliloti*. MBio. 2020;11: 1–21.

19. Tatusov RL, Koonin E V., Lipman DJ. A genomic perspective on protein families. Science. 1997;278: 631–637.

20. Galperin MY, Kristensen DM, Makarova KS, Wolf YI, Koonin E V. Microbial genome analysis: the COG approach. Brief Bioinform. 2019;20: 1063.

21. Chaoprasid P, Dokpikul T, Johnrod J, Sirirakphaisarn S, Nookabkaew S, Sukchawalit R, et al. *Agrobacterium tumefaciens* Zur regulates the high-affinity zinc uptake system TroCBA and the putative metal chaperone YciC, along with ZinT and ZnuABC, for survival under zinc-limiting conditions. Appl Environ

Microbiol. 2016;82: 3503.

22. Yuan Z-C, Edlind MP, Liu P, Saenkham P, Banta LM, Wise AA, et al. The plant
signal salicylic acid shuts down expression of the *vir* regulon and activates
quormone-quenching genes in *Agrobacterium*. 2007;104: 11790-11795

23. Haryono M, Cho ST, Fang MJ, Chen AP, Chou SJ, Lai EM, et al. Differentiations
in gene content and expression response to virulence induction between two
Front Microbiol. 2019;10: 1554.

24. Wu CF, Lin JS, Shaw GC, Lai EM. Acid-induced Type VI Secretion System is
regulated by ExoR-ChvG. PLoS Pathog. 2012;8.

25. Figueroa-Cuilan W, Daniel JJ, Howell M, Sulaiman A, Brown PJB. Mini-Tn7
insertion in an artificial attTn7 site enables depletion of the essential master
regulator CtrA in the phytopathogen *Agrobacterium tumefaciens*. Appl Environ
Microbiol. 2016;82: 5015–5025.

26. Rotter C, Mühlbacher S, Salamon D, Schmitt R, Scharf B. Rem, a new
transcriptional activator of motility and chemotaxis in *Sinorhizobium meliloti*. J
Bacteriol. 2006;188: 6932.

27. Wu CF, Smith DA, Lai EM, Chang JH. The *Agrobacterium* Type VI Secretion
System: a contractile nanomachine for interbacterial competition. 2018;418: 215–
231.

28. Wu CF, Weisberg AJ, Davis EW, Chou L, Khan S, Lai EM, et al. Diversification of
the Type VI Secretion System in Agrobacteria. MBio. 2021;12.

29. Lin JS, Ma LS, Lai EM. Systematic Dissection of the *Agrobacterium* Type VI
Secretion System reveals machinery and secreted components for subcomplex
formation. PLoS One. 2013;8: e67647.

30. Dilanji GE, Teplitski M, Hagen SJ. Entropy-driven motility of *Sinorhizobium meliloti*
on a semi-solid surface. Proc R Soc B Biol Sci. 2014;281.

31. Mendis HC, Madzima TF, Queiroux C, Jones KM. Function of succinoglycan
polysaccharide in *Sinorhizobium meliloti* host plant invasion depends on
succinylation, not molecular weight. MBio. 2016;7.

32. Schmid J, Sieber V, Rehm B. Bacterial exopolysaccharides: Biosynthesis
pathways and engineering strategies. Front Microbiol. 2015;6: 496.

33. Curtis NAC, Orr D, Ross GW, Boulton MG. Affinities of penicillins and
cephalosporins for the penicillin-binding proteins of *Escherichia coli* K-12 and their
antibacterial activity. Antimicrob Agents Chemother. 1979;16: 533.

34. Yuan Y, Fuse S, Ostash B, Sliz P, Kahne D, Walker S. Structural analysis of the
contacts anchoring moenomycin to peptidoglycan glycosyltransferases and
implications for antibiotic design. 2008;3: 429–436.

35. Quintero-Yanes A, Coppine J, Mayard A, Hallez R. The two-component system
ChvGI maintains cell envelope homeostasis in *Caulobacter crescentus*. bioRxiv.
2022.

36. Piepenbreier H, Diehl A, Fritz G. Minimal exposure of lipid II cycle intermediates
triggers cell wall antibiotic resistance. Nat Commun 2019 101. 2019;10: 1–13.

37. Erbs G, Silipo A, Aslam S, De Castro C, Liparoti V, Flagiello A, et al.
Peptidoglycan and muropeptides from pathogens *Agrobacterium* and
Xanthomonas elicit plant innate immunity: Structure and Activity. Chem Biol.
2008;15: 438–448.

38. Zupan JR, Grangeon R, Robalino-Espinosa JS, Garnica N, Zambryski P.
GROWTH POLE RING protein forms a 200-nm-diameter ring structure essential
for polar growth and rod shape in *Agrobacterium tumefaciens*. Proc Natl Acad Sci
U S A. 2019;166: 10962–10967.
39. Schäper S, Yau HCL, Krol E, Skotnicka D, Heimerl T, Gray J, et al. Seven-
transmembrane receptor protein RgsP and cell wall-binding protein RgsM
promote unipolar growth in Rhizobiales. PLOS Genet. 2018;14: e1007594.
40. Howell M, Aliashkevich A, Salisbury AK, Cava F, Bowman GR, Brown PJB.
Absence of the Polar Organizing Protein PopZ Results in Reduced and
Asymmetric Cell Division in *Agrobacterium tumefaciens.* Journal of Bacteriology.
2017;199.
41. Grangeon R, Zupan J, Jeon Y, Zambryski PC. Loss of PopZAt activity in
Agrobacterium tumefaciens by deletion or depletion leads to multiple growth
poles, minicells, and growth defects. MBio. 2017;8.
42. Vallet SU, Hansen LH, Bistrup FC, Laursen SA, Chapalay JB, Chambon M, et al.
Loss of bacterial cell pole stabilization in *Caulobacter crescentus* sensitizes to
outer membrane stress and peptidoglycan-directed antibiotics. MBio. 2020;11.
43. Rosen R, Büttner K, Schmid R, Hecker M, Ron EZ. Stress-induced proteins of
Agrobacterium tumefaciens. FEMS Microbiol Ecol. 2001;35: 277–285.
44. Hecker M, Völker U. General stress response of Bacillus subtilis and other
bacteria. Adv Microb Physiol. 2001;44: 35–91.
45. Hayden JD, Ades SE. The extracytoplasmic stress factor, σE, is required to
maintain cell envelope integrity in *Escherichia coli*. PLoS One. 2008;3: e1573.
46. Morton ER, Fuqua C. UNIT 3D.1 Laboratory Maintenance of *Agrobacterium*. Curr
Protoc Microbiol. 2012.
47. Beringer JE. R factor transfer in *Rhizobium leguminosarum*. J Gen Microbiol.
1974;84: 188–198.
48. Poindexter P, J. BIOLOGICAL PROPERTIES AND CLASSIFICATION OF THE
CAULOBACTER GROUP. Bacteriol Rev. 1964;28: 231.
49. Morton ER, Fuqua C. Genetic manipulation of *Agrobacterium*. Curr Protoc
Microbiol. 2012;Chapter 3.
50. Howell M, Daniel JJ, Brown PJB. Live cell fluorescence microscopy to bbserve
essential processes during microbial cell growth. J Vis Exp. 2017;2017: 56497.
51. Ducret A, Quardokus EM, Brun Y V. MicrobeJ, a tool for high throughput bacterial
cell detection and quantitative analysis. Nat Microbiol. 2016;1.
52. Martin M. Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet.journal. 2011;17: 10–12.
53. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat
Methods. 2012;9: 357–359.
54. Dobin A, Gingeras TR. Mapping RNA-seq reads with STAR. Curr Protoc
Bioinforma. 2015;51: 11.14.1-11.14.19.
55. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15: 1–21.
56. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al.
Database resources of the national center for biotechnology information. Nucleic
Acids Res. 2022;50: D20–D26.

57. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook
H, et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated
orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids
Res. 2019;47: D309.
58. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL,
et al. Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49:
D412–D419.
59. Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, Mcginnis S, Madden TL.
NCBI BLAST: a better web interface. Nucleic Acids Res. 2008;36: 5–9.
60. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 2004;32: 1792.
61. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version
2—a multiple sequence alignment editor and analysis workbench. Bioinformatics.
2009;25: 1189–1191.
62. Bateman A, Martin MJ, Orchard S, Magrane M, Agivetova R, Ahmad S, et al.
UniProt: the universal protein knowledgebase in 2021. Nucleic Acids Res.
2021;49: D480–D489.
63. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary
genetics analysis across computing platforms. Mol Biol Evol. 2018;35: 1547–
1549.
64. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. The Phyre2 web
portal for protein modeling, prediction and analysis. Nat Protoc. 2015;10: 845–
858.
65. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly
accurate protein structure prediction with AlphaFold. Nat 2021 5967873.
2021;596: 583–589.
66. Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al.
UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. Protein Sci. 2021;30: 70–82.

SUPPORTING INFORMATION

Movie S1. Growth and morphological changes during 8 hours of PBP1a depletion. Cells
were washed to remove inducer and spotted immediately on an ATGN pad. Images
were acquired every ten minutes and movie is played at 16 frames per second for a
total of 48 frames.

Movie S2. Growth and division of D*chvI* during PBP1a depletion. Cells were washed to
remove inducer and spotted immediately on an ATGN pad. Images were acquired every
five minutes and movie is played at 40 frames per second for a total of 200 frames.

Fig S1. Pelleting of PBP1a-depleted and repleted cells. Conical tubes show turbidity
after pelleting cells grown for 16 hours in PBP1a replete (+PBP1a) or depleted (-PBP1a)
conditions. Cells were centrifuged at 1690 x g (3000 rpm in TX-400 rotor in a Sorvall
Legend X1R centrifuge) for 10 minutes. Supernatants were spotted on a 1.25% ATGN
agarose pad. $\bar{x}$ = average number of cells from 10 fields of view.

Fig S2. Analysis of the control transcriptomes by RNA-seq. A. Plots comparing Log2Fold Change of the WT 6-hour transcriptome to that of the WT +ITPG 6-hour transcriptome. Gray dots represent a single transcript, and the dotted lines represent +/- 2.0 Log2Fold Change threshold. Plots are delimited by chromosomes and mega plasmids. B. Plots comparing Log2Fold Change of the WT +IPTG transcriptome to that of the PBP1a depletion strain with ITPG present to drive PBP1a expression. Comparisons shown are of the 6-hour transcriptomes. Gray dots represent a single transcript, and the dotted lines represent +/- 2.0 Log2Fold Change threshold. Plots are delimited by chromosomes and mega plasmids.

Fig S3. Transcriptional changes of TCS regulators and kinases during PBP1a depletion. The fold change in expression level of TCS regulators and kinases are shown following 6 hours (gray) and 16 hours (black) of PBP1a depletion. The *virAG* and *chvGI* TCS pairs are labeled.

Fig S4. The response to the depletion of PBP1a mimics transcriptional changes associated with host invasion. A. Correlation scatterplots depicting relationships between the log2fold-change (L2FC) values in the 16-hour PBP1a depletion and transcriptomic data sets taken under simulated virulence-inducing conditions (AS) and under simulated host-invading conditions (Δ*exoR*). Each point represents a unique transcript. AS, acetosyrinogone; Rho, Spearman correlation coefficient. B. Correlation scatterplots comparing L2FC values of transcripts in the pH 5.5 microarray, a condition known to induce the *chvG-chvI* regulon, to either the 6-hour (red) or 16-hour (blue) PBP1a depletion. Rho, Spearman correlation coefficient.

Fig S5. Western blot of proteins expressed from the two type VI secretion system operons in ΔT6SSpro strains. Top panel, diagram of the two operons encoding elements of Type VI Secretion in *A. tumefaciens.* T6SSpro labels the intergenic gap that is deleted in ΔT6SSpro strains. Middle panel, western blots using anti-Hcp and anti-TssB in each of the indicated strains. Protein sizes (kDa) are shown on the right. Bottom panel, Coomassie stained gel showing total protein from each strain.

Fig S6. Impact of decreased PG synthesis on *A. tumefaciens* and *C. crescentus* microcolony formation. A. Micrographs of PBP1a depletion Δ*chvl* with (+PBP1A) or without (-PBP1A) IPTG. Cells was grown to exponential phase in ATGN media containing IPTG, spotted on an ATGN agar pad with or without IPTG, allowed to grow for 16 hours, and imaged by DIC microscopy. B. Micrographs of WT *A. tumefaciens* cells growth with or without moenomycin. Cells was grown to exponential phase in ATGN media, spotted on an ATGN agar pad with or without moenomycin, allowed to grow for 16 hours, and imaged by DIC microscopy. C. Micrographs of WT *C. crescentus* cells growth with or without cefsulodin. Cells were grown to exponential phase in PYE media, spotted on a PYE agar pad with or without cefsulodin, allowed to grow for 16 hours, and imaged by DIC microscopy. All scale bars depict 2µm.

Fig S7. Alignment of periplasmic regions of ChvG orthologs. Partial MUSCLE
alignment of ChvG ortholog periplasmic domains. Highlighted columns represent strong
conservation across aligned sequences. Atu0033 (ChvG of *A. tumefaciens)* is the
reference sequence for this analysis. L1 and L2 correspond to two conserved structural
loops. Conservation, quality, and consensus scores for each site are represented as bar
graphs under the alignment. Shading indicates order of the bacterium containing the
ChvG ortholog: Orange, Rhizobiales; Purple, Rhodobacterales; Green,
Caulobacterales; Gold, Rhodospirales.
842
Fig S8. Structure predictions for the periplasmic regions of ChvG orthologs.
Phyre2 structural predictions for each organism displayed in Figure 6A of this work.
Genus and species names as well as locus tags for each ChvG ortholog are provided.
Range of numbers following the back slash are the amino acid sites used in structure
prediction. Colors indicates order of the bacterium containing the ChvG ortholog:
Orange, Rhizobiales; Purple, Rhodobacterales; Green, Caulobacterales; Gold,
Rhodospirales.
850
851   **Table S1. Selected differentially expressed genes**
852
853   **Table S2. Bacterial strains and plasmids.**
854
855   **Table S3. Synthesized DNA primers.**
856
857   **Supplementary Methods**
858
859   **Supplementary References**

860
Fig 1. The PBP1a depletion fails to form microcolonies independent of flagellar
motility. A. Micrographs of wildtype, PBP1a depletion, and RgsM depletion with or
without 1mM IPTG inducer. Each strain was grown to exponential phase, spotted
on an ATGN agar pad, allowed to grow for 16 hours, and imaged by DIC
microscopy. Scale bar depicts 2µm. The graphic depicts the working model that
RgsM $_{DD}$-endopeptidase activity is required for incorporation of nascent glycan strands
into the preexisting peptidoglycan (PG) macromolecule by PBP1a. i) RgsM cleaves $_{DD}$-
crosslinks, ii) PBP1a glycosyltransferase activity incorporates lipid II into the PG glycan
strand, iii) PBP1a $_{DD}$-transpeptidase activity crosslinks the peptide stem of the nascent
PG, fully incorporating it into the macromolecule. B. Micrographs of wild type, Δ*pbp1b1*,
Δ*pbp1b2*, Δ*pbp1c*, and Δ*mtgA*. Each strain was grown to exponential phase, spotted on
an ATGN agar pad, allowed to grow for 16 hours, and imaged by DIC microscopy.
Scale bar depicts 2µm. C. Time-lapse microscopy of the PBP1a depletion grown on an
agar pad with or without 1mM IPTG inducer. DIC images were acquired every 10
minutes. Time is shown in hours. For the - PBP1a strain, cells were washed 3X with
ATGN media and grown at 28 C with shaking for 4 hours before cells were spotted on
an agar pad for imaging.

Fig 2. Analysis of the PBP1a depletion transcriptomes by RNA-seq. A. Plots
comparing Log2Fold Change of the + PBP1a transcriptome to that of the - PBP1a 6-
hour transcriptome and to that of the 16-hour depletion. Gray dots represent a single
transcript, and the dotted lines represent +/- 2.0 Log2Fold Change threshold. Plots are
delimited by chromosome and mega plasmid. B. COG categorical analysis of the 16-
hour depletion of PBP1a. Pink, downregulated; Cyan, upregulated.

886
Fig 3. The response to the depletion of PBP1a mimics transcriptional changes
associated with host invasion. A. Correlation scatterplots depicting relationships
between the log2fold-change (L2FC) values in the 6-hour PBP1a depletion and
transcriptomic data sets taken under simulated virulence-inducing conditions (AS and
AB+AS) and under simulated host-invading conditions (ΔexoR). Each point represents a
unique transcript. AS, acetosyrinogone; AB, *Agrobacterium* minimal media; rho,
Spearman correlation coefficient. B. Correlation scatterplots comparing L2FC values of
transcripts in the ΔexoR microarray to either the 6-hour or 16-hour PBP1a depletion.
Each transcript is colored according to its change in L2FC values from 6 hours of
PBP1a depletion to 16 hours of depletion. Gray, no change; Blue, transcript has |L2FC|
> 2.0 in the 6-hour but not in the 16-hour depletion; Red, |L2FC| > 2.0 in the 16-hour but
not in the 6-hour depletion; Purple, |L2FC| > 2.0 in both the 6-hour and 16-hour
depletion.
900
901

902
Fig 4. Succinoglycan overproduction is a conserved response to PBP1a depletion
and results in failed microcolony formation. A. Scatter plots depicting RPKM values
of the 6-hour and 16-hour compared to wild type. Each point represents a unique
transcript. Points are colored by category. Gold, ChvG-ChvI regulon; Lavender, Motility
and Chemotaxis; Green, Type VI Secretion; Blue, Succinoglycan Biosynthesis; Black,
mrcA (encoding PBP1a). B. Micrographs of wild type, Δ*rem*, PBP1a replete Δ*rem*,
PBP1a depleted Δ*rem*, ΔT6SSpro, PBP1a replete ΔT6SSpro, and PBP1a depleted
ΔT6SSpro. Each strain was grown to exponential phase, spotted on a 1% ATGN agar
pad containing 1mM IPTG if inducing *mrcA*, allowed to grow for 16 hours, and imaged
by DIC microscopy. The scale bar depicts 2µm. C. Micrographs of wild type, and PBP1a
depletion with or without IPTG inducer. Each strain was grown to exponential phase and
spotted on a 1% ATGN agar pad containing 25µg/mL calcofluor white and 1mM IPTG if
inducing *mrcA*. Each was allowed to grow for 16 hours and imaged by phase
microscopy with and without the DAPI filter for visualizing calcofluor-stained
succinoglycan. D. Micrographs of Δ*exoA* and PBP1a depletion Δ*exoA*, with or without
IPTG inducer. Strains were grown and imaged as described for panel C.

920
Fig 5. The ChvG-ChvI TCS is conditionally essential under treatment with β-
lactam antibiotics. A. Micrographs of untreated and cefsulodin-treated cells. Wild-type
cells were grown to exponential phase, spotted on a 1% ATGN agar pad with or without
20 μg/mL cefsulodin and allowed to grow for 16 hours. Each strain was imaged by DIC
microscopy. B. Box plots comparing cell length and width between wild-type, PBP1a-
depleted, and cefsulodin-treated cells. ns, not significant; ****, p < 0.00005. C. Cell
viability of each wild type, Δ*chvG,* Δ*chvI,* and Δ*exoR* spotted on an ATGN agar plate
containing 10 μg/mL of cefsulodin. Ten-fold serial dilutions are indicated. D. Graph
depicting the change in zone of inhibition from wildtype in Δ*chvG* and Δ*chvI* against ten
different antibiotic disks. Error bars represent +/- 1 standard deviation from the mean.
931

932
Fig 6. Conservation constraints of ExoR suggest conserved ChvG-ChvI response
is independent of ExoR. A. Maximum parsimony tree constructed using MUSCLE
sequence alignment [60] on the periplasmic regions of ChvG orthologs. In clades that
don't have a ChvG ortholog, the protein with the highest sequence similarity to ChvG
was used instead. Conservation of ExoR was calculated using blast max scores from
top hits when protein blasting [59] ExoR from *Agrobacterium tumefaciens* against each
species in the tree. Phyre2 predicted structures of periplasmic domains of ChvG
orthologs from representatives (bold) in each genus are shown. Conserved structural
loops are denoted as L1 and L2. B. Micrographs of *Sinorhizobium meliloti* Rm2011
PBP1a replete, PBP1a depleted, and a strain with deletions of genes encoding all other
high molecular weight PBPs (Δ5pbp). Each strain was grown to exponential phase,
spotted on a 1% TY agar pad containing 1mM IPTG if inducing *mrcA*, allowed to grow
for 16 hours, and imaged with phase microscopy. C. Working model of activation of
ChvG-ChvI in *A. tumefaciens*. H^+, free proton representing an acidic environment; β, β-
lactam antibiotic.