The evolution of manipulative cheating

Ming Liu[1*], Stuart A. West[1], Geoff Wild[2]

[1]Department of Biology, University of Oxford

[2]Department of Mathematics, University of West Ontario

Abstract

A social cheat is typically assumed to be an individual that does not perform a

cooperative behaviour, or performs less of it, but can still exploit the cooperative

behaviour of others. However, empirical data suggests that cheating can be more

subtle, involving evolutionary arms races over the ability to both exploit and resist

exploitation. These complications have not been captured by evolutionary theory,

which lags behind empirical studies in this area. We bridge this gap with a mixture of

game-theoretical models and individual-based simulations, examining what

conditions favour more elaborate patterns of cheating. We found that as well as

adjusting their own behaviour, individuals can be selected to manipulate the

behaviour of others, which we term 'manipulative cheating'. Further, we found that

manipulative cheating can lead to dynamic oscillations (arms races), between

selfishness, manipulation, and suppression of manipulation. Our results can help

explain both variation in the level of cheating, and genetic variation in the extent to

which individuals can be exploited by cheats.

Keywords: phenotypic diversity, tragedy of the commons, social evolution, public

goods, dynamic oscillation.

[*]Correspondence: ming.liu@ biology.ox.ac.uk

Introduction

Many social interactions involve a trade-off between investing in behaviours or traits that increase the productivity of the group, versus those that increase an individual's share of group productivity (Frank 1998; Bourke 2011; Queller & Strassmann 2018)(Fig. 1a). A female social insect larva could become a queen or a worker (Hamilton 1972). A slime mould cell could become a stalk or a spore within a fruiting body (Strassmann *et al.* 2000). A bacterial cell could invest costly resources into producing some 'public good' that benefits the local group of cells, such as iron scavenging siderophore molecules (Griffin *et al.* 2004).

A standard assumption is that individuals can behave more selfishly by investing less in cooperative behaviours that benefit the group, and by taking a larger share of the available resources (Ghoul *et al.* 2014). For example, a slime mould lineage could 'cheat' by having a larger proportion of its cells develop into spore cells, or a bacterial cell could cheat by producing fewer siderophores (Strassmann *et al.* 2000; Jiricny *et al.* 2010). When cheats can invade, theory predicts that this will lead to either uniform cooperation at a lower level or the coexistence of cheats and cooperators (Queller 1984; Frank 1994, 1998; West & Buckling 2003; MacLean & Gudelj 2006; Ross-Gillespie *et al.* 2007; Gore *et al.* 2009; Frank 2010c; Frank 2010a, b; Patel *et al.* 2019). Coexistence is only expected when some factor leads to the fitness of cheats being higher when they are less common (frequency-dependent selection).

In contrast, the empirical data suggests that there can be more complex forms of cheating and resistance to cheating within social groups. For example, in the slime *Dictyostelium discoideum*: (i) lineages appear to be able to evolve resistance to cheating, where they prevent other lineages from obtaining a higher proportion of spore cells (Buttery *et al.* 2009; Khare *et al.* 2009; Levin *et al.* 2015); (ii) genomic data suggests an ongoing evolutionary arms race, and not just selection towards an optimal level of cooperation (Ostrowski *et al.* 2015; Noh *et al.* 2018; Ostrowski 2019). In some ant species, genetic variation appears to be maintained in the extent to which different patrilines contribute to the production of workers in a colony, a phenomenon termed 'royal cheating' (Hughes *et al.* 2003; Linksvayer 2006; Anderson *et al.* 2008; Hughes & Boomsma 2008; Mitchell *et al.* 2012; Stürup *et al.* 2014). In *Pseudomonas* bacteria, strains appear to vary in both the extent to which they can cheat by exploiting the siderophores produced by other strains and the extent to which they can be cheated by other strains (Kümmerli *et al.* 2009; Bruce *et al.* 2017; Butaitė *et al.* 2017). In all of these cases, cheating appears to involve doing more than just cooperating at a lower level. Other examples of more complex forms of

cheating can be found in bacteria, viruses, and replicator RNA (Pollak *et al.* 2016;
Furubayashi *et al.* 2020; Bruce *et al.* 2021; Mizuuchi *et al.* 2022).

Our aim is to explain how these more complex forms of cheating can be
maintained in a population. As mentioned above, most evolutionary models assume
that an individual can control their own investment into selfish and cooperative
behaviours – put simply, an individual cheats by investing less in cooperation (Queller
1984; MacLean & Gudelj 2006; Ross-Gillespie *et al.* 2007; Gore *et al.* 2009; Frank
2010c; Patel *et al.* 2019). We investigate another possibility, that individuals can
manipulate the behaviour of others. For example, an ant patriline may increase the
likelihood that its own larvae are raised as queens, while reducing the likelihood that
the larvae of other patrilines are reared. Or a bacteria cell lineage could increase the
public goods produced by another lineage. Such 'manipulative cheating' could
involve a variety of mechanisms, ranging from coercive communication to more
direct influencing. In addition, we explore how once manipulation has evolved, there
could be selection to evolve resistance or suppression of that cheating.

We develop a series of models to examine whether manipulative cheating can
be favoured, and if it will lead to variation in the extent to which individuals can
exploit others. We use a mixture of game theory and simulation, to examine
scenarios where resistance can evolve relatively simply, or where there are multiple
mechanisms of manipulative cheating that each require a different form of
resistance. Our model assumes an asexual haploid population, but we also examine
the consequences of recombination. We find that natural selection can maintain a
variety of strategies at equilibrium, as well as the possibility for ongoing oscillating
dynamics, such as evolutionary arms races. While our focus is on 'manipulation', our
more general aim is to investigate scenarios where cheating is more than just
'cooperate less'.

Methods
Model overview
Our aim is to investigate the general feasibility of manipulation, in a way that could
be applicable across a range of species, and does not rely on the specifics of any
single species. Consequently, we have chosen a deliberately simple approach to
maintain tractability. We use a combination of modelling approaches to examine the
evolutionary stability and dynamics of manipulative cheating.

First, we use an equilibrium game theoretical approach to examine under what

conditions manipulative cheating could be favoured. Our model is based on the
classic tragedy of the commons model (Frank 1994, 1996, 1998; Dionisio & Gordo
2006; Frank 2010a, b), where individuals can invest in their own selfish reproduction,
or the productivity of the group (Fig. 1a; Appendix 1—1.2). We extend this model by
assuming that: (i) individuals can invest effort into manipulating the behaviour of
others to make them cooperate at a higher rate (behave less selfishly; Fig. 1b); and
(ii) individuals can also invest effort into suppressing (blocking) the extent to which
they are manipulated by others (Fig. 1c). We then test the robustness of our game-
theoretical results by comparing them with individual-based simulations.
115
Second, we examine the consequence of manipulative cheating when multiple
different mechanisms of manipulation can evolve, which each requires a different
mechanism to suppress. We consider cases where traits are introduced at the same
time point or different time points (mutation) and focus on the temporal trait
dynamics. As multiple mechanisms could generate complications between traits
which cannot be easily presented by equilibrium analysis, we focus on dynamic
results from individual-based simulations.
123
Our general aim is to investigate scenarios where cheating can be more than
just 'cooperate less'. We do this by developing models focused on the special case of
manipulation and its suppression. Other possibilities would include developing
coevolutionary genetic models to look at genotype-specific cheating-resistance,
analogous to gene-for-gene or matching alleles models of host-parasite coevolution,
or to model a different form of coercion (Sasaki 2000; Agrawal & Lively 2002).
130
Model design
We assume a haploid and asexual population inhabiting an infinite number of
identical habitat patches, where each patch supports a single social group and lets
social interactions take place. There is no recombination, but our model is equivalent
to one that assumes a sexual diploid with additive interactions between alleles at a
given locus, with alleles at different loci segregating independently (Taylor 1996; Day
& Taylor 1998). We also assume the fitness of each individual is proportional to the
productivity of the social group multiplied by the individual's share of group
productivity, and all offspring compete globally for patches in the next generation.
Within each social group, each individual divides its effort into cooperation with
group members, which aids group productivity or selfishly reproducing (gaining a
greater share of the group productivity). We consider three coevolving traits:
143

Figure 1. Schematic view of the three evolving traits in our model. Each individual can put
effort into three actions: (a) invest resources into reproduction (z_i; represented by stripe texture)
or group productivity (average cooperation; $1 - \bar{z}$); (b) manipulate others (x_i; represented by
the grid texture), to make them invest more into group productivity; and (c) suppress (block)
the manipulation from others (u_i; represented by horizontal dashed texture).

Selfishness (z): We assume that when individuals put more effort into

reproduction (z), they: (i) obtain a larger share of the group productivity; (ii) put less

effort into cooperation, which reduces group productivity. Specifically, if a focal

individual i invests z_i, into reproduction, and the group average of all individuals in

the group is $\bar{z}$ ($0 < z < 1$). In other words, the parameter z measures the level of

'selfishness' (Fig. 1a)(Frank 1994, 1996, 1998; Dionisio & Gordo 2006; Frank 2010a,

b).

Manipulative cheating (x): We assume that a focal individual can invest x in

manipulation, which causes other individuals to invest fewer resources into their

own reproduction, and instead invest those resources into the reproduction of the

focal individual (manipulator; $0 < x < 1$; Fig. 1b). This manipulation requires effort

and so reduces the resources that the manipulator contributes to group productivity

(by $\bar{x}$). Manipulation x therefore takes reproduction from others, whereas

selfishness z shifts an individual's allocation between cooperation and

reproduction. Consequently, manipulation is redistributing the share of all

individual's reproduction while z is changing the effort spent in reproduction. For

simplicity, we have assumed there is no 'loss' in reproductive effort during the

process of manipulation, but we found similar results when manipulation also leads

to a personal reproduction cost (Appendix 1—3).

Suppression of manipulative cheating (u): We add the possibility for individuals

to suppress (block) manipulation ($0 < u < 1$; Fig. 1c). When suppression exists, a

manipulator can only take a proportion $1 - u^b$ of reproduction, where *b* is a shape

parameter (see below). Like the other two traits, suppression takes effort and

reduces the effort put into group productivity (by $\bar{u}$).

Combining the effects of all three traits lead to:

180
$$w_i = \frac{z_i + \left(1 - u_{-i}^b\right)x_i^a z_{-i} - \left(1 - u_i^b\right)x_{-i}^a z_i}{\bar{z}}(1 - \bar{z} - \bar{x} - \bar{u}),$$

181   (1)

where z_i is the selfishness of the focal individual, $\left(1 - u_{-i}^b\right)x_i^a z_{-i}$ is the gain from

manipulating others, $\left(1 - u_i^b\right)x_{-i}^a z_i$ is the loss from being manipulated, and $1 -$

$\bar{z} - \bar{x} - \bar{u}$ is the group productivity. Because the sum of gain and loss across all

individuals in the group is zero, the average effort in reproduction is just $\bar{z}$. Thus, the

first term is the relative share of group productivity of the focal individual (i.e.,

$\frac{z_i + \left(1 - u_{-i}^b\right)x_i^a z_{-i} - \left(1 - u_i^b\right)x_{-i}^a z_i}{\bar{z}}$). In other words, our manipulative cheating trait involves a

individual an actor causing a recipient (other group members) to behave less selfishly

(more cooperatively), in a way that is beneficial to the actor, but costly to the

recipient. This contrasts with previous theory which examined the consequences of

changing the benefit and cost of social behaviour (Frank 1995), or when relatives are

manipulated to perform a social behaviour that provides an indirect benefit to the

actor (González-Forero & Gavrilets 2013)(Appendix 1—1.3). We provide a more

detailed derivation of equation 1 in Appendix 1—1.1. In the special case of no

manipulation or suppression, equation 1 simplifies to the classic tragedy of the

commons model (Frank 1994, 1996, 1998; Dionisio & Gordo 2006; Frank 2010a, b).

We allow the effects of manipulation and suppression to be nonlinear by the

shape parameters, a and b. Nonlinearities can arise for a number of reasons, such

as mechanisms of molecular diffusion, or interference between the behaviour of

different individuals (Müller *et al.* 2006; Wiley 2013; Rogers *et al.* 2017). The

parameter a determines whether the benefit from increased investment into

manipulation is decelerating ($a < 1$), linear ($a = 1$) or accelerating ($a > 1$). Similarly,

the shape parameter b determines the shape of the benefit of increased

investment into suppression. We have focused on the decelerating forms because

this will often be the case for biological traits (Appendix 1—5.1).

We find it useful to think about manipulation, suppression, and equation 1 with

a lottery-ticket metaphor. Imagine reproduction as the buying of a lottery ticket –

producing an offspring, that has some chance of 'winning' (reproducing) in the next

generation. Individuals can either invest resources into buying the lottery tickets

produced by the group (z; selfish), or invest resources into helping the group

produce more lottery tickets ($1 - z$; cooperation). Each group produces its own

lottery tickets and more tickets means a higher chance for the focal group to win.

Being more selfish (higher z) means obtaining a larger fraction of the lottery tickets

produced by the group, but also putting fewer resources into producing tickets (and

so the group procures fewer tickets). In contrast, manipulation (higher x) allows

individuals to steal lottery tickets from others within their group. Suppression would

involve blocking lottery tickets from being stolen. We consider biological examples of

manipulation and suppression in the discussion section.

Results

Model analysis

We seek the evolutionarily stable strategy (ESS), which is the individual strategy that

cannot be invaded by any other rare strategy. Using the neighbour-modulated fitness

approach of Taylor & Frank (1996), we find:

228
$$\begin{cases} \Delta W_z = \left(\bar{R}_{-i} \frac{-(\frac{n-1}{n} - x^a(1-u^b))}{z} + \frac{\frac{n-1}{n} - x^a(1-u^b)}{z} \right)(1-z-x-u) - \bar{R} \\ \Delta W_x = (\bar{R}_{-i}(-ax^{a-1})(1-u^b) + ax^{a-1}(1-u^b))(1-z-x-u) - \bar{R} \\ \quad\quad \Delta W_u = (\bar{R}_{-i}(-x^a)bu^{b-1} + x^a bu^{b-1})(1-z-x-u) - \bar{R} \end{cases}.$$

229   (2)

where traits are analysed one at a time, where ΔW_z means the change in fitness

due to a slight increase in selfishness in the focal individual, and analogous for ΔW_x

and ΔW_u. In addition, $\bar{R}_{-i}$ denotes the relatedness of focal individual to all other

group members, $\bar{R}$ indicate the whole-group relatedness (Pepper 2000)(Appendix

1—1.2). The two relatedness terms can be linked— for example, if the group size is

n, then $\bar{R} = 1/n + ((n-1)/n)\,\bar{R}_{-i}$. For simplicity, we assume all traits only

operate within the group so that $\bar{R}_{-i}$ does not differ between traits. More general

form allowing for cases when different traits are at different social scales is

considered in the derivation in Appendix 1—1. We also assume each trait evolves

independently and there is no allele-level interaction between each trait.

Each of the three equations in (2) represents the change to the inclusive fitness

of an individual that occurs when the individual increases the level at which it

expresses the focal trait (Hamilton 1964a, b). Because all three expressions share

similar structure, we can simplify them into

$$\Delta W_{trait} = (\bar{R}_{-i}(-A) + A)(1-z-x-u) - \bar{R},$$

247   (3)

where A is different for each trait and subscript 'trait' means the type of trait we

are analysing, which could be z, x, or u. The first term on the right, $-A$, represents

the relatedness-weighted reduction in neighbours' share of group productivity due to

an individual's selfishness or manipulation (an inclusive-fitness cost; e.g.,

$\frac{-\left(\frac{n-1}{n}-x^a\left(1-u^b\right)\right)}{z}$ for selfishness). The second term, A, represents the increase in

personal share of group productivity made by a selfish or manipulative individual (an

inclusive-fitness benefit; e.g., $\frac{\frac{n-1}{n}-x^a\left(1-u^b\right)}{z}$ for selfishness). The third term, group

productivity, represents a relatedness-weighted reduction in fitness of the average

group member, self-included, owing to selfishness, manipulation, or suppression (i.e.,

$1-z-x-u$). The cost in group productivity is spread equally among all individuals

in the group, including the actor. The last term, $\bar{R}$, represents the indirect cost of

reducing the share of group productivity obtained by relatives.

Expression 2 could not be solved analytically, and so we found solutions with an

iterative numerical method (Fig. 2; Appendix 1—6.2). We tested the robustness of

our solutions with an individual-based simulation, which includes mutation and drift.

Our simulations and numerical predictions arising showed close agreement (Fig. 2;

Appendix 1—7).

Figure 2. Selfishness, manipulation, and suppression of manipulation. The panels show the ESS
predictions and simulation results for our three-trait model. The solid lines show the ESS
predictions, and the dots show the results of our individual-based simulations ($a = b = 0.5$).
The black segments within each dot show the standard deviation of trait values of the last 10%
duration of 10 repeated simulations (Appendix 1—7).

We found that the ESS selfishness, manipulation, and suppression are all

predicted to show negative relationships with relatedness (Fig. 2). At a lower

relatedness, individuals are predicted to: (i) selfishly keep a higher fraction of their

resources for themselves (as predicted by the classic tragedy of the commons

model); (ii) invest more heavily in manipulating others to help them (as also

predicted in a model without suppression; Appendix 1—2); (iii) invest more heavily in

suppressing manipulation perpetrated by others (when manipulation is at higher

levels). These predictions qualitatively hold for a broad range of parameter space,

with linear or non-linear returns on increased investment in manipulation and

suppression, and show that manipulative cheating as well as its suppression can be

favoured by natural selection.

We predicted that the level of selfishness was greater than the level of

manipulation, which was greater than the level of suppression ($z > x > u$). To a

large extent this reflects that benefit of manipulation depends upon the level of

selfishness, and the benefit of suppression depends upon the level of manipulation.

However, the quantitative predictions depend on details such as the specific values

of the shape parameters (see below), and these relations need not always hold. Our

simulations, which involved mutations, led to relatively static predictions, in

agreement with our analytical results, supporting the results as ESSs (dots versus

lines in Fig. 2).

The quantitative ESS predictions depend upon the shape parameters (Fig. 3).

When the shape parameter of manipulation, a, is more decelerating, selfishness

becomes lower, suppression becomes higher, and manipulation becomes higher until

suppression evolved (Fig. 3a-c). Secondly, when the shape parameter of suppression,

b, is more decelerating, selfishness becomes higher, manipulation becomes lower,

and suppression becomes higher until there is no manipulation to suppress (Fig. 3g-

i). Thirdly, when the two shape parameters covary together, the impact on ESS trait

values is a combination of the two forementioned effects (Fig. 3d-f). Both

manipulation and suppression can be favoured across a wide range of parameter

values.

Figure 3. Nonlinear returns of manipulative cheating and its suppression. Panels (a), (b), and
(c) show the predicted ESS level of selfishness (z^*), manipulation (x^*), and suppression (u^*)
respectively, for different values of relatedness (R, x-axis). From top to bottom, each row of
panels presents varying different parameters on y-axis: the shape parameter for manipulative
cheating (a-c; a), both parameters (d-f; a and b), and the shape parameter for suppression
(g-i; b). The black triangle indicates the horizontal row where the parameter settings are
identical to each other and to Fig. 2. We have focused on decelerating function for shape
parameters because it resembles the nature of communication (Appendix 1—6.1).

Multiple mechanisms for manipulative cheating and suppressing

In our above model, we have focused on evolutionary equilibrium, assuming there is

only a single mechanism to manipulate others. In nature, manipulation could take

many forms, and each requires a different form of suppression. This could lead to

complicated dynamics, such as a genetic arms race, where different forms of

manipulation arise and are then suppressed over time. Because the rise and fall of

trait values are potentially transient, we move away from game-theoretical analysis

and focus on non-equilibrium scenarios through individual-based simulations. We

present two different cases here: one with multiple mechanisms where all are

introduced at the same time, and the other with multiple mechanisms where new

mechanisms are introduced to the population at different time points (via mutation).

Assuming there are k mechanisms to manipulate and suppress, fitness is given

by

$$w_i = \left(z_i + \left(1 - u^b_{1,-i}\right)z_{-i}x^a_{1,i} - \left(1 - u^b_{1,i}\right)z_i x^a_{1,-i} + \left(1 - u^b_{2,-i}\right)z_{-i}x^a_{2,i} - $$

$$\left(1 - u^b_{2i}\right)z_i x^a_{2,-i} + \cdots + \left(1 - u^b_{k,-i}\right)z_{-i}x^a_{k,i} - \left(1 - u^b_{k,i}\right)z_i x^a_{k,-i} \right) \bar{z}^{-1} \left(1 - \bar{z} - \right.$$

$$\left. \bar{x}_1 - \bar{u}_1 - \bar{x}_2 - \bar{u}_2 - \cdots - \bar{x}_k - \bar{u}_k \right).$$

(4)

where $x_1, x_2, \ldots, x_n$ denote the n mechanisms of manipulation and $u_1, u_2, \ldots, u_n$

represent suppression. Additional terms of gain and loss from manipulation and

being manipulated are added to the first parenthesis, and group productivity can

potentially be decreased by all of the $2k + 1$ traits.

We first examine this potentially non-equilibrium scenario by assuming all

mechanisms are present from the beginning and consider the simplest case of two

mechanisms ($k = 2$). By simulating various combinations of shape parameters and

relatedness, we found that levels of selfishness and manipulation do not always tend

to a single equilibrium. Instead, they can oscillate in a periodic style (Fig. 4a). Though

the oscillation in suppression is less profound, it also fluctuates in a smaller

amplitude in response to the changes in levels of manipulation. We used harmonic

regression to test for periodic oscillations, as opposed to random noise (Fig. 4b and

Appendix 1—4.2).

We found oscillations occur when relatedness is relatively low ($R < 1/6$) and

when shape parameters are intermediate decelerating ($0.6 < a, b < 0.85$). In these

intermediate settings, the optimal levels of manipulation and selfishness are similar,

which could lead to a similar pay-off between investing in them. Thus, any small

perturbation in trait values could disturb the dynamics and generate oscillation. In

support of the hypothesis that oscillation is a consequence of interactions between

selfishness and manipulation, we found that oscillation can be prevented when

selfishness cannot mutate (Appendix 1—4.3) or when there is no linkage between

traits (Appendix 1—4.4).

We also found that oscillations can occur when there are more than two

mechanisms to manipulate and supress (Appendix 1—4.5), becomes slower when

mutation rate is smaller (Appendix 1—4.6), and less likely when there is a personal

reproduction cost to manipulation and suppression (Appendix 1—4.7). In addition,

we consider the case where recombination of trait values can occur (Appendix 1—
4.8). The dynamics between cheats and cooperators can involve frequency
dependent selection (Ross-Gillespie *et al.* 2007; Gore *et al.* 2009; Patel *et al.* 2019). In
this model, we were examining evolution in a continuous trait space and so we were
unable to test for frequency dependent selection in the way that can be done with
interactions between cheats and cooperator (i.e., 2 types, as opposed to continuous)
(Allison 2005; Yurtsev *et al.* 2013).

Figure 4. Selfishness and manipulative cheating traits can oscillate when there are two
mechanisms for manipulation and suppression. (a) An example of oscillating dynamics when
shape parameters are intermediate, and relatedness is relatively low ($R = 1/8$; $a = b = 0.7$).
Broader analysis and more details are provided in Appendix 1—4. (b) The estimated amplitude
of the time series of selfishness by harmonic regression. Harmonic regression is an analysis of
time series to show the significance of periodic oscillation, where low period means frequent
recurrent pattern, and high period means recurrence ager a long span. Here, 10 repeated
simulations were used and only the last 75% of the 3000 generations was analysed, after
selfishness is roughly stabilised. The analysis shows maximal amplitude is found at period of
815 generations with p-value of 1.8035×10^{-5}. Mean estimation is shown in black curve whereas
the 95% confidence interval is illustrated as the grey area (F-test; $\mu \pm 2\sigma$).

We now consider a slightly different scenario. In the above analysis, we
assumed that all manipulation and suppression mechanisms were available at the
start of the simulation. An alternative possibility is that different mechanisms can
arise over time, by mutation. When a manipulation trait arises, it could be selected
for. Furthermore, once a form of manipulation becomes sufficiently common, then a
mutation that allows suppression of that manipulation could be selected for. These
different forms of manipulation could involve very different traits, or new forms of an
old trait, that would require a different suppression mechanism. This scenario could
potentially lead to an evolutionary arms race where new forms of manipulation
evolve and are then suppressed. We modelled this scenario by introducing mutations
for new manipulative traits at random time points, and then mutations for the
suppression of manipulation 400 generations afterwards for illustrative purpose.

We found that this scenario also leads to a form of arms race (Fig. 5). When a

new form of manipulation arises, it initially spreads. This spread of manipulation

favours lower levels of selfishness, because the manipulation reduces the benefit of

selfishly investing in reproduction. In addition, once a manipulation is common, any

mutation that allows suppression of that manipulation will be favoured, and hence

increase in frequency. As suppression spreads, this selects for reduced investment in

manipulation (which would just be blocked), and hence also increased selfish

investment in reproduction (which won't be manipulated anymore) (Fig. 5). Overall,

this leads to a pattern of recurring dynamics whenever a new manipulative trait

arises, and where individuals can show variable levels of selfishness, manipulation

and suppression.

For illustrative purposes, our example in Figure 5 shows the gradual evolution of

10 different manipulation mechanisms and focuses on the parameter space where

suppression can almost completely block manipulation in a very effective way ($a =$

$0.5, b = 0.01$)(Scott & West 2019). We examine other scenarios in Appendix 1—5

(Appendix 1—figure 14 & Appendix 1—figure 15). The repeated invasion of different

forms of manipulation, and their suppression, can be seen as a form of trait

substitution sequence (Champagnat 2006; Méléard & Tran 2008).

Altogether, our results from the multiple mechanisms of manipulation suggest

that: (i) multiple types of manipulation can arise and / or be maintained in a

population; (ii) the manipulation traits and their suppression can sometimes show

oscillating dynamics, where the levels of selfishness, manipulation, and/ or

suppression oscillate. The oscillation originates either from interactions between

coevolving traits (endogenous; Fig 4) or from the introduction of new mechanisms

(exogenous; Fig. 5). These results provide possible explanations for cases where

there appears to be genetic variation for the extent or form of cheating and/or

cheating suppression, such as the variance between ant patrilines (Hughes &

Boomsma 2008), slime moulds strains (clones)(Buttery *et al.* 2009; Khare *et al.* 2009;

Levin *et al.* 2015), or different bacteria strains (Kümmerli *et al.* 2009; Bruce *et al.*

2017; Butaitė *et al.* 2017).

Figure 5. Time series of trait dynamics of the simulations when new mechanisms for manipulation and suppression are introduced at different time points. Selfishness is shown in the upper panel, the ten manipulative traits are shown in the middle panel, and the ten suppressive traits are shown in the bottom panel. The y-axis is scaled differently across the panels, to match the levels of variation. Each manipulative and suppressive trait are mutated to non-zero values in arbitrary time points for better visualisation. The shape parameters are set to $a = 0.5$ for manipulation and $b = 0.01$ for suppression, and the relatedness is $R = 1/3$. All data are from a single representative simulation.

Discussion

We found that: (1) manipulative cheating can be favoured across a wide variety of

conditions; (2) manipulation is more likely to be favoured when relatedness is

relatively low; (3) suppression of manipulation can also be favoured. In addition,

when we allowed for multiple mechanisms (types) of manipulation, we found that
non-equilibrium dynamics could occur, such as oscillations. Oscillations could be
generated in either (i) a coevolutionary interaction between selfishness and
manipulation; or (ii) a genetic arms race between different forms of manipulative
cheating and their specific suppressors. These non-equilibrium dynamics could help
explain the genetic diversity that has been observed for both cheating and the
response to cheating across the biological world (Hughes *et al.* 2003; Linksvayer
2006; Anderson *et al.* 2008; Hughes & Boomsma 2008; Buttery *et al.* 2009; Khare *et*
al. 2009; Kümmerli *et al.* 2009; Mitchell *et al.* 2012; Stürup *et al.* 2014; Levin *et al.*
2015; Ostrowski *et al.* 2015; Pollak *et al.* 2016; Bruce *et al.* 2017; Butaitė *et al.* 2017;
Noh *et al.* 2018; Ostrowski 2019; Furubayashi *et al.* 2020; Bruce *et al.* 2021; Mizuuchi
et al. 2022).
457
Our general aim was to investigate scenarios where cheating is characterised by
more than just 'cooperate less'. We did this by developing a theoretical model where
individuals could manipulate others, to make them cooperate more. There is a range
of other possibilities for more complex forms of cheating, such as obtaining a
disproportionate share of some public good, where we suggest that a similar logic
would apply (i.e., the evolution of 'more efficient' cheating). A key distinction here is
that we are not modelling some evolutionarily stable signalling system which
benefits the signaller and receiver (Maynard Smith & Harper 2003). We have instead
modelled coercion, which provides no benefit to the recipient of that coercion.
Consequently, the response would be to evolve to suppress or ignore that coercion,
rather than evolve to some evolutionarily stable mutually beneficial response. In
Appendix 1—4.8 we show that recombination does not change the predictions of our
model so we expect our general conclusions to apply more widely, such as in a
diploid population. This finding agrees with previous literature that a haploid asexual
population is equivalent to a diploid and sexual population (Taylor 1996; Day & Taylor
1998). It would, however, be useful to examine the consequences of different genetic
systems, especially those with asymmetric inheritance such as haplodiploidy.
475
Manipulative cheating could take many different forms in different species. In
bacteria and other microorganisms, individuals could produce coercive molecules
that make other cells invest more into the production of public goods, and invest less
into reproduction. In species that form fruiting bodies, manipulation could cause the
cells of other lineages to be more likely to become stalk cells (Buttery *et al.* 2009;
Khare *et al.* 2009; Levin *et al.* 2015). In organisms such as social insects, the method
of manipulation would depend upon how caste is determined (Schwander *et al.*

2010). In species where workers control the caste of offspring, manipulation could
cause workers from other lineages (different queens or patrilines) to be less likely to
rear their own offspring as reproductive, and be more likely to rear the offspring of
the manipulating lineage as reproductive (Hamilton 1972). In species where larvae
control their own caste, manipulation could decrease the likelihood that other
individuals develop as reproductive (Wenseleers & Ratnieks 2004). Alternatively,
manipulation could occur through more direct approaches, such as selectively killing
the eggs from other lineages. These mechanisms all share the same concept that the
'more manipulative' lineage gains a greater share of group productivity.

Similarly, suppression could involve a variety of different mechanisms, from
evolving to ignore coercion, to blocking behaviours that steal resources or kill
individuals, to making fewer public goods. Experiments on slime moulds have found
that strains can evolve to become more resistant to cheats (Levin *et al.* 2015).
Bacteria appear to be able to produce more specific and less exploitable public goods
(Kümmerli *et al.* 2009; Bruce *et al.* 2017; Butaitė *et al.* 2017). Different lineages of
cooperative RNA replicators can survive better against different cheating replicators
(Mizuuchi *et al.* 2022). These cases share the same concept that cooperators can
somewhat defend against or suppress the cheats. Other possible mechanisms are
discussed elsewhere, for bacteria and viruses.

Previous theoretical work has focused on conflict suppression and hence how
the tragedy of the commons could be avoided (Frank 1995; West *et al.* 2002; Frank
2003; Wenseleers *et al.* 2003; Wenseleers *et al.* 2004; Ratnieks *et al.* 2006). In these
previous papers, selfishness could be selected against, by mechanisms such as
policing, sanctions, and social reputation, which alter the costs and benefits of
behaving selfishly (Frank 1995; Milinski *et al.* 2002; Dionisio & Gordo 2006, 2007; El
Mouden *et al.* 2010). In contrast, we have examined a different route to increased
selfish behaviour – the manipulation of others via manipulative cheating.
Furthermore, we have shown that if there are multiple manipulation traits, then
genotypic diversity can be maintained for selfishness, manipulative cheating, and/or
its suppression. This diversity result links to previous work on quorum sensing in
bacteria, which has shown how cycles of 'cheating' and 'cheating immunity' can arise
(Eldar 2011; Pollak *et al.* 2016). Several theoretical models have examined the
coevolution of offense and defence traits in ecological interactions (Press & Dyson
2012; Hilbe *et al.* 2013) and in sexual conflict (Gavrilets 2000; Gavrilets *et al.* 2001;
Rowe *et al.* 2005). These models examined specific ecological scenarios, such as
memory of previous interactions or runaway processes. Other recent models have

examined a different manipulation question- how individuals can be favoured to manipulate relatives when this leads to an indirect fitness benefit to the manipulator, and the manipulated individual is not selected to resist (González-Forero & Gavrilets 2013; González-Forero 2014; González-Forero 2015; González-Forero & Peña 2021). Manipulation can also be favoured in mutualisms if it leads to better terms of trading (Wyatt *et al.* 2016). More generally, manipulation and counteradaptations to prevent elimination can arise in many forms of communication (Dawkins *et al.* 1979; Maynard Smith & Harper 2003).

To conclude, we have shown how cheating can take more complex forms than just lower levels of cooperation. We have shown that individuals can also be selected to manipulate the behaviour of others, for their own selfish gain. Furthermore, this can lead to evolutionary arms races between attempts to manipulate and to suppress that manipulation. A major task for the future is to determine the extent to which more complex forms of cheating occur in the natural world (Pollak *et al.* 2016; Meir *et al.* 2020; Leeks *et al.* 2021). In many cases, this would require elucidation of the underlying mechanisms, which could be completely different across different species. The potential for manipulation and its suppression could arise at all levels of life, from RNA replicators to complex animal societies.

Code and data availability

All results are generated using C and Python. The codes and data used for this study are available at: https://github.com/mingpapilio/Codes_Manipulative_Cheat

References

1. Agrawal, A. & Lively, C.M. (2002). Infection genetics: gene-for-gene versus matching-alleles models and all points in between. *Evolutionary Ecology Research*, 4, 91-107.

2. Allison, S.D. (2005). Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecology Letters*, 8, 626-635.

3. Anderson, K.E., Linksvayer, T.A. & Smith, C.R. (2008). The causes and consequences of genetic caste determination in ants (Hymenoptera: Formicidae). *Myrmecological News*, 11, 119-132.

4. Bonner, J.T. (1982). Evolutionary strategies and developmental constraints in the cellular slime molds. *The American Naturalist*, 119, 530-552.

5. Bourke, A.F.G. (2011). *Principles of social evolution*. Oxford University Press.

6. Bruce, J.B., Cooper, G.A., Chabas, H., West, S.A. & Griffin, A.S. (2017). Cheating and

resistance to cheating in natural populations of the bacterium *Pseudomonas*
fluorescens. *Evolution*, 71, 2484-2495.

7. Bruce, J.B., Lion, S., Buckling, A., Westra, E.R. & Gandon, S. (2021). Regulation of
prophage induction and lysogenization by phage communication systems.
Current Biology, 31, 5046-5051.e5047.

8. Butaitė, E., Baumgartner, M., Wyder, S. & Kümmerli, R. (2017). Siderophore
cheating and cheating resistance shape competition for iron in soil and
freshwater *Pseudomonas* communities. *Nature Communications*, 8, 414.

9. Buttery, N.J., Rozen, D.E., Wolf, J.B. & Thompson, C.R.L. (2009). Quantification of
social behavior in *D. discoideum* reveals complex fixed and facultative
strategies. *Current Biology*, 19, 1373-1377.

10. Champagnat, N. (2006). A microscopic interpretation for adaptive dynamics trait
substitution sequence models. *Stochastic Processes and their Applications*,
116, 1127-1160.

11. Dawkins, R., Krebs, J.R., Maynard Smith, J. & Holliday, R. (1979). Arms races
between and within species. *Proceedings of the Royal Society B: Biological*
Sciences, 205, 489-511.

12. Day, T. & Taylor, P.D. (1998). Unifying Genetic and Game Theoretic Models of Kin
Selection for Continuous Traits. *Journal of Theoretical Biology*, 194, 391-407.

13. Dionisio, F. & Gordo, I. (2006). The tragedy of the commons, the public goods
dilemma, and the meaning of rivalry and excludability in evolutionary biology.
Evolutionary Ecology Research, 8, 321-332.

14. Dionisio, F. & Gordo, I. (2007). Controlling excludability in the evolution of
cooperation. *Evolutionary Ecology Research*, 9, 365-373.

15. El Mouden, C., West, S.A. & Gardner, A. (2010). The enforcement of cooperation
by policing. *Evolution*, 64, 2139-2152.

16. Eldar, A. (2011). Social conflict drives the evolutionary divergence of quorum
sensing. *Proceedings of the National Academy of Sciences*, 108, 13635-13640.

17. Frank, S.A. (1994). Kin selection and virulence in the evolution of protocells and
parasites. *Proceedings of the Royal Society B: Biological Sciences*, 258, 153-
161.

18. Frank, S.A. (1995). Mutual policing and repression of competition in the evolution
of cooperative groups. *Nature*, 377, 520-522.

19. Frank, S.A. (1996). Models of parasite virulence. *The Quarterly Review of Biology*,
71, 37-78.

20. Frank, S.A. (1998). *Foundations of Social Evolution*. Princeton University Press.

21. Frank, S.A. (2003). Repression of competition and the evolution of cooperation.
Evolution, 57, 693-705.

22. Frank, S.A. (2010a). Demography and the tragedy of the commons. *Journal of*
Evolutionary Biology, 23, 32-39.

23. Frank, S.A. (2010b). A general model of the public goods dilemma. *Journal of*
Evolutionary Biology, 23, 1245-1250.

24. Frank, S.A. (2010c). Microbial secretor- cheater dynamics. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, 365, 2515-2522.

25. Furubayashi, T., Ueda, K., Bansho, Y., Motooka, D., Nakamura, S., Mizuuchi, R. *et*
al. (2020). Emergence and diversification of a host-parasite RNA ecosystem
through Darwinian evolution. *eLife*, 9, e56038.

26. Gardner, A. & West, Stuart A. (2004). Cooperation and punishment, especially in
humans. *The American Naturalist*, 164, 753-764.

27. Gardner, A., West, Stuart A. & Barton, Nicholas H. (2007). The relation between
multilocus population genetics and social evolution theory. *The American*
Naturalist, 169, 207-226.

28. Gavrilets, S. (2000). Rapid evolution of reproductive barriers driven by sexual
conflict. *Nature*, 403, 886-889.

29. Gavrilets, S., Arnqvist, G. & Friberg, U. (2001). The evolution of female mate
choice by sexual conflict. *Proceedings of the Royal Society B: Biological*
Sciences, 268, 531-539.

30. Ghoul, M., Griffin, A.S. & West, S.A. (2014). Toward an evolutionary definition of
cheating. *Evolution*, 68, 318-331.

31. Gill, J. & Johnson, P. (2002). *Research methods for managers*. Sage.

32. González-Forero, M. (2014). An evolutionary resolution of manipulation conflict.
Evolution, 68, 2038-2051.

33. González-Forero, M. (2015). Stable eusociality via maternal manipulation when
resistance is costless. *Journal of Evolutionary Biology*, 28, 2208-2223.

34. González-Forero, M. & Gavrilets, S. (2013). Evolution of manipulated behavior.
The American Naturalist, 182, 439-451.

35. González-Forero, M. & Peña, J. (2021). Eusociality through conflict dissolution.
Proceedings of the Royal Society of London. Series B: Biological Sciences, 288,
20210386.

36. Gore, J., Youk, H. & van Oudenaarden, A. (2009). Snowdrift game dynamics and
facultative cheating in yeast. *Nature*, 459, 253-256.

37. Griffin, A.S., West, S.A. & Buckling, A. (2004). Cooperation and competition in
pathogenic bacteria. *Nature*, 430, 1024-1027.

38. Hamilton, W.D. (1964a). The genetical evolution of social behaviour. I. *Journal of*
Theoretical Biology, 7, 1-16.

39. Hamilton, W.D. (1964b). The genetical evolution of social behaviour. II. *Journal of*

Theoretical Biology, 7, 17-52.

40. Hamilton, W.D. (1972). Altruism and related phenomena, mainly in social insects.
Annual Review of Ecology and Systematics, 3, 193-232.

41. Hilbe, C., Nowak, M.A. & Traulsen, A. (2013). Adaptive dynamics of extortion and
compliance. *PLOS ONE*, 8, e77886.

42. Hudson, R.E., Aukema, J.E., Rispe, C. & Roze, D. (2002). Altruism, cheating, and
anticheater adaptations in cellular slime molds. . *The American Naturalist*,
160, 31-43.

43. Hughes, W.O.H. & Boomsma, J.J. (2008). Genetic royal cheats in leaf-cutting ant
societies. *Proceedings of the National Academy of Sciences*, 105, 5150-5153.

44. Hughes, W.O.H., Sumner, S., Van Borm, S. & Boomsma, J.J. (2003). Worker caste
polymorphism has a genetic basis in Acromyrmex leaf-
cutting ants. *Proceedings of the National Academy of Sciences*, 100, 9394.

45. Jiricny, N., Diggle, S.P., West, S.A., Evans, B.A., Ballantyne, G., Ross-Gillespie, A. *et
al.* (2010). Fitness correlates with the extent of cheating in a bacterium.
Journal of Evolutionary Biology, 23, 738-747.

46. Khare, A., Santorelli, L.A., Strassmann, J.E., Queller, D.C., Kuspa, A. & Shaulsky, G.
(2009). Cheater-resistance is not futile. *Nature*, 461, 980-982.

47. Kümmerli, R., Griffin, A.S., West, S.A., Buckling, A. & Harrison, F. (2009). Viscous
medium promotes cooperation in the pathogenic bacterium *Pseudomonas
aeruginosa*. *Proceedings of the Royal Society B: Biological Sciences*, 276,
3531-3538.

48. Leeks, A., West, S.A. & Ghoul, M. (2021). The evolution of cheating in viruses.
Nature Communications, 12, 6928.

49. Lehmann, L., Rousset, F., Roze, D. & Keller, L. (2007). Strong reciprocity or strong
ferocity? A population genetic view of the evolution of altruistic punishment.
The American Naturalist, 170, 21-36.

50. Levin, S.R., Brock, D.A., Queller, D.C. & Strassmann, J.E. (2015). Concurrent
coevolution of intra-organismal cheaters and resisters. *Journal of Evolutionary
Biology*, 28, 756-765.

51. Linksvayer, T.A. (2006). Direct, maternal, and sibsocial genetic effects on
individual and colony traits in an ant. *Evolution*, 60, 2552-2561.

52. Liu, M., West, S.A. & Cooper, G.A. (2021). Relatedness and the evolution of
mechanisms to divide labor in microorganisms. *Ecology and Evolution*, 11,
14475-14489.

53. MacLean, R.C. & Gudelj, I. (2006). Resource competition and social conflict in
experimental populations of yeast. *Nature*, 441, 498-501.

54. Maynard Smith, J. (1964). Group selection and kin selection. *Nature*, 201, 1145-

1147.

55. Maynard Smith, J. & Harper, D. (2003). *Animal signals*. Oxford University Press.

56. Meir, M., Harel, N., Miller, D., Gelbart, M., Eldar, A., Gophna, U. *et al.* (2020). Competition between social cheater viruses is driven by mechanistically different cheating strategies. *Science Advances*, 6, eabb7990.

57. Méléard, S. & Tran, V.C. (2008). Trait substitution sequence process and canonical equation for age-structured populations. *Journal of Mathematical Biology*, 58, 881.

58. Milinski, M., Semmann, D. & Krambeck, H.-J. (2002). Reputation helps solve the 'tragedy of the commons'. *Nature*, 415, 424-426.

59. Mitchell, R.E., Frost, C.L. & Hughes, W.O.H. (2012). Size and asymmetry: are there costs to winning the royalty race? *Journal of Evolutionary Biology*, 25, 522-531.

60. Mizuuchi, R., Furubayashi, T. & Ichihashi, N. (2022). Evolutionary transition from a single RNA replicator to a multiple replicator network. *Nature Communications*, 13, 1460.

61. Müller, J., Kuttler, C., Hense, B.A., Rothballer, M. & Hartmann, A. (2006). Cell–cell communication by quorum sensing and dimension-reduction. *Journal of Mathematical Biology*, 53, 672-702.

62. Noh, S., Geist, K.S., Tian, X., Strassmann, J.E. & Queller, D.C. (2018). Genetic signatures of microbial altruism and cheating in social amoebas in the wild. *Proceedings of the National Academy of Sciences*, 115, 3096-3101.

63. Ostrowski, E.A. (2019). Enforcing cooperation in the social amoebae. *Current Biology*, 29, R474-R484.

64. Ostrowski, Elizabeth A., Shen, Y., Tian, X., Sucgang, R., Jiang, H., Qu, J. *et al.* (2015). Genomic signatures of cooperation and conflict in the social amoeba. *Current Biology*, 25, 1661-1665.

65. Patel, M., Raymond, B., Bonsall, M.B. & West, S.A. (2019). Crystal toxins and the volunteer's dilemma in bacteria. *Journal of Evolutionary Biology*, 32, 310-319.

66. Pepper, J.W. (2000). Relatedness in trait group models of social evolution. *Journal of Theoretical Biology*, 206, 355-368.

67. Pollak, S., Omer-Bendori, S., Even-Tov, E., Lipsman, V., Bareia, T., Ben-Zion, I. *et al.* (2016). Facultative cheating supports the coexistence of diverse quorum-sensing alleles. *Proceedings of the National Academy of Sciences*, 113, 2152-2157.

68. Press, W.H. & Dyson, F.J. (2012). Iterated prisoner's dilemma contains strategies that dominate any evolutionary opponent. *Proceedings of the National Academy of Sciences*, 109, 10409-10413.

69. Queller, D.C. (1984). Kin selection and frequency dependence: a game theoretic
approach. *Biological Journal of the Linnean Society*, 23, 133-143.

70. Queller, D.C. & Strassmann, J.E. (2018). Evolutionary conflict. *Annual Review of
Ecology, Evolution, and Systematics*, 49, 73-93.

71. Ratnieks, F.L.W., Foster, K.R. & Wenseleers, T. (2006). Conflict resolution in insect
societies. *Annual Review of Entomology*, 51, 581-608.

72. Rogers, D.W., McConnell, E., Miller, E.L. & Greig, D. (2017). Diminishing returns on
intragenic repeat number expansion in the production of signaling peptides.
Molecular Biology and Evolution, 34, 3176-3185.

73. Ross-Gillespie, A., Gardner, A., West, Stuart A. & Griffin, Ashleigh S. (2007).
Frequency dependence and cooperation: theory and a test with bacteria. *The
American Naturalist*, 170, 331-342.

74. Rowe, L., Cameron, E. & Day, T. (2005). Escalation, retreat, and female
indifference as alternative outcomes of sexually antagonistic coevolution. *The
American Naturalist*, 165, S5-S18.

75. Sasaki, A. (2000). Host-parasite coevolution in a multilocus gene-for-gene system.
Proceedings of the Royal Society B: Biological Sciences, 267, 2183-2188.

76. Schwander, T., Lo, N., Beekman, M., Oldroyd, B.P. & Keller, L. (2010). Nature
versus nurture in social insect caste differentiation. *Trends in Ecology &
Evolution*, 25, 275-282.

77. Scott, T.W. & West, S.A. (2019). Adaptation is maintained by the parliament of
genes. *Nature Communications*, 10, 5163.

78. Strassmann, J.E., Zhu, Y. & Queller, D.C. (2000). Altruism and social cheating in the
social amoeba Dictyostelium discoideum. *Nature*, 408, 965-967.

79. Stürup, M., Nash, D.R., Hughes, W.O.H. & Boomsma, J.J. (2014). Sperm mixing in
the polyandrous leaf-cutting ant Acromyrmex echinatior. *Ecology and
Evolution*, 4, 3571-3582.

80. Taylor, P.D. (1996). Inclusive fitness arguments in genetic models of behaviour.
Journal of Mathematical Biology, 34, 654-674.

81. Taylor, P.D. & Frank, S.A. (1996). How to make a kin selection model. *Journal of
Theoretical Biology*, 180, 27-37.

82. Tibbetts, E.A., Liu, M., Laub, E.C. & Shen, S.-F. (2020). Complex signals alter
recognition accuracy and conspecific acceptance thresholds. *Philosophical
Transactions of the Royal Society B: Biological Sciences*, 375, 20190482.

83. Wenseleers, T., Helanterä, H., Hart, A. & Ratnieks, F.L.W. (2004). Worker
reproduction and policing in insect societies: an ESS analysis. *Journal of
Evolutionary Biology*, 17, 1035-1047.

84. Wenseleers, T. & Ratnieks, F.L.W. (2004). Tragedy of the commons in *Melipona*

bees. *Proceedings of the Royal Society B: Biological Sciences*, 271, S310-S312.

85. Wenseleers, T., Ratnieks, F.L.W. & Billen, J. (2003). Caste fate conflict in swarm-
founding social *Hymenoptera*: an inclusive fitness analysis. *Journal of*
Evolutionary Biology, 16, 647-658.

86. West, S.A. & Buckling, A. (2003). Cooperation, virulence and siderophore
production in bacterial parasites. *Proceedings of the Royal Society B:*
Biological Sciences, 270, 37-44.

87. West, S.A., Kiers, E.T., Simms, E.L. & Denison, R.F. (2002). Sanctions and
mutualism stability: why do rhizobia fix nitrogen? *Proceedings of the Royal*
Society B: Biological Sciences, 269, 685-694.

88. Wild, G. (2011). Inclusive fitness from multitype branching processes. *Bulletin of*
Mathematical Biology, 73, 1028-1051.

89. Wiley, R.H. (2013). Signal detection, noise, and the evolution of communication.
In: *Animal Communication and Noise* (ed. Brumm, H). Springer Berlin
Heidelberg Berlin, Heidelberg, pp. 7-30.

90. Wyatt, G.A.K., Kiers, E.T., Gardner, A. & West, S.A. (2016). Restricting mutualistic
partners to enforce trade reliance. *Nature Communications*, 7, 10322.

91. Yurtsev, E.A., Chao, H.X., Datta, M.S., Artemova, T. & Gore, J. (2013). Bacterial
cheating drives the population dynamics of cooperative antibiotic resistance
plasmids. *Molecular Systems Biology*, 9, 683.

Appendix 1

Our appendix contains seven sections. In section 1, we formally analyse our model

with single mechanisms to manipulate and suppress manipulation. Section 2 contains

the non-suppression model, which is a reduced model without suppressive traits.

This model bridges the gap between the single-mechanism model and the classic

tragedy of the commons (Frank 1994, 1996; Taylor & Frank 1996; Frank 1998;

Dionisio & Gordo 2006; Frank 2010a, b). Further, the non-suppression model also

illustrates a biologically possible case where manipulation cannot be suppressed. We

consider the effects of personal cost in manipulation and suppression in section 3.

Section 4 provides additional analyses for the scenario with two mechanisms to

manipulate and suppress, where we found oscillations in selfishness and

manipulation (Fig. 4). We examine the origin of the oscillation dynamics and compare

the results with game-theoretical analysis. We subsequently present similar arms

race dynamics in section 5 for the scenario with multiple mechanisms introduced at

different time points (Fig. 5), with different biological settings assumed in each case.

In section 6, we discuss some modelling details and justification. Lastly, section 7 has

the details for the individual-based simulations.

1. Formal analysis of the scenario with a single mechanism to manipulate and

suppress manipulation

1.1 Derivation of the fitness equation

We track a population of haploid asexual individuals by observing it at discrete points

spaced evenly in time. The model population itself is subdivided into groups of size

n. Groups are re-constituted at the very end of a given time step, through among

group competition. Individuals compete within their own group for their share of the

group's productivity.

We use y_i to denote the (absolute) competitive effort realised by focal

individual i. We build up the fitness expression by assuming that the focal individual

initially commits to a within-group level of competition equal to z_i. Likewise, each

non-focal individual $k \neq i$ commits to a within-group level of competitive effort

equal to z_k.

Once each group member has committed to an initial level of competitive

effort, the group enters a phase in which members can manipulate effort (i.e.,

investment) from one another. We model the manipulation phase by assuming that

the focal group member is assigned a non-self partner, uniformly at random with

replacement. The assumption is equivalent to letting the focal individual manipulate

every neighbour to the same degree and we use a probabilistic sampling narrative

for the ease of explaining each process (Wild 2011). When manipulation takes place,

the focal individual attempts to manipulate a fraction x_i^a of the competitive effort

z_k committed to by the partner. Here, we are treating x_i as a new trait (i.e., effort

devoted to manipulation), and a as a constant that modulates the efficacy of the

trait (i.e., determines how effective the effort is). The partner individual can also

suppress manipulation. Specifically, we assume that the partner individual prevents a

fraction $(1 - u_k^b)$ of manipulation that would have otherwise occurred. Here, we are

thinking of u_k as a trait, and b as a constant that modulates the efficacy of the

trait.

The focal individual is not only a manipulator, it can also be a the target of

manipulation. In keeping with the description above, the focal individual is chosen to

be manipulated by each of its non-self neighbours with probability $1/(n-1)$. When

chosen, the focal individual loses $x_k^a(1 - u_i^b)$ of the competitive effort z_i

committed to by itself.

Putting the details together, we arrive at the following expression for the

competitive effort realised by focal individual i:

$$y_i = z_i + x_i^a \frac{1}{n-1}\sum_{k \neq i}(1 - u_k^b)z_k - (1 - u_i^b)z_i \frac{1}{n-1}\sum_{k \neq i}x_k$$

$$= z_i + x_i^a(1 - \bar{u}_{-i}^b)\bar{z}_{-i} - \bar{x}_{-i}(1 - u_i^b)z_i$$

831   (A1)

where a bar paired with a $-i$ subscript denotes the arithmetic mean of non-focal

individuals in the focal group. It is important to note that the share of group

productivity of the average non-focal individual can be expressed as

$$\bar{y}_{-i} = \frac{1}{n-1}\sum_{k \neq i}\left(z_k + x_k^a \frac{1}{n-1}(1 - u_i^b)z_i - \frac{1}{n-1}x_i^a(1 - u_k^b)z_k + \right.$$

$$\left. x_k^a \frac{1}{n-1}\sum_{j \neq i,k}(1 - u_j^b)z_j - \frac{1}{n-1}(1 - u_k^b)z_k \sum_{j \neq i,k}x_j^a\right)$$

$$= \bar{z}_{-i} + \frac{1}{n-1}\bar{x}_{-i}(1 - u_i^b)z_i - \frac{1}{n-1}x_i^a(1 - \bar{u}_{-i}^b)\bar{z}_{-i} + \frac{1}{(n-1)^2}(\sum_{k \neq i}\sum_{j \neq i,k}x_k^a(1 -$$

$$u_j^b)z_j - \sum_{k \neq i}\sum_{j \neq i,k}x_j^a(1 - u_k^b)z_k)$$

840   (A2)

841   Further simplification can be achieved by recognising that the term contained in

842   parenthesis immediately above is zero. To see this let $\delta_{kj} = \delta_{jk}$ take a value of 1 if

843     $k = j$ and a value of 0 otherwise. Then,

845     $\left(\sum_{k\neq i,} \sum_{j\neq i,k} x_k^a \left(1 - u_j^b\right)z_j - \sum_{k\neq i,} \sum_{j\neq i,k} \left(1 - u_k^b\right)z_k x_j^a \right) = \left(\sum_{k,j\neq i} x_k^a \left(1 - \right.\right.$

846     $\left.\left. u_j^b\right)z_j\left(1 - \delta_{kj}\right) - \sum_{k,j\neq i} x_j^a \left(1 - u_k^b\right)z_k\left(1 - \delta_{jk}\right) \right) = 0.$

847   (A3)

In words, the term in question simplifies to zero because the double sums both

represent the total sum of all entries of the same $(n-1) \times (n-1)$ matrix.

Therefore, the average competitive effort in the focal group is:

$\bar{y} = \frac{1}{n}y_i + \frac{n-1}{n}\bar{y}_{-i}$

$= \frac{1}{n}z_i + \frac{n-1}{n}\bar{z}_{-i} + \frac{1}{n}x_i^a(1 - \bar{u}_{-i}^b)\bar{z}_{-i} - \frac{n-1}{n}\frac{1}{n-1}x_i^a(1 - \bar{u}_{-i}^b)\bar{z}_{-i} - $

$\frac{1}{n}\bar{x}_{-i}\left(1 - u_i^b\right)z_i + \frac{n-1}{n}\frac{1}{n-1}\bar{x}_{-i}\left(1 - u_i^b\right)z_i$

$= \bar{z}$

856   (A4)

Where bars with no subscript denote an arithmetic mean taken over the entire group

of n individuals. The average competitive effort in the group is not changed by the

manipulation phase. Manipulation only serves to redistribute the initial commitment

to competitive effort within a given group. It follows that the relative share of group

productivity of the focal individual can be expressed as $y_i/\bar{z}$.

We next assume, for simplicity, that all traits impose a cost on group performance at

the same rate, and we propose that group productivity is proportional to $1 - \bar{z} - $

$\bar{x} - \bar{u}$. Thus, we express fitness of the focal individual as

$w_i = \frac{y_i}{\bar{z}}(1 - \bar{z} - \bar{x} - \bar{u}).$

868   (A5)

Technically, w_i in the previous line is more accurately described as an inclusive-

fitness-generating function, because differentiating it following the steps outline in

(Taylor & Frank 1996) will lead to the correct generalisation of Hamilton's rule

(Hamilton 1964a, b).

1.2 Kin selection analysis

Following Taylor& Frank (1996), we assume that traits z, x, and u are controlled by

a separate locus. We use, g_z, g_x, and g_u, respectively, to denote the genotypic

value of the focal individual at the z, x, and u loci.

To capture the effect of competition on an individual's inclusive fitness we treat
the traits z_i, $\bar{z}_{-i}$, and $\bar{z}$ as a function of g_z. We then differentiate w_i with respect
to g_z and evaluate the result at a point where all individuals in the population have
the same genotypic value at every locus. Evaluating at such a point allows us to
assert that $z_i = \bar{z}_{-i} = \bar{z} = z$, $x_i = \bar{x}_{-i} = \bar{x} = x$, and $u_i = \bar{u}_{-i} = \bar{u} = u$. It also
justifies us replacing dz_i/dg_z with relatedness between the focal individual and
itself ($= 1$), replacing $d\bar{z}_{-i}/dg_z$ with relatedness between the focal individual and
its average non-self group mate ($= \bar{R}_{-i,z}$, where the subscript reminds us that
relatedness is assessed at the z locus), and replacing $d\bar{z}/dg_z$ with relatedness
between the focal individual and the average member of its group, self-included ($=$
$\bar{R}_z$). Following through this entire multi-step procedure, we arrive at

$$\Delta W_z = \left(\bar{R}_{-i,z} \frac{-(\frac{n-1}{n} - x^a(1-u^b))}{z} + \frac{\frac{n-1}{n} - x^a(1-u^b)}{z} \right)(1 - z - x - u) - \bar{R}_z$$

892   (A6)
893 as the inclusive fitness effect of a small increase in selfishness, z. When ΔW_z is
894 zero, z is at equilibrium with respect to the action of selection (ESS).

896   Treating traits x_i, $\bar{x}_{-i}$, and $\bar{x}$ as a function of g_x we can go through
897 analogous steps to obtain

899    $$\Delta W_x = (\bar{R}_{-i,x}(-ax^{a-1})(1 - u^b) + ax^{a-1}(1 - u^b))(1 - z - x - u) - \bar{R}_x$$
900   (A7)
901 where the subscript reminds us that relatedness is now assessed at the x locus.
902 Treating traits u_i, $\bar{u}_{-i}$, and $\bar{u}$ as a function of g_u, we get

904    $$\Delta W_u = (\bar{R}_{-i,u}(-x^a)bu^{b-1} + x^a bu^{b-1})(1 - z - x - u) - \bar{R}_u$$
905   (A8)
as the inclusive fitness effect of suppression, where relatedness is assessed at the u
locus.

1.3 Mathematical criterion of manipulation in other theoretical papers
We compare our model with other theoretical papers, which examine related
concepts. We start with Frank's (1995) policing model:

$$w_i = (\bar{a} - ca_i + (1 - \bar{a}) z_i/\bar{z})(1 - (1 - \bar{a})\bar{z}),$$

914   (A9)

where a_i and z_i are the degree of policing and selfishness of the focal individual,

$\bar{a}$ and $\bar{z}$ are the averages in the focal group. c is the cost of policing. Policing

changes the benefit and cost of selfishness for all group members, with each

multiplied by $1 - \bar{a}$. In contrast, our manipulative cheating involves the actor

coercing the recipient into being less selfish (more cooperative).

González-Forero and Gavrilets (2013) examined a different form of

'manipulation' . In their model, fitness is depending on the behaviour of three types:

manipulator that performs manipulation, recipient of manipulation, and a benefactor

that is not involved in manipulation but benefit from its social partners' behaviour:

$$w = 1 - pc_{manipulate}k_{manipulator} - \left(c_{acquiesces}P(1 - q) - \right.$$
$$\left. c_{resistance}Pq\right)k_{recipient} + b\Pi(1 - Q)k_{benefactor},$$

928   (A10)

where k are the reproductive values, c are the cost for taking each option, p is

the probability for a manipulator to act on a reachable subject, q is the probability

for a subject to resist, P is the average manipulation probability, Q is the average

resistance probability, and Π is the average manipulation probability that

manipulation occurs and successfully reaches a recipient. In addition to the more

complicated pay-off setting, their model assumes that social interactions are tiggered

by manipulation. They are examining when an manipulator can be favoured to

manipulate a related recipient, to behave in a way that provides an indirect benefit

to the recipient, and that this benefit is sufficient that the recipient is not favoured to

resist the manipulator. This contrasts with our model, where manipulative cheating

involves manipulation of selfishness (cooperation) – manipulation of an already

occurring social behaviour to provide a selfish benefit. Therefore, although both

models use the term 'manipulation', they are modelling very different biological

scenarios.

2. The non-suppression model: model with only selfishness and one mechanism for

manipulation

We assume suppression is absent in this model, but manipulation can occur.

Consequently, this gives:

$$w_i = \frac{z_i + z_{-i}x_i^a - z_i x_{-i}^a}{\bar{z}}(1 - \bar{z} - \bar{x}),$$

950   (A11)

where x_i is the level of manipulative cheating of the focal individual, z_{-i} is the

average level of selfishness (excluding the focal individual), a is the shape

parameter for manipulative cheating, and $1 - \bar{z} - \bar{x}$ is the group's productivity. As

above, the parameter *a* determines whether the benefit from increased investment

into manipulation is decelerating ($a < 1$), linear ($a = 1$) or accelerating ($a > 1$).

To facilitate comparison with previous work, the relations between the classic

tragedy of the commons, the non-suppression model, and the model with

suppression are illustrated in Appendix 1—figure 1. The first model is the classic

tragedy of the commons model, which predicts a single optimal level of investment

into cooperation / selfish reproduction (Appendix 1—figure 1a)(Frank 2010a). The

next model is the non-suppression model, which allows for manipulative cheating,

where individuals increase their own investment into reproduction, by reducing that

of others (Appendix 1—figure 1b). Lastly, in the model with suppression, we allow

suppression as a defensive mechanism against manipulative cheating (Appendix 1—

figure 1c).

(a) Classic tragedy of the commons

(b) The non-suppression model:
tragedy+ manipulation

(c) The model with suppression (main model):
tragedy+ manipulation+ suppression

	Classic tragedy of the commons	Non-suppression model	The model with suppression
Group productivity	$1 - \bar{z}$	$1 - \bar{z} - \bar{x}$	$1 - \bar{z} - \bar{x} - \bar{u}$
Individual share of group productivity	$\dfrac{z_i}{\bar{z}}$	$\dfrac{z_i + x_i^a z_{-i} - x_{-i}^a z_i}{\bar{z}}$	$\dfrac{(z_i + (1-u_{-i}^b)x_i^a z_{-i} - (1-u_i^b)x_{-i}^a z_i)}{\bar{z}}$

968
969 Appendix 1—figure 1. Comparison between the classic tragedy of the commons, the non-
970 suppression model, and the model with suppression. Here, we illustrate three models with

gradually increasing complexity. (a) In the classic tragedy of the commons, individuals can choose to invest resources into reproduction (z_i) or group productivity (cooperation; $1 - \bar{z}$). (b) In the non-suppression model, individuals can also manipulate others (x_i), to make them invest more in group productivity. (c) In the model with suppression, individuals can also suppress (block) the manipulation by others (u_i). Within each model, arrows indicate manipulation, the blockings indicate suppression, and the outcome of the two actions are expressed below the arrows. The fitness of an individual is divided into two components: group productivity and share of group productivity, both shown at the bottom of the figure.

We find the change of inclusive fitness effect for selfishness, and manipulation:

$$\begin{cases} \Delta W_z = \left(\bar{R}_{-i,z} \dfrac{-(\frac{n-1}{n} - x^a)}{z} + \dfrac{\frac{n-1}{n} - x^a}{z} \right)(1 - z - x - u) - \bar{R}_z \\ \Delta W_x = (\bar{R}_{-i,x}(-ax^{a-1}) + ax^{a-1})(1 - z - x - u) - \bar{R}_x \end{cases}.$$

(A12)

Expression A12 could not be solved analytically, and so we found solutions with an iterative numerical analysis. We also tested the predictions of our analytical model with an individual-based simulation, which showed close agreement (Appendix 1—figure 2b).

The non-suppression model predicts that manipulation can be favoured, and that greater levels of manipulation will be favoured when relatedness is lower (Appendix 1—figure 2b). This prediction arises because there is an indirect fitness cost to manipulating relatives and reducing group productivity. Further, since a function with smaller shape parameter means the manipulative effective is higher for a given trait value, as shown in the three panels on the left of Appendix 1—figure 2, we find greater intermediate values of manipulation when the shape is more decelerating (smaller a have more shaded area, Appendix 1—figure 3b).

This model also predicts a very different relationship between selfishness and relatedness, compared with the classic tragedy of the commons model and the model with suppression (compare Appendix 1—figure 2b with other panels). Specifically, rather than selfishness increasing as relatedness decreases, the non-suppression model predicts that selfishness shows a non-monotonic (domed) pattern, peaking at intermediate relatedness (Appendix 1—figure 2b). The difference in the non-suppression model is that, relative to tragedy of the commons, selfishness is not as advantageous when relatedness is low. At low relatedness, there is selection for high levels of manipulation – this dilutes the benefit of selfish reproduction because that effort will be taken away by manipulation. Within our lottery ticket metaphor in the main text, the benefit of buying tickets is reduced if those tickets are just going to be stolen.

1010

Comparing the non-suppression model and the model with suppression, a

consequence of adding in suppression is that lower levels of manipulation are

selected for, because the benefit of increased manipulation is reduced if it will be

suppressed (compare Appendix 1—figure 2b & c). This lower level of manipulation,

and the fact that some of it is suppressed, lead to selfishness being beneficial at low

relatedness, as in the classic tragedy of the commons model (compare Appendix 1—

figure 2a & c), and in contrast to when there is no suppression (Appendix 1—figure

2b).

1019

By comparing all three models, we also found the group productivity did not

vary depending upon whether or not manipulation and suppression were included in

our model, compared to the classic tragedy of the commons model (Appendix 1—

figure 2). This is because adding manipulation and suppression does not change the

basic structure of the model: share of productivity multiplied by group productivity.

Thus, the relation between relatedness and group productivity is not affected: the

two additional traits are sharing the 'loss' in group productivity with selfishness,

where the sum of loss is the same for a given relatedness. In other words,

manipulation and suppression do not change the degree of cooperation across the

group and this finding confirms the design in expression A4.

1030

1031

Appendix 1—figure 2. Selfishness, manipulation, and suppression of manipulation. The panels
show the ESS predictions and simulation results for: (a) tragedy of the commons (only
selfishness evolves); (b) the non-suppression model (selfishness and manipulation evolve); (c)
the model with suppression (selfishness, manipulation, and suppression of manipulation
evolve). The solid lines show the ESS predictions, and the dots show the results of our
individual-based simulations ($a = b = 0.5$).

1038

1039
1040 Appendix 1—figure 3. Nonlinear returns in the non-suppression model. Panels (a) and (b) show
1041 the predicted ESS level of selfishness (z^*) and manipulation (x^*) respectively, for different
1042 values of relatedness (R) and the shape parameter (a). The black triangle indicates the
1043 horizontal row where the parameter settings are identical to each other and to Appendix 1—
1044 figure 2b.

1045

3. The models with personal costs: adding cost to relative shares of group
productivity

In the main model, investment in manipulation or suppression led to reduced

investment in group productivity. As an alternative, we also investigated a scenario

where manipulation and selfishness have a personal cost, reducing the share that an

individual obtains of the group productivity, but not influencing the overall success of

the group. In this case, manipulative cheating and suppression of cheating now have

impacts on the average level of competition, and they are not just redistributing the

share of productivity. We first consider the case where suppression is absent and

manipulation has a personal cost, and then we move on to the case where both

manipulation and suppression are allowed.

1057

3.1 The non-suppression model with personal costs

The personal cost of manipulation can be added in many ways, and one easy way is

to multiply the benefit of selfishness by $(1 - x)$, where x is the level of

manipulation. For simplicity, we assume cost is multiplicative and there is no

coefficient before x. This modification changes the competitive effort through

selfishness from z_i to $(1 - x_i)z_i$, and changes the competitive effort gained

through manipulation from $z_{-i}x_i^a$ to $(1 - x_{-i})z_{-i}x_i^a$. Similarly, the average effort

of the group is $(1 - \bar{x})\bar{z}$, making the fitness of the focal individual,

1066

1067
$$w_i = \frac{(1-x_i)z_i + (1-x_{-i})z_{-i}x_i^a - (1-x_i)z_i x_{-i}^a}{(1-\bar{x})\bar{z}}(1 - \bar{z} - \bar{x}),$$

1068
<div align="right">(A13)</div>

where $(1 - \bar{z} - \bar{x})$ is the group productivity and the terms before it is the relative

share of group productivity. By comparing expressions A13 and A11, it is apparent

that we are multiplying $(1 - x)$ for both the numerator and denominator in A11.

However, it is meaningful because each term is multiplied by different x: some are

the level of manipulation of the focal individual, while others are the average of the

social neighbours or the average across the entire group.

1075

We found that adding a personal cost of manipulation leads to a reduced level

of manipulation, especially when $R < 0.1$ (compare Appendix 1—figure 2b& A4b).

Furthermore, because both selfishness and manipulation have a personal cost, the

net benefit obtained from investing in manipulation is reduced, and so we found no

dramatic increase in the predicted level of manipulation when $R < 0.1$ (Appendix

1—figure 4b). In contrast, when manipulation is cost-free at the individual level, the

predicted strategy is to put all selfish effort into manipulation when relatedness of

the population is very low (Appendix 1—figure 2b).

1084

1085   *3.2 The model with suppression and personal cost*

1086   We then allow suppression, with a personal cost, that reduces the share of group

1087   productivity obtained. We multiply each term in the relative share of productivity in

1088   expression 1 with $(1 - x - u)$, where x and u are the ones with same identity as

1089 z they are multiplying (e.g., z_i multiplied by $(1 - x_i - u_i)$. This gives:

1090

$w_i = \dfrac{(1-x_i-u_i)z_i + \left(1-u_{-i}^b\right)(1-x_{-i}-u_{-i})z_{-i}x_i^a - \left(1-u_i^b\right)(1-x_i-u_i)z_i x_{-i}^a}{(1-\bar{x}-\bar{u})\bar{z}}(1 - \bar{z} - \bar{x} - \bar{u}).$

(A14)

We found the predicted level of suppression is lower when a personal cost of

suppression is also included to the model (compare Appendix 1—figure 2c&

Appendix 1—figure 4c). Because there is (i) a lower need for suppression from a

lower predicted level of manipulation, and (ii) a more costly design of suppression,

we only find small levels of suppression when relatedness is low (Appendix 1—figure

4c). In addition, given the presence of suppression, the predicted level of

manipulation is lower than the model without suppression (compare Appendix 1—

figure 4b& c). Despite the lower predicted ESS values, these models with personal

cost for manipulation and suppression still predict both traits can be favoured,

especially when relatedness is low.

1103

Appendix 1—figure 4. Selfishness, manipulation, and suppression of manipulation. The panels show the ESS predictions and simulation results for: (a) tragedy of the commons (only selfishness evolves); (b) the non-suppression model with personal cost (both selfishness and manipulation evolve); (c) the model with suppression and personal cost (selfishness, manipulation, and suppression evolve). All results are generated with the same shape parameters as other figures ($a = b = 0.5$).

4. Additional analysis of the scenario with two mechanisms where all traits are simultaneously introduced

4.1 The relation between ESS and relatedness

In order to understand the oscillating dynamics (Fig. 4) better, we perform game-theoretical analysis for the two-mechanisms case and examine when do the two approaches disagree with each other. We begin by analysing the model along relatedness and found it gives very similar predictions to the scenario with only one mechanism to manipulate (compare Fig. 2 & Appendix 1—figure 5). Selfishness, manipulation, and suppression are all greater when relatedness is lower. Both the two manipulative traits, and the two suppression traits converge to the same ESS values. Notably, because of the constraints on group productivity, where the sum of traits cannot exceed 1, a lower level of selfishness is selected in the scenario with two mechanisms while the levels of manipulation and suppression is roughly the same between the scenario with single mechanism and two mechanisms. In other words, because the additional manipulative mechanism is the same worthy to invest as the other one, we find a lower level of selfishness in the scenario with two mechanisms.

Appendix 1—figure 5. Selfishness, manipulations, and suppressions of manipulation in the
scenario with two mechanisms to manipulate and suppress manipulation, where all traits are
introduced at the same time point. Solid lines show the ESS predictions, and dots show the
results of our individual-based simulations ($a = b = 0.5$).

4.2 The occurrence of oscillation is related to shape parameters

In contrast to the close agreement shown above, we found some disagreement

between simulation and game-theoretical analyses when we analyse along shape

parameters (Appendix 1—figure 6). We used harmonic regression to test for

oscillating dynamics in our simulation data. In a harmonic regression, white noise

(randomly time series) would be a flat line because all frequencies have similar

predicted amplitude; but an oscillating pattern would lead to peak or peaks. For

example, a harmonic regression of sine wave would be mainly a single peak because

it only consists of a wave at single period. From the time series of selfishness, we

found oscillating dynamics at intermediate shape parameters ($a = b = 0.7$;

Appendix 1—figure 6c), but not at low and high shape parameters (Appendix 1—

figure 6b& d; $a = b = 0.4$ or 1.0). Specifically, a harmonic regression of selfishness

under low and high shape parameter values gave almost horizontal lines, whereas

intermediate values showed several peaks at periods around 500-1500 generations

(insets of Appendix 1—figure 6). These peaks suggest that there is a periodic

component to the evolutionary dynamics, with stochastic cycle that repeats about

every 500-1500 generations, and the dynamic is not random fluctuations.

The oscillations could occur because the benefits of manipulation and

selfishness become similar when shape parameters are at intermediate values, which

subsequently can be influenced by the interactions between traits. The decreased

difference in benefits is supported by a broader analysis, showing the discrepancy

between game-theoretical and simulative results only occurs when the former

predicts selfishness is lower than the level of manipulative cheating (Appendix 1—

figure 6a; $0.6 < a, b < 0.8$). When shape parameters are at these intermediate

values, the level of selfishness in simulations does not go below the level of

manipulation, as predicted by game-theoretical analysis (Fig. 5a, dots& curves).

Moreover, the simulated trait values are a lot more variable as the error bars become

visible (Fig. 5a; black vertical lines over dots).

1165

1166

Appendix 1—figure 6. Oscillation dynamics of selfishness and manipulation occur in some
ranges of shape parameters in the scenario with two mechanisms when relatedness is low ($R =$
1/8). (a) ESS predictions and simulation results from the model, where disagreement between
two approaches is found when shape parameters are between 0.6 and 0.8. (b-d) Time series of
simulations under various relatedness settings. Each panel shows the time series of traits, and
the analysis from harmonic regression is shown in the inset (same as Fig. 4). Harmonic
regression found the greatest amplitude in the period of 1385 generations and p-value of
1.5333×10^{-2} when shape parameters are 0.4; period of 815 generations and p-value of
1.8035×10^{-5} when shape parameters are 0.7; period of 1760 and p-value of 2.5202×10^{-3} when
shape parameters are 1.0.

1177

We can further examine the oscillations by checking the changes in maximal

amplitude along with shape parameters or evolved trait values (Appendix 1—figure

7). In agreement with our previous results (Appendix 1—figure 6b-d), the maximal

amplitude shows a peak when shape parameters are at intermediate values and

amplitude is low when the shape is either high or low (> 0.8 or <0.6). We are

interested in the difference in evolved trait values because oscillation seems to take

place when the evolved level of selfishness is close to the level at which

manipulation is expressed (Appendix 1—figure 6a). After we visualised the

relationship between trait difference and the maximal amplitude of oscillation, we

found a consistent negative correlation between the difference and amplitude. This

finding suggests oscillation might have resulted from interactions between traits

when their evolved levels are similar.

1190

1191

Appendix 1—figure 7. The relation of maximal amplitude from harmonic regression in response to (a) shape parameters, and (b) distance in evolved levels of selfishness and manipulation. Relatedness is set to 1/8 and the error bars are the 95% confidence intervals. Details of the analysis are described in Appendix 1—7.5.

4.3 Oscillating trait dynamics disappear when selfishness is not coevolving with

manipulation

In order to better understand the origin of oscillation, we reduce the complexity in

simulations by reducing the number of coevolving traits. In Appendix 1—figure 6 and

Appendix 1—figure 7, there are 5 traits coevolving: selfishness, manipulation 1,

suppression 1, manipulation 2, and suppression 2. We make 1 of the 5 traits non-

mutable at a time and analyse the evolving traits through harmonic regression. We

found oscillation disappears when selfishness is held constant, while holding one of

the two manipulative traits constant can still produce oscillating dynamics (Appendix

1—figure 8b& c). Moreover, the results are consistent regardless of whether the

traits are fixed at the average of coevolving dynamics or the prediction from game-

theoretical analysis (only the latter is plotted in Appendix 1—figure 7 to avoid

redundancy).

1210

Appendix 1—figure 8. Oscillation disappears when selfishness is controlled and fixed at a value. We use harmonic regression to instead analyse the dynamics of manipulation 2 in this figure, because selfishness and manipulation 1 are fixed in some panels. The level of fixed selfishness and manipulation are determined by the ESS predicted in game-theoretical analysis, where relatedness is 1/8 and shape parameters are 0.7. Harmonic regression found greatest amplitude in the period of 825 generations and p-value of 2.1045×10^{-4} when all traits coevolve; period of 1450 generations and p-value of 2.5239×10^{-4} when selfishness is fixed; period of 1575 and p-value of 5.5099×10^{-4} when manipulation 1 is fixed.

The above findings suggest the oscillation is likely to be an outcome from the linkage (interaction) between selfishness and manipulation. That is, some stochasticity may lead to deviations in trait values, for instance, increased selfishness and decreased manipulation. This perturbation would generate selective forces that favours decreased level of selfishness and increased level of manipulation. However, because the optimal trait values of both traits are similar to each other, the selective force would be weak, which allows trait values to fluctuate within a range of levels. Finally, when the optimal trait values are more differed and selective forces are strong, oscillation would only take place in a small range of parameter or even disappear. Such effects of linkage (interactions) are not discovered in game-theoretical analysis because inclusive fitness effects are estimated as independent factor (i.e., no linkage between traits). Similar complication in coevolving traits have also been observed in other studies involving individual-based simulations (Tibbetts *et al.* 2020; Liu *et al.* 2021).

4.4 Oscillation also disappears when traits are independently inherited without linkage
To further test our linkage hypothesis for the oscillations, we examined an artificial scenario where there could be no possible linkage. Specifically, we assumed that each offspring inherits each trait independently, possibly from different individuals (multiple asexual parents, a different one for each trait). In support of our hypothesis, we found that when there can be no linkage, that oscillations do not occur (Appendix 1—figure 9a & b). In addition, in this scenario, where there can be no linkage, the results of the simulation where in quantitative agreement with the

1244 analytical prediction (Appendix 1—figure 9c). This contrasts with our simulations

1245 results in Figure A6a, where oscillations occurred, and there was a greater

1246 quantitative disagreement between our analytical and simulation results. It also

1247 suggests that some disagreement between analytical and simulation results were

1248 due to the simulation allows for correlation among inherited traits to build over time.

1249

1250

1251 Appendix 1—figure 9. Oscillation does not occur when traits are inherited independently
1252 without linkage. (a) Time series of coevolving traits when shape parameters are intermediate,
1253 and relatedness is low ($R = 1/8$; $a = b = 0.7$). (b) The estimated amplitude of the time series
1254 of selfishness by harmonic regression. (c) analytical predictions and simulation results from the
1255 model without linkage of traits.

1256

4.5 Can oscillation happen when there are fewer or more mechanisms?

We then analyse whether oscillation can occur in cases with different number of

mechanisms. The idea is if the oscillation was caused by interacting selfishness and

manipulation, can oscillation still occur when there is only one mechanism to

manipulate, or when there are 3 mechanisms. As shown in Appendix 1—figure 10,

we found single-mechanism simulation cannot generate oscillation while the three-

mechanism simulation still has oscillations. This makes sense because multiple

mechanisms of manipulation can amplify the initially small perturbations and help

oscillation to be maintained in the trait dynamics.

1266

1267

Appendix 1—figure 10. Oscillation happens when there is more than one mechanism to manipulate and suppress. Here, relatedness is 1/8 and shape parameters are 0.75. Harmonic regression found greatest amplitude in the period of 1800 generations and p-value of 4.7899×10^{-3} in single-mechanism simulations; period of 1475 generations and p-value of 2.7475×10^{-5} in two-mechanism simulations; period of 390 and p-value of 1.4544×10^{-2} in three-mechanism simulations.

4.6 Does changing mutation rate lead to a different oscillation period?

In section 4.3 and 4.4 we have showed that the oscillation results from interactions and linkages between traits, consequently, we expect mutation rate should change the behaviour of oscillation. This is because mutation rate controls the speed of introducing new genotype to the population and the oscillation can be slower if mutation rate is lower. We verify this thought by running simulation with half of the original mutation rate (i.e., 5×10^{-3}, Appendix 1—7). We found supporting evidence as the oscillation of traits last longer per each cycle and have the greatest amplitude at a larger period close to 1500 generations (compare Appendix 1—figure 11 and Fig. 4).

1286

Figure A11. Selfishness and manipulative cheating traits oscillate slower when mutation rate is lower. In this figure, we set mutation rate is half the size of which in all other simulation ($R = 1/8$; $a = b = 0.7$).

1290

4.7 Can we find oscillation in the personal cost model?

In section 3, we have discussed that implementing personal cost to manipulation and

suppression would not qualitatively change the predictions, but can we still get

oscillation in these models? We made a two-mechanism model with personal cost

and found out there is no oscillation (Appendix 1—figure 12b). The underlying

reason is the evolved level of selfishness is much higher than the level of

manipulation (Appendix 1—figure 12a). Despite the model design is different, the

relation between maximal amplitude and trait difference seems to remain

unchanged by comparing Appendix 1—figure 12c and Appendix 1—figure 7b.

1300

1301 Appendix 1—figure 12. (a) The simulated level of selfishness, manipulative cheating, and
1302 suppression, (b) the relation between maximal amplitude from harmonic regression and shape
1303 parameters ($a = b$), and (c) the relation between maximal amplitude and trait difference in
1304 evolved selfishness and evolved manipulation.

1305

4.8 Does recombination change the simulative results?

So far, we have assumed that each set of trait values are carried by single individual

across generations, but it might not be the case when recombination take place,

where genes may be exchanged between individual cells. We address this potential

biological complexity by setting an additional recombination phase in the life cycle of

individual-based simulations. During the recombination phase, individuals are

selected according to relative fitness in the previous generation, paired with another,

and then recombine one trait value depending upon the recombination probability

(externally set parameter). Only one trait value is exchanged, and individual

recombines only once. After the possible recombination, the two individuals are put

into a pool of founders. Once all individuals are in the founder pool, and a specific

number of individual (m, Appendix 1—7) are sampled as founder to start the new

population.

1319

1320
1321 Appendix 1—figure 13. Effects of recombination on the simulated level of selfishness,
1322 manipulation, and suppression. Here, each panel has a specific relatedness setting (R= 0.5, 0.25,
1323 or 0.125), and we simulated five values of recombination rate (0, 0.25, 0.5, 0.75, and 1).

1324

We examined the effects of recombination at the parameter setting the same as

Fig. 2 and found recombination does not affect the results ($a = b = 0.5$; Appendix

1—figure 13). This is likely because all social interactions, including manipulation and

suppression, are reset at the beginning of each generation that nothing in the past

generations is passed on the future generation. Recombination could matter in

scenarios where associations between traits can be important and potentially build

up over time, such as whether punishment can favour cooperation (Gardner & West

2004; Gardner *et al.* 2007; Lehmann *et al.* 2007; Press & Dyson 2012; Hilbe *et al.*

2013).

1334

5. Arms race under different biological settings

We construct two different scenarios to understand if alternative cost design can still

generate arms-race-like dynamics. In particular, we modified group productivity so

that the cost of suppression is closer to a switch-on/switch-off design. We assume

suppression becomes costly only when the group average is above 0.05, and the cost

is the same for all suppression above 0.05. This modification allows the share of

group productivity and group productivity, to disentangle with each other and relax

the linkage assumed in the main model. On top of whether the cost depends

continuously on the degree of suppression, the cost can be conditional and only

present when there are sufficient levels of manipulation. Thus, we assume the cost

of suppression is unconditional in one case, and conditional to the average level of

manipulation in the other case. For simplicity, we do not randomly introduce each

new mechanism but mutate them at fixed time points with equal intervals.

1348

We found the genetic arms race of manipulation and suppression still occur in

both cases, but with different dynamics of suppressive traits. When the cost is

unconditional and fixed, we found all of the suppressions go to almost 1 when the

shape parameter for manipulation and suppression are equal (a=b=0.5; Appendix 1—

figure 14). This is because the effect of suppression is relatively linear and it does not

cost extra if suppression has higher values (compare Appendix 1—figure 14c & Fig.

5c). When the cost is conditional to manipulation, we found a drift-like dynamics for

all suppressive traits (a=0.5, b=0.01; Appendix 1—figure 15). The underlying reason is

suppressions are cost-free and no longer under selection when manipulations are

completely suppressed (Appendix 1—figure 15b & c). The cost-free suppressions are

support by the dynamics of selfishness, which is not gradually decreasing like other

cases (compare Appendix 1—figure 15a with Fig. 5a& Appendix 1—figure 14a).

Nevertheless, the dynamics of manipulation are very similar in all cases, thus

suggesting genetic arms race is a robust result which can be found in different

biological settings (Fig. 5, Appendix 1—figure 14, and Appendix 1—figure 15).

1364

1365

Appendix 1—figure 14. Arms race with an unconditional and fixed amount of cost for suppressing manipulative cheating. Suppression is costly in group productivity when it exceeds 0.05, and the cost is fixed at 0.01 regardless of the trait value of suppression. Shape parameters are equal for manipulation and suppression ($a = b = 0.5$) and relatedness is $1/3$.

Appendix 1—figure 15. Arms race with a conditional and fixed cost for suppression. Suppression is costly only when both manipulative cheating and suppression are above 0.05, and the cost of suppression is fixed at 0.05 regardless of the trait value. Shape parameters are the same as Fig. 5 ($a = 0.5$ and $b = 0.01$) and relatedness is 1/3.

6.   Additional discussion

6.1 Decelerating return and the nature of communication

Contrary to the linear effects assumed in the classic tragedy of the commons, empirical studies have shown that nonlinear effects might be common in communications between cells and individuals. This is because the limitation of diffusion in communicative molecules, making the benefit of increased production in those molecules diminishes (Müller *et al.* 2006; Wiley 2013; Rogers *et al.* 2017);

same principle can be applied to social insects, as increased time in communicating

would also lead to increased possibility in encountering previously-interacted

individuals (Gill & Johnson 2002). As manipulation and suppression require such

short-distance interaction, we have considered the nonlinear impact of traits on

fitness to make the model more realistic. In particular, manipulative, and suppressive

traits may have larger gains when they evolved from absence to some intermediate

extent of effects, whereas the benefit of complete manipulation or suppression is

diminishing. Similar saturating property has been found in other social traits, such as

the benefit in increased stalk/spore ratio in slime moulds (Bonner 1982; Hudson *et*

al. 2002). Thus, we focus on cases where nonlinear effects have linear or diminishing

returns ($0 < a \leq 1$ and $0 < b \leq 1$). For simplicity, we only add nonlinearity effect

to individual reproduction component and not the group productivity component

(i.e., $1 - \bar{z} - \bar{x} - \bar{u}$).

1396

6.2 Methodological details of numerical analysis in game theory results

For a given number of iteration t, we update trait values by $\{z_{t+1}, x_{t+1}, u_{t+1}\} =$

$\{z_t + c_{scale}\Delta W_z, x_t + c_{scale}\Delta W_x, u_t + c_{scale}\Delta W_u\}$ and find the trait values that

converge to fixed values. All traits are initialised at 0.1. We set the scaling coefficient

c_{scale} to 2.5×10^{-3} for better precision. Finally, due to slow convergence when

relatedness is low, we iterated 4×10^6 times for $0.1 \geq R$ and 4×10^6 times for

the rest of relatedness values.

1404

7. Individual-based simulations

7.1 Overview

We use individual-based simulations following genetic algorithm to understand the

evolution of selfishness, manipulative cheating, and suppression of manipulation. In

contrast to the game-theoretical analysis, individual-based simulations allow us to

explicitly model mutations and trait dynamics under a continuous trait space.

Following the formal analysis (section 1), we employ similar assumptions that the

population size is constant, and generations do not overlap.

1413

After initialisation, the biological processes in each generation take place in the

following order: First, groups are formed from several (l) founders where the number

of founders determines the relatedness ($R = 1/l$). Each group has the same number

of founders, and it is a parameter for population structure. Second, the founders

expand the social group to a size that social interactions take place, where each

founder replicate the same number of times. Third, we calculate the group averages

of trait values and the relative fitness of each individual. Fourth, the new-born

founders are randomly drawn from the current population according to the relative
fitness. Each new-born founder inherits its parent's trait values with independent
chances of mutation for each of the three traits.
1424
1425   We use the design with founders and group growth to avoid the potential
1426 effects of small integers and small group sizes (i.e., demographic stochasticity in
1427 social groups). This simulation framework is known as haystack models (Maynard
1428 Smith 1964).
1429
7.2 Initialisation
We initialise the population by creating a population with uniform group size (n) and
uniform properties of no selfish traits (i.e., $z = x = u = 0$). In order to reduce the
effect of demographic stochasticity, we hold the size of the whole population and let
the number of groups change because group size (n) is explicitly modelled in the
model. Specifically, we set population size parameter (N) and obtain the number of
groups as $M = \lceil N/n \rceil$, the least integer larger than or equal to N/n. Thus, the
population size, Mn, is relatively constant while the number of groups increases as
the group size decreases.
1439
1440   Each individual has three variables: the degree of selfishly increase own fitness
1441 share at the cost of entire group (z), the degree of manipulative cheating (x) and the
1442 degree of suppressing manipulative cheating (u). Within one group, there are $m =$
1443 n/l individuals that have identical trait values because they are replicated from the
1444 same founder. Further, trait values could only be changed in *mutation* process.
1445
7.3 Fitness calculation and the coevolving traits
We perform two levels of selection to choose founders for the next generation: (i)
group selection according to group productivity from average trait values of all group
members, and (ii) individual selection according to each group members' relative
fitness (Fig. 1). In other words, the probability that a founder comes from a group
depends on the group's productivity, $1 - \bar{z} - \bar{x} - \bar{u}$, and the chance that it is an
offspring of the focal individual is determined by its relative fitness in the group (or
share or productivity), $\frac{z_i + (1-u_{-i})z_{-i}x_i - (1-u_i)z_i x_{-i}}{\bar{z}}$. Although the simulations calculate
fitness in the same way as the game-theoretical analysis, the process for each event
is explicitly modelled in the simulations and individuals can have any trait values,
whereas the game-theoretical analysis tracks on the trait dynamics of the entire
population under weak selection assumption.

1458

1459   *7.4 Mutation of the coevolving traits*

1460   We assumed the traits can mutate between generations. For each trait of a new

1461   founder, the mutation rate is $p_{mut} = 0.01$. If a mutation occurs, we perturb the

1462   parental trait value by a normally distributed deviation with mean of 0 and standard

1463   deviation of 0.1, constrained at the boundaries [0,1].

1464

7.5 Analysis of the simulations

In all cases, we repeated the simulation 10 times, and let social interactions take

place at the population around 10^5 individuals (i.e., $lm\lceil 10^5/lm \rceil$) to ensure that the

trait values converged to their evolutionary optima. We introduced mutation to

selfishness from the start of simulation, manipulative cheating from 100 generation,

and suppression from 200 generation to make sure the latter traits are not evolved in

the absence of selfishness. We tracked the trait dynamics for 3×10^3 generations

(unless specified elsewhere) and average over the last 10% to calculate the average

and standard deviation of trait values. Except for the ones with oscillation dynamics,

the coevolving traits usually stabilise within 500 generations (e.g., Appendix 1—

figure 6b & d).

1476