1 **The inner nuclear membrane protein NEMP1 is required for nuclear envelope**

2 **openings and enucleation of erythroblasts during erythropoiesis**

3

4 <u>**Short title**</u>**: NEMP1 supports nuclear envelope openings of erythroblasts**

5

Didier Hodzic[1, #,*], Jun Wu[2,*], Karen Krchma[2], Andrea Jurisicova[3], Yonit Tsatskis[3], Yijie

Liu[4], Peng Ji[4], Kyunghee Choi[2, #] and Helen McNeill[1, #]

[1]Department of Developmental Biology, [2]Department of Pathology and Immunology,

Washington University School of Medicine, 660 S. Euclid Ave, St Louis, MO, 63110;

[3]Lunenfeld-Tanenbaum Research Institute, Sinai Health Systems, 25 Orde St., Toronto,

ON; [4]Department of Pathology, Feinberg School of Medicine, Northwestern University,

303 East Chicago Avenue, Chicago, IL, 60611.

[*]Equal contributions

[#]Corresponding authors

[#]Helen McNeill**,** Department of Developmental Biology, Washington University School of

Medicine, 660 S. Euclid Ave, St Louis, MO, 63110. <u>mcneillh@wustl.edu</u>, phone: 314-

273-3050

[#]Kyunghee Choi, Department of Pathology and Immunology, Washington University

School of Medicine, 660 S. Euclid Ave, St Louis, MO, 63110, <u>kchoi@wustl.edu</u> , phone:

314-362-8716

#Didier Hodzic, Department of Developmental Biology, Washington University School of

Medicine, 660 S. Euclid Ave, St Louis, MO, 63110, dhodzic@wustl.edu, phone: 314-

273-3051

AUTHOR CONTRIBUTIONS

H.M., K.C., P.J. A.J., D.H. and J.W. conceived and designed experimental approaches

and participated in manuscript writing and review; D.H., J.W., K.K, Y.B. and Y.L

performed experiments, data collection and assembly. All authors participated in data

analyses and have proved the final version of the manuscript.

ABSTRACT

Nuclear Envelope Membrane Proteins (NEMP) are a conserved family of nuclear

envelope proteins that reside within the inner nuclear membrane. Even though *Nemp1*

knockout (KO) mice are overtly normal, they display a pronounced splenomegaly. This

phenotype and recent reports describing a requirement for nuclear envelope openings

during erythroblasts terminal maturation led us to examine a potential role for Nemp1 in

erythropoiesis. Here, we report that *Nemp1* knockout (KO) mice show peripheral blood

defects, anemia in neonates, ineffective erythropoiesis, splenomegaly and stress

erythropoiesis. The erythroid lineage of *Nemp1* KO mice is overrepresented until the

pronounced apoptosis of polychromatophilic erythroblasts. We show that NEMP1

localizes to the nuclear envelope of erythroblasts and their progenitors. Mechanistically,

we discovered that NEMP1 accumulates into aggregates that localize near or at the

edge of nuclear envelope openings and *Nemp1* deficiency leads to a marked decrease

of both nuclear envelope openings and ensuing enucleation. Together, our results for

the first time demonstrate that NEMP1 is essential for nuclear envelope openings and

erythropoietic maturation *in vivo* and provide the first mouse model of defective

erythropoiesis directly linked to the loss of an inner nuclear membrane protein.

INTRODUCTION

Nuclear Envelope Membrane Protein1 (NEMP1, encoded by *Tmem194a*) is a highly conserved multipass transmembrane protein that resides within the inner nuclear membrane (INM) of the nuclear envelope (NE). We recently showed that genetic inactivation of *Nemp1* leads to a loss of fertility in worm, fish, and flies [1]. In mice, *Nemp1* is required for female fertility but dispensable for male fertility [1]. Except for the presence of a conserved domain of unknown function (DUF 2215) that encompasses its transmembrane and proximal nucleoplasmic C-terminal region, NEMP1 does not harbor any known functional motif. However, its nucleoplasmic region has been shown to interact with barrier to autointegration factor (BAF) and RAN GTPase to mediate *Xenopus* eye development [2,3]. NEMP orthologs have recently been identified in plants (PNET2a, b and c) where they play an essential role in chromatin architecture [4]. Using BioID as well as affinity purification followed by mass spectrometry, we recently showed that NEMP1 interacts with LEM domain proteins EMERIN, MAN1 and LAP2, known to physically link the NE to chromatin and support mechanical stiffness. Accordingly, we showed that loss of *Nemp1* expression drastically affects NE mechanical stiffness in cultured cells and oocytes [1].

Mammalian erythropoiesis consists of the differentiation of hematopoietic stem cells into megakaryocyte-erythrocyte progenitors (MEPs) that generate burst forming unit-erythroid (BFU-E) that in turn differentiate into colony forming unit-erythroid (CFU-E). The latter generate proerythroblasts (ProE) that correspond to the first recognizable erythroid cell. During terminal erythropoiesis, ProE undergo 4-5 mitoses that generate

basophilic (EryA), polychromatophilic (EryB) and orthochromatic (EryC) erythroblasts.

Erythroblast differentiation is characterized by chromatin condensation which is required

for enucleation, the ultimate step of erythropoiesis that generates pyrenocytes and

reticulocytes [5-8]. Interestingly, recent studies have established that recurrent NE

openings in maturing erythroblasts allow for the partial and selective release of histones

in the cytoplasm, a biological process that is essential for chromatin condensation and

final enucleation[9-12]. However, the role of NE proteins in this remarkable biological

process remains to be established.

Adult *Nemp1* knockout (KO) mice are overtly normal. However, both *Nemp1* KO males

and females display strikingly enlarged spleens. This phenotype and the involvement of

NE openings in terminal erythropoiesis led us to examine the biological function of *Nemp1*

in erythropoiesis. We show that *Nemp1* KO mice display erythroid lineage differentiation

defects. Polychromatophilic erythroblasts displayed reduced frequencies of NE openings

and of enucleation as well as increased apoptosis, leading to erythroid maturation

defects. These data show that NEMP1 is required for NE openings and enucleation during

the late stages of erythroblast maturation.

RESULTS

***Nemp1* KO mice have splenomegaly and abnormal erythropoiesis**

Nemp1 KO mice displayed significantly enlarged spleens with increased cellularity

compared to heterozygous *Nemp1* or wild type (WT) mice (Figure 1A, B, C). Wright-

Giemsa staining of *Nemp1* KO blood smears showed red blood cells (RBCs) with irregular

shapes and spiky membranes (Figure 1D, red arrows). Complete blood count (CBC)

analyses of peripheral blood (PB) using a Hemavet further showed decreased RBC

counts in neonates suggestive of anemia although adult mice showed no such obvious

phenotype (Figure 1E and S1). RBCs showed decreased hemoglobin content and higher

red cell distribution widths (RDW), reflecting irregular RBC membrane shapes, two

phenotypes that persisted throughout life (Figure 1E and S1). Finally, FACS analysis of

PB also showed increased percentage of immature RBCs in the circulation (Figure 1F).

The bone marrow (BM) of *Nemp1* KO mice also appeared more densely packed and

displayed increased cellularity by comparison to WT BM (Figure 1G).

To understand the nature and origin of erythroid defects, we quantified the erythroid

population of BM and spleens from adult WT and *Nemp1* KO mice by using Ter119 and

CD71 markers. Compared to WT BM, *Nemp1* KO BM showed increased Ter119+ cell

population (Figure 1H) that was mostly accounted for by a significantly increased

population of Ter119+CD71+ erythroblasts (Figure 1I). The increase in erythroblasts was

also detected by flow imaging of Hoechst+Ter119+ populations (Figure 1J). In the adult

spleen, FACS and flow imaging data showed similar increases in the representation of

erythroblast population (Figure 1 K, L, and M). Taken together, these data show that loss

of *Nemp1* leads to a significant increase of the erythroid lineage in BM and spleens.

Figure 1: Nemp1 KO mice display splenomegaly, blood defects and erythroid lineage

expansion. *Representative spleen size (**A**) and spleen/body weight ratios (**B**) across genotypes.*

*Scale bar: 1 cm. **C**. H&E staining of WT and Nemp1KO spleen. **D**. Wright-Giemsa staining of WT*

and Nemp1 KO blood smears. Red arrows point to cytoplasmic membrane spikes of Nemp1 KO

*RBCs. **E**. Hemavet measurement of red blood cells (RBC), red blood cells distribution width*

(RDW) and hemoglobin (Hg) content of Nemp1KO blood from postnatal day 4 (P4) and P10. 6,

3, 5 and 4 mice were respectively analyzed for RBC, Hb and RDW measurements for P4 WT, P4

*KO, P10 WT and P10 KO mice. **F**. Percentage of immature nucleated erythroblasts in peripheral*

*blood of indicated genotypes. **G**. H&E staining and quantification of cellularity of WT and*

*Nemp1KO BM. **H, I**. Percentage of Ter119+ (H) and Ter119+/CD71+ (I) cells in WT and Nemp1*

*KO BM assessed by FACS. **J**. Percentage of nucleated Ter119+/Hoechst+ cells in WT and*

*Nemp1 KO BM measured by flow imaging. **K,L**. Percentage of Ter119+ (H) and Ter119+/CD71+*

*(I) cells in WT and Nemp1 KO spleen assessed by FACS. **M**. Percentage of nucleated*

Ter119+/Hoechst+ cells in WT and Nemp1 KO spleen measured by flow imaging. Data are shown

*as mean ± SD. Student's t-test. ns. not significant, *p<0.05. **p<0.01. ***p<0.001. ****p<0.0001.*

Erythroid progenitors and early erythroblast populations are expanded in *Nemp1*

KO mice

To trace the origin of erythroid lineage overrepresentation in *Nemp1* KO mice, we

quantified hematopoietic progenitors. Long-term hematopoietic stem cells (LT-HSCs,

SLAM-KSL), hematopoietic stem and progenitor cells (HSPCs, KSL, ckit+/Sca1+/Lin-) as

well as common myeloid progenitors (CMP, CD34$^+$CD16/32$^-$Lin$^-$c-Kit$^+$Sca1$^-$) were equally

represented in WT and *Nemp1* KO BM (Figure 2 A,B,C). In contrast, the megakaryocyte-

erythrocyte progenitor (MEP, CD34$^-$CD16/32$^-$Lin$^-$c-Kit$^+$Sca1$^-$) population was significantly

increased with a concomitant decrease of the granulocyte-macrophage progenitor (GMP,

CD34$^+$CD16/32$^+$Lin$^-$c-Kit$^+$Sca1$^-$) population (Figure 2C) in *Nemp1* KO BM. In agreement

with this increase in MEPs, *Nemp1* KO BM cells consistently showed a higher capacity

to generate BFU-E (Figure 2D). Together, these results indicate that the earliest

phenotype resulting from the lack of *Nemp1* expression is the expansion of MEPs in the

hematopoietic cascade.

To better understand the erythroid differentiation defects in *Nemp1* KO mice, we next

examined the representation of ProE, EryA, EryB, and EryC erythroblasts using the gating

strategy shown in Figure 2E. Consistent with increased MEP population and higher BFU

formation capacity in *Nemp1* KO BM (Figure 2C, D), the ProE and EryA populations were

increased, with EryA showing a more significant increase (Figure 2F, G). By contrast, the

EryB and EryC were decreased, with EryB showing a more significant decrease, in the

Nemp1 KO BM. Whereas apoptosis was mildly reduced in EryA, the apoptotic EryB

population was significantly higher in *Nemp1* KO BM (Figure 2H). Similar trends were

also observed in the spleen (Figure 2I, J, and K). Collectively, these data indicate that

genetic ablation of *Nemp1* leads to the expansion of MEPs to EryA populations and to a

decrease of EryB and EryC populations. Increased apoptosis in EryB probably accounts

for the loss of RBC.

Figure 2: Loss of Nemp1 leads to the expansion of erythroid progenitors and early

erythroblasts*. Quantification of LT-HSCs (**A**), HSPCs (**B**), CMPs, GMPs and MEPs progenitors*

*(**C**). **D**. BFU-E formation capacity of WT and Nemp1 KO BM. **E.** FACS sorting strategy used to*

quantify ProE and Ery A, B, C populations from whole BM and spleen. Representative FACS

*analyses and quantification of ProE, EryA, B and C populations in WT and Nemp1KO BM (**F, G**)*

*and spleens (**I, J**). Apoptotic indexes of the same respective populations in WT and Nemp1 KO*

*BM (**H**) and spleen (**K**). Data are shown as individual subject and the mean ± SD. Student's t-*

test. ns. not significant, *p<0.05. **p<0.01. ***p<0.001. ****p<0.0001.

Given the compromised erythropoiesis in the BM, we suspected that spleen enlargement

in *Nemp1* KO mice might be due to stress erythropoiesis. Accordingly, c-

kit+/CD71+/Ter119- stress erythroid progenitors (SEPs) and Epo-responsive BFU-E and

CFU-E were significantly increased in Nemp1KO spleen and splenectomy experiments

also pointed to a significant contribution of splenic SEPs to ongoing erythropoiesis in

Nemp1 KO mice (S2 text and S2 Fig). Collectively, these data show that genetic ablation

of NEMP1 affects erythroid maturation, especially from EryA to EryB, and elicit stress

erythropoiesis.

NEMP1 is required for erythroblast NE openings, nuclear compaction and nuclear extrusion

Using a NEMP1 antibody directed towards a stretch of 15 amino acids from the C-terminal

region of NEMP1 (Figure S3A) [1], NEMP1 (49.8 kDa) was detected both in lysates of

sorted WT CD11b+ myeloid and Ter119+ erythroid cells but not in their KO counterparts

(Figure 3A). In immunofluorescence confocal microscopy, NEMP1 was detected at the

NE of ProE, EryA, B and C erythroblasts where it colocalized with LAP2, a well-

established NE marker (Figure 3B). However, NEMP1 noticeably formed occasional NE

puncta of higher intensity whereas LAP2 was homogenously distributed on the NE of

erythroblasts (Figure 3B). NEMP1 was also detected at the NE of cKit+/Ter119-

progenitors (Figure S3B). NEMP1 was undetectable at the NE of BM cells isolated from

Nemp1 KO mice demonstrating the specificity of the NEMP1 antibody (Figure S3C).

Taken together, these results show that NEMP1 is ubiquitously expressed at the NE of

the erythroid lineage.

Late stages of erythroblast maturation are characterized by the progressive compaction

of chromatin and its partial release into the cytoplasm via transient openings of the NE,

which is most prominent in the polychromatophilic stage [9,10]. As shown in Figure 3C,

NE openings were clearly identified with NEMP1 and LAP2 antibodies in Ter119+

erythroblasts. This phenomenon is distinct from nuclear extrusion as the cytoplasmic

membrane labelled with Ter119 remains intact (Figure 3C). 3D reconstruction of confocal

Z-stacks clearly emphasized NE openings delineated by NEMP1 through which

chromatin protrudes into the cytoplasm (Figure 3D and S1 Video).

Interestingly, the close examination of successive confocal planes encompassing NE

openings revealed the accumulation of NEMP1 into higher intensity aggregates (Figure

3E). Maximum intensity projections also showed that NEMP1 aggregates preferentially

accumulated near or close to NE openings (Figure 3E, bottom panel). In contrast, the

localization of LAP2 remained uniform and did not accumulate into NEMP1 aggregates.

Intensity profiles showed that NEMP1 was 2 to 3 times more abundant in aggregates at

NE openings by comparison to intact NE (Figure S3D). To better appreciate the spatial

distribution of Nemp1 aggregates relative to NE openings in three dimensions (3D), we

performed intensity thresholding of confocal slices followed by 3D reconstruction and α-

blending. As shown in Figure 3F, multiple NEMP1 aggregates localized close to or at the

edges of NE opening sites delineated by Lap2 (Figure 3F, top and video S2). Interestingly,

small NE openings already displayed a few NEMP1 aggregates at their edges (Figure 3F,

bottom panel and video S3). NEMP1 aggregates were undetectable at the NE of *Nemp1*

KO erythroblasts thereby confirming the specificity of these structures (Figure S3E).

To determine if NEMP1 is required for NE openings, WT and *Nemp1* KO BM were

immunostained with Lamin B1. Loss of *Nemp1* expression led to a significant decrease

of NE opening frequencies suggesting a role for NEMP1 in NE openings (Figure 3G). We

also observed a significant increase of nuclear size in *Nemp1* KO BM cells that may be

indicative of decreased chromatin compaction (Figure 3H).

Figure 3: Nemp1 is required for NE openings in erythroblasts. A. *Immunodetection of*

endogenous NEMP1 in lysates from WT and Nemp1 KO myeloid (CD11b+) and erythroid

(Ter119+) cells. Histone3 was used as a loading control and Ter119 to confirm myeloid/erythroid

sorting specificity. ***B****. Nemp1 immunostaining of WT BM erythroblasts. Lap2 was used as a control*

for NE labeling. ProE were identified based on low Ter119 expression and low nuclear circularity.

Ery A, B and C were distinguished based on increasing Ter119 intensity and decreasing nuclear

diameter (7.4, 6.4 and 5.4 μm for EryA, B and C, respectively). The arrow points to discrete

Nemp1 aggregates. ***C.*** *Immunolocalization of Nemp1 and Lap2 in a WT erythroblast undergoing*

NE opening and chromatin release into the cytoplasm. ***D.*** *3D reconstruction (non-thresholded) of*

single confocal planes of a NE opening delineated by Nemp1 after denoising and α-blending.

Note the extrusion of chromatin through the NE openings within the cytoplasm. ***E.*** *Successive*

confocal planes (0.5 μm apart) of an erythroblast undergoing NE opening showing Nemp1

aggregates at NE opening sites. Note the presence of higher intensity aggregates of Nemp1 at

or near NE openings delineated by Lap2. Bottom panel: Maximum intensity projection (Max I) of

*a Z-stack encompassing the NE opening. **F**. 3D reconstruction after α-blending of confocal slices*

thresholded for Nemp1 intensity showing Nemp1 aggregates at or near to NE openings. The

*bottom panel shows a small NE opening already decorated with Nemp1 aggregates. **G**. Confocal*

imaging of LaminB1 in WT and Nemp1 KO BM cells and quantification of NE opening frequencies.

***H**. Quantification of nuclear area in WT and Nemp1 KO BM cells.*

Because chromatin compaction is required for enucleation [9], we next examined whether

NEMP1 is required for enucleation. As shown in Figure 4A, NEMP1 decorated the NE

with occasional puncta at all described stages of enucleation [13]. We used two

approaches to determine whether genetic ablation of *Nemp1* affects nuclear extrusion.

First, lineage negative cells from wild type and *Nemp1* KO mice were purified from BM,

cultured for 2 days in erythropoietin-containing medium and analyzed to distinguish

nucleated vs non-nucleated Ter119+ cells. As shown in Figure 4B, although erythroblast

differentiation was not affected, the ratio of nucleated erythroblasts (DNA+/Ter119+) vs

RBC (DNA-/Ter119+) was increased in cultures derived from *Nemp1* KO BM. In

agreement with these data and using flow imaging as a second approach, we consistently

measured a higher ratio of nucleated erythroblasts (Hoechst+/Ter119+) vs RBC (DNA-

/Ter119+) in *Nemp1* KO BM by comparison to WT BM cells (Figure 4C). Taken together,

these results indicate that NEMP1 plays a role in NE openings and nuclear extrusion.

***Figure 4: Nemp1 is required for erythroblasts enucleation. A**. Immunolocalization of Nemp1*

*at different stages of enucleation. Scale bars: 5 μm. **B**. DNA and Ter119 FACS analysis of lineage-*

negative cells purified from WT and KO BM and subsequently cultured in erythropoietin-

*containing medium for 2 days. Note the higher nucleated/enucleated ratio in Nemp1 KO cells. **C**.*

Quantification of nucleated and enucleated Ter119+ cells in WT and Nemp1 KO BM measured

by flow imaging. Right: representative quantification graphs of nucleated (Hoechst+Ter119+) and

enucleated (Hoechst-Ter119+) in WT and Nemp1KO BM in flow imaging.

DISCUSSION

In this work, we show that loss of the integral transmembrane NE protein NEMP1 results

in erythropoietic defects consisting of the expansion of the erythroid lineage in BM and

spleens and erythroid maturation defects during terminal erythropoiesis (Figure 5). As a

consequence, *Nemp1* KO mice display anemia in neonates and splenomegaly associated

with stress erythropoiesis in adult mice.

Figure 5: Graphical depiction of Nemp1 phenotypes and functions during erythropoiesis.

Green and red arrows respectively depict increased or decreased populations in Nemp1KO by

comparison to WT cells or biological processes affected by the lack of expression of NEMP1.

Blue arrows depict increased apoptosis. See text for details.

The higher capacity of *Nemp1* KO BM and spleens to generate BFU-E and CFU-E and

the significant expansion of ProE and EryA populations together show that erythroid

expansion is taking place throughout the EryA stage in *Nemp1* KO mice. MEP expansion

was notably associated with a significant decrease in GMPs, suggesting that *Nemp1* may

have an additional biological function in non-erythroid lineages. Alternatively, GMP

decrease may reflect compensatory mechanisms in response to erythroid lineage

expansion.

We speculate that the overrepresentation of erythroid progenitors and EryA as well as the

stress erythropoiesis we observed in spleen originate from a feedback loop due to the

massive apoptotic loss of EryB erythroblasts. Indeed, in contrast to their progenitors and

EryA precursors, EryB and EryC populations were markedly decreased in *Nemp1* KO BM

and RBC counts were significantly lower in the peripheral blood of neonate and young

Nemp1 KO mice. This decrease of EryB and EryC populations was accompanied by a

marked increase of apoptosis that was especially pronounced in EryB from BM. We

reason that this apoptosis stems from key biological functions of NEMP1 in NE openings

during terminal erythropoiesis. First, *Nemp1* transcripts are significantly upregulated

during terminal differentiation [14] and *Nemp1* expression level peaks in

EryB/polychromatophilic erythroblasts [15]. Second, in erythroblasts undergoing NE

openings, NEMP1 accumulates into aggregates that preferentially localized near or at NE

openings. Importantly, this aggregation is specific to NEMP1 as LAP2 remains uniformly

distributed at NE openings. It is possible that increased levels of NEMP1 expression

reported in proteomic screens at that differentiation stage [15] reflects the accumulation

of Nemp1 into aggregates. Finally, we directly show that lack of *Nemp1* expression in

erythroblasts is linked to reduced frequencies of NE openings. Taken together, we

propose that the increased apoptosis measured in *Nemp1* KO EryB and EryC stems from

the requirement of NEMP1 aggregates for efficient NE openings. In support of this idea,

inhibition of NE openings is directly linked to induction of cell death in G1ER cells [9].

An alternative but not exclusive origin of elevated apoptosis in EryB erythroblasts may

also stem from a biological function of NEMP1 in chromatin organization. Indeed, we

recently reported the interaction of NEMP1 with LEM domain-containing proteins

EMERIN, LAP2 and MAN1 [1] that are also expressed in erythroblasts [15] and play

essential roles in chromatin organization in cooperation with BAF [16]. In addition, the

genetic ablation of NEMP proteins in plants (PNET2) leads to major disruptions of higher-

order chromatin organization [4] and *Nemp1* KO erythroblasts displayed larger nuclear

sizes. To that regard, it is interesting to note that the genetic inactivation of LAP2α, a

LAP2 isoform devoid of transmembrane domain and involved in the stabilization of higher

order chromatin [17], also leads to the overrepresentation of the erythroid lineage [18].

Impaired nuclear extrusion in *Nemp1* KO erythroblasts may also contribute to increased

apoptosis levels but future studies are needed to determine whether NEMP1 intrinsically

affects nuclear extrusion or if decreased enucleation in *Nemp1* KO BM is a mere

consequence of impaired NE openings.

We detected the expression of NEMP1 at the NE of erythroblasts and cKit+ progenitors

by immunofluorescence microscopy and in purified Ter119+ erythroid cells by

immunoblotting which is in agreement with its identification in proteomic analyses of the

human erythroid cells[15]. By contrast, the same proteomic analyses show that human

erythroblasts do not express *Nemp2*, a homolog of *Nemp1* [15]. Accordingly, *Nemp2* KO

mice do not display any splenomegaly or blood defects, and *Nemp1* phenotypes were not

worsened in *Nemp1/2* double knockout (DKO) mice (data not shown). Taken together,

we conclude that *Nemp1* is specifically required for the normal homeostasis of the

erythroid lineage.

The precise function of NEMP1 aggregates at NE openings requires more investigation.

Biochemical studies showing that Nemp1 oligomerizes via its transmembrane domains

[3] suggest that NEMP1 may aggregate through multimerization upon increased

expression levels detected in transcriptomic and proteomic studies [14,15]. Because we

know that NEMP1 supports NE stiffness in the germline and cultured cells[1], it is possible

that Nemp1 aggregates mechanically support the nuclear envelope during the physical

stresses of NE openings.

To our knowledge, this is the first report describing the requirement for an integral

transmembrane protein of the NE in erythropoiesis. Indeed, pathologies linked to

mutations of NE proteins or the nuclear lamina, globally termed "nuclear envelopathies"

and "laminopathies", have not previously been associated to erythropoietic

defects[19,20].   Similarly to the involvement of multiple NE proteins in specific

pathologies, Nemp1 deficiency specifically underlies female infertility and erythropoietic

defects in mice. This further fuels the concept of tissue-specific composition of NE

proteins whose mutations results in tissue-specific pathologies[21].

Repetitive NE openings during terminal differentiation provides a relatively little-known

yet powerful opportunity to study NE dynamics and remodeling *in vivo*. To that regard,

Zhao et al.[7,9] have shown that NE openings are blocked by inhibition of caspase-3 or

through the expression of a caspase-3 non-cleavable Lamin B1 mutant. As a result,

histone release from the nucleus, chromatin condensation and the terminal differentiation

of erythroid cells are also affected *in vitro*. These data and our current findings therefore

further stress the biological relevance of NE openings during terminal erythropoiesis.

Finally, because Caspase-3 KO mice show relatively mild erythroid defects most likely

due to *in vivo* compensatory pathways[9], *Nemp1* KO mice provide the first mouse model

of acute erythropoietic defects linked to NE openings deficiency. In conclusion, our results

uncovered the involvement of *Nemp1* in NE openings and enucleation in erythroblasts

and its requirement for normal erythropoiesis.

MATERIAL AND METHODS

Animals

Animal protocols used in this study strictly adhered to the ethical and sensitive care and

use of animals in research and were approved by the Washington University School of

Medicine Animal Studies Committee (Animal Welfare Insurance Permit #A-3381-01,

protocol#21-0206). *mNemp1* (Nemp1$^{em\# (TCP) McNeill}$) CRISPR KO allele was obtained by

CRISPR-Cas9–mediated deletion of exon3 that is present in all *mNemp1* transcripts

(Toronto Center for Phenogenomics). Mice were generated and maintained on a

C57Bl6N background [1].

Antibodies

Rat Ter119-Alexa647 (Biolegend, #116218, 1:200), rabbit Nemp1 (1:1,000), mouse Lap2

(BD Biosciences, #611000, 1:200), Rat ckit-Alexa647 (#105817, 1:200), goat LaminB1

(Santa Cruz Biotechnology) were used for immunofluorescence microscopy. Fluotag-X4

anti-Rabbit IgG-Atto488 (Nanotag, #N2404, 1:500) and FluotagX2 anti-mouse IgG-

AberioSTAR-580 (Nanotag, #N1202, 1:500) were used as secondary antibodies. Rat Ter-

119-PE (Biolegend, #116208, 1:200), Rat CD71-APC (Biolegend, #113820, 1:200) were used for FACS and flow imaging. Histone3 (Abcam, #1791, 1:2000), Nemp1 (1:1000) and Ter119-Biotin (Biolegend, #116204, 1:500) were used for immunoblotting.

Bone marrow and spleen cells collection

Two to four month-old mice were euthanized by CO_2 inhalation. Legs were separated at the pelvic-hip joint and femurs and tibia cleaned off from tendons and muscle tissues in cold PBS. Bones were cut on their extremities and transferred into collection units consisting of 0.5 ml tubes with pre-perforated (18 gauge needle) bottoms inserted in 1.5 ml collection tubes. Collections units were centrifuged 4 min at 6000 RPM at 4C. Pellets of BM cells were then resuspended in 1 ml of FACS buffer (0.5% BSA and 2mM EDTA in PBS), transferred to a 40uM cell strainer and washed with 10 ml of FACS buffer. Cell suspensions were then spinned down for 3 min at 3,000 RPM. Pellets were resuspended in 2 ml of FACS buffer and fixed by adding 3.5 ml of fixation/permeabilization buffer (BD Biosciences) and rocking overnight at 4C. Cells were then washed in permeabilization buffer (BD Biosciences) through three cycles of centrifugation for 3 min at 850g and stored at 4C for further use.

Flow imaging

One million fixed BM and spleen cells were immunolabelled for one hour at room temperature with Ter119-PE (1:200, Biolegend) and 1ug/ml Hoechst 33342 (Thermofisher) in permeabilization buffer and then washed in PBS through three cycles of centrifugation for 3 min at 850g. The final pellet was resuspended in 40 µl PBS for

image flow analysis. Data were acquired on an AMNIS ImageStreamX multispectral

imaging flow cytometer (Luminex) using the Inspire software package. All images were

acquired with the 60X objective with Hoechst (405 laser line), Ter119PE (488 laser line)

and brightfield imaged in channels 1, 3 and 4, respectively. Laser intensities were

adjusted to avoid signal saturation. Single fluorophore labeling were used to build a

compensation matrix. Post-acquisition data analyses were performed with the IDEAS

software package. For measurements of Ter119+ erythroid populations, cells in focus

(gradient RMS) were gated on Hoechst+ cells that were subsequently plotted for Ter119

intensity.

Immunofluorescence confocal microscopy

For immunostaining of intracellular epitopes, fixed BM cells were washed and

permeabilized three times in Perm/wash buffer (BD Biosciences) supplemented with

0.1% TritonX100 and further incubated in the same buffer with primary antibodies

overnight at 4C. After three washes, cells were incubated with secondary antibodies and

with fluorescently labeled antibodies against extracellular epitopes for 2 hours at room

temperature and then washed three times with three cycles of centrifugation for 3 min at

850g. Cells pellet were resuspended in 20ul of PBS. 3 μl of cell suspension were mixed

with 10ul of fluorescence mounting medium (Dako) and mounted for downstream

confocal imaging. All images were acquired on a Nikon confocal microscope with a 1.4

NA 100X objective. Images denoising and 3D reconstruction were performed with the

NIS-Element software package suite.

Analysis of mouse peripheral blood

Whole blood was collected by venipuncture of the facial vein and immediately transferred in blood collection tubes (BD Microtainer). Blood samples were mixed and placed under the Hemavet HV950 probe (Drew Scientific, Inc.) for analysis using reagents from the LV-PAK (Drew Scientific Inc.). Multi-Trol mouse serum controls (Drew Scientific, Inc.) were used for calibration of the Hemavet HV950. Collected blood was also spread on glass slides for Wright-Giemsa staining according to the manufacturer's protocol (Wright-Giemsa Stain Modified, Sigma-Aldrich).

In vitro colony-forming assay

Methylcellulose colony-forming assay were performed using Epo-only MethoCult 3334 (Stem Cell Technologies) according to the manufacturer's instructions. BM ($6x10^4$ cells/mL) or spleen ($1x10^5$ cells/mL) cells were mixed with M3334 methylcellulose and plated in triplicates using 35mm Petri-dishes. Cultures were maintained in a humidified incubator at 37°C, 5% CO_2. CFU-E colonies were counted after 2-3 days of culture. BFU-E colonies were counted 5 days after culturing.

Acute Anemia and Splenectomy

Acute hemolytic anemia was induced by intraperitoneal injection of phenylhydrazine (PHZ) (Sigma) with a single dose of 100mg/kg. Splenectomy was performed at the Hope Center for Neurological Disorder at Washington University in St. Louis. Approximately 1 month later, the splenectomized mice were induced by PHZ for hemolytic anemia. Blood samples were collected by venipuncture of the facial vein at different time point, and

hematologic parameters measured on a HemaVet HV950 complete blood count

instrument.

Flow cytometry and cell sorting

BM and spleen cells from WT or Nemp1 KO mice were dissociated, resuspended in

PBS/0.5% BSA and passed through a 40uM cell strainer to obtain single-cell

suspension before antibody staining. Analysis of erythroid maturation using CD71 and

Ter119 was conducted as previously described [22]. Freshly isolated cells were stained

at 4°C in PBS/0.5% BSA with purified anti-mouse CD16/32 to block Fc receptors and

incubated with PE-Ter119 (TER-119) and APC-CD71 (R17217) antibodies for 30

minutes at 4°C. DAPI was used to exclude dead cells from analysis. Where apoptosis

was measured, immunostaining for Ter119 and CD71 was followed by a 15-minute

incubation with FITC-conjugated Annexin-V and propidium iodide following the

manufacturer's protocol (BD Biosciences). Flow cytometry was carried out on BD LSR II

machine and BD FACSAria II was used for cell sorting. Gate strategy was performed as

previously described [22]. The ProE gate contains $CD71^{high}Ter119^{intermediate}$. The

$Ter119^{high}$ cells are further analyzed. Here $CD71^{high}$ cells are subdivided into less

mature, large EryA erythroblasts ($CD71^{high}Ter119^{high}FSC^{high}$) and smaller, more mature

EryB erythroblasts ($CD71^{high}Ter119^{high}FSC^{low}$). The most mature erythroblasts subset is

EryC ($CD71^{low}Ter119^{high}FSC^{low}$).

HSPC and committed progenitors staining were conducted as described previously [23].

Cells from BM were harvested in PBS/0.5% BSA, and quickly lysed with RBC lysis

buffer for 1 min at 4°C. Cells were then stained with PE-Cy7 conjugated anti-Gr-1 (RB6-

8C5), -Mac1 (M1/70), -B220 (RA3-6B2), -Ter119 (TER-119), -CD3 (17A2), in

combination with APC-e780-c-Kit (2B8), PerCP-Cy5.5-Sca1 (D7), APC-CD48 (HM48-1),

PE-CD150 (TC-12F12.2), FITC-CD34 (RAM34) and BV421-CD16/32 (93) antibodies for

30min on ice. Flow cytometry was carried out on BD Symphony A3 machine. Data were

analyzed on FlowJo software (FlowJo, LLC). Different committed progenitors were

defined as granulocyte-monocyte progenitor (GMP, $CD34^+CD16/32^+Lin^-c\text{-}Kit^+Sca1^-$)

cells, common myeloid progenitor (CMP, $CD34^+CD16/32^-Lin^-c\text{-}Kit^+Sca1^-$) cells and

megakaryocyte-erythrocyte progenitor (MEP, $CD34^-CD16/32^-Lin^-c\text{-}Kit^+Sca1^-$).

ACKNOWLEDGMENTS

The authors would like to thank the Centre for Phenogenomics (TCP), Sinai Health

System, Toronto, Ontario, for assistance with blood analysis and the Mouse Genetics

Core at Washington University School of Medicine in St Louis (Dr. Mia Wallace) for mouse

husbandry and genotyping.

REFERENCES

1. Tsatskis Y, Rosenfeld R, Pearson JD, Boswell C, Qu Y, et al. (2020) The NEMP family supports metazoan fertility and nuclear envelope stiffness. Sci Adv 6: eabb4591.

2. Mamada H, Takahashi N, Taira M (2009) Involvement of an inner nuclear membrane protein, Nemp1, in Xenopus neural development through an interaction with the chromatin protein BAF. Dev Biol 327: 497-507.

3. Shibano T, Mamada H, Hakuno F, Takahashi S, Taira M (2015) The Inner Nuclear Membrane Protein Nemp1 Is a New Type of RanGTP-Binding Protein in Eukaryotes. PLoS One 10: e0127271.

4. Tang Y, Dong Q, Wang T, Gong L, Gu Y (2022) PNET2 is a component of the plant nuclear lamina and is required for proper genome organization and activity. Dev Cell 57: 19-31 e16.

5. Menon V, Ghaffari S (2021) Erythroid enucleation: a gateway into a "bloody" world. Exp Hematol 95: 13-22.

6. Migliaccio AR (2010) Erythroblast enucleation. Haematologica 95: 1985-1988.

7. Zhao B, Yang J, Ji P (2016) Chromatin condensation during terminal erythropoiesis. Nucleus 7: 425-429.

8. Barminko J, Reinholt B, Baron MH (2016) Development and differentiation of the erythroid lineage in mammals. Dev Comp Immunol 58: 18-29.

9. Zhao B, Mei Y, Schipma MJ, Roth EW, Bleher R, et al. (2016) Nuclear Condensation during Mouse Erythropoiesis Requires Caspase-3-Mediated Nuclear Opening. Dev Cell 36: 498-510.

10. Hattangadi SM, Martinez-Morilla S, Patterson HC, Shi J, Burke K, et al. (2014) Histones to the cytosol: exportin 7 is essential for normal terminal erythroid nuclear maturation. Blood 124: 1931-1940.

11. Zhao B, Liu H, Mei Y, Liu Y, Han X, et al. (2019) Disruption of erythroid nuclear opening and

histone release in myelodysplastic syndromes. Cancer Med 8: 1169-1174.

12. Zhen R, Moo C, Zhao Z, Chen M, Feng H, et al. (2020) Wdr26 regulates nuclear condensation

in developing erythroblasts. Blood 135: 208-219.

13. Nowak RB, Papoin J, Gokhin DS, Casu C, Rivella S, et al. (2017) Tropomodulin 1 controls

erythroblast enucleation via regulation of F-actin in the enucleosome. Blood 130: 1144-

1155.

14. Choi J, Baldwin TM, Wong M, Bolden JE, Fairfax KA, et al. (2019) Haemopedia RNA-seq: a

database of gene expression during haematopoiesis in mice and humans. Nucleic Acids

Res 47: D780-D785.

15. Gautier EF, Ducamp S, Leduc M, Salnot V, Guillonneau F, et al. (2016) Comprehensive

Proteomic Analysis of Human Erythropoiesis. Cell Rep 16: 1470-1484.

16. Segura-Totten M, Wilson KL (2004) BAF: roles in chromatin, nuclear structure and retrovirus

integration. Trends Cell Biol 14: 261-266.

17. Gesson K, Vidak S, Foisner R (2014) Lamina-associated polypeptide (LAP)2alpha and

nucleoplasmic lamins in adult stem cell regulation and disease. Semin Cell Dev Biol 29:

116-124.

18. Naetar N, Korbei B, Kozlov S, Kerenyi MA, Dorner D, et al. (2008) Loss of nucleoplasmic

LAP2alpha-lamin A complexes causes erythroid and epidermal progenitor

hyperproliferation. Nat Cell Biol 10: 1341-1348.

19. Somech R, Shaklai S, Amariglio N, Rechavi G, Simon AJ (2005) Nuclear envelopathies--

raising the nuclear veil. Pediatr Res 57: 8R-15R.

20. Janin A, Bauer D, Ratti F, Millat G, Mejat A (2017) Nuclear envelopathies: a complex LINC

between nuclear envelope and pathology. Orphanet J Rare Dis 12: 147.

21. Worman HJ, Schirmer EC (2015) Nuclear membrane diversity: underlying tissue-specific

pathologies in disease? Curr Opin Cell Biol 34: 101-112.

22. Koulnis M, Pop R, Porpiglia E, Shearstone JR, Hidalgo D, et al. (2011) Identification and

analysis of mouse erythroid progenitors using the CD71/TER119 flow-cytometric assay. J

Vis Exp.

23. Xu CX, Lee TJ, Sakurai N, Krchma K, Liu F, et al. (2017) ETV2/ER71 regulates hematopoietic

regeneration by promoting hematopoietic stem cell proliferation. J Exp Med 214: 1643-

1653.

540
541

SUPPLEMENTAL FIGURE CAPTIONS

S1 Fig: Nemp1 KO mice have peripheral blood defects: *Hemavet measurement of red blood cells (RBC), red blood cells distribution width (RDW) and hemoglobin (Hg) content of Nemp1KO and WT blood from P4 to P150. N: number of biological replicates for each time point and genotype. Data are shown as mean ± SD. Student's t-test. ns. not significant, *p<0.05. **p<0.01. ***p<0.001. ****p<0.0001.*

S2 Fig: Nemp1 KO mice display increased splenic stress erythropoiesis: *Representative FACS analysis (**A**) and quantification (**B**) of SEPs in dissociated WT and Nemp1 KO spleens. Comparisons of BFU-E (**C**) and CFU-E (**D**) formation capacity between WT and Nemp1KO dissociated spleens. Data are shown as individual subject and the mean ± SD. Student's t-test. ****p<0.0001. Complete blood count (CBC) analysis of WT (n=5) and KO (n=5) mice during the entire recovery period following PHZ treatment (100mg/kg). Data are show as geometric mean ± 95% of difference. Using two-way ANOVA test: *, p=0.0112. (**E**). CBC analysis of WT (n=6) and KO (n=6) mice after splenectomy. Data are show as geometric mean ± 95% of difference. Using* two-way ANOVA test: **, p=0.0012 for RBC comparison; p=0.0022 for Hb comparison; p=0.0045 for MCV comparison; ***, p=0.0007*(**F**). CBC analysis of WT (n=6) and KO (n=6) mice after splenectomy plus PHZ treatment (100mg/kg). Data are show as geometric mean ± 95% of difference. Using two-way ANOVA test: *, p=0.0102; **, p=0.0078 for RBC comparison; p=0.0013 for Hb comparison; ****, p < 0.0001; (**G**).*

S3 Fig: Nemp1 localizes at the NE of hematopoietic cells and aggregates at NE openings. ***A****. Topology of NEMP1 at the NE with the Nemp1 antibody epitope denoted in green. INM, ONM: inner and outer nuclear membrane, respectively. **B.** Immunolocalization of endogenous Nemp1 in cKit+ progenitors by confocal microscopy (top panel). A Ckit-/Ter119low ProE is shown for comparison (Bottom panel). **C.** Intensity profile lines showing Nemp1 peaks (green,*

nuclear envelope) that are distinct from Ter119 peaks (cytoplasmic membrane) in WT BM.

*Nemp1 peaks are not present in Nemp1 KO BM. **D.** Intensity profile lines comparing the*

intensity of Nemp1 in aggregates at the edge of NE openings to the intensity of Nemp1 in intact

NE. Note the lack of accumulation of either Lap2 (red trace) or chromatin (blue trace) in Nemp1

aggregates. Right panel: Maximum intensity projection (Max I) of the same cell showing the

*preferential accumulation of Nemp1aggregates near or at the edge of the NE opening. **E.***

Maximum intensity projection of WT (top) or Nemp1 KO (bottom) erythroblasts undergoing NE

opening. Note the absence of Nemp1 aggregates in Nemp1KO erythroblasts thereby confirming

their specificity.

***S1 Video:** Non-tresholded 3D reconstruction from confocal planes of an erythroblast*

undergoing nuclear envelope opening

***S2, S3 Videos**: 3D reconstruction from confocal planes (thresholded for Nemp1 intensity to*

emphasize Nemp1 aggregates, top panels) of erythroblasts undergoing a large (top) or small

(bottom) nuclear envelope opening. Note the presence of Nemp1 aggregates near or close to

NE openings.

Figure 1

Figure 2

Figure 3

Figure 4

A

Time

DNA

Ter119

Nemp1

DIC

Merge

B

WT

KO

BV-421-DNA

44.1% 44.8% 58.9% 50.5%

40.4% 41.8% 28.8% 35.1%

APC-Ter119

C

% in WT and Nemp1 KO

**

**

Nucleated   Enucleated

WT   KO

Hoechst

75.8%   87.7%

24.1%   11.9%

Ter119

Figure 5