ICAT: A Novel Algorithm to Robustly Identify Cell States

Following Perturbations in Single Cell Transcriptomes

Dakota Y. Hawkins[1,2], Daniel T. Zuch[2,3], James Huth[2], Nahomie Rodriguez-Sastre[2], Kelley R.

McCutcheon[2], Abigail Glick[2,3], Alexandra T. Lion[2,3], Christopher F. Thomas[2], Abigail E.

Descoteaux[2,3,5], W. Evan Johnson[1,6], and Cynthia A. Bradham[1,2,3,5*]

[1]Bioinformatics Program, [2]Biology Department, [3]Program in Molecular and Cellular Biology and

Biochemistry, and [5]Biological Design Center, Boston University, Boston, MA, USA

[6]Division of Computational Biomedicine, Boston University School of Medicine, Boston, MA,

USA

*author for correspondence; e-mail: cbradham@bu.edu

Author e-mails: DYH, dyh0110@bu.edu; DTZ, dtzuch@bu.edu; JH, jhuth@bu.edu; NRS,

chantal1@bu.edu; KRM, kmccutch@bu.edu; AG, aglick18@bu.edu; CFT: cfthomas@bu.edu;

AED: abigaild@bu.edu; WEJ, wej@bu.edu

KEY WORDS: single cell, scRNA-seq, clustering, cell states, perturbation, cell type, semi-

supervised, self-supervised

Abstract

The detection of distinct cellular identities is central to the analysis of single-cell RNA sequencing experiments. However, in perturbation experiments, current methods typically fail to correctly match cell states between conditions or erroneously remove population substructure. Here we present the novel, unsupervised algorithm ICAT that employs self-supervised feature weighting and semi-supervised clustering to accurately resolve cell states across heterogeneous conditions. Using simulated and various real datasets, we show ICAT is superior in identifying and resolving cell states compared to current integration workflows. ICAT is robust to low signal strength, high perturbation severity, and disparate cell type proportions. We empirically validate ICAT in a developmental model and find that only ICAT identifies a perturbation-unique cellular response. Taken together, our results demonstrate that ICAT offers a significant improvement in defining cellular responses to perturbation in single-cell RNA sequencing data.

Introduction

From deconstructing tumor cell compositions [1], to reconstructing developmental pathways [2], single-cell RNA sequencing (scRNA-seq) has revolutionized scientists' ability to explore complex tissues and systems. Clustering of individual cells is central to the analysis of scRNA-seq data, in which cell states are identified by grouping cells with similar gene expression profiles [3]. Whether to ascertain the effect of drug treatment on cancer viability, or to infer the mechanistic role of a given gene in tissue specification, perturbation experiments have long been used to dissect and understand complex biological systems. Oftentimes this analysis is formalized by identifying changes in the abundance of cell identities between treatments [3–5]. Inherently, such analyses are based on the assumption that cell states are readily identified and matched between treatments. However, matching identities across perturbing treatments presents particular issues with scRNA-seq datasets due to dominating treatment effects (Fig. S1), where it is typical that scRNA-seq datasets segregate by treatments rather than cellular identities (Fig. S1) [3,4,6,7]. This challenge presents significant difficulties in correctly evaluating cellular responses to perturbation in scRNA-seq experiments.

Common techniques for identifying cell states in scRNA-seq perturbation experiments rely on either reference datasets to map cells onto known cell identities [8,9], or an integration algorithm to minimize differences between control and perturbed cells by transforming cells into a shared space prior to clustering cells [3,10]. The first scenario is problematic for non-model systems, where such atlases may not exist, and is inappropriate when the goal of a given experiment is discovery of novel cell identities or subtypes; for these reasons, the latter approach is often preferred. Toward that end, scRNA-seq algorithms have been developed that broadly integrate heterogeneous datasets [3,11,12]. Briefly, integration algorithms identify a shared space between separate datasets, then subsequently project cells into this embedding. While promising for combining datasets across modalities, integration methods can smooth over

and eliminate biological signals, leading to incorrect identification and possibly wholesale

erasure of captured cell states [3,13–15].

We therefore developed a two-step algorithm to robustly and sensitively **I**dentify **C**ell

states **A**cross **T**reatments (ICAT) in scRNA-seq data. By employing self-supervised feature

weighting, followed by semi-supervised clustering, ICAT accurately identifies cell states across

scRNA-seq perturbation experiments with high accuracy. ICAT does not require prior

knowledge of marker genes or extant cell states, is robust to perturbation severity, and identifies

cell states with higher accuracy than leading integration workflows within both simulated and

real scRNA-seq perturbation experiments, while operating at lower signal resolutions.

Results

ICAT discriminates cell states across perturbations in single-cell RNA-seq data

To overcome the inability of current clustering algorithms to accurately detect cell

identities in perturbation experiments (Fig. S1), we developed a two-step algorithm (Fig. 1A).

Using the Louvain method [16] by default, the first step separately clusters control cells to

produce initial cluster labels for each cell. Cluster labels are then used to weight genes by their

ability to predict the previously generated labels using Neighborhood Component Feature

Selection (NCFS) [17]. NCFS weights genes by using gradient ascent to maximize a regularized

leave-one-out classification accuracy metric. By the nature of regularization, most gene weights

converge to zero, leaving only the most predictive genes with weights greater than 1. Applying

the learned gene weights to the original expression matrix with both control and treated cells

transforms the matrix into a "cluster-defined" space, where the distances between cells are

dominated by highly weighted marker genes. After this transformation, ICAT retains previously

identified cluster labels for control cells, then performs semi-supervised Louvain community detection [16,18], whereby control labels are held immutable (Fig. 1A).

This strategy achieves two goals: 1. ICAT makes comparing treatment effects on cell states straightforward. By implementing semi-supervised clustering with immutable control labels, ICAT ensures identities discovered in control cells are retained when treated cells are introduced. 2. ICAT produces easily interpretable embeddings. Because NCFS is a sparse method, it overtly separates informative and non-informative genes. Thus, the expression patterns of a few highly weighted genes directly account for

Figure 1. Overview of the ICAT algorithm. A. The schematic illustrates the ICAT$_C$ implementation of ICAT. To identify cell states across treatments, ICAT first performs self-supervised feature weighting to find genes that discriminate cell identities among control cells alone, followed by semi-supervised clustering using the newly transformed expression matrix. To learn feature weights, ICAT clusters control cells using Louvain community detection, then uses the newly generated cluster labels as input into NCFS to weight genes by their predictiveness. After applying the learned gene weights to the original gene expression matrix, ICAT clusters both treated and control cells together using a semi-supervised implementation of Louvain community detection. During this process, ICAT holds the previously identified cluster labels for the control cells immutable. **B.** The schematic illustrates the ICAT$_{C+T}$ implementation, which expands feature weighting to treated cells to identify asymmetrical populations between treatments. Cells are split along treatments and independently clustered using the Louvain method, then cluster labels are used to learn gene weights using NCFS in each treatment set independently. To retain asymmetrically informative genes, weights for each treatment are concatenated row-wise and subsequently reduced to the maximum weight using a row-wise maxpool. The reduced weight vector is then used to transform the original count matrix.

cell location in the NCFS-transformed space. This is a noted benefit over typical integration

methods, where transformations are obscured behind neighbor-based approaches [11,12].

These two features together produce interpretable results that ease compositional analysis

between treatments.

While similar in approach, ICAT provides several advantages compared to identifying

shared cell states via marker gene mapping: first, ICAT does not require previously known

marker genes, making ICAT appropriate for situations with previously uncharacterized cell

states. Second, by clustering control and treated cells together, ICAT takes advantage of the

structure in both datasets compared to clustering both independently. This alleviates issues with

imperfect matches between marker genes. Third, by foregoing common marker gene presence-

absence comparisons to map cell types, ICAT accurately identifies activated states, which is

otherwise difficult. Finally, by generating cell state labels through a clustering step rather than a

classification approach, ICAT is able to group cells with treatment-specific phenotypes and

identify out-of-sample states.

One caveat of our approach is that if asymmetrical populations exist in treated cells, but

not controls, their identifying genes will likely be collapsed by the control-only feature-weighting

step. To avoid this artifact, ICAT can be optionally enabled to perform feature-weighting in all

treatment types to identify asymmetrical populations across conditions (Fig. 1B, Fig. 2A). To

reduce gene weights to a single gene-weight vector, only the maximum gene weight produced

across the conditions is retained. During semi-supervised clustering, labels for control cells are

retained during label initialization, while treated cells are re-initialized as singleton clusters.

ICAT exhibits superior performance across conditions in simulated datasets

To assess ICAT, we initially tested performance in simulated single-cell RNAseq data

(Fig. 2A-E). We simulated data using a zero-inflated Negative Binomial distribution to model

gene counts per Büttner and colleagues [14]. To simulate perturbations, we sampled a scalar

multiplier from a $\Gamma(2,2)$ distribution, then applied this multiplier to randomly selected genes as perturbation targets. To generate distinct cell types, we randomly selected a separate set of genes to act as markers for each population. Selected marker genes had their average expression shifted by a scalar multiplier following a $\Gamma(3,3)$ distribution to

Figure 2. ICAT outperforms current methods for cell state identification and is robust to experimental conditions. A. UMAP projections of raw (left), ICAT$_C$ processed (middle), and ICAT$_{C+T}$ processed count matrices for the simulated data. Projections show ICAT$_C$ and ICAT$_{C+T}$ correctly mix shared populations (red and green), while only ICAT$_{C+T}$ isolates asymmetrical populations (purple, yellow). ICAT performance for simulated data was further evaluated using the ICAT$_{C+T}$ implementation only. **B.** Select marker and perturbed gene expression patterns are displayed as violin plots for three simulated control cell types (1-3) under normal (C) and perturbed (P) conditions. P(C1)+ is a stimulated and perturbed version of cell type 1; perturbation-specific cell types P4 and P5 express distinct marker genes. **C.** The percentage of perturbed genes used to assess robustness to perturbation severity is shown; values range from 1-25%. **D.** The average number of marker genes per cell identity used to assess robustness to signal strength is shown; values range from 10 to 105 (0.7 - 7% of total genes). **E.** The set of cell identity proportions used to test the ability to identify rare cell states is shown. The number of cells per treatment-label pair ranges from 50 to 175. **F.** Method performance is compared as the fraction of perturbed genes increases (left), and as the average number of marker genes per population increases (right). Results are depicted as averages with 95% confidence intervals shown by shading. **G.** Method performance is compared as the proportion of cell types is varied (left). The Gini coefficient reflects the degree of population size disproportion among cell states. **H.** The performance of integration methods alone or in combination with ICAT (as ICAT$_{Seurat}$ or ICAT$_{Scan}$) is compared across the same metrics for the perturbation (H1) and signal strength experiments (H2).

create population substructure. To test ICAT's ability to detect both asymmetrical populations

and activated states, we simulated five populations of cells: three that are present in both

control and treated cells, with one of these exhibiting elevated marker gene expression upon

treatment and referred to as "stimulated" (population C1 and P(C1)+), and two treatment-only

populations (populations P4 and P5) (Fig. 2A-B).

Using simulated data, we first assessed ICAT's robustness against increasing

perturbation severity, testing five conditions with varying proportions of perturbed genes,

ranging from 1% to 25% (Fig. 2C, Table S1). Next, we assessed performance with changing

resolutions between cell types by simulating 10 different experimental conditions with a varying

number of average marker genes per cell type. Mean marker gene numbers ranged from 10-

105 per population (0.67-7.0% of total genes) (Fig. 2D, Table S2). In both cases, each

experiment was simulated 15 times, for a total of 90 independent datasets in the first case (Fig.

2C) and 150 in the second case (Fig. 2D). Finally, to test performance as cell type proportions

deviate from equality, we simulated three different experimental conditions with varying

distributions of each cell type. Experiments ranged from all cell types having 100 cells per

treatment, to the most disparate where the minimum and maximum population size per

treatment were 50 and 150, respectively (Fig. 2E, Table S3). To isolate the effect of differential

abundance, proportion simulations did not include an activated cell type, and instead were

composed of three shared populations and two treatment-unique populations (Fig. 2E). Each

experimental proportion was simulated 15 times.

When assessing ICAT's performance in simulated data, we considered three criteria

reflecting "good" clustering between control and treated cells: 1. Global label conservation: the

extent of mapping between cluster labels and known cell identities, which is ideally one-to-one

to reflect accurate detection of cell identities, was assessed with the Adjusted Rand Index [19]

(ARI). 2. Treatment mixing: the local homogeneity of shared cell types without undo isolation by

treatment status was assessed with the Local Inverse Simpson's Index [20] (LISI); 3. Unique

label separation: the accurate detection of populations-of-interest was measured using

population-specific F1 measures [13]. Such metrics were used to measure the ability to

accurately detect asymmetric populations that exist only in single conditions (F1-Unique), the

ability to discriminate stimulated cell states from non-stimulated analogues (F1-Stim), and the

identification of rare cell types (F1-Rare). ARI, LISI, and all F1 metrics were standardized

between 0 and 1, where 1 indicated stronger performance.

ICAT is robust to perturbation severity and better resolves cell states at lower

resolutions than leading integration workflows

We compared ICAT against two state-of-the-art integration methods: Seurat [11,21] and

Scanorama [12]. Seurat was chosen for its ubiquity in single-cell analysis, while Scanorama was

chosen due to its strong performance in benchmarks comparisons [13]. In both cases, datasets

were integrated between treatments, then the results were clustered via Louvain community

detection. To establish baseline performance, we also tested the "No Integration" scenario in

which cells were clustered using Louvain community detection without accounting for treatment

status. During clustering, we standardized nearest neighbor and resolution parameters for each

method.

When testing perturbation severity or variable marker gene numbers (Fig. 2C-D), ICAT

outperformed both Scanorama and Seurat across all metrics while exhibiting highly stable

performance across both increasing perturbation intensities and at varying cell-state resolutions

(Fig. 2F). ICAT provides near-perfect treatment mixing, with LISI scores nearing 1 for all signal

levels. In contrast, Scanorama and Seurat only approach 0.5 LISI scores at the highest signal

levels, indicating that ICAT exhibits a substantially improved ability to correctly and robustly mix

cell identities across treatments (Fig. 2F).

Scanorama offered comparable performance to ICAT in global label conservation (ARI)

and the ability to detect asymmetrical populations (F1-Unique) (Fig. 2F), while ICAT was

uniformly better at accurately distinguishing the stimulated population from its non-stimulated

analogue (Fig. 2F, F1-Stim). Seurat underperformed both Scanorama and ICAT in label

conservation and asymmetrical population detection (ARI, F1-Unique), and offered comparable

treatment-mixing performance to Scanorama (LISI), indicating that Seurat does not accurately

separate known populations relative to Scanorama or ICAT (Fig. 2F).

For each of the perturbation severity tests and with low marker gene numbers, Seurat

was surprisingly outperformed by the no-integration scenario for ARI (Fig. 2F) which can be

explained by Seurat's general under-clustering of cells (Fig. S2A). Scanorama exhibited a

similar drop in ARI performance at low marker numbers, but consistently outperformed no

integration (Fig. 2F). These results show ICAT more accurately and robustly resolves cell

identities across experimental conditions than either Seurat or Scanorama integration

workflows.

ICAT was unaffected by differences in cell identity proportions as assessed by global

and rare population label conservation via ARI and the F1-Rare metric, respectively, while

Seurat and Scanorama each exhibited decreased performance as the disparity in proportions

increased (Fig. 2G, left). The performance of ICAT was robust and stronger than both

Scanorama and Seurat in each measure.

ICAT improves and stabilizes the performance of integration workflows

Outside of requiring a data matrix, ICAT is agnostic to input data. Accordingly, we also

evaluated clustering performance for two extended methods: $ICAT_{Seurat}$ and $ICAT_{Scan}$, whereby

cells were first integrated using Seurat or Scanorama, respectively, then clustered with ICAT. In

this way, ICAT can be utilized either as an alternative to integration methods, or as an additional

step after integration.

Compared to the typical Seurat workflow, $ICAT_{Seurat}$ led to significant improvements in all

three simulated experiments, leading to greater robustness to perturbation (Fig. 2H1, left),

increased signal sensitivity (Fig. 2H2, left), and better isolation of rare cells (Fig. 2G, middle).

When compared to Scanorama, $ICAT_{Scan}$ better isolate rare cells (Fig. 2G, right), and improved

performance in some metrics in perturbation and signal experiments. Gains were more modest

compared to ICAT$_{Seurat}$, primarily since Scanorama alone already exhibits strong performance

(Fig. 2H1-2, right). In particular, ICAT$_{Scan}$ substantially improved treatment-mixing (LISI) across

conditions compared to Scanorama alone (Fig. 2H1, right). Interestingly, at both signal

extremes, stand-alone ICAT scored higher in the F1-Stim metric than either combined or

original integration methods (Fig. 2H2), indicating that ICAT better preserves discriminatory

signals compared to either integration algorithms.

In summary, the results from simulated data demonstrate that ICAT offers robust and

accurate cell state identification that exceeds both Seurat and Scanorama, two current state-of-

the-art integration algorithms, across a range of simulated conditions including severe

perturbations, low marker gene numbers, and variations in relative cell abundance. Further

analysis shows that identifying cell states with ICAT following integration produces higher quality

clusters compared to either current integration workflow alone.

ICAT produces higher quality clusters in real datasets than integration methods

To assess ICAT's ability to correctly cluster cells in real perturbation experiments, we

made use of three publicly available datasets (Table S4). While no true benchmark datasets

currently exist for defining cell identities in perturbation experiments, we selected datasets that

either contain genotyped cell lines, or are from well-studied cell types with confidently labeled

cells. The first dataset, by Tian et al., was created as a standard to evaluate methods developed

for scRNA-seq data analysis [22]. To evaluate ICAT, we used their CellMix dataset that was

produced by creating various mixtures of nine cells from combinations of H2228, HCC827, and

H1975 adenocarcinoma cell lines, sequencing the cell mixtures, then downsampling the

resulting counts to produce pseudo-cells that are in-line with true single-cell data. While the Tian

dataset was developed to test integration across batch effects, we reasoned that variance

introduced by batch effects is similar to perturbation-induced variation. To better mimic distinct

phenotypes, we selected only four cell "types", consisting of pure H2228, HCC827, and H1975

cells, along with a single mixture with equal proportions of each cell line, denoted "Mixed" (Fig.

S3, Table S4). The second dataset from Kagohara et al. [7] offers a simple perturbation

experiment in which three genotyped cell lines of head and neck squamous carcinoma cells

were treated with either PBS (control) or the chemotherapeutic cetuximab (Table S4). Finally,

we assessed performance in a peripheral blood mononuclear cell dataset collected by Kang et

al.[4] with eight cell types in control and IFN-β-stimulated conditions (Table S4). Prior to analysis,

each dataset was preprocessed by normalizing cell counts and selecting highly variable genes

using *Scanpy*. To ensure each tested algorithm analyzed the same input data, we provided the

same *Scanpy* preprocessed data to each method.

Because NCFS feature-weighting is a computationally intensive task, and since the

Kagohara and Kang datasets each contain > 20,000 cells, it was necessary to subselect cells

for feature weighting. To this end, ICAT selects a representative sample of cells via submodular

optimization using the *apricot* [23] Python package. Submodular optimization via *apricot* was

able to effectively represent the data space using only a small number of cells (Fig. S4).

To measure ICAT's performance in these datasets, we again evaluated clustering quality

using ARI and LISI metrics to measure global label conservation and treatment status mixing,

respectively. Because cell identity is resolution dependent and likely imperfect in real datasets

[3,6], we also calculated the label-free Davies-Bouldin (DB) metric to assess the density of

clusters produced by each method [24]. The DB metric measures similarity between a cluster

and the next most similar cluster. To facilitate comparison, each score was scaled between 0

and 1 such that 1 represents better clustering while 0 represents worse clustering.

In the Tian dataset, ICAT outperformed both existing integration methods in LISI and DB metrics

and was comparable to both in ARI, in which ICAT was slightly worse than Seurat and

Scanorama (Fig. 3A-B, Table S5). ICAT outperformed both integration methods in the Kagohara

and Kang datasets (Fig. 3A-B). Aside from ARI in the Kagohara dataset, in which Scanorama

and ICAT scored equally well (Fig. 3B), ICAT produced the highest scores for ARI, LISI, and DB metrics (Fig. 3B). Overall, ICAT generated higher quality clusters in real datasets compared to existing integration algorithms (Fig. 3A-B, Table S5).

Compared to Scanorama, ICAT_Scan led to greater ARI, LISI and DB scores across the three analyzed datasets (Fig. 3C, Table S5), aside from ARI in the Kagohara dataset where Scanorama and ICAT_Scan scored equally well (Fig. 3C). For Seurat, ICAT_Seurat resulted in improvements that exhibit dataset and metric dependence: cell states were more readily identified in the Kagohara dataset, shown by the dramatic increase in ARI scores, and clusters were tighter and better separated as shown by large increases in DB scores (Fig. 3C). In contrast, ARI scores were maintained or marginally

Figure 3: ICAT outperforms current integration methods in identifying cell states across treatments in real datasets. A. Spider plots compare the performance for each algorithm within each dataset for ARI, LISI, and DB quality metrics. **B.** Lollipop plots highlight the differences in the metrics for each method across the same datasets. **C.** Spider plots comparing Seurat to ICAT_Seurat and Scanorama to ICAT_Scan, respectively. All metrics are scaled from 0 to 1, where 1 is best (closest to the apex of each corner). Abbreviations: DB Davies-Bouldin metric, and otherwise as in Fig. 2. ICAT performance for the Tian dataset was evaluated using the ICAT_{C+T} implementation, while Kagohara and Kang datasets were evaluated using the ICAT_C implementation.

308   reduced with ICAT_Seurat compared to standalone Seurat for the Tian and Kang datasets. The

309   results demonstrate that ICAT offers a marked improvement in cell state identification in multi-

310   condition scRNA-seq experiments compared to the current integration workflows Seurat and

311   Scanorama.

ICAT alone accurately predicts subpopulation response to perturbation *in vivo*

To empirically test ICAT predictions on cell state compositions following perturbation, we

generated scRNA-seq data using the SMART-Seq2 protocol [25]. Specifically, we isolated and

sequenced the skeletogenic lineage known as primary mesenchyme cells (PMCs) at 18 hours

post-fertilization, comparing cells from embryos subjected to chemical treatments that perturb

skeletal patterning, specifically, MK-886 (MK) and chlorate [26–28]. PMCs are of interest

because this population of cells receives patterning cues during development that cause their

diversification into subsets; since MK and chlorate each perturb distinct patterning cues, we

reasoned that these differences would be reflected in PMC subpopulation composition [28–31].

After quality control and cell filtering, we retained expression data for 435 cells (147 control, 134

MK886, and 154 chlorate-treated cells).

ICAT clustered the combined PMC expression profiles into five distinct cell states (Fig.

4A1). The impacts of MK exposure were more dramatic than those of (Fig. 4A2); we therefore

focused on the effects of MK. Two clusters showed stark compositional differences between

control and MK-treated cells, in that cluster 3 lacks MK cells compared to controls, while

clusters 2 and 4 show significant enrichment for MK cells (G-test fdr < 0.05, post hoc Fisher's

Exact test fdr < 0.05; Fig. 4A2, Table S6-7). Since MK-treated cells dominated cluster 4 such

that it appears to be an asymmetric population, we more closely examined clusters 3 and 4.

Investigating highly

predictive genes identified by ICAT found that clusters 3 and 4 were distinguished by inverse

expression of genes *pks2* and *sm50* (Fig. 4A3-4), such that cluster 3 displayed a *sm50+/pks2-*

phenotype, while cluster 4 exhibited a reciprocal *sm50-/pks2+* pattern. To experimentally

validate ICAT's predicted enrichment of the *sm50-/pks2+* cell state and depletion of *sm50+/pks2*

cells, in MK886-treated PMCs, we performed single molecule fluorescence *in situ* hybridization

(FISH) [32] to quantify expression levels for both genes in DMSO (control) and MK-treated embryos [32] (Fig. 4B, Fig. S5). To accurately map FISH signals to PMCs, embryos were stained for PMCs using the PMC-specific 6a9 antibody. Using Ilastik [33], we trained a random forest classifier (out-of-bag error = 0.14) to predict and segment individual PMCs in each confocal image stack (Fig. 4B1-2, Fig. S6). Following methods for relative FISH quantification from the original paper [32], we calculated cell-level expression values for each PMC by first restricting fluorescence signal to PMC-labeled voxels, then calculated

Figure 4: ICAT most accurately defines subpopulation response to perturbation. SMART-seq2 was performed on cells isolated from controls or from embryos treated with perturbants chlorate or MK-886. **A.** ICAT predicts five clusters from the combined data (A1), with treatment-dependent subpopulation compositions (pairwise Fisher's Exact test, fdr < 0.05) (A2). ICAT-calculated gene weights plotted by rank show *sm50* and *pks2* as highly informative for cell-state membership, with 37 total genes being considered informative (weight > 1) (A3). MK-induced differences are defined by reciprocal expression patterns of ICAT-identified genes, *sm50* and *pks2,* in subpopulations 3 (sm50+/pks2-) and 4 (sm50-/pks2+) (A4). **B.** ICAT predictions for control (B1,3,5) and MK-866 (B2,4,6) are validated by HCR FISH analysis. **B1-2.** Automated detection and segmentation of PMCs *in vivo* from 3-D image data is shown as various colors; each color represents an individual cell. **B3-4.** The expression of *sm50* (red) and *pks2* (cyan) transcripts are shown in the same embryos; arrowheads indicate PMCs that express only *sm50* or only *pks2*. **B5-6.** MK-866 treated embryos exhibit statistically significantly fewer *sm50+/pks2-* PMCs (B5) and more *sm50-/pks2+* PMCs (B6) than controls, consistent with the predictions from ICAT (A). Dot plot error bars denote 95% confidence intervals of the mean with each dot representing an individual embryo (n_{DMSO} = 26, n_{MK886} = 37; B5 (Binomial GLM, μ_{DMSO} = 3.59%, μ_{MK886} = 1.91%; β_{MK886} = -0.71, p < 10^{-3}; B6 Binomial GLM, μ_{DMSO} = 0.19%, μ_{MK886} = 2.71%; β_{MK886} = 2.60, p < 10^{-4}). **C.** Integration methods fail to identify the MK886-enriched *sm50+/pks2-* subpopulation (ICAT cluster 4). UMAP projections show clusters generated by Seurat (C1) or Scanorama (C2) for the same data, showing with the cells comprising the MK866-specific asymmetric *sm50+/pks2-* population (ICAT cluster 4, black dots) distributed across the multiple clusters generated by each method. The clusters produced by each integration method are indicated by colored regions.

the average relative intensity for each gene in every PMC (Fig. 4B3-4, Fig. S5). Across

embryos, we labeled individual PMCs as *sm50+* or *sm50-* if their standardized *sm50* expression

values were above the 75th or below the 25th percentiles of all PMCs, respectively (Fig. S7).

The procedure was repeated for *pks2* to determine *pks2+* and *pks2-* PMCs. We found that MK-

treated embryos displayed both a significant loss of the *sm50+/pks2-* phenotype, as well as a

significant enrichment of the *sm50-/pks2+* cell state relative to controls (adjusted p < 0.05; Fig.

4B5-6, Table S8-9): verifying the ability of ICAT to correctly identify affected cell states in

scRNA-seq datasets from perturbed samples.

When the PMC scRNA-seq dataset were integrated using either Seurat or Scanorama,

each method failed to identify the enriched *sm50-/pks2+* cell state in MK-treated embryos, and

instead incorrectly merged those cells with other cell states (Fig. 4C, Fig. S8). In fact, Seurat

produced incorrect conclusions by failing to identify any dependence of cell state composition

on treatment status (G-test fdr > 0.05; Fig. S8A2, Table S10). Scanorama, on the other hand,

correctly identified the depletion of *sm50+/pks2-* PMCs in MK embryos (Fig S8B2-3, Table S11-

12; G-test fdr < 0.05, post hoc Fisher's Exact test fdr < 0.05), but importantly, failed to identify

sm50-/pks2+ enrichment in MK embryos (Fig. S8B2-3, Table S12-13; G-test fdr > 0.05). ICAT is

the only method among these three to correctly identify each cell state that is perturbed by MK

exposure. These results demonstrate that ICAT significantly improves the computational

capability to accurately detect and describe cellular responses to perturbation in scRNA-seq

data.

Discussion

One of the key promises of scRNA-seq technology is to more accurately define complex

biological systems by offering a more finely resolved perspective on transcriptional landscapes

than was previously possible. However, due to technical and biological noise, taking advantage

of such a powerful tool in diverse biological conditions has proven to be challenging. We

therefore developed the ICAT algorithm to improve the accuracy of cell state identification from

scRNA-seq data that includes perturbation treatments. By combining sparse feature weighting

with semi-supervised Louvain community detection, ICAT not only correctly identifies shared

and disparate cell states, but also produces interpretable results that aid in understanding

analytical outcomes and selecting the most suitable downstream analyses.

When comparing the performance of ICAT to state-of-the-art integration algorithms in

simulated datasets, we found that ICAT performs more robustly in response to perturbation

severity, readily identifies cell states at lower signal levels, better isolates asymmetrical and

activated cell states, and more readily identifies rare cell populations when compared to either

Seurat or Scanorama. ICAT also outperforms Seurat and Scanorama in three benchmark

datasets, further solidifying ICAT's ability to better identify cell states across biological

conditions compared to either integration method alone. By testing ICAT on datasets with >

20,000 cells [4,7], we demonstrated ICAT's ability to analyze the large datasets, necessary for

any practical scRNA-seq tool. Depending on the experimental system, researchers may still

elect to use integration algorithms. Excitingly, the choice between ICAT and integration methods

is not mutually exclusive; we show that following Scanorama or Seurat with ICAT produces

higher quality clusters compared to either of these integration workflows alone.

By experimentally validating two uniquely predicted cell states in a developmental model

using single molecule FISH, we empirically demonstrated that the capability of ICAT to

accurately identify perturbed cell states surpasses current state-of-the-art integration workflows.

ICAT was the only method among the three tested that correctly identified these distinct states

and their perturbation responses, whereas both Seurat and Scanorama integration workflows

failed, highlighting ICAT as a substantial advancement in identifying and describing cell state

behaviors across biological conditions and perturbations.

With the growing ubiquity of scRNA-seq experiments and the development of ever more

sophisticated experimental designs, we anticipate that ICAT will play an important role in highly

resolving cell state-specific responses to treatments and differing biological conditions. In the

future, the ICAT framework can easily be extended to operate on other single-cell modalities,

such as scATAC-seq, and will offer a potential route to intelligently cluster multi-modal data from

heterogeneous conditions.

Materials and Methods

Initial clustering

Prior to feature weighting, ICAT identifies initial clusters using Louvain community

detection via the Scanpy Python package [16,34]. The neighbor graph is built using default

parameters of *scanpy.pp.neighbors* with the exception of *n_neighbors* and *resolution*

parameters, which are optimized for each dataset. Once the neighbor graph is constructed,

ICAT performs Louvain community detection using the *scanpy.tl.louvain* function.

Neighborhood Component Feature Selection

Neighborhood Component Feature Selection (NCFS) was implemented in Python

following the methods outlined in the original publication [17]. The implementation takes

advantage of the just-in-time compiler library, NUMBA, to produce highly performant code [35].

NCFS optimizes feature weights to maximize the score function:

$$\xi(\mathbf{w}) = \sum_i \sum_j y_{ij} p_{ij} - \lambda \sum_{l=1}^{d} w_l^2$$

422   where $\mathbf{w}$ is the weight vector; $y_{ij} = 1$ if samples i and j belong to the same class, and is 0

423   otherwise; p_{ij} is the probability of selecting j as a reference to i in a kNN classifier; and $\lambda > 0$ is

424   a regularization parameter. We alter the score function by including a class weight term, c_i, for

425   each sample to account for class imbalance:

$$\xi(\mathbf{w}) = \sum_i c_i \sum_j y_{ij} p_{ij} - \lambda \sum_{l=1}^{d} w_l^2$$

Sample weights are calculated using methods outlined in the scikit-learn Python

package [36,37]. Gradient ascent is then performed using the line method described in the

original publication.

Semi-supervised Louvain community detection

To identify partitions in a graph, traditional Louvain community detection initializes each

node into singleton clusters. Nodes are then merged into other communities if doing so would

optimize a specified quality score [16]. To perform semi-supervsied clustering, ICAT initializes

communities with previously identified control cluster labels. Nodes representing perturbed cells

are placed into their own singleton clusters. During optimization, initialized control labels are set

immutable, while labels for perturbed cells are free to change. By keeping control labels

immutable, we simultaneously allow perturbed cells to merge into control communities — to

identify mixed cell states — while also allowing isolated nodes to form treatment-specific

communities. By default, ICAT optimizes graph partitions using the Reichardt-Burnholdt metric

[38]. Semi-supervised Louvain was implemented by extending the popular louvain-igraph

package (github.com/vtraag/louvain-igraph) to allow for immutable nodes.

Subsampling by submodular optimization

Due to the computational intensity of NCFS feature weighting, for large datasets it is

beneficial to learn feature weights on a subset of the overall cells. To select cells that correctly

capture the dataset variation, ICAT employs submodular optimization via the *Apricot* Python

package [23]. By default, ICAT uses the *facilityLocation* model in *Apricot* to successively select

cells that represent underrepresented regions of the data. The number of cells to select is

chosen by the user; this was set to 1,500 for both Kagohara and Kang datasets.

451   **Simulating scRNA-seq perturbation experiment data**

452   Simulated data was generated using a zero-inflated negative binomial distribution to

453   model gene counts. Our approach follows methods outlined by Büttner and colleagues [14],

454   used to simulate batch effects in scRNA-seq data, adopted to provide for multiple cell

455   populations as well as different cell identity resolutions and perturbation severity.

456   For each dataset, we simulated 1,500 genes. Letting $G = \{g_1, \ldots, g_{1,500}\}$ be the set of

457   all genes, to model M distinct cell identities, we simulate K marker genes such that

458 $G_k = \{g^1, \ldots, g^K\} \subset G$ and $K \sim B(n = 1500, p = p_{marker})$, where p_{marker} is set for each

459   simulation. For simplicity, marker genes were made to be mutually exclusive, such that a gene

460   could not be selected as a marker for multiple populations. Count matrices were constructed as

461   follows:

462   Denote each cell identity y^k, where $Y = \{y^1, \ldots, y^M\}$ is the set of all identity labels.

463   Assuming N cells, let y_i be the cell identity label for cell i such that $y_i \in Y$ and G_{y_i} is the set of

464   marker genes associated with the given identity. For control cells, to generate the count matrix

465 X, we simulate the cell-gene expression values $X_{ij} \sim NB(\mu_{ij}, r = r_j | p_{ij})$, where $r_j \sim U(1, 4)$

466   is the gene-wide dispersion factor and p_{ij} is the cell-gene specific dropout probability. Prior to

467   marker gene application, gene-wide expected values are simulated using a Beta distribution,

468   where $\mu_j \sim Beta(a = 2, b = 5) \cdot c$, where $c = 100$ is a scalar constant. For marker genes, this

469   scalar c is modified to a cell-identity specific value where $c_{y_i} \sim \Gamma(a = 3, b = 3) \cdot 100$. Thus, cell-

470   gene expected values are modeled:

471
$$\mu_{i,j} = \begin{cases} \beta(a = 2, b = 5) \cdot 100, & g_j \notin G_l \\ \beta(a = 2, b = 5) \cdot c_{y_i}, & g_j \in G_l \end{cases}$$

472   Letting $\mu_{y_i} = \{\mu_{y_i,1}, \ldots, \mu_{y_i,M}\}$ be the set expected values for all gene expression

473   values for cells in population y_i, the cell-gene specific dropout probability, p_{ij} is set using the

474   sigmoid function $p_{ij} = sigm\left(-\beta_0 + \beta_{y_ij} \cdot \mu_{ij}\right)$, where $\beta_0 = -1.5$ and $\beta_{y_ij} = median(\mu_{y_i})^{-1}$.

To simulate the count matrix for perturbed cells, X^{pert}, we use the general framework

described above, but apply a scalar shift to gene-wide expected values for perturbed genes.

First, genes to perturb are randomly sampled from G without replacement, such that

$G^{pert} \subset G^{sim}$ is the set of all genes to perturb, and G^{sim} is the set of all allowable genes to

perturb. By default, G^{sim} is the set of all non-marker genes such that $G^{sim} = G \setminus \bigcup\limits_{k=1}^{k=M} G_k$. When

simulating a stimulated population, we specifically included all marker genes for target

population y_{stim} in G^{pert} where the rest of the genes were again sampled from G^{sim}. Assuming

gene $g_j \in G^{pert}$, a perturbation scalar is sampled from a Gamma distribution such that

$s_j \sim \Gamma(a=2, b=2)$. To simulate expression changes due to perturbation, this scalar is

multiplied by cell-gene expected values. Letting $\mu_{i,j}^{pert}$ be the expected value for gene g_j of cell

i in X^{pert},

$$\mu_{i,j}^{pert} = \begin{cases} \mu_{ij}, & g_j \notin G^{pert} \\ \mu_{ij} \cdot s_j, & g_j \in G^{pert} \end{cases}$$

By plugging in μ_{ij}^{pert} into the described count simulation framework, we simulate both

cell-gene dropout probabilities and the final counts. An exception is made for marker genes: in

order to ensure expected population structure exists in both control and perturbed count

matrices, dropout probabilities of marker genes were forced to be the same between the two

conditions.

Simulating experimental conditions

To increase the severity of perturbation and cell identity separation, we varied $|G^{sim}|$ and

p_{marker}, respectively. For perturbation severity experiments, $|G^{sim}|$ was set to 1%, 5%, 10%,

15%, 20%, or 25% of the total number of genes. To increase cell identity signal strength,

p_{marker} was chosen from the range [0.007, 0.07] with a step size of 0.07. Given a binomial

distribution, the average number of marker genes ranged from 10.5 - 105 genes per population.

Assessing performance in real datasets

Raw count matrices of publicly available datasets [4,7,22] were downloaded from GEO.

All datasets were preprocessed using Scanpy, following typical workflows: first, cell counts were

normalized by library size using the *scanpy.pp.normalize_total* function. Normalized counts

were then log-transformed using the *scanpy.pp.log1p* function. Finally, treatment-specific highly

variable genes were identified using the *scanpy.pp.highly_variable_genes* function, with

"flavor=seurat" setting, and with each dataset's treatment variable set as the batch key. The

union of highly variable genes was kept, and genes that were not highly variable in at least one

treatment set were removed.

Tian processing

The Tian CellMix count matrices and metadata files were downloaded from GEO with ID

GSE118767. To isolate more discrete populations, all cells not belonging to pure H2228,

HCC827, and H1975 cell mixtures, or the cell mixture with equal proportions, were filtered out.

Count matrices were preprocessed using *Scanpy* to remove genes expressed in less than three

cells, and the dataset of origin (originally the batch confounder) was set to the treatment

variable.

Kagohara processing

The Kagohara count matrices and metadata files were downloaded from GEO with ID

GSE137524. Genes present in less than 50 cells were removed, along with cells with reads for

less than 50 genes.

Kang processing

The Kang count matrices and metadata files were downloaded from GEO with ID

GSE96583. Cells described as doublets in the cell metadata files were removed from

downstream analysis. Further, genes present in less than 50 cells were removed, along with

cells with reads for less than 50 genes.

Louvain parameter selection

In Louvain community detection, two parameters k, the number of neighbors, and γ, a

resolution parameter, play important roles in identifying clusters. To optimize these parameters

for each dataset (both simulated and real), we filtered each unintegrated dataset to control cells,

and performed a grid search to find parameter pairs that optimized the Calinski-Harabarz score

as implemented in scikit-learn[21,28]. Neighbor values were searched on the interval [5, 50] with a

step size of 5. Likewise, we searched resolution values spanning [0.2, 1.25] with a step size of

0.05.

ICAT feature weighting

For the Kagohara and Kang datasets, feature weights were learned using only control

cells (via the ICAT$_C$ implementation). Meanwhile, when analyzing all simulated datasets along

with the Tian dataset, feature weights were jointly learned and subsequently collapsed across

conditions using the ICAT$_{C+T}$ implementation. Genes were considered "informative" if they had a

learned weight > 1. NCFS parameters used for each dataset are listed in Table S13.

Seurat integration

Seurat integration was performed using version 3.1 as outlined in the vignette

(https://satijalab.org/seurat/articles/integration_introduction.html). Datasets were integrated over

the treatment variable specified for each dataset. Default parameters were used with the

exception of specifying calculation of 50 principal components, and explicitly passing k

neighbors and γ resolution parameters to standardize parameters between methods. To follow

the expected Seurat workflow as best as possible, clusters were identified using the Seurat

function, FindClusters, with the "algorithm=1" to perform Louvain community detection.

Scanorama integration

Scanorama integration was performed using version 1.4, and passing AnnData objects

to the *run_scanorama()* function. Default parameters were used, with the exception of k and γ

parameters being specified for each dataset. After Scanorama integration, clusters were

identified using the *scanpy.tl.louvain* function from Scanpy

Quality metrics

All metrics, with the exception of LISI, were calculated using the scikit-learn Python

package. Metrics that rely on distance measures, Davies-Bouldin (DB) and LISI scores, were

calculated using the euclidean distance over the first 50 principal components.

Davies-Bouldin

For DB, scores closer to zero represent a tighter cluster that is more separated from

other clusters. To facilitate comparison between metrics for each dataset, the negative of DB

scores was scaled between [0, 1] using the MinMaxScaler preprocessor from scikit-learn, where

1 now represents the best performing method.

LISI

LISI metrics were calculated using the "lisi" R package [20]. The LISI score naturally falls

between 0 and the number of batches/treatments in the dataset (representing total separation

and perfect mixing, respectively). Because we expect mixing in only shared cell identities, PCA

was first run on the complete data matrix, then cells from non-overlapping cell identities were

removed, and LISI scores were then calculated using PCs from shared identities. Because LISI

calculates a score for each cell, cell scores were averaged for each dataset-method

comparison. To facilitate comparison to other metrics, LISI scores were scaled between 0 and 1

using the formula:

$$LISI_{scaled} = \frac{LISI - 1}{P - 1}$$

where P is the number of treatment conditions in the dataset.

Population specific F1-scores

To assess each method's ability to detect stimulated and asymmetrical populations, we

created population-specific F1-scores. For each population of interest, we identified the cluster

where a plurality of cells was assigned, and denoted this cluster as c. For each cell within the

population of interest, we created a binary vector, $\mathbf{v}$, denoting whether the cell was assigned to

cluster c.

$$v_i = \begin{cases} 1 & y_i = c \\ 0 & y_i \neq c \end{cases}$$

where y_i is the cluster label for cell i. Assuming perfect isolation where all cells belong to the

same cluster, all entries would be 1. Creating a vector of ones: $\mathbf{u} = \vec{1}$, serves as the basis for

comparison. We passed $\mathbf{v}$ and $\mathbf{u}$ to the *sklearn.metrics.f1_score* function to calculate the

population-specific F1-score [36]. In the case of F1-Unique, where multiple populations of

interest exist, we calculate F1-scores for each population independently and average the scores

between populations. F1-Stim was calculated based solely on the activated population.

Animals and perturbations

Adult *Lytechinus variegatus* sea urchins were obtained from the Duke University Marine

Laboratory (Beaufort, NC) or from Pelagic (Miami, FL). Gamete harvesting and embryo culturing

was performed as described [31,39]. Embryos were treated with chlorate to inhibit the formation

of sulfated proteoglycans as described [28]. MK-886 (Tocris) was reconstituted in DMSO, and

dose–response experiments were performed to obtain the optimal working doses. The optimal

MK-886 concentration was determined to be 1.5-2 µM.

PMC preparation and FACS for scRNA-seq analysis

Embryos were disaggregated at 18 hours post-fertilization (hpf) according to McClay and

Fink [40], with modification as follows: Approximately 3000 embryos at 18 hpf were collected in

artificial seawater (ASW) supplemented with pronase (0.66 mg/mL, Sigma) and immediately

centrifuged for 1 minute at 800 x g, then resuspended in calcium-free seawater (CFSW: 450

mM sodium chloride, 9 mM potassium chloride, 48 mM magnesium sulfate, 6 mM sodium

bicarbonate) containing 20 mg/mL BSA, followed by the addition of 1 volume of hyaline

extraction medium (HEM) [40] and allowed to incubate for 4 minutes at room temperature (RT).

Embryos were then centrifuged for 1 minute at 800 x g, resuspended in CFSW [40], triturated

with decreasing diameter pipettes for 30 seconds total, then centrifuged at 1000 x g for 2

minutes to pellet the dissociated cells. Single cells were then resuspended in CFSW,

centrifuged for 1 minute, then incubated in ASW containing 1 mg/mL BSA with 6a9 anti-PMC

primary antibody (1:5) and anti-mouse Alexa-Fluor 488 labeled secondary antibody (1:500) for 5

minutes. Following washes, cells were suspended in CFSW containing Sytox Blue Dead Cell

Stain (1:1000, ThermoFisher) to label dead cells for exclusion, then sorted using a Sony SH800

Cell Sorter. Cells were gated to exclude doublets and dead cells and collected into 96-well

plates preloaded with RNA lysis buffer at one cell per well, for processing for single cell library

preparation.

scRNA-seq library preparation and sequencing

RNA quantitation and integrity were determined using a Qubit® 2.0 Fluorometer (Life

Technologies) and a 2100 Bioanalyzer (Agilent Technologies). scRNA-seq libraries were

generated using Smart-seq2 on FACS-sorted PMCs as described [25,41]. Libraries were

sequenced on an Illumina NXT platform (BU Microarray and Sequencing Resource Core

Facility) using 75 bp paired end reads. We obtained a median of 1.4M reads per cell over a total

of 416 cells. 144 of these cells were untreated (ASW, artificial sea water), 66 were treated with

DMSO (vehicle), 206 with chlorate, and 192 with MK-886. For all reads, quality control was

performed prior to alignment with fastp [42] using default parameters. Roughly 97% of reads

passed quality control measures with a median of ~1.3M reads per cell. Quality-controlled reads

were aligned to the *L. variegatus* genome [43] using STAR [44], and alignments were quantified

using featureCounts [45]. A median of 148,142 reads per cell were mapped and assigned to

annotated regions within the genome. To ensure only well-sequenced cells and genes were

considered, we removed cells with less than 25,000 mapped reads, then genes with reads

assigned in less than five cells, using Scanpy [34]. Once the final count matrix was generated,

we normalized reads between cells and performed batch-correction using the SCRAN package

[46,47]. Assuming some non-PMCs escaped FACS, we evaluated the identity of both the sorted

and remainder cells from the FACS analysis for the expression of known marker genes for each

major cell type [48]. We then performed a final filtering step that removed sorted cells that lack

expression of canonical PMC marker genes. After filtering, 119 ASW cells, 26 DMSO-treated

cells, 154 chlorate-treated cells, and 134 MK-treated cells remained. For analysis, ASW and

DMSO vehicle controls were combined into a single control group since the number of DMSO

cells were too small to consider them as a representative sample for these comparisons and

since comparisons of DMSO and ASW cells did not reveal significant expression differences.

Prior to cluster analysis, genes were subsetted to highly variable genes identified in control cells

using the *highly_variable_genes* function in *Scanpy*. For clustering, neighbor parameter k=15

and resolution parameter γ = 1 were used for both ICAT clustering, and subsequent Louvain

Community Detection in Seurat and Scanorama workflows.

Assessing treatment effect on cell state composition

To assess whether cell state membership was dependent upon treatment, we performed

G-tests on each cell state using the scipy.stats.*power_divergence* function in the *scipy* Python

package [49,50]. P-values were then adjusted using the *statsmodels.multitest.multipletests*

function from *statsmodel* in Python, with *method="fdr_bh"* [51]. Associations were considered

significant with α = 0.05. Cell states with compositions dependent upon treatment status were

further evaluated with post-hoc pairwise Fisher's Exact test to determine which treatments

affected cell state membership compared to control. Tests were performed using

scipy.stats.*fisher_exact* function from scipy, with p-values again adjusted using the *multipletests*

function with the same parameters.

Single Molecule FISH: probe sets, amplifiers and buffers

Embryos were collected and fixed in 4 % paraformaldehyde at 18 hpf. Hybridization

chain reaction (HCR)-FISH was performed as described [32] using probes, hairpins and buffers

obtained from Molecular Instruments (Los Angeles, CA). Embryos were incubated with hairpins

for 2.5 hours, then signals were captured as 3-D z-stacks using confocal microscopy.

Immunostaining and confocal microscopy

Immunolabeling was performed as described [31,52,53]. PMCs were detected using

primary antibody 6a9 (1:50; from Prof. Charles Ettensohn, Carnegie Mellon University,

Pittsburgh, PA, USA) and secondary goat-anti mouse antibody labeled with Alexa 546 (1:500;

ThermoFisher). Hoechst 33342 (1:1000; Thermo Fisher) was included as a nuclear

counterstain. Confocal microscopy was performed using a Nikon-C2 confocal & Ti-E Spectral

imaging system. Confocal z-stacks were projected using Napari [54].

PMC segmentation

To isolate individual PMCs from the 3-D image stack, a random forest PMC classifier

was trained using the "pixel classification" workflow from Ilastik [33]. The model was trained

using DMSO and MK-treated embryos (n=70) to produce "PMC" and "Background" probabilities.

Prior to training, images were Z-cropped to include only z-slices with PMC signal. Each slice

was then contrast and intensity adjusted using the functions *exposure.adapt_hist* and

exposure.rescale_intensity functions from *scikit-image (skimage)* [55] library in Python. PMC

probabilities were exported from Ilastik for post-processing and PMC segmentation by a custom

Python script.

Briefly, exported Ilastik probabilities were first segmented using

filters.apply_hysteresis_threshold from *skimage* with parameters "low"=0.45 and "high"=0.5. The

resulting binary mask was then refined via morphological opening using

morphology.binary_opening, again from *skimage*. The opened mask was subsequently used as

seed points for watershed segmentation (*skimage.segmentation.watershed*) using the image

gradients from the original data (*skimage.filters.sobel, axis=0*). To restrict PMC labels to typical

PMC sizes and to better separate dense PMC areas, individual regions with areas exceeding

600 voxels were further segmented using a stricter threshold workflow: first, PMC probabilities

were smoothed using a gaussian filter (*skimage.filters.gaussian*). Second, starting with

threshold t = 0.8, a binary mask of the smoothed probabilities was generated by finding all

voxels > t and performing a morphological opening. Isolated objects were then detected using

measure.label from *skimage.* If the number of identified objects did not exceed one, the process

was repeated after incrementing t by 0.025 until the original region was split. The split labels

were then filled back into the original larger region using watershed. To split cell segmentations

with unrealistic 2-D diameters, this process was repeated for any labeled region having a

diameter greater than 15 voxels as well as an area greater than 500 voxels. To correctly

segment vertically stacked PMCs, any label that spanned more than seven z-slices was split

after calculating the label diameter at each slice. Given that PMCs are spherical, it was

reasoned local minima in diameter would indicate the end of a cell, and thus these served as

break points between labels. Finally, to remove unrealistically small labels, the final

segmentation was produced by removing any label that occupied less than 55 voxels.

HCR FISH quantification and analysis

Prior to quantification, images were filtered to remove embryos that displayed either poor

stain contrast or failed to be entirely captured by the z stack. After quality control, 26 DMSO and

37 MK treated embryos were retained from the three fertilizations performed. Individual PMCs

were then labeled using the previously described methodology. After label generation, *pks2* and

sm50 HCR FISH image channels were preprocessed using FISH-quant v2 [56] to scale and

normalize intensities and to perform background subtraction and autofluorescence removal.

Gene expression was quantified for each predicted PMC by calculating the average signal

intensity as described [32]. Prior to analysis, PMC-specific intensities were standardized to a [0,

1] scale to allow for inter-embryo comparisons. Cells were considered to have decreased (-) or

increased (+) expression for a given gene if its standardized expression fell below the 25th or

above the 75th percentile, respectively. We then calculated the proportion of *sm50+/pks2-* and

sm50-/pks2+ PMCs for each embryo. Statistical analysis was performed using a general linear

model to perform binomial logistic regression to assess the effect of treatment status on

phenotype proportion while also accounting for the fertilization batch. The model was fit using

the *statsmodels* [51] Python package using the formula *phenotypic + not_phenotypic ~*

treatment + fertilization, where *phenotypic* is the number of cells displaying the phenotype of

interest, *not_phenotypic* is the number of cells not displaying the phenotype of interest,

treatment is treatment status, and *fertilization* is the fertilization batch for the given embryo.

Availability of data and materials.

ICAT, *ncfs*, and *sslouvain* Python packages are freely available on pip and github

(github.com/BradhamLab). Scripts to perform simulated data generation and benchmark

comparisons are available as a SnakeMake workflow at

(github.com/BradhamLab/ICAT_manuscript). Archived data files for both simulated and real

data are also available at the same repository. The pipeline to align and quantify SMART-seq2

reads is freely available at (github.com/BradhamLab/scPipe), and the raw sequencing data is

available at GEO accession GSE164240. Likewise, all scripts to segment PMCs and quantify

HCR FISH signal are available at (github.com/BradhamLab/hcr_quantify) and the image data is

available upon request.

Author Contributions

719 The study was conceived by CAB and DYH. The computational work was performed by DYH,

720 with advice from WEJ. JH and DTZ collected the urchin scRNA-seq data. NRS, CFT, KRM,

721 ATL, and AED performed the imaging experiments. DYH, KRM, and AG trained the PMC

722   prediction model. The manuscript was written by DYH, CAB, DTZ and NRS, and edited by all

723   coauthors.

724   Competing interests

725   The authors declare no competing interests.

Acknowledgements

We thank Prof. Masanao Yajima (Dept. of Mathematics and Statistics, Boston University) for

helpful discussions. We thank Prof. Charles Ettensohn (Carnegie Mellon University, PA, USA)

for the anti-PMC antibody. We thank Jose Ordovas-Montanes and Sanjay Prakadan of the

Shalek lab (MIT) for help with scRNA-seq library preparation. We thank Dr. Todd Blute for his

advice and help with FACS and microscopy imaging.

Funding

This work was supported by NSF IOS 1656752 (CAB).

References

1. Schelker M, Feau S, Du J, Ranu N, Klipp E, MacBeath G, et al. Estimation of immune cell content in tumour tissue using single-cell RNA-seq data. Nat Commun. 2017 Dec 11;8(1):2032.
2. Wagner DE, Weinreb C, Collins ZM, Briggs JA, Megason SG, Klein AM. Single-cell mapping of gene expression landscapes and lineage in the zebrafish embryo. Science. 2018 Jun 1;360(6392):981–7.
3. Luecken MD, Theis FJ. Current best practices in single-cell RNA-seq analysis: a tutorial. Mol Syst Biol. 2019 Jun 1;15(6):e8746.
4. Kang HM, Subramaniam M, Targ S, Nguyen M, Maliskova L, McCarthy E, et al. Multiplexed droplet single-cell RNA-sequencing using natural genetic variation. Nat Biotechnol. 2017 Dec 11;36(1):89–94.
5. Haber AL, Biton M, Rogel N, Herbst RH, Shekhar K, Smillie C, et al. A single-cell survey of the small intestinal epithelium. Nature. 2017 Nov;551(7680):333–9.
6. Lähnemann D, Köster J, Szczurek E, McCarthy DJ, Hicks SC, Robinson MD, et al. Eleven grand challenges in single-cell data science. Genome Biol. 2020 Feb 7;21(1):31.
7. Kagohara LT, Zamuner F, Davis-Marcisak EF, Sharma G, Considine M, Allen J, et al. Integrated single-cell and bulk gene expression and ATAC-seq reveals heterogeneity and early changes in pathways associated with resistance to cetuximab in HNSCC-sensitive cell lines. Br J Cancer. 2020 Jul;123(1):101–13.
8. Regev A, Teichmann SA, Lander ES, Amit I, Benoist C, Birney E, et al. The Human Cell Atlas. eLife. 2017 Dec 5;6.
9. Schaum N, Karkanias J, Neff NF, May AP, Quake SR, Wyss-Coray T, et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. Nature. 2018 Oct;562(7727):367–72.
10. Perillo M, Oulhen N, Foster S, Spurrell M, Calestani C, Wessel G. Regulation of dynamic pigment cell states at single-cell resolution. Newmark PA, Bronner ME, Rast JP, editors.

763   eLife. 2020 Aug 19;9:e60388.

764 11. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
765   Comprehensive Integration of Single-Cell Data. Cell. 2019 Jun 13;177(7):1888-1902.e21.

766 12. Hie B, Bryson B, Berger B. Efficient integration of heterogeneous single-cell transcriptomes
767   using Scanorama. Nat Biotechnol. 2019 Jun 6;37(6):685–91.

768 13. Luecken MD, Büttner M, Chaichoompu K, Danese A, Interlandi M, Mueller MF, et al.
769   Benchmarking atlas-level data integration in single-cell genomics. Nat Methods. 2022
770   Jan;19(1):41–50.

771 14. Büttner M, Miao Z, Wolf FA, Teichmann SA, Theis FJ. A test metric for assessing single-
772   cell RNA-seq batch correction. Nat Methods. 2019 Jan 20;16(1):43–9.

773 15. Tyler SR, Bunyavanich S, Schadt EE. PMD Uncovers Widespread Cell-State Erasure by
774   scRNAseq Batch Correction Methods [Internet]. 2021 Nov [cited 2021 Nov 29] p.
775   2021.11.15.468733. Available from:
776   https://www.biorxiv.org/content/10.1101/2021.11.15.468733v1

777 16. Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. Fast unfolding of communities in large
778   networks. J Stat Mech Theory Exp. 2008 Oct 9;2008(10):P10008.

779 17. Yang W, Wang K, Zuo W. Neighborhood Component Feature Selection for High-
780   Dimensional Data. J Comput [Internet]. 2012 Jan 1 [cited 2018 Nov 12];7(1). Available
781   from: http://ojs.academypublisher.com/index.php/jcp/article/view/5076

782 18. Traag V. vtraag/louvain-igraph [Internet]. 2021 [cited 2021 Mar 15]. Available from:
783   https://github.com/vtraag/louvain-igraph

784 19. Hubert L, Arabie P. Comparing partitions. J Classif. 1985 Dec 1;2(1):193–218.

785 20. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and
786   accurate integration of single-cell data with Harmony. Nat Methods. 2019
787   Dec;16(12):1289–96.

788 21. Satija R, Farrel JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell
789   gene expression data. Nat Biotechnol. 2015;33(5).

790 22. Tian L, Dong X, Freytag S, Lê Cao KA, Su S, JalalAbadi A, et al. Benchmarking single cell
791   RNA-sequencing analysis pipelines using mixture control experiments. Nat Methods. 2019
792   Jun 27;16(6):479–87.

793 23. Schreiber J, Bilmes J, Noble WS. apricot: Submodular selection for data summarization in
794   Python. J Mach Learn Res. 2020;21(161):1–6.

795 24. Davies DL, Bouldin DW. A Cluster Separation Measure. IEEE Trans Pattern Anal Mach
796   Intell. 1979 Apr;PAMI-1(2):224–7.

797 25. Picelli S, Faridani OR, Björklund ÅK, Winberg G, Sagasser S, Sandberg R. Full-length
798   RNA-seq from single cells using Smart-seq2. Nat Protoc. 2014 Jan 2;9(1):171–81.

799 26. Pidgeon GP, Lysaght J, Krishnamoorthy S, Reynolds JV, O'Byrne K, Nie D, et al.
800   Lipoxygenase metabolism: roles in tumor progression and survival. Cancer Metastasis
801   Rev. 2007 Oct 18;26(3):503.

802 27. Simionato GB, da Silva ACR, Oliva AH de, de Abreu Furquim EM, Couto MCR, Shinohara
803   AL, et al. Lack of 5-lipoxygenase in intramembranous and endochondral 129Sv mice
804   skeleton and intramembranous healing. Arch Oral Biol. 2021 Nov 1;131:105266.

805 28. Piacentino ML, Zuch DT, Fishman J, Rose S, Speranza EE, Li C, et al. RNA-Seq identifies
806   SPGs as a ventral skeletal patterning cue in sea urchins. Dev Camb Engl. 2016 Feb
807   15;143(4):703–14.

808 29. Lyons DC, Martik ML, Saunders LR, McClay DR. Specification to biomineralization:
809   following a single cell type as it constructs a skeleton. Integr Comp Biol. 2014
810   Oct;54(4):723–33.

811 30. Sun Z, Ettensohn CA. Signal-dependent regulation of the sea urchin skeletogenic gene
812   regulatory network. Gene Expr Patterns. 2014;16:93–103.

813 31. Piacentino ML, Ramachandran J, Bradham CA. Late Alk4/5/7 signaling is required for

anterior skeletal patterning in sea urchin embryos. Development. 2015 Mar 1;142(5):943–
52.

32.   Choi HMT, Schwarzkopf M, Fornace ME, Acharya A, Artavanis G, Stegmaier J, et al.
Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive,
versatile, robust. Development. 2018 Jun 26;145(12):dev165753.

33.   Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, et al. ilastik: interactive
machine learning for (bio)image analysis. Nat Methods 2019. 2019 Sep 30;1–7.

34.   Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data
analysis. Genome Biol. 2018 Dec 6;19(1):15.

35.   Lam SK, Pitrou A, Seibert S. Numba: a LLVM-based Python JIT compiler. In: Proceedings
of the Second Workshop on the LLVM Compiler Infrastructure in HPC [Internet]. New York,
NY, USA: Association for Computing Machinery; 2015 [cited 2021 Mar 3]. p. 1–6. (LLVM
'15). Available from: https://doi.org/10.1145/2833157.2833162

36.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn:
Machine Learning in Python. J Mach Learn Res. 2011;12(85):2825–30.

37.   King G, Zeng L. Logistic Regression in Rare Events Data. Polit Anal. 2001;9:137–63.

38.   Reichardt J, Bornholdt S. Statistical mechanics of community detection. Phys Rev E. 2006
Jul 18;74(1):016110.

39.   Bradham CA, McClay DR. p38 MAPK is essential for secondary axis specification and
patterning in sea urchin embryos. Development. 2006 Jan 1;133(1):21–32.

40.   McClay DR, Fink RD. Sea urchin hyalin: Appearance and function in development. Dev
Biol. 1982 Aug 1;92(2):285–93.

41.   Trombetta JJ, Gennert D, Lu D, Satija R, Shalek AK, Regev A. Preparation of Single-Cell
RNA-Seq Libraries for Next Generation Sequencing. Curr Protoc Mol Biol. 2014 Jul
1;107:4.22.1-17.

42.   Chen S, Zhou Y, Chen Y, Gu J. Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018;34(17):i884–90.

43.   Li Y, Omori A, Flores RL, Satterfield S, Nguyen C, Ota T, et al. Genomic insights of body
plan transitions from bilateral to pentameral symmetry in Echinoderms. Commun Biol.
2020 Jul 10;3(1):1–10.

44.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

45.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014 Apr 1;30(7):923–30.

46.   L. Lun AT, Bach K, Marioni JC. Pooling across cells to normalize single-cell RNA
sequencing data with many zero counts. Genome Biol. 2016 Dec 27;17(1):75.

47.   Haghverdi L, Lun ATL, Morgan MD, Marioni JC. Batch effects in single-cell RNA-
sequencing data are corrected by matching mutual nearest neighbors. Nat Biotechnol.
2018 Apr 2;36(5):421–7.

48.   Hogan JD, Keenan JL, Luo L, Ibn-Salem J, Lamba A, Schatzberg D, et al. The
developmental transcriptome for Lytechinus variegatus exhibits temporally punctuated
gene expression changes. Dev Biol. 2020 Apr 15;460(2):139–54.

49.   Sokal R, Rohlf F. Biometry : the principles and practice of statistics in biological research /
Robert R. Sokal and F. James Rohlf. SERBIULA Sist Libr 20. 2013 Apr;

50.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy
1.0: Fundamental Algorithms for Scientific Computing in Python. Nat Methods.
2020;17:261–72.

51.   Seabold S, Perktold J. statsmodels: Econometric and statistical modeling with python. In:
9th Python in Science Conference. 2010.

52.   Bradham CA, McClay DR. p38 MAPK is essential for secondary axis specification and
patterning in sea urchin embryos. Dev Camb Engl. 2006 Jan;133(1):21–32.

53. Gross JM, Peterson RE, Wu SY, McClay DR. LvTbx2/3: a T-box family transcription factor
involved in formation of the oral/aboral axis of the sea urchin embryo. Dev Camb Engl.
2003 May;130(9):1989–99.
54. Nicholas Sofroniew, Talley Lambert, Kira Evans, Juan Nunez-Iglesias, Kevin Yamauchi,
Ahmet Can Solak, et al. napari/napari: 0.3.7rc3 [Internet]. Zenodo; 2020 [cited 2020 Sep
3]. Available from: https://zenodo.org/record/4012782#.X1EdIPEpCHs
55. van der Walt S, Schönberger JL, Nunez-Iglesias J, Boulogne F, Warner JD, Yager N, et al.
scikit-image: image processing in Python. PeerJ. 2014 Jun;2:e453.
56. Imbert A, Ouyang W, Safieddine A, Coleno E, Zimmer C, Bertrand E, et al. FISH-quant v2:
a scalable and modular analysis tool for smFISH image analysis [Internet]. 2021 Jul [cited
2021 Oct 25] p. 2021.07.20.453024. Available from:
https://www.biorxiv.org/content/10.1101/2021.07.20.453024v1
877

Figure Legends

Figure 1. Overview of the ICAT algorithm. A. The schematic illustrates the $ICAT_C$ implementation of ICAT. To identify cell states across treatments, ICAT first performs self-supervised feature weighting to find genes that discriminate cell identities among control cells alone, followed by semi-supervised clustering using the newly transformed expression matrix. To learn feature weights, ICAT clusters control cells using Louvain community detection, then uses the newly generated cluster labels as input into NCFS to weight genes by their predictiveness. After applying the learned gene weights to the original gene expression matrix, ICAT clusters both treated and control cells together using a semi-supervised implementation of Louvain community detection. During this process, ICAT holds the previously identified cluster labels for the control cells immutable. **B.** The schematic illustrates the $ICAT_{C+T}$ implementation, which expands feature weighting to treated cells to identify asymmetrical populations between treatments. Cells are split along treatments and independently clustered using the Louvain method, then cluster labels are used to learn gene weights using NCFS in each treatment set independently. To retain asymmetrically informative genes, weights for each treatment are concatenated row-wise and subsequently reduced to the maximum weight using a row-wise maxpool. The reduced weight vector is then used to transform the original count matrix.

Figure 2. ICAT outperforms current methods for cell state identification and is robust to experimental conditions. A. UMAP projections of raw (left), $ICAT_C$ processed (middle), and $ICAT_{C+T}$ processed count matrices for the simulated data. Projections show $ICAT_C$ and $ICAT_{C+T}$ correctly mix shared populations (red and green), while only $ICAT_{C+T}$ isolates asymmetrical populations (purple, yellow). ICAT performance for simulated data was further evaluated using the $ICAT_{C+T}$ implementation only. **B.** Select marker and perturbed gene expression patterns are displayed as violin plots for three simulated control cell types (1-3) under normal (C) and perturbed (P) conditions. P(C1)+ is a stimulated and perturbed version of cell type 1; perturbation-specific cell types P4 and P5 express distinct marker genes. **C.** The percentage of perturbed genes used to assess robustness to perturbation severity is shown; values range from 1-25%. **D.** The average number of marker genes per cell identity used to assess robustness to signal strength is shown; values range from 10 to 105 (0.7 - 7% of total genes). **E.** The set of cell identity proportions used to test the ability to identify rare cell states is shown. The number of cells per treatment-label pair ranges from 50 to 175. **F.** Method performance is compared as the fraction of perturbed genes increases (left), and as the average number of

marker genes per population increases (right). Results are depicted as averages with 95%

confidence intervals shown by shading. **G.** Method performance is compared as the proportion

of cell types is varied (left). The Gini coefficient reflects the degree of population size

disproportion among cell states. **H.** The performance of integration methods alone or in

combination with ICAT (as ICAT$_{Seurat}$ or ICAT$_{Scan}$) is compared across the same metrics for the

perturbation (H1) and signal strength experiments (H2).

Figure 3: ICAT outperforms current integration methods in identifying cell states across

treatments in real datasets. A. Spider plots compare the performance for each algorithm

within each dataset for ARI, LISI, and DB quality metrics. **B.** Lollipop plots highlight the

differences in the metrics for each method across the same datasets. **C.** Spider plots comparing

Seurat to ICAT$_{Seurat}$ and Scanorama to ICAT$_{Scan}$, respectively. All metrics are scaled from 0 to 1,

where 1 is best (closest to the apex of each corner). Abbreviations: DB Davies-Bouldin metric,

and otherwise as in Fig. 2. ICAT performance for the Tian dataset was evaluated using the

ICAT$_{C+T}$ implementation, while Kagohara and Kang datasets were evaluated using the ICAT$_C$

implementation.

Figure 4: ICAT most accurately defines subpopulation response to perturbation. SMART-

seq2 was performed on cells isolated from controls or from embryos treated with perturbants

chlorate or MK-886. **A.** ICAT predicts five clusters from the combined data (A1), with treatment-

dependent subpopulation compositions (pairwise Fisher's Exact test, fdr < 0.05) (A2). ICAT-

calculated gene weights plotted by rank show *sm50* and *pks2* as highly informative for cell-state

membership, with 37 total genes being considered informative (weight > 1) (A3). MK-induced

differences are defined by reciprocal expression patterns of ICAT-identified genes, *sm50* and

pks2, in subpopulations 3 (sm50+/pks2-) and 4 (sm50-/pks2+) (A4). **B.** ICAT predictions for

control (B1,3,5) and MK-866 (B2,4,6) are validated by HCR FISH analysis. **B1-2.** Automated

detection and segmentation of PMCs *in vivo* from 3-D image data is shown as various colors;

each color represents an individual cell. **B3-4.** The expression of *sm50* (red) and *pks2* (cyan)

transcripts are shown in the same embryos; arrowheads indicate PMCs that express only *sm50*

or only *pks2*. **B5-6.** MK-866 treated embryos exhibit statistically significantly fewer *sm50+/pks2-*

PMCs (B5) and more *sm50-/pks2+* PMCs (B6) than controls, consistent with the predictions

from ICAT (A). Dot plot error bars denote 95% confidence intervals of the mean with each dot

representing an individual embryo (n_{DMSO} = 26, n_{MK886} = 37; B5 (Binomial GLM, μ_{DMSO}= 3.59%,

μ_{MK886}= 1.91%; β_{MK886}= -0.71, p < 10^{-3}; B6 Binomial GLM, μ_{DMSO}= 0.19%, μ_{MK886}= 2.71%; β_{MK886}=

2.60, p < 10^{-4}). **C.** Integration methods fail to identify the MK886-enriched *sm50+/pks2-*

subpopulation (ICAT cluster 4). UMAP projections show clusters generated by Seurat (C1) or

Scanorama (C2) for the same data, showing with the cells comprising the MK866-specific

asymmetric *sm50+/pks2-* population (ICAT cluster 4, black dots) distributed across the multiple

clusters generated by each method. The clusters produced by each integration method are

indicated by colored regions.