An amino acid transporter AAT1 plays a pivotal role in chloroquine resistance evolution in malaria parasites

Alfred Amambua-Ngwa[1†], Katrina A. Button-Simons[2†], Xue Li[3†], Sudhir Kumar[4†], Katelyn Vendrely Brenneman[2†], Marco Ferrari[3], Lisa A. Checkley[2], Meseret T. Haile[4], Douglas A. Shoue[2], Marina McDew-White[3], Sarah M. Tindall[5], Ann Reyes[3], Elizabeth Delgado[3], Haley Dalhoff [2], James K. Larbalestier[2], Roberto Amato[6], Richard D. Pearson[6], Alexander B. Taylor[7], François H. Nosten[8], Umberto D'Alessandro[1], Dominic Kwiatkowski [6], Ian H. Cheeseman[9], Stefan H. I. Kappe[4,10,11], Simon V. Avery[5], David J. Conway[12], Ashley M. Vaughan[4,10]*, Michael T. Ferdig[2]*, Timothy J. C. Anderson[3]*

[1]MRC Unit The Gambia at London School of Hygiene and Tropical Medicine, Banjul, The Gambia

[2]Eck Institute for Global Health, Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA

[3]Disease Intervention and Prevention Program, Texas Biomedical Research Institute, San Antonio, TX, USA

[4]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA, USA

[5]School of Life Sciences, University of Nottingham, Nottingham, UK

[6]Wellcome Sanger Institute, Hinxton, UK

[7]Department of Biochemistry & Structural Biology, University of Texas Health Science Center at San Antonio, TX, USA

[8]Shoklo Malaria Research Unit, Mahidol-Oxford Tropical Medicine Research Unit, Mahidol University, Mae Sot, Thailand; Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research building, University of Oxford Old Road campus, Oxford, UK

[9]Host Pathogen Interactions Program, Texas Biomedical Research Institute, San Antonio, TX, USA

[10]Department of Pediatrics, University of Washington, Seattle, WA, USA

[11]Department of Global Health, University of Washington, Seattle, WA, USA

[12]Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK

[†]These authors contributed equally to this work

*Corresponding authors

Malaria parasites break down host hemoglobin into peptides and amino acids in the digestive vacuole for export to the parasite cytoplasm for growth: interrupting this process is central to the mode of action of several antimalarial drugs. Mutations in the chloroquine (CQ) resistance transporter, *pfcrt*, located in the digestive vacuole membrane, confer CQ resistance in *Plasmodium falciparum*, but typically affect parasite fitness. However, the role of other parasite loci in the evolution of CQ resistance is unclear. Here we use a combination of population genomics, genetic crosses and gene editing to demonstrate that a second vacuolar transporter plays a key role in both resistance and compensatory evolution. Longitudinal genomic analyses of the Gambian parasites revealed temporal signatures of selection on an amino acid transporter (*pfaat1*) S258L variant which

increased from 0-87% in frequency between 1984 and 2014 in parallel with *pfcrt1* K76T.
Parasite genetic crosses then identified a chromosome 6 quantitative trait locus containing
***pfaat1* that is selected by CQ treatment. Gene editing demonstrated that *pfaat1* S258L**
potentiates CQ-resistance but at a cost of reduced fitness, while *pfaat1* F313S, a common
Southeast Asian polymorphism, reduces CQ-resistance while restoring fitness. Our
analyses reveal hidden complexity in CQ-resistance evolution, suggesting that *pfaat1* may
underlie regional differences in the dynamics of resistance evolution, and modulate parasite
resistance or fitness by manipulating the balance between both amino acid and drug
transport.

Drug resistance in microbial pathogens complicates control efforts. Therefore, understanding the genetic architecture and the complexity of resistance evolution is critical for resistance monitoring and the development of improved treatment strategies. In the case of malaria parasites, deployment of five classes of antimalarial drugs over the last half century have resulted in well-characterized hard and soft selective sweeps associated with drug resistance, with both worldwide dissemination and local origins of resistance driving drug resistance alleles across the range of *P. falciparum* [1-3]. Chloroquine (CQ) monotherapy played a central role in an ambitious plan to eradicate malaria in the last century. Resistance to CQ was first observed in 1957 in Southeast Asia (SEA), and subsequently arrived and spread across Africa from the late 1970s, contributing to the end of this ambitious global eradication effort [4].

Resistance to CQ has been studied intensively. The CQ resistance transporter (*pfcrt*, chromosome [chr.] 7) was originally identified using a *P. falciparum* genetic cross conducted between a CQ-resistant SEA parasite and a CQ-sensitive South American parasite generated in a

chimpanzee host [5,6]. Twenty years of intensive research revealed the mechanistic role of CRT in drug resistance [7,8], its location in the digestive vacuole membrane, and its natural function transporting short peptides from the digestive vacuole into the cytoplasm [9]. CQ kills parasites by interfering with hemoglobin digestion in the digestive vacuole, preventing conversion of heme, a toxic by-product of hemoglobin digestion, into inert hemozoin crystals. Parasites carrying CQ-resistance mutations at CRT transport CQ out of the food vacuole, away from the site of drug action [7,8]. The *pfcrt* K76T single nucleotide polymorphism (SNP) is widely used as a molecular marker for CQ resistance [10], while additional variants within *pfcrt* modulate levels of resistance to CQ [11] and other quinoline drugs [12], and determine associated fitness costs [13]. While mutations in a second transporter located in the food vacuole membrane, the multidrug resistance transporter (*pfmdr1*), have been shown to modulate CQ resistance in some genetic backgrounds [14], the role of other genes in CQ resistance evolution remains unclear. In this work we sought to understand the contribution of additional parasite loci to CQ-resistance evolution using a combination of population genomics, experimental genetic crosses, and gene editing.

Results

Strong signatures of selection on pfaat1

Longitudinal population genomic data can provide compelling evidence of the evolution of drug resistance loci [15]. We conducted a longitudinal whole genome sequence analysis of 600 *P. falciparum* genomes collected between 1984 and 2014 in The Gambia to examine signatures of selection under drug pressure (Supplementary Table 1). Following filtration using genotype missingness (<10%) and minor allele frequency (>2%), we retained 16,385 biallelic SNP loci from 321 isolates (1984 [134], 1990 [13], 2001 [34], 2008 [75], and 2014 [65]. The *pfcrt* K76T allele associated with CQ-resistance increased from 0% in 1984 to 70% in 2014. Intriguingly,

there was also rapid allele frequency change on chr. 6: the strongest differentiation is seen at an

S258L mutation in a putative amino acid transporter, *pfaat1* (PF3D7_0629500, chr. 6), which

increased during the same time period from 0% to 87% (Fig. 1a). Assuming a generation time

(mosquito to mosquito) of 6 months for malaria parasites, these changes were driven by selection

coefficients of 0.17 for *pfaat1* S258L, and 0.11 for *pfcrt* K76T (Supplementary Fig. 1). Both

pfaat1 S258L and *pfcrt* K76T mutations were absent in 1984 samples, but present in 1990,

suggesting that they arose and spread in a short time window. Both *pfaat1* and *pfcrt* showed

similar temporal haplotype structures in The Gambia (Supplementary Fig. 2). These were

characterized by almost complete replacement of well differentiated haplotypes at both loci

between 1984 and 2014. During this period, we also observed major temporal changes in another

known drug resistance locus (*pfdhfr*) (chr. 4) (Fig. 1b) [16]. That these rapid changes in allele

frequency occur at *pfcrt*, *pfaat1* and *pfdhfr*, but not elsewhere in the genome (Fig. 1b), provides

unambiguous evidence for strong directional selection.

Further evidence of strong selection on *pfaat1* and *pfcrt* came from the analysis of identity-by-

descent (IBD) in Gambian parasite genomes. We saw the strongest signals of IBD in the genome

around both *pfaat1* and *pfcrt* (Fig. 1c). These signals are dramatic, because there is minimal IBD

elsewhere in the genome, with the exception of a strong signal centering on *pfdhfr* after 2008.

Interestingly, the strong IBD is observed in all four temporal samples examined including 1984,

prior to the spread of either *pfaat1* S258L or *pfcrt* K76T. However, only a single synonymous

variant at *pfaat1* (I552I) and none of the CQ-resistant associated mutant variants in *pfcrt* were

present at that time. CQ was the first line treatment across Africa from the 1950s. These results

are consistent with the possibility of CQ-driven selective sweeps conferring low level CQ-

resistance prior to 1984, perhaps targeting promotor regions of resistance-associated genes.

pfaat1 has also been selected in global locations: this is evident from prior population genomic

analyses from Africa [17], SEA [18] and South America [19]. Plots summarizing IBD in these regions

are provided in Supplementary Fig. 3.

Patterns of linkage disequilibrium (LD) provide further evidence for functional linkage between

pfcrt and *pfaat1*. The strongest genome-wide signal of interchromosomal LD was found between

these two loci both in our Gambian data (Supplementary Fig. 4) and in samples from across

Africa [20]. LD between *pfaat1* and *pfcrt* was strongest in 2001, and then decayed in 2008 and

2014 (Supplementary Fig. 4&5), consistent with maintenance of LD during intensive CQ usage,

and subsequent LD decay after CQ monotherapy was replaced by sulfadoxine-pyrimethamine +

CQ combinations in 2004, and then with artemisinin combinations in 2008 [16].

Correlations in allele frequencies are expected between *pfcrt* and *pfaat1* if these loci are

interacting or are co-selected. Frequencies of the CVIET haplotype for amino acids 72-76 in

CRT are significantly correlated with allele frequencies of *pfaat1* S258L in West Africa (R^2 =

0.65, *p* = 0.0017) or across all African populations (R^2 = 0.44, *p* = 0.0021) (Supplementary Fig.

6). This analysis further strengthens the argument for co-evolution or epistasis between these two

genes.

Divergent selection on pfaat1 in Southeast Asia

We examined the haplotype structure of *pfaat1* from *P. falciparum* genomes (MalariaGEN

release 6 [21]) (Fig. 2, Supplementary Table 2). The *pfaat1* S258L SNP is at high frequency in

SEA (58%) but is found on divergent flanking haplotypes suggesting an independent origin from

the *pfaat1* S258L in The Gambia and elsewhere in Africa (Fig. 2 c & d, Supplementary Fig. 7).

Hendon et al [18] reached the same conclusion for the chr. 6 region using an identity-by-descent

analysis of parasites from global locations. Convergent evolution of *pfaat1* S258L provides

further evidence for selection, and contrasts with *pfcrt* and *pfdhfr*, where resistance alleles that

spread in Africa had an Asian origin [1,2]. *Pfaat1* evolution is more complex in SEA than

elsewhere in the world. There are three additional common derived amino acid changes in SEA.

Pfaat1 F313S has spread close to fixation in SEA (total 96%, F_{ST} = 0.91 compared with African

samples) paired with *pfaat1* S258L (55%), Q454E (15%), or K541N (22%). We speculate that

these geographically localized *pfaat1* haplotypes have played an important role in CQ-resistance

evolution in SEA and could also reflect geographic differences in the historical use of other

quinoline drugs (mefloquine, quinine, piperaquine, lumefantrine) in this region [22].

Parasite genetic crosses using humanized mice identify a QTL containing pfaat1

P. falciparum genetic crosses can be achieved with human-liver chimeric mice, reviving and

enhancing this powerful tool for malaria genetics [23,24], after use of great apes for research was

banned. We conducted three independent biological replicates of a cross between the CQ-

sensitive West African parasite, 3D7, and a recently isolated CQ-resistant parasite from the

Thailand-Myanmar border, NHP4026 (Supplementary Table 3). We then compared genome-

wide allele frequencies in CQ-treated and control-treated progeny pools to identify quantitative

trait loci (QTL). This bulk segregant analysis (BSA) [25] of progeny parasites robustly identified

the chr. 7 locus containing *pfcrt* as expected, validating our approach (Fig. 3a and Supplementary

Fig. 8 & 9). We were also intrigued to see a significant QTL on chr. 6 in each of the replicate

crosses (Fig. 3, Supplementary Fig. 8-10). We prioritized genes within the 95% confidence

interval of each QTL (Supplementary Table 4) by inspecting the SNPs and indels that

differentiated the two parents (Supplementary Table 5). The chr. 6 QTL spanned from 1,013 kb

to 1,283 kb (270 kb) and contained 60 genes. Of these, 54 are expressed in blood stages, and 27

have non-synonymous mutations that differentiate 3D7 from NHP4026. *pfaat1* was located at the

peak of the chr. 6 QTL (Fig. 3c). NHP4026 carried two derived non-synonymous mutations in

pfaat1 (S258L and F313S) compared to 3D7, which carries the ancestral allele at each of these

codon sites. We thus hypothesized that one or both of these *pfaat1* SNPs may be driving the chr. 6 QTL.

We isolated individual clones from the bulk 3D7 $\times$ NHP4026 F_1 progeny to recover clones with all combinations of parental alleles at the chr. 6 and chr. 7 QTL loci. We cloned parasites both from a bulk progeny culture that was CQ-selected (96 hours at 250 nM CQ) and from a control culture. This generated 155 clonal progeny: 100 from the CQ-selected culture, 62 of which were genetically unique and 55 from the untreated control culture, of which 47 were unique (Fig. 4a). We compared allele frequencies between these two progeny populations (Fig. 4b), revealing significant differences at both chr. 6 and chr. 7 QTL regions, paralleling the BSA results. We observed a dramatic depletion of the NHP4026 CQ-resistant allele at the chr. 7 QTL in control treated cultures, consistent with strong selection against CQ resistant *pfcrt* alleles in the absence of CQ-selection. Conversely, all progeny isolated after CQ treatment harbored the NHP4026 CQ-resistant *pfcrt* allele. The inheritance of the *pfcrt* locus (chr. 7) and the *pfaat1* locus (chr. 6) was tightly linked in the isolated clones (Fig. 4c). To further examine whether the cross data were consistent with epistasis or co-selection, we examined a larger sample of recombinant clones isolated from five independent iterations of this genetic cross in the absence of CQ selection. This revealed significant under-representation of clones with genotype *pfcrt* 76<u>T</u> and *pfaat1* 258S/313F (WT) (Supplementary Table 6, $X^2 = 12.295$, *p*-value = 0.0005). These results are consistent with the strong LD between these loci observed in nature (Supplementary Fig. 4) [20] and suggest a functional relationship between the two loci. A role for *pfaat1* S258<u>L</u>/F313<u>S</u> in compensating for the reduced fitness of parasites bearing *pfcrt* K76<u>T</u> is one likely explanation for the observed results.

We next measured *in vitro* CQ IC_{50} values for 18 parasites (a set of 16 progeny and both parents), carrying all combinations of the chr. 6 and chr. 7 QTL alleles (Supplementary Fig. 11,

Supplementary Table 7). The NHP4026 parent was the most CQ-resistant parasite tested. All

progeny that inherited NHP4026 *pfcrt* showed a CQ-resistant phenotype while all progeny that

inherited 3D7 *pfcrt* were CQ-sensitive, consistent with previous reports. The effect *of pfcrt*

alleles on parasite CQ resistance was significant based on a two-way ANOVA test (p=7.52 × 10^{-}

^{11}). We did not see an impact of the *pfaat1* genotypes on IC$_{50}$ in clones carrying *pfcrt* 76T ($p =$

0.06) or *pfcrt* 76K ($p = 0.19$).   This analysis has limited power because only two progeny

parasites were recovered with *pfaat1* 258S/313F (WT) in combination with *pfcrt* 76T (Fig.4a,

Supplementary Fig. 11), but is consistent with the *pfAAT1* QTL being driven by parasite fitness

in our genetic crosses. We therefore focused on gene manipulation of isogenic parasites for

functional analysis.

Functional validation of the role of pfaat1 in CQ resistance

We utilized CRISPR/Cas9 modification of the NHP4026 CQ-resistant parent to investigate the

effects of mutations in *pfaat1* on CQ IC$_{50}$ drug response and parasite fitness (Fig. 5). NHP4026

pfaat1 carries the two most common SEA non-synonymous changes (S258L and F313S) (Fig.

2), relative to the sensitive 3D7 parent. We edited these positions back to the ancestral state both

singly and in combination and confirmed the modifications in three clones isolated from

independent CRISPR experiments for each allelic change (Fig. 5a). We then determined CQ IC$_{50}$

and measured fitness using pairwise competition experiments for parental NHP4026$^{258L/313S}$, the

single mutations NHP4026$^{258L/313F}$, NHP4026$^{258S/313S}$ and the ancestral allele NHP4026$^{258S/313F}$.

This revealed a highly significant impact of the S258L mutation, which increased CQ IC$_{50}$ 1.5-

fold, and a more moderate but significant impact of F313S and the double mutation

(S258L/F313S) (Fig. 5b, Supplementary Table 8). The observation that 258L shows an elevated

IC$_{50}$ only in combination with the ancestral 313F allele reveals an epistatic interaction between

these amino acid variants (Fig 5B).

We also examined the impact of the S258$\underline{L}$ and F313$\underline{S}$ substitutions on responses to other quinoline drugs. The results revealed significant impacts of PfAAT1 substitutions on quinine, amodiaquine and lumefantrine IC_{50} responses, and no impact on mefloquine IC_{50} (Supplementary Fig. 12). Notably, these IC_{50} value shifts were well below the threshold associated with clinical resistance. Consequently, although mutations in *pfaat1* can subtly impact susceptibly to a range of compounds, these results are consistent with CQ treatment being the primary selective force that drove the *pfaat1* S258$\underline{L}$ and F313$\underline{S}$ mutations along with those in *pfcrt*.

Mutations conferring drug resistance often carry fitness costs in the absence of drug treatment. We thus examined parasite fitness by conducting pairwise competition experiments with the parental NHP4026 parasite against the same mutant *pfaat1* parasites created above. This revealed significant differences in fitness (Fig. 5c). The 258$\underline{L}$/313F allele that showed a selective sweep in The Gambia, was the least fit of all genotypes, the ancestral allele (258S/313F) carried by the 3D7 parent was the most fit, while the 258S/313$\underline{S}$ mutation, showed a similar fitness to the NHP4026 parent (258$\underline{L}$/313$\underline{S}$). These results also revealed strong epistatic interactions in fitness. While the 258$\underline{L}$/313F allele that conferred high CQ IC_{50} (Fig. 5b), carried a heavy fitness penalty (Fig. 5c), fitness was partially restored by addition of the 313$\underline{S}$ mutation to generate the 258$\underline{L}$/313$\underline{S}$ allele that predominates in SEA. Together these results show that the *pfaat1* S258$\underline{L}$ substitution underpins a 1.5-fold increase in CQ-resistance that likely drove its selective spread in The Gambia. However, S258$\underline{L}$ carries a high fitness cost that in SEA parasites was likely mitigated by addition of the compensatory substitution, F313$\underline{S}$. Overall, these results demonstrate a large impact of *pfaat1* mutations on fitness of parasites carrying *pfcrt* K76$\underline{T}$ resistance alleles.

The editing experiments reveal that clones carrying the ancestral *pfaat1* allele in combination with *pfcrt* K76$\underline{T}$ show the highest fitness. In contrast, the close association of *pfaat1*

S258L/F313S with *pfcrt* K76T in progeny from the genetic crosses revealed the opposite relationship. We speculate that these opposing results may reflect differing selection pressures in blood stage parasites in the case of CRISPR experiments, or in the mosquito and liver stages of the life cycle in the case of genetic crosses.

To further understand how *pfaat1* S258L impacts parasite phenotype, we used a yeast heterologous expression system. WT *pfaat1* is expressed in the yeast plasma membrane [26], where it increases quinine and CQ uptake conferring sensitivity to quinoline drugs, resulting in reduced growth. CQ uptake can be competitively inhibited by the aromatic amino acid tryptophan, suggesting a role for *pfaat1* in drug and amino acid transport [26]. We therefore expressed *pfaat1* S258L in yeast, which restored yeast growth in the presence of high levels of CQ (Supplementary Fig. 13). Interestingly, expression of another amino acid variant (T162E), responsible for CQ-resistance in rodent malaria parasites (*Plasmodium chabaudi*) [27], also prevents accumulation of quinoline drugs within yeast cells and restores cell growth in the presence of CQ [26]. Together, these new and published results suggest that yeast expression of *pfaat1* mutations impact resistance and fitness by altering the rates of amino acid and CQ transport.

We evaluated three-dimensional structural models based on the 3D7 PfAAT1 amino acid sequence using AlphaFold [28] and I-TASSER [29] (Supplementary Fig. 14). While CRT has ten membrane-spanning helices [30], AAT1 has eleven; this was corroborated using the sequence-based membrane topology prediction tool TOPCONS [31]. The common AAT1 mutations S258L, F313S and Q454E are situated in membrane-spanning domains, while K541L is in a loop linking domains 9 and 10. The location of these high frequency non-synonymous changes in membrane spanning domains has strong parallels with CRT evolution [30] and is consistent with a functional role for these amino acids in transporter function.

Discussion

Identification of *pfcrt* as the major determinant of CQ resistance was a breakthrough that transformed the malaria drug resistance research landscape, but the contribution of additional genetic factors in the evolution and maintenance of CQ resistance remained unclear [32,33]. By combining longitudinal population genomic analysis spanning the emergence of CQ resistance in The Gambia, analysis of bulk populations and progeny from controlled genetic crosses, and functional validation using both yeast and *P. falciparum*, we find clear evidence that a second locus, *pfaat1*, has played a central role in CQ resistance. This powerful combination of approaches allowed us to examine critical *pfaat1* variants that contribute to the architecture of CQ-resistance and interactions between *pfcrt* and *pfaat1*.

Our results provide compelling evidence that consolidates disparate observations from several systems suggesting a role for *pfaat1* in drug resistance evolution. In the rodent malaria parasite *P. chabaudi*, a mutation (T162E) in the orthologous amino acid transporter (*pcaat1*) was found to be a determinant of low level CQ resistance in laboratory-evolved resistance [27]. In *P. falciparum* genome-wide association studies (GWAS), the S258L mutation of *pfaat1* was associated with CQ resistance in field isolates collected along the China-Myanmar border [34], while *pfcrt* K76T and *pfaat1* S258L show the strongest LD between physically unlinked chromosomes genome-wide [20]. In addition, mutations in *pfaat1* have been linked to the *in vitro* evolution of resistance in *P. falciparum* to three different drug scaffolds [35]. Previous work identified strong signatures of recent selection in parasites in Africa at regions surrounding *pfcrt*, *pfaat1*, and other drug resistance loci [16,17,36]; similar signatures of selection are seen in Asia and South America [18,19], while *pfaat1* was highlighted in a list of *P. falciparum* genes showing extreme geographical differentiation [21].

The different *pfaat1* haplotypes in Africa and Asia may be partly responsible for the contrasting evolution of CQ-resistance in these two continents. CQ-resistant parasites, carrying both *pfcrt* K76T and *pfaat1* S258L spread across Africa, but after removal of CQ as the first line drug, the prevalence of CQ resistant parasites declined in many countries [37-39]. This is consistent with the low fitness of parasites carrying *pfcrt* K76T and *pfaat1* S258L in the absence of drug pressure, and intense competition within malaria parasite infection in Africa [40].

In contrast, *pfcrt* K76T has remained at or near fixation in many SEA countries [21,41] (Fig. 2). On the Thailand-Myanmar border, CQ-resistance has remained at fixation since 1995, when CQ was removed as first line treatment of *P. falciparum* malaria [41]. Our *pfaat1* mutagenesis results demonstrate that parasites bearing *pfaat1* 258L/313S show reduced IC_{50}, but elevated fitness relative to *pfaat1* 258L/313F. The restoration of fitness by F313S may help to explain retention of CQ-resistant *pfcrt* K76T alleles in SEA. The alternative hypothesis – that high frequencies of F313S mutations are driven by widespread use of other quinoline partner drugs in SEA [42] is not supported, because we see only minor impacts of this substitution on response to lumefantrine, quinine, mefloquine and amodiaquine (Supplementary Fig. 12).

Mutations in *pfcrt* confer CQ-resistance by enabling efflux of CQ across the digestive vacuole membrane, away from its site of action [8]. PfAAT1 is also located in the digestive vacuole membrane, where it likely acts as a bidirectional transporter of aromatic amino acids [9,43]. Given the structural similarity of quinoline drugs and aromatic amino acids, *pfaat1* mutations may modulate the ability of PfAAT1 to transport CQ and/or amino acids [26,43]. The *pfaat1* S258L mutation could potentiate resistance by either increasing efflux of CQ out of the digestive vacuole or reducing the rate of entry into the vacuole. Given that this *pfaat1* mutation blocks entry of quinoline drugs into yeast cells when heterologously expressed in the yeast cell membrane [26], we hypothesize that the *pfaat1* S258L mutation reduces CQ uptake into the food

vacuole (Fig. 6). Our mutagenesis analyses show that the S258L allele has a high fitness cost, perhaps due to a decreased capacity for the amino acid transport from the vacuole. Interestingly, comparison of the *pfaat1* S258L/F313S haplotype segregating in our genetic cross with the WT *pfaat1* allele generated using gene editing, revealed only marginal increases in IC_{50} and limited reductions in fitness. This is consistent with the F313S mutation restoring the natural *pfaat1* function of transporting amino acids, thereby reducing osmotic stress and starvation, while also partially reducing levels of CQ-resistance (Fig. 6). That this haplotype has reached high frequency in SEA, may contribute to the maintenance of *pfcrt* K76T alleles long after the removal of CQ as a first line drug.

Our results reveal hidden complexity in CQ-resistance evolution: drug treatment has driven global selective sweeps acting on mutations in an additional transporter (PfAAT1) located in the *P. falciparum* digestive vacuole membrane, which fine tune the balance between nutrient and drug transport, revealing evidence for epistasis and compensation, and impacting both drug resistance and fitness.

References

Wootton, J. C. *et al.* Genetic diversity and chloroquine selective sweeps in Plasmodium falciparum. *Nature* **418**, 320-323, doi:10.1038/nature00813 (2002).

Roper, C. *et al.* Intercontinental spread of pyrimethamine-resistant malaria. *Science* **305**, 1124, doi:10.1126/science.1098876 (2004).

Ariey, F. *et al.* A molecular marker of artemisinin-resistant Plasmodium falciparum malaria. *Nature* **505**, 50-55, doi:10.1038/nature12876 (2014).

Wellems, T. E. & Plowe, C. V. Chloroquine-resistant malaria. *J Infect Dis* **184**, 770-776, doi:10.1086/322858 (2001).

Su, X., Kirkman, L. A., Fujioka, H. & Wellems, T. E. Complex polymorphisms in an approximately 330 kDa protein are linked to chloroquine-resistant P. falciparum in Southeast Asia and Africa. *Cell* **91**, 593-603, doi:10.1016/s0092-8674(00)80447-x (1997).

Wellems, T. E., Walker-Jonah, A. & Panton, L. J. Genetic mapping of the chloroquine-resistance locus on Plasmodium falciparum chromosome 7. *Proc Natl Acad Sci U S A* **88**, 3382-3386, doi:10.1073/pnas.88.8.3382 (1991).

Fidock, D. A. *et al.* Mutations in the P. falciparum digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance. *Mol Cell* **6**, 861-871, doi:10.1016/s1097-2765(05)00077-8 (2000).

Ecker, A., Lehane, A. M., Clain, J. & Fidock, D. A. PfCRT and its role in antimalarial drug resistance. *Trends Parasitol* **28**, 504-514, doi:10.1016/j.pt.2012.08.002 (2012).

Shafik, S. H. *et al.* The natural function of the malaria parasite's chloroquine resistance transporter. *Nat Commun* **11**, 3922, doi:10.1038/s41467-020-17781-6 (2020).

Djimde, A. *et al.* A molecular marker for chloroquine-resistant falciparum malaria. *N Engl J Med* **344**, 257-263, doi:10.1056/NEJM200101253440403 (2001).

Pelleau, S. *et al.* Adaptive evolution of malaria parasites in French Guiana: Reversal of chloroquine resistance by acquisition of a mutation in pfcrt. *Proc Natl Acad Sci U S A* **112**, 11672-11677, doi:10.1073/pnas.1507142112 (2015).

Ross, L. S. *et al.* Emerging Southeast Asian PfCRT mutations confer Plasmodium falciparum resistance to the first-line antimalarial piperaquine. *Nat Commun* **9**, 3314, doi:10.1038/s41467-018-05652-0 (2018).

Dhingra, S. K. *et al.* Global Spread of Mutant PfCRT and Its Pleiotropic Impact on Plasmodium falciparum Multidrug Resistance and Fitness. *mBio* **10**, doi:10.1128/mBio.02731-18 (2019).

Shafik, S. H., Richards, S. N., Corry, B. & Martin, R. E. Mechanistic basis for multidrug resistance and collateral drug sensitivity conferred to the malaria parasite by polymorphisms in PfMDR1 and PfCRT. *PLoS Biol* **20**, e3001616, doi:10.1371/journal.pbio.3001616 (2022).

Cerqueira, G. C. *et al.* Longitudinal genomic surveillance of Plasmodium falciparum malaria parasites reveals complex genomic architecture of emerging artemisinin resistance. *Genome Biol* **18**, 78, doi:10.1186/s13059-017-1204-4 (2017).

Nwakanma, D. C. *et al.* Changes in malaria parasite drug resistance in an endemic population over a 25-year period with resulting genomic evidence of selection. *J Infect Dis* **209**, 1126-1135, doi:10.1093/infdis/jit618 (2014).

Amambua-Ngwa, A. *et al.* Major subpopulations of Plasmodium falciparum in sub-Saharan Africa. *Science* **365**, 813-816, doi:10.1126/science.aav5427 (2019).

Henden, L., Lee, S., Mueller, I., Barry, A. & Bahlo, M. Identity-by-descent analyses for measuring population dynamics and selection in recombining pathogens. *PLoS Genet* **14**, e1007279, doi:10.1371/journal.pgen.1007279 (2018).

Carrasquilla, M. *et al.* Resolving drug selection and migration in an inbred South American Plasmodium falciparum population with identity-by-descent analysis. *bioRxiv*, 2022.2002.2018.480973, doi:10.1101/2022.02.18.480973 (2022).

Band, G. *et al.* Malaria protection due to sickle haemoglobin depends on parasite genotype. *Nature* **602**, 106-111, doi:10.1038/s41586-021-04288-3 (2022).

MalariaGen *et al.* An open dataset of Plasmodium falciparum genome variation in 7,000 worldwide samples. *Wellcome Open Res* **6**, 42, doi:10.12688/wellcomeopenres.16168.2 (2021).

Martin, R. E., Shafik, S. H. & Richards, S. N. Mechanisms of resistance to the partner drugs of artemisinin in the malaria parasite. *Curr Opin Pharmacol* **42**, 71-80, doi:10.1016/j.coph.2018.07.010 (2018).

Vendrely, K. M., Kumar, S., Li, X. & Vaughan, A. M. Humanized Mice and the Rebirth of Malaria Genetic Crosses. *Trends Parasitol* **36**, 850-863, doi:10.1016/j.pt.2020.07.009 (2020).

Vaughan, A. M. *et al.* Plasmodium falciparum genetic crosses in a humanized mouse model. *Nat Methods* **12**, 631-633, doi:10.1038/nmeth.3432 (2015).

Brenneman, K. V. *et al.* Optimizing bulk segregant analysis of drug resistance using Plasmodium falciparum genetic crosses conducted in humanized mice. *iScience* **25**, 104095, doi:10.1016/j.isci.2022.104095 (2022).

Tindall, S. M. *et al.* Heterologous Expression of a Novel Drug Transporter from the Malaria Parasite Alters Resistance to Quinoline Antimalarials. *Sci Rep* **8**, 2464, doi:10.1038/s41598-018-20816-0 (2018).

Modrzynska, K. K. *et al.* Quantitative genome re-sequencing defines multiple mutations conferring chloroquine resistance in rodent malaria. *BMC Genomics* **13**, 106, doi:10.1186/1471-2164-13-106 (2012).

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

Yang, J. *et al.* The I-TASSER Suite: protein structure and function prediction. *Nat Methods* **12**, 7-8, doi:10.1038/nmeth.3213 (2015).

Kim, J. *et al.* Structure and drug resistance of the Plasmodium falciparum transporter PfCRT. *Nature* **576**, 315-320, doi:10.1038/s41586-019-1795-x (2019).

Tsirigos, K. D., Peters, C., Shu, N., Kall, L. & Elofsson, A. The TOPCONS web server for consensus prediction of membrane protein topology and signal peptides. *Nucleic Acids Res* **43**, W401-407, doi:10.1093/nar/gkv485 (2015).

Callaghan, P. S., Hassett, M. R. & Roepe, P. D. Functional Comparison of 45 Naturally Occurring Isoforms of the Plasmodium falciparum Chloroquine Resistance Transporter (PfCRT). *Biochemistry* **54**, 5083-5094, doi:10.1021/acs.biochem.5b00412 (2015).

Patel, J. J. *et al.* Chloroquine susceptibility and reversibility in a Plasmodium falciparum genetic cross. *Mol Microbiol* **78**, 770-787, doi:10.1111/j.1365-2958.2010.07366.x (2010).

Wang, Z. *et al.* Genome-wide association analysis identifies genetic loci associated with resistance to multiple antimalarials in Plasmodium falciparum from China-Myanmar border. *Sci Rep* **6**, 33891, doi:10.1038/srep33891 (2016).

Cowell, A. N. *et al.* Mapping the malaria parasite druggable genome by using in vitro evolution and chemogenomics. *Science* **359**, 191-199, doi:10.1126/science.aan4472 (2018).

36   Amambua-Ngwa, A. *et al.* SNP genotyping identifies new signatures of selection in a
deep sample of West African Plasmodium falciparum malaria parasites. *Mol Biol Evol*
29, 3249-3253, doi:10.1093/molbev/mss151 (2012).
37   Laufer, M. K. *et al.* Return of chloroquine-susceptible falciparum malaria in Malawi was
a reexpansion of diverse susceptible parasites. *J Infect Dis* **202**, 801-808,
doi:10.1086/655659 (2010).
38   Kublin, J. G. *et al.* Reemergence of chloroquine-sensitive Plasmodium falciparum
malaria after cessation of chloroquine use in Malawi. *J Infect Dis* **187**, 1870-1875,
doi:10.1086/375419 (2003).
39   Mharakurwa, S. *et al.* Steep Rebound of Chloroquine-Sensitive Plasmodium falciparum
in Zimbabwe. *J Infect Dis* **223**, 306-309, doi:10.1093/infdis/jiaa368 (2021).
40   Anderson, T. J. *et al.* Microsatellite markers reveal a spectrum of population structures in
the malaria parasite Plasmodium falciparum. *Mol Biol Evol* **17**, 1467-1482,
doi:10.1093/oxfordjournals.molbev.a026247 (2000).
41   Anderson, T. J. *et al.* Geographical distribution of selected and putatively neutral SNPs in
Southeast Asian malaria parasites. *Mol Biol Evol* **22**, 2362-2374,
doi:10.1093/molbev/msi235 (2005).
42   Ashley, E. A. & Phyo, A. P. Drugs in Development for Malaria. *Drugs* **78**, 861-879,
doi:10.1007/s40265-018-0911-9 (2018).
43   Martin, R. E. The transportome of the malaria parasite. *Biol Rev Camb Philos Soc* **95**,
305-332, doi:10.1111/brv.12565 (2020).
438

439 **Figures**

441 **Fig. 1. Rapid allele frequency change and strong signals of selection around *pfaat1* in The**

442 **Gambia.** (**a**) Temporal allele frequency change at SNPs coding for *pfaat1* S258L and *pfcrt* K76T

443 between 1984 and 2014. The map and expanded West African region show the location of The

Gambia. (**b**) Significance of haplotype differentiation across temporal populations of *P. falciparum* parasites determined using hapFLK. Significance thresholds at -$\log_{10}$(FDR-corrected *p*-value) = 5 are indicated with red dotted horizontal lines. Regions within the top 1% tail of FDR-corrected *p*-values are marked with gene symbols. The strongest signals genome-wide seen are around *pfcrt*, *pfaat1*, and *pfdhfr* (which is involved in pyrimethamine resistance). (**c**) Identity-by-descent (IBD), quantified with the isoRelate (iR) statistic, for temporal populations sampled from The Gambia. Consistently high peaks of IBD around *pfcrt* and *pfaat1* are seen for parasite populations in all years of sampling. The 1990 sample (n=13) is not shown in panel C.

Fig. 2. Distinctive trajectory of *pfaat1* evolution in Southeast Asia. (**a**) Global distribution of

pfaat1 alleles. (**b**) Comparable maps showing percentages of *pfcrt* haplotypes for amino acids

72-76. The colored segments show the major *pfcrt* haplotypes varying the K76T mutation. We

used dataset from MalariaGEN release 6 for *pfaat1* and *pfcrt* allele frequency analysis. Data used

for the figure is contained in Supplementary Table 2. Only samples with monoclonal infections

(N = 4051) were included (1233 from West Africa [WAF], 415 from East Africa [EAF], 170

from Central Africa [CAF], 994 from east Southeast Asia [ESEA], 998 from west Southeast Asia [WSEA], 37 from South Asia [SA], 37 from South America [SM], and 167 from the Pacific Ocean region [PO]). (C and D) show minimum spanning networks of haplotypes colored by *pfaat1* allele (**c**) and geographical location (**d**), respectively. Networks were constructed from 50 kb genome regions centered by *pfaat1* (25 kb up- and down-stream. This spans the genome regions showing LD around *pfaat1* (Fig S5). 581 genomes with the highest sequence coverage were used to generate the network. The networks were generated based on 1847 SNPs (at least one mutant in the full dataset - MalariaGEN release 6). Circle size indicates number of samples represented (smallest, 1; largest, 87). Haplotypes from the same region (Asia or Africa) were clustered together, indicating independent origin of *pfaat1* alleles.

Fig. 3. Genetic crosses and bulk segregant analysis reveal two QTL after CQ selection. (**a**) Allele frequency plots across the genome before and after CQ treatment. Lines with the same color indicate results from technical replicates. (**b**) QTLs identified using the G' approach. Lines with the same color indicate results from technical replicates. Panels (**a**) and (**b**) include results from BSA with 48 hour CQ treatment with samples collected at day 4. See Figures S8 and S9 for

the complete BSA from different collection time points and drug treatment duration under

different CQ concentrations. (**c**) Fine mapping of the chr. 6 QTL. The 95% confidence intervals

(CIs) were calculated from the 250 nM CQ treated samples, including data from different

collection time points (day 4 for 48 hour CQ treatment and day 5 for 96 hour CQ treatment),

pools (pool 1 and pool 2), and drug treatment duration (48 hour and 96 hour). Light cyan shadow

shows boundaries of the merged CIs of all the QTLs. Each line indicates one QTL; black dashed

line indicates threshold for QTL detection (*G prime* = 20). The vertical red dashed line indicates

pfaat1 location.

Fig. 4. Analysis of cloned progeny reveals linkage and epistatic interactions between *pfcrt* and *pfaat1*. (**a**) Allelic inheritance of 109 unique recombinant progeny. Black and red blocks indicate alleles from 3D7 and NHP4026, separately. Vertical grey lines show non-core regions where no SNPs were genotyped. Clones isolated from recombinant progeny pools with or without CQ treatment are labeled on the left side of the panel. *pfaat1* and *pfcrt* alleles are labeled on the right side of the panel. WT indicates *pfaat1* and *pfcrt* alleles from 3D7 and MUT indicates alleles from NHP4026. The location of *pfaat1* and *pfcrt* are marked using black triangles on the top of the panel. (**b**) Genome-wide 3D7 allele frequency plot of unique progeny cloned from

494   pools after 96-hours of CQ (250 nM) treatment (blue) or from control pools (gold). (**c**) Linkage

495   between loci on different chromosomes measured by Fisher's exact test. The dotted vertical line

496   marks the Bonferroni corrected significance threshold, while points shown in red are

497   comparisons between SNPs flanking *pfaat1* and *pfcrt*. Supplementary Table 6 shows non-

498   random associations between genotypes in parasite clones recovered from untreated cultures.

Fig. 5. Allelic replacement impacts drug response and parasite fitness. (**a**) CRISPR/Cas9

gene editing. Starting with the NHP4026 parent, we generated all combinations of the SNP-states

at *pfaat1*. (**b**) Drug response. Each dot indicates one replicate IC_{50} measurement: we used 2-4

independent CRISPR edited clones for each haplotype examined. Haplotypes are shown on the

x-axis with derived amino acids shown in red. Bars show means (±1 s.e.), while significant

506   differences between haplotypes are marked. (**c**) Fitness. The bars show relative fitness measured

507   in replicated competition experiments conducted in the absence of CQ. See Fig S17 for allele

508   frequency changes for each competition experiment. n.s., not significant.

Fig. 6. Model for involvement of *pfaat1* haplotypes in chloroquine resistance and fitness.

PfCRT (red) and PfAAT1 (blue) are both situated in the digestive vacuole (DV) membrane. (**a**)

Wildtype PfCRT and PfAAT1 transport peptides and aromatic amino acids respectively as well

as chloroquine (CQ). (**b**) PfCRT K76<u>T</u> exports CQ from the DV away from its site of action,

leading to elevated resistance but transports peptides inefficiently leading to a loss of fitness.

PfAAT1 S258<u>L</u> reduces entry of CQ into the DV, leading to elevated resistance, but amino acid

flux is affected leading to a loss of fitness. (**c**) The PfAAT1 S258<u>L</u>/F313<u>S</u> double mutation

increases CQ influx in comparison to the S258<u>L</u> alone but the amino acid transport function is

restored, leading to reduced IC$_{50}$ and increased fitness in the absence of drug treatment.

Methods

<u>Ethics approval and consent to participate</u>

The study was performed in strict accordance with the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health (NIH), USA. The Seattle Children's Research Institute (SCRI) has an Assurance from the Public Health Service (PHS) through the Office of Laboratory Animal Welfare (OLAW) for work approved by its Institutional Animal Care and Use Committee (IACUC). All of the work carried out in this study was specifically reviewed and approved by the SCRI IACUC.

<u>Project design</u>

The project design is summarized in Supplementary Fig. 15. In brief, we use (i) population genomic analyses (ii) genetic crosses and quantitative genetics analysis followed by (iii) functional analyses to investigate the role of additional loci in CQ resistance.

<u>Gambia population analysis</u>

P. falciparum genome sequences. *P*. *falciparum* infected blood samples collected from central (Farafenni) and coastal (Serrekunda) Gambia in 1984 and 2001 respectively, were processed for whole blood DNA and *P. falcip*arum genomes and deep sequenced at the Wellcome Trust Sanger Institute. Data from isolates collected from coastal Gambia in 2008 and 2014 had been published previously [44,45] (see Supplementary Table 1 for details). Prior to sequencing, *P. falciparum* genomes were amplified from whole blood DNA of each sample from 1984 and 2001 using selective whole genome amplification (sWGA) and then subjected to paired-end short-read sequencing on the Illumina HiSeq platform [46]. Short sequencing reads were mapped to the *P. falciparum* 3D7 reference genome using *bwa mem* (http://bio-bwa.sourceforge.net/). Mapping files (BAM) were sorted and deduplicated by Picard tools v2.0.1 (http://broadinstitute.github.io/picard/), and single-nucleotide polymorphism (SNP) and indel

were called with GATK Haplotype Caller (https://software.broadinstitute.org/gatk/) following

the best practices endorsed by the Pf3K project (https://www.malariagen.net/data/pf3K-5). Only

SNP genotypes were used for this study. VCF files were generated by chromosome, and then

were merged using *bcftools* (https://samtools.github.io/bcftools/bcftools.html) and filtered using

vcftools (https://vcftools.sourceforge.net/). After filtration, only biallelic SNP variants with a

VQSLOD score of at least 2, a map quality >30, and supported by no less than 5 reads per allelic

variant were remained. SNPs with minor allele frequency <2% were further removed from our

analysis. We also removed samples with > 10% genotypes missing. In the final dataset, there

were in total of 16,385 biallelic SNP loci and 321 isolates (1984 [134], 1990 [13], 2001 [34],

2008 [75], and 2014 [65]) remained for further analysis. The complexity of infection

(monogenomic or polygenomic) was estimated as the inbreeding coefficient *Fws* from the

merged VCF file using R package *Biomix*. The short-read sequence data for all isolates and those

from across Africa that have been previously published are available from the European

Nucleotide Archive, and sample identifiers are given in Supplementary Table 1.

Allele frequencies and pairwise differentiation. For each sample with a complexity of infection

greater than 1, the allele with most reads was retained for mixed-allele genotypes to create a

virtual haploid genome variation dataset. Allele frequencies were calculated in plink and

pairwise differences between temporal populations and genetic clusters were estimated by *Fst*

using Weir and Cockerham's method applied in the *hierfstat* package in R. The likelihood ratio

test for allele frequency difference pFST was further calculated using *vcflib*. For a combined

pFST *p*-value, the fisher method was performed in R *metaseq* package. The summary *p*-values

were corrected for multiple testing using Benjamini-Hochberg (BH) method.

Genome scans for selection. We considered samples collected in the same year as a single

population irrespective of the location of collection. We used the hapFLK approach to detect

signatures of positive selection through haplotype differentiation following hierarchical

clustering of Gambian temporal population groups compared to an outgroup from Tanzania, as

described by Fariello et al., 2013 [47]. *P*-values were computed for each SNP-specific value using

the Python script provided with the hapFLK program, and values were corrected for multiple

testing using the BH method. Secondly, we used pairwise relatedness based on identity by

descent to derive an iR statistic for each SNP as implemented by the *IsoRelate* [18] package in R.

Regions with overlapping iR and hapFLK -log10 *p*-values > 5 were considered as regions of

interest.

<u>Population analysis on *pfaat1* and *pfcrt* evolution</u>

Datasets. We included two datasets in this study: **1)** genotypes of 7,000 world-wide *P.*

falciparum samples from MalariaGEN Pf community project (version 6.0) [21]. This dataset

includes samples from South America (SM), West Africa (WA), Central Africa (CA), East

Africa (EA), South Asia (SA), the western part of Southeast Asia (WSEA), the eastern part of

Southeast Asia (ESEA) and Pacific Oceania (PO). **2)** We also included 194 Thailand samples

with whole genome sequencing data available from Cerqueira et al. [15], and merged them into the

WSEA population. Duplicate sequences were removed according to the sample's original ID

(Hypercode). Only samples with single parasite infections (within-host diversity $F_{WS} > 0.90$) and

>50% of SNP loci genotyped were included for further analysis. A total of 4051 samples

remained after filtration (Supplementary Table 2). Non-biallelic SNPs and heterozygous variant

calls were further removed from the dataset. We then extracted genotype data at *pfaat1* and *pfcrt*

gene regions and calculated the allele frequencies (Fig 2A).

Pfaat1 haplotypes and evolutionary relationships. To minimize the effect from recombination,

we extracted 1847 SNPs distributed within 25kb upstream and 25kb downstream of the *pfaat1*

gene. For clarity, only samples with all 1847 SNPs genotyped (581/4051) were used for evolutionary analysis. To visualize the population structure, we calculated the genetic distance between every two samples and generated a minimum spanning network (MSN, Fig 2B and Supplementary Fig. 4), using R package *poppr*. We compared genome sequences (PlasmoDB, version 46) between *P. falciparum* and *Plasmodium reichenowi* and extracted genotypes at 1803/1847 common loci. We then built a UPGMA tree rooted by *P. reichenowi* using the 581 haplotypes and 1803 SNPs (Supplementary Fig. 7), using the R packages *ape* and *phangorn* under default parameters. MSN network and UPGMA tree were plotted with ggplot2.

Genetic cross and bulk segregant analysis

Genetic cross preparation. We generated genetic crosses between parasite 3D7 and NHP4026 [48], using FRG NOD huHep mice with human chimeric livers and *A. stephensi* mosquitoes as described previously [23-25,49,50]. 3D7 has been maintained in the lab for decades and is CQ sensitive; while NHP4026 was recently cloned from a patient visiting the Shoklo Malaria Research Unit (SMRU) clinic on the Thailand-Myanmar border in 2007 and is CQ resistant (Supplementary Table 3). For the genetic cross 3D7×NHP4026, we generated three recombinant pools using independent cages of infected mosquitoes: these are independent pools of recombinants [48]. The estimated number of recombinant genotypes in each pool was ~2800 [48]. We used two of the pools (pool 1 and pool 2) that were maintained in culture media containing AlbuMAX for this study.

Drug treatment and sample collection. For each recombinant pool, the parasite culture was expanded under standard culture conditions as described in Brenneman et al. [25]. Briefly, cultures were maintained in complete media (CM) at 5% hematocrit in O^+ red blood cells (RBC) (Biochemed Services, Winchester, VA) at 37°C between a pH of 7.0-7.5 in an atmosphere of 5%

CO$_2$, 5% O$_2$ and 90% N$_2$. Media changes were performed every 48 hours and cultures were expanded to keep the parasitemia at ~1%. Once expanded sufficiently, each recombinant pool was divided into sixteen 0.5 ml aliquots while diluting to 1% parasitemia. The aliquots were then maintained in 48-well plates and treated with CQ (Supplementary Fig. 16). In total, we had 32 cultures: 2 pools × 4 CQ concentrations (0 [control], 50, 100 or 250 nM) × 2 drug duration time (48 hour or 96 hour) × 2 technical replicates. We define the day when drug was applied as day 0. After two days (48 hour) of drug treatment, the infected RBCs were washed with Phosphate-buffered saline (PBS) solution twice to remove residual drug. For the plate assigned for 48 hour CQ treatment (48-well plate 1), cultures were then maintained in complete media (CM); and samples were collected at day 0, 4 and 7. For the plate assigned for 96 hour CQ treatment (48-well plate 2), fresh CQ was added back to the culture media and treated for another 48 hour; and after a total of 96 hour CQ treatment, drug was removed and samples were collected at day 0, 5 and 10. CQ was dissolved in H$_2$O and diluted in incomplete media (ICM) (Gibco, Life Technologies). Culture medium was changed every 48 hours. Parasitemia was monitored using 20% Giemsa-stained slides and cultures were diluted to 1% parasitemia if the parasitemia was higher than 1%. Approximately 15 μl packed RBCs was collected per sample.

Library preparation and sequencing. We prepared Illumina libraries and sequenced both parents and the 96 segregant pools collected above. We extracted and purified genomic DNA using the Qiagen DNA mini kit and quantified the amount of DNA with Quant-iT™ PicoGreen® Assay (Invitrogen). For samples with less than 50 ng DNA obtained, we performed whole genome amplification (WGA) following Nair et al. [51]. WGA products were cleaned with KAPA Pure Beads (Roche Molecular Systems, Inc.) at a 1:1 ratio. We prepared sequencing libraries using 50-100 ng DNA or WGA product using KAPA HyperPlus Kit following the instruction with 3-

cycles of PCR. All libraries were sequenced at 150bp pair-end using Illumina Novaseq S4 or

Hiseq X sequencers, to get >100× genome coverage per sample.

Mapping and genotyping. We mapped the sequencing reads against the 3D7 reference genome

(PlasmoDB version 46) using BWA mem (http://bio-bwa.sourceforge.net/), and deduplicated

and   trans-formatted   the   alignment   files   using   picard   tools   v2.0.1

(http://broadinstitute.github.io/picard/). We recalibrated the base quality score based on a set of

verified known variants [52] using *BaseRecalibrator*, and called variants through *HaplotypeCaller*.

Both   functions   were   from   Genome   Analysis   Toolkit   GATK   v3.7

(https://software.broadinstitute.org/gatk/). Only variants located in the core genome regions

(defined in [52]) were called and used for further analysis.

Genotype of parents. We merged calls from the two parents using *GenotypeGVCFs* in GATK,

and applied *standard* filtration to the raw variant dataset as described in [53]. We recalibrated the

variant quality scores (VQSR) and removed loci with VQSR < 1. The final variants in VCF

format were annotated using snpEff v4.3 (https://pcingola.github.io/SnpEff/) with 3D7

(PlasmoDB, release46) as the reference. After filtration and annotation, we selected SNP loci

that are distinct in the two parents and used those SNPs for further bulk segregant analysis.

Bulk segregant analysis. We used statistical methods described previously in [25,48,50] for bulk

segregant analysis. The variant calls from segregant progeny pools were merged together.

Additionally, SNP loci with coverage < 30× were removed. We counted reads with genotypes of

each parent and calculated allele frequencies. Allele frequencies of 3D7 were plotted across the

genome, and outliers were removed following Hampel's rule [54] with a window size of 100 loci.

We performed the bulk segregant analysis using the R package *QTLseqr* [55]. Extreme-QTLs were

defined as regions with G prime > 20 [56]. Once a QTL was detected, we calculated an

approximate 95% confidence interval using Li's method [57] to localize causative genes.

Progeny cloning and phenotyping

Progeny cloning. Individual progeny were cloned via limiting dilution at 0.3 cells per well from

bulk cultures on day 10 after 96 hours of control/250nM CQ treatment. Individual wells with

parasites were determined by qPCR (as described in Button-Simons et al.[49]) and expanded to

larger cultures under standard culture conditions to obtain enough material for both

cryopreservation and genome sequencing.

Sequencing and genotyping. Cloned progeny were sequenced and genotyped as described above

in the Genetic cross and bulk segregant analysis section, with these modifications: 1) the cloned

progeny were sequenced at $25\times$ genome coverage; 2) SNP calls were filtered out if the coverage

was < 3 reads per sample.

Cloned progeny analysis. Unique recombinant progeny were identified from all cloned progeny

using a pipeline described in Button-Simons et al.[49]. Non-clonal progeny were identified based

on the number and distribution of heterozygous SNP calls. Selfed progeny were identified as

having greater than 90% sequence similarity to either parent. Unique recombinant progeny that

were sampled multiple times during cloning were identified as clusters of individual clonal

progeny with greater than 90% sequence similarity. We plotted frequencies of 3D7 alleles across

the genome in progeny populations with and without CQ treatment. Heatmaps were generated to

visualize inheritance patterns in individual unique recombinant progeny (Fig 4A). We selected

sixteen unique recombinant progeny with different allele combination at chromosome 6 and

chromosome 7 QTL regions for further CQ IC_{50} measurement (Fig S11).

Genome-wide linkage analysis on pfaat1 in cloned progeny. Fisher's exact test was used to test

for linkage between all inter-chromosomal pairs of loci across the set of 109 unique recombinant

progeny. The distribution of the -log of the resulting *p*-values was plotted in Fig 4C and the

significance cut-off was calculated based on a Bonferroni correction for the number of loci.

IC$_{50}$ measurement for cloned progeny. Cryopreserved stocks of 3D7, NHP4026,

3D7×NHP4026 progeny were thawed and grown in CM under standard culture conditions as

described above. Cultures were kept below 3% parasitemia with media changes every 48 hours.

Parents and progeny IC$_{50}$ was assessed via a standard 72 hour SYBR Green 1 fluorescence assay

[58]. Cultures were assessed daily for parasitemia and stage. Cultures that were at least 70% ring

were loaded into CQ dose-response assays of a series of 2 fold drug dilutions across 10 wells at

0.15% parasitemia. Drug stocks (1mg/ml) for CQ were prepared in H$_2$O as single-use aliquots

and stored at -20°C until use. Drug dilutions were prepared in incomplete media. Biological

replicates were conducted with at least two cycles of culturing between load dates. IC$_{50}$ values

were calculated in GraphPad Prism 8 using a 4 parameter curve from two technical replicates

loaded per plate.

CRISPR/Cas9 editing at *pfaat1* and parasite phenotyping

CRISPR/Cas9 editing. We designed plasmids for CRISPR/Cas9 editing using the approach

described in [59]. The guide RNA (*GAAATTAAATACATAAAAGA*) was designed to target *pfaat1* in

NHP4026. Edits (258L/313F, 258S/313S and 258S/313F, Fig. 5A) were introduced to NHP4026

through homology arm sequence with target and shield mutations. The parasites were transfected

at ring stages with 100 µg plasmid DNA, and successful transfectants were selected by treatment

with 24 nM WR99210 (gifted by Jacobus Pharmaceuticals, Princeton, NJ) for 6 days. The

parasites were recovered after ~3 weeks. To determine if recovered parasites contained the

expected   mutations,   we   amplified   the   target   region   (forward   primer,

AGTACGGTACTTTTTATATGTACAGCT;   reverse   primer,

TGCATTTGGTTGTTGAGAGAAGG) and confirmed the mutation with Sanger sequencing.

We cloned parasites from successful transfection experiments: independent edited parasites

(from different transfection experiments) were recovered for each *pfaat1* genotype. Edited

parasites were then genome sequenced to identify off-target edits elsewhere in the genome. We

were not able to find any SNP or indel changes between the original NHP4026 and any CRISPR

edited parasites other than the target and shield mutations.

IC_{50} measurement for CRISPR/Cas9 edited parasites. Parasite IC_{50s} for CQ, AMD, LUM, MQ

and QN were measured for 3-5 clones per CRISPR/Cas9 modified line and for NHP4026 across

multiple load dates as described above for cloned progeny, except that each plate included two

NHP4026 technical replicates as controls. This replication of genotype within each load date

allowed for detection of batch effects due to load date.

Batch correction for IC_{50} data. Analysis of Variance (ANOVA) was used to account for batch

effects and to test for differences in IC_{50} between all genotype groups and for each contrast

between each CRISPR/Cas9 modified line and NHP4026 for each drug tested [60]. A linear model

with load date (batch) and genotype as explanatory variables was utilized to generate batch

corrected IC_{50} values for visualization of the impact of CRISPR/Cas9 modifications (Fig 5B and

Supplementary Fig. 12).

Measurement of parasite fitness using competitive growth assays. Parasites were synchronized

to late stage schizonts using a density gradient [61]. The top layer of late stage schizonts was

removed and washed twice with RPMI. Synchronized cultures were suspended in 5 ml of CM at

5% hematocrit and allowed to reinvade overnight with gentle shaking. Parasitemia and parasite

stage were quantified using flow cytometry. Briefly, 80 μl of culture and an RBC control were

stained with SYBR Green I and SYTO 61 and measured on a guava easyCyte HT (Luminex

Co.). 50,000 events were recorded to determine relative parasitemia and stage. When 80% of

parasites were in the ring stage, the head-to-head competition experiments were set-up as

previously described [62]. Competition assays were set up between CRISPR/Cas9 edited parasites

and NHP4026 in a 1:1 ratio at a parasitemia of 1% in a 96-well plate (200 μl per well) and

maintained for 30 days. Each of the assays contained three biological replicates (three

independent clones from different CRISPR/Cas9 editing experiments) and two technical

replicates (two wells of culture). Every two days, the parasitemia was assessed by microscopy

using Giemsa-stained slides, samples were taken and stored at -80°C and the cultures were

diluted to 1% parasitemia with fresh RBCs and media. The proportion of parasite in each

competition (Supplementary Fig. 17) was measured using a rhAmp SNP Assay (IDT, Integrated

DNA Technologies, Inc.) with primers targeting the CRISPR/Cas9 edited region in *pfaat1*.

Selection coefficient. We measured selection coefficient (*s*) by fitting a linear model between the

natural log of the allele ratio (freq [allele edited parasite]/freq [NHP4026]) and time (measured in

48 hour parasite asexual cycles). The slope of the linear model provides a measure of the driving

s of each mutation [63]. To compare relative fitness of parasites carrying different *pfaat1* alleles,

we normalized the fitness of NHP4026 to 1 and used slope + 1 to quantify the fitness of

CRISPR/Cas9 edited parasites (Fig 5C).

<u>Overexpression of PfAAT1 in yeast</u>

To generate pfAAT1 expressing yeast, plasmid carrying the *pfaat1* coding sequence was

transformed into yeast *Saccharomyces cerevisiae* (BY4743) as described in [26]. The doubling

time (h) was measured for strains carry empty vector, wild-type pfAAT1, or S258<u>L</u> mutant

pfAAT1. We measured doubling time under two culture conditions: control or with 5 mM CQ.

Three independent experiments were performed for each assay.

   <u>PfAAT1 protein structure analysis</u>

Three-dimensional homology models for PfAAT1 were predicted using AlphaFold [28,64] and I-

TASSER [29,65,66] and analyzed with PyMol software (v2.3.0; Schrödinger, LLC). At the primary

sequence level, we used TOPCONS [31] to predict transmembrane helix topology for comparison.

We plotted a cartoon version of the protein transmembrane topology based on the

computationally predicted structures and membrane topology (Supplementary Fig. 14). Models

were truncated to exclude amino-terminal residues 1-166, likely positioned outside of the

membrane, because AlphaFold assigns low confidence to this N-terminal stretch. Furthermore,

mutations of interest map only to transmembrane helices according to both 3-D models and

TOPCONS. I-TASSER generated models with topology similar to AlphaFold with the highest

variations in AlphaFold low-confidence regions 1-166 and 475-516. The top five I-TASSER

models superimpose on the AlphaFold model with a root-mean-square-deviation (RMSD) range

of 2.4-2.8 Å over 303-327 of 440 aligned residues using the PDBeFold Server

(http://www.ebi.ac.uk/msd-srv/ssm). The four common SNPs (S258<u>L</u>, F313<u>S</u>, Q454<u>E</u> and

K541<u>L</u>) overlay closely between the homology models. We evaluated the impact of different

mutations on protein stability using the mutagenesis function in PyMol.

<u>Data Availability</u>

All raw sequencing data have been submitted to the NCBI Sequence Read Archive (SRA,

https://www.ncbi.nlm.nih.gov/sra) or European Nucleotide Archive (ENA) with accession

numbers available in supplementary Table 1. All other data are available in the main text or

supplementary materials. The code used in analysis and data analyzed are available at GitHub

through the following links: https://github.com/emilyli0325/CQ.AAT1.git (XL),

782   https://github.com/MPB-mrcg?tab=repositories   (AAN),   and

783   https://github.com/kbuttons/CQ.AAT1.progeny.git (KBS).

References for Methods

Amambua-Ngwa, A. *et al.* Consistent signatures of selection from genomic analysis of pairs of temporal and spatial Plasmodium falciparum populations from The Gambia. *Sci Rep* **8**, 9687, doi:10.1038/s41598-018-28017-5 (2018).

Amambua-Ngwa, A. *et al.* Population genomic scan for candidate signatures of balancing selection to guide antigen characterization in malaria parasites. *PLoS Genet* **8**, e1002992, doi:10.1371/journal.pgen.1002992 (2012).

Manske, M. *et al.* Analysis of Plasmodium falciparum diversity in natural infections by deep sequencing. *Nature* **487**, 375-379, doi:10.1038/nature11174 (2012).

Fariello, M. I., Boitard, S., Naya, H., SanCristobal, M. & Servin, B. Detecting signatures of selection through haplotype differentiation among hierarchically structured populations. *Genetics* **193**, 929-941, doi:10.1534/genetics.112.147231 (2013).

Kumar, S. *et al.* A Malaria Parasite Cross Reveals Genetic Determinants of Plasmodium falciparum Growth in Different Culture Media. *Front Cell Infect Microbiol* **12**, 878496, doi:10.3389/fcimb.2022.878496 (2022).

Button-Simons, K. A. *et al.* The power and promise of genetic mapping from Plasmodium falciparum crosses utilizing human liver-chimeric mice. *Commun Biol* **4**, 734, doi:10.1038/s42003-021-02210-1 (2021).

Li, X. *et al.* Genetic mapping of fitness determinants across the malaria parasite Plasmodium falciparum life cycle. *PLoS Genet* **15**, e1008453, doi:10.1371/journal.pgen.1008453 (2019).

Nair, S. *et al.* Single-cell genomics for dissection of complex malaria infections. *Genome Res* **24**, 1028-1038, doi:10.1101/gr.168286.113 (2014).

Miles, A. *et al.* Indels, structural variation, and recombination drive genomic diversity in Plasmodium falciparum. *Genome Res* **26**, 1288-1299, doi:10.1101/gr.203711.115 (2016).

McDew-White, M. *et al.* Mode and Tempo of Microsatellite Length Change in a Malaria Parasite Mutation Accumulation Experiment. *Genome Biol Evol* **11**, 1971-1985, doi:10.1093/gbe/evz140 (2019).

Davies, L. & Gather, U. The identification of multiple outliers. *Journal of the American Statistical Association* **88**, 782-792 (1993).

Mansfeld, B. N. & Grumet, R. QTLseqr: An R Package for Bulk Segregant Analysis with Next-Generation Sequencing. *Plant Genome* **11**, doi:10.3835/plantgenome2018.01.0006 (2018).

Magwene, P. M., Willis, J. H. & Kelly, J. K. The statistics of bulk segregant analysis using next generation sequencing. *PLoS Comput Biol* **7**, e1002255, doi:10.1371/journal.pcbi.1002255 (2011).

Li, H. A quick method to calculate QTL confidence interval. *J Genet* **90**, 355-360 (2011).

Rason, M. A., Randriantsoa, T., Andrianantenaina, H., Ratsimbasoa, A. & Menard, D. Performance and reliability of the SYBR Green I based assay for the routine monitoring of susceptibility of Plasmodium falciparum clinical isolates. *Trans R Soc Trop Med Hyg* **102**, 346-351, doi:10.1016/j.trstmh.2008.01.021 (2008).

Goswami, D. *et al.* A replication-competent late liver stage-attenuated human malaria parasite. *JCI Insight* **5**, doi:10.1172/jci.insight.135589 (2020).

Taioli, E. *et al.* Application of reliability models to studies of biomarker validation. *Environ Health Perspect* **102**, 306-309, doi:10.1289/ehp.94102306 (1994).

61 Davis, S. Z. *et al.* The extended recovery ring-stage survival assay provides a superior
association with patient clearance half-life and increases throughput. *Malar J* **19**, 54,
doi:10.1186/s12936-020-3139-6 (2020).
62 Tirrell, A. R. *et al.* Pairwise growth competitions identify relative fitness relationships
among artemisinin resistant Plasmodium falciparum field isolates. *Malar J* **18**, 295,
doi:10.1186/s12936-019-2934-4 (2019).
63 Dykhuizen, D. & Hartl, D. L. Selective neutrality of 6PGD allozymes in E. coli and the
effects of genetic background. *Genetics* **96**, 801-817, doi:10.1093/genetics/96.4.801
(1980).
64 Varadi, M. *et al.* AlphaFold Protein Structure Database: massively expanding the
structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids*
Res **50**, D439-D444, doi:10.1093/nar/gkab1061 (2022).
65 Roy, A., Kucukural, A. & Zhang, Y. I-TASSER: a unified platform for automated protein
structure and function prediction. *Nat Protoc* **5**, 725-738, doi:10.1038/nprot.2010.5
(2010).
66 Zhang, Y. I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics* **9**,
40, doi:10.1186/1471-2105-9-40 (2008).
67 Ringwald, P., Bickii, J. & Basco, L. K. In vitro activity of antimalarials against clinical
isolates of Plasmodium falciparum in Yaounde, Cameroon. *Am J Trop Med Hyg* **55**, 254-
258, doi:10.4269/ajtmh.1996.55.254 (1996).
68 Nzila, A., Okombo, J., Ohuma, E. & Al-Thukair, A. Update on the in vivo tolerance and
in vitro reduced susceptibility to the antimalarial lumefantrine. *J Antimicrob Chemother*
67, 2309-2315, doi:10.1093/jac/dks252 (2012).
854

Acknowledgments

We thank past and present staff and visiting researchers at the MRC Unit in The Gambia involved in studies from which samples were archived, including those under the earlier directorships of Prof Brian Greenwood and Prof Tumani Corrah. Work at Texas Biomedical Research Institute was conducted in facilities constructed with support from Research Facilities Improvement Program grant C06 RR013556. SMRU is part of the Mahidol Oxford University Research Unit supported by the Wellcome Trust of Great Britain. We thank the staff of the Wellcome Trust Sanger Institute Sample Logistics, Sequencing and Informatics facilities for their contributions. The Structural Biology Core at The University of Texas Health Science Center at San Antonio is a part of Institutional Research Cores supported by the Office of the Vice President for Research and the Mays Cancer Center Drug Discovery and Structural Biology Shared Resource (National Institutes of Health grant P30 CA054174).

Author Contributions

AAN, KBS, XL, SK, and KVB contributed equally to this work.

Conceptualization: MTF, TA, IC, AV, AAN, DC

Methodology: AV, SK, SVA, ABT

Investigation: AAN, UDA, DK, DC, RA, RDP, KBS, KVB, LC, HD, JKL, AR, ED, SK, DS, ST, MF, ABT, TA

Analysis: XL, KBS, AAN

Visualization: XL, KBS, AAN

Funding acquisition : MTF, TA, DC, AAN, DK, SVA, FN

Project administration: MF, DK, AAN, DC, SVA

Supervision: MF, AV, TA, IC, SHIK, SVA, DK, DC

Writing – original draft: KVB, TA, KBS, XL

Writing – review & editing: MTF, AV, IC, SK, AAN, DC, DK, UDA, RDP, SHIK, FN, ABT,

TA

Competing Interests

Authors declare that they have no competing interests.

Additional Information

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to Timothy J. C. Anderson

(tanderso@txbiomed.org).

Reprints and permission information is available at www.nature.com/reprints.

Funding:

National Institutes of Health grant P01 12072248 (MTF)

National Institutes of Health grant R37 AI048071 (TA)

MRC Career Development Fellowship MC_EX_MR/K02440X/1 (AAN)

MRC malaria program grant MC_EX_MR/J002364/1 (UDA)

European Research Council grant AdG-2011-294428 (DC and AAN)

MRC grant MR/S009760/1 (DC)

Wellcome Trust (206194, 204911) and the Bill & Melinda Gates Foundation

(OPP1204628, INV-001927) (DK)

Biotechnology and Biological Sciences Research Council grant BB/M022161/1 (SVA)

Wellcome Trust [Grant number 220221] (FN); For the purpose of open access, the author

has applied a CC BY public copyright license to any Author Accepted Manuscript

version arising from this submission.

Supplementary Figures

Figure S1

Fig. S1. Estimation of selection coefficient (*s*) for *pfaat1 (S258L̲)* and *pfcrt (K76T̲)* alleles. *p*

is the frequency of mutant alleles (*pfaat1* S258L̲ or *pfcrt* K76T̲) as indicated in Fig. 1A, and q

(=1-p) is the inferred frequency of wild-type (3D7) alleles. The x-axis indicates parasite

generations (labeled with sample collection year). We estimated selection coefficients (*s*) based

on allele frequency from year 1990 and 2001, as CQ monotherapy was stopped in Gambia in

2004. *s* indicates the changes in relative growth per parasite generation (i.e. the duration of the

complete lifecycle in both mosquito and human host). The calculation was based on estimates of

2, 4, or 6 generations per year.

Figure S2

 Fig. S2. Haplotype structure at the *pfcrt* (left panel) and *pfaat1* (right panel) regions. Each

 point depicts an isolate with point colors representing the years from which they were sampled.

 Square points represent complex infections and circles represent monoclonals. MOI, multiplicity

 of infection.

Figure S3

Fig. S3. The proportion of pairs identical by descent (IBD) within populations from global

locations. For samples with >90% of the genomes are IBD, only one representative sample with

the highest genotype rate was selected and used for IBD analysis. Sample numbers are showed in

each panel. Chromosome boundaries are indicated with grey dashed vertical lines. The location

of *pfaat1* and *pfcrt* are indicated with red arrows on top of each panel. See also analysis by

Amambua-Ngwa et al [17], Hendon et al [18], and Carrasquilla et al [19].

Figure S4

Fig. S4. Interchromosomal linkage disequilibrium (LD) analysis. A. R^2 between *pfaat1* (SNP pfaat258 located at Pf3D7_06_v3:1215233) and a 200 kb region around the *pfcrt* gene on chr. 7. B. R^2 between *pfcrt* (SNP pfcrt76 located at Pf3D7_07_v3:403625) and a 200kb region around the *pfaat1* gene on chr. 6. R^2 values against SNPs within gene *pfcrt* (in panel A) or *pfaat1* (in panel B) are shown as green or red points, while the red points are LDs between SNPs pfaat258 and pfcrt76. For both panel A and B, from top to bottom, rows represent LD analysis for populations from year 2001, 2008, and 2014, respectively. Samples from 1984 are not shown, because neither *pfaat1* S258<u>L</u> nor *pfcrt* K76<u>T</u> were present. 1990 was omitted because only 13 samples were available. The highest LDs were observed between gene *pfaat1* and *pfcrt*.

949   **Figure S5**

Fig. S5. Extended haplotype homozygosity (EHH) analysis of Gambian samples. Graphs show

EHH surrounding *pfaat1* S258L̲ (left panels), or *pfcrt* K76T̲ (right panels) in samples collected in

1984, 2001, 2008 and 2014. 1990 was omitted because only 13 samples were available. Samples

from 1984 show EHH around the ancestral allele only (blue), because neither *pfaat1* S258L̲ nor

pfcrt K76T̲ were sampled at that time.

Figure S6

Fig. S6. *pfcrt* and *pfaat1* allele frequency distributions and correlations in African countries. A.

pfcrt allele distribution in African countries. B. *pfaat1* allele distribution in African countries. C.

Correlations in allele frequencies between *pfcrt* (CVIET) and *pfaat1* (S258$\underline{L}$). Frequencies of the

CVIET haplotype for amino acids 72-76 in *pfcrt* are significantly correlated with allele

frequencies of *pfaat1* S258$\underline{L}$ in West Africa ($R^2 = 0.65$, $p = 0.0017$, red dashed line) or across all

African populations ($R^2 = 0.44$, $p = 0.0021$). Point size indicates sample numbers, while color

indicates sampling locations.

Figure S7

Fig. S7. UPGMA tree showing the relationship of 581 haplotypes based on SNPs inside the

50 kb region surrounding *pfaat1***.** The tree was rooted with *Plasmodium reichenowi* (not shown

in the tree). WAF: west Africa, EAF: east Africa, CAF: central Africa, SM: south America,

ESEA: east Southeast (SE) Asia, SA: south Asia, WSEA: west SE Asia, PO: Pacific Ocean.

Figure S8

Fig. S8. Mapping CQ resistance loci with cross 3D7×NHP4026 using recombinant progeny

pool 1. Grey, orange, and black lines indicate BSAs from 50 nM, 100 nM, and 250 nM CQ

treatments, separately. Lines with the same color are BSAs from technical replicates. Red dashed

lines are the threshold ($G\ prime$ = 20) for QTL detection. Red arrows indicate the location of the

chromosome 6 QTL.

Figure S9

Fig. S9. Mapping CQ resistance loci with cross 3D7×NHP4026 using recombinant progeny

pool 2. Grey, orange, and black lines indicate BSAs from 50 nM, 100 nM, and 250 nM CQ

treatments, separately. Lines with the same color are BSAs from technical replicates. Red dashed

lines are the threshold (*G prime* = 20) for QTL detection. Red arrows indicate the location of the

chromosome 6 QTL.

Figure S10

Fig. S10. 3D7 allele frequency for QTLs at chr.6, chr.7 and chr.14. CQ treatments were

applied to the pools on day 0 at 0 (control), 50, 100, or 250 nM. CQ was removed on day 2 (48

hour treatment) or day 4 (96 hour treatment), as shaded with light blue. For 48 hour CQ

treatments, samples were collected at day 0, 4, and 7; while for 96 hour CQ treatment, samples

were collected at day 0, 5, and 10. Solid or dashed lines are results from different technical

996   replicates. The 3D7 allele frequencies in CQ treated pools decrease at both chr.6 and chr.7 QTL

997   regions. At the chr.14 QTL region, allele frequencies are unchanged following drug treatment,

998   suggesting this QTL is unrelated to drug treatment.

999 **Figure S11**

1000

1001 **Fig. S11**. Mean IC$_{50}$ (± S.D.) of parents and progeny grouped by combinations of *pfcrt* and

1002 *pfaat1* allele. IC$_{50}$ for each clone is calculated from 4-13 biological replicates (Supplementary

1003 Table 7). Only two progeny carrying *pfaat1* 258S/313F (WT) and *pfcrt* K76T̲ were recovered. *P*

1004 values indicate significance levels are based on two-way ANOVA analysis.

1005

1006   **Figure S12**

1007

Fig. S12. Impact of CRISPR/Cas9 substitutions on IC_{50} of quinine, lumefantrine,

mefloquine and amodiaquine. CRISPR/Cas9 gene editing resulted in small differences in IC_{50}

for quinine (QN), lumefantrine (LUM) and Amodiaquine (AMD), but no significant changes for

mefloquine (MQ). However, all IC_{50}'s were below levels of clinical significance for these drugs

(clinical thresholds: QN = 600 nM [67]; MQ = 30 nM [67], AMD = 60 nM [67]), or at the lower end of

the *in vitro* range (0-150 nM) in the case of LUM [68]. *P* values indicate significance levels are

based on two-way ANOVA analysis.

1015

1016 **Figure S13**

1017

Fig. S13. Introduction of S258L results in chloroquine resistance in yeast. Yeast doubling

time was calculated from the linear portion of exponential growth. Data was shown as means

from 3 independent experiments ± SEM, and significance and was calculated according to

multiple comparisons (with Turkey corrections) of two-way ANOVA. Growth of yeast cells

expressing wild type *pfaat1* (WT) is severely impacted by CQ treatment (5 mM CQ through the

experiments) but is recovered in yeast expressing *pfaat1* S258L. Published results demonstrate

that AAT1 is expressed in the yeast cell membrane [26].

Figure S14

1026

1027   **Fig. S14. Topology structure of pfAAT1 protein.** (A) AlphaFold model of PfAAT1 (left),

1028   representative I-TASSER model of PfAAT1 (center), structural superposition of the AlphaFold

1029   model (teal) and I-TASSER model (gray, right). TOPCONS transmembrane (TM) helix topology

1030   predictions are mapped onto the models in dark blue (left, center). AlphaFold and I-TASSER

models align with a RMSD of 2.5 Å over 327 of 440 residues. Amino-terminal residues 1-166 were excluded from all models due to low confidence in structure prediction. (B) Detailed view of the mutations on the predicted PfAAT1 3-D structure using the AlphaFold model. The right view is related to the left by a 45° rotation about the axis looking down at the figure followed by a 90° rotation about the horizontal axis. The four SNPs shown as space-filling models are all arranged within a plane at one side of the model, perpendicular to the membrane. S258L (helix 3) and F313S (helix 5) are located opposite each other with helix 8 in between. Given the epistatic interactions between the PfAAT1 S258L and F313S SNPs evident from our functional analyses, the F313S substitution of the bulky, hydrophobic phenylalanine with the smaller, polar serine may compensate for a disruption in the transmembrane region that includes helices 3, 5, and 8 allowing for partial restoration of amino acid transport activity. Q454E is located on helix 8 near the TM surface and K541N is located in a loop connecting helix 9 and 10. (C) Topology of PfAAT1 inferred using 3D structure. There are eleven transmembrane (TM) helices. Three of the mutations are located at the TM helices, while K541N is located at a loop connecting helix 9 and 10. The color scheme matches the schematic in Panel B. The blue triangle indicates amino-terminal residues 1-166 that were excluded from structure prediction.

1048 **Figure S15**

1049

1050 **Fig. S15.** Project design. We use (i) population genomic analyses, (ii) genetic crosses and

1051 quantitative genetics analysis followed by (iii) functional analyses to investigate the role of

1052 additional loci in CQ resistance.

1053

1054   **Figure S16**

1055

1056   **Fig. S16.** Experiment design for CQ bulk segregant analysis.

 Figure S17

Fig. S17. Allele frequency changes in head-to-head competition experiments between different
CRISPR/Cas9 edited parasites and NHP4026. We used three independent CRISPR/Cas9 edited
clones for each genotype, and two technical replicates for each competition experiment (average
values plotted).