1 # *Pseudomonas aeruginosa* contracts mucus to rapidly
2 # form biofilms in tissue-engineered human airways
3

Authors:

Tamara Rossy[1,2], Tania Distler[1,2], Joern Pezoldt[2], Jaemin Kim[3], Lorenzo Talà[1,2], Nikolaos
Bouklas[3], Bart Deplancke[2], Alexandre Persat[1,2]

Affiliations:

[1] Global Health Institute, School of Life Sciences, École Polytechnique Fédérale de Lausanne
(EPFL), Lausanne, Switzerland

[2] Institute of Bioengineering, School of Life Sciences, École Polytechnique Fédérale de
Lausanne (EPFL), Lausanne, Switzerland

[3] Sibley School of Mechanical and Aerospace Engineering, Cornell University, Ithaca, NY, USA

Correspondence: alexandre.persat@epfl.ch

Abstract

Bacteria commonly protect themselves by forming multicellular structures called biofilms. The opportunistic pathogen *Pseudomonas aeruginosa* causes antibiotic-recalcitrant pneumonia by forming biofilms in the respiratory tract. Despite extensive *in vitro* experimentation, how *P. aeruginosa* forms biofilms at the airway mucosal surface is unresolved. We investigated *P. aeruginosa* biofilm biogenesis in optically-accessible tissue-engineered human lung models that emulate the airway mucosal environment. We found that *P. aeruginosa* forms mucus-associated biofilms within hours, much faster than previously observed in *in vitro* experiments. Early during colonization, *P. aeruginosa* induces contractions of luminal mucus which accelerates bacterial aggregation. We show that *P. aeruginosa* uses retractile type IV pili to actively compress mucus. Our results suggest that, while protecting epithelia, mucus constitutes a breeding ground for biofilms.

Main text

Bacteria predominantly colonize their environments in the form of biofilms, dense communities

of contiguous cells embedded in a self-secreted polymeric matrix[1]. The mechanisms of biofilm

formation have been extensively studied on abiotic surfaces and in laboratory conditions[2,3]. In

contrast, our understanding of biofilm formation and physiology in a realistic context of human

infections is limited[4,5]. Biofilms from the pathogen *P. aeruginosa* epitomize this disparity. *P.*

aeruginosa causes acute and chronic pneumoniae in immunocompromised individuals by

forming airway-associated biofilms[6]. The architecture of *P. aeruginosa* biofilms sampled from

patient sputum and lung explants is quite different from *in vitro* ones[5]. This divergence indicates

that airway mucosal factors that contribute to *P. aeruginosa* biofilm formation *in vivo* are

omitted from laboratory investigations.

Epithelial tissues secrete a hydrogel substance called mucus, the first line of defense of the

airway against all respiratory pathogens. Mucus is composed of gel-forming mucin

glycoproteins, which cross-link into a viscoelastic substance upon exocytosis. It is commonly

assumed that mucus is a passive physical barrier. However, bacterial pathogens and

commensals physiologically adapt to mucins, suggesting mucus plays a regulatory role in host-

microbe interactions. For example, glycans that decorate mucins regulate *P. aeruginosa*

physiology, including virulence and adhesion[7]. Less is known about the regulatory role of

mucus in its hydrogel form[8]. Individuals with underlying respiratory conditions such as chronic

obstructive pulmonary disease (COPD) and cystic fibrosis (CF) have aberrant mucus. At the

same time, they are at risk of specifically developing chronic *P. aeruginosa* pneumonia[9].

Despite this common association, how mucus mechanics contribute to the onset and

persistence of *P. aeruginosa* infection remains unresolved. Mucus in its hydrogel form is fragile

as it loses integrity upon mechanical and chemical perturbations[10,11]. As a result, mucus is

largely omitted from investigations of *P. aeruginosa* biofilm formation. Consequently, we still

ignore how *P. aeruginosa* interacts with native mucus to form biofilms during infection.

To bridge the gap between *in vitro* biofilm studies and clinical observations, we emulated

the mucosal environment of the airway in the lab. We used a tissue-engineering approach to

develop AirGels (<u>air</u>way in <u>gels</u>): human lung epithelial tissue in a molded collagen/Matrigel

extracellular matrix (ECM) scaffold[12,13]. AirGels grow from primary human bronchial epithelial

(HBE) cells of healthy and CF donors, which expand to confluence on the cylindrical cavity of

the ECM scaffold (Figure 1A). An elastomeric microfluidic chip maintains AirGels and allows

for luminal access while preserving epithelial integrity. The ECM geometry guides epithelial

architecture, enabling morphological customization of AirGels. Here, we designed and

optimized AirGels to enable high resolution fluorescence microcopy to monitor infection

dynamics at the single bacterium level in live tissue. Maintaining an air-liquid interface in the
lumen promotes epithelial cell differentiation and reproduces the physiological conditions
encountered in the airway. We therefore optimized the matrix formulation so that AirGels
remain stable at the air-liquid interface, thereby biologically and physically replicating the
airway environment (Figure 1B).

Single-plane illumination microscopy images show that mature AirGels form tubular
epithelial tissue, recapitulating the architecture and dimensions of a human small bronchus
(Figure 1C)[14,15]. AirGel epithelia are tight and impermeable (Supplementary Figure 1). We
characterized HBE cell differentiation in 34-day-old AirGels. Immunofluorescence highlighted
an abundant population of mucus-producing goblet cells and ciliated cells (Figure 1C). To
quantify the abundance of each cell type, we performed single-cell RNA sequencing (scRNA-
seq) of mature AirGels. We identified five main clusters (Figure 1D and Supplementary Figure
2): basal cells (8%), ciliated cells (41%), secretory cells (34%), as well as immature ciliated
(7%) and immature secretory cells (10%). AirGels therefore reproduce the cellular composition
and histological signature of human airway epithelia[16–19].

Given its prominent function in host-microbe interactions, we carefully characterized the
architecture of mucus in AirGels. Immunofluorescence against the airway gel-forming mucins
MUC5AC and MUC5B showed the presence of extracellular mucus in the form of thick luminal
filaments (Figure 1E). We also observed similar fiber-like mucus architecture in live AirGels by
staining with the fluorescently-labeled lectin jacalin[20]. These fibers recapitulate the mucus
architecture observed in porcine and murine tracheal explants[20–22]. We then characterized
AirGel mucociliary clearance functions. Measurements of cilia beating frequency in AirGels
were indistinguishable from previous *ex vivo* measurements (Supplementary Figure 3 &
Supplementary Video 1)[23–26]. In addition, AirGel cilia generated a directional flow whose
clearance velocity matched the physiological range (Figure 1F)[22,24,27]. Overall, AirGels
reproduce biological, physical and dynamic parameters of the human airway including its tube-
shape, all in a system allowing for live imaging of host-pathogen interactions at high resolution.

To visualize biofilm formation in a realistic airway mucosal context, we inoculated *P.
aeruginosa* constitutively expressing the fluorescent protein mScarlet in the lumen of AirGels
maintained at the air-liquid interface. After 13h of incubation, we observed that bacteria had
extensively colonized the mucosal surface. *P. aeruginosa* formed interconnected bacterial
clusters colocalized with mucus within the airway surface liquid (ASL) between epithelial cells
and the air-liquid interface (Figure 1G). In dynamic visualizations, bacteria remained attached
to mucus despite mucus movements induced by beating cilia (Supplementary Video 2). Since
P. aeruginosa takes days to form biofilms *in vitro*, we were surprised to see these communities

form only within a few hours in AirGels[28]. We therefore went on to investigate the mechanisms
of rapid biofilm formation on mucus.

We imaged biofilm biogenesis in AirGels at the single cell level using confocal spinning disk
microscopy. *P. aeruginosa* already formed aggregates a few hours after inoculation (Figure
2A). While the mucus surface was initially largely devoid of bacteria, half of it was covered by
P. aeruginosa multicellular structures after 5.5h of infection (Figure 2B). Bacterial clusters with
the same architecture also formed in the absence of jacalin staining, confirming these biofilms
do not form through labeling artefacts (Supplementary Figure 4B). Moreover, *P. aeruginosa*
rapidly formed biofilms on mucus of diseased AirGels engineered from primary cells of CF
donors (Supplementary Figure 5). To confirm the pivotal role of mucus in rapid biofilm
formation, we infected a non-differentiated AirGel which does not produce mucus. In the
absence of a protective mucus layer, epithelial cells were more vulnerable to *P. aeruginosa*
infection (Supplementary Figure 6). Bacteria breached through the epithelial barrier and
invaded the underlying ECM. *P. aeruginosa* did not form three-dimensional multicellular
structures in the ASL. This further demonstrates the role of mucus hydrogel as a substrate for
biofilm formation in differentiated AirGels, and at the same time highlights its protective
function.

Our data suggests that *P. aeruginosa* forms biofilms in the airway by attaching to mucus at
early stages of infection. To further explore the biophysical mechanisms of biofilm formation,
we harvested mucus to perform *ex situ* visualizations. However, we could not observe the
formation of *P. aeruginosa* biofilms on mucus extracted from AirGels (Supplementary Figure 7
& Supplementary Video 3). We attribute this discrepancy to perturbations in mucus mechanical
integrity when extracted from the epithelium. This difference highlights the importance of
investigating microbe-mucus interactions in a native mucosal context such as the one
established in AirGels.

To understand how biofilms rapidly form on native mucus, we therefore inspected the
different steps of their formation in AirGels. To nucleate *in vitro* biofilms, *P. aeruginosa* cells
navigate the surface of abiotic materials using twitching motility, which promotes the formation
of aggregates[29]. Fast imaging of single cells shows that *P. aeruginosa* moves with twitching-
like trajectories at the surface of mucus fibers (Supplementary Video 4). As expected from
axenic experiments, these single cells aggregate into small clusters within 2h of colonization
(Figure 2A). In contrast with axenic conditions, these small multicellular clusters subsequently
moved along mucus fibers to eventually fuse (Figure 2C). This caused a cascade of cluster
fusion events that sped up biofilm formation (Figure 2C & Supplementary Video 5). We tracked
aggregate fusion in kymographs highlighting the correlation between mucus and bacterial

displacements (Figure 2D). The size of individual clusters remains approximately constant
during motion and fusion, showing aggregate fusion predominates over bacterial growth. After
only 6h of aggregation and fusion, dense biofilms are formed.

We found that during biofilm formation, the mucus surface area tends to decrease
compared to an uninfected control (Figure 3A & Supplementary Figure 8). We therefore
hypothesized that mucus gel remodeling could drive *P. aeruginosa* aggregation and aggregate
fusion, thereby speeding up biofilm formation. We envisioned two mechanisms for bacteria-
induced mucus deformations: degradation or direct mechanical contraction. *P. aeruginosa*
secretes mucinases capable of degrading gel-forming mucins[30]. Enzymatic mucus degradation
could release polymers that generate entropic depletion forces promoting bacterial
aggregation or that generate osmotic forces compressing mucus[31,32]. To test whether mucus
degradation could drive contraction, we colonized AirGels with a mutant in the type II secretion
system locus *xcp* that is necessary for mucin utilization[30,33]. The *Δxcp* mutant however formed
biofilms similar to WT, ruling out the hypothesis of polymer-induced forces driving the formation
of multicellular structures (Figure 3B).

Could *P. aeruginosa* remodel mucus by directly and actively applying force on the surface?
P. aeruginosa can generate extracellular forces using flagella and type IV pili (T4P), motorized
filaments that also play a role during *in vitro* biofilm biogenesis. In addition, T4P and flagella
mediate single cell interactions with mucins[34–38]. To investigate their functions in the context of
rapid biofilm formation on mucus, we infected AirGels with *P. aeruginosa* mutants lacking
flagella (*ΔfliC*) and T4P (*ΔpilA*). The *ΔfliC* mutant formed biofilms that were indistinguishable
from wild type (WT) (Figure 3B). By contrast, *ΔpilA* cells did not form multicellular structures,
indicating T4P play a role in rapid biofilm formation. Since T4P may bind to glycans present on
mucins[37,38], weaker cell attachment to mucus could cause a decrease in aggregation of *ΔpilA*.
Yet, colocalization analysis shows that the *ΔpilA* mutant is still able to attach efficiently to
mucus (Supplementary Figure 9 & Supplementary Video 6). We therefore envisioned a
mechanism where T4P generate retractile forces that contract mucus from the surface,
ultimately speeding up *P. aeruginosa* aggregation and cluster fusion.

To physically explore this scenario, we ran non-linear finite element simulations wherein
mucus is treated as a hyperelastic material[39]. The mechanical action of *P. aeruginosa* T4P at
the mucus surface is considered through the introduction of an active surface stress. The
simulations recapitulated the experimental observations of mucus hydrogel contraction during
P. aeruginosa colonization (Figure 3C). Simulations also predict that the steady-state mucus
area decreases with the magnitude of the surface contractile modulus. This suggests that the
more T4P retract, the more *P. aeruginosa* compresses mucus. To experimentally validate this

model, we visualized AirGels colonization by a *ΔpilT* mutant which produces T4P that cannot
retract, mimicking conditions of zero contractile modulus. *P. aeruginosa ΔpilT* could still
associate with mucus and form a few small clusters, but clearly failed to form mucus-
associated biofilms to the same extent as WT (Figures 3D & E), which was coherent with
simulations. These results show that T4P retraction is necessary for rapid biofilm formation,
and is consistent with a scenario where retraction compresses the mucus substrate.

To further support the surface contraction model, we tested the prediction that deformations
increase further with surface contractility. We imaged AirGel colonization by the hyperpiliated
P. aeruginosa mutant *ΔpilH,* whose T4P retraction frequency is approximately twice the one
of WT (Supplementary Figure 10). *ΔpilH* formed biofilms more rapidly than WT: we observed
dense aggregates as early as 2h, while we only did after 4h for WT (Figure 3F & G). In addition,
ΔpilH induced more rapid mucus contraction than WT (Figure 3H & Supplementary Videos 7
& 8), consistent with simulations. After 5.5h, WT and *ΔpilH* biofilms had similar morphologies
and size, suggesting biofilm fusion reaches a physical limit most likely due to packing at the
mucus surface (Figure 3F and G). Overall, our results support a model where *P. aeruginosa*
contracts the surface of mucus by active T4P retraction. Single cells initially twitch on mucus
to form small aggregates. The static aggregate collective generate force from T4P that are
sufficient to deform their substrate, driving large-scale mucus contraction. By contracting,
mucus brings aggregates closer. They eventually fuse and form biofilms (Supplementary
Figure 11).

Active mucus contraction by *P. aeruginosa* speeds up biofilm formation compared to axenic
conditions. Although the classical view of airway infections associates biofilms with chronic
infections and planktonic cells with acute infections, recent work has demonstrated the
coexistence of these bacterial lifestyles in sputum samples from both acutely and chronically
infected patients[40]. Our observations of rapid biofilm formation in both healthy and CF AirGels
is consistent with these clinical observations.

Rapid biofilm formation may provide a fitness advantage in the non-hospitable airway
environment. Forming biofilms rapidly could reduce *P. aeruginosa*'s susceptibility to
neutrophils and macrophages, which are rapidly recruited to the mucosal surface during
infection[41,42]. At the same time, forming biofilms increases *P. aeruginosa* tolerance to
subsequent antibiotic treatment and promotes the emergence of resistant mutants[43]. There is
however an upside: mucus adsorbs a large proportion of the planktonic *P. aeruginosa*, thereby
shielding the epithelium from acute virulence. Our results therefore demonstrate the duality of
mucus: protecting the airway epithelium from acute infections, while providing a fertile ground

for biofilm formation and chronic infections. Altogether, our study emphasizes the importance

of investigating bacterial physiology in more realistic infection contexts.

Acknowledgements

We thank Zaïnebe Al-Mayyah, Jeremy Wong, the Bioimaging and Optics Core Facility (BIOP) and the Gene Expression Core Facility (GECF) at EPFL for technical assistance. Marco Kühn, Johannes Bues and Sophia Hsin-Jung Li for insightful discussions, Romé Voulhoux for strains and Formlabs forum user Telliria for suggestions on 3D-printing process. We acknowledge the Swiss National Science Foundation for funding this project through the Project grant number 310030_189084 and NCCR AntiResist.

References

1. Flemming, H.-C. *et al.* Biofilms: an emergent form of bacterial life. *Nature Reviews Microbiology* **14**, 563–575 (2016).

2. Teschler, J. K. *et al.* Living in the matrix: assembly and control of Vibrio cholerae biofilms. *Nat Rev Microbiol* **13**, 255–268 (2015).

3. Yan, J. & Bassler, B. L. Surviving as a Community: Antibiotic Tolerance and Persistence in Bacterial Biofilms. *Cell Host & Microbe* **26**, 15–21 (2019).

4. Costerton, J. W., Stewart, P. S. & Greenberg, E. P. Bacterial biofilms: a common cause of persistent infections. *Science* **284**, 1318–1322 (1999).

5. Bjarnsholt, T. *et al.* The in vivo biofilm. *Trends in Microbiology* **21**, 466–474 (2013).

6. Rossi, E. *et al.* Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis. *Nat Rev Microbiol* **19**, 331–342 (2021).

7. Wheeler, K. M. *et al.* Mucin glycans attenuate the virulence of Pseudomonas aeruginosa in infection. *Nature Microbiology* **4**, 2146–2154 (2019).

8. Matsui, H. *et al.* A physical linkage between cystic fibrosis airway surface dehydration and Pseudomonas aeruginosa biofilms. *Proceedings of the National Academy of Sciences* **103**, 18131–18136 (2006).

9. Folkesson, A. *et al.* Adaptation of Pseudomonas aeruginosa to the cystic fibrosis airway: an evolutionary perspective. *Nature Reviews Microbiology* **10**, 841–851 (2012).

10.   Wagner, C. E., Wheeler, K. M. & Ribbeck, K. Mucins and Their Role in Shaping the

Functions of Mucus Barriers. *Annual Review of Cell and Developmental Biology* **34**, 189–

215 (2018).

11.   Perez-Vilar, J. & Mabolo, R. Gel-forming mucins. Notions from in vitro studies. *Histol.*

Histopathol. **22**, 455–464 (2007).

12.   Jiménez-Torres, J. A., Peery, S. L., Sung, K. E. & Beebe, D. J. LumeNEXT: A Practical

Method to Pattern Luminal Structures in ECM Gels. *Advanced Healthcare Materials* **5**, 198–

204 (2016).

13.   Barkal, L. J. *et al.* Microbial volatile communication in human organotypic lung models.

Nature Communications **8**, 1770 (2017).

14.   Chen, Y. *et al.* Validation of human small airway measurements using endobronchial

optical coherence tomography. *Respiratory Medicine* **109**, 1446–1453 (2015).

15.   Horsfield, K. & Cumming, G. Morphology of the bronchial tree in man. *Journal of*

Applied Physiology **24**, 373–383 (1968).

16.   Dvorak, A., Tilley, A. E., Shaykhiev, R., Wang, R. & Crystal, R. G. Do Airway Epithelium

Air–Liquid Cultures Represent the In Vivo Airway Epithelium Transcriptome? *Am J Respir*

Cell Mol Biol **44**, 465–473 (2011).

17.   Montoro, D. T. *et al.* A revised airway epithelial hierarchy includes CFTR-expressing

ionocytes. *Nature* **560**, 319–324 (2018).

18.   Plasschaert, L. W. *et al.* A single-cell atlas of the airway epithelium reveals the CFTR-

rich pulmonary ionocyte. *Nature* **560**, 377–381 (2018).

19.   Travaglini, K. J. *et al.* A molecular cell atlas of the human lung from single-cell RNA

sequencing. *Nature* **587**, 619–625 (2020).

20.   Ostedgaard, L. S. *et al.* Gel-forming mucins form distinct morphologic structures in

airways. *PNAS* **114**, 6842–6847 (2017).

21.   Hoegger, M. J. *et al.* Impaired mucus detachment disrupts mucociliary transport in a

piglet model of cystic fibrosis. *Science* **345**, 818–822 (2014).

22.   Fakih, D. *et al.* Normal murine respiratory tract has its mucus concentrated in clouds

based on the Muc5b mucin. *American Journal of Physiology-Lung Cellular and Molecular*

Physiology **318**, L1270–L1279 (2020).

23.   Hoegger, M. J. *et al.* Assessing mucociliary transport of single particles in vivo shows

variable speed and preference for the ventral trachea in newborn pigs. *PNAS* **111**, 2355–

2360 (2014).

24.   Fahy, J. V. & Dickey, B. F. Airway mucus function and dysfunction. *New England*

Journal of Medicine **363**, 2233–2247 (2010).

25.   Bermbach, S. *et al.* Mechanisms of Cilia-Driven Transport in the Airways in the

Absence of Mucus. *Am J Respir Cell Mol Biol* **51**, 56–67 (2014).

26.   Rayner, R. E., Makena, P., Prasad, G. L. & Cormet-Boyaka, E. Optimization of Normal

Human Bronchial Epithelial (NHBE) Cell 3D Cultures for in vitro Lung Model Studies.

Scientific Reports **9**, 500 (2019).

27.   Kudo, E. *et al.* Low ambient humidity impairs barrier function and innate resistance

against influenza infection. *PNAS* **116**, 10905–10910 (2019).

28.   Ma, L. *et al.* Assembly and Development of the Pseudomonas aeruginosa Biofilm

Matrix. *PLOS Pathogens* **5**, e1000354 (2009).

29.   Zhao, K. *et al.* Psl trails guide exploration and microcolony formation in Pseudomonas

aeruginosa biofilms. *Nature* **497**, 388–391 (2013).

30.   Alrahman, M. A. & Yoon, S. S. Identification of essential genes of Pseudomonas

aeruginosa for its growth in airway mucus. *J Microbiol.* **55**, 68–74 (2017).

31.   Datta, S. S., Preska Steinberg, A. & Ismagilov, R. F. Polymers in the gut compress the

colonic mucus hydrogel. *Proceedings of the National Academy of Sciences* **113**, 7041–7046

(2016).

32.   Secor, P. R., Michaels, L. A., Ratjen, A., Jennings, L. K. & Singh, P. K. Entropically

driven aggregation of bacteria by host polymers promotes antibiotic tolerance in

Pseudomonas aeruginosa. *Proceedings of the National Academy of Sciences* **115**, 10780–

10785 (2018).

33.   Chapon-Hervé, V. *et al.* Regulation of the *xcp* secretion pathway by multiple quorum-

285   sensing modulons in *Pseudomonas aeruginosa*. *Molecular Microbiology* **24**, 1169–1178

286   (1997).

287   34.   Co, J. Y. *et al.* Mucins trigger dispersal of Pseudomonas aeruginosa biofilms. *NPJ*

288   *Biofilms Microbiomes* **4**, (2018).

289   35.   Arora, S. K., Ritchings, B. W., Almira, E. C., Lory, S. & Ramphal, R. The Pseudomonas

290   aeruginosa Flagellar Cap Protein, FliD, Is Responsible for Mucin Adhesion. *Infect Immun*

291   **66**, 1000–1007 (1998).

292   36.   Landry, R. M., An, D., Hupp, J. T., Singh, P. K. & Parsek, M. R. Mucin–Pseudomonas

293   aeruginosa interactions promote biofilm formation and antibiotic resistance. *Molecular*

294   *Microbiology* **59**, 142–151 (2006).

295   37.   Comolli, J. C., Waite, L. L., Mostov, K. E. & Engel, J. N. Pili Binding to Asialo-GM1 on

296   Epithelial Cells Can Mediate Cytotoxicity or Bacterial Internalization by Pseudomonas

297   aeruginosa. *Infect Immun* **67**, 3207–3214 (1999).

298   38.   Bucior, I., Pielage, J. F. & Engel, J. N. Pseudomonas aeruginosa Pili and Flagella

299   Mediate Distinct Binding and Signaling Events at the Apical and Basolateral Surface of

300   Airway Epithelium. *PLOS Pathogens* **8**, e1002616 (2012).

301   39.   Kim, J., Mailand, E., Ang, I., Sakar, M. S. & Bouklas, N. A model for 3D deformation

302   and reconstruction of contractile microtissues. *Soft Matter* **17**, 10198–10209 (2021).

303   40.   Kolpen, M. *et al.* Bacterial biofilms predominate in both acute and chronic human lung

304   infections. *Thorax* (2022) doi:10.1136/thoraxjnl-2021-217576.

305   41.   Montefusco-Pereira, C. V. *et al.* P. aeruginosa Infected 3D Co-Culture of Bronchial

306   Epithelial Cells and Macrophages at Air-Liquid Interface for Preclinical Evaluation of Anti-

307   Infectives. *JoVE (Journal of Visualized Experiments)* e61069 (2020) doi:10.3791/61069.

308   42.   Yonker, L. M. *et al.* Development of a Primary Human Co-Culture Model of Inflamed

309   Airway Mucosa. *Sci Rep* **7**, 8182 (2017).

310   43.   Ciofu, O., Moser, C., Jensen, P. Ø. & Høiby, N. Tolerance and resistance of microbial

311   biofilms. *Nat Rev Microbiol* 1–15 (2022) doi:10.1038/s41579-022-00682-4.

Figures

Figure 1: A tissue-engineered airway as a novel infection model. A. Schematic of an
AirGel chip. **B.** Brightfield image of an AirGel on the day of HBE cell were seeding (left) and at
the air-liquid interface after 30 days in culture (right). **C.** Longitudinal cross-sectional images
of immunostained differentiated AirGel chips. Confocal images show the gel-forming mucin
MUC5AC (green) and acetylated α-tubulin labeling cilia (orange) along with the actin dye

phalloidin (pink) and nuclear dye DAPI (blue). The transverse cross section 3D image was
reconstituted from SPIM data for actin fluorescence. The bottom right panel is a maximal
intensity projection of a z-stack acquired in the curved lumen. **D.** scRNA-seq identifies cell type
diversity of AirGels. Uniform Manifold Approximation and Projection (UMAP) embedding of
cells pooled from three differentiated AirGels (35-days old), subjected to scRNA-seq profiling.
E. Extracellular luminal mucus in AirGels. Immunofluorescence of methacarn-fixed AirGels
shows the presence of extracellular MUC5AC and MUC5B gel-forming mucins (left). The
fluorescent lectin jacalin, which targets glycans, labels extracellular mucus in AirGels (right).
F. Mucociliary clearance in AirGels. i. trajectories of fluorescent microparticles in the lumen of
7-day-old and 34-day-old AirGels. ii. corresponding velocity distributions. Black lines indicate
the median velocity. iii. median particle velocities for three differentiated and undifferentiated
AirGels show the contribution of cilia beating in clearance. Each data point corresponds to the
median in each experiment; the gray bar shows the median of triplicates. Statistics:
independent samples Student t-test with Bonferroni correction ($p < 10^{-7}$). **G.** *P. aeruginosa*
infection of a 62-day-old cystic fibrosis AirGel. Confocal images were acquired 13h after
inoculation. *P. aeruginosa* constitutively expresses the fluorescent protein mScarlet. The
plasma membrane of epithelial cells was stained with CellMask Deep Red (pink). Mucus was
stained with jacalin (green) shortly before infection.

Figure 2: *P. aeruginosa* rapidly forms mucus-associated biofilms. **A.** i. maximal intensity projection images of *P. aeruginosa* in a 35-day-old AirGel (healthy donor) shows biofilm formation within hours. ii. mean biofilms cluster area for 3 AirGels. The bar indicates the range between the maximum and minimum of the three means. The circle represents the mean of the means. iii. percentage of clusters that were larger than 100 µm^2 in each replicate (black dots). The bars represent the mean across replicates. Statistics: paired samples Student t-test with Bonferroni correction ($p = 0.051$ and $p = 0.01$). **B.** *P. aeruginosa* rapidly colonizes mucus surfaces. Images show maximal intensity projection of confocal stacks of a 33-day-old infected AirGel at $t = 0$ and $t = 5.5$h post-inoculation. The graph quantifies the proportion of mucus not occupied by bacteria. Gray bars show the mean of triplicates. Statistics: paired samples Student t-test with Bonferroni correction ($p = 0.02$). **C.** Dynamic visualization of *P. aeruginosa* cluster fusion on mucus in a 33-day-old AirGel ($t_1 = 6.2$h). The blue and pink arrowheads show two aggregates that fuse within the first 15 min. The resulting cluster is indicated by a yellow arrowhead. This new cluster then moves closer to the one indicated by the white arrowhead. All images are maximal intensity projections from z-stacks. **D.** Kymographs showing the displacement of two clusters along their axis of motion. The bacterial aggregate and underlying mucus traveled together at an approximate speed of 0.5 µm/min.

353 **Figure 3: Mucus contraction drives rapid biofilm formation. A.** Time course visualization

of a mucus patch in a 35-day-old AirGel infected with WT *P. aeruginosa* (t_1 = 1.2h). Reference features are indicated by colored arrowheads. Dashed lines show their trajectories. **B.** Biofilm formation of PAO1 mutants unable to degrade mucus or to generate force (*t* = 5.5h). The *Δxcp* mutant that lacks type II secretion system necessary for secretion of mucinases forms biofilms similar to WT. The *ΔfliC* mutant lacking flagella also formed WT-like biofilms. In contrast, the *ΔpilA* mutant lacking T4P was unable to form luminal biofilms in AirGels. **C.** Finite element simulations of mucus deformation during surface contraction. i. schematic representation of *P. aeruginosa* applying contractile force on mucus by retracting T4P. ii. finite element simulation of deformation of a mucus cylinder at rest (top) and under active surface stress (bottom). Colormap indicate displacement of surface elements. iii. relative mucus area as a function of surface contractile modulus. As the surface contractile modulus increases, the relative area of mucus decreases. **D.** T4P retraction is necessary for biofilm formation. Images compare biofilms from WT *P. aeruginosa* and from the *ΔpilT* mutant unable to retract T4P (*t* = 5.5h). *ΔpilT* cluster area and percentage of large clusters is much smaller than WT (*N* = 3). Statistics: independent samples Student t-test with Bonferroni correction (*p* = 0.035 and *p* = 0.015). **E.** Mucus does not rearrange during *ΔpilT* colonization (*t* = 5.5h). Most of the mucus surface remains free of bacteria during *ΔpilT* colonization (N = 3). Statistics: independent samples Student t-test with Bonferroni correction (*p* = 0.01). **F.** Increased T4P activity speeds up biofilm formation on mucus. Comparison of biofilm formation by the *ΔpilH* mutant with hyperactive T4P with WT, at *t* = 2h and *t* = 5.5h after inoculation. *ΔpilH* already forms small biofilms after 2h. **G.** i. kinetics of biofilm size for WT, *ΔpilT* and *ΔpilH*. For each strain, we infected three AirGels from a healthy donor. Bars represent the range between the maximum and minimum of the means from triplicates, circles represent the overall mean. ii. comparison of percentage of large clusters for WT, *ΔpilT* and *ΔpilH*, over time. Statistics: one-way ANOVA for each time point, followed by a post-hoc Tukey test if the null hypothesis was rejected. At *t* = 4h, the differences between WT and *ΔpilH* (*p* = 0.003) and between *ΔpilH* and *ΔpilT* (*p* = 0.001) were significant. At *t* = 5.5h, the differences between WT and *ΔpilT* (*p* = 0.02) and between *ΔpilH* and *ΔpilT* (*p* = 0.001) were significant. **H.** *ΔpilH* dramatically contracts mucus. i. timelapse confocal images showing a dynamic event of mucus contraction by *ΔpilH*. ii. orthogonal views of the bacteria-covered mucus cluster at t_1 + 85 min, showing that PAO1 *ΔpilH* cells surround mucus. iii. relative mucus area changes measured during a 30 min time interval for WT and *ΔpilH*; since *ΔpilH* starts aggregating and remodeling mucus earlier than WT, the starting points of the recording differed (*ΔpilH*: 2.5h, WT: 6.2h, negative control: 8.1h). Images are maximal intensity projections of z-stacks throughout the figure except for the orthogonal projection in H.