Determination of a target image of freshwater fish composition in a mitigation project using

eDNA metabarcoding methods

Gen Ito[1,2], Hiroshi Yamauchi[3], Miwa Shigeyoshi[3], Kousuke Ashino[3], Chie Yonashiro[3], Maki

Asami[2], Yuko Goto[2], Hiroki Yamanaka[1, 2]

[1] Faculty of Advanced Science and Technology, Ryukoku University, 1-5 Yokotani, Seta

Oe-cho, Otsu, Shiga 520-2194, Japan

[2] Center for Biodiversity Science, Ryukoku University, 1-5 Yokotani, Seta Oe-cho, Otsu,

Shiga 520-2194, Japan

[3] Chuo Fukken Consultants, 4-11-10 Higashinakajima, Higashiyodogawa, Osaka, Osaka

533-0033, Japan

Corresponding author: Gen Ito

sakurahayabusa6647@gmail.com

Center for Biodiversity Science, Ryukoku University, 1-5 Yokotani, Seta Oe-cho, Otsu, Shiga

520-2194, Japan

Keywords: eDNA, Freshwater fishes, NMDS, Japan, intraspecific lineages, Shigenobu River

system

Abstract

Short-term target species for mitigation should be realistically and effectively selected for

restoration within a few years. Current short-term target species selection is based on spatial

distribution data of past and present biota in the neighborhood based on capture surveys.

However, current methods have the following problems: If historical spatial distribution data

are used, the risk of including large areas, where target species cannot be dispersed, exists, as

well as the risk of including extinct species. If recent spatial distribution data are used, the survey area may be narrowed due to time constraints, thereby increasing the possibility of missing species that should be selected as the target species. We examined the possibility of using the environmental DNA (eDNA) metabarcoding method, which can obtain spatial distribution data quickly, sensitively, and extensively, as a new short-term target species selection method. The study site was a restoration mitigation area (Kaihotsu–Kasumi) in the Shigenobu River system, Ehime Prefecture, Japan, where freshwater fish target species were selected based on capture surveys and past literature information. Furthermore, eDNA surveys were conducted at 17 sites in the Kaihotsu–Kasumi and its inflow and outflow rivers in 2020, and metabarcoding analysis using MiFish primers was conducted to determine the current freshwater fish fauna at each site. The visualization of the fish community composition obtained from the eDNA survey using NMDS showed that the Kaihotsu–Kasumi and surrounding watersheds were significantly divided into three clusters: upper reaches, middle and lower reaches, and estuarine reaches. Kaihotsu–Kasumi was included in Cluster 2, comprising the middle and lower reaches of the inflow and outflow rivers. Twenty-six species (including five non-native species) were detected in Cluster 2. Twenty-one fish species, excluding non-native species, were considered suitable as short-term target species as taxa with high potential for dispersal into Kaihotsu–Kasumi. The selection of short-term target species using the eDNA metabarcoding method covered all short- and medium-term target species (14 species) selected based on the capture surveys, and seven additional species were also proposed. The study also showed that intraspecific lineages of *Misgurnus anguillicaudatus* (Clades A and B), which were missed by the capture surveys, can be identified. These results indicate that the eDNA metabarcoding method can provide more comprehensive and realistic short-term target species selection than capture surveys, providing higher resolution monitoring through intraspecific lineage detection.

Introduction

The selection of restorable organisms—hereafter referred to as "target species"—is imperative for biodiversity restoration (Morimoto and Kameyama, 2001). In restoration mitigation in rivers, aquatic organisms naturally disperse from connecting waters. However, it is difficult to objectively and realistically select which species inhabiting which range should be the target species. An approach that relies on past to present spatial distribution data of organisms in neighboring areas based on capture surveys has the following problems. If historical spatial distribution data are used, the risk of including large areas where target species cannot be dispersed exists, as well as the risk of including extinct species. If recent spatial distribution data are used, the survey area may be narrowed due to time constraints, thereby increasing the possibility of missing species that should be selected as the target species. In addition, there may be no available spatial distribution data for nearby areas. The selection of target species often involves the establishment of species that can be restored in a short time (hereafter referred to as "short-term target species") and in a long time (hereafter referred to as "long-term target species"). Especially for short-term target species that are intended to be restored in a short time, selection should be more realistic, avoiding the selection of organisms that may not be distributed.

eDNA methods can obtain information on the spatial distribution of species more sensitively and extensively at a lower cost than capture surveys (Jerde et al., 2013; Sigsgaard et al., 2015; Sakata et al., 2017; Yamamoto et al., 2017; Oka et al., 2021). Many studies have shown the usefulness of eDNA methods for determining the spatial distribution and diversity of fish species (e.g., Yamamoto et al., 2016, 2017; Nakagawa et al., 2018; Doi et al., 2020; Oka et al., 2021) and that eDNA methods can be used to monitor rare species with low densities and to provide greater sensitivity in monitoring after removing artificial instream barriers (e.g., Sakata et al., 2017; Duda et al., 2021; Muha et al., 2021). In particular, the eDNA metabarcoding method can simultaneously detect many fish species at each location

(e.g., Yamamoto et al., 2017; Nakagawa et al., 2018). Because the eDNA metabarcoding

method can easily obtain spatial distribution data for current fish species, it can provide

objective and realistic targets for selecting target species after restoration mitigation.

Freshwater fishes are a major target species for restoration mitigation in rivers, but the

presence of cryptic non-native populations that cannot be distinguished by morphology is a

problem for biodiversity conservation (Mabuchi et al., 2008; Watanabe et al., 2017). eDNA is

a promising tool for detecting cryptic non-native populations (Uchii et al., 2016). MiFish

primers, commonly used in eDNA metabarcoding methods (Miya et al., 2015), can detect

many fishes at the species level, but the resolution of intraspecific lineages is not well

discussed. The ability to distinguish between native and non-native lineages using eDNA

metabarcoding methods would be useful in target species selection. Three lineages of

Misgurnus anguillicaudatus in the Japanese archipelago, which have been difficult to

morphologically identify, are now identified: two native lineages (Clades A and C sensu

Koizumi et al., 2009) and a non-native lineage from China (Clade B sensu Koizumi et al.,

2009). The native *M. anguillicaudatus* is categorized as *near threatened* by the Ministry of

the Environment Government of Japan (2020). Native and non-native lineages of *M.*

anguillicaudatus can be distinguished in the cytochrome *b* and control regions of

mitochondrial DNA (Morishima et al., 2008; Koizumi et al., 2009). Therefore, MiFish for the

12S rRNA region of mitochondrial DNA may also distinguish them. In this paper, we

examined whether eDNA metabarcoding methods using MiFish primers can distinguish

intraspecific lineages of *M. anguillicaudatus*.

The Shigenobu River flows 36 km into the Seto Inland Sea, and wetlands with springs as

their source of water exist along the banks of the middle reaches of the river. In the middle

reaches of the Shigenobu River, including wetlands, good habitats have decreased or

deteriorated due to urbanization and levee construction in recent years (Ministry of Land,

Infrastructure, Transport and Tourism. Shikoku Regional Development Bureau,   2008).

Therefore, restoration mitigation was conducted by the Ministry of Land, Infrastructure,

Transport and Tourism from 2014 to 2019 for Kaihotsu–Kasumi, one of the wetlands located

in the middle reaches of the Shigenobu River, to restore the spring area and improve its

connectivity with the mainstream. On the Shigenobu River, there were records of capture

surveys conducted every five years from 1994 to 2021 (Ministry of Land, Infrastructure,

Transport and Tourism, http://www.nilim.go.jp/lab/fbg/ksnkankyo/index.html, accepted 1

April 2022), as well as information on the distribution of endangered species (Ehime

Prefecture, Japan, 2014). Based on the above data on distribution, advice from experts, and

discussions with local residents, short-, medium-, and long-term target species for restoration

mitigation of Kaihotsu–Kasumi were selected. In addition, monitoring was conducted in 2020

using capture and eDNA metabarcoding surveys. Kaihotsu–Kasumi of the Shigenobu River is

a suitable field for testing whether the application of eDNA methods provides additional

benefits for target species selection by restoration mitigation. In this study, we examined the

usefulness of eDNA metabarcoding methods as a short-term target species selection using

eDNA metabarcoding monitoring data conducted in Kaihotsu–Kasumi in 2020.

Materials and methods

Water sampling

Water samples were collected from Kaihotsu–Kasumi (5 sites: K1−5), the outflow river

Shigenobu River (9 sites: S1−9), and the rivers (Uchi and Sagawa rivers) flowing into the

Kaihotsu–Kasumi (3 sites: U1−3) from October 26 to October 27, 2020 (Fig. 1; Table S1).

The basic water sampling protocol followed Minamoto et al. (2021), with some modifications.

Disposable plastic cups were used to collect water samples, and 500 ml of surface water was

collected per site. To prevent contamination, plastic cups were replaced with new plastic cups

at each water sampling site. After water sampling, 500 μl of 10% benzalkonium chloride

(BAC) was added immediately to inhibit DNA degradation (Yamanaka et al., 2017). Each

water sample was filtered in the field by loading a syringe with a 0.45 μm pore size Sterivex

filter. At the end of the filtration, we left the sample water containing BAC and plugged it. As

a field negative control (FNC), 500 ml of ultrapure water was filtered using the same method

as above after water sampling from three areas. After filtration, the samples were transported

to the laboratory at room temperature, and the Sterivex was stored in a –20°C freezer on

October 28, 2020.

DNA extraction and amplification

DNA extraction was performed using the DNeasy Blood and Tissue Kit (Qiagen, Hilden,

Germany), basically following Miya et al. (2016), with some modifications following

Hirohara et al. (2021).

PCR was performed according to Miya et al. (2015), and the partial 12S rRNA region of

mitochondrial DNA was amplified by the first and second PCRs using the MiFish-6N-U-F/R

primer pair. For the 1st PCR, in a 12 ul reaction volume containing 0.24 μl of KOD Fx Neo

(Toyobo, Osaka, Japan), 6 μl of 2 × Buffer for KOD Fx Neo, 2.4 μl of 2 mM dNTPs, 0.7 μl

each of 5 μM MiFish-6N-U-F/R, and 2 μl of 10 × diluted template DNA were mixed. The

thermal cycle profile was 94°C for 2 min; 40 cycles of 98°C for 10 s, 65°C for 30 s, and 68°C

for 30 s; and 68°C for 5 min. Eight PCR replications were used for each sample and a PCR

non-template control (PCR–NTC) using ultrapure water. The PCR amplicons were treated

with ExoSAP-IT (ThermoFisher Scientific, Waltham, USA). The PCR amplicons from each

of the eight replications were aggregated by 5 μl to a total of 40 μl, and 3.6 μl of ultrapure

water and 0.4 μl of ExoSAP-IT were added per sample and treated at 37°C for 15 min and

80°C for 15 min. The 2nd PCR was performed to add MiSeq adaptor sequences and eight bp

index sequences to both amplicon ends. The total reaction volume of the 2nd PCR was 12 μl

containing 0.24 μl of KOD Fx Neo, 6 μl of 2 × Buffer for KOD Fx Neo, 2.4 μl of 2 mM dNTP,

0.7 μl each of 5 μM MiFish-6N-U-F/R, 1 μl of ultrapure water, and 1 μl of the 1st PCR

amplicons. The thermal cycle profile was 95°C for 3 min; 10 cycles of 98°C for 20 s, 65°C

for 15 s, and 72°C for 15 s; and 72°C for 5 min. A 2% agarose gel electrophoresis was

performed to confirm the amplification of the 2nd PCR amplicons.

Amplicon library and Miseq sequencing

The samples were aggregated in volumes ranging from 2 or 3 µl per sample, depending on

the results of the electrophoresis (Table S2). AMPure XP (Beckman Coulter, Brea CA, USA)

was added to 0.8 µl per 1 µl of each sample and treated according to the manual. DNA was

eluted with 60 µl of TE buffer. Then, 25 µl of the treated DNA solution was added to a 2%

E-Gel Size Select (Thermo Fisher Scientific, Waltham, USA) to retrieve the target band (ca.

370 bp). The collected DNA concentration was measured using a Qubit dsDNA HS assay kit

(Thermo Fisher Scientific, Waltham, USA). Finally, a bioanalyzer (Agilent Technologies,

Santa Clara, CA, USA) was used to confirm that PCR amplicons of the desired size were

obtained.

The libraries were sequenced using an Illumina MiSeq with a reagent kit v2 nano for 2 ×

150 bp PE (Illumina, San Diego, CA, USA). Raw reads were demultiplexed into each sample

by a preinstalled program in MiSeq, and the outputted fastq files were processed by an

in-house bioinformatic pipeline that uses USEARCH v11.0.667 (Edgar 2010) as its core

program as follows. 1) Pair-end reads were merged using the *fastq_mergepairs* function. 2)

Primer sequences were removed from both ends of the reads using *stripleft* and *stripright*

functions. 3) Short reads less than 100 bp and those with a Q-Score less than 30 were

removed using *fastq_minlen* and *fastq_truncqual* functions, respectively. 4) Identical reads

were dereplicated using the *fastx_uniques* function and treated as operational taxonomic units

(OTUs) with read numbers gathered up (i.e. size) appended to the OTU information. 5)

Denoising was performed using the *unoise3* function, and the OTUs with size less than 4

were removed. Moreover, the OTUs that were judged as chimera were removed. 6)

Taxonomic identity was assigned to the OTUs as follows: The sequences of each OTU were

used as queries against a local sequence database at the threshold level of 0.95 using the

usearch_global function. The database was a compilation of mitochondrial 12S sequences of

fish downloaded from the National Center for Biotechnology Information website

(https://www.ncbi.nlm.nih.gov/) on January 15, 2020. The hit sequences retrieved from the

database for each OTU were converted into phylogenic trees and visually checked to assign a

taxonomic identity for the focal OTU. The granularity of the taxonomic assignment to the

OTUs was mainly at the species level, but some OTUs were identified at the genus level

when a sequence cluster, including the query sequence, included more than one species. Taxa

that were unlikely to be distributed around the Shigenobu River (mainland of the Japanese

archipelago) based on known geographic distribution information (Hosoya, 2019) were

excluded (e.g., *Acanthopagrus sivicolus* and *Luciogobius_ryukyuensis*, *Tridentiger kuroiwae*).

Distinguishing between intraspecific lineages

For *M. anguillicaudatus*, we tried to distinguish between intraspecific lineages using the

following methods. First, a 124 bp 12S rRNA (MiFish) tree was constructed using the

sequence data output as *M. anguillicaudatus* OTU and *M. mizolepis* OTU. In addition, we

used *Lefua echigonia*, the related species, as an outgroup (LC069488). Next, we constructed a

1015 bp cyt *b* tree, referring to Koizumi et al. (2009), in which molecular phylogenetic

analyses of *M. anguillicaudatus* in Japan were conducted. For phylogenetic analysis,

nucleotide sequence data of *M. anguillicaudatus* and *M. dabryanus* registered in DDBJ,

EMBL, and GenBank by Koizumi et al. (2009) were used: Clade A (native lineage,

AB473261, AB473312), Clade B (non-native lineage, AB473360, AB473346), Clade C

(native lineage, AB473381, AB473354, AB473382), and *M. dabryanus* (AB473408,

AY625701). The complete mitochondrial genome sequences included in the MiFish

phylogenetic tree (AP011291, AP017654, HM856629, KC509900, KC823274, MF579257,

KF771003, KJ939363, KM186181, KM186183, MG938589, MG938590, NC027448) were

also added to the cyt b phylogenetic tree. In addition, we used *L. echigonia* as an outgroup

(LC079000, Ito et al., 2019). Multiple alignments of nucleotide sequences were performed

using CLUSTAL W (Thompson et al., 1994). Phylogenetic analyses were conducted using the

neighbor-joining (NJ) method. The NJ analyses were performed using MEGA X (Kumar et al.,

2018), with the *p*-distance. The reliability of each internal branch was evaluated using a

bootstrap test with 1000 replicates.

Community analysis

The differences in community composition were visualized using nonmetric

multidimensional scaling (NMDS). The dissimilarity of the community in NMDS was

calculated based on the Jaccard index using the vegdist function in the vegan package. The

simprof function in the clustsig package was used to cluster the community, and

PERMANOVA was conducted using the adonis function in the vegan package used to test

whether the clusters obtained significantly differed from each other. All of the above statistics

were conducted using R version 4.1.2 (R Core Team 2021).

Capture survey

Capture surveys in Kaihotsu–Kasumi were conducted on August 27 and 28, 2020, and on

October 26 and 27, 2020 at the same five sites as the eDNA sampling sites. The capture

survey was conducted using hand nets, net pots, and bottle traps, with permission from Ehime

Prefecture. The capture survey on October 26 was conducted after the eDNA survey.

Results

Miseq sequencing

A total of 1,435,754 reads were obtained by Miseq run from 17 sampling sites, 3 of FNCs,

and 1 of PCR–NTC. After the data processing noted in the Material and Methods section, 314

OTUs were identified as fish species comprising 817,797 reads. Species assignment

automatically conducted by Usearch was manually confirmed to check the independence of

each OTU from the other. If the representative sequence of the focal OTU could not be

identified at the species level, the OTU was identified at the genus level. Five reads of

Opsariichthys platypus were detected from an FNC of the Shigenobu River; therefore, the

equivalent read number was subtracted from the *O. platypus* reads detected from the other

sites in the Shigenobu River. No read was detected from the PCR–NTC.

Fish species detected in the samples

Fish species from 38 taxa were identified in total. Common saltwater fishes for food use

were excluded from the results: *Etrumeus teres*, *Trachurus japonicus*, *Pseudocaranx dentex*,

Seriola quinqueradiata, *Pagrus major*, and *Pleurogrammus azonus*. However, peripheral

freshwater fishes (*Plotosus japonicus*, *Mugil cephalus*, *Plectorhinchus cinctus*,

Acanthopagrus schlegelii, *Omobranchus punctatus*, *Luciogobius pallidus*, *Luciogobius* spp.,

Acanthogobius flavimanus, *Tridentiger trigonocephalus*, *Favonigobius gymnauchen*,

Gymnogobius breunigii, *Gymnogobius scrobiculatus*, *Takifugu* spp.) were included in the

results. Fish species of 36, 10, and 19 taxa were identified from the Shigenobu River (nine

sites), Uchi and Sagawa River (three sites), and Kaihotsu–Kasumi (five sites), respectively

(Table S3).

MiFish and cyt *b* trees are shown in Fig. 2. In the MiFish tree, the sequences of *M.*

anguillicaudatus OTU and *M. mizolepis* OTU differed. The *M. anguillicaudatus* OTU

included three sequences, OTU 57, 259, and 283, and 1 mitogenome sequence (AP011291).

The *M. mizolepis* OTU included three sequences, OTU 40, 123, and 190, and 12 mitogenome

sequences. In the cyt *b* tree, three intraspecific lineages of *M. anguillicaudatus* (Clades A, B,

and C), and *M. dabryanus* were distinguished with high bootstrap probability (100%), similar

to Koizumi et al. (2009) and Perdices et al. (2016). Clade C (native lineage) contained one

mitogenome sequence (AP011291), and Clade B (non-native lineage) contained the other 12

mitogenome sequences. No mitogenome sequences were identified in *M. dabryanus* or *M.*

anguillicaudatus Clade A.

Clustering based on community analysis

The community analysis results showed that the community was significantly differentiated

into three clusters (Fig. 3; $P < 0.001$). Cluster 1 comprised three sites (S1−3) in the upper

stream of the Shigenobu River (elevation 178.7−419.9 m), where five taxa were detected,

including *Phoxinus oxycephalus jouyi*, *Oncorhynchus* spp, and *Rhinogobius flumineus*.

Cluster 2 included five sites in the middle stream of the Shigenobu River (elevation 6.7−99.3

m), all sites in the Kaihotsu–Kasumi, and all sites in the Uchi and Sagawa River (K1−5, S4−8,

U1−3), where 26 taxa were detected, including *Pseudorasbora parva*, *Gnathopogon*

elongatus, *Oryzias latipes*, and *Odontobutis obscura*. Fourteen of these taxa were detected

only in Cluster 2. Cluster 3 comprised only one site (S9) in the estuary of the Shigenobu

River (elevation −0.8 m), where 21 taxa were detected, including *Plotosus japonicus*, *Mugil*

cephalus, and *Acanthopagrus schlegelii*. Twelve of these taxa were detected only in Cluster 3

(Table 1).

For Cluster 2, 23 taxa were identified from the Shigenobu River, 10 taxa from the Uchi and

Sagawa River, and 19 taxa from Kaihotsu–Kasumi. Of these, *Tanakia limbata* and *Takifugu*

spp. were detected only in Kaihotsu–Kasumi. Seven taxa, including *P. parva* and *Cobitis*

shikokuensis, were detected only in the Shigenobu, Uchi, and Sagawa rivers (Table 1).

Fish species detected by capture surveys

The capture surveys in 2020 identified 15 taxa in Kaihotsu–Kasumi (Table 2). *C.*

shikokuensis and *Lepomis macrochirus* were identified only in the capture surveys.

Discussion

All of the 38 taxa in 16 families of fishes detected by eDNA in this study were recorded in

previous capture surveys in the Shigenobu River (Ministry of Land, Infrastructure,

Transport and Tourism, http://www.nilim.go.jp/lab/fbg/ksnkankyo/index.html, accepted 1

April 2022). Therefore, the eDNA survey results were determined to be reasonable. The six

taxa that could not be distinguished by eDNA were compared with previous capture surveys

(Ministry of Land, Infrastructure, Transport and Tourism,

http://www.nilim.go.jp/lab/fbg/ksnkankyo/index.html, accepted 1 April 2022), and the

following taxa were identified: *Carassius* spp. as *Carassius* sp. (Silver crucian carp) or

Carassius buergeri (Giant golden crucian carp), *Oncorhynchus* spp. as *Oncorhynchus masou*

ishikawae, *Luciogobius* spp. as *Luciogobius guttatus* or *Luciogobius martellii*, *Tridintiger* spp.

as *Tridintiger obscurus* or *Tridintiger brevispinis*, *Rhinogobius* spp. as *Rhinogobius nagoyae*,

Rhinogobius fluviatilis, *Rhinogobius* sp. OR or *Rhinogobius tyoni*, and *Takifugu* spp. as

Takifugu niphobles or *Takifugu poecilonotus*.

In this paper, we attempted to identify intraspecific lineages in loach using the eDNA

metabarcoding method with MiFish primers. Based on the results of the Cyt *b* phylogenetic

tree, including the mitogenome sequence, *M. anguillicaudatus* OTU is the native lineage

(Clade C) and *M. mizolepis* OTU is the non-native lineage (Clade B). *M. anguillicaudatus*

Clade B has also been recorded from the Shigenobu River system as a non-native lineage by

capture surveys (Shimizu and Takagi, 2010). No matching sequences were found between *M.*

anguillicaudatus and *M. mizolepis* OTUs; thus, both OTUs are genetically distinct. However,

M. dabryanus (including *Paramisgurnus dabryanus* and *M. mizolepis*) and *M.*

anguillicaudatus sequences were mixed in the *M. mizolepis* OTU. Molecular phylogenetic

studies of mtDNA indicate that the *M. dabryanus* is not contained within *M. anguillicaudatus*

Clades A and B, and shows a phylogenetic position closely related to *M. anguillicaudatus*

Clade A and *M. nikolskyi* (Koizumi et al., 2009; Perdices et al., 2016). The mtDNA of some *M.*

dabryanus populations is known to be contained within *M. anguillicaudatus* Clade B, but this

M. dabryanus mtDNA sequence is likely to have originated from hybridization with *M.*

anguillicaudatus (Okada et al., 2020). Intentional exclusion of sequences derived from

hybridization is difficult. Therefore, a phylogenetic tree that includes mitogenomic sequences

would help identify intraspecific lineages.

Native lineage (Clade C) and non-native lineage (Clade B) of *M. anguillicaudatus* were

identified from Kaihotsu–Kasumi using eDNA metabarcoding methods. The successful

identification of native and non-native lineages of *M. anguillicaudatus* using eDNA is one of

the advantages of introducing eDNA in monitoring. Future studies on the feasibility of

identifying intraspecific lineages for a larger number of fishes will contribute to improving

the resolution of monitoring using the eDNA metabarcoding method.

Community analysis showed that the fish community of the Shigenobu River was

significantly divided into three clusters (Clusters 1–3). Each cluster has an upstream (Cluster

1), middle and downstream (Cluster 2), and estuarine (Cluster 3) fish community, suggesting

that changes in the fish community in response to the river environment were a factor in

separating the clusters. The communities in Kaihotsu–Kasumi (K1–5) were included in

Cluster 2, which showed a community structure in the middle and lower reaches of the river.

Thus, the fish community at Kaihotsu–Kasumi represents the community structure of the

middle and lower reaches of the Shigenobu River; 26 taxa of fish were detected at the Cluster

2 sites, 17 of which were common between Kaihotsu–Kasumi and neighboring waters (S4–8

and U1–3). This indicated a high degree of commonality between the fish communities in

Kaihotsu–Kasumi and neighboring waters, probably because of the natural dispersal of fishes

inhabiting Kaihotsu–Kasumi from neighboring waters. Kaihotsu–Kasumi is considered to

provide a good fish habitat due to its high connectivity with neighboring waters.

Nineteen taxa were detected by eDNA in Kaihotsu–Kasumi, five of which were not

detected in the capture surveys conducted during the same period. Fish surveys using eDNA

tend to detect more species than capture surveys (e.g., Oka et al., 2021), and this study

showed similar results. *Plecoglossus altivelis altivelis*, *C. shikokuensis*, and *L. macrochirus*

were detected only in the capture surveys. This suggests that a combination of capture surveys

may be effective for a more accurate understanding of the fish community.

The community structure of Kaihotsu–Kasumi resembled that of Cluster 2, which showed a

community structure in the middle and lower reaches of the river. Therefore, 21 taxa could be

selected as short-term target species for Kaihotsu–Kasumi, excluding non-native species

(*Cyprinus carpio*, *Tanakia limbata*, *M. anguillicaudatus* non-native lineage, *L. macrochirus*,

Micropterus salmoides; Hosoya, 2019) out of the 26 taxa included in Cluster 2. The target

species not identified in Kaihotsu–Kasumi were *Pseudorasbora parva*, *Oncorhynchus* spp.,

and *Plectorhinchus cinctus*. *P. parva* is categorized as Near Threatened by Ehime Prefecture,

Japan (2014). The slow flow of Kaihotsu–Kasumi is considered a suitable habitat and

breeding environment for *P. parva*. In the future, efforts should be made to facilitate the

natural dispersal of *P. parva*. The main habitats for the remaining two species are the upper

reaches of rivers for *O.* spp. and estuarine and coastal areas for *P. cinctus* (Hosoya, 2019).

Therefore, it is rare for the above two species to be recorded in Kaihotsu–Kasumi in the

middle reaches of the river.

In this paper, we attempted to select short-term target species after restoration mitigation

using the eDNA metabarcoding method. As a result, 21 short-term target species were

selected. For target species based on historical capture surveys in Kaihotsu–Kasumi, 9, 5, and

2 species were short-term (within 3 years), medium-term (within 20 years), and long-term

(>20 years), respectively (Table S4). Of these, 14 of the short- and medium-term target

species were among the short-term target species selected by eDNA. Seven new species were

included in the short-term target species established by eDNA. Therefore, the selection of

short-term target species using the eDNA metabarcoding method is more effective than

capture surveys in selecting fish species that are expected to recover in the short term in a

more comprehensive manner, probably because eDNA is more sensitive to the current fish

community than capture surveys. In addition, fish such as *O.* spp. and *P. cinctus*, which are

not considered to inhabit the Kaihotsu–Kasumi habitat, were also selected as short-term target

species by eDNA. Future studies of selection methods, including fish preference by habitat,

will bring us closer to a more realistic selection of short-term target species.

Funding

This work was supported by Environment Research and Technology Development Fund

(JPMEERF20204004) of the Environmental Restoration and Conservation Agency, Japan.

Credit author statement

Gen Ito: Data analysis, Writing – original draft, Writing – review and editing. Hiroshi

Yamauchi: Planning and conducting field surveys, Writing – review and editing. Miwa

Shigeyoshi, Kousuke Ashino and Chie Yonashiro: Planning and conducting field surveys.

Maki Asami and Yuko Goto: Experiments in the laboratory. Hiroki Yamanaka: Planning and

conducting field surveys, Data analysis, Writing – review and editing, Conceptualization,

Funding acquisition.

Declaration of Competing Interest

The authors declare no competing financial interests.

Acknowledgements

We express sincere thanks to members of the Civil Engineering Division1, Matsuyama

river National Highway Office, Shikoku regional development bureau for their help with the

field surveys.

References

Doi, H., Matsuoka, S., Matsuzaki, S. S., Nagano, M., Sato, H., Yamanaka, H., Matsuhashi, S., Yamamoto, S., Minamoto, T., Araki, H., Ikeda, K., Kato, A., Kumei, K., Maki,N., Mitsuzuka, T., Takahara, T., Toki, K., Ueda, N., Watanabe, T., Yamazoe, K., Miya, M., 2020. Effects of species traits and ecosystem characteristics on species detection by eDNA metabarcoding in lake fish communities. Bio Rxiv. doi.org/10.1101/2020.09.25.314336.

Duda, J.J., Hoy, M.S., Chase, D. M., Pess, G.R., Brenkman, S. J., McHenry, M.M., Ostberg, C.O., 2021. Environmental DNA is an effective tool to track recolonizing migratory fish following large-scale dam removal. Environmental DNA. 3, 121–141. https://doi.org/10.1002/edn3.134.

Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics. 26, 2460–2461. https://doi.org/10.1093/bioin forma tics/btq46 1.

Ehime Prefecture, Japan., 2014. Freshwater fishes. Red Data Book Ehime. Ehime Prefecture, Matsuyama, pp. 107–136.

Hirohara, T., Tsuri, K., Miyagawa, K., Paine, R.T.R., Yamanaka, H., 2021. The application of PMA (propidium monoazide) to different target sequence lengths of Zebrafish eDNA: a new approach aimed toward improving environmental DNA ecology and biological surveillance. Frontiers in Ecology and Evolution. 9, 632973. https://doi.org/10.3389/fevo.2021.632973.

Hosoya, K. (Ed), 2019. Freshwater fishes of Japan, extended and revised edition. Yama-kei Publishers, Tokyo, Japan.

Ito, G., Koya, Y., Kitanishi, S., Horiike, T., Mukai, T., 2019. Genetic population structure of the eight-barbel loach *Lefua echigonia* in the Ise Bay region, a single paleo-river basin in central Honshu, Japan. Ichthyological Research. 66, 411–416. https://doi.org/10.1007/s10228-019-00683-z

Jerde, C.L., Chadderton, W.L., Mahon, A.R., Renshaw, M.A., Corush, J., Budny, M.L.,

Mysoorekar, S., Lodge, M., 2013. Detection of Asian carp DNA as part of a Great Lakes

basin-wide surveillance program. Canadian Journal of Fisheries and Aquatic Sciences. 70,

522–526. https://doi.org/10.1139/cjfas-2012-0478.

Koizumi, N., Takemura, T., Watabe, K., Mori, A., 2009. Genetic variation and diversity of

Japanese loach inferred from mitochondrial DNA -Phylogenetic analysis using the

cytochrome *b* gene sequences-. Trans. of JSIDRE. 259, 7–16.

https://doi.org/10.11408/jsidre.77.7.

Kumar, S., Stecher, G., Li, M. Knyaz, C., Tamura, K., 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platform. Molecular Biology and

Evolution. 35, 1547–1549. doi: 10.1093/molbev/msy096.

Mabuchi, K, Senou, H., Nishida, M., 2008. Mitochondrial DNA analysis reveals cryptic

large-scale invasion of non-native genotypes of common carp (*Cyprinus carpio*) in Japan.

Molecular Ecology. 17, 796–809. https://doi.org/10.1111/j.1365-294X.2007.03626.x.

Minamoto, T., Miya, M., Sado, T., Seino, S., Doi, H., Kondoh, M., Nakamura, K., Takahara,

T., Yamamoto, S., Yamanaka, H., Araki, H., Iwasaki, W., Kasai, A., Masuda, R., Uchii, K.,

2021. An illustrated manual for environmental DNA research: water sampling guidelines

and experimental protocols. Environmental DNA. 3, 8–13.

https://doi.org/10.1002/edn3.121.

Ministry of Land, Infrastructure, Transport and Tourism. Shikoku regional development

bureau., 2008. The river management program of the Shigenobu River system. Ministry of

Land, Infrastructure, Transport and Tourism. Shikoku regional development bureau.

Takamatsu, Japan.

https://www.skr.mlit.go.jp/matsuyam/river/seibikeikaku/files/plan/plan_all.pdf. (accessed 1

April 2022).

Ministry of the Environment Government of Japan., 2020. Ministry of the environment red

list 2020 (Translated from Japanese). http://www.env.go.jp/press/files/jp/114457.pdf.

(accessed 1 April 2022).

Miya, M., Minamoto, T., Yamanaka, H., Oka, S., Sato, K., Yamamoto, S., Sado, T., Doi, H.,

2016. Use of a filter cartridge for filtration of water samples and extraction of

environmental DNA. Journal of Visualized Experiments. 117, e54741.

https://doi.org/10.3791/54741.

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J.Y., Sato, K., Minamoto, T. Yamamoto,

S., Yamanaka, H. Araki, H. Kondoh M., Iwasaki,W., 2015. MiFish, a set of universal PCR

primers for metabarcoding environmental DNA from fishes: detection of more than 230

subtropical marine species. Royal Society Open Science. 2, 150088.

http://dx.doi.org/10.1098/rsos.150088.

Morimoto, Y., Kameyama A. (Eds), 2001. Ecological mitigation in theory and practice. Soft

Science, Inc, Tokyo, Japan.

Morishima, K., Nakamura-Shiokawa, Yuka., Bando, E., Li, Y.-J., Boron´, A., Khan, M. R.,

Arai, K., 2008. Cryptic clonal lineages and genetic diversity in the loach *Misgurnus*

anguillicaudatus (Teleostei: Cobitidae) inferred from nuclear and mitochondrial DNA

analyses. Genetica. 132, 159–171. https://doi.org/10.1007/s10709-007-9158-1.

Muha, T.P., Rodriguez-Barreto, D., O'Rorke, R., Garcia de Leaniz, C., Consuegra, S., 2021.

Using eDNA Metabarcoding to Monitor Changes in Fish Community Composition After

Barrier Removal. Frontiers in Ecology and Evolution. 9, 629217.

https://doi.org/10.3389/fevo.2021.629217.

Nakagawa, H., Yamamoto, S., Sato, Y., Sado, T., Minamoto, T., Miya, M., 2018. Comparing

local- and regional-scale estimations of the diversity of stream fish using eDNA

metabarcoding and conventional observation methods. Freshwater Biology. 63, 569–580.

https://doi.org/10.1111/fwb.13094.

Oka, S., Doi, H., Miyamoto, K., Hanahara, N., Sado, T., Miya, M., 2021. Environmental DNA

metabarcoding for biodiversity monitoring of a highly diverse tropical fish community in a

coral reef lagoon: Estimation of species richness and detection of habitat segregation.

Environmental DNA. 3, 55–69. https://doi.org/10.1002/edn3.132.

Okada, R., Shimizu, T., Kitagawa, T., 2020. Evidence of a secondary interspecific

mitochondrial DNA introgression in the pond loach *Misgurnus dabryanus* (Teleostei:

Cobitidae) population introduced in Japan. Applied Ichthyology. 36, 655–667.

https://doi.org/10.1111/jai.14079.

Perdices, A., Bohlen, J., Šlechtová, V., Doadrio, I., 2016. Molecular Evidence for Multiple

Origins of the European Spined Loaches (Teleostei, Cobitidae). Plos One. 11, e0144628.

https://doi.org/10.1371/journal. pone.0144628.

R Core Team, 2021. R: A language and environment for statistical computing. R Foundation

for Statistical Computing. https://www.R-project.org.

Sakata, M.K., Maki, N., Sugiyama, H., Minamoto T., 2017. Identifying a breeding habitat of a

critically endangered fish, Acheilognathus typus, in a natural river in Japan. Science Nature.

104, 100. https://doi.org/10.1007/s00114-017-1521-1.

Shimizu, T., Takagi, M., 2010. Genetic structure of natural populations of *Misgurnus*

anguillicaudatus in Ehime Prefecture impacted by recent exotic introductions, inferred

from mitochondrial DNA analysis. Japan. Journal of Ichthyology. 57, 13 – 26.

https://doi.org/10.11369/jji.57.13

Sigsgaard, E.E., Carl, H., Møller P.R., Thomsen P.F., 2015. Monitoring the near-extinct

European weather loach in Denmark based on environmental DNA from water samples.

Biological Conservation. 183, 46–52. https://doi.org/10.1016/j.biocon.2014.11.023.

Thompson, J.D., Higgins, D.G., Gibson, T.J., 1994. CLUSTAL W: improving the sensitivity

of progressive multiple sequence alignment through sequence weighting, position-specific

gap penalties and weight matrix choice. Nucleic Acids Research. 22, 4673–4680.

https://doi.org/10.1093/nar/22.22.4673.

Uchii, K., Doi, H., Minamoto, T., 2016. A novel environmental DNA approach to quantify the

cryptic invasion of non-native genotypes. Molecular Ecology Resources. 16, 415–422.

https://doi.org/10.1111/1755-0998.12460.

Yamamoto, S., Masuda, R., Sato, Y., Sado, T., Araki, H., Kondoh, M., Minamoto, T., Miya,

M., 2017. Environmental DNA metabarcoding reveals local fish communities in a

species-rich coastal sea. Scientific Reports. 7, 40368. https://doi.org/10.1038/srep40368.

Yamamoto, S., Minami, K., Fukaya, K., Takahashi, K., Sawada, H., Murakami, H., Tsuji, S.,

Hashizume, H., Kubonaga, S., Horiuchi, T., Hongo, M., Nishida, J., Okugawa, Y., Fujiwara,

A., Fukuda, M., Hidaka, S., Suzuki, K.W., Miya, M., Araki, H., Yamanaka, H., Maruyama,

A., Miyashita, K., Masuda, R., Minamoto, T., Kondoh, M., 2016. Environmental DNA as a

'Snapshot' of fish distribution: a case study of Japanese Jack Mackerel in Maizuru Bay, Sea

of Japan. PLoS One. 11, e0149786. https://doi.org/10.1371/journal.pone.0149786.

Yamanaka, H., Minamoto, T., Matsuura, J., Sakurai, S., Tsuji, S., Motozawa, H., Hongo, M.,

Sogo, Y., Kakimi, N., Teramura, I., Sugita, M., Baba, M., Kondo, A., 2017. A simple

method for preserving environmental DNA in water samples at ambient temperature by

addition of cationic surfactant. Limnology. 18, 233–241.

https://doi.org/10.1007/s10201-016-0508-5.

Watanabe, K., Tominaga, K., Nakajima, J., Kakioka, R., Tabata, R., 2017. Japanese freshwater

fishes: Biogeography and cryptic diversity. In M. Motokawa & H. Kajihara (Eds.), Species

diversity of animals in Japan. Springer, Tokyo, Japan. pp. 183–227.

Figure legends

Fig. 1 Water sampling sites in the Shigenobu River system. Each circle symbol indicates

sampling sites. This altitude map was used with permission from the Geospatial Information

Authority of Japan (https://maps.gsi.go.jp/).

Fig. 2 Neighbor-joining trees of 124 bp 12S rRNA (MiFish) gene (left) and 1015 bp

cytochrome *b* gene (right) sequences of *M. anguillicaudatus*. The numbers at the nodes

indicate bootstrap probabilities. The complete mitochondrial genome sequences are indicated

by open boxes.

Fig. 3 NMDS plot of fish species community compositions at different eDNA sampling sites.

The community was significantly differentiated into three clusters (PERMANOVA: $P <$

0.001). Green, red, and blue indicate Clusters 1, 2, and 3, respectively.

Table 1 Fish species in each cluster.

Table 2 Fish species confirmed by eDNA and capture surveys in Kaihotsu–Kasumi.

Table S1 Latitude, longitude, and elevation of each sampling site.

Table S2 The volume of aggregated 2nd PCR amplicons in each sample.

Table S3 Number of reads of detected species by MiFish metabarcoding.

Table S4 Target species by eDNA and capture surveys.

12S (MiFish)

Cyt *b*

Table 1 Fish species in each cluster

Family	Species	Cluster 1	Cluster 2			Cluster 3
		S1−3	S4−8	U1−3	K1−5	S9
Anguillidae	*Anguilla japonica*		+		+	+
Cyprinidae	*Cyprinus carpio**		+	+	+	+
	Carassius spp.		+		+	
	*Tanakia limbata**				+	
	Opsariichthys platypus	+	+	+	+	+
	Candidia temminckii	+	+	+	+	+
	Phoxinus oxycephalus jouyi	+	+	+	+	
	Pseudorasbora parva		+			
	Gnathopogon elongatus elongatus		+		+	
	Pseudogobio esocinus		+	+	+	+
Cobitidae	*Misgurnus anguillicaudatus* (native)		+		+	
	Misgurnus anguillicaudatus (non-native)*		+		+	
	Cobitis shikokuensis		+			
	Cobitis sp. BIWAE type A		+		+	
Siluridae	*Silurus asotus*		+		+	
Plotosidae	*Plotosus japonicus*					+
Plecoglossidae	*Plecoglossus altivelis altivelis*		+			+
Salmonidae	*Oncorhynchus* spp.	+	+			
Mugilidae	*Mugil cephalus*					+
Adrianichthyidae	*Oryzias latipes*		+		+	
Centrarchidae	*Lepomis macrochirus**		+			
	*Micropterus salmoides salmoides**			+		
Haemulidae	*Plectorhinchus cinctus*		+			
Sparidae	*Acanthopagrus schlegelii*					+
Blenniidae	*Omobranchus punctatus*					+
Odontobutidae	*Odontobutis obscura*		+	+	+	
Gobiidae	*Luciogobius pallidus*					+
	Luciogobius spp.					+
	Acanthogobius flavimanus					+
	Tridentiger spp.		+	+	+	+
	Tridentiger trigonocephalus					+
	Rhinogobius flumineus	+	+	+	+	
	Rhinogobius similis					+
	Rhinogobius spp.		+	+	+	+
	Favonigobius gymnauchen					+
	Gymnogobius breunigii					+
	Gymnogobius scrobiculatus					+
Tetraodontidae	*Takifugu* spp.				+	+
	Total	5	23	10	19	21

* non-native species

Table 2 Fish species confirmed by eDNA and capture surveys in the Kaihotsu-Kasumi

Family	Species	eDNA	Capture
Anguillidae	*Anguilla japonica*	+	
Cyprinidae	*Cyprinus carpio**	+	+
	Carassius spp.	+	+
	*Tanakia limbata**	+	
	Opsariichthys platypus	+	+
	Candidia temminckii	+	+
	Phoxinus oxycephalus jouyi	+	+
	Pseudorasbora parva		
	Gnathopogon elongatus elongatus	+	
	Pseudogobio esocinus	+	+
Cobitidae	*Misgurnus anguillicaudatus* (native)	+	+
	Misgurnus anguillicaudatus (non-native)*	+	
	Cobitis shikokuensis		+
	Cobitis sp. BIWAE type A	+	+
Siluridae	*Silurus asotus*	+	
Plotosidae	*Plotosus japonicus*		
Plecoglossidae	*Plecoglossus altivelis altivelis*		+
Salmonidae	*Oncorhynchus* spp.		
Mugilidae	*Mugil cephalus*		
Adrianichthyidae	*Oryzias latipes*	+	+
Centrarchidae	*Lepomis macrochirus**		+
	*Micropterus salmoides salmoides**		
Haemulidae	*Plectorhinchus cinctus*		
Sparidae	*Acanthopagrus schlegelii*		
Blenniidae	*Omobranchus punctatus*		
Odontobutidae	*Odontobutis obscura*	+	+
Gobiidae	*Luciogobius pallidus*		
	Luciogobius spp.		
	Acanthogobius flavimanus		
	Tridentiger spp.	+	+
	Tridentiger trigonocephalus		
	Rhinogobius flumineus	+	+
	Rhinogobius similis		
	Rhinogobius nagoyae	+	+
	Rhinogobius sp. OR		+
	Favonigobius gymnauchen		
	Gymnogobius breunigii		
	Gymnogobius scrobiculatus		
Tetraodontidae	*Takifugu* spp.	+	
	Total	19	17

* non-native species