Integrating proteomic data with metabolic modelling provides insight into key

pathways of *Bordetella pertussis* biofilms

Hiroki Suyama[1], Laurence Don Wai Luu[1], Ling Zhong[2], Mark J. Raftery[2] and Ruiting Lan[1*]

[1]School of Biotechnology and Biomolecular Sciences, University of New South Wales,

Sydney, New South Wales, Australia

[2]Bioanalytical Mass Spectrometry Facility, University of New South Wales, Sydney, New

South Wales, Australia

*Corresponding Author

Email: r.lan@unsw.edu.au

Phone: 61-2-9385 2095

Fax: 61-2-9385 1483

Abstract

Pertussis, commonly known as whooping cough is a severe respiratory disease caused by the bacterium, *Bordetella pertussis*. Despite widespread vaccination, pertussis resurgence has been observed globally. The development of the current acellular vaccine (ACV) has been based on planktonic studies. However, recent studies have shown that *B. pertussis* readily forms biofilms. A better understanding of *B. pertussis* biofilms is important for developing novel vaccines that can target all aspects of *B. pertussis* infection. This study compared the proteomic expression of biofilm and planktonic *B. pertussis* cells to identify key changes between the conditions. Major differences were identified in virulence factors including an upregulation of toxins (adenylate cyclase toxin and dermonecrotic toxin) and downregulation of pertactin and type III secretion system proteins in biofilm cells. To further dissect metabolic pathways that are altered during the biofilm lifestyle, the proteomic data was then incorporated into a genome scale metabolic model using the integrated metabolic analysis tool (iMAT). The analysis revealed that planktonic cells utilised the glyoxylate shunt while biofilm cells completed the full tricarboxylic acid cycle. Differences in processing aspartate, arginine and alanine were identified as well as unique export of valine out of biofilm cells which may have a role in inter-bacterial communication and regulation. Finally, increased polyhydroxybutyrate accumulation and superoxide dismutase activity in biofilm cells may contribute to increased persistence during infection. Taken together, this study modelled major proteomic and metabolic changes that occur in biofilm cells which helps lay the groundwork for further understanding *B. pertussis* pathogenesis.

Introduction

Whooping cough is a re-emerging severe respiratory disease caused by *Bordetella pertussis*. Following the change from the whole cell vaccine (WCV) to the acellular vaccine (ACV) in many developed countries, there has been an increase in the incidence of whooping cough [1-4]. Although most likely multifaceted, waning immunity of the ACV and vaccine driven selection of non-ACV genotypes or strains not expressing one of the ACV antigens have been previously reported as major factors contributing to the re-emergence of pertussis [5, 6]. It is evident that an improved ACV is needed to control the infections.

Recent studies have shown that *B. pertussis* readily forms biofilms *in vivo* [7-10]. Development of the vaccine has been based on planktonic studies and may not be entirely representative of the infection cycle. Although proteomic comparisons have been performed between biofilm and planktonic *B. pertussis* cells [11-13], little is known about the metabolic reactions that are altered while in the biofilm state. In response to changes in environment, the most widely studied regulator of gene expression in *B. pertussis* is the *Bordetella* virulence gene (Bvg) system [14]. The Bvg system controls the expression of most of the virulence factors in *B. pertussis* but has also been implicated in the regulation of metabolism [15]. The Bvg system exists in 3 states, Bvg^+, Bvg^- and Bvg^i where the expression of virulence genes is active, inactive or intermediately expressed, respectively. There are conflicting studies surrounding the role of the Bvg system in the biofilm process, but studies have linked *Bordetella* biofilm with the Bvg^i phase [11]. Further studies identifying key changes in protein expression and metabolism in biofilm cells can provide an insight into the capabilities of the pathogen and help with understanding the role of biofilms in *B. pertussis* pathogenesis.

Genome scale metabolic models (GSMM) have emerged as a powerful tool in understanding the metabolic capabilities of an organism. The creation of a GSMM begins as a draft network based on annotated enzyme data and the genome. This yields a model with a network of reactions and metabolites within a mathematical matrix. The movement of metabolites through the network is defined as flux i.e. the rate at which the metabolites are consumed or produced. Based on stoichiometric and thermodynamic constraints, permissible minimum and maximum flux values for each reaction can be calculated [16].

GSMMs have been successfully utilised in several pathogens such as *Salmonella enterica* serovar Typhimurium (17)*, Listeria monocytogenes* (18, 19)*, Staphylococcus aureus* (20) and *Mycobacterium tuberculosis* (21) to predict important pathways for virulence and growth. Currently, there have been over 6,000 organisms that have been metabolically reconstructed either manually or automatically (22). There have been two GSMMs extensively curated for *B. pertussis* (23, 24). These models showed the metabolic versatility of the organism by identifying minimal nutrient requirements. Additionally, both models were utilised to validate key pathways essential for infection (25). The major drawback of these models is the assumption that all protein products are simultaneously expressed. Many physical and chemical constraints make this assumption unlikely to be true. Additionally, the different conditions in which the organism is grown strongly affects the metabolic processes (26). To predict metabolic reactions reflecting a specific phenotype, such as biofilms, a context specific model should be created. This can be done through the integration of 'omics' expression data into the GSMM to enrich pathways reflective of the context (i.e. biofilm). Integrative Metabolic Analysis Tool (iMAT) (27) is a method that has been developed to incorporate expression data within a metabolic model. The iMAT algorithm uses a mixed integer linear programming (MILP) problem to enrich pathways based on the expression data while maintaining a steady flux distribution and the stoichiometric and thermodynamic constraints. The iMAT model not only creates a phenotype specific model but post-transcriptional regulation or inaccuracies in expression due to experimental limitations can be predicted through this model as the surrounding fluxes would indicate the relative activity of an enzyme in a pathway (27). By comparing the flux distributions of two context specific models, changes in metabolic reactions between phenotypes can be predicted (28).

In this study, proteomic expression data was used to compare biofilm and planktonic *B. pertussis* cells from a representative current circulating strain. Furthermore, the protein expression data was incorporated into a GSMM to create context specific iMAT models to elucidate key metabolic changes that may allow biofilm cells to persist in the host.

Methods

Bacterial strains and biofilm growth

A clinical *Bordetella pertussis* strain, L1423 isolated from the 2008-2012 Australian epidemic with genotype *ptxP3/ptxA1/fim3A/prn2*, was used as a representative of the

predominant cluster I strains. The genome has been previously sequenced, and the strain has

been used in two separate infection studies in mice (29, 30). *B. pertussis* planktonic and

biofilm cells were grown using previously established methods and proteins were extracted

(31). Briefly, the *B. pertussis* strain was grown on Bordet-Gengou agar (BG, BD Scientific)

for 3-5 days at 37°C. A loopful of pure Bvg$^+$ colonies were suspended into Thalen-Ijssel

(THIJS) media (32) supplemented with 1% heptakis ((2,6-O-dimethyl) β-cyclodextrin) and

1% THIJS supplement and grown for 24 h shaking at 180 rpm at 37°C. For planktonic

growth, the OD_{600} of the starter culture was adjusted to 0.05 in 20 mL THIJS and incubated

for 12 h, shaking at 180 rpm. For biofilms, the OD_{600} was adjusted to 0.1 and 1 mL of this

adjusted culture was seeded into each well of a 24 well plate. The 24 well plate was incubated

statically for 5 h at 37°C for attachment of cells before the media was refreshed to remove

non-adherent cells (11). After 96 h of incubation, the wells were washed with PBS and then

the plate was water bath sonicated at 37 kHz for 2 min to detach cells. The planktonic and

biofilm cells were then probe sonicated and proteins extracted as described by Luu et al. (31).

Six biological replicates per condition were performed.

Confocal laser scanning microscopy analysis

To confirm biofilm maturity, confocal laser scanning microscopy (CLSM) was used.

This method was adapted from Cattelan et al. (33). The *B. pertussis* cells were grown on a

glass coverslip angled at 45° in the same method as described above. The biofilm was imaged

at 24, 48, 72 and 96 h. The biofilm was fixed with 4% paraformaldehyde and stained with

SYTO9 (Thermo Fisher Scientific) fluorescent dye. The coverslips were imaged on the

FluoView FV1200 inverted confocal microscope (Olympus Life Sciences) at the UNSW

Katharina Gaus Light Microscopy Facility (KG-LMF). Three biological replicates were

performed per time point and 3 field of views per replicate were randomly selected for z-

stack 3D imaging. The biomass was calculated using the COMSTAT2 ImageJ plugin (34).

Protein preparation and LC-MS/MS

The protein extracts (10 μg) were reduced with dithiothreitol, alkylated with

iodoacetamide and then digested with trypsin as described in Luu et al. (35). The peptides

were analysed on the LTQ-Orbitrap Velos mass spectrometer (Thermo Fisher Scientific) at

the UNSW Bioanalytical Mass Spectrometry Facility (BMSF) with the settings described in

Luu et al. (35). The output spectra were matched against a custom *B. pertussis* database on

the MaxQuant (v2.0.3.1) proteomics software with the following parameters: digestion mode

– specific, enzyme – Trypsin/P, max missed cleavages – 1, Label free quantification – LFQ,

Protein identification false discovery rate – 0.01 and min peptides per protein – 2. All other

parameters were set as the recommended default values. Student's *t*-test was calculated and a

false discovery rate (FDR) *q*-value multiple test correction was performed using the Storey-

Tibshirani method on R (v4.1.1) (36). Proteins were considered upregulated if the fold

change (FC) was > 1.2, *q* < 0.05 and downregulated if FC < 0.8, *q* < 0.05 based on previous

studies (31). Functional categories were assigned to proteins based on Bart et al. (37). The

mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium

via the PRIDE (38) partner repository with the dataset identifier PXD033664 and

10.6019/PXD033664.

Integrated Metabolic Analysis Tool (iMAT) model generation

The proteomic expression data was used to generate context specific metabolic

models for the planktonic and biofilm cells. The iMAT (39) method, available in the COBRA

Toolbox (v3.0) (40), was used. Processing was done in MATLAB (R2020a) using the IBM

CPLEX optimiser (v12.10.0). The iMAT algorithm extracts a simplified model based on the

trade-off between high expression and low expression reactions. The comprehensive,

manually curated *B. pertussis* metabolic model (iBP1870) generated by Branco dos Santos et

al. (23) was used as the base model. Subsystems from the *Escherichia coli* genome scale

metabolic model (iAF1260) were assigned to each of the reactions in the *B. pertussis* model

(41). The *E. coli* (iAF1260) model was used as a template for the original *B. pertussis* model

(iBP1870) and therefore, most of the subsystems are transferrable. The original *B. pertussis*

iBP1870 model had the tricarboxylic acid cycle partially dysfunctional (no flux from

oxaloacetate to α-ketoglutarate), as this model was based on previous studies which stated *B.*

pertussis has an incomplete TCA cycle (32). However, recent studies have shown that the

TCA cycle is fully functional and therefore the disabled pathways were activated before

generating the iMAT models (42). After refining the *B. pertussis* model, the biofilm and

planktonic proteins were assigned to the gene-protein-reaction (GPR) associations listed in

the *B. pertussis* model (iBP1870).

Label free quantification (LFQ) intensity values designated by the MaxQuant

software were used to estimate relative protein abundance. To increase confidence in the

imported data, only proteins identified in all six biological replicates per condition were

incorporated. Proteins were designated as uniquely identified if found in all 6 biological

replicates in one condition and none in the other. The expression data was defined as highly

(1), lowly (-1) or moderately (0) expressed based on the threshold of mean expression $\pm$ 0.5 x

STD of the proteins included in the model (39). After generating the planktonic and biofilm

context specific iMAT models, the reactions in the biofilm and planktonic models were

compared (Figure 1).

Additional infection specific models were generated with proteomic expression data

from two separate studies. The proteomic study by de Gouw et al. (11) compared *B. pertussis*

biofilm and planktonic (mid-log and stationery) cytosolic and membrane proteins. The

averaged expression values of the cytosolic and membrane fractionated proteins were

combined, and mid-log planktonic expression values were used to generate the models. The

study by Lamberti et al. (43) compared the proteomes of *B. pertussis* cells extracted from

within macrophages at 3 h and 48 h and planktonic cells. iMAT models were generated for

all three conditions. All models have been deposited to the BioModels database (44) in

SBML L3V1 format under the model identifier MODEL2205270001. Differences between

the growth conditions are listed in Supplementary Table S1.

Flux Analysis

A flux variability analysis (FVA) was performed on the models which provides

minimum and maximum permissible flux bounds for each reaction. Flux is recorded as a

millimolar concentration of metabolite per gram of dry cell weight per hour (mmol $\cdot$ g_{DCW}^{-1} $\cdot$

h^{-1}). The flux bounds in common reactions between the biofilm and planktonic models were

compared using the Jaccard index to calculate the similarity between the two models for each

reaction. The FVA is also important to determine which direction the reaction runs in the

model. Furthermore, to identify the most important reactions in each model, each reaction

was deleted in turn and an FVA calculation was performed to deem whether the model would

still be feasible without the reaction.

A flux balance analysis (FBA) was performed to identify changes in flux in reactions

shared by the planktonic and biofilm models. The FBA provides a single value for each

reaction based on a predefined metabolic goal so that the models may be compared. The FBA

method has been extensively developed to predict metabolic fluxes within metabolic models

(16). Typically, the objective function (*c*) of the linear optimisation equation is set as an

artificial biomass reaction as the metabolic goal of the organism. As generating biomass may

not be the metabolic objective of biofilm cells, an alternate method for defining the objective

function was used in this study. The objective function was defined based on proteomic

expression data as described in Montezano et al. (45). For all proteins that were identified

with GPR associations in the model, the intensities were normalised by the maximum

intensity value for each condition (planktonic and biofilm) and these values were input as c.

This leads the FBA to push flux towards the reactions which have higher protein expression

and considerably shrinks the solution space to increase prediction accuracy (45).

Results

Key proteomic changes identified between biofilm and planktonic cells

To identify changes that occur in biofilm conditions, label free quantification mass

spectrometry (LFQ-MS) was performed on biofilm and planktonic cells. Analysis was

performed on L1423, a clinical isolate representative of the current circulating *B. pertussis*

strains. Confocal microscopy confirmed biofilm formation and mature structure at 96 h

(Supplementary Figure S1). Furthermore, there was a polysaccharide biosynthesis protein,

WbpO (FC = 2.98, q = 1.17E-5) and a phosphoglucomutase enzyme, pgm (FC = 1.23, q =

0.041), that were seen to be upregulated in biofilm cells. These proteins have been previously

linked with *B. pertussis* biofilm reinforcing the biofilm phenotype achieved in this study (46).

There were 948 proteins identified in total, of which, 571 were proteins identified in all 6

biological replicates (Figure 2A). There were 478 proteins with significantly differential

expression (q < 0.05) between the two conditions. In biofilm cells, there were 242 proteins

downregulated and 236 proteins upregulated (Figure 2B).

There were many proteins related to virulence with altered expression identified in

this study (Table 1). BipA, an outer membrane protein associated with biofilm formation was

upregulated (11). Adenylate cyclase toxin (CyaA) and dermonecrotic toxin (Dnt) were also

upregulated in biofilm. There were 10 proteins from the Type III secretion system (T3SS)

that were downregulated in biofilm cells or uniquely identified in planktonic cells. One

protein (Bcr4) involved in the T3SS was upregulated in biofilm cells (47). Tracheal

colonisation factor (TcfA) and virulence associated gene 8 (Vag8) were downregulated in

biofilm cells while *Bordetella* resistance to killing (BrkA) protein was upregulated. Of the

ACV antigens, there was no change in expression for filamentous haemagglutinin (FhaB) but

the filamentous haemagglutinin outer membrane transporter protein (FhaC) was uniquely identified in biofilm cells. The membrane bound pertussis toxin subunit 4 (PtxD) was also uniquely identified in biofilm cells, however, pertussis toxin subunit 1 (PtxA) was downregulated in biofilms. Furthermore, fimbriae protein (Fim2) was downregulated in biofilm. Finally, pertactin (Prn) was strongly downregulated in biofilm (FC = 0.16, q = 2.42E-6). These changes demonstrate a strongly altered virulence profile in *B. pertussis* biofilm cells compared to planktonic cells.

When the proteins were grouped into functional categories based on Bart et al. (37), there was a significant increase in proteins in the categories of transport/binding and miscellaneous proteins under biofilm conditions. There was also a downregulation of the functional groups: ribosome constituents and cell processes (Figure 3).

iMAT model generation

For an in-depth analysis of the metabolic changes between biofilm and planktonic cells, iMAT metabolic models were generated based on protein expression from both conditions. Of the 571 proteins identified using mass spectrometry, 228 were annotated with GPR associations in the *B. pertussis* (iBP1870) model (23). These are proteins which have assigned reactions in the base model. The overlap between the identified proteins and the reactions in the model constitutes 29.61% of the total GPR associations listed in the base model. Context specific models were created using the iMAT algorithm implemented in the COBRA Toolbox (40). The biofilm iMAT model consisted of 198 metabolites, 206 reactions and 188 genes. The planktonic iMAT model had 213 metabolites, 219 reactions and 195 genes (Table 2). To identify changes between the planktonic and biofilm iMAT models, the number of unique and common reactions between the two models were compared. There were 168 reactions that were common between the two models, 51 reactions that were unique to the planktonic model and 38 that were unique to the biofilm model (Figure 4A). The reactions were grouped into subsystems and total number of reactions in each group were similar between the two models (Figure 4B). It is notable however that there was a high proportion of reactions that were unique to the individual models.

B. pertussis **biofilm cells complete the tricarboxylic acid cycle**

An FBA was performed to reveal changes in flux between reactions that were common between the planktonic and biofilm models. A major difference was seen in the

tricarboxylic acid (TCA) cycle. The biofilm cells pushed flux to complete the TCA cycle

while the planktonic cells pushed flux towards the glyoxylate shunt (Figure 5). While there

was slightly higher levels of flux from oxaloacetate through to isocitrate in the planktonic

model (planktonic: 10.40 mmol $\cdot$ g_{DCW}^{-1} $\cdot$ h^{-1}, biofilm: 8.92 mmol $\cdot$ g_{DCW}^{-1} $\cdot$ h^{-1}), the

planktonic cells pushed all flux (11.31 mmol $\cdot$ g_{DCW}^{-1} $\cdot$ h^{-1}) into the glyoxylate shunt to

convert isocitrate to glyoxylate and succinate while comparatively, only a small flux (0.57

mmol $\cdot$ g_{DCW}^{-1} $\cdot$ h^{-1}) was pushed for the same reactions for biofilm cells. Instead, the flux

moved towards α-ketoglutarate (AKG) to succinyl-CoA and succinate to complete the TCA

cycle (Figure 5). The planktonic cells comparatively had no flux through these reactions.

Despite these changes, there was an increased level of flux from succinate to fumarate and

malate in the planktonic model which was fed from the glyoxylate shunt, branched chain

amino acid degradation and the arginine biosynthesis cycle. Finally, there was also increased

conversion of pyruvate to acetyl-CoA in the planktonic model which was then fed back into

the TCA cycle as citrate. Most of the acetyl-CoA synthesis/utilisation pathways were either

unique to the planktonic model or downregulated in the biofilm model. However, the biofilm

model had unique reactions to convert acetyl-CoA to polyhydroxybutyrate (PHB).

Cardiolipin synthesis active in planktonic cells

There were many reactions included in both models that were grouped into the

glycerophospholipid metabolism subsystem. Flux moves through the gluconeogenesis

pathway in both the models from pyruvate through to dihydroxyacetone phosphate and then

glycerol 3-phosphate (Figure 5). The pathways for most of the glycerophospholipids and fatty

acid synthesis were the same between the two models. However, the planktonic model had

pathways producing cardiolipin (tetradodecanoyl, n-C12:0) while the biofilm model lacked

these pathways. The production of cardiolipin is through the pathway of conversion of 1,2-

didodecanoyl-sn-glycero-3-cytidine 5'-diphosphate (CDP-DAG) to

phosphatidylglycerophosphate (didodecanoyl, n-C12:0) and therefore there is a higher level

of flux from the cytidine monophosphate (CMP) to cytidine triphosphate (CTP) in the

planktonic model. The models suggest that there would be a higher level of 1-dodecanoyl-*sn*-

glycerol 3-phosphate, 1-hexadec-9-enoyl-*sn*-glycerol 3-phosphate and glycerol 3-phosphate

in the periplasm of the biofilm cells. In the planktonic cells, there was a higher level of

cardiolipin, phosphatidylglycerol (didodecanoyl, n-C12:0), 2-dodecanoyl-*sn*-glycerol 3-

phosphate and 2-hexadec-9-enoyl-*sn*-glycerol 3-phosphate. The biofilm model uniquely

exported glycerol 3-phosphate into the extracellular space through these pathways while

planktonic model uniquely exported glycerol into the extracellular space (Figure 5).

Arginine biosynthesis decreased in biofilm cells

The metabolic models also revealed many altered amino acid metabolism pathways.

There was an equal number of reactions grouped into the arginine and proline metabolism

subsystem between the two models. However, the processes within the pathways were varied

as seen by the number of unique reactions within the subsystem (Figure 4B). Both models

feed into the arginine pathways through conversion of L-glutamate (GLU) to L-glutamine.

Subsequently, both models have reactions to produce carbamoyl phosphate (CBP). While the

planktonic model pushed the CBP into the arginine biosynthesis pathway, the biofilm model

had reactions to convert CBP through to orotate which was then exported out of the cell. The

planktonic cells complete the arginine biosynthesis cycle, exporting urea out of the cell as a

by-product. Furthermore, in the planktonic model, arginine succinate is converted to arginine

and fumarate which is fed back into the TCA cycle (Figure 5). When each of these reactions

were deleted in turn and tested for model feasibility, it led to infeasible models for the

planktonic model but were still feasible for the biofilm model (Supplementary Table S2).

Both models produced ornithine surrounding the arginine metabolism pathways. The

planktonic cells produced ornithine through the arginine biosynthesis pathway while the

biofilm cells synthesised ornithine through AKG to acetyl-ornithine and then to ornithine.

The fate of ornithine also differed greatly between the two models. When the FVA values

were compared, the smallest Jaccard index value (lowest similarity) between the models were

for the reactions acetylornithine transaminase (R_ACOTA) and glutamate N-acetyl

transferase (R_ORNTA). R_ACOTA is a reversible reaction of acetyl-ornithine and AKG to

N-acetyl-L-glutamate 5-semialdehyde and GLU. The FVA revealed that the biofilm model

only had capabilities to run this reaction in reverse. The opposite was seen for R_ORNTA, a

reversible reaction of acetyl-ornithine and GLU to ornithine and N-acetyl-L-glutamate. The

planktonic model ran this reaction in reverse. Overall, the ornithine in the planktonic model

was either utilised to produce acetate or pushed back through the arginine biosynthesis

pathway. The biofilm model pushed ornithine through the butanoate metabolism pathway

into the TCA cycle as succinate with the production of NADPH and NADH. This process

utilised GLU and produced AKG.

Aspartate metabolism downregulated in biofilm cells

Aspartate that was fed into the arginine biosynthesis pathway was decreased in biofilm cells. Furthermore, when the reaction was deleted in the planktonic model, it led to an infeasible model while the biofilm model remained feasible without the reaction. The flux in the biofilm model pushed from aspartate through to threonine synthesis and CBP metabolism. Threonine was then degraded through the branched chain amino acid (BCAA) degradation pathway. Additionally, aspartate was uniquely converted with CBP to N-carbamoyl-L-aspartate and ultimately into orotate as mentioned above for the biofilm model. The planktonic model had flux flow from aspartate to homoserine, but rather than following through to threonine, the homoserine was converted to O-acetyl-L-homoserine towards acetate and L-cystathionine (Figure 5). L-cystathionine was then converted to L-cysteine and 2-oxobutanoate which was fed through the BCAA (specifically the isoleucine pathway) degradation pathway that was upregulated in the planktonic model. The BCAA degradation pathway leads to the production of acetyl-CoA, NADH, NADPH and $FADH_2$. There was further production of these molecules through the valine degradation pathway in the planktonic model. These pathways lead flux being moved back into the TCA cycle as succinate and pyruvate through propanoate metabolism. Although the planktonic model moved flux through the valine degradation pathway, the actual production of valine was unique to the biofilm model. The valine was either exported out of the cell or converted to AKG in the biofilm model.

Cysteine was transported into the cell at the same rate between the two models. Both models source cysteine through diffusion, active ABC transport and the conversion of glutathione and L-cysteinylglycine. The biofilm model pushed all flux of cysteine to L-alanine while the planktonic model also has pathways to convert the cysteine through to acetyl-CoA as mentioned above. The reactions to move L-alanine in and out of the periplasm were unique to the biofilm model. The main source of L-alanine for the biofilm cells was from the conversion of pyruvate and glutamine to L-alanine and AKG. The planktonic model creates L-alanine through the conversion of β-alanine. This reaction also creates malonate-semialdehyde and was strongly downregulated in the biofilm model. The L-alanine is converted to D-alanine and then to GLU and pyruvate in both models.

Biofilm cells have increased superoxide dismutase activity

Both models converted ubiquinol to ubiquinone through cytochrome ubiquinol oxidase. This reaction was downregulated in biofilm model. However, the biofilm model also had an additional reaction that converted ubiquinol to ubiquinone with the by-product of superoxide anions. The superoxide was converted to hydrogen peroxide and O_2 by superoxide dismutase. The hydrogen peroxide was converted to H_2O by thioredoxin. The other reactions involving ubiquinone were downregulated in biofilm cells including succinate dehydrogenase and NADH dehydrogenase. However, ubiquinone was converted to ubiquinol through the conversion S-dihydroorotate to orotate in a reaction unique to the biofilm cells.

Comparison with other metabolic models

We further used proteomics data from two previous *B. pertussis* studies to create iMAT models and compare the reactions. The first study performed by de Gouw et al. (11) grew *B. pertussis* biofilms on flat polypropylene beads in a glass column reactor for 72 h with THIJS media refreshed every 24 h. The planktonic cells were extracted at both mid-exponential phase at 17 h and stationary phase at 40 h. That study identified 729 – 825 proteins from the three different conditions with an overlap of 645 proteins (11). While the biofilm iMAT model generated using that proteomic data had a higher number of reactions, metabolites and genes to the planktonic iMAT model generated from that same data, the overall values were comparable to the models of this study (Supplementary Table S1). The second study that was compared was proteomic data from *B. pertussis* cells isolated from within macrophages performed by Lamberti et al. (43). *B. pertussis* cells were extracted from within macrophages at both 3 h and 48 h after infection in addition to a planktonic counterpart grown in Roslyn Park Memorial Institute (RPMI)-1640-BSA 0.2% (*w/v*). A total of 762 proteins were identified with 300 with differential expression between the conditions (43). iMAT models were generated for these conditions and compared (Supplementary Table S1). As with the additional models generated from the study by de Gouw et al. (11), the three models generated by the Lamberti et al. (43) intracellular macrophage study yielded similar numbers of reactions, metabolites and genes as with this present study

In line with our study, the de Gouw et al. (11) biofilm model completed the full TCA cycle while the planktonic model pushed flux through the glyoxylate shunt. A similar trend was identified for the models created from the Lamberti et al. (43) intracellular macrophage study. The planktonic model showed the same preference for the glyoxylate shunt and the *B. pertussis* cells extracted from the macrophages at 3 h had a similar flux through the TCA

cycle as the biofilm models, however, the 48 h model pushed flux through the glyoxylate

shunt.

Further similarities were identified between the intracellular macrophage and biofilm

models in NADH dehydrogenase, aspartate transaminase and CO_2 and O_2 exchange. All

intracellular macrophage and biofilm models had decreased flux in these reactions compared

to their planktonic counterpart. The NADH dehydrogenase reaction produces NAD+ from

NADH with the conversion of ubiquinone to ubiquinol. The aspartate transaminase reaction

(R_ASPTA - reversed) had decreased flux for all intracellular macrophage and biofilm

models. This reaction converted GLU and oxaloacetate to AKG and aspartate. The reaction

from aspartate to CBP and subsequent reactions to orotate were unique to the biofilm models

(Supplementary Table S2).

Discussion

Recent studies have shown that biofilms are an important aspect of *B. pertussis*

pathogenesis (33, 48, 49). Biofilm cells are more resilient against antimicrobials and

environmental stresses (12). While previous studies of *B. pertussis* biofilms have made

important discoveries related to growth and virulence, there has been less research focused on

biofilm metabolism. Biofilms are readily formed by *B. pertussis* and has been defined as an

integral part of its pathogenesis (7). Investigating the metabolic changes that occur within a

biofilm community may help understand the pathogenesis of *B. pertussis*. Therefore, this

study integrated proteomic expression data from a currently circulating epidemic strain into a

metabolic model of *B. pertussis* to identify major changes that occur in metabolism between

the biofilm and planktonic states. To our knowledge this is the first extensive study into

specific metabolic pathways within biofilms of *B. pertussis*. Although utilised metabolic

pathways differ from species to species, many of the changes in this study have been

identified and confirmed experimentally in other species reinforcing the strength of the

metabolic models. The major metabolic differences identified in this study relate to the TCA

cycle, amino acid metabolism and virulence.

The TCA cycle is the major central metabolic pathway for many aerobic organisms,

therefore it was surprising to find that there was altered flux for the TCA cycle. There was a

major shift in planktonic cells to utilise the glyoxylate shunt instead of completing the full

TCA cycle. The TCA reactions with increased flux in biofilms leads to the production of NADH, NADPH and ATP. Variation in the glyoxylate shunt has been observed in the biofilms of other species. When the glyoxylate shunt was disabled in *Pseudomonas aeruginosa,* there was an increase in biofilm formation (50), which is reflected in this study. It was suggested that the increased extracellular polymeric substances (EPS) produced when the glyoxylate shunt was disabled could lead to higher survival in the microaerobic conditions of the cystic fibrosis lung environment (50). Additionally, there was an increased level of glyoxylate activity in *Candida albicans* cells dispersed from biofilm (51). It was suggested that as the glyoxylate shunt is activated to increase nutrient versatility, it may be an anticipatory reaction for low nutrient levels while searching for a new colonisation location (51). The dispersed cells may reflect a planktonic lifestyle while established mature biofilm cells may utilise the network of cells to share nutrients and hence have a decreased requirement of the glyoxylate shunt. Targeting the TCA cycle has been suggested as a potential therapeutic strategy against biofilms (52). Additional models that were created using *B. pertussis* biofilm and planktonic proteomic expression data from de Gouw et al. (11) had similar differences in the TCA cycle to our results (Supplementary Figure S2).

The glyoxylate shunt was also identified as important in intracellular macrophage survival. Additional iMAT models created with *B. pertussis* proteomic expression data from Lamberti et al. (43) showed that the glyoxylate shunt was inactive at 3 h inside macrophages but active at 48 h (Supplementary Figure S2). This is reflective of previous studies that have shown that inactivation of the glyoxylate shunt led to decreased macrophage intracellular survival (53, 54). The differences seen may be due to the need to scavenge nutrients in the hostile environment and the time it takes for the bacteria to adapt to the new environment. The additional models had limited altered pathways reflecting the other significant changes identified in this study which may be due to the many differences between conditions (Supplementary Table S1). However, it is noteworthy that the same pattern was seen with the glyoxylate shunt in the biofilm and intracellular macrophage models compared to the planktonic cells. This highlights the importance of the glyoxylate shunt as a key determinant of virulence during *B. pertussis* infection.

Increased polyhydroxybutyrate (PHB) and superoxide dismutase activity may explain the increased survivability of *B. pertussis* biofilm cells. Reactions regarding PHB synthesis were identified as unique in the biofilm model and superoxide dismutase activity was

increased in the biofilm model. While most reactions surrounding acetyl-CoA were

downregulated in the biofilm model or unique in the planktonic model, one reaction that

involved the production of PHB was unique in biofilm. The PHB reactions led to an increase

in cytoplasmic PHB demand. It has been previously reported that cytoplasmic PHB

inclusions exist in *B. pertussis* cells. These inclusions are used as an energy reserve for when

cells are in harsh environments (32). As the PHB demand reactions are increased in the

biofilm cells, this may lead to increased cell survivability that has been observed in *B.*

pertussis biofilms (12). Furthermore, an increase in the superoxide dismutase activity is

protective as it leads to a decrease in superoxide which, as a free radical, may lead to cellular

damage (55). Put together, these two factors may be a reason for increased cell survivability

of *B. pertussis* biofilm cells and may lead to persistent infections.

Many pathways for glycerophospholipid metabolism were seen to be active in both

metabolic models. Lipid remodelling can occur when cells are exposed to harsh environments

or antimicrobials (56). The changes in the lipids have been linked to increased survivability

and persistence in pathogenic bacterial biofilms (57). The models in this study suggest that

there would be differing levels of certain glycerophospholipids in the periplasm between

planktonic and biofilm cells. Of particular interest is cardiolipin, identified as a unique lipid

synthesis pathway in the planktonic model. Previous studies have shown that cardiolipin

depletion led to decreased biofilm in *E. coli* (58). Cardiolipins have been identified in *B.*

pertussis previously however, studies have not remarked on biofilm cells (59). It is important

to note that the proteomic data captures enzymes that are active at a particular time point

rather than a measurement of lipids, therefore, the changes identified in these reactions

merely provide a snapshot of the time from which the proteins were harvested. The

planktonic cells may continuously synthesise cardiolipin while mature biofilm cells either

synthesise cardiolipin at an earlier stage of biofilm or have a lower requirement for

cardiolipin. To further investigate the changes in lipid composition, future lipidomic analysis

of biofilm cells would be necessary. Nevertheless, the evidence from the metabolic models in

this study suggest differing lipid composition in biofilm and planktonic *B. pertussis* cell

membranes. Another interesting change that was linked to the cardiolipin pathway was the

exportation of glycerol out of the cell. While the planktonic cells exported glycerol, the

biofilm cells exported glycerol 3-phosphate. The reason for this is unexplored and it would be

interesting to identify whether varying concentrations of glycerol 3-phosphate or glycerol

would have a regulatory role in *B. pertussis*.

Many regulatory processes are involved in *B. pertussis* biofilm formation (60). This study identified multiple amino acid processing pathways that varied between biofilm and planktonic cells. It has been reported previously that amino acids and their intermediates can act as regulatory molecules (61-63). The planktonic cells showed increased arginine biosynthesis and aspartate transaminase activity. It has been shown in other species that arginine may increase biofilm formation although higher concentrations have an inhibitory effect (62) and increase antibiotic killing (64). In addition, arginine may disrupt bacterial coaggregation to prevent biofilm formation (65). Finally, the arginine biosynthesis process leads to urea by-products which have been previously shown to disrupt the EPS matrix of biofilms (66). The evidence in these studies combined with results from this present study points to a tightly regulated arginine synthesis pathway which is partially fed by the aspartate transaminase reaction. While the planktonic model pushed aspartate to the arginine biosynthesis pathway, the biofilm cells pushed aspartate to the threonine degradation pathway and towards the synthesis of N-carbamoyl-L-aspartate (through to orotate) leading to an overall reduction of available arginine. The orotate was uniquely exported out of the biofilm cells rather than processed for pyrimidine metabolism. Orotate therefore has the potential to act as a signalling molecule for *B. pertussis* biofilm cells. The pathways regarding carbamoyl phosphate synthesis which leads to orotate synthesis has been implicated in biofilm formation in other species (67, 68). This orotate production pathway was specific to both the biofilm model in this study and the biofilm model created from the data of de Gouw et al. (11), reinforcing the possible importance of orotate export for *B. pertussis* biofilms.

While orotate metabolism may be specific to *B. pertussis* biofilm models, other aspartate processing changes identified may be specific to broader infection environments. Aspartate conversion from AKG (aspartate transaminase reaction) was seen to be a downregulated pathway in all intracellular macrophage and biofilm metabolic models compared to their planktonic counterparts. Aspartate feeds directly into the arginine biosynthesis pathway as mentioned previously. Aspartate has been seen to inhibit biofilm formation in multiple *Staphylococcus* species (69). Furthermore, the specific aspartate transaminase reaction was downregulated in *Streptococcus pneumoniae* biofilms (70). Interestingly, in *Mycobacterium tuberculosis* it was shown that the aspartate pathways are important for intracellular survival in macrophages. It was suggested that intracellular *M. tuberculosis* requires exogenous aspartate for intracellular survival and that it is a potential target for novel therapeutics (71). The intracellular models generated in this study suggest

that exogenous aspartate scavenging is not as important for *B. pertussis* intracellular survival.

However, as it is important for the planktonic model, it may still provide a potential target to

prevent initial colonisation.

Finally, there was a change in the processing of alanine and valine between the

biofilm and planktonic models. The models suggest that β-alanine is important for planktonic

cells while L-alanine is more vital for biofilm cells. The biofilm model had unique reactions

transporting L-alanine into the periplasm. A recent study showed that alanine metabolism

activity displays a unique pattern based on spatial distribution of cells within a heterogenous

biofilm (72). The alanine metabolism changes identified in the present study may relate to a

particular layer of the biofilm. It would be interesting to see the specific active reactions

between the layers of biofilm. Proteomic analysis on layers of biofilm have previously been

performed using laser ablation sample transfer (LAST) and label free mass spectrometry (73).

A combination of LAST and the methods used in this study may be able to confirm the

changes in alanine metabolism. Furthermore, valine was uniquely exported out of the biofilm

model. This confirms what has been observed in other Gram-negative species that had

increased valine secretion in biofilm conditions (74). It was suggested that within the low

oxygen and reduced growth rate conditions of biofilms, there would be an excess of pyruvate

leading to an increase in valine and alanine biosynthesis (74). The discussed changes in

amino acid metabolism pathways are biofilm specific and therefore it is possible that the

secreted molecules could work as signalling molecules potentially triggering biofilm

formation.

Major expression differences were also identified for virulence factors. The Bvg

system has been shown to regulate most of the virulence factors in *B. pertussis* (75). In

addition, there are also secondary regulators that alter virulence factor expression in biofilm

cells (76). This study identified increased expression of the BvgA regulator and BvgS in

biofilms while the expression of BvgR was similar between the two conditions (Table 1).

Unexpectedly, there was a wide variability in protein expression within Bvg regulated

proteins. For example, Dnt and CyaA were upregulated while many proteins of the type III

secretion system (T3SS) and Prn were found to be downregulated in biofilm cells. To our

knowledge, this is the first report of a major downregulation of Prn in *B. pertussis* biofilm

cells. As one of the three ACV components, the change identified supports the hypothesis

that the planktonic based ACV may not protect as efficiently against cells in the biofilm

condition. To further reinforce this idea, the biofilm membrane derived protein that has been

suggested as a potential new addition to the vaccine, BipA was identified as upregulated in

biofilm cells in this study (11). Upregulation of BipA also underscores the importance of the

Bvg^i phase in the biofilm existence. There were also changes in both Bvg^+ and Bvg^- protein

regulation. Within both the upregulated and downregulated proteins, 13-15% of the proteins

were Bvg^+. Furthermore, there was a significantly higher proportion (14%) of proteins

upregulated in biofilms that were Bvg^- compared to the proportion (4%) of proteins

downregulated that were Bvg^- (Fishers exact test, $p = 0.0002$). The change in the expression

of these Bvg regulated proteins may be related to the 8 regulatory proteins with differential

expression in biofilm cells (Supplementary Table S2). The findings in this study point to

other major regulatory systems that affect the regulation of virulence proteins in biofilm

conditions.

The flux balance analysis identified reactions that have increased or decreased flux

within the models. There was a number of reactions with $0 \text{ mmol} \cdot g_{DCW}^{-1} \cdot h^{-1}$ flux. This is a

common issue that occurs with metabolic models (45). This is understandable as when the

objective function is set relative to the expression data, the mixed integer linear programming

calculation prioritises the reactions in the models with increased protein expression. This

leads to reactions that were not identified in the proteomic data recorded under the minimum

value, which is often $0 \text{ mmol} \cdot g_{DCW}^{-1} \cdot h^{-1}$ as revealed by the FVA flux bounds. Overall, the

$0 \text{ mmol} \cdot g_{DCW}^{-1} \cdot h^{-1}$ flux reactions may lead to decreased accuracy in the mM

measurements of flux from the FBA but it does not invalidate the relative flux through

pathways when they are compared between the models. Additionally, the unique reactions

with $0 \text{ mmol} \cdot g_{DCW}^{-1} \cdot h^{-1}$ fluxes are still important for that model as designated by the

iMAT algorithm.

Traditional analysis of proteomic data involves grouping proteins into broad

functional categories to identify major changes, but it lacks the precision to examine specific

changes in metabolism. The incorporation of the proteomic data in the GSMM in this study

provided new insights into the metabolic pathways that were altered between biofilm and

planktonic cells. It is important to note that the biofilm cells were grown in an artificial media

designed to optimise *B. pertussis* growth. It would be interesting to see how growth in

different media such as media more representative of the respiratory environment or co-

culture with epithelial cells would affect the metabolic model (77). Additionally, it has been

shown that there is a wide range of heterogeneity of cells in a biofilm (78). This is not
surprising as cells within the biofilm would be exposed to varying levels of nutrient gradients
(79). As it is, the model in this study would have taken an average of overall expression and
there may have been important differences between the layers of biofilm that could improve
our understanding of the overall biofilm process. Additionally, it would be interesting to
create temporal metabolic models to identify the changes that occur throughout biofilm
development (80).

In conclusion, this study demonstrated major metabolic changes that occur during
biofilm conditions in *B. pertussis*. Many similar changes in biofilm pathways from other
species were also found in *B. pertussis,* reinforcing the utility of integrated expression data
into metabolic modelling. Notably, it was found that biofilm cells utilised the full TCA cycle
while planktonic cells pushed flux through the glyoxylate shunt. There was an increase in
PHB accumulation and superoxide dismutase activity which may lead to increased
persistence of biofilm cells. Major changes were identified in amino acid pathways including
exportation of valine out of biofilm cells which may have a role in cell-to-cell
communication. Finally, there were changes in virulence factor expression including an
increase in Dnt and CyaA and a strong downregulation of T3SS and Prn in biofilm cells.
Overall, the changes identified in this study helps lay the groundwork for further studies into
B. pertussis biofilms and their role in pathogenesis.

References

1.   Galanis, E., A. S. King, P. Varughese, and S. A. Halperin. 2006. Changing epidemiology and
emerging risk groups for pertussis. *Can. Med. Assoc. J.* 174: 451-452.
2.   de Melker, H. E., M. A. Conyn-van Spaendonck, H. C. Rümke, J. K. van Wijngaarden, F. R.
Mooi, and J. F. Schellekens. 1997. Pertussis in The Netherlands: an outbreak despite high
levels of immunization with whole-cell vaccine. *Emerg. Infect. Dis.* 3: 175-178.
3.   Güriş, D., P. M. Strebel, B. Bardenheier, M. Brennan, R. Tachdjian, E. Finch, M. Wharton,
and J. R. Livengood. 1999. Changing epidemiology of pertussis in the United States:
increasing reported incidence among adolescents and adults, 1990-1996. *Clin. Infect. Dis.* 28:
1230-1237.
4.   Campbell, P. T., J. M. McCaw, P. McIntyre, and J. McVernon. 2015. Defining long-term
drivers of pertussis resurgence, and optimal vaccine control strategies. *Vaccine* 33: 5794-
5800.
5.   Octavia, S., R. P. Maharjan, V. Sintchenko, G. Stevenson, P. R. Reeves, G. L. Gilbert, and R.
Lan. 2011. Insight into Evolution of *Bordetella pertussis* from Comparative Genomic
Analysis: Evidence of Vaccine-Driven Selection. *Molecular Biology and Evolution* 28: 707-
715.
6.   Mooi, F. R., N. A. Van Der Maas, and H. E. De Melker. 2014. Pertussis resurgence: waning
immunity and pathogen adaptation - two sides of the same coin. *Epidemiology and infection*
142: 685-694.
7.   Conover, M. S., G. P. Sloan, C. F. Love, N. Sukumar, and R. Deora. 2010. The Bps
polysaccharide of *Bordetella pertussis* promotes colonization and biofilm formation in the
nose by functioning as an adhesin. *Mol. Microbiol.* 77: 1439-1455.
8.   Goldsmith, C. S., J. Guarner, J. L. Montague, M. T. Eliason, P. W. Greer, S. R. Zaki, W.-J.
Shieh, C. D. Paddock, R. C. Holman, G. N. Sanden, K.-H. Wu, K. M. Tatti, A. A. Gal, J. D.
Cherry, and C. Langston. 2008. Pathology and Pathogenesis of Fatal *Bordetella pertussis*
Infection in Infants. *Clin. Infect. Dis.* 47: 328-338.
9.   Soane, M. C., A. Jackson, D. Maskell, A. Allen, P. Keig, A. Dewar, G. Dougan, and R.
Wilson. 2000. Interaction of *Bordetella pertussis* with human respiratory mucosa *in vitro*.
Respir. Med. 94: 791-799.
10.   Serra, D. O., M. S. Conover, L. Arnal, G. P. Sloan, M. E. Rodriguez, O. M. Yantorno, and R.
Deora. 2011. FHA-mediated cell-substrate and cell-cell adhesions are critical for *Bordetella*
pertussis biofilm formation on abiotic surfaces and in the mouse nose and the trachea. *PLoS*
One 6: e28811.
11.   de Gouw, D., D. O. Serra, M. I. de Jonge, P. W. Hermans, H. J. Wessels, A. Zomer, O. M.
Yantorno, D. A. Diavatopoulos, and F. R. Mooi. 2014. The vaccine potential of *Bordetella*
pertussis biofilm-derived membrane proteins. *Emerg Microbes Infect* 3: e58.
12.   Dorji, D., R. M. Graham, P. Richmond, A. Keil, and T. K. Mukkur. 2016. Biofilm forming
potential and antimicrobial susceptibility of newly emerged Western Australian *Bordetella*
pertussis clinical isolates. *Biofouling* 32: 1141-1152.
13.   Carriquiriborde, F., P. Martin Aispuro, N. Ambrosis, E. Zurita, D. Bottero, M. E. Gaillard, C.
Castuma, E. Rudi, A. Lodeiro, and D. F. Hozbor. 2021. Pertussis Vaccine Candidate Based on
Outer Membrane Vesicles Derived From Biofilm Culture. *Front. Immunol.* 12: 730434.
14.   Moon, K., R. P. Bonocora, D. D. Kim, Q. Chen, J. T. Wade, S. Stibitz, and D. M. Hinton.
2017. The BvgAS Regulon of *Bordetella pertussis*. *mBio* 8: e01526-01517.
15.   Belcher, T., I. MacArthur, J. D. King, G. C. Langridge, M. Mayho, J. Parkhill, and A.
Preston. 2020. Fundamental differences in physiology of *Bordetella pertussis* dependent on
the two-component system Bvg revealed by gene essentiality studies. *Microb Genom* 6:
mgen000496.
16.   Orth, J. D., I. Thiele, and B. Ø. Palsson. 2010. What is flux balance analysis? *Nat. Biotechnol.*
28: 245-248.
17.   Fong, N. L., J. A. Lerman, I. Lam, B. O. Palsson, and P. Charusanti. 2013. Reconciling a
Salmonella enterica metabolic model with experimental data confirms that overexpression of

645		the glyoxylate shunt can rescue a lethal ppc deletion mutant. *FEMS Microbiol. Lett.* 342: 62-
646		69.
647	18.	Metz, Z. P., T. Ding, and D. J. Baumler. 2018. Using genome-scale metabolic models to
648		compare serovars of the foodborne pathogen *Listeria monocytogenes*. *PLoS One* 13:
649		e0198584.
650	19.	Lobel, L., N. Sigal, I. Borovok, E. Ruppin, and A. A. Herskovits. 2012. Integrative Genomic
651		Analysis Identifies Isoleucine and CodY as Regulators of *Listeria monocytogenes* Virulence.
652		*PLoS Genet.* 8: e1002887.
653	20.	Lee, D.-S., H. Burd, J. Liu, E. Almaas, O. Wiest, A.-L. Barabási, Z. N. Oltvai, and V.
654		Kapatral. 2009. Comparative Genome-Scale Metabolic Reconstruction and Flux Balance
655		Analysis of Multiple *Staphylococcus aureus* Genomes Identify Novel Antimicrobial Drug
656		Targets. *J. Bacteriol.* 191: 4015-4024.
657	21.	Rienksma, R. A., P. J. Schaap, V. A. P. Martins Dos Santos, and M. Suarez-Diez. 2018.
658		Modeling the Metabolic State of *Mycobacterium tuberculosis* Upon Infection. *Frontiers in*
659		*cellular and infection microbiology* 8: 264-264.
660	22.	Gu, C., G. B. Kim, W. J. Kim, H. U. Kim, and S. Y. Lee. 2019. Current status and
661		applications of genome-scale metabolic models. *Genome Biol.* 20: 121.
662	23.	Branco Dos Santos, F., B. G. Olivier, J. Boele, V. Smessaert, P. De Rop, P. Krumpochova, G.
663		W. Klau, M. Giera, P. Dehottay, B. Teusink, and P. Goffin. 2017. Probing the Genome-Scale
664		Metabolic Landscape of *Bordetella pertussis*, the Causative Agent of Whooping Cough.
665		*Applied and environmental microbiology* 83: e01528-01517.
666	24.	Fyson, N., J. King, T. Belcher, A. Preston, and C. Colijn. 2017. A curated genome-scale
667		metabolic model of *Bordetella pertussis* metabolism. *PLOS Computational Biology* 13:
668		e1005639.
669	25.	Gonyar, L. A., P. E. Gelbach, D. G. McDuffie, A. F. Koeppel, Q. Chen, G. Lee, L. M.
670		Temple, S. Stibitz, E. L. Hewlett, J. A. Papin, F. H. Damron, and J. C. Eby. 2019. *In Vivo*
671		Gene Essentiality and Metabolism in *Bordetella pertussis*. *mSphere* 4: e00694-00618.
672	26.	Åkesson, M., J. Förster, and J. Nielsen. 2004. Integration of gene expression data into
673		genome-scale metabolic models. *Metab. Eng.* 6: 285-293.
674	27.	Zur, H., E. Ruppin, and T. Shlomi. 2010. iMAT: an integrative metabolic analysis tool.
675		*Bioinformatics* 26: 3140-3142.
676	28.	Stempler, S., K. Yizhak, and E. Ruppin. 2014. Integrating Transcriptomics with Metabolic
677		Modeling Predicts Biomarkers and Drug Targets for Alzheimer's Disease. *PLoS One* 9:
678		e105383.
679	29.	Safarchi, A., S. Octavia, L. D. W. Luu, C. Y. Tay, V. Sintchenko, N. Wood, H. Marshall, P.
680		McIntyre, and R. Lan. 2016. Better colonisation of newly emerged *Bordetella pertussis* in the
681		co-infection mouse model study. *Vaccine* 34: 3967-3971.
682	30.	Safarchi, A., S. Octavia, L. D. W. Luu, C. Y. Tay, V. Sintchenko, N. Wood, H. Marshall, P.
683		McIntyre, and R. Lan. 2015. Pertactin negative *Bordetella pertussis* demonstrates higher
684		fitness under vaccine selection pressure in a mixed infection model. *Vaccine* 33: 6277-6281.
685	31.	Luu, L. D. W., S. Octavia, L. Zhong, M. J. Raftery, V. Sintchenko, and R. Lan. 2018.
686		Proteomic Adaptation of Australian Epidemic *Bordetella pertussis*. *Proteomics* 18: 1700237.
687	32.	Thalen, M., I. J. van den, W. Jiskoot, B. Zomer, P. Roholl, C. de Gooijer, C. Beuvery, and J.
688		Tramper. 1999. Rational medium design for *Bordetella pertussis*: basic metabolism. *J.*
689		*Biotechnol.* 75: 147-159.
690	33.	Cattelan, N., O. M. Yantorno, and R. Deora. 2018. Structural Analysis of *Bordetella pertussis*
691		Biofilms by Confocal Laser Scanning Microscopy. *Bio Protocol* 8: e2953.
692	34.	Heydorn, A., A. T. Nielsen, M. Hentzer, C. Sternberg, M. Givskov, B. K. Ersboll, and S.
693		Molin. 2000. Quantification of biofilm structures by the novel computer program
694		COMSTAT. *Microbiology* 146 (Pt 10): 2395-2407.
695	35.	Luu, L. D. W., S. Octavia, L. Zhong, M. Raftery, V. Sintchenko, and R. Lan. 2017.
696		Characterisation of the *Bordetella pertussis* secretome under different media. *J. Proteomics*
697		158: 43-51.

36.   Storey, J. D., and R. Tibshirani. 2003. Statistical significance for genomewide studies.
Proceedings of the National Academy of Sciences 100: 9440-9445.

37.   Bart, M. J., S. R. Harris, A. Advani, Y. Arakawa, D. Bottero, V. Bouchez, P. K. Cassiday, C.-
S. Chiang, T. Dalby, N. K. Fry, M. E. Gaillard, M. van Gent, N. Guiso, H. O. Hallander, E. T.
Harvill, Q. He, H. G. J. van der Heide, K. Heuvelman, D. F. Hozbor, K. Kamachi, G. I.
Karataev, R. Lan, A. Lutyńska, R. P. Maharjan, J. Mertsola, T. Miyamura, S. Octavia, A.
Preston, M. A. Quail, V. Sintchenko, P. Stefanelli, M. L. Tondella, R. S. W. Tsang, Y. Xu, S.-
M. Yao, S. Zhang, J. Parkhill, and F. R. Mooi. 2014. Global Population Structure and
Evolution of *Bordetella pertussis* and Their Relationship with Vaccination. *mBio* 5: e01074-
01014.

38.   Perez-Riverol, Y., J. Bai, C. Bandla, D. García-Seisdedos, S. Hewapathirana, S.
Kamatchinathan, D. J. Kundu, A. Prakash, A. Frericks-Zipper, M. Eisenacher, M. Walzer, S.
Wang, A. Brazma, and J. A. Vizcaíno. 2022. The PRIDE database resources in 2022: a hub
for mass spectrometry-based proteomics evidences. *Nucleic Acids Res.* 50: D543-d552.

39.   Zur, H., E. Ruppin, and T. Shlomi. 2010. iMAT: an integrative metabolic analysis tool.
Bioinformatics 26: 3140-3142.

40.   Heirendt, L., S. Arreckx, T. Pfau, S. N. Mendoza, A. Richelle, A. Heinken, H. S.
Haraldsdóttir, J. Wachowiak, S. M. Keating, V. Vlasov, S. Magnusdóttir, C. Y. Ng, G.
Preciat, A. Žagare, S. H. J. Chan, M. K. Aurich, C. M. Clancy, J. Modamio, J. T. Sauls, A.
Noronha, A. Bordbar, B. Cousins, D. C. El Assal, L. V. Valcarcel, I. Apaolaza, S. Ghaderi,
M. Ahookhosh, M. Ben Guebila, A. Kostromins, N. Sompairac, H. M. Le, D. Ma, Y. Sun, L.
Wang, J. T. Yurkovich, M. A. P. Oliveira, P. T. Vuong, L. P. El Assal, I. Kuperstein, A.
Zinovyev, H. S. Hinton, W. A. Bryant, F. J. Aragón Artacho, F. J. Planes, E. Stalidzans, A.
Maass, S. Vempala, M. Hucka, M. A. Saunders, C. D. Maranas, N. E. Lewis, T. Sauter, B. Ø.
Palsson, I. Thiele, and R. M. T. Fleming. 2019. Creation and analysis of biochemical
constraint-based models using the COBRA Toolbox v.3.0. *Nat. Protoc.* 14: 639-702.

41.   Feist, A. M., C. S. Henry, J. L. Reed, M. Krummenacker, A. R. Joyce, P. D. Karp, L. J.
Broadbelt, V. Hatzimanikatis, and B. Ø. Palsson. 2007. A genome-scale metabolic
reconstruction for *Escherichia coli* K-12 MG1655 that accounts for 1260 ORFs and
thermodynamic information. *Mol. Syst. Biol.* 3: 121-121.

42.   Izac, M., D. Garnier, D. Speck, and N. D. Lindley. 2015. A Functional Tricarboxylic Acid
Cycle Operates during Growth of *Bordetella pertussis* on Amino Acid Mixtures as Sole
Carbon Substrates. *PLoS One* 10: e0145251.

43.   Lamberti, Y., J. H. Cafiero, K. Surmann, H. Valdez, J. Holubova, B. Večerek, P. Sebo, F.
Schmidt, U. Völker, and M. E. Rodriguez. 2016. Proteome analysis of *Bordetella pertussis*
isolated from human macrophages. *J. Proteomics* 136: 55-67.

44.   Malik-Sheriff, R. S., M. Glont, T. V. N. Nguyen, K. Tiwari, M. G. Roberts, A. Xavier, M. T.
Vu, J. Men, M. Maire, S. Kananathan, E. L. Fairbanks, J. P. Meyer, C. Arankalle, T. M.
Varusai, V. Knight-Schrijver, L. Li, C. Dueñas-Roca, G. Dass, S. M. Keating, Y. M. Park, N.
Buso, N. Rodriguez, M. Hucka, and H. Hermjakob. 2019. BioModels—15 years of sharing
computational models in life science. *Nucleic Acids Res.* 48: D407-D415.

45.   Montezano, D., L. Meek, R. Gupta, L. E. Bermudez, and J. C. M. Bermudez. 2015. Flux
Balance Analysis with Objective Function Defined by Proteomics Data—Metabolism of
Mycobacterium tuberculosis Exposed to Mefloquine. *PLoS One* 10: e0134014.

46.   Serra, D. O., G. Lucking, F. Weiland, S. Schulz, A. Gorg, O. M. Yantorno, and M. Ehling-
Schulz. 2008. Proteome approaches combined with Fourier transform infrared spectroscopy
revealed a distinctive biofilm physiology in *Bordetella pertussis*. *Proteomics* 8: 4995-5010.

47.   Goto, M., A. Abe, T. Hanawa, and A. Kuwae. 2021. Bcr4 is a Chaperone for the Inner Rod
Protein in the *Bordetella* Type III Secretion System. *bioRxiv*: 2021.2009.2028.462275.

48.   Arnal, L., T. Grunert, N. Cattelan, D. de Gouw, M. I. Villalba, D. O. Serra, F. R. Mooi, M.
Ehling-Schulz, and O. M. Yantorno. 2015. *Bordetella pertussis* Isolates from Argentinean
Whooping Cough Patients Display Enhanced Biofilm Formation Capacity Compared to
Tohama I Reference Strain. *Front. Microbiol.* 6: 1352.

49. Bosch, A., D. Serra, C. Prieto, J. Schmitt, D. Naumann, and O. Yantorno. 2005.
Characterization of *Bordetella pertussis* growing as biofilm by chemical analysis and FT-IR
spectroscopy. *Applied Microbiology and Biotechnology* 71: 736.

50. Ahn, S., J. Jung, I.-A. Jang, E. L. Madsen, and W. Park. 2016. Role of Glyoxylate Shunt in
Oxidative Stress Response. *Journal of Biological Chemistry* 291: 11928-11938.

51. Uppuluri, P., M. Acosta Zaldívar, M. Z. Anderson, M. J. Dunn, J. Berman, J. L. Lopez Ribot,
and J. R. Köhler. 2018. *Candida albicans* Dispersed Cells Are Developmentally Distinct from
Biofilm and Planktonic Cells. *mBio* 9: e01338-01318.

52. Yahya, M., U. Hamid, M. Norfatimah, and R. Kambol. 2014. *In silico* analysis of essential
tricarboxylic acid cycle enzymes from biofilm-forming bacteria. *Trends in Bioinformatics* 7:
19-26.

53. Rivera, I., B. Linz, K. K. Dewan, L. Ma, C. A. Rice, D. E. Kyle, and E. T. Harvill. 2019.
Conservation of Ancient Genetic Pathways for Intracellular Persistence Among Animal
Pathogenic Bordetellae. *Front. Microbiol.* 10: 2839.

54. McKinney, J. D., K. Höner zu Bentrup, E. J. Muñoz-Elías, A. Miczak, B. Chen, W. T. Chan,
D. Swenson, J. C. Sacchettini, W. R. Jacobs, Jr., and D. G. Russell. 2000. Persistence of
Mycobacterium tuberculosis in macrophages and mice requires the glyoxylate shunt enzyme
isocitrate lyase. *Nature* 406: 735-738.

55. Suo, Y., Y. Huang, Y. Liu, C. Shi, and X. Shi. 2012. The expression of superoxide dismutase
(SOD) and a putative ABC transporter permease is inversely correlated during biofilm
formation in *Listeria monocytogenes* 4b G. *PLoS One* 7: e48467-e48467.

56. Seydlova, G., J. Beranova, I. Bibova, A. Dienstbier, J. Drzmisek, J. Masin, R. Fiser, I.
Konopasek, and B. Vecerek. 2017. The extent of the temperature-induced membrane
remodeling in two closely related *Bordetella* species reflects their adaptation to diverse
environmental niches. *J. Biol. Chem.* 292: 8048-8058.

57. Dubois-Brissonnet, F., E. Trotier, and R. Briandet. 2016. The Biofilm Lifestyle Involves an
Increase in Bacterial Membrane Saturated Fatty Acids. *Front. Microbiol.* 7: 1673.

58. Nepper, J. F., Y. C. Lin, and D. B. Weibel. 2019. Rcs Phosphorelay Activation in Cardiolipin-
Deficient *Escherichia coli* Reduces Biofilm Formation. *J. Bacteriol.* 201: e00804-00818.

59. Kawai, Y., and A. Moribayashi. 1982. Characteristic lipids of *Bordetella pertussis*: simple
fatty acid composition, hydroxy fatty acids, and an ornithine-containing lipid. *J. Bacteriol.*
151: 996-1005.

60. Cattelan, N., P. Dubey, L. Arnal, O. M. Yantorno, and R. Deora. 2016. *Bordetella* biofilms: a
lifestyle leading to persistent infections. *Pathogens and Disease* 74: ftv108.

61. Cava, F., H. Lam, M. A. de Pedro, and M. K. Waldor. 2011. Emerging knowledge of
regulatory roles of D-amino acids in bacteria. *Cell Mol Life Sci* 68: 817-831.

62. Goh, S.-N., A. Fernandez, S.-Z. Ang, W.-Y. Lau, D.-L. Ng, and E. S. G. Cheah. 2013. Effects
of Different Amino Acids on Biofilm Growth, Swimming Motility and Twitching Motility in
Escherichia coli BL21. *Journal of Biology and Life Science* 4: 13.

63. Aliashkevich, A., L. Alvarez, and F. Cava. 2018. New Insights Into the Mechanisms and
Biological Roles of D-Amino Acids in Complex Eco-Systems. *Front. Microbiol.* 9: 683.

64. Borriello, G., L. Richards, G. D. Ehrlich, and P. S. Stewart. 2006. Arginine or Nitrate
Enhances Antibiotic Susceptibility of *Pseudomonas aeruginosa* in Biofilms. *Antimicrob.*
Agents Chemother. 50: 382-384.

65. Kolderman, E., D. Bettampadi, D. Samarian, S. E. Dowd, B. Foxman, N. S. Jakubovics, and
A. H. Rickard. 2015. L-Arginine Destabilizes Oral Multi-Species Biofilm Communities
Developed in Human Saliva. *PLoS One* 10: e0121835.

66. Brindle, E. R., D. A. Miller, and P. S. Stewart. 2011. Hydrodynamic deformation and removal
of *Staphylococcus epidermidis* biofilms treated with urea, chlorhexidine, iron chloride, or
DispersinB. *Biotechnology and bioengineering* 108: 2968-2977.

67. Zhuo, T., W. Rou, X. Song, J. Guo, X. Fan, G. G. Kamau, and H. Zou. 2015. Molecular study
on the *carAB* operon reveals that *carB* gene is required for swimming and biofilm formation
in *Xanthomonas citri* subsp. *citri*. *BMC Microbiol.* 15: 225-225.

68. Pisithkul, T., J. W. Schroeder, E. A. Trujillo, P. Yeesin, D. M. Stevenson, T. Chaiamarit, J. J.
Coon, J. D. Wang, D. Amador-Noguez, J.-M. Ghigo, and E. P. Greenberg. 2019. Metabolic
Remodeling during Biofilm Development of *Bacillus subtilis*. *mBio* 10: e00623-00619.
69. Yang, H., M. Wang, J. Yu, and H. Wei. 2015. Aspartate inhibits *Staphylococcus aureus*
biofilm formation. *FEMS Microbiol. Lett.* 362: fnv025.
70. Allan, R. N., P. Skipp, J. Jefferies, S. C. Clarke, S. N. Faust, L. Hall-Stoodley, and J. Webb.
2014. Pronounced metabolic changes in adaptation to biofilm growth by *Streptococcus*
pneumoniae. *PLoS One* 9: e107015.
71. Hasenoehrl, E. J., D. Rae Sajorda, L. Berney-Meyer, S. Johnson, J. M. Tufariello, T. Fuhrer,
G. M. Cook, W. R. Jacobs, and M. Berney. 2019. Derailing the aspartate pathway of
Mycobacterium tuberculosis to eradicate persistent infection. *Nature Communications* 10:
4215.
72. Díaz-Pascual, F., M. Lempp, K. Nosho, H. Jeckel, J. K. Jo, K. Neuhaus, R. Hartmann, E.
Jelli, M. F. Hansen, A. Price-Whelan, L. E. P. Dietrich, H. Link, and K. Drescher. 2021.
Spatial alanine metabolism determines local growth dynamics of *Escherichia coli* colonies.
eLife 10: e70794.
73. Pulukkody, A. C., Y. P. Yung, F. Donnarumma, K. K. Murray, R. P. Carlson, and L. Hanley.
2021. Spatially resolved analysis of *Pseudomonas aeruginosa* biofilm proteomes measured by
laser ablation sample transfer. *PLoS One* 16: e0250911.
74. Valle, J., S. Da Re, S. Schmid, D. Skurnik, R. D'Ari, and J. M. Ghigo. 2008. The amino acid
valine is secreted in continuous-flow bacterial biofilms. *J. Bacteriol.* 190: 264-274.
75. Cotter, P. A., and A. M. Jones. 2003. Phosphorelay control of virulence gene expression in
Bordetella. *Trends Microbiol.* 11: 367-373.
76. Bone, M. A., A. J. Wilk, A. I. Perault, S. A. Marlatt, E. V. Scheller, R. Anthouard, Q. Chen,
S. Stibitz, P. A. Cotter, and S. M. Julio. 2017. Bordetella PlrSR regulatory system controls
BvgAS activity and virulence in the lower respiratory tract. *Proceedings of the National*
Academy of Sciences 114: E1519-E1527.
77. Palmer, K. L., L. M. Aye, and M. Whiteley. 2007. Nutritional cues control *Pseudomonas*
aeruginosa multicellular behavior in cystic fibrosis sputum. *J. Bacteriol.* 189: 8079-8087.
78. Williamson, K. S., L. A. Richards, A. C. Perez-Osorio, B. Pitts, K. McInnerney, P. S. Stewart,
and M. J. Franklin. 2012. Heterogeneity in *Pseudomonas aeruginosa* Biofilms Includes
Expression of Ribosome Hibernation Factors in the Antibiotic-Tolerant Subpopulation and
Hypoxia-Induced Stress Response in the Metabolically Active Population. *J. Bacteriol.* 194:
2062-2073.
79. Salgar-Chaparro, S. J., K. Lepkova, T. Pojtanabuntoeng, A. Darwin, L. L. Machuca, and A. J.
M. Stams. 2020. Nutrient Level Determines Biofilm Characteristics and Subsequent Impact
on Microbial Corrosion and Biocide Effectiveness. *Applied and environmental microbiology*
86: e02885-02819.
80. Siriwach, R., F. Matsuda, K. Yano, and M. Y. Hirai. 2020. Drought Stress Responses in
Context-Specific Genome-Scale Metabolic Models of *Arabidopsis thaliana*. *Metabolites* 10:
159.

Tables

Table 1 Virulence proteins differentially expressed or uniquely identified between planktonic

and biofilm cells

Locus	Gene	Product	Fold change (Biofilm/Planktonic)	t-Test ($p \leq 0.05$)	q-Value ($q \leq 0.05$)
BP0499	btcA	Type III secretion chaperone	0.594	0.002	0.0005
BP0500	bteA	Type III secretion toxin, effector	0.131	9.02E-08	2.52E-07
BP0760	cyaA	Bifunctional hemolysin-adenylate cyclase	1.530	0.042	0.008
BP1054	prn	Pertactin autotransporter	0.163	2.19E-06	2.42E-06
BP1112	bipA	Outer membrane ligand binding protein	2.073	3.65E-06	3.4E-06
BP1119	fim2	Serotype 2 fimbrial subunit	0.590	0.002	0.0006
BP1201	tcfA	Tracheal colonisation factor	0.070	2.94E-05	1.64E-05
BP1251	-	Putative toxin	Planktonic unique		
BP1877	bvgS	Virulence sensor protein BvgS	1.876	5.63E-05	2.77E-05
BP1878	bvgA	Virulence factors transcription regulator BvgA	3.101	1.69E-06	2.02E-06
BP1884	fhaC	Filamentous hemagglutinin transporter protein FhaC	Biofilm unique		
BP2233	bspR	Type III secretion chaperone	0.430	1.66E-06	2.02E-06
BP2236	bscW	Type III secretion chaperone	Planktonic unique		
BP2250	bcr4	Type III secretion protein	1.533	0.027	0.005
BP2251	bcrH2	Type III secretion chaperone	0.324	1.38E-08	8.98E-08
BP2253	bopD	Type III secretion system outer protein D	0.116	7.61E-10	1.49E-08
BP2255	btc22	Type III secretion protein chaperone	Planktonic unique		
BP2256	bsp22	Type III secretion tip protein	Planktonic unique		
BP2257	bopN	Type III secretion outer protein N, effector	Planktonic unique		
BP2263	bscE	Type III secretion protein	0.282	3.24E-08	1.73E-07
BP2315	vag8	Autotransporter	0.429	1.39E-05	9.03E-06
BP3439	dnt	Dermonecrotic toxin	1.429	0.135	0.022
BP3494	brkA	BrkA autotransporter	1.218	0.139	0.023
BP3654	cyaY	Iron-sulfur cluster assembly protein CyaY	0.618	0.188	0.030
BP3783	ptxA	Pertussis toxin subunit 1	0.299	1.37E-06	1.75E-06
BP3785	ptxD	Pertussis toxin subunit 4	Biofilm unique		

Table 2 Number of metabolites, reactions and genes in the iMAT context specific models for *B. pertussis* biofilm and planktonic cells.

	Biofilm	Planktonic
High expression (+1)	36	36
Moderate expression (0)	117	86
Low expression (-1)	65	78
Total proteins incorporated	218	200
iMAT models		
Metabolites	198	213
Reactions	206	219
Genes	188	195

Figures

Figure 1 Experimental design for iMAT model generation. Proteins were extracted from planktonic and biofilm *B. pertussis* cells. The expression values were then used to generate context specific metabolic models based on the extensively curated iBP1870 *B. pertussis* model (23). Flux balance analysis was run on the models and the altered reactions compared between the planktonic and biofilm models.

Figure 2 Volcano plot of total protein expression changes between biofilm and planktonic cells. A Expression profile of all proteins identified through the LC-MS/MS analysis of planktonic and biofilm *B. pertussis* cells. Proteins are plotted on a volcano plot displaying the $\log_2(q\text{-value})$ on the y-axis and $\log_2(\text{fold change [biofilm/planktonic]})$ on the x-axis. The dashed vertical grey lines mark a fold change of 0.8 and 1.2 and the horizontal line marks the threshold of q-value = 0.05. Highlighted in red are the proteins that were identified in all 6 biological replicates that were incorporated into the iMAT models. **B** Expression profile of the subset of proteins incorporated into the iMAT model. Red markers are proteins that were designated significantly differentially downregulated and the blue markers are proteins that were significantly upregulated in biofilm cells. The dashed vertical

 grey lines mark a fold change of 0.8 and 1.2 and the horizontal line marks the threshold of *q*-

value = 0.05.

Figure 3 Proteins up and downregulated in *B. pertussis* biofilm cells compared to

planktonic cells identified using LC-MS/MS. Proteins significantly up and downregulated

in biofilm cells were categorised in functional categories based on Bart et al. (37). Red and

blue bars represent the total number of proteins within the functional category significantly

up or downregulated, respectively. Asterisk (*) denotes functional categories significantly up

or downregulated based on Fisher's exact test with Benjamini-Hochberg multiple test

correction.

Figure 4 Comparison of reactions of *B. pertussis* planktonic and biofilm iMAT models.

A Venn diagram highlighting common and unique reactions between planktonic and biofilm

iMAT models generated by incorporating proteomic expression data. **B** Planktonic and

biofilm model reactions grouped into subsystems based on the *Escherichia coli* iAF1260

metabolic model. Unique reactions identified in one model but not the other are highlighted

in different colours on the graph.

Figure 5 Summarised pathways with major changes between planktonic and biofilm

iMAT models. Major changed core metabolic reactions were summarised and represented in

a model. The model includes the reactions that were unique in either model as well as

reactions with altered flux. Ac, Acetate; Ac-CoA, Acetyl-CoA; Akg, α-ketoglutarate; Ala,

Alanine; Arg, Arginine; Asp, Aspartate; BCAA, Branched chain amino acid degradation;

Cbp, Carbamoyl phosphate; Cgly, Cysteinylglycine; Cit, Citrate; CO_2, Carbon dioxide; Cys,

Cysteine; Fum, Fumarate; Gln, Glutamine; Glu, Glutamate; Glx, Glyoxylate; Gly, Glycine;

Gly3p, Glycine 3-phosphate; Glyc, Glycine; GNG, Gluconeogenesis; GPL,

Glycerophospholipid metabolism; Gthrd, Glutathione; Hom, Homoserine; Lac, Lactate; Mal,

Malate; O_2, Oxygen; OAA, Oxaloacetate; Orn, Ornithine; Orot, Orotate; Phb,

Polyhydroxybutyrate; Prop, Propanoate metabolism; Ptrc, Putrescine; Pyr, Pyruvate; SO_2,

Sulfur dioxide; Suc, Succinate; TCA, Tricarboxcylic acid cycle; Thr, Threonine; Val, Valine.

B. pertussis iBP180
metabolic model

Planktonic protein expression

m/z

Biofilm protein expression

m/z

Planktonic iMAT model

Biofilm iMAT model

Model comparison