Title: Don't get a big head: *Centris pallida* (Hymenoptera: Apidae) male head widths decrease across

five decades

Running Title: Declines in male *C. pallida* head width

Authors: Meghan Barrett[1], Meredith G. Johnson[2]

[1]Department of Biology, California State University Dominguez Hills, Carson, CA 90747

[2]School of Life Sciences, Arizona State University, Tempe, AZ 85281

Authorship Contributions: MB – conceptualization, data curation, formal analysis, investigation,

methodology, writing – original draft; MGJ – investigation, funding acquisition, writing – review &

editing

Conflict of Interests: None.

Abstract

Historical data suggest that many bee species have declined in body size. Larger-bodied bees with

narrow phenological and dietary breadth are most prone to declines in body size over time. This

may be especially true in solitary, desert-adapted species that are vulnerable to climate change – such

as *Centris pallida* (Hymenoptera: Apidae). In addition, body size changes in species with size-linked

behaviors could threaten the prevalence of certain behavioral phenotypes long-term. *C. pallida*

solitary bees are found in the Sonoran Desert. Males use alternative reproductive tactics (ARTs) and

are dimorphic in both morphology and behavior. *C. pallida* male body size has been studied since the

1970s in the same population. We collected body size data in 2022 and combined it with published

records from 1974-2022. We find a persistent decline in the mean head width of patrolling males,

and shifts towards smaller body sizes in the populations of males found foraging and hovering.

Mating males did not experience a decline in mean body size over time. We discuss hypotheses

related to the decline in *C. pallida* male head width. Finally, we advocate for *C. pallida* as an excellent

study system for understanding the stability of ARTs with size-linked behavioral phenotypes.

Introduction

Anthropogenic changes in climate and habitat have caused a decline in abundance, a shift in

geographic range or phenology, and changes in the morphology, physiology, or behavior of many

species, sometimes leading to alterations in life history and ecological relationships (Bartomeus et al.

2011, Biesmeijer et al. 2006, Jacobson et al. 2018, Béltran et al. 2021, Burraco et al. 2020, Chou et al.

2019, Chung & Schulte 2020, Duffy et al. 2015, Huey & Kingsolver 2019, Kuhlmann et al. 2012,

Ockendon et al. 2014, Parmesan 2006, Reed 2018, Sinervo et al. 2010, Walters & Hassall 2006,

Zattara & Aizen 2021, Sánchez-Bayo & Wyckhuys 2019, Burkle et al. 2013).

Bees may be particularly vulnerable to climate change or habitat modifications. As heterothermic

ectotherms they rely on environmental conditions to maintain non-lethal body temperatures (Wieser

1973, Huey & Stevenson 1979). This is compounded by a reliance on resources (e.g., nesting

resources, plants) that are often disturbed by human activity (Grab et al. 2019, McCabe et al. 2021).

Documented changes in bee populations over time can help determine how human activity may

impact bees with different life history characteristics, with knock-on effects for their ecological

relationships. (e.g., Cane et al. 2006, Burkle et al. 2013). For example, studies in Europe and the

Northeastern regions of North America have demonstrated that larger-bodied bee species are more

likely to experience declines in population (Bartomeus et al. 2013, Scheper et al. 2014, Oliveira et al.

2016), or stronger intraspecific declines in body size (Nooten & Rehan 2020, but see Gérard et al.

2019), perhaps due to greater nutritional requirements (Müller et al. 2006).

Bee body sizes have not changed uniformly over time across species, sexes, or regions (Nooten &

Rehan 2020, Bartomeus et al. 2013, Scheper et al. 2014, Oliveira et al. 2016, Kleijn & Raemakers

2008), suggesting life history traits and ecosystem characteristics (e.g., habitat fragmentation,

Warzecha et al. 2016) may play important roles in determining the impact of human activities on

body size. For most bees, body size is determined by resource provisioning in the larval stage (Chole

et al. 2019, Kukuk 1996, Alcock 1984, Lawson et al. 2017), and can thus be strongly influenced by

changes in resource availability (Chown & Gaston 2010). Bees with narrow phenological windows

and dietary specialists are likely to be more susceptible to declines, given their reliance on a smaller

pool of available resources (Bartomeus et al. 2013). Solitary and desert-adapted bees are expected to

be particularly vulnerable to climate change (Sala et al. 2000, Loarie et al. 2009, Vale & Brito 2015,

Hamblin et al. 2017, Burdine & McCluney 2019, McCabe et al. 2021, but see Silva et al. 2018), given

the thermal and hygric stressors already evident in their environment. To date, no studies have

tracked changes in morphology of any solitary bees in desert ecosystems.

Declines in body size could also generate species-level behavioral changes whenever size contributes

to behavior, such as in many alternative reproductive tactic systems (ARTs). ARTs occur when

categorical variation in the morphological/behavioral traits is associated with mating across

individuals of the same sex within the same population (Oliveira et al. 2008, Paxton 2005, Shuster

2010). In many such systems, larger- and smaller-bodied individuals use different strategies for

accessing mates (e.g., fighters vs. sneakers; Oliveira et al. 2008). Declines in body size may alter the

occurrence of size-linked morphs, and their behaviors, resulting in a loss of important intraspecific

variation over time.

Centris pallida (Hymenoptera: Apidae) are widespread, common solitary digger bees are found in the

Sonoran Desert of the Southwestern United States and Northern Mexico. This species often forms

dense nesting aggregations, where many thousands of individuals occupy several hectares. Males

emerge first from their natal nests at these sites and use ARTs to find mates. Males are behaviorally

and morphologically dimorphic, using different sensory mate location strategies and microclimates

(Alcock 1976, Alcock et al. 1977a, Snelling 1984, Barrett et al. 2021, Barrett 2022). Large-morph,

'metandric' males with pale gray coloration patrol in sinuous loops ~10 cm over the emergence site

and use scent to locate females emerging from natal nests; males then engage in fights for the

opportunity to dig up and mate with emerging females (Alcock et al. 1976b). Small-morph males,

with dark brown coloration, are more behaviorally flexible; they may patrol, but are often found

hovering near plants where they use visual cues to locate females or mating pairs flying away from

the aggregation site that may be interrupted. Large-morph males have a fitness advantage in

situations where mating aggregations are densely populated (Alcock et al. 1977a, 1977b, Alcock

1984, Alcock 1995, Alcock 2013a). *C. pallida* mating aggregations may persist in the same locations

over multiple decades – allowing for the continuous resampling of a population (Barrett 2022). *C.*

pallida are large-bodied, desert-adapted, solitary bees with an extremely narrow phenological and

dietary breadth. Bees nest for less than six weeks in the late spring, and females utilize four species

of flowering trees for larval nectar/pollen provisions (*Parkinsonia microphylla, P. aculeata, Olneya tesota,*

and, rarely, *Psorothamnus (Dalea) spinosa,* Alcock et al. 1976a). *P. microphylla* is the predominant host

plant for *C. pallida* over its entire geographic range (Buchmann, pers. comm). The body sizes of male

C. pallida have been studied at the same site since for 48 years (Alcock et al. 1977a). By combining

historical data on *C. pallida* body sizes from 1974 – 2018 (Alcock et al. 1977a, Alcock 1984, Alcock

1989, Alcock 2013a, Barrett 2022) with data we collected in 2022 from the same population, we

aimed to determine if male *C. pallida* bees have experienced persistent changes or persistent stability

in body size. Our results demonstrate that the population of males found foraging at trees

(representative of the total population of males), hovering, and patrolling have all declined in body

size since 1974. However, the population of mating males has not persistently declined in body size

over time. In addition, we present the first body size data on female *C. pallida* from this site, for use

in future historical comparisons. Finally, we discuss several hypotheses of causes for the decline in

male body size, and the implications of declining male body size on the stability of the *C. pallida*

ART.

Methods

<u>Head width measurements</u>

We collected head widths at the same site where most previous research on *C. pallida* body size has

been conducted (Alcock et al. 1977a, Alcock 1984, Alcock 1989, Alcock 2013a), in the floodplain by

Blue Point Bridge/Saguaro Lake over the Salt River north of Mesa, Arizona (33.552, -111.566).

Alcock et al. (1977a) demonstrated that head width is correlated with body weight in *C. pallida* males,

and serves as a reliable field indicator of body size.

We collected bees (n = 921 males, 114 females) in the same manner as described in Alcock (1984,

2013), to be sure our data were comparable. Briefly: for patrolling males, we made rapid sweeps low

to the ground with an insect net through open areas of searching males. Digging/fighting males

were approached and collected by hand. Foraging males and females were collected with a

telescoping insect net as they visited paloverde trees (*Parkinsonia* sp.). Mating males and females were

collected while engaged in copulation on the ground or vegetation. Hovering males were collected

near mesquite or palo verde trees, wherever they had established stable aerial stations. We measured

head widths to 0.01 mm using digital calipers (Wen 10761), before releasing the bees. We collected

bees daily when the aggregation was most active between April 20 and May 8, 2022, generally

between the hours of 0700 and 1130, and noted the time of collection and behavior for each

individual.

<u>Historical data for comparisons</u>

To obtain historical male head width data, we used reported means and standard deviations in

Alcock et al. 1977a, Alcock 1984, Alcock 2013a, and Barrett 2022. Together, these studies report

data collected in: 1974, 1975, 1976, 1982, 1988, 2011, 2012, and 2018. As prior surveys of the

population by Alcock (et al. 1977; Alcock 1984, 1989, 2013) generally only captured bees between

roughly 0700 and 0930, we also used only those bees captured before 0930 in our sampling for the

historical comparison with prior literature. Data were also reported for mating, hovering, and

patrolling bees from this site in 2018 (Barrett 2022); however, these bees were collected between

0600 and 1100, with no individual data on time of collection. Patrolling bees in 2018 (Barrett 2022)

were not captured in a manner consistent with Alcock's prior surveys, and time of day was shown to

impact body size (see results) in patrollers, so we elected not to use the 2018 dataset for patrolling

males in the comparison. However, we did include hovering and mating males as they were collected

using methods consistent with Alcock's prior surveys and time of day was shown not to impact

body size in these populations (see results).

<u>Statistical analysis</u>

Data were analyzed using GraphPad Prism 9.3.1 (GraphPad Prism for Windows 2021). When

analyzing data on the total population in 2022 (as compared to the before-0930 dataset used for the

historical data comparison), we include all bees captured in 2022 (including males for which we had

not taken data on time of capture). Because head width data were not normally distributed for all

behavior categories, we used a Kruskal-Wallis test with post-hoc Dunn's MCT to analyze differences

between hovering, patrolling, and mating male populations. We used a Mann-Whitney test to

compare mating and foraging male populations, and an unpaired t-test to compare digging and

fighting males and mean hovering male head widths from 1974/1975 and 2022. We used a linear

regression to analyze the effect of time of day on patrolling and hovering male head widths. We also

used a linear regression to analyze the effect of year on mean patrolling and mating male head width

across studies. All data can be found archived on Dryad.

Results

C. pallida head widths in 2022

We present data on the mean ± SD and ranges of male head widths for the 2022 population of

males engaged in: foraging, patrolling, hovering, mating, digging, and fighting behaviors in Table 1.

We also present mean female head widths in this population, for females caught while mating,

foraging, or nesting (distribution of female head widths: Supplemental Figure 1).

Table 1. Head widths of *C. pallida* collected in April - May 2022 at Blue Point Bridge before

1130 am (n = 921 male, 114 female).

Type	*Mean ± SD (n)*	*Range*
Foraging Males	4.85 ± 0.36 (203)	4.01 – 5.77
Patrolling Males	5.03 ± 0.34 (410)	4.03 – 5.84
Hovering Males	4.76 ± 0.20 (91)	4.29 – 5.16
Mating Males	5.50 ± 0.31 (71)	4.66 – 6.04
Digging Males	5.48 ± 0.25 (66)	4.74 – 6.00
Fighting Males	5.41 ± 0.25 (71)	4.74 – 5.83
Females	5.23 ± 0.28 (114)	4.33 – 5.95

Hovering males were significantly smaller than patrolling males (Figure 1; Kruskal-Wallis: K-W =

140.4, $p < 0.0001$; Dunn's MCT: $Z = 6.51$, $p < 0.0001$), and both types of males were smaller than

mating males (patrolling vs. mating: $Z = 8.70$, $p < 0.0001$; hovering vs. mating: $Z = 11.83$, $p <$

0.0001). Mating males skewed larger than the total population of foraging males (Mann-Whitney

test: $U = 1439$, $p < 0.0001$). There was no difference in the mean head widths of digging and

fighting males (unpaired t-test: $t = 1.55$, $df = 135$, $p = 0.12$).

Figure 1. Distribution of male head widths by behavioral category in 2022. Distribution of

male head widths (n = 921) collected while foraging (n = 203), hovering (n = 91), patrolling (n =

410), mating (n = 71), and digging/fighting (n = 66/71) before 1130 on April 20 – May 8, 2022 at

Blue Point Bridge, Arizona.

Patrolling and foraging male head widths increased over the course of the morning (Supplementary

Figure 2; linear regression, patrolling: [head width] = 0.095 [time of day] + 4.19, F = 23.65, df =

408, R^2 = 0.05, p < 0.0001; foraging: [head width] = 0.097 [time of day] + 4.08, F = 7.90, df = 201,

R^2 = 0.03, p < 0.0001); hovering and mating male head widths were constant across the morning

(hovering: F = 1.57, df = 88, p = 0.21; mating: F = 0.43, df = 57, p = 0.51).

<u>Historical comparison of mean male head widths from 1974 to 2022</u>

Reported head widths for males foraging, hovering, patrolling, or mating at the Blue Point Bridge

site from 1974 to 2022 can be found in Table 2, with associated references for the historical data.

Table 2. Reported means ± SD of males at Blue Point Bridge before 930 from 1974 – 2022.

	Mean ± SD (n)				
Year	Foraging	Hovering	Patrolling	Mating	Reference
1974 + 1975	no mean reported; Supp Fig 3	4.92 ± 0.15 (50)	5.14 ± 0.29 (100)	5.51 ± 0.36 (50)	Alcock et al. 1977a
1976			5.20 ± 0.28 (168)	5.44 ± 0.33 (102)	Alcock 1984, 1989
1981			5.15 ± 0.23 (162)		Alcock 1984
1982			5.11 ± 0.25 (265)	5.42 ± 0.29 (140)	Alcock 1984, 1989
1988			5.09 ± 0.29 (400)	5.47 ± 0.29 (144)	Alcock 1989
2011			5.02 ± 0.28 (214)		Alcock 2013a
2012	4.88 ± 0.31 (132)		4.95 ± 0.25 (285)	5.32 ± 0.29 (11[*])	Alcock 2013a
2018		4.86 ± 0.21 (11[*†])		5.40 ± 0.31 (23[*†])	Barrett 2022
2022	4.84 ± 0.36 (200)	4.77 ± 0.19 (57)	4.98 ± 0.34 (300)	5.50 ± 0.31 (56)	

169 [*]small sample size.

170 †these bees were not all collected before 0930, as in the other years, but no individual data on time of capture was recorded. However,
171 body size was not affected by time of day in hovering and mating male populations, so we included these males.
172

Patrolling male head widths decreased by 3.72% at the Blue Point Bridge site from 1974 to 2022

(Figure 2A, Supplementary Figure 3B; linear regression, [head width] = -0.004344 [year] + 13.74, F

= 131.2, df = 1892, R^2 = 0.06, p < 0.0001). Although there were only two years to compare

hovering males with a large sample size, there was a significant decrease in mean hovering male head

width in 2022 compared to 1974/1975 (Supplementary Figure 3A, unpaired t-test; t = 4.49, df =

105, p < 0.0001). The mean of the foraging males (n = 100) measured in 1974/1975 was not

reported in Alcock et al. 1977a; however, comparing the distribution of head widths from

1974/1975 to our 2022 data suggests significant declines in the head width of the overall population

of foraging males (Figure 2B & 2C). Mating male head widths did not change from 1974 to 2022

(Figure 2A, Supplementary Figure 4; F = 0.04, df = 524, p = 0.84).

Figure 2. Changes in head widths of patrolling, foraging, and mating males from 1974 to

2022. A) Patrolling male head widths (gray dots, solid line, n = 300) decreased over time at the Blue

Point Bridge site from 1974 to 2022 (Figure 2; linear regression, [head width] = -0.004 [year] +

13.74, F = 131.2, df = 1892, R^2 = 0.06, p < 0.0001). Mating male head widths (black diamonds,

dashed line to demonstrate non-significance, n = 56) did not change from 1974 to 2022 (F = 0.36,

df = 501, p = 0.55). Black dotted lines = 95% confidence interval. *note: small sample size (n = 11)

for this year (Alcock 2013a). B) Distribution of head widths of foraging males (n = 100) reported in

Alcock et al. 1977a shows larger median head widths than C) the distribution of foraging males (n =

200) collected in 2022. All males collected before 930 am.

Discussion

The mean head width of *C. pallida* males engaged in foraging (e.g., total population), hovering, and

patrolling has declined at Blue Point Bridge over the past 50 years. The most frequent head width

class has shifted from 5.0 – 5.2 mm in 1974 to 4.6 – 4.8 mm in 2022 (foraging males, Figure 2B,C).

Head widths of the mating male population have remained constant over time, despite a significant

decline in the frequency of males of this larger size class in the population overall, suggesting that

the large-male fitness advantage continues to hold (Alcock et al. 1977a, 1977b, Alcock 1984, Alcock

1995, Alcock 2013a).

Changes in body size over time have been reported for female bees in temperate regions (Bartomeus

et al. 2013, Scheper et al. 2014, Oliveira et al. 2016, Nooten & Rehan 2020). Unlike females, males

were not reported to decline in body size in the Netherlands; Oliveira et al. (2016) propose that

fitness advantages associated with larger male body size may prevent similar declines in males.

However, the fitness advantages of larger male body sizes are unclear or nonexistent in many bee

species, particularly those without male-male competition (Alcock 2013b). Thus, in combination

with our results on *C. pallida,* this seems like an unlikely explanation for the trends observed across

bees so far. Further studies in other systems with large-male fitness advantages (Kukuk 1996,

Danforth 1991, Alcock 1983, Alcock 1994, Alcock 1997, Paxton 2005) would be beneficial to test

this hypothesis.

Sex-biased resource allocation is common in many Hymenoptera (e.g., O'Neill & O'Neill 2009),

generating male bees that are typically smaller than the females of their species (Shreeves & Field

2008). This may help explain the differences observed in male body size trends between *C. pallida*

(where this is not the case) and the other studied species in Oliveira et al. (2016). Low quantity or

quality larval nutrition significantly reduces survival and impacts adult physiology (Lawson et al.

2020, Nicholls et al. 2021), resulting in reduced mating and foraging success (Muller et al. 2015, Xie

et al. 2015). The negative fitness consequences of small body sizes may cause a species-specific

lower bound on resource allocation per offspring. As male bees are already closer to that species-

specific lower bound in resource allocation, this may prevent male bees of most species from

declining as much in body size relative to females. *C. pallida* is unique in that males can be the same

size or larger than females, allowing for more significant size declines in the male population.

Alternately, other life history or ecosystem characteristics between the bees studied in Oliveira et al.

(2016) and *C. pallida* may be responsible for variation in reported male body size declines. As a large-

bodied, solitary, desert-adapted bee species with narrow phenological and dietary breadth, *C. pallida*

may be particularly susceptible to human-activity induced declines. Hypotheses for bee body size

declines include: 1) habitat simplification or agricultural intensification, 2) climate-induced

phenological mismatches between bees and host plants, or changes in total resource availability (e.g.,

lower floral abundance due to more persistent droughts), and 3) increasing temperatures during

development (Chole et al. 2019; Sánchez-Bayo et al. 2019). Although we do not test these

hypotheses in this report, we provide contextual information that may support particular causes of

C. pallida body size declines.

The area where *C. pallida* have been studied is surrounded by land managed by the United States

Forest Service, with minimal development since the 1970s. It is unlikely that habitat simplification or

agricultural intensification is responsible for body size declines. Soil significantly buffers ground

temperature fluctuations at 10 cm depth (Parton & Logan 1981, Cane & Neff 2011; approximately

C. pallida nest depth, Alcock et al. 1976a), which should lessen the impact of increasing temperatures

on development and thus body size.

Resource limitation may occur via a combination of phenological mismatch and climate-induced

reductions in floral availability. Phenological mismatch, which may arise more frequently for ground-

nesting bees with short foraging periods (Stemkovski et al. 2020), is a possible driver of reduced

body sizes in *C. pallida*. Though inconsistently documented, it is likely that the emergence dates of *C.*

pallida have shifted earlier since the 1970-80s. The female flight season was documented as late May

to mid-June in Alcock et al. (1976a); this year (2022), the beginning of the flight season was late

April.

More data are available on dates of mating aggregations than female foraging behavior at this site

and support the idea of a two week advance in phenology since the 1980s. In 1982, peak emergence

is between April 30 – May 14 (Alcock 1984). Peak emergence is even earlier in 2012: Alcock (2013a)

suggests peak activity around April 25-May 4. By 2022, we found peak mating between April 20-May

4, an additional five days sooner. Generalist bees in the Northeastern United States have advanced

in phenology by ~10 days over the last 130 years, with most of that advance occurring after 1970

(Bartomeus et al. 2011); this aligns with shifts observed in the *C. pallida* mating aggregation activity.

However, phenological mismatch is unlikely to be the sole cause of body size declines – studies of

native bees have demonstrated that they largely advance at a pace similar to their host plants

(Bartomeus et al. 2011). In addition, data on the flowering periods of the three most common host

plants for *C. pallida*, collected by the Arizona-Sonora Desert Museum in Tucson from 1983-2009,

does not suggest an obvious mismatch in flowering and emergence dates (Arizona-Sonora Desert

Museum, 2022).

Resource limitation could affect male body size by 1) decreasing female body sizes (maternal and

offspring body sizes are linked in *C. pallida*; Alcock 1979) or 2) altering female resource allocation

decisions. Unfortunately, no data are available on female *C. pallida* body sizes at Blue Point Bridge

prior to 2022, and this relationship is expected to be population-dependent (Alcock 1979). However,

body size can affect female flight velocity and pollen carrying capacity (Müller et al. 2006, Everaars

et al. 2018), which may have dramatic effects on the number or size of offspring of different types in

C. pallida. Alcock et al. (1977; Alcock 1979) propose that a females' assessment of the likelihood of

nest failure due to parasitism or resource limitation may also affect her decision to allocate for more,

smaller (risk-averse strategy) versus fewer, larger male offspring. Notably, resource limitation seems

more likely than parasitism: rates of parasitism by bee flies, blister beetles, and mutillids are notably

low (Alcock 1979), and no kleptoparasite bees are known to parasitize *C. pallida* nests (Rozen &

Buchmann 1990).

Irrespective of the mechanism responsible for declines in male *C. pallida* body size, reductions in the

frequency of large-morph males over time may have many important consequences for the species.

First, the opportunity to pass on genetic information may become increasingly limited to an ever-

smaller pool of males capable of winning competitions (the largest males may mate multiple times;

Alcock 1995). Second, the stability of the *C. pallida* male ART system, which likely relies on

competing selective forces related to nesting density, female provisioning behavior, and male mating

success (Alcock 1979), may be threatened by declines in the species's mean body size. The impacts

of human activities often consider biodiversity loss at the level of the species, however losses in

intraspecific diversity (variation in behavior or morphology) are currently underexplored (see

Bolnick et al. 2003). ARTs represent the functional potential a species has to adapt to a changing

environment (Oliviera et al. 2016); losses in ecologically functional intraspecific diversity, which

might occur if size-based ARTs are destabilized, should be considered when evaluating the impact of

human activities. Additional work following *C. pallida* male and female body sizes at the Blue Point

Bridge population may allow for the mechanisms of body size decline in bees broadly, and *C. pallida*

specifically, to be further tested, as human-induced climate and landscape modifications continue to

drive morphological and behavioral changes in a variety of species.

Acknowledgements

We thank John Alcock for his record of *C. pallida* head widths at Blue Point over the years, and

Stephen Buchmann for his kindness and valuable *C. pallida* wisdom, which have made *C. pallida* an

accessible system for many early career bee researchers; his comments also greatly improved an early

draft of this paper. We thank Victoria Morgan for helping collect head widths on May 8[th]. Support

was provided to MGJ by Friends of the Sonoran Desert. MB is an NSF postdoctoral fellow

(2109399): any opinions, findings, conclusions, or recommendations expressed in this manuscript

are the authors, and do not necessarily reflect the views of the NSF.

References

Alcock J (1976) The social organization of male populations of *Centris pallida* (Hymenoptera,

Anthophoridae). Psyche 83:121-131.

Alcock J (1983) Male behaviour in two bumblebees, *Bombus nevudensis auricomus* and *B. griseicollis*

(Hymenoptera: Apidae). J Zool 200 561–570.

Alcock J (1984) Long-term maintenance of size variation in populations of *Centris pallida*

(Hymenoptera: Anthophoridae). Evol 38:220-223.

Alcock J (1989) Size variation in the anthophorid bee *Centris pallida:* New evidence on its

long-term maintenance. J Kansas Entomol Society 62:484-489.

Alcock J (1994) Body size and its effects on male-male competition in *Hylaeus alcyoneus*

(Hymenoptera: Colletidae). J Insect Behav 8: 149-159.

Alcock J (1995) Persistent size variation in the anthophorine bee *Centris pallida* (Apidae) despite a

large male mating advantage. Ecol Entomol 20: 1-4.

Alcock J (1997) Competition from large males and the alternative mating tactics of Dawson's

308   burrowing bee, *Amegilla dawsoni* (Apidae, Apinae, Anthophorini). Journal Insect Behav 10:

309   99e113.

310   Alcock J (2013a) Role of body size in the competition for mates by males of *Centris pallida*

311   (Anthophorinae: Hymenoptera). Southwest Nat 58:427-430.

312   Alcock J (2013b) Sexual selection and the mating behavior of solitary bees. In: Advances in the

313   Study of Behavior 45: 1-48.

314   Alcock J, Jones CE, Buchmann SL (1976a) The nesting behavior of three species of *Centris* bees

315   (Hymenoptera: Anthophoridae). J Kansas Entomol Soc 49: 469-474.

316   Alcock J, Jones CE, Buchmann SL (1976b) Location before emergence of the female bee, *Centris*

317   *pallida*, by its male (Hymenoptera: Anthophoridae). J Zool Lond 179: 189-199.

318   Alcock J, Jones CE, Buchmann SL (1977a) Male mating strategies in the bee *Centris pallida* Fox

319   (Anthophoridae: Hymenoptera). The American Naturalist 111:145-155.

320   Alcock J, Eickwort GC, Eickwort KR (1977a) The reproductive behavior of *Anthidium maculosum*

321   (Hymenoptera: Megachilidae) and the evolutionary significance of multiple copulation by

322   females. Behav Ecol Sociobiol 2: 385e396.

323   Arizona-Sonora Desert Museum (2022, May 18) Phenology – Species. *Desert Museum Digital Library*.

324   Retrieved from: http://www.desertmuseumdigitallibrary.org/public/phenology.php

325   Barrett M (2022) Alternative reproductive tactics associated with variable sensory biology and

326   thermal physiology in male *Centris pallida* (Hymenoptera: Apidae) bees. PhD Dissertation.

327   Drexel University.

328   Barrett M, Schneider S, Sachdeva P, Gomez A, Buchmann S, O'Donnell S (2021)

329   Neuroanatomical differentiation associated with alternative reproductive tactics in

male arid land bees, *Centris pallida* and *Amegilla dawsoni*. J Comp Physiol A 207:497-504.

Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R (2011) Climate-

associated phenological advances in bee pollinators and bee-pollinated plants. PNAS 108:

20645-20649.

Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R (2013)

Historical changes in northeastern US bee pollinators related to shared ecological traits.

PNAS 110:4656–4660.

Beltrán I, Herculano-Houzel S, Sinervo B, Whiting MJ (2021) Are ectotherm brains vulnerable

to global warming? Trends Ecol Evol 36:691-699.

Bolnick D, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML (2003) The

ecology of individuals: Incidence and implications of individual specialization. Am Nat

161: 1-28.

Burdine JD, McCluney KE (2019) Differential sensitivity of bees to urbanization-driven changes in

body temperature and water content. Sci Reports 9: 1643.

Burkle LA, Marlin JC, Knight TM (2013) Plant-pollinator interactions over 120 years: Loss of

species, co-occurrence, and function. Science 339: 1611-1615.

Burraco P, Orizaola G, Monaghan P, Metcalfe NB (2020) Climate change and ageing in

ectotherms. Glob Change Biol 26: 5371-5381.

Cane JH, Minckley RL, Kervin LJ, Roulston TH, Williams NM (2006) Complex responses

within a desert bee guild (Hymenoptera: Apiformes) to urban habitat fragmentation.

Ecological Applications 16: 632-644.

Cane JH, Neff JL (2011) Predicted fates of ground-nesting bees in soil heated by wildfire: Thermal

tolerances of life stages and a survey of nesting depths. Biological Conservation 144: 2631-

2636.

Chole H, Woodard SH, Bloch G (2019) Body size variation in bees: Regulation, mechanisms, and

relationship to social organization. Current Opinion in Insect Science, 35: 77-87.

Chou CC, Perez DM, Johns S, Gardner R, Kerr RA, Head ML, McCullough EL, Backwell PRY

(2019) Staying cool: The importance of shade availability for tropical ectotherms. Behav

Ecol Sociobiol 73:106.

Chown SL, Gaston KJ (2010) Body size variation in insects: a macroecological perspective. Biol Rev

Camb Philos Soc 85:139–169.

Chung DJ, Schulte PM (2020) Mitochondria and the thermal limits of ectotherms. J Exp Biol

223:jeb227801.

Danforth B (1991) The morphology and behavior of dimorphic males in *Perdita portalis*

(Hymenoptera: Andrendiae). Behav Ecol Sociobiol 29: 235-247.

Duffy GA, Coetzee BW, Janion-Scheepers C, Chown SL (2015) Microclimate-based

macrophysiology: Implications for insects in a warming world. Curr Opin Insect Sci 11:84-

89.

Everaars J, Settele J, Dormann CF (2018) Fragmentation of nest and foraging habitat affects time

budgets of solitary bees, their fitness and pollination services, depending on traits: Results

from an individual-based model. PLoS One 13: e0188269.

Gérard M, Martinet B, Maebe K, Marshall L, Smagghe G, Vereecken NJ, Vray S, Rasmont P,

Michez D (2019) Shift in size of bumblebee queens over the last century. Glob Change Biol

26: 1185-1195.

Grab H, Brokaw J, Anderson E, Gedlinske L, Gibbs J, Wilson J, Loeb G, Isaacs R, Poveda K (2019)

Habitat enhancements rescue bee body size from the negative effects of landscape

simplification. J App Ecol 56: 2144 – 2154.

GraphPad Software. 2021. GraphPad Prism v 9.3.1 for Windows. La Jolla, CA.

Hamblin AL, Youngsteadt E, López-Uribe MM, Frank SD (2017) Physiological thermal limits

predict differential responses of bees to urban heat-island effects. Biol Lett 13:20170125.

Huey RB, Kingsolver JG (2019) Climate warming, resource availability, and the metabolic

meltdown of ectotherms. Am Nat 194:E140-E150

Huey RB, Stevenson RD (1979) Integrating thermal physiology and ecology of ectotherms: A

discussion of approaches. Am Zool 19:357-366.

Kleijn D, Raemakers I (2008) A retrospective analysis of pollen host plant use by stable and

declining bumble bee species. Ecology 89:1811–1823.

Kuhlmann M, Guo D, Veldtman R, Doanldson J (2012) Consequences of warming up a hotspot:

Species range shifts within a centre of bee diversity. Diversity Distrib 18: 885-897.

Kukuk PF (1996) Male dimorphism in *Lasioglossum (Chilalictus) hemichalceum*: the role of larval

nutrition. J Kans Entomol Soc 69:147-157.

Lawson SP, Helmreich SL, Rehan SM (2017) Effects of nutritional deprivation on development and

behavior in the subsocial bee *Ceratina calcarata* (Hymenoptera: Xylocopinae). J Exp Biol 220:

4456-4462.

Lawson SP, Kennedy KB, Rehan SM (2020) Pollen composition significantly impacts the

development and survival of the native small carpenter bee, *Ceratina calcarata*. Ecol Entomol

46: 232-239.

Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of climate

change. Nature 462:1052-1055.

McCabe LM, Aslan CE, Cobb NS (2021) Decreased bee emergence along an elevation gradient:

Implications for climate change revealed by a transplant experiment. Ecology 103: e03598.

Nooten SS, Rehan SR (2020) Historical changes in bumble bee body size and range shift of

declining species. Biodiversity and Conservation 29: 451-467.

Muller K, Thiéry D, Moret Y, Moreau J (2015) Male larval nutrition affects adult reproductive

success in wild European grapevine moth (*Lobesia botrana*). Behav Ecol Sociobiol 69: 39-47.

Müller A, Diener S, Schnyder S, Stutz K, Sedivy C, Dorn S (2006) Quantitative pollen requirements

of solitary bees: implications for bee conservation and the evolution of bee-fower

relationships. Biol Conserv 130:604–615.

Nicholls E, Rossi M, Niven JE (2021) Larval nutrition impacts survival to adulthood, body size and

the allometric scaling of metabolic rate in adult honeybees. J Exp Biol 224: jeb242393.

Oliveira MO, Freitas BM, Scheper J, Kleijn D (2016) Size and sex-dependent shrinkage of dutch

bees during one-and-a-half centuries of land-use change. PLoS ONE 11:e0148983.

Ockendon N, Baker DJ, Carr JA, White EC, Almond REA, Amano T… Pearce-Higgins JW (2014)

Mechanisms underpinning climatic impacts on natural populations: altered species

interactions are more important than direct effects. Glob Change Biol 20:2221–2229.

Oliveira RF, Taborsky M, Brockmann HJ (2008) Alternative reproductive tactics: an integrative

approach. Cambridge University Press, Cambridge.

O'Neill KM, O'Neill RP (2009) Prey, nest associates, and sex ratios of *Isodontia mexicana*

(Saussure) (Hymenoptera: Sphecidae) from two sites in New York. Entomologica

Americana 115: 90-94.

Parmesan C (2006) Ecological and evolutionary responses to recent climate change. Ann Rev

Ecol Evol Systematics 37:637-669.

Parton WJ, Logan JA (1981) A model for diurnal variation in soil and air temperature.

Agricultural Meteorology 23: 205-216.

Paxton RJ (2005) Male mating behaviour and mating systems of bees: an overview. Apidologie

36:145-156.

Rozen JG, Buchmann SL (1990) Nesting biology and immature stages of the bees *Centris caesalpiniae,*

C. pallida, and the cleptoparasite *Ericrosis lata* (Hymenoptera: Apoidea: Anthophoridae).

American Museum Novitates, 2985: 30 pp.

Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, et al. (2000) Global biodiversity

scenarios for the year 2100. Science 287: 1770-1774.

Sánchez-Bayo F, Wyckhuys KAG (2019) Worldwide decline of the entomofauna: A review of its

drivers. Biological Conservation 232: 8-27.

Scheper J et al (2014) Museum specimens reveal loss of pollen host plants as key factor driving wild

bee decline in the Netherlands. Proc Natl Acad Sci USA 111:17552–17557.

Shreeves G, Field J (2008) Parental care and sexual size dimorphism in wasps and bees. Behav Ecol

Sociobiol 62: 843-852.

Shuster SM (2010) Alternative mating strategies. In: C. Fox and D. F. Westneat (eds.) *Evolutionary*

behavioral ecology. Cambridge University Press, Cambridge.

Silva DP, Dew RM, Vilela B, Stevens MI, Schwarz MP (2018) No deaths in the desert: Predicted

responses of an arid-adapted bee and its two nesting trees suggest resilience in the face of

warming climates. Insect Conservation and Diversity 11: 449-463.

Sinervo B, Mendez-de-la-Cruz F, Miles DB, Heulin B, Bastiaans E, Villagran-Santa Cruz M, et al.

(2010) Erosion of lizard diversity by climate change and altered thermal niches. Science

328:894–899.

Snelling RR (1984) Studies on the taxonomy and distribution of American Centridine bees

(Hymenoptera: Anthophoridae). Contributions in Science 347:1-69.

Stemkovski M, Pearse WD, Griffin SR, Pardee GL, Gibbs J, Griswold T, Neff JL, Oram R, et al.

(2020) Bee phenology is predicted by climatic variation and functional traits. Ecology Letters

23: 1589-1598.

Vale CG, Brito JC (2015) Desert-adapted species are vulnerable to climate change: Insights from the

warmest region on Earth. Global Ecol Conserv 4: 369-379.

Walters RJ, Hassall M (2006) The temperature-size rule in ectotherms: May a general explanation

exist after all? Am Nat 167:510-523.

Warzecha D, Diekötter T, Wolters V, Jauker F (2016) Intraspecific body size increases with habitat

fragmentation in wild bee populations. Landscape Ecology 31: 1449-1455.

Wieser W (1973) "Temperature Relations of Ectotherms: A Speculative Review." In: Wieser W.

(eds) *Effects of Temperature on Ectothermic Organisms*. Springer, Berlin, Heidelberg.

Xie J, de Clercq P, Pan C, Li H, Zhang Y, Pang H (2015) Larval nutrition-induced plasticity affects

reproduction and gene expression of the ladybeetle, *Cryptolaemus montrouzieri*. BMC Ecol Evol

15:276.

Zattara EE, Aizen MA (2021) Worldwide occurrence records suggest a global decline in bee species

richness. One Earth 4: 114-123.

Supplemental Figures

Supplementary Figure 1. Distribution of female head widths in 2022. Distribution of head

widths of females (n = 114) collected while mating, foraging, and nesting from before 1130 on April

20 – May 8, 2022, at Blue Point Bridge, Arizona.

Supplementary Figure 2. Increase in patrolling or foraging male head widths from 0700 to

1130. Head width in the populations of males patrolling (grey squares, dashed line, n = 410) and

foraging (black dots, solid line, n = 203) increased over the course of the morning (Supplementary

Figure 2; linear regression, patrolling: [head width] = 0.095 [time of day] + 4.19, F = 23.65, df =

408, R^2 = 0.05, p < 0.0001; foraging: [head width] = 0.097 [time of day] + 4.08, F = 7.90, df = 201,

R^2 = 0.03, p < 0.0001).

Supplementary Figure 3. Changes in head width distributions of hovering and patrolling

males from 1974 to 2022. Head width distributions for hovering (n = 57) and patrolling (n = 300)

males at Blue Point Bridge are skewed towards smaller body sizes in 2022 as compared to

1974/1975 (n = 50 hoverers, 100 patrollers; data reproduced from Alcock et al. 1977a). All males

collected before 930 am.

Supplementary Figure 4. Similar head width distributions of mating males from 1974 to

2022. Head width distributions for mating males at Blue Point Bridge demonstrates the population

of males that obtain mating opportunities have relatively similar body sizes in 2022 (n = 56) as

compared to 1974/1975 (n = 50; data reproduced from Alcock et al. 1977a). All males collected

before 0930 am.