Title: *Centris pallida* (Hymenoptera: Apidae) male body size decreases across five decades

Running Title: Declines in male *C. pallida* body size

Authors: Meghan Barrett[1], Meredith G. Johnson[2]

[1]Department of Biology, California State University Dominguez Hills, Carson, CA 90747

[2]School of Life Sciences, Arizona State University, Tempe, AZ 85281

Authorship Contributions: MB – conceptualization, data curation, funding acquisition, formal

analysis, investigation, methodology, writing – original draft; MGJ – investigation, funding

acquisition, writing – review & editing

Conflict of Interests: None.

Abstract

Historical data suggest that many bee species have declined in body size. Larger-bodied bees with narrow phenological and dietary breadth are most prone to declines in body size over time. This may be especially true in solitary, desert-adapted species that are vulnerable to climate change – such as *Centris pallida* (Hymenoptera: Apidae). In addition, body size changes in species with size-linked behaviors could threaten the prevalence of certain behavioral phenotypes long-term. *C. pallida* solitary bees are found in the Sonoran Desert. Males use alternative reproductive tactics (ARTs) and are dimorphic in both morphology and behavior. *C. pallida* male body size has been studied since the 1970s in the same population. We collected body size data in 2022 and combined it with published records from 1974-2022. We find a persistent decline in the mean head width of patrolling males, and shifts towards smaller body sizes in the populations of males found foraging and hovering. Both morphs declined in average body size, and the proportion of large-morph males in the population decreased by 8%. Mating males did not decline in mean body size over the last five decades. We discuss hypotheses related to the decline in *C. pallida* male head width. Finally, we advocate for *C. pallida* as an excellent study system for understanding the stability of ARTs with size-linked behavioral phenotypes.

Introduction

Anthropogenic changes in climate and habitat have caused a decline in abundance, a shift in

geographic range or phenology, and changes in the morphology, physiology, or behavior of many

species, sometimes leading to alterations in life history and ecological relationships (Bartomeus et al.

2011, Biesmeijer et al. 2006, Jacobson et al. 2018, Béltran et al. 2021, Burraco et al. 2020, Chou et al.

2019, Chung & Schulte 2020, Duffy et al. 2015, Huey & Kingsolver 2019, Kuhlmann et al. 2012,

Ockendon et al. 2014, Parmesan 2006, Sinervo et al. 2010, Walters & Hassall 2006, Zattara & Aizen

2021, Sánchez-Bayo & Wyckhuys 2019, Burkle et al. 2013, Turley et al. 2022).

Bees may be particularly vulnerable to climate change or habitat modifications. As heterothermic

ectotherms they rely on environmental conditions to maintain non-lethal body temperatures (Wieser

1973, Huey & Stevenson 1979). This is compounded by a reliance on resources (e.g., nesting

resources, plants) that are often disturbed by human activity (Grab et al. 2019, McCabe et al. 2021).

Documented changes in bee populations over time can help determine how human activity may

impact bees with different life history characteristics, with knock-on effects for their ecological

relationships. (e.g., Cane et al. 2006, Burkle et al. 2013). For example, studies in Europe and the

Northeastern regions of North America have demonstrated that larger-bodied bee species are more

likely to experience declines in population (Bartomeus et al. 2013, Scheper et al. 2014, Oliveira et al.

2016), or stronger intraspecific declines in body size (Nooten & Rehan 2020, but see Gérard et al.

2019), perhaps due to greater nutritional requirements (Müller et al. 2006).

Bee body sizes have not changed uniformly over time across species, sexes, or regions (Nooten &

Rehan 2020, Bartomeus et al. 2013, Scheper et al. 2014, Oliveira et al. 2016, Kleijn & Raemakers

2008), suggesting life history traits and ecosystem characteristics (e.g., habitat fragmentation,

Warzecha et al. 2016) may play important roles in determining the impact of human activities on

body size. For most bees, body size is determined by resource provisioning in the larval stage (Chole

et al. 2019, Kukuk 1996, Alcock 1984, Lawson et al. 2017), and can thus be strongly influenced by

changes in resource availability (Chown & Gaston 2010). Bees with narrow phenological windows

and bees that are dietary specialists are likely to be more susceptible to declines, given their reliance

on a smaller pool of available resources (Bartomeus et al. 2013). Solitary and desert-adapted bees are

expected to be particularly vulnerable to climate change (Sala et al. 2000, Loarie et al. 2009, Vale &

Brito 2015, Hamblin et al. 2017, Burdine & McCluney 2019, McCabe et al. 2021, but see Silva et al.

2018), given the thermal and hygric stressors already evident in their environment. To date, no

studies have tracked changes in morphology of any solitary bees in desert ecosystems.

Declines in body size could also generate species-level behavioral changes whenever size contributes

to behavior, such as in many alternative reproductive tactic systems (ARTs). ARTs occur when

categorical variation in the morphological/behavioral traits is associated with mating across

individuals of the same sex within the same population (Oliveira et al. 2008, Paxton 2005, Shuster

2010). In many such systems, larger- and smaller-bodied individuals use different strategies for

accessing mates (e.g., fighters vs. sneakers; Oliveira et al. 2008). Declines in body size may alter the

occurrence of size-linked morphs, and their behaviors, resulting in a loss of important intraspecific

variation over time.

Centris pallida (Hymenoptera: Apidae) are widespread, common solitary digger bees are found in the

Sonoran Desert of the Southwestern United States and Northern Mexico. This species often forms

dense nesting aggregations, where many thousands of individuals occupy several hectares. *C. pallida*

mating aggregations may persist in the same locations over multiple decades – allowing for the

continuous resampling of a population (Barrett 2022). Males emerge first from their natal nests at

these sites and use ARTs to find mates. Males are behaviorally and morphologically dimorphic, using

different sensory mate location strategies and microclimates (Alcock 1976, Alcock et al. 1977a,

Snelling 1984, Barrett et al. 2021, Barrett 2022). Large-morph, 'metandric' males with pale gray

coloration patrol in sinuous loops ~10 cm over the emergence site and use scent to locate females

emerging from natal nests; males then engage in fights for the opportunity to dig up and mate with

emerging females (Alcock et al. 1976b). Small-morph males, with dark brown coloration, are more

behaviorally flexible; they may patrol, but are often found hovering near plants where they use visual

cues to locate females or mating pairs flying away from the aggregation site that may be interrupted.

Large-morph males have a clear fitness advantage in situations where mating aggregations are

densely populated (Alcock et al. 1977a, 1977b, Alcock 1984, Alcock 1995, Alcock 2013a), as is

always the case in this specific *C. pallida* population.

C. pallida are large-bodied, desert-adapted, solitary bees with an extremely narrow phenological and

dietary breadth. Bees nest for less than six weeks in the late spring, and females utilize four species

of flowering trees for larval nectar/pollen provisions (*Parkinsonia microphylla, P. aculeata, Olneya tesota,*

and, rarely, *Psorothamnus (Dalea) spinosa*; Alcock et al. 1976a). *P. microphylla* is the predominant host

plant for *C. pallida* over its entire geographic range (Buchmann, pers. comm). The body sizes of male

C. pallida have been studied at the same site since for 48 years (Alcock et al. 1977a). By combining

historical data on *C. pallida* body sizes from 1974 – 2018 (Alcock et al. 1977a, Alcock 1984, Alcock

1989, Alcock 2013a, Barrett 2022) with data we collected in 2022 from the same population, we

aimed to determine if male *C. pallida* bees have experienced persistent changes or persistent stability

in body size. Our results demonstrate that the population of males found foraging at trees

(representative of the total population of males), hovering, and patrolling have all declined in body

size since 1974. However, the population of mating males has not persistently declined in body size

over time. In addition, we present the first body size data on female *C. pallida* from this site, for use

in future historical comparisons. Finally, we discuss several hypotheses of causes for the decline in

male body size, and the implications of declining male body size on the stability of the *C. pallida*

ART.

Methods

Head width measurements

We collected head widths at the same site where most previous research on *C. pallida* body size has

been conducted (Alcock et al. 1977a, Alcock 1984, Alcock 1989, Alcock 2013a), in the floodplain by

Blue Point Bridge/Saguaro Lake over the Salt River north of Mesa, Arizona (33.552, -111.566).

Alcock et al. (1977a) demonstrated that head width is correlated with body weight in *C. pallida* males,

and serves as a reliable field indicator of body size.

We collected bees (n = 921 males, 114 females) in the same manner as described in Alcock (1984,

2013), to be sure our data were comparable. Briefly: for patrolling males, we made rapid sweeps low

to the ground with an insect net through open areas of searching males. We approached and

collected digging/fighting males by hand. We collected foraging males and females with a

telescoping insect net as they visited paloverde trees (*Parkinsonia* sp.). We collected mating males and

females while engaged in copulation on the ground or vegetation. We collected hovering males near

mesquite or palo verde trees, wherever they had established stable aerial stations. We measured head

widths to 0.01 mm using digital calipers (Wen 10761), before releasing the bees. We collected bees

daily when the aggregation was most active between April 20 and May 8, 2022, generally between

the hours of 0700 and 1130, and noted the time of collection and behavior for each individual.

Historical data for comparisons

To obtain historical male head width data, we used reported means and standard deviations in

Alcock et al. 1977a, Alcock 1984, Alcock 2013a, and Barrett 2022. Together, these studies report

data collected in: 1974, 1975, 1976, 1982, 1988, 2011, 2012, and 2018. As prior surveys of the

population by Alcock (et al. 1977; Alcock 1984, 1989, 2013a) generally only captured bees between

roughly 0700 and 0930, and did not note the specific individual time of capture. We therefore also

used only those bees captured before 0930 in our sampling (2022) for the historical comparison with

prior literature. Barrett (2022) reported data for mating, hovering, and patrolling bees from this site

in 2018, but patrolling bees were not captured in a manner consistent with Alcock's prior surveys

and so we elected to only use the mating and hovering male datasets in our current comparison.

<u>Changes in the Proportion and Size of Each Morph</u>

To determine changes in the proportion of large- and small-morph males in the total population

(represented by males caught foraging, Alcock et al. 1977a), we used the head width frequency

distributions. These distributions are bimodal, with the two local maxima representing the two

morphs; there is thus a local minimum in frequency between the small- and large-morph males (see

Fig 2, where an arrow indicates the local minimum). The proportion of all males in the foraging

population that were smaller in head width than the local minimum was assumed to be the

proportion of small-morph males in the population. Those males larger in head width than the local

minimum were assumed to be large-morph males. The proportion within the local minimum were

considered the rare intermediate males that occur in all *Centris* species with male dimorphism (only

5-6% in both the 1974/5 and 2022 samples).

To determine morph-specific changes in body size, we assessed shifts in the head width frequency

distributions on either side of the local minimum. We corroborated this shift in small-morph male

body size by looking at changes in the head widths of hovering males, specifically. Hovering is

completed *only* by small-morph males, and therefore can be used as a secondary method for

assessing body size changes within this morph.

<u>Statistical analysis</u>

Data were analyzed using GraphPad Prism 9.3.1 (GraphPad Prism for Windows 2021). When

analyzing data on the total population in 2022 (as compared to the before-0930 dataset used for the

historical data comparison), we include all bees captured in 2022 (including males for which we had

not taken data on time of capture). Because head width data were not normally distributed for all

behavior categories, we used a Kruskal-Wallis test with post-hoc Dunn's MCT to analyze differences

between hovering, patrolling, and mating male populations. We used a Mann-Whitney test to

compare mating and foraging male populations, and an unpaired t-test to compare digging and

fighting males and mean hovering male head widths from 1974/1975 and 2022. We used a linear

regression to analyze the effect of time of day on patrolling and hovering male head widths. We also

used a linear regression to analyze the effect of year on mean patrolling and mating male head width

across studies, using the reported mean, standard deviation (SD), and population size (n) of each

study. All data can be found archived on Dryad (Barrett & Johnson 2022).

Results

C. pallida head widths in 2022

The ranges and frequencies of male head widths for the 2022 population of males engaged in

foraging, patrolling, hovering, mating, digging, and fighting behaviors are shown in Figure 1 (mean

± SD and n, reported in Supplementary Table 1). We also present mean female head widths in this

population, for females caught while mating, foraging, or nesting in Supplementary Table 1

(distribution of female head widths: Supplemental Figure 1).

Hovering males were significantly smaller than patrolling males (Figure 1; Kruskal-Wallis: K-W =

140.4, $p < 0.0001$; Dunn's MCT: $Z = 6.51$, $p < 0.0001$), and both types of males were smaller than

mating males (patrolling vs. mating: $Z = 8.70$, $p < 0.0001$; hovering vs. mating: $Z = 11.83$, $p <$

0.0001). Mating males skewed larger than the total population of foraging males (Mann-Whitney

test: $U = 1439$, $p < 0.0001$). There was no difference in the mean head widths of digging and

fighting males (unpaired t-test: t = 1.55, df = 135, p = 0.12).

Figure 1. Distribution of male head widths by behavioral category in 2022. Distribution of

male head widths (n = 921) collected while foraging (n = 203), hovering (n = 91), patrolling (n =

410), mating (n = 71), and digging/fighting (n = 66/71) before 1130 on April 20 – May 8, 2022 at

Blue Point Bridge, Arizona.

Patrolling and foraging male head widths increased over the course of the morning (Supplementary

Figure 2; linear regression, patrolling: [head width] = 0.095 [time of day] + 4.19, F = 23.65, df =

408, R^2 = 0.05, p < 0.0001; foraging: [head width] = 0.097 [time of day] + 4.08, F = 7.90, df = 201,

R^2 = 0.03, p < 0.0001); hovering and mating male head widths were constant across the morning

(hovering: F = 1.57, df = 88, p = 0.21; mating: F = 0.43, df = 57, p = 0.51).

<u>Historical comparison of mean male head widths from 1974 to 2022</u>

Reported head widths for males foraging, hovering, patrolling, or mating at the Blue Point Bridge

site from 1974 to 2022 can be found in Table 1, with associated references for the historical data.

Table 1. Reported means ± SD of males at Blue Point Bridge before 930 from 1974 – 2022.

	Mean ± SD (n)				
Year	Foraging	Hovering	Patrolling	Mating	Reference
1974 + 1975	no mean reported; Supp Fig 3	4.92 ± 0.15 (50)	5.14 ± 0.29 (100)	5.51 ± 0.36 (50)	Alcock et al. 1977a
1976			5.20 ± 0.28 (168)	5.44 ± 0.33 (102)	Alcock 1984, 1989
1981			5.15 ± 0.23 (162)		Alcock 1984
1982			5.11 ± 0.25 (265)	5.42 ± 0.29 (140)	Alcock 1984, 1989
1988			5.09 ± 0.29 (400)	5.47 ± 0.29 (144)	Alcock 1989
2011			5.02 ± 0.28 (214)		Alcock 2013a
2012	4.88 ± 0.31 (132)		4.95 ± 0.25 (285)	5.32 ± 0.29 (11[*])	Alcock 2013a
2018		4.86 ± 0.21 (11[*†])		5.40 ± 0.31 (23[*†])	Barrett 2022
2022	4.84 ± 0.36 (200)	4.77 ± 0.19 (57)	4.98 ± 0.34 (300)	5.50 ± 0.31 (56)	

[*]small sample size.
[†]these bees were not all collected before 0930, as in the other years, but no individual data on time of capture was recorded. However
body size was not affected by time of day in hovering and mating male populations, so we included these males.

The mean head width of the foraging males (n = 100; representing the total population of

males), was not reported in Alcock et al. 1977a. However, comparing the distribution of head widths

from 1974/1975 to our 2022 data suggests significant declines in overall male head width (Figure 2B

& 2C). The most frequent head width class declines from 5.0 – 5.22 mm to 4.6 – 4.8 mm (which was

previously the smallest size class). The smallest size class observed in the male population decline

from 4.6 – 4.8 mm to 4.0 – 4.2 mm. The two largest size classes (5.8 – 6.2 mm) were lost entirely

from the foraging male distribution despite surveying twice the number of foraging males in 2022

compared to 1974/1975. These two size classes decreased from ~30% of all mating events in

1974/1975 to ~10% in 2022, demonstrating a significant decrease in the largest size large-morph

males.

Additionally, there were within-morph body size declines for both morphs. The most

frequent head width for small-morph males (to the left of the arrow in Figure 2B/2C) in 1974/1975

was 5.0 – 5.2 mm; by 2022, the peak was 4.6 – 4.8 mm (previously, the smallest size category for all

males). For large-morph males (to the right of the arrow) the peak frequency also decreased, from

5.8 – 6.0 mm to 5.4 – 5.6 mm.

Behaviorally-linked data corroborate this decrease in body size. Patrolling male head widths,

which represent males from both morphs (but is skewed towards larger small-morph, and large-

morph, males compared to the overall population of foraging males), decreased by 3.72% at the

Blue Point Bridge site from 1974 to 2022 (Figure 2A, Supplementary Figure 3B; linear regression,

[head width] = -0.004344 [year] + 13.74, F = 131.2, df = 1892, R^2 = 0.06, p < 0.0001). There was a

3.05% decrease in mean hovering male head width from 1974/1975 to 2022 (Supplementary Figure

3A, unpaired t-test; t = 4.49, df = 105, p < 0.0001); hovering is exclusively performed by small-

morph males, corroborating the decrease in head width shown in the foraging male frequency

distribution data.

Small-morph males increased from 73% of the total male (foraging) population in

1974/1975 to 82% in 2022. Large-morph males (to the right of the arrow) decreased from 21% to

13%; intermediates were rare and represented only 5-6% of all males in each year.

Figure 2. Changes in head widths of patrolling, foraging, and mating males from 1974 to

2022. A) Patrolling male head widths (gray dots, solid line, n = 300) decreased over time at the Blue

Point Bridge site from 1974 to 2022 (Figure 2; linear regression, [head width] = -0.004 [year] +

13.74, F = 131.2, df = 1892, R^2 = 0.06, p < 0.0001). Mating male head widths (black diamonds,

dashed line to demonstrate non-significance, n = 56) did not change from 1974 to 2022 (F = 0.36,

df = 501, p = 0.55). Black dotted lines = 95% confidence interval. *note: small sample size (n = 11)

for this year (Alcock 2013a). B) Distribution of head widths of foraging males (n = 100) reported in

Alcock et al. 1977a shows larger median head widths than C) the distribution of foraging males (n =

200) collected in 2022. All males collected before 930 am. The arrow represents the rare

intermediate males (5-6%; gray column, local minimum); to the left of this head width class are the

small-morph males, to the right are the large-morph males.

Discussion

The mean head width of *C. pallida* males engaged in foraging (e.g., total population), hovering, and

patrolling has declined at Blue Point Bridge over the past five decades. The most frequent head

width class among all males has shifted from 5.0 – 5.2 mm in 1974 to 4.6 – 4.8 mm in 2022

(foraging males, Figure 2B,C). Within-morph, both large- and small-morph males have seen a similar

decline of 0.4 mm in their most frequent head width.

Time-of-day affects body size for patrolling and foraging male populations, but not mating or

hovering males – foraging and patrolling males were smaller early in the morning. This may relate to

the thermal adaptations of the males: the darker coloration of small-morph males may allow them to

heat up their flight muscles faster in the cool early mornings, while the lighter coloration of large-

morph males may keep them cooler later in the morning (Barrett et al. 2022, Barrett & O'Donnell

2022). A similar coloration effect, which keeps the flight muscles at an optimal temperature while

balancing convective cooling and shortwave radiative heat gain, is seen in male butterflies with

different mating behaviors (Van Dyck & Matthysen 1998). Alternately, small-morph males may

simply emerge earlier in the day to avoid displacement by large-morph males in fights, as Alcock

(2009) hypothesized for size-dimorphic *Amegilla dawsoni* (Hymenoptera: Apidae) males that use

similar ARTs (but do not differ in coloration between the morphs). However, our data suggest this

is unlikely in *C. pallida* – males mating earlier in the morning are not any smaller in body size,

suggesting that small-morph males are not avoiding displacement by larger males simply by 'getting

up' early. Therefore, rather than sexually-selected early emergence, we favor the physiological

limitations on flight muscle temperature hypothesis. Future studies may also look for temporal

variation in male body size across the emergence season (Alcock 2009).

We also found that small-morph males were a larger proportion of the overall population in 2022

compared to 1974/1975 (from 73% to 82% of the total foraging population). Large-morph males

appear to be slowly declining within the overall population, representing a potential threat to the

longevity of the ART system in this species. Despite their numerical decline, the large-morph fitness

advantage continues to hold: head widths of the mating male population have remained constant

over time, despite a significant decline in the frequency of males of this larger size class in the

population overall (Alcock et al. 1977a, 1977b, Alcock 1984, Alcock 1995, Alcock 2013a). However,

the largest large-morph male size class (5.8 – 6.2 mm) decreased from nearly 30% of mating events

to only 10% in 2022, providing further evidence that the large-morph males are declining.

Changes in body size over time have been reported for female bees in temperate regions (Bartomeus

et al. 2013, Scheper et al. 2014, Oliveira et al. 2016, Nooten & Rehan 2020). Unlike females, males

were not reported to decline in body size in the Netherlands; Oliveira et al. (2016) propose that

fitness advantages associated with larger male body size may prevent similar declines in males.

However, the fitness advantages of larger male body sizes are unclear or nonexistent in many bee

species, particularly those without male-male competition (Alcock 2013b). Thus, in combination

with our results on *C. pallida* (where large-morph males are declining in frequency), this seems like an

unlikely explanation for the trends observed across bees so far. Further studies in other systems with

large-male fitness advantages (Kukuk 1996, Danforth 1991, Alcock 1983, Alcock 1994, Alcock 1997,

Paxton 2005) would be beneficial to test this hypothesis.

Sex-biased resource allocation is common in many Hymenoptera (e.g., O'Neill & O'Neill 2009),

generating male bees that are typically smaller than the females of their species (Shreeves & Field

2008). This may help explain the differences observed in male body size trends between *C. pallida*

(where this is not the case) and the other studied species in Oliveira et al. (2016). Low quantity or

quality larval nutrition significantly reduces survival and impacts adult physiology (Lawson et al.

2020, Nicholls et al. 2021), resulting in reduced mating and foraging success (Muller et al. 2015, Xie

et al. 2015). The negative fitness consequences of small body sizes may cause a species-specific

lower bound on resource allocation per offspring. As male bees are already closer to that species-

specific lower bound in resource allocation, this may prevent male bees of most species from

declining as much in body size relative to females. *C. pallida* is unique in that males can be the same

size or larger than females, allowing for more significant size declines in the male population. This

phenomenon may also explain the shift in the proportion of large- versus small-morph males, as

there is more room in the *C. pallida* system to decrease the high end of the male body size spectrum.

Alternately, other life history or ecosystem characteristics between the bees studied in Oliveira et al.

(2016) and *C. pallida* may be responsible for variation in reported male body size declines. As a large-

bodied, solitary, desert-adapted bee species with narrow phenological and dietary breadth, *C. pallida*

may be particularly susceptible to human-activity induced declines. Hypotheses for bee body size

declines include: 1) habitat simplification or agricultural intensification, 2) climate-induced

phenological mismatches between bees and host plants, or changes in total resource availability (e.g.,

lower floral abundance due to more persistent droughts), and 3) increasing temperatures during

development (Chole et al. 2019; Sánchez-Bayo and Wyckhuys 2019). Although we do not test these

hypotheses in this report, we provide contextual information that may support particular causes of

C. pallida body size declines.

The area where *C. pallida* have been studied is surrounded by land managed by the United States

Forest Service, with minimal development since the 1970s. It is unlikely that habitat simplification or

agricultural intensification is responsible for body size declines. Soil significantly buffers ground

temperature fluctuations at 10 cm depth (Parton & Logan 1981, Cane & Neff 2011; approximately

C. pallida nest depth, Alcock et al. 1976a), which should lessen the impact of increasing temperatures

on development and thus body size.

Resource limitation may occur via a combination of phenological mismatch and climate-induced

reductions in floral availability. Phenological mismatch, which may arise more frequently for ground-

nesting bees with short foraging periods (Stemkovski et al. 2020), is a possible driver of reduced

body sizes in *C. pallida*. Though inconsistently documented, it is likely that the emergence dates of *C.*

pallida have shifted earlier since the 1970-80s. The female flight season was documented as late May

to mid-June in Alcock et al. (1976a); this year (2022), the beginning of the flight season was late

April.

More data are available on dates of mating aggregations than female foraging behavior at this site

and support the idea of a two week advance in phenology since the 1980s. In 1982, peak emergence

was between April 30 – May 14 (Alcock 1984). Peak emergence is even earlier in 2012: Alcock

(2013a) suggests peak activity around April 25-May 4. By 2022, we found peak mating between April

20-May 4, an additional five days sooner. Generalist bees in the Northeastern United States have

advanced in phenology by ~10 days over the last 130 years, with most of that advance occurring

after 1970 (Bartomeus et al. 2011); this aligns with shifts observed in the *C. pallida* mating

aggregation activity. However, phenological mismatch is unlikely to be the sole cause of body size

declines – studies of native bees have demonstrated that they largely advance at a pace similar to

their host plants (Bartomeus et al. 2011). In addition, data on the flowering periods of the three

most common host plants for *C. pallida*, collected by the Arizona-Sonora Desert Museum in Tucson

from 1983-2009, does not suggest an obvious mismatch in flowering and emergence dates (Arizona-

Sonora Desert Museum, 2022).

Resource limitation could affect male body size by 1) decreasing female body sizes (maternal and

offspring body sizes are linked in *C. pallida*; Alcock 1979) or 2) altering female resource allocation

decisions. Unfortunately, no data are available on female *C. pallida* body sizes at Blue Point Bridge

prior to 2022, and this relationship is expected to be population-dependent (Alcock 1979). However,

body size can affect female flight velocity and pollen carrying capacity (Müller et al. 2006, Everaars

et al. 2018), which may have dramatic effects on the number or size of offspring of different types in

C. pallida. Alcock et al. (1977a; Alcock 1979) propose that a females' assessment of the likelihood of

nest failure due to parasitism or resource limitation may also affect her decision to allocate for more,

smaller (risk-averse strategy) versus fewer, larger male offspring. Notably, resource limitation seems

more likely than parasitism: rates of parasitism by bee flies, blister beetles, and mutillids are notably

low (Alcock 1979), and no kleptoparasite bees are known to parasitize *C. pallida* nests (Rozen &

Buchmann 1990).

Irrespective of the mechanism responsible for declines in overall male *C. pallida* body size,

reductions in the frequency of large-morph males, specifically, over time may have many important

consequences for the species. First, the opportunity to pass on genetic information may become

increasingly limited to an ever-smaller pool of males capable of winning competitions (the largest

males may mate multiple times; Alcock 1995). However, given that small-morph males will patrol

and mate with females whenever large-morph males are not around to compete with them, this

outcome may be unlikely in the *C. pallida* mating system.

Second, the stability of the *C. pallida* male ART system, which likely relies on competing selective

forces related to nesting density, female provisioning behavior, and male mating success (Alcock

1979), may be threatened by declines in the species's mean body size and the decreasing frequency

of large-morph males. The impacts of human activities often consider biodiversity loss at the level of

the species, however losses in intraspecific diversity (variation in behavior or morphology) are

currently underexplored (see Bolnick et al. 2003). ARTs represent the functional potential a species

has to adapt to a changing environment (Oliviera et al. 2016); losses in ecologically functional

intraspecific diversity, which might occur if size-based ARTs are destabilized, should be considered

when evaluating the impact of human activities. Additional work following *C. pallida* male and

female body sizes at the Blue Point Bridge population may allow for the mechanisms of body size

decline in bees broadly, and *C. pallida* specifically, to be further tested, as human-induced climate and

landscape modifications continue to drive morphological and behavioral changes in a variety of

species.

Acknowledgements

We thank John Alcock for his record of *C. pallida* head widths at Blue Point over the years, and

Stephen Buchmann for his kindness and valuable *C. pallida* wisdom, which have made *C. pallida* an

accessible system for many early career bee researchers; his comments also greatly improved an early

draft of this paper. We thank Victoria Morgan for helping collect head widths on May 8[th], 2022. We

also thank the editor, and two anonymous reviewers, for thoughtful comments that improved this

manuscript significantly. Support was provided to MGJ by Friends of the Sonoran Desert and MB

by the National Science Foundation. MB is an NSF postdoctoral fellow (2109399): any opinions,

findings, conclusions, or recommendations expressed in this manuscript are the authors, and do not

necessarily reflect the views of the NSF.

References

Alcock J (1976) The social organization of male populations of *Centris pallida* (Hymenoptera,

Anthophoridae). Psyche 83:121-131.

Alcock J (1979) The relation between female body size and provisioning behavior in the bee *Centris*

pallida Fox (Hymenoptera: Anthophoridae). J Kansas Entomol Soc 52:623-632.

Alcock J (1983) Male behaviour in two bumblebees, *Bombus nevudensis auricomus* and *B. griseicollis*

(Hymenoptera: Apidae). J Zool 200 561–570.

Alcock J (1984) Long-term maintenance of size variation in populations of *Centris pallida*

(Hymenoptera: Anthophoridae). Evol 38:220-223.

Alcock J (1989) Size variation in the anthophorid bee *Centris pallida:* New evidence on its

long-term maintenance. J Kansas Entomol Society 62:484-489.

Alcock J (1994) Body size and its effects on male-male competition in *Hylaeus alcyoneus*

(Hymenoptera: Colletidae). J Insect Behav 8: 149-159.

Alcock J (1995) Persistent size variation in the anthophorine bee *Centris pallida* (Apidae) despite a

large male mating advantage. Ecol Entomol 20: 1-4.

Alcock J (1997) Competition from large males and the alternative mating tactics of Dawson's

burrowing bee, *Amegilla dawsoni* (Apidae, Apinae, Anthophorini). Journal Insect Behav 10:

99e113.

Alcock J (2009) Small males emerge earlier than large males in Dawson's burrowing bee (*Amegilla*

dawsoni) (Hymenoptera: Anthophorini). J Zool 242: 453-462.

Alcock J (2013a) Role of body size in the competition for mates by males of *Centris pallida*

(Anthophorinae: Hymenoptera). Southwest Nat 58:427-430.

Alcock J (2013b) Sexual selection and the mating behavior of solitary bees. In: Advances in the

Study of Behavior 45: 1-48.

Alcock J, Jones CE, Buchmann SL (1976a) The nesting behavior of three species of *Centris* bees

(Hymenoptera: Anthophoridae). J Kansas Entomol Soc 49: 469-474.

Alcock J, Jones CE, Buchmann SL (1976b) Location before emergence of the female bee, *Centris*

pallida, by its male (Hymenoptera: Anthophoridae). J Zool Lond 179: 189-199.

Alcock J, Jones CE, Buchmann SL (1977a) Male mating strategies in the bee *Centris pallida* Fox

(Anthophoridae: Hymenoptera). The American Naturalist 111:145-155.

Alcock J, Eickwort GC, Eickwort KR (1977b) The reproductive behavior of *Anthidium maculosum*

(Hymenoptera: Megachilidae) and the evolutionary significance of multiple copulation by

females. Behav Ecol Sociobiol 2: 385e396.

Arizona-Sonora Desert Museum (2022, May 18) Phenology – Species. *Desert Museum Digital Library*.

Retrieved from: http://www.desertmuseumdigitallibrary.org/public/phenology.php

Barrett M (2022) Alternative reproductive tactics associated with variable sensory biology and

thermal physiology in male *Centris pallida* (Hymenoptera: Apidae) bees. PhD Dissertation.

Drexel University.

Barrett M, Johnson MG (2022). 2022 *Centris pallida* head widths. Dryad, Dataset.

https://doi.org/10.5061/dryad.bvq83bkc1

Barrett M, Tigreros N, Davidowitz G, O'Donnell S (2022) Adaptive variation in sex and male size

morph critical thermal maxima in *Centris Pallida* desert bees. SSRN [Preprint]. DOI:

http://dx.doi.org/10.2139/ssrn.4085466

Barrett M, O'Donnell S (2022) Individual reflectance of solar radiation as a thermoregulatory

adaptation to microclimate: A test in dimorphic male bees (*Centris pallida*). BioRxiv

[Preprint]. DOI: https://doi.org/10.1101/2022.06.28.497958

Barrett M, Schneider S, Sachdeva P, Gomez A, Buchmann S, O'Donnell S (2021)

Neuroanatomical differentiation associated with alternative reproductive tactics in

male arid land bees, *Centris pallida* and *Amegilla dawsoni*. J Comp Physiol A 207:497-504.

Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R (2011) Climate-

associated phenological advances in bee pollinators and bee-pollinated plants. PNAS 108:

20645-20649.

Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R (2013)

Historical changes in northeastern US bee pollinators related to shared ecological traits.

PNAS 110:4656–4660.

Beltrán I, Herculano-Houzel S, Sinervo B, Whiting MJ (2021) Are ectotherm brains vulnerable

to global warming? Trends Ecol Evol 36:691-699.

Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP, Potts

SG, Kleukers P, Thomas CD, Settele J, Kunin WE (2006) Parallel declines in pollinators and

insect-pollinated plants in Britain and the Netherlands. Science 313: 5785.

Bolnick D, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML (2003) The

ecology of individuals: Incidence and implications of individual specialization. Am Nat

161: 1-28.

Burdine JD, McCluney KE (2019) Differential sensitivity of bees to urbanization-driven changes in

body temperature and water content. Sci Reports 9: 1643.

Burkle LA, Marlin JC, Knight TM (2013) Plant-pollinator interactions over 120 years: Loss of

species, co-occurrence, and function. Science 339: 1611-1615.

Burraco P, Orizaola G, Monaghan P, Metcalfe NB (2020) Climate change and aging in

ectotherms. Glob Change Biol 26: 5371-5381.

Cane JH, Minckley RL, Kervin LJ, Roulston TH, Williams NM (2006) Complex responses

within a desert bee guild (Hymenoptera: Apiformes) to urban habitat fragmentation.

Ecological Applications 16: 632-644.

Cane JH, Neff JL (2011) Predicted fates of ground-nesting bees in soil heated by wildfire: Thermal

tolerances of life stages and a survey of nesting depths. Biological Conservation 144: 2631-

2636.

Chole H, Woodard SH, Bloch G (2019) Body size variation in bees: Regulation, mechanisms, and

relationship to social organization. Current Opinion in Insect Science, 35: 77-87.

Chou CC, Perez DM, Johns S, Gardner R, Kerr RA, Head ML, McCullough EL, Backwell PRY

(2019) Staying cool: The importance of shade availability for tropical ectotherms. Behav

Ecol Sociobiol 73:106.

Chown SL, Gaston KJ (2010) Body size variation in insects: a macroecological perspective. Biol Rev

Camb Philos Soc 85:139–169.

Chung DJ, Schulte PM (2020) Mitochondria and the thermal limits of ectotherms. J Exp Biol

223:jeb227801.

Danforth B (1991) The morphology and behavior of dimorphic males in *Perdita portalis*

(Hymenoptera: Andrendiae). Behav Ecol Sociobiol 29: 235-247.

Duffy GA, Coetzee BW, Janion-Scheepers C, Chown SL (2015) Microclimate-based

macrophysiology: Implications for insects in a warming world. Curr Opin Insect Sci 11:84-

89.

Everaars J, Settele J, Dormann CF (2018) Fragmentation of nest and foraging habitat affects time

budgets of solitary bees, their fitness and pollination services, depending on traits: Results

from an individual-based model. PLoS One 13: e0188269.

Gérard M, Martinet B, Maebe K, Marshall L, Smagghe G, Vereecken NJ, Vray S, Rasmont P,

Michez D (2019) Shift in size of bumblebee queens over the last century. Glob Change Biol

26: 1185-1195.

Grab H, Brokaw J, Anderson E, Gedlinske L, Gibbs J, Wilson J, Loeb G, Isaacs R, Poveda K (2019)

Habitat enhancements rescue bee body size from the negative effects of landscape

simplification. J App Ecol 56: 2144 – 2154.

GraphPad Software. 2021. GraphPad Prism v 9.3.1 for Windows. La Jolla, CA.

Hamblin AL, Youngsteadt E, López-Uribe MM, Frank SD (2017) Physiological thermal limits

predict differential responses of bees to urban heat-island effects. Biol Lett 13: 20170125.

Huey RB, Kingsolver JG (2019) Climate warming, resource availability, and the metabolic

meltdown of ectotherms. Am Nat 194:E140-E150.

Huey RB, Stevenson RD (1979) Integrating thermal physiology and ecology of ectotherms: A

discussion of approaches. Am Zool 19:357-366.

Jacobson MM, Tucker EM, Mathiasson ME, Rehan SM (2018) Decline of bumble bees in

northeastern North America, with special focus on Bombus terricola. Biol Conservation

217:437-445.

Kleijn D, Raemakers I (2008) A retrospective analysis of pollen host plant use by stable and

declining bumble bee species. Ecology 89:1811–1823.

Kuhlmann M, Guo D, Veldtman R, Doanldson J (2012) Consequences of warming up a hotspot:

Species range shifts within a centre of bee diversity. Diversity Distrib 18: 885-897.

Kukuk PF (1996) Male dimorphism in *Lasioglossum (Chilalictus) hemichalceum*: the role of larval

nutrition. J Kans Entomol Soc 69:147-157.

Lawson SP, Helmreich SL, Rehan SM (2017) Effects of nutritional deprivation on development and

behavior in the subsocial bee *Ceratina calcarata* (Hymenoptera: Xylocopinae). J Exp Biol 220:

4456-4462.

Lawson SP, Kennedy KB, Rehan SM (2020) Pollen composition significantly impacts the

development and survival of the native small carpenter bee, *Ceratina calcarata*. Ecol Entomol

46: 232-239.

Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of climate

change. Nature 462:1052-1055.

McCabe LM, Aslan CE, Cobb NS (2021) Decreased bee emergence along an elevation gradient:

Implications for climate change revealed by a transplant experiment. Ecology 103: e03598.

Muller K, Thiéry D, Moret Y, Moreau J (2015) Male larval nutrition affects adult reproductive

success in wild European grapevine moth (*Lobesia botrana*). Behav Ecol Sociobiol 69: 39-47.

Müller A, Diener S, Schnyder S, Stutz K, Sedivy C, Dorn S (2006) Quantitative pollen requirements

of solitary bees: implications for bee conservation and the evolution of bee-fower

relationships. Biol Conserv 130:604–615.

Nicholls E, Rossi M, Niven JE (2021) Larval nutrition impacts survival to adulthood, body size and

the allometric scaling of metabolic rate in adult honeybees. J Exp Biol 224: jeb242393.

Nooten SS, Rehan SR (2020) Historical changes in bumble bee body size and range shift of

declining species. Biodiversity and Conservation 29: 451-467.

Ockendon N, Baker DJ, Carr JA, White EC, Almond REA, Amano T… Pearce-Higgins JW (2014)

Mechanisms underpinning climatic impacts on natural populations: altered species

interactions are more important than direct effects. Glob Change Biol 20:2221–2229.

Oliveira MO, Freitas BM, Scheper J, Kleijn D (2016) Size and sex-dependent shrinkage of Dutch

bees during one-and-a-half centuries of land-use change. PLoS ONE 11:e0148983.

Oliveira RF, Taborsky M, Brockmann HJ (2008) Alternative reproductive tactics: an integrative

approach. Cambridge University Press, Cambridge.

O'Neill KM, O'Neill RP (2009) Prey, nest associates, and sex ratios of *Isodontia mexicana*

(Saussure) (Hymenoptera: Sphecidae) from two sites in New York. Entomologica

Americana 115: 90-94.

Parmesan C (2006) Ecological and evolutionary responses to recent climate change. Ann Rev

Ecol Evol Systematics 37:637-669.

Parton WJ, Logan JA (1981) A model for diurnal variation in soil and air temperature.

Agricultural Meteorology 23: 205-216.

Paxton RJ (2005) Male mating behaviour and mating systems of bees: an overview. Apidologie

36:145-156.

Rozen JG, Buchmann SL (1990) Nesting biology and immature stages of the bees *Centris caesalpiniae,*

C. pallida, and the cleptoparasite *Ericrosis lata* (Hymenoptera: Apoidea: Anthophoridae).

American Museum Novitates, 2985: 30 pp.

Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, et al. (2000) Global biodiversity

scenarios for the year 2100. Science 287: 1770-1774.

Sánchez-Bayo F, Wyckhuys KAG (2019) Worldwide decline of the entomofauna: A review of its

drivers. Biological Conservation 232: 8-27.

Scheper J et al (2014) Museum specimens reveal loss of pollen host plants as key factor driving wild

bee decline in the Netherlands. Proc Natl Acad Sci USA 111:17552–17557.

Shreeves G, Field J (2008) Parental care and sexual size dimorphism in wasps and bees. Behav Ecol

Sociobiol 62: 843-852.

Shuster SM (2010) Alternative mating strategies. In: C. Fox and D. F. Westneat (eds.) *Evolutionary*

behavioral ecology. Cambridge University Press, Cambridge.

Silva DP, Dew RM, Vilela B, Stevens MI, Schwarz MP (2018) No deaths in the desert: Predicted

responses of an arid-adapted bee and its two nesting trees suggest resilience in the face of

warming climates. Insect Conservation and Diversity 11: 449-463.

Sinervo B, Mendez-de-la-Cruz F, Miles DB, Heulin B, Bastiaans E, Villagran-Santa Cruz M, et al.

(2010) Erosion of lizard diversity by climate change and altered thermal niches. Science

328:894–899.

Snelling RR (1984) Studies on the taxonomy and distribution of American Centridine bees

(Hymenoptera: Anthophoridae). Contributions in Science 347:1-69.

Stemkovski M, Pearse WD, Griffin SR, Pardee GL, Gibbs J, Griswold T, Neff JL, Oram R, et al. (2020) Bee phenology is predicted by climatic variation and functional traits. Ecology Letters 23: 1589-1598.

Turley NE, Biddinger DJ, Joshi NK, López-Uribe MM (2022) Six years of wild bee monitoring shows changes in biodiversity within and across years and declines in abundance. Ecology and Evolution 12: e9190.

Vale CG, Brito JC (2015) Desert-adapted species are vulnerable to climate change: Insights from the warmest region on Earth. Global Ecol Conserv 4: 369-379.

van Dyck H, Matthysen E (1998) Thermoregulatory differences between phenotypes in the speckled wood butterfly: Hot perchers and cold patrollers? Oecologia 114:326-334.

Walters RJ, Hassall M (2006) The temperature-size rule in ectotherms: May a general explanation exist after all? Am Nat 167:510-523.

Warzecha D, Diekötter T, Wolters V, Jauker F (2016) Intraspecific body size increases with habitat fragmentation in wild bee populations. Landscape Ecology 31: 1449-1455.

Wieser W (1973) "Temperature Relations of Ectotherms: A Speculative Review." In: Wieser W. (eds) *Effects of Temperature on Ectothermic Organisms*. Springer, Berlin, Heidelberg.

Xie J, de Clercq P, Pan C, Li H, Zhang Y, Pang H (2015) Larval nutrition-induced plasticity affects reproduction and gene expression of the ladybeetle, *Cryptolaemus montrouzieri*. BMC Ecol Evol 15:276.

Zattara EE, Aizen MA (2021) Worldwide occurrence records suggest a global decline in bee species richness. One Earth 4: 114-123.

Supplemental Figures

Supplementary Figure 1. Distribution of female head widths in 2022. Distribution of head

widths of females (n = 114) collected while mating, foraging, and nesting from before 1130 on April

20 – May 8, 2022, at Blue Point Bridge, Arizona.

Supplementary Figure 2. Increase in patrolling or foraging male head widths from 0700 to

1130. Head width in the populations of males patrolling (gray squares, dashed line, n = 410) and

foraging (black dots, solid line, n = 203) increased over the course of the morning (Supplementary

Figure 2; linear regression, patrolling: [head width] = 0.095 [time of day] + 4.19, F = 23.65, df =

408, R^2 = 0.05, p < 0.0001; foraging: [head width] = 0.097 [time of day] + 4.08, F = 7.90, df = 201,

R^2 = 0.03, p < 0.0001).

Supplementary Figure 3. Changes in head width distributions of hovering and patrolling males from 1974 to 2022. Head width distributions for hovering (n = 57) and patrolling (n = 300) males at Blue Point Bridge are skewed towards smaller body sizes in 2022 as compared to 1974/1975 (n = 50 hoverers, 100 patrollers; data reproduced from Alcock et al. 1977a). All males collected before 930 am.

Supplementary Figure 4. Similar head width distributions of mating males from 1974 to 2022. Head width distributions for mating males at Blue Point Bridge demonstrates the population of males that obtain mating opportunities have relatively similar body sizes in 2022 (n = 56) as compared to 1974/1975 (n = 50; data reproduced from Alcock et al. 1977a). All males collected before 0930.

Supplementary Table 1. Head widths of *C. pallida* collected in April - May 2022 at Blue Point Bridge before 1130 am (n = 921 male, 114 female).

Type	Means ± SD (n)	Range
Foraging Males	4.85 ± 0.36 (203)	4.01 – 5.77
Patrolling Males	5.03 ± 0.34 (410)	4.03 – 5.84
Hovering Males	4.76 ± 0.20 (91)	4.29 – 5.16
Mating Males	5.50 ± 0.31 (71)	4.66 – 6.04
Digging Males	5.48 ± 0.25 (66)	4.74 – 6.00
Fighting Males	5.41 ± 0.25 (71)	4.74 – 5.83
Females	5.23 ± 0.28 (114)	4.33 – 5.95

577