Deciphering Hierarchical Chromatin Domains and Preference of Genomic Position Forming Boundaries in Single Mouse Embryonic Stem Cells

Yusen Ye[1*], Shihua Zhang[2,3,4], Lin Gao[1], Yuqing Zhu[5], Jin Zhang[6,7,8,9]

1School of Computer Science and Technology, Xidian University, Xi'an 710071, Shaanxi, China.

2NCMIS, CEMS, RCSDS, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China.

3School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China.

4Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences, Kunming 650223, China.

5Center for Stem Cell and Translational Medicine, School of Life Sciences, Anhui University, Hefei, 230601, Anhui, China.

6Center for Stem Cell and Regenerative Medicine, Department of Basic Medical Sciences, and Bone Marrow Transplantation Center of the First Affiliated Hospital, Hangzhou, Zhejiang, China.

7Zhejiang Laboratory for Systems and Precision Medicine, Zhejiang University Medical Center, Hangzhou, Zhejiang, China.

8Institute of Hematology, Zhejiang University, Hangzhou, Zhejiang, China.

9Center of Gene/Cell Engineering and Genome Medicine, Hangzhou, Zhejiang, China.

*To whom correspondence should be addressed. Tel: +86-15802963380; Email: ysye@xidian.edu.cn

Abstract

The exploration of single-cell 3D genome maps reveals that chromatin domains are indeed physical structures presenting in single cells and domain boundaries vary from cell to cell. However, exhaustive analysis of regulatory factor binding or elements for preference of the formation of chromatin domains in single cells has not yet emerged. To this end, we first develop a **hi**erarchical **c**hromatin domain **s**tructure identification algorithm (named as **HiCS**) from individual single-cell Hi-C maps, with superior performance in both accuracy and efficiency. The results suggest that in addition to the known CTCF-cohesin complex, Polycomb, TrxG, pluripotent protein families and other multiple factors also contribute to shaping chromatin domain boundaries in single embryonic stem cells. Different cooperation patterns of regulatory factors decipher the preference of genomic position categories forming boundaries. And the most extensive six types of retrotransposons differentially distributed in these genomic position categories with preferential localization.

Introduction

Genome across a wide range of eukaryotic organisms is efficiently packaged and organized into hierarchical chromatin architecture via ubiquitous architectural features, which is critical to gene regulation and dynamical changes in development and disease [1-3]. These basic features consist of chromatin fibers, which fold into chromatin loops, such as enhancer-promoter interactions and architectural loops mediated by CCCTC-binding factor (CTCF) [4]. These fibers further fold into chromatin domains, referred to as topologically associating domains (TADs) or sub-TADs, which are associated with each other to generate chromosomal compartments. Each chromosome occupies a distinct volume or chromosome territory within the nucleus [4-6]. Genome architecture is an integral part of the chromatin landscape that transcription factors (TFs) must navigate to exert their regulatory roles [7]. Although most loci and chromosomes are characterized by a high degree of order and non-randomness, the precise functional roles and formation mechanism of these features remains obscure [8].

Some TFs, cofactors, and histone modifications that correlate with the chromatin structures have been identified to study features of chromatin organization [9-11]. Particularly, TAD boundaries are enriched with multiple factors including CTCF, cohesin, H3K4me3, H3K36me3, transcription start sites, housekeeping genes, etc, suggesting that CTCF binding, high levels of transcription activity, multiple histone modifications, and other regulatory factors may contribute to the formation of chromatin domains in mammals [8]. Although TADs and their associated regulatory factors have been widely identified in multiple species and are highly conserved and stable across different cell types [6, 12], single-cell 3D genome analysis indicated that they display substantial cell-to-cell variation [13, 14]. Therefore, the bulk analysis only reflects properties of ensemble average structures from millions of cells, which may mask chromatin features appearing in a few cells or a single cell and limit our understanding of chromatin structures.

A recent study reveals that domain structures often adopt globular conformation with strongly physical segregation of neighboring domains, and domain boundaries are preferentially located at CTCF- and cohesin-binding sites with a super-resolution chromatin tracing method [13]. More surprisingly, single-cell domain structures persist even after cohesin degradation [13]. These results suggest other TFs or epigenetic factors may contribute to the formation of chromatin domains. Currently, mouse embryonic stem cells (mESCs) have been served as a specific model cell system to elucidate the mechanisms of 3D genome organization and explore the relationship between chromatin domains and gene regulation. Hundreds of TFs and epigenetic modification profiles have been identified in mESCs [9-11]. The above observations push us to investigate the formation of chromatin domains and their relationship with functional elements in single mESCs systematically.

Here, we first develop a **hi**erarchical **c**hromatin domain **s**tructure identification algorithm (named as **HiCS**) from single-cell Hi-C maps, which shows superior performance in both accuracy and efficiency. We reorganize atlas of ChIP-seq for

85 mESCs, and reveal the patterns of hundreds of regulatory factors are significantly either
86 enriched or absented in domain boundaries of single cells, suggesting that, in addition
87 to known CTCF-cohesin complex, Polycomb, TrxG, pluripotent protein families,
88 different types of histone modifications, and other multiple factors could promote the
89 formation of chromatin domains in single mESCs. To further elaborate on cooperation
90 patterns between different types of regulatory factors, and genomic position categories
91 with differential preference forming boundaries drive by these cooperation patterns, we
92 cluster 13 large genomic position categories (consisting of 29 sub-categories) annotated
93 by 7 different regulatory factor clusters (consisting of 27 sub-clusters). The clear
94 patterns provide a detailed view of the preference of these genomic position categories
95 forming domain boundaries. Furthermore, we discover that these genomic position
96 categories are enriched by different cooperation of retrotransposons with preferential
97 localization. Last but not the least, we find that genomic positions enriched by
98 Alu/B2/B4 retrotransposons have higher preference scores for forming boundaries in
99 G1 and ES phases in comparison with MS and LS/G2 phases, whereas genomic
100 positions enriched by L1/ERVK retrotransposons display opposite tendency. In
101 summary, we reveal that multiple types of regulatory factors interplaying with each
102 other in specific genomic positions could affect focal chromatin interactions, thereby
103 changing interaction density or insulation strength of these regions. This further
104 navigates the preference of genomic position forming boundaries, shape hierarchical
105 chromatin domains, and thus regulate gene expression and cell functions, even cell
106 identity in single embryonic stem cells.
107
108 **Results**
109 **Overview of HiCS.** The key design of HiCS is to convert the problem of the
110 identification of hierarchical chromatin domains into finding peaks of insulation
111 strength at different genome scales. The domain boundaries usually have higher
112 insulation strength than their neighbors and a relatively large distance from any regions
113 with higher strength (**Fig. 1a**). HiCS calculates two metrics for each bin including the
114 insulation strength ρ and the minimum distance between the bin and any other bin with
115 higher strength δ, and controls the number of peaks to obtain the hierarchical chromatin
116 domains at different scales by α (**Fig. 1b and c**). HiCS is super-fast to identify a
117 chromatin hierarchy that the domain of a higher level embraces the multiple smaller
118 ones of a lower level (**Fig. 2b**). Note that high-level boundaries with high δ in local
119 regions may have lower insulation strength than low-level ones, and the boundaries in
120 the same level may have different local insulation strengths, HiCS can automatically
121 detect different level domain boundaries based on local background (**Fig. 2b**).
122   Specifically, HiCS consists of three steps (**Methods**): (1) Preprocesses single-cell
123 Hi-C data and calculates the two metrics to find the peaks at a given scale α, and
124 determines the domain boundaries for each chromosome of an individual cell (**Fig. 1a-**
125 **c**); (2) Determines the hierarchical domains of a chromatin by adjusting the optimal
126 structural identification parameter ($\alpha = 1$) (**Fig. 1c**, **Fig. 2a, b**, and **Supplementary**

127 **Fig. S1a**), Noting that the optimal parameter is automatically determined by the
128 algorithm; (3) Applies a bi-clustering method to group chromatin positions and
129 regulatory factors respectively and analyzes the genomic structure-function
130 relationship by combining hundreds of TFs and epigenetic factors with chromatin
131 domains (**Fig. 4**).
132
HiCS shows superior performance. We benchmark the performance of HiCS for
domain detection against two methods: one is the commonly used method for bulk data,
insulation score (IS) [15], and another is a recent single-cell TAD detection method,
deTOKI [16]. We apply these methods to the preprocessed single-cell Hi-C data
generated from mESCs [17]. To compare the performance of them fairly, we adjust the
scaling parameter of HiCS to obtain similar number and size of domains with IS and
deTOKI, respectively (**Fig. 1e**). Actually, the insulation strengths of domains obtained
from HiCS is significantly higher than those of IS and deTOKI respectively with similar
number of domains, suggesting its superiority to competing methods (**Fig. 1f**). The
running time of HiCS is significantly less than both algorithms under the same
hardware condition (**Fig. 1g**). Moreover, the domain boundaries detected by HiCS are
more significantly enriched in multiple common factors, including CTCF, H3K4me3,
Housekeeping (HK) genes (**Fig. 1h**), as well as RNA polymerase II (PolII), promoters,
highly expressed genes, and average phastcon score (**Supplementary Fig. S1b**). Also,
the boundaries detected by HiCS are more pronounced disappearance of the enhancers
or the super enhancers (SEs), which unfavorably form boundaries in single cells as
reported on analysis of bulk-cells [9] (**Supplementary Fig. S1b**). With an example, we
can see that HiCS can obtain more accurate chromatin domain boundaries at single-cell
resolution (**Fig. 1a and d**). Taken together, HiCS shows superior performance in both
accuracy and efficiency.
153
The existence of hierarchical chromatin domains. We adjust the optimal structural
identification parameter to generate multiple-scale chromatin domains at different
genomic scales for 1315 single mESCs at 40kb resolution. We clearly observe four
peaks of domain size and insulation strength distribution with different scaling
parameters of 0.2, 1, 4, and 8, which we choose for the downstream analysis (**Fig. 2a,
b** and **Supplementary Fig. S1a**). The domain scales of these four levels are
approximately 200Kb~600Kb, 800Kb~1Mb, 2Mb~3Mb, and ~5Mb respectively. We
show an example for hierarchical chromatin domains, which well match the local
insulation strength of chromatin regions (**Fig. 2b**).
The median size of chromatin domains increases and the median boundaries'
insulation strength enhances with the increase of genomic position level
(**Supplementary Fig. S2a-c**). The boundaries show obvious cell-to-cell heterogeneity
with a nonzero probability of being located at any genomic positions. 99.8% of genomic
positions form boundaries in at least 1% of cells, while only 4.1% of genomic positions
form boundaries in more than 14% of cells (**Fig. 3a**). We also observe that the

probability forming boundaries enhances with the levels of genomic position increasing
(**Supplementary Fig. S2d**). The above observations suggest that domain boundaries
vary from cell to cell with nonzero probability at all genomic positions as reported in
[13], and the preference of genomic position forming boundaries may shape the
formation of hierarchical chromatin domains in single cells.
174
Regulatory factors navigate the preference of genomic position forming
boundaries. Recent study has shown that domain boundaries are preferentially located
at CTCF- and cohesin-binding sites with a super-resolution chromatin tracing method
[13]. CTCF and cohesin have been proven to be key factors controlling the functional
architecture of mammalian chromosomes forming TADs or sub-TADs by "loop
extrusion" [18]. Indeed, we observe that both CTCF and cohesin show similar
enrichment patterns in boundaries of single cells in different genomic levels, and as the
level of boundaries increases, the degree of enrichment gradually increases (**Fig. 3d**
and e).
Previous study has illustrated that high levels of transcription activity may
contribute to TAD formation in bulk analysis [19]. Here, we observe that the
accessibility of genomic positions, the enrichment degree of HK genes and
differentially expressed genes (DEGs), and the expression of genes gradually
increases along with the increasing levels of genomic positions (**Fig. 3b, c and**
Supplementary Fig. S2e). Conversely, enhancers and super enhancers (SEs) greatly
become more absented with the increase of levels (**Fig. 3b and c**). It suggests that the
emergence of highly transcriptional activity, especially DEGs, and the absence of
enhancers, especially SEs, improve the probability of genomic positions forming
boundaries of single cells (**Fig. 3b and c**).
Although the CTCF-cohesin complex is critical for the formation of TADs in
mammalian cells, a substantial number of boundaries remain unaffected after cohesin
degradation in single cells, suggesting other modulators exist on domain boundaries [7,
20]. We indeed observe that mediators (*Med1*, *Med12*, *Med26*, *Cdk8*, *Cdk9*), *Nipbl*,
PolII, TFIIB are all enriched in the boundaries of single cells, and as the level of
boundaries increases, the degree of enrichment gradually increases (**Supplementary**
Fig. S2g-f). Mediators are essential coactivators that are recruited to the regulatory
regions of active genes and facilitate the ability of enhancer-bound TFs to recruit PolII
to the promoters of target genes, and *Nipbl* has been proven to bind mediator to load
cohesin [9, 10, 21, 22]. The above results suggest that these factors may play important
roles in shaping preference of genomic position forming boundaries of single cells.
The master factors of Polycomb repressive complex 1 (PRC1) and PRC2 just
exhibit two different enrichment patterns at the domain boundaries of single cells. One
type (*Aebp2*, *Rybp*, *Ring1b*) forms obvious single peaks at the domain boundaries, and
the degree of enrichment gradually increases with the boundary level increasing
(**Supplementary Fig. S3a**). Another type (*Ezh2*, *Pcl2*, *Suz12*, *Eed*) shows double peaks
around the domain boundaries (**Supplementary Fig. S3b**). These two types of

complexes have been proved to have distinct catalytic activities, but both are generally associated with transcriptional silencing [23]. We also observe that TrxG associated proteins (COMPASS: *Set1a*, *Mll2*, *Mll3/4,* and SWI/SNP: *Brg1*) are all enriched in domain boundaries, except for *Mll3/4* (**Supplementary Fig. S3c**). In mammalian, *Set1a* reportedly contributes to most of the H3K4me3, and *Mll2* mediates H3K4me2 and H3K4me3 at developmental genes, while *Mll3/4* implements monomethylation of H3K4 at enhancers [24]. *Brg1* is an ATP-dependent chromatin remodeler, contributing to the maintenance of pluripotency and self-renewal in ESCs [9]. The above results suggest that the PRC and TrxG protein families could change focal chromatin interactions in different ways.

We find that the core TFs (*Oct4*, *Sox2*, and *Nanog*) controlling the pluripotent state do not prefer to appear in domain boundaries, which may be related to chromatin hubs occupied by super-enhancers/enhancers (**Supplementary Fig. S4a**) as reported in [25]. In addition, we also collect 14 additional TFs that may contribute to the pluripotent state of mESCs and investigate whether they are enriched in domain boundaries of single cells [9] (**Supplementary Fig. S4b and c**). The results indicate that 9 additional TFs (*Esrrb*, *Nr5a2*, *Klf4*, *Zfp281*, *Tcf3*, *Tcfcp2l1, Stat3*, *Prdm14*, and *Smad2/3*) that were previously shown to occupy both typical enhancers and SEs do not prefer to appear in domain boundaries, while 5 factors (*c-Myc*, *n-Myc*, *Zfx*, *Tbx3*, and *Yy*1) that were previously shown to occupy promoter-proximal sites are enriched in domain boundaries [9]. Among them, it is particularly interesting that *Smad2/3*, *Stat3*, and *Tcf3* signaling pathways were considered as key modulators controlling mESCs pluripotent state transition by modifying chromatin states and shaping chromatin domains [26-28].

In addition, we also observe additional 45 proteins are either enriched or absented in domain boundaries of single cells with different enrichment patterns (**Supplementary Figs. S5-7**), which suggests their potential in shaping chromatin domains.

The above TFs and chromatin regulators have the most profound impact on cell states through collaborative control of chromatin states and spatial structures. 14 different histone-modifying enzymes show a variety of enrichment patterns around domain boundaries of single cells (**Supplementary Fig. S8**). For example, ZC3H11A shares consistent enrichment patterns with CTCF-cohesin (**Fig. 3e and Supplementary Fig. S8**). H3K27me3 shares similar enrichment patterns with PRC2, while H2AK119ub1 shares similar enrichment patterns with PCR1 (**Supplementary Figs. S3a and b**), which are consistent with their function in participating chromatin modifications [23]. H3K4me1 marking of enhancers is not enriched in domain boundaries. And H3K4me2 shares similar enrichment patterns with H3K4me3 around domain boundaries, which all are associated with TrxG protein family [24] (**Supplementary Fig. S3c**). It suggests different histone modifications may cooperate with different TFs and chromatin regulators, to modify chromatin states, shape the local chromatin interaction status and organize chromatin domains of single cells.

To sum up, we have observed hundreds of TFs, chromatin regulators, and histone
modifications are significantly either enriched or absented in domain boundaries of
single cells with differential enrichment patterns. The occupancy of these regulatory
factors in specific genomic positions will affect focal chromatin interactions, thereby
changing the interaction density or insulation strengths of these regions. These
processes may navigate the preference of genomic position forming boundaries, and
then shape hierarchical chromosome domains of single cells.
259
Cooperation among regulatory factors differentiate genomic position categories.
To further elaborate on cooperation patterns between different types of regulatory
factors, and genomic position categories with differential preference forming
boundaries drive by these cooperation patterns, we grouped 13 large spatially organized
genomic position categories (consisting of 29 sub-categories), which were annotated
by 7 different regulatory factor clusters (consisting of 27 sub-clusters). The result helps
to explain the preference of genomic position forming boundaries in single cells, and
providing an increasingly complex view of the genomic structure-function relationship
(**Fig. 4 and Supplementary Fig. S9**).
The categories of regulatory factors. The result of hierarchical clustering indicates
that these categories mainly consist of the cluster of core regulatory factors of mESC
(CoreRF, particularly *Oct4*, *Sox2*, and *Nanog)*, the clusters associated with highly
active activators for SEs and enhancers (HASE&E), the clusters of highly transcribed
activators (HTA), the clusters of transcriptional repressor factors (TRF), the cluster
associated with heterochromatin factors (Hetero), and the clusters of architectural
proteins (AP). There are also a few single factors (Other) (**Fig. 4a and Supplementary
Fig. S9**). For a specific example with less prior studies among these clusters, *Sumo2* is
required to play critical roles in the canonical *Zfp809/Trim28/Eset* complex via post-
translational sumoylation of *Trim28*, which enhances the recruitment of *Trim28* to the
proviral DNA, resulting in the modification of proviral chromatin with repressive
histone H3K9me3 marks in turn [29]. These four factors were grouped with H3K9me3
together in the analysis below, which may organize the formation of heterochromatin
(**Fig. 4a**). The annotation information and supporting materials of these all categories
were summarized in **Supplementary Table S2**.
We also observe some interesting cooperative patterns among different clusters
(**Supplementary Fig. S9b**). For example, the CoreRF cluster is strongly associated
with the HASE&E cluster, but not the HTG cluster, while the HASE&E cluster is
intimated to the HTG cluster. It suggests the HASE&E cluster may be a bridge between
CoreRF and HTG clusters, which can be linked to various signaling pathways involved
in the transcriptional network in mESCs.
The division of genomic position categories. We further analyze the preference of
genomic position categories with a ratio>1% (**Supplementary Fig. S10a**). These
genomic position categories can be divided into three families, including the categories
with high accessibility preferentially forming boundaries, the categories with high

accessibility unfavorably forming boundaries, and the categories with low accessibility
unfavorably forming boundaries (**Fig. 5a** and **Supplementary Fig. S10b**).

Of note, the largest type with weak open chromatin accessibility (13.7% of all
positions, consisting of 12.6%, 1.1% from the 37^{th}, 34^{th} classes, respectively, defined
as AP1-Positions) preferentially forming boundaries only are significantly occupied by
AP1 factors (CTCF, cohesin, etc.) comparing with other types of factors (**Fig. 4b, c** and
Fig. 5a). Previous studies have indicated that loop extrusion executed by CTCF and
cohesin is a leading factor governing domain formation and facilitating chromatin
folding. We find that the other subunits (*Sa1* and *Sa2*) of cohesin which occupy the
same genomic locus and present similar enrichment patterns at boundaries of single
cells, suggesting that both *Sa1* and *Sa2* may also participate in the maintenance of
domain boundaries (**Fig. 3e**). ZC3H11A, a zinc finger protein, shows a uniform pattern
as *Sa1* and *Sa2,* implying its novel role in shaping boundaries of chromatin domains
(**Supplementary Fig. S8**). The chromatin positions that AP1 proteins occupy may be
directly related to the macroscopic architecture of chromatin within the nucleus, and
indirectly change the local chromatin context to exert regulatory functions.

The second major category of open chromatin positions preferentially forming
boundaries (7.5%, 2.7% from the 38^{th}, 39^{th} classes, defined as HTG-Positions) is
associated with highly transcribed genes (**Fig. 4b, c** and **Fig. 5a**). These regions are
mainly occupied by the HTA (particularly the TrxG class, playing important roles in
orchestrating the stable activation of gene expression) and HASE&E clusters, but not
the CoreRF cluster, which indicates that the CoreRF cluster hardly participates in the
regulation of genes occupying in domain boundaries of single cells.

The third major category of open chromatin positions preferentially forming
boundaries (3.1%, 1.7% from the 41^{th}, 33^{th} classes, defined as TrxG-PRC-Positions,
1.5% from the 35^{th} class, defined as TrxG-AP1-Positions) is associated with PRC, TrxG,
and AP1 proteins (**Fig. 4b, c** and **Fig. 5a**). The 33^{th} class of genomic regions is also
associated with AP1 proteins with a higher probability of forming domain boundaries
of single cells compared with the 41^{th} class, which suggests CTCF-cohesin complexes
may help these regions form a more stable structure. In addition, the 35^{th} class of
genomic region with higher enrichment of TrxG proteins and lower enrichment of PRC
proteins compared with the 33^{th} class, which results in higher open chromatin
accessibility, enhancers occupation, and probability forming domain boundaries in
single cells. The above results indicate that chromatin modifiers (TrxG and PRC
proteins) can provide an additional layer of regulation by changing chromatin structures,
and balance the formation of domain boundaries by repressing and activating chromatin
states, respectively.

The major categories of the open chromatin positions unfavorably forming
boundaries are associated with highly active SEs or enhancers, which mainly contain
three categories (**Fig. 4b, c** and **Fig. 5a**). The first category (4.8% from the 0^{th} class,
defined as SEs-Positions) almost encompasses more than 75% of SEs, which are
strongly related with the HASE&E, CoreRF, TrxG, and Med-PolII clusters. The second

category (5.5%, 1% from the 21^{th}, 9^{th} classes, defined as CoreRF-Positions) is mainly
associated with enhancers occupied by CoreRF. The slight difference between the 21^{th}
and 9^{th} classes is that the 9^{th} one shows higher chromatin accessibility, while the 21^{th}
one shows a stronger DamID enrichment, which may be related to their locations in the
nucleus. The third category (2.7%, 1.3% from the 7^{th}, 5^{th} classes, H3K4me1-Positions)
are related to enhancers enriched by H3K4me1. These regions taking SEs or enhancers
associated with CoreRF and H3K4me1 as focal regions generate dense chromatin
structures mediated by different regulatory factors in an orientation-independent
manner, and unfavorably form boundaries in single cells with weakened insulation
strengths [30].
We also check the classes of genomic positions with low chromatin accessibility,
all of which unfavorably form boundaries in single cells. These chromatin positions are
mainly divided into the following categories (**Fig. 4b, c** and **Fig. 5a**). The first category
(19% of all positions, consisting of 9.2%, 3.7%, 3.5%, 1.6%, and 1% from the 29^{th}, 28^{th},
20^{th}, 31^{th}, and 27^{th} classes, respectively, defined as Hetero-Positions) are associated
with the Hetero cluster. The second category (18% of all positions, consisting of 2.8%,
2.3%, 1.9%, and 1.7% from the 16^{th}, 15^{th}, 12^{th}, and 13^{th} classes, respectively, defined
as DamID-Positions) maintains high enrichment of the DamID signal. The above
observations suggest that the presence of heterochromatin factors reduce the probability
of single-cell domain boundaries formation. Finally, we observe an interesting and
specific category (4.6%, 1.2% from the 45^{th}, 44^{th} classes, defined as AP2-Repressor-
Positions), which are associated with the AP2 (*Yy1* and H2AZ) and H3K36me3 classes
(**Fig. 4a** and **Fig. 5a**). Both *Yy1* and H2AZ facilitate the organization of genome
architecture (**Supplementary Table S2**) [31-34].
To summarize, we obtain the following key results by the above analysis: (1) The
genomic positions occupied by architectural proteins (CTCF and cohesin), highly
transcribed genes, and TrxG proteins preferentially form boundaries in the single cells
analysis. (2) The genomic positions taking SEs or enhancers associated with CoreRF
or marked by H3K4me1, heterochromatin factors, and repressing factors unfavorably
form boundaries in single cells with weakened insulation strengths.
366
Retrotransposons are associated with these genomic position categories. In a recent
paper, Shen and colleagues find that retrotransposons embedded in 3D genome
architecture, regulates the formation of euchromatin and heterochromatin respectively,
particularly the separation of compartments A/B [35, 36], which are consistent with our
observations. Notably, our study focuses on the effect of retrotransposons on the
preference of genomic position forming boundaries in single cells. In order to explain
the preference of genomic position categories forming boundaries in single cells with
more detail, we further analyze the influences of the largest six types of
retrotransposons on genomic architecture.
We observe that the division of genomic position categories is strongly associated
with the regulation of retrotransposons. The genomic positions are clearly divided into

five functional units based on categories of retrotransposons, including Highly-Alu/B2-
Positions, Other-Alu/B2/B4-Positions, MaLR-dominant-Positions, L1-Positions, and
ERVK/L1-Positions (**Fig. 4d** and **Fig. 5a**).

Firstly, all genomic regions (Hetero-Positions, DamID-Positions, AP2-Repressor-
Positions) with low accessibility and unfavorably forming boundaries are enriched by
L1 elements, which tend to occupy gene-poor, heterochromatic B compartments that
interact with lamina-associated domains in previous studies [35]. Among the defined
regions, we are surprised to find that Hetero-Positions are specially associated with
ERVK elements, which may indicate that ERVK acts as specific roles in regulating
embryonic development as reported in [37, 38]. The above observation also implies
that the previously unannotated 18^{th} and 43^{th} genomic positions may be related to
heterochromatin organization.

We also observe that genomic regions (HTG-Positions, SEs-Positions, TrxG-AP1-
Positions, TrxG-PRC-Positions, H3K4me1-Positions, the 9^{th} class of CoreRF-Positions)
are enriched by Alu/B2/B4, which may be related to euchromatin organization [35].
HTG-Positions, SEs-Positions as well as TrxG-AP1-Positions with higher Alu/B2
enrichment than others suggest that the enrichment of Alu/B2 may indicate the
transcription level of genes, and promote the formation of hierarchical chromatin
structures by regulating gene transcription and SEs/Enhancer activation.

Besides the above genomic positions, what is interesting is that MaLR elements
are enriched in AP1-Positions, the 21^{th} class of CoreRF-Positions, and other
unannotated regions (the 6^{th}, 3^{th}, 8^{th}, and 22^{th} classes). We first observe that the 21^{th}
class occurs a stronger heterochromatin factors (the Hetero cluster) enrichment
compared to the 9^{th} (CoreRF-Positions), which may result in uncertain chromatin states
in the 21^{th} positions (**Fig. 4b and c**). In addition, previous studies have shown that
domain boundaries mediated by AP1 proteins (e.g., CTCF and cohesin) may block the
spread of chromatin states [4]. It may suggest that the regions dominantly enriched by
MaLR elements may often undergo switches between euchromatin and
heterochromatin.

In summary we find that: (1) L1-Positions with low accessibility and unfavorably
forming boundaries are associated with heterochromatin organization, and Alu/B2/B4-
Positions are associated with euchromatin chromatin, which are consistent with a recent
study [35], while MaLR-Positions may result in switches between euchromatin and
heterochromatin, which is yet to be proven. (2) ERVK elements acts as specific roles
heterochromatin formation, while Alu/B2 may promote highly transcription of genes
and highly activation of SEs/Enhancers. These retrotransposons contribute to the
maintaining of chromatin states, and interplay with other types of regulatory factors, to
navigate the preference of genomic positions forming boundaries and gene regulation
in single cells.

Genomic landscape regulates cellular states. To investigate the preference of
genomic positions in the above functional groups along the cell cycle process, we check

the dynamics of each functional group forming boundaries in single cells among four different cycle phases, including G1, early-S (ES), mid-S (MS), and late-S/G2 (LS/G2). We first reveal that genomic positions enriched by SEs show a significant preference for forming boundaries in the G1 phase (**Fig. 5b**). The observation suggests that highly activation of SEs in the phase may promote gene regulation and transcription for cell growth in size, and ensure biomaterials for DNA synthesis. In addition, the functional group (HTG-Positions) accompanied with highly transcribed genes exhibits a significant loss of boundaries in the MS phase, which may be because the rates of transcription and protein synthesis are low during DNA replication (**Fig. 5b**). We also observe that functional groups occupied by both CTCF-cohesin and TrxG-PRC complexes prefer to form boundaries in ES phases (**Fig. 5b**). The observation implicates in the clearest segmentation of chromatin structures at the beginning of DNA replication [39, 40]. Both complexes have been proven to modify local chromatin structure and regulate higher-order chromatin organization [7, 41]. And functional groups (Hetero-Positions and AP2-Repressor-Positions) associated with heterochromatin organization prefers to form boundaries in both MS and LS/G2 phases (**Fig. 5b**). Both phases may prepare for everything entering the mitosis phase with condensing chromatin states.

In general, genomic positions (HTG-Positions, SEs-Positions, TrxG-AP1-Positions, TrxG-PRC-Positions, H3K4me1-Positions, and the 9th class of CoreRF-Positions) enriched by Alu/B2/B4 retrotransposons have higher preference scores for forming boundaries of single cells in G1 and ES phases in comparison with MS and LS/G2 phases, whereas genomic positions (Hetero-Positions, AP2-Repressor-Positions, and DamID-Positions) enriched by L1/ERVK retrotransposons display opposite tendency, following by high preference scores for forming boundaries in both MS and LS/G2 phases (**Fig. 5b** and **Supplementary Fig. S11**). The above observations further expound that the dynamic interplay among different types of regulatory factors, retrotransposons, and chromatin structures could navigate gene regulation and cell functions, even cell identity in single embryonic stem cells.

Discussion
Several decades of research have shown that eukaryotic chromatin adopts a complex hierarchical architecture within the nucleus, which plays a key role in functional implications for almost all nuclear processes. Thus, the spatially organized chromatin architecture interplaying with multiple types of regulatory factors shape focal chromatin landscapes and then exert gene regulatory functions. Single-cell 3D genome analysis extends the limitation of bulk analysis to show substantial cell-to-cell variation and promote our understanding of chromatin structures in the individual cell. Recent discoveries on single-cell 3D genome have shed light on the relationship between CTCF-cohesin complexes and domain formation, but the more molecular details associated with regulatory factors remain to be investigated [13].

461   Here, we develop HiCS to detect hierarchical chromatin domains from single-cell
462 Hi-C maps, and observe hundreds of regulatory factors, including TFs, chromatin
463 regulators, and histone modifications, are significantly either enriched or absented in
464 domain boundaries of single cells, which presents several different enrichment patterns.
465 The results suggest their potential cooperative associations in shaping focal chromatin
466 interactions, thereby changing interaction density or insulation strength of these regions,
467 and drive different genomic position categories. We further group chromatin position
468 categories and different regulatory factor clusters, explaining the emergence and
469 functionality of different chromatin landscapes and providing a comprehensive view of
470 the genomic structure-function relationship. We also find that different
471 retrotransposons exactly match the above genomic position categories. The above
472 results indicate that these regulatory factors interplaying with each other exert gene
473 regulatory processes and control cell functions, even cell identity.

474   The chromatin structures within the nucleus operate in an obvious dynamic process
475 driven by both "loop extrusion" and attractive process induced by regulatory factors
476 (associated with compartmentalization). The process may condense or loose local
477 chromatin landscapes in an overlapping and concerted manner accompanied by
478 adjusting insulation strength of chromatin position and generating chromatin loops, and
479 then shape gene expression programs during cell-fate specification [39]. Further work
480 is needed to leverage more specific chromatin structures, particularly chromatin loops,
481 of single cells with more abundant regulatory factors (TFs, chromatin regulators,
482 histone modifications, retrotransposons, RNA, even structural variations) to understand
483 structure-function relationships in complex tissues or diseases, particularly cancers [42,
484 43]. It will promote our understanding of how multiple types of regulatory factors
485 interact with chromatin topological engines (such as loop extraction and
486 compartmentalization) to regulate the gene-repression program, determine cell
487 functions and identity, and further explain tissue complexity and disease development.
488

489 **Materials and Methods**
490 **Single-cell Hi-C and other genomic data processing**
491 ***Single-cell Hi-C data generated from mESCs.*** The single-cell Hi-C dataset used in this
492 study consists of 1992 diploid cells of mESCs grown in 2i media without feeder cells
493 with stringent quality control filter. This dataset involves a median number of 393506
494 restriction fragments, and 127233 distinct >1 kb contacting pairs on average per cell
495 [17]. The top 1315 cells with >250000 contacts per cell were selected for downstream
496 analysis. Among them, 317, 341, 303, and 354 cells belong to G1, early-S (ES), mid-S
497 (MS), and late-S/G2 (LS/G2) phases labeled by fluorescence-activated cell sorting
498 (FACS) sort criterion, respectively.
499

500 ***An atlas of ChIP-seq for mESCs.*** We organized an atlas of ChIP-seq for hundreds of
501 regulatory factors of mESCs (**Supplementary Table S1**), including CTCF, cohesin
502 (*Smc1, Smc3, Rad21, Sa1, Sa2*), mediators (*Med1, Med2, Med26, Cdk8, Cdk9*),

codensin (*Capd3, Nipbl*), PolII, TFIIB, polycomb repressive complex (*Aebp2, Rybp,*
Ring1b, Rnf2, Suz12, Ezh2, Eed, Pcl2), Trithorax protein family (*Set1a, Mll2, Mll3/4,*
Brg1), the core regulatory factors of mESC (*Oct4, Sox2, Nanog*), the regulatory factors
of ESC occupied on enhancers or SEs (*Esrrb, Nr5a2, Klf4, Stat3, Prdm14, Zfp281, Tcf3,*
Tcfcp2l1, Smad2/3), the regulatory factors of ESC occupied on promoter-proximal sites
or sites that border topological domains (*c-Myc, n-Myc, Zfx, Tbx3, Yy1*), and additional
45 TFs. We collected 14 histone modification factors, including H3K4me1, H3K4me2,
H3K4me3, H3K27me3, H3K36me3, H3K79me2, H3k9me3, H3K9ac, H3K122ac,
H3K64ac, H3K27ac, H2AZ, ZC3H11A, and H2AK119ub1. In addition, we also
collected two chromatin accessibility datasets (ATAC-seq and Dnase-seq) and the
DamID-seq dataset.
For the ChIP-seq of regulatory factors, peaks were called using MACS2 software
with q-value cut-off 1×10^{-5} [44]. The source information and supporting materials
of these factors were summarized in **Supplementary Table S1 and S2**.
517
Regulatory elements and genes. Enhancers/SEs and gene expression datasets of
mESCs were downloaded in GSE29278 [45]. Housekeeping genes were downloaded
in Housekeeping and Reference Transcript Atlas (HRT Atlas v1.0,
www.housekeeping.unicamp.br) [46]. PhastCons scores were downloaded from the
UCSC Genome Browser via
ftp://hgdownload.cse.ucsc.edu/goldenPath/mm9/phastCons30way/vertebrate [47].
Mouse cell-cycle annotated genes were obtained from the mouse genome informatics
(MGI) (http://www.informatics.jax.org/), containing 891 genes relating to cell cycle
process and regulation. We adopted Seurat to detect DEGs and the variable score of
genes using *FindAllMarkers* and *FindVariableFeatures* functions based on single-cell
RNA-seq data of ESCs, which consists of 182 cells labeled by FACS sort criterion,
including 59, 58, and 65 cells belonging to "G1" phase, "S" phase, and "G2M" phase,
respectively [48, 49].
531
532 ***Retrotransposons.*** Retrotransposons built from RepeatMasker annotations were
533 downloaded from the UCSC Table Browser (http://genome.ucsc.edu/). We kept the top
534 six categories of counts for the downstream analysis, including Alu, B2, B4, MaLR, L1,
535 and ERVK.
536
HiCS
Preprocess contact probability for each chromosome of the individual cell. We first
divided each chromosome into bins of specific size (40kb in this study), and counted
the contact for each bin pair. Next, we modeled each chromosome as an unweighted
network (each bin is one node, and each bin pair with non-zero contacts is added as one
edge), and implemented a classic graph embedding method node2vec, which applies a
biased random walks procedure, to compute the contact probability of edges by
computing the cosine similarity of any two node embedding vectors, and obtained the

preprocessed matrix A (**Fig. 1a**) [50]. We only kept the top 5% pairs for downstream
analysis and the diagonal pairs were removed in our study.
547
Detect domain boundaries for each individual cell. Inspired by a fast density-based
clustering method designed for grouping data points [51, 52], we take advantage of
finding the cluster centers to detect domain boundaries for each chromosome of
individual cells (**Fig. 1b and c**). Specifically, we define two indexes for each 40kb bin:
(1) insulation strength $\rho(i)$ of the i^{th} genomic position is defined as the ratio of
$(I_{i,intra} - I_{i,inter})$ and $(I_{i,intra} + I_{i,inter})$ [53] using a 800kb sliding window size:

$$I_{i,intra} = I_a + I_b,$$
$$I_{i,inter} = I_c,$$
$$\rho(i) = \frac{I_{i,intra} - I_{i,inter}}{I_{i,intra} + I_{i,inter}},$$

558 where I_a, I_b, and I_c respectively represent the summation of interaction frequencies for
559 region a, b, and c, and (2) minimum distance between the bin i and any other bin j with
560 higher insulation strength is defined as $\delta(i)$:

$$\delta(i) = \min_{j:\rho(j)>\rho(i)} |i - j|, \quad (\delta_{max} < MAX).$$

We search for higher insulation strength of bin i in the range of $MAX = 500$ (20M

genomic distance at 40kb resolution). Next, we define $\rho^{'}(i) = \rho^{'}(i)/\rho_{max}$ and

$\delta^{'}(i) = \delta(i)/\delta_{max}$ such that both $\rho^{'}(i)$ and $\delta^{'}(i)$ are within range [0,1]. Then, we

generate the rank γ of all bins for each chromosome by their $\eta(i) = \rho^{'}(i) \times \delta^{'}(i)$ in

the descending and normalized the rank of each bin by $\gamma^{'}(i) = \gamma(i)/\gamma_{max}$. We define

the optimal reflection point with $r = \underset{i=1:len}{\mathrm{argmin}}\, \eta(i)^2 + \gamma^{'}(i)^2$, where len is defined as

the number of bins for a specific chromosome. The boundaries of the optimal structure
are assigned by bins with $\eta(i) > \alpha \times \eta(r)$, $(\alpha = 1)$. The gap regions are defined by
$I_{i,inter} = 0$ or have no contact with any other bins.
571
572 ***Determine the hierarchical chromatin domains.*** We ran the above procedures of
573 detecting domain boundaries multiple times by define $\eta(i) > \alpha \times \eta(r)$ as a screening
574 selection for different genomic levels, where the range of α are set as (0.1, 10) (**Fig.**
575 **2a**). In our study, we selected multiple scaling parameters α as {0.2, 1, 4, 8} to
576 obtain the hierarchical domains of chromatin, according to the distribution of the

domain sizes and insulation strengths under different scaling parameters (**Fig. 2a and b**).

Clustering and annotating genomic positions with different types of regulatory factors. We applied the hierarchical clustering method to group regulatory factors and chromatin positions, respectively (**Fig. 4a**). The classes of chromatin positions with the ratio>1% of all chromatin positions were selected for downstream analysis, which led to 29 classes of chromatin positions, along with 27 different transcriptional factor classes.

We further applied the hierarchical clustering method to merge the 27 classes into 7 large clusters, based on the Pearson correlation of the normalization ratio of the mean counts for peaks of regulatory factor classes (**Supplementary Fig. S9b**). These clusters or sub-clusters were manually annotated based on the annotation information (**Supplementary Table S2**). We then merged different chromatin position classes into 12 large categories based on hierarchical clustering of correlation of regulatory factor classes, and manually annotated these categories based on the ratios of regulatory factor classes. We obtained and annotated 12 large chromatin position categories (consisting of 29 sub-categories), and 7 different regulatory factor clusters (consisting of 27 sub-clusters).

Data availability

All datasets analyzed in this study were published previously. The corresponding descriptions and preprocessing steps can be found in Supplementary Materials.

Software availability

The open-source HiCS python package and tutorial are available at GitHub (https://github.com/YusenYe/HiCS).

Acknowledgements

This work was supported by the National Natural Science Foundation of China [No. 62002275 to Y.Y., Nos. 62132015 & 61873198 to L.G., and Nos. 12126605&61621003 to S.Z.], the National Key Research and Development Program of China [No. 2019YFA0709501 to S.Z.], the Strategic Priority Research Program of the Chinese Academy of Sciences (CAS) [Nos. XDA16021400, XDPB17 to S.Z.].

Author Contributions

Y.Y. conceived the idea, implemented the algorithm and performed the analyses. Y.Y. interpreted the results. S.Z. and L.G. provided scientific insights on the applications. Y.Y. wrote the manuscript with feedback from all other authors. All of the authors read and approved the final manuscript.

Completing interests

The authors declare no completing interests.

References

1.   Kempfer, R. and A. Pombo, *Methods for mapping 3D chromosome*
architecture. Nature Reviews Genetics, 2020. **21**(4): p. 207-226.
2.   Misteli, T., *The Self-Organizing Genome: Principles of Genome*
Architecture and Function. Cell, 2020. **183**(1): p. 28-45.
3.   Dixon, J.R., et al., *Integrative detection and analysis of structural variation*
in cancer genomes. Nature Genetics, 2018. **50**(10): p. 1388-1398.
4.   Rao, Suhas S.P., et al., *A 3D Map of the Human Genome at Kilobase*
Resolution Reveals Principles of Chromatin Looping. Cell, 2014. **159**(7): p.
1665-1680.
5.   Lieberman-Aiden, E., et al., *Comprehensive Mapping of Long-Range*
Interactions Reveals Folding Principles of the Human Genome. Science,
2009. **326**(5950): p. 289-293.
6.   Dixon, J.R., et al., *Topological domains in mammalian genomes identified*
by analysis of chromatin interactions. Nature, 2012. **485**(7398): p. 376-380.
7.   Stadhouders, R., G.J. Filion, and T. Graf, *Transcription factors and 3D*
genome conformation in cell-fate decisions. Nature, 2019. **569**(7756): p.
345-354.
8.   Rowley, M.J. and V.G. Corces, *Organizational principles of 3D genome*
architecture. Nature Reviews Genetics, 2018. **19**(12): p. 789-800.
9.   Hnisz, D., et al., *Super-Enhancers in the Control of Cell Identity and Disease.*
Cell, 2013. **155**(4): p. 934-947.
10.   Young, Richard A., *Control of the Embryonic Stem Cell State.* Cell, 2011.
144(6): p. 940-954.
11.   Zepeda-Martinez, J.A., et al., *Parallel PRC2/cPRC1 and vPRC1 pathways*
silence lineage-specific genes and maintain self-renewal in mouse
embryonic stem cells. Science Advances. **6**(14): p. eaax5692.
12.   Zuin, J., et al., *Cohesin and CTCF differentially affect chromatin architecture*
and gene expression in human cells. Proceedings of the National Academy
of Sciences, 2014. **111**(3): p. 996.
13.   Bintu, B., et al., *Super-resolution chromatin tracing reveals domains and*
cooperative interactions in single cells. Science, 2018. **362**(6413): p.
eaau1783.
14.   Tan, L., et al., *Three-dimensional genome structures of single diploid*
human cells. Science, 2018. **361**(6405): p. 924.
15.   Crane, E., et al., *Condensin-driven remodelling of X chromosome topology*
during dosage compensation. Nature, 2015. **523**(7559): p. 240-244.
16.   Li, X., et al., *DeTOKI identifies and characterizes the dynamics of chromatin*
TAD-like domains in a single cell. Genome Biology, 2021. **22**(1): p. 217.
17.   Nagano, T., et al., *Cell-cycle dynamics of chromosomal organization at*

single-cell resolution. Nature, 2017. **547**(7661): p. 61-67.

18.   Schwarzer, W., et al., *Two independent modes of chromatin organization*
revealed by cohesin removal. Nature, 2017. **551**(7678): p. 51-56.

19.   Yu, M. and B. Ren, *The Three-Dimensional Organization of Mammalian*
Genomes. Annual Review of Cell and Developmental Biology, 2017. **33**(1):
p. 265-289.

20.   Nora, E.P., et al., *Targeted Degradation of CTCF Decouples Local Insulation*
of Chromosome Domains from Genomic Compartmentalization. Cell, 2017.
169(5): p. 930-944.e22.

21.   Whyte, Warren A., et al., *Master Transcription Factors and Mediator*
Establish Super-Enhancers at Key Cell Identity Genes. Cell, 2013. **153**(2): p.
307-319.

22.   Kagey, M.H., et al., *Mediator and cohesin connect gene expression and*
chromatin architecture. Nature, 2010. **467**(7314): p. 430-435.

23.   Piunti, A. and A. Shilatifard, *The roles of Polycomb repressive complexes in*
mammalian development and cancer. Nature Reviews Molecular Cell
Biology, 2021. **22**(5): p. 326-345.

24.   Douillet, D., et al., *Uncoupling histone H3K4 trimethylation from*
developmental gene expression via an equilibrium of COMPASS,
Polycomb and DNA methylation. Nature Genetics, 2020. **52**(6): p. 615-625.

25.   Buecker, C., et al., *Reorganization of Enhancer Patterns in Transition from*
Naive to Primed Pluripotency. Cell Stem Cell, 2014. **14**(6): p. 838-853.

26.   Cole, M.F., et al., *Tcf3 is an integral component of the core regulatory*
circuitry of embryonic stem cells. Genes & development, 2008. **22**(6): p.
746-755.

27.   Niwa, H., et al., *Self-renewal of pluripotent embryonic stem cells is*
mediated via activation of STAT3. Genes & development, 1998. **12**(13): p.
2048-2060.

28.   Beattie, G.M., et al., *Activin A maintains pluripotency of human embryonic*
stem cells in the absence of feeder layers. Stem cells, 2005. **23**(4): p. 489-
495.

29.   Yang, Bin X., et al., *Systematic Identification of Factors for Provirus*
Silencing in Embryonic Stem Cells. Cell, 2015. **163**(1): p. 230-245.

30.   Furlong Eileen, E.M. and M. Levine, *Developmental enhancers and*
chromosome topology. Science, 2018. **361**(6409): p. 1341-1345.

31.   Rege, M., et al., *Chromatin Dynamics and the RNA Exosome Function in*
Concert to Regulate Transcriptional Homeostasis. Cell Reports, 2015. **13**(8):
p. 1610-1622.

32.   Surface, Lauren E., et al., *H2A.Z.1 Monoubiquitylation Antagonizes BRD2*
to Maintain Poised Chromatin in ESCs. Cell Reports, 2016. **14**(5): p. 1142-
1155.

33.   Thomas, M.J. and E. Seto, *Unlocking the mechanisms of transcription factor*

YY1: are chromatin modifying enzymes the key? Gene, 1999. **236**(2): p. 197-208.

34. Weintraub, A.S., et al., *YY1 Is a Structural Regulator of Enhancer-Promoter Loops.* Cell, 2017. **171**(7): p. 1573-1588.e28.

35. Lu, J.Y., et al., *Homotypic clustering of L1 and B1/Alu repeats compartmentalizes the 3D genome.* Cell Research, 2021. **31**(6): p. 613-630.

36. Liang, Z. and X.-D. Fu, *3D genome encoded by LINE and SINE repeats.* Cell Research, 2021. **31**(6): p. 603-604.

37. Hanna, C.W., et al., *Endogenous retroviral insertions drive non-canonical imprinting in extra-embryonic tissues.* Genome Biology, 2019. **20**(1): p. 225.

38. Guo, F., et al., *The Transcriptome and DNA Methylome Landscapes of Human Primordial Germ Cells.* Cell, 2015. **161**(6): p. 1437-1452.

39. Nuebler, J., et al., *Chromatin organization by an interplay of loop extrusion and compartmental segregation.* Proceedings of the National Academy of Sciences, 2018. **115**(29): p. E6697-E6706.

40. Ye, Y., L. Gao, and S. Zhang, *Circular Trajectory Reconstruction Uncovers Cell-Cycle Progression and Regulatory Dynamics from Single-Cell Hi-C Maps.* Advanced Science, 2019. **6**(23): p. 1900986.

41. Schuettengruber, B., et al., *Genome Regulation by Polycomb and Trithorax: 70 Years and Counting.* Cell, 2017. **171**(1): p. 34-57.

42. Jerkovic´, I. and G. Cavalli, *Understanding 3D genome organization by multidisciplinary methods.* Nature Reviews Molecular Cell Biology, 2021. **22**(8): p. 511-528.

43. Wang, X., Y. Luan, and F. Yue, *EagleC: A deep-learning framework for detecting a full range of structural variations from bulk and single-cell contact maps.* Science Advances. **8**(24): p. eabn9215.

44. Zhang, Y., et al., *Model-based Analysis of ChIP-Seq (MACS).* Genome Biology, 2008. **9**(9): p. R137.

45. Shen, Y., et al., *A map of the cis-regulatory sequences in the mouse genome.* Nature, 2012. **488**(7409): p. 116-120.

46. Hounkpe, B.W., et al., *HRT Atlas v1.0 database: redefining human and mouse housekeeping genes and candidate reference transcripts by mining massive RNA-seq datasets.* Nucleic Acids Research, 2021. **49**(D1): p. D947-D955.

47. Siepel, A., et al., *Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes.* Genome research, 2005. **15**(8): p. 1034-1050.

48. Buettner, F., et al., *Computational analysis of cell-to-cell heterogeneity in single-cell RNA-sequencing data reveals hidden subpopulations of cells.* Nature Biotechnology, 2015. **33**(2): p. 155-160.

49. Hao, Y., et al., *Integrated analysis of multimodal single-cell data.* Cell, 2021. **184**(13): p. 3573-3587.e29.

50. Zhou, J., et al., *Robust single-cell Hi-C clustering by convolution- and*

745     *random-walk–based imputation.* Proceedings of the National Academy of
746     Sciences, 2019. **116**(28): p. 14011.
747 51. Rodriguez, A. and A. Laio, *Clustering by fast search and find of density*
748     *peaks.* Science, 2014. **344**(6191): p. 1492-1496.
749 52. Ye, Y., L. Gao, and S. Zhang, *MSTD: an efficient method for detecting multi-*
750     *scale topological domains from symmetric and asymmetric 3D genomic*
751     *maps.* Nucleic Acids Research, 2019. **47**(11): p. e65-e65.
752 53. Luo, X., et al., *3D Genome of macaque fetal brain reveals evolutionary*
753     *innovations during primate corticogenesis.* Cell, 2021. **184**(3): p. 723-
754     740.e21.
755
756

     Figures and Tables

759
Figure 1. Illustration and efficiency of HiCS for determining the chromatin
domains. a. An illustrative example of the preprocessed single-cell Hi-C contact map
(top) and the insulation strength (ρ) of genomic positions (bottom). HSMD (δ)
represents the minimum distance between the bin and any other bins with higher
strengths. **b, c.** The decision graph (b) and the normalization value of $\eta = \rho \times \delta$ in a
decreasing order (c) for the domain boundaries (colored in red) in the optimal structural
identification parameter. The decision graph of optimal structural identification
parameter (zoom box in c). **d.** A local example of domain boundaries from map in **a** at

the region Chr2:80.8Mb-108Mb with different methods. **e, f.** Comparison of different

methods for domain sizes and insulation strengths. **g.** Runtime (seconds) of different

methods or parameters. **h.** The average number of CTCF peaks, H3K4me3 peaks, and

HK genes at domain boundaries of single cells. The above results (d-h) are implemented

by different methods across all single cells.

774
Figure 2. Identification of hierarchical chromatin domains. a. Density distribution
of domain sizes at different multi-scale parameters. **b.** An example for hierarchical
chromatin domains at the region Chr2:80.8Mb-108Mb. Circles mark three boundaries
(two boundaries in level 1 have a magnitude difference in local insulation strength, and
the boundary in level 3 has lower insulation strength than one of the boundaries in level
1).

Figure 3. Regulatory factors navigate the preference of genomic position forming boundaries. a. The ratio distribution of genomic positions forming boundaries at less than a given ratio of single cells in the left y-axis (such as 99.8% of genomic positions form boundaries in at least 1% of cells, respectively), and the number distribution of genomic positions forming boundaries in the right y-axis as the ratio of single cells increases. **b-c.** The concentration scores (b) and the average number of ATAC-seq peaks and multiple regulatory elements (ATAC: ATAC-seq peaks, HK genes, Exp: gene expression value, DEGs: differentially expressed genes, E: enhancers, SEs) (c) around domain boundaries. **d.** The concentration scores for CTCF, cohesin, mediator- and Pol II-associated factors. **e.** The average number of peaks for CTCF and cohesin around domain boundaries. In b-e, the results were detected in the different genomic

scales across all single cells. The concentration score is defined in **Supplementary Methods**.

Figure 4. The systematic analysis on genomic position categories. a. The bi-cluster of chromatin positions and regulatory factors. **b.** The column normalization ratio and the average number of peaks for regulatory factor-classes on genomic position

categories (different dot color represents different categories in Fig. 4a). **c.** The column

normalization ratio of different elements or factors, including SC_ratio (the ratio of

single cells forming boundaries), ATAC, DamID (DamID-seq), HK genes, Cyc (mark

genes for cell cycle), Exp (gene expression value), DEGs, Variable (variable scores for

genes), E (Enhancers), and SEs, on genomic position categories. **d.** The heatmap shows

the column normalization ratio of different retrotransposons based on genomic position

categories.

Figure 5. A schematic showing different genomic position categories. a. The graph presents the different clusters based on retrotransposons in the area marked by different colors of dotted line containing multiple functional groups associated with different regulatory factor-classes (the sub-categories serial number in above and the ratio occupying from all positions are marked for each functional group). The shades of purple on the background of circles indicate the preference of genomic positions forming boundaries in single cells for each functional group, and the shades of green on the background of these areas marked by dotted line indicate the degree of chromatin accessibility of genomic positions for each cluster based on retrotransposons. **b.** Preference scores of different chromatin landscapes categories across different cell states. Statistical significance is calculated by Welch's t-test (*P < 0.05, **P < 0.01, and ***P < 0.001). The preference score is defined in **Supplementary Methods**.