SMALL NONCODING RNA CONTENTS OF RAT EPIDIDYMAL EXTRACELLULAR VESICLES INCLUDING A PUTATIVE NOVEL SMALL RNA

Running Head: Small noncoding RNAs in epididymal EVs

Keywords: sncRNA, extracellular vesicles, epididymis, epididymosomes, miRNA, tRNA fragments, CpG islands, rat

Ross Gillette[1], Shyamal Waghwala[2], Andrea C. Gore[1†]

[1] The Center for Molecular Carcinogenesis and Toxicology, Division of Pharmacology and Toxicology, College of Pharmacy, The University of Texas at Austin, Austin, TX, 78712, USA

[2] College of Natural Sciences, The University of Texas at Austin, Austin, TX, 78712, USA

AUTHOR CONTRIBUTIONS

Conceptualization: R.G.; Methodology: R.G.; Investigation: R.G.; Formal Analysis: R.G., S.W.; Writing – Original Draft: R.G.; Writing – Review & Editing: R.G., A.C.G.; Funding Acquisition: R.G., A.C.G.

[†] Corresponding Author: andrea.gore@austin.utexas.edu

ABSTRACT

Extracellular vesicles (EVs) released from the epididymal epithelium (epididymosomes)

impart functional competence on sperm as they transit the epididymis by merging with sperm and

releasing a complex repertoire of molecules. The cargo of epididymosomes includes small

noncoding RNAs (sncRNAs) that are modified by external factors such as stress, nutrition, and

drug use. If incorporated into sperm, the EV sncRNA cargo can affect offspring and lead to

heritable phenotypes. In the current study we characterized the RNA contents of EVs collected

from the caput epididymis of adult male rats in order to fill a gap in knowledge in this species and

to establish a sncRNA profile. Small RNAs of EVs were isolated from the caput portion of the

epididymis of adult male rats, and sequenced on a NovaSeq 6000 on a SP flow cell in a single-end

50 bp configuration. The resulting reads were checked for quality, trimmed for adapter sequences,

aligned to the unmasked rat genome (Rnor 6), and assigned an annotation designation. The

majority of RNA reads aligned to either tRNA fragments (79.1%) or piRNA (18.1%) loci. Micro

RNAs (miRNAs) accounted for a surprisingly small proportion of reads (0.18%). The third largest

category of aligned reads (1.5%) was in intergenic space and not strictly associated with canonical

sncRNA loci. In-depth investigation determined these latter reads (~19 nt) aligned strictly within

the boundaries of known CpG islands (CpGi), which have not previously been reported to express

any form of sncRNA. These newly described "CpGi sRNAs" could not consistently be accounted

for by overlaying features of any other annotation type (including rRNA and piRNA). The CpGi

sRNAs have characteristics of RNA fragments that can associate with the Argonaute/PIWI family

of proteins and therefore could have regulatory function via RNA induced silencing or de novo

DNA methylation. We propose that CpGi sRNAs constitute a new family of sncRNA that may

represent an important and unreported class of regulatory RNA in gametes.

INTRODUCTION

Spermatozoa are incapable of fertilization as they exist the testes [1] and only gain motility and the capacity for fertilization as they transit the epididymis [2]. This process requires epididymosomes [3–6], a type of extracellular vesicle (EV) within the microvesicle category [7–9]. Epididymosomes are released from the epididymal epithelium via apocrine secretion [10], fuse to spermatozoa [11], aid in their maturation and functionalization [4,12,13], and mark dead or dying spermatozoa for elimination [14,15]. Epididymosomes carry a complex repertoire of ions, proteins and small non-coding RNAs (sncRNA) [4,12,16,17] that are transferred to [18,19] and presumably alter the function of the spermatozoa or impart functional consequences on the fertilized embryo [20]. Importantly, sncRNA molecules transferred to spermatozoa have the capacity to alter the epigenetic landscape of either the spermatozoa or the fertilized embryo and thereby influence subsequent generations. Furthermore, the sncRNA contents of epididymosomes are subject to modification by the hormonal milieu [21,22] and environmental challenges such as psychosocial or metabolic stress [22–27], making them a potential vector for intergenerational effects in offspring. This key role was illuminated by studies showing that early life stress caused behavioral and physiological changes in the offspring, phenotypes that were recapitulated by microinjection of a small number of microRNAs into a fertilized egg subsequently used for *in vitro* fertilization [28,29].

The sncRNA repertoire of epididymosomes is complex and diverse [30]. The most abundant sncRNA found in epididymosomes is microRNA (miRNA) followed by smaller proportions of transfer RNA (tRNA) and ribosomal RNA (rRNA) [25,30,31]. The remainder consists of piwi-interacting RNA (piRNA), small nuclear RNA (snRNA), and small nucleolar RNA (snoRNA), as well as long non-coding RNA (lncRNA) [25]. Other species of sncRNA, such as YRNA and Vault RNA, have not yet been reported in epididymosomes but should not be ruled out for inclusion in these complex profiles. Furthermore, there are likely additional sncRNAs yet to be identified [32].

Much of the knowledge we have on the sncRNA contents of caput epididymosomes, which

are crucial for the final steps of spermatozoa maturation, is from mouse models. Very little has

been reported on the epididymosomes of rats, which are an important model system for

endocrinology, reproduction, and epigenetic transgenerational inheritance, and in which

behavioral work is considered to better translate to humans [33–35]. Here, we characterized the

contents of EVs derived from the caput epididymis of the male rat and describe what we believe

to be a previously undefined class of sncRNA.

MATERIALS AND METHODS

Animals and treatment

All animal experiments were conducted using humane procedures that were pre-approved

by the Institutional Animal Care and Use Committee at The University of Texas at Austin and in

Accordance with NIH guidelines. Three-month old male and female Sprague Dawley rats were

purchased (Envigo, Indianapolis, IN) and shipped to the Animal Resource Center at the University

of Texas at Austin and allowed two weeks to acclimate to the housing facility. All animals in the

colony were housed in a room with consistent temperature (~22C) and light cycle (14:10

light:dark) and had *ad libitum* access to filtered tap water and a rat chow with minimal

phytoestrogens (Teklad 2019: Envigo, Indianapolis, IN). After acclimation, female rats were

observed for vaginal cytology indicating proestrus, and therefore receptivity. Receptive female

rats were randomly paired with an experienced breeder male, receptivity was confirmed, and the

pair was left together overnight. The following morning, vaginal cytology was checked for the

presence of sperm and if present marked as embryonic day (E1). The pregnant dams were single-

housed and provided nesting material on E18.

On E8, the dams (N = 6) were randomly split into two groups (each N = 3) and fed ¼ Nilla

Wafers™ with either 3% DMSO or 1 mg/kg Aroclor 1221 (A1221) (#C-221N, Accustandard, Lot

#072-202-01 - an estrogenic polychlorinated biphenol) from E8 – E18 and postnatal day (PND) 1

– 21. Treatment was not considered or analyzed in any of the data presented in this manuscript as

the goal was to first characterize the composition of EV sncRNAs, with greater statistical power

attained by combining the 6 litters. Treatment details are provided here for transparency. At PND

21, all pups were weaned and housed in cages of two or three. Only males were used in the present

manuscript and were otherwise unmanipulated until euthanasia besides weekly handling to

acclimate each rat to the experimenters in order to reduce stress. Females were used for other

projects.

Sample Collection

At PND 105, 6 male rats were randomly selected (N = 1 per litter) and euthanized by rapid

decapitation. The testis and epididymis were removed via a small incision in the scrotum. The

epididymis was separated from the testis and segmented into three portions (caput, corpus, and

cauda). The caput of the epididymis was minced into small pieces with scissors and placed in warm

(37 C) M2 culture media (M7167, Millipore Sigma) with HEPES and 2% exosome depleted fetal

bovine serum (A2720803, ThermoFisher). The slurry was placed on a rocker for 30 minutes to

allow sperm and epididymal fluid to suspend in solution, the supernatant was removed and large

tissue chunks were excluded. The resulting supernatant was centrifuged at 500 X g for 5 minutes

to pellet and remove sperm. The supernatant was again removed, immediately frozen on dry ice,

and stored at -80 C until use.

Extracellular Vesicle Purification and Nucleic Acid Extraction

The buffer supernatant containing extracellular vesicles from the caput of the epididymis

was thawed on ice for 30 minutes prior to isolation. After it was fully thawed the buffer was mixed

by pipetting and then filtered with a 0.8 um syringe filter (Millex-AA, SLAA033DD, Millipore

Sigma) to remove any cells or large cellular debris. Extracellular vesicles and RNA were

sequentially isolated from the media using the Qiagen exoRNeasy Midi kit (#77144, Qiagen,

Germantown, MD) according to the manufacturer's protocol. Briefly, 200 ul of media from each

sample was passed through a spin column that selectively binds exosomes. The column was

washed to remove contaminants and debris, and then eluted in Qiazol lysis buffer (#79306, Qiagen,

Germantown, MD). Chloroform (AAJ67241AP, FischerScientific) was added, mixed by shaking,

and incubated at room temperature to allow phase separation. Phase separation was aided by

centrifugation at 12,000 X g for 15 minutes at 4 C and the aqueous phase was aspirated and passed

through a second membrane spin column that selectively binds RNA. The membrane was washed

to remove contaminants and the sample eluted in 13 ul RNase free water. M2 culture media and

2% exosome depleted fetal bovine serum was used as a negative control to determine exogenous

contamination and subjected to the same extracellular vesicle and RNA isolation procedures

described above. These negative controls were analyzed via particle analysis and for resulting

nucleic acids to ensure there were no exogenous or contaminating exosomes or nucleic acids. None

were found.

Extracellular Vesicle and RNA Quality Control

An aliquot of the extracellular vesicles from the same samples described above were

isolated using the exoEasy Maxi Kit (76064, Qiagen) according to the manufacturer's protocols.

The resulting EVs from two samples were analyzed using a NanoSight 300 (Malvern Panalytical)

in 5 replicates to establish a size distribution of the resulting EVs and to ensure there were no

contaminating particles. RNA extracted from isolated EVs was first quantified (ND-1000,

ThermoScientific) and diluted for size distribution analysis and quality control using the small

RNA (5067-1548, Agilent) and pico RNA kits (5067-1514, Agilent) on a BioAnalyzer 2100

(Agilent). We found there was no cellular contamination indicated by a lack of 18s and 28s

ribosomal RNA and that the majority (~80%) of extracted RNA was in the 20-40 bp range,

indicative of small RNA molecules.

Library Preparation, Sequencing, and Quality Control

Library preparation was performed at the University of Texas Genomic Sequencing

Facility using the NEBNext Small RNA library preparation kit (E7330, New England

Biolabs). Samples were prepared with 14 cycles of PCR and the final product was size selected

using a 3% gel cassette on the Blue Pippin instrument (Sage Sciences), with the parameters set to

105-165 bp. Final size selected libraries were checked for quality on the Agilent BioAnalyzer with

High Sensitivity DNA analysis kit (5067-4626, Agilent) to confirm proper size selection. The

Kapa Library Quantification kit for Illumina libraries (KK4602) was used to determine the loading

concentrations prior to sequencing. Samples were sequenced on a NovaSeq 6000 Single Read,

SR50 run, with reads counts ranging from 25 to 32 million per sample.

Analysis Pipeline

Raw RNA reads were first passed through quality control (FastQC) and checked for read

quality and size distribution which showed excellent average read quality (avg. PHRED > 36) and

expected read length (50 bp). The raw reads were trimmed for Illumina adaptors (QuasR,

BioConductor) and again passed through quality control (FastQC) to determine per base read

quality and size distribution, which again showed excellent average read quality (avg. PHRED >

36) and an expected read length distribution (~30 nt) based on electrophoresis performed before

library prep. The trimmed reads were aligned (RBowtie, Bioconductor) to the unmasked rat

genome (Rnor v6) with strict matching parameters to ensure reads were aligned to best-hit

locations but were allowed multiple mapping locations (n = 12) due to the promiscuous nature of

small RNA origins. If the alignment parameters were met but a read mapped to multiple locations,

the read was randomly assigned to one of the similar best hit locations. Alignment efficiency was

analyzed for cross-species contamination or PCR primer amplification artifacts and was found to

be highly efficient (~85%). The aligned reads were assigned a feature annotation (Seqmonk, R,

BioConductor), visualized, and analyzed for read count and length distribution. Our annotation

pipeline considered reads aligned to microRNA (miRNA), piwi-interacting RNA (piRNA),

ribosomal RNA (rRNA), small nucleolar RNA (snoRNA), small nuclear RNA (snRNA), transfer

RNA (tRNA), long non-coding RNA (lncRNA), vault RNA, and YRNA. We also obtained

annotation tracks for nested and simple repeats (UCSC genome database) and microsatellites

(UCSC genome database). Reads were assigned an annotation designation only after considering

all other annotation types co-occurring in the same region.

RESULTS

The size distribution of rat caput epididymosomes

Particle tracking (NanoSight 300) was used to determine the size distribution of the

extracellular vesicles isolated from the caput of the rat epididymis using the Qiagen exoEasy Maxi

kit as described above. The isolated EVs from two separate samples were analyzed 5 times each.

The first sample had a mean EV size of 181.4 nm (SE 3), mode of 152 nm (SE 2.5), and standard

deviation of 40.6 nm (SE 1.1) and the second a mean of 178.7 (SE 0.5), mode of 156.1 (SE 3.8),

and standard deviation of 45.7 nm (SE 1.8). The size distribution and density of the extracellular

vesicles analyzed are shown in **Figure 1**. Approximately ~93% of all particles tracked were

between 50 and 250 nm in size, which are characteristic of epididymosomes [4].

The small RNA contents of rat caput EVs

The sncRNA contents of rat epididymosomes were dominated by reads that aligned to

tRNA (79.1%) or piRNA loci (18.1% - **Figure 2A**). We were surprised at the relatively low

abundance of miRNA in rats considering that mice models show substantial miRNA cargo in caput

epididymosomes [17]. All other annotation types, including miRNAs, accounted for the remaining

2.8% of the aligned reads. When tRNA and piRNA reads were removed from analysis and aligned

reads were analyzed as a proportion of the remaining reads (**Figure 2B**), the largest remaining

category belonged to reads that aligned to CpG islands (55.2%), which is not a known unit of

sncRNA expression (discussed below). The remaining reads aligned to YRNA (24.6%), lncRNA

(7.5%), miRNA (6.4%), rRNA (4.3%), and snoRNA (1.8%).

We analyzed the length of the reads aligned and assigned to each respective sncRNA

category to determine if our annotation pipeline performed well and to describe the subspecies of

various sncRNAs (**Table 1**). The two canonical categories with known read lengths were checked

first. The average read length for miRNA (22 nt) was exactly as expected and showed a 90-

percentile range of 19 – 27 nt. We set a cut-off of 10 reads per million (RPM) to determine which

miRNA were loaded into caput EVs and identified 15 miRNAs, the majority of which are known

to exist either in sperm or EVs in various models, and 5 uncharacterized miRNAs (**Table 2**).

We identified 81 piRNA loci that were substantially (> 10 RPM) expressed across the

genome, with clusters on chromosome 1, 2, 10, 13, 14 and 17. Approximately 60% (49/81) of the

expressed piRNA loci overlapped with known tRNA loci; these accounted for a significant portion

of all piRNA annotated reads (82.3%). These are a known subclass of piRNAs termed tRNA-

derived piRNA (td-piRNAs -- [36]) so we further analyzed their origin and determined that

virtually all of these reads aligned with the 5' end of tRNA loci (98.98%). Finally, we categorized

the tRNA type from which piRNAs were generated and determined that the majority were derived

from either 5'-tRNA^{Gly} (53.3%) or 5'-tRNA^{Glu} (32.6%) loci. 5'-tRNA^{Lys} (8.5%) and 5'-tRNA^{Val}

(2.3%) originating reads were also identified. The average read length for piRNA was also as

expected (30 nt) with a 90-precentile range of 26 – 32 nt.

Small RNA fragments are prevalent in rat epididymosomes

We then analyzed other sncRNA categories where read lengths can be variable depending

on how the RNA is processed. We found that reads aligned to tRNAs, which are ~70 – 100 nt [37],

had an average read length of 30 nt and a 90-precentile range of 28 – 31 nt (**Table 1**). This result

suggests that rat epididymosomes do not carry full length tRNAs, but rather carry either tRNA

fragments (tRFs) or tRNA-halves, which are 14 – 30 nt and 30 – 40 nt in length respectively [38].

We further categorized tRNA reads by aligning them to either the 5' or 3' end of a given tRNA

loci and discovered that 99.35% of all reads aligned to the 5' end (**Figure 3A**). Given their size

distribution, we were able to further categorize these reads. The tRNA reads show a 28 – 30 nt size

distribution indictive of tRF-5c fragments, the longest of the three known 5'-tRFs, or tRNA-halves

[39]. Alignment of reads to the tRNA structure (tRNAvis) confirmed their identity as tRF-5c

fragments because reads did not align to the anti-codon stem loop, which would be characteristic

of 5' tRNA-halves [38]. Because of the significant overlap of reads identified between piRNAs

and tRNA loci (presented above), we analyzed the proportion of tRNA reads that could be

accounted for by piRNA loci. We found that the majority tRNA reads (~81%) were unique to

tRNAs and could not be accounted for by overlapping piRNA loci. Finally, we analyzed the amino-

acid feature that tRNA reads were derived from and found that there was an over-representation

of reads derived from Glycine-tRNAs (81.4%). Reads were also found to originate from glutamate

(7.7%), histidine (6.9%), and lysine (1.7%) tRNA loci across the genome (**Figure 3B**). The size

distribution for 3' tRNA reads (**Figure 3C**) was similar but more broad compared to 5' tRNA

reads (**Figure 3D**).

We then looked at reads aligned to rRNA loci, which typically have a broad length range

depending on their rRNA type of origin (5s: ~120 nt [40], 5.8s: ~ 150 nt [41], 18s: 1874 nt [42],

and 28s: 4,802 nt [43]). In our samples, rRNAs had an average length of 22 nt and a 90-percentile

range of 17 – 31 nt (**Table 1**). These results suggest that rat caput epididymosomes do not carry

full length rRNA but instead carry rRNA fragments (rRFs). rRNA is expressed as a 45s pre-rRNA,

which contains the transcripts for 18s, 5.8s, and 28s rRNA, organized as cassettes of tandem

repeats on the short arms of chromosomes 3, 11, and 12 of the rat genome [44]. Reference genomes

are poorly annotated for these repeating cassettes because the number of repeats often varies

between individuals. Hence, these regions are typically masked in alignment reference genomes.

Nonetheless, there are a number of 5s rRNAs, which are expressed separately from the 45s pre-

rRNA, and 5.8s rRNA loci that are annotated in the rat reference genome that we analyzed.

Approximately 80% of rRNA derived reads aligned to 5.8s loci (**Figure 4A**). Subcategories of

rRFs have been described from human samples where short (18-19 nt), intermediate (24-25 nt),

and long (32-33 nt) rRFs are expressed depending on the rRNA type (e.g. 5s vs 5.8s) [45]. Our

data generally fit the categories described in human samples; 5s rRNA resulted in either short or

long transcripts (**Figure 4B**) whereas 5.8s rRNA resulted in intermediate fragments (**Figure 4C**).

Finally, we observed that the Y RNA was present in caput EVs at levels ~4 times that of

miRNA, which is surprising because we are not aware of any reports of Y RNA included in

epididymal EVs. Full length Y RNAs are ~80 – 110 nt in length [46]. We identified 13 (of 28) Y

RNA loci with substantial expression (> 10 RPKM). These reads demonstrated a sharp peak at 31

nt and a smaller peak at 22 nt with a 90-percentile range of 22 – 31 nt (**Table 1**). These results

suggest that rat caput EVs include fragments of Y RNA as a part of their sncRNA repertoire.

Small RNA molecules are expressed from within CpG islands

A subset of reads could not be accounted for by known and annotated sncRNA. We

identified dense clusters of expression that appeared to be expressed within the boundaries of GC-

rich CpG islands that were not fully accounted for by any other overlapping features (**Figure 5A**).

In total we found significant expression (>10 RPKM) within the boundaries of 50 CpG islands.

We visually inspected each of these CpG islands and were able to rule out the majority of them

due to overlapping features that had canonical expression patterns expected from features like

rRNA and piRNA (**Figure 5B**) or tRNA (**Figure 5C**). These loci were removed from further

analyses and the remaining 12 CpG islands were treated as an annotation category for sncRNAs.

We found this category to be the third most abundant behind tRNA and piRNA reads (**Figure 2A**

and **B**) and that the reads derived from CpG islands had a length distribution pattern that was

distinct from all of the other analyzed categories (**Figure 2C**) with a median length of 19 nt and a

90-percentile range of 15 – 28 nt (**Table 1**). As expected, the GC content of reads aligned to CpG

islands was substantial (~78%) compared to the average of all aligned reads (~50%).

We calculated a signal-to-noise ratio to determine if the reads we observed within CpG

islands were due to random chance [45]. When compared to the number of reads per kilobase

million (RPKM) over a 10 kb rolling window across the entire genome, our identified CpG islands

had a 609 s/n ratio. When compared to all CpG islands across the entire genome, our identified

CpG islands had a 92 s/n ratio. Put simply, the reads contained with our identified CpG islands

were 609 times more abundant than a random 10kb window in the genome and 92 times more

abundant than an average CpG island, indicating that the reads we observed were very unlikely to

be due to random chance.

Because CpG islands are not known units of small RNA expression, nor should they be

capable of such expression, we then tried to systematically demonstrate that the observed reads

were due to overlying features. We observed that CpG islands that contained more piRNA loci

appeared to be associated with more reads (**Figure 6A**), but when normalized for the read depth

and length of the CpG island (RPKM) the relationship was virtually non-existent ($R^2 = 0.077$,

Table 3). We quantified the reads that aligned under known piRNA loci and found that ~48% of

the total reads within the boundaries of CpG islands were also associated with piRNA loci.

However, we are hesitant to classify these reads as piRNA because they do not follow the canonical

length distribution associated with piRNA (~30nt) identified here and elsewhere [47]. None of the

reads (0.0%) within CpG islands were expressed from known tRNA loci. In a subset of the CpG

islands we observed, there was an abundance of predicted (eponine) transcriptional start sites

(**Figure 6B**), but this was not always the case (e.g. **Figure 5A**).

Finally, half (6/12) of the identified CpG islands were associated with annotated 5.8s rRNA

loci (**Table 3**). Functional rRNAs are expressed in tandem repeating cassettes that are processed

from a 45s precursor transcript that contains the functional 5.8s, 18s, and 28s rRNA after

processing. The rat genome is known to express these repeats from the short arms of chromosomes

3, 11, and 12 [44] on which none of our identified CpG islands reside. Notwithstanding, the rat

genome is poorly annotated for rRNA, so to overcome this shortcoming in the available

annotations we used BLAST to align the raw sequences expressed from CpG islands to the

available rRNA sequences: 45s, 32s (which includes the 28s rRNA and 5' sequence between the

transcription start site and 28s), 28s, 18s, and 5.8s. Approximately one third (~34.3%) of all reads

derived from CpG islands aligned to some form of rRNA or precursor rRNA. Of those reads that

aligned to any form of rRNA, the vast majority were derived from the 28s sequence (86.1% --

29.6% of all CpG island reads), while the 18s (11.15% -- 3.9% of all CpG island reads), and 5.8s

(0.16% -- 0.05% of all CpG island reads) accounted for a small proportion of the reads. The

remaining reads (2.6% -- 0.74% of all reads) aligned to the external or internal transcribed spacers

within the full length rRNA precursor transcript. In summary, while it appears that the CpG islands

we identified as expressing 19 nt small RNAs were in the vicinity of heretofore unannotated rRNA

or piRNA loci, the reads observed expressed within CpG islands cannot be fully accounted for by

either of these designations nor are their expression profiles congruent.

DISCUSSION

Epididymosomes from the caput are an essential component of sperm maturation and

acquisition of function [5,48,49]. They are also implicated in the control of heritable non-genetic

phenotypes, as their sncRNA cargo is altered by stress [19,22,25], diet [23,26,50], and alcohol

consumption [27]. Here, we provide the first comprehensive characterization of small RNAs

derived from caput EVs in the rat. We show that EVs isolated from the caput epididymis match

the size range of epididymosomes and that their small RNA contents are dominated by tRFs and

piRNA and contain far fewer miRNAs than expected from other organisms (Mice - [25,30,31],

Humans - [51]). We also identify Y RNA fragments in caput EVs for the first time in any organism,

although they are expressed in EVs of other organs [52–54]. Finally, we identify a potentially

novel small RNA molecule that is expressed from GC-rich CpG islands that cannot be accounted

for by known overlying small RNA features, and have a unique size distribution that is distinct

from other small RNAs analyzed.

The sncRNA composition of rat caput EVs differs from that in mice

The small RNA contents of EVs from the mouse caput epididymis are primarily miRNAs

(~60% - [25,30,31]), with over 350 expressed [17]. This is in contrast to the rat in which miRNAs

were a small portion (0.18%) of the small RNA complement. Of these, we identified 15 miRNAs,

7 of which overlapped with those in mice (miR-143, let-7c, let-7i, miR-26a, miR-99a, miR-143,

miR-148 [17]). The most abundant miRNA we identified in rats, miR-184, did not meet the

threshold for abundance in mice [17]. MiR-143, which is a hallmark of EVs in mice [17] and

humans [30], was also expressed in rats. There were also substantial species differences in tRFs,

which are the second most abundant category (~30%) in mouse caput EVs while piRNAs are

present at very low levels (<0.05% - [30,31]). Here, we show in rats that tRFs are by far the most

abundant category (~79%) and piRNAs are second most abundant (~18%). Further work is needed

to validate these findings and to understand why there are these dramatic species differences in the

EV sncRNA cargo.

tRFs dominate the cargo of rat caput EVs

tRFs, particularly from 5'-tRNA^{Gly} loci, dominated the cargo of caput EVs in the rat. As

sperm exit the testis, they contain few tRFs but their abundance gradually increases as sperm transit

from the proximal caput to the cauda [19,55]. In a mouse model, paternal diet changed the

abundance of 5'-tRF^{Gly} in sperm, which is implicated in the repression of endogenous

retroelements in pre-implantation embryos [19]. The authors suggested that EVs could deliver

these fragments, and we provide evidence that they are well poised to do so, and we confirmed the

exact species they identified (5'-tRF^{Gly}). We also identify 5'-tRF^{Glu} and 5'-tRF^{His} as prevalent in rat

caput EVs. The former (5'-tRF^{Glu}) has been implicated in playing a role in a high fat diet

intergenerational phenotype [26] via transcriptional regulation, directing alternate splicing, and

acetylation and phosphoprotein activation [27]. The latter (5'-tRF^{His}) has been implicated in

responses to low protein diets [19]. Further work is needed to elucidate their functional roles in

our rat model.

We found that tRFs in caput EVs are almost exclusively derived from the 5' end of tRNA

loci. There are two possible explanations for the overrepresentation of 5' tRFs. First, rat caput

epididymosomes may be selectively loaded with 5'-tRF fragments; these have actions similar to

RNA interference as governed by miRNAs, which act through the Argonaute pathway [56] to

silence endogenous retroelements in embryonic stem cells and embryos [19]. Alternatively, our

observation may be due to a technical bias in the sequencing library preparation procedure [38].

Libraries are amplified with PCR during preparation, which can be prematurely aborted if an RNA

molecule contains modified tRNA nucleobases that would be too short for sequencing, thereby

excluded, and not detected during analysis. In order to confirm the dominating presence of 5'-tRFs

in rat caput epididymosomes, a specific analysis pipeline (streamlined platform for observing

tRNA – SPOt) would need to be used to prevent observation bias [57].

piRNAs are abundant and coincide with 5' tRNA fragments

piRNAs represented a substantial proportion (18.1%) of EVs in the rat caput epididymis.

This is contrary to what is found in mouse models (< 0.05% -- [30,31]) but not entirely unexpected

as rat pachytene sperm are densely populated with piRNAs [58] depending on the stage of

development [59,60] presumably to control the expression of transposable elements in the

germline [61] by directing the catalyzation of de novo DNA methylation to suppress their

expression [62]. In the rat, unlike in the mouse, piRNAs appear poised to be delivered to sperm

via EVs in the caput epididymis. The reasons for this discrepancy between rats and mice could be

due to a number of reasons. First, this may represent a fundamental difference in the reproductive

biology between the two species. Second, it may be because the piRNAs that do exist in mouse

caput EVs are typically categorized as tRNA-halves instead of piRNA. We found that a majority

of reads that aligned with piRNA loci (~80%) also align to the 5' end of tRNA loci, which could

explain why they are missed during analysis in the mouse. The overlap of these two features

(piRNAs and tRFs) and expression from these loci has precedence. tRNA derived-piRNA (td-

piRNA) have been described in the testes of marmosets where Piwi proteins were found to bind

reads mapping to the 5'-tRNA^{Glu}, 5'-tRNA^{Gly}, and 5'-tRNA^{Val} loci [63]. It is intriguing that these

three identified td-piRNAs account for three of the four td-piRNAs found here and we believe we

are the first to identify these td-piRNA in a position poised to be delivered to sperm in the

epididymis. Additionally, we append 5'-tRNA^{Lys}, the third most abundant td-piRNA in our

analysis, as a potentially significant td-piRNA.

rRNA fragments are present in three distinct sizes

rRFs have long been considered RNA degradation or apoptotic by-products and have been

disregarded for any functional value. Compared to miRNA, piRNA, and tRFs, little is known about

the function or categories of rRFs and they have only recently been described [45] with nearly no

information existing in rats that we could find. Interestingly, rRFs appear in immunoprecipitations

with the Argonaute complex in mice and humans, which suggests a role for translational regulation

[64]. In humans, three distinct categories of rRFs are known with differential expression in each

of the rRNA categories (e.g. 5s, 5.8s, 18s, and 28s); short (18-21 nt), intermediate (24-25 nt), and

long (26-33 nt) [45]. Our ability to map rRFs to the reference genome was limited by the lack of

annotations for rRNA in the rat genome, but given the annotations we did have, we identified a

similar expression profile to Cherlin et al. (2020) (in human samples -- [45]) in which three distinct

subtypes of rRFs were present: 18 nt & 30 nt rRFs originating from 5s loci and 22 nt rRFs from

5.8s loci. Furthermore, rRFs are sparsely reported in reproductive tissue (Humans -- [65,66],

Bovine -- [67]) and we believe we are the first to report their presence in caput EV samples, which

suggests a novel role for rRFs in reproduction.

19-nt small RNAs are expressed from within the boundaries of CpG Islands

There were 12 identified loci in the rat genome that corresponded with substantial

expression of 19-nt small RNAs that are GC rich and expressed within the boundaries of annotated

CpG islands. We postulate that this is a novel unit of expression for sncRNAs. CpG islands are

generally annotated based on the characteristics of their sequence; longer than 200-bp with a GC

content higher than 50% and an expected to observed ratio greater than 0.6 [68]. Many of the first

CpG islands were identified at the 5' end of "housekeeping" genes [69,70] but have since been

computationally and experimentally predicted across the genome, including in inter- and

intragenic space [71]. Parts of CpG islands are sometimes transcribed on the 5' or 3' end of

expressed genes and removed during splicing unless they extend into an exon, including sncRNAs

[71,72]. This expression pattern we observed within CpG islands does not appear to have a

relationship with any other annotation type; although we considered other nearby or overlying

rRNA cassettes or piRNA clusters, neither accounted for all of the reads or loci, and some loci had

neither feature. Nevertheless, we carefully inspected the involvement of both types of features.

For rRNAs, it is difficult to determine the precise location of cassettes because they are not

well annotated in the rat genome. By using sequence similarity via BLAST alignment, we

determined that ~34% of the small RNA expressed from CpG islands are found within the 45s pre-

rRNA transcript, the majority of those from the 28s sequence (~86%). Because of their 19 nt

length, we can say with certainty these reads are not from mature rRNAs. Furthermore, we are

hesitant to term these rRNA fragments for two reasons; first, the length of the reads (19 nt) from

annotated 5.8s rRNA within the identified CpG islands do not correspond to other 5.8s rRNA reads

(22 nt) found elsewhere in this data set; and second, the expression profile of the 19 nt reads are

bounded by the CpG island and not by the 5.8s rRNA or predicted position of 45s rRNA. It is

possible that the observed CpG islands may be a part of the ~30 kb non-transcribed spacer that

separates 45s repeats, but as the name implies, these portions are up or downstream of rRNA and

should not be transcribed.

A second possibility is that the reads we identified arise from piRNA clusters. The function

of piRNA is RNA-guided silencing mediated by Piwi proteins (Riwi in the rat) that are particularly

active in the germline of the testes [73,74]. piRNAs regulate the expression of transposable

elements [59,75] by inducing de novo DNA methylation [76]. Primary piRNA can be amplified

by "ping-pong amplification" [77] by binding to expressed transposable elements which are

cleaved to generate a secondary guide transcript that ensure the lack of transposition [59,78].

Primary piRNA are observed in developing germ cells [76] while secondary piRNAs, generated

by the ping-pong cycle, are thought only to be present before pachytene stages in the basal

compartment of the testes [59,76] after which only primary piRNAs should be present [76,79].

The transcripts we identified in caput EVs are well outside of the window of canonical piRNA

functionality and while they appear to be associated with piRNA clusters, when read depth is

corrected for by the length of a CpG island the association becomes non-existent. Furthermore,

their distinct 19-nt signature differentiates them from canonical piRNAs which are typically 26-

35 nt in length [80].

A final possibility we considered is that the 19-nt sRNA we observe are byproducts from

the generation of secondary piRNAs during which exonuclease activity should generate fragments

of a consistent size due to the portion of the transcript that is shielded by Piwi-complex itself [76].

There have been three reports we could find that have identified such byproducts [81–83].

Berninger et al. (2011) were the first to identify "19mers" expressed from both the sense and anti-

sense strands of piRNA clusters solely in the testis of rats, mice, and platypuses, but did not find

them in any other tissue type [81]. Oey et al. (2011) also described 19mer byproducts of piRNA

with a focus on repeat elements (LINEs, SINEs, LTRs) and IAPs where ping-pong amplification

is predicted [82]. They also found that 19mers that were equally present from both the sense and

antisense strands, were exclusive to the testes, and while they didn't specifically analyze or

mention it, their data show no GC content bias. Ichiyanagi et al. (2014) also observed 19mers

immediately upstream of piRNAs but dismissed them as piRNA amplification byproducts as per

Berninger et al. and Oey et al. (2011) [83]. The commonality of the piRNA amplification

byproducts identified in these reports is that they are 19nt fragments expressed from a broad

distribution on both strands relative to the piRNA cluster, a profile that does not fit our current

observation.

Based on this evidence, we do not believe that the alternatives (rRNA fragments,

piRNAs, or piRNA amplification byproducts) can account for our observation of GC-rich 19-nt

small RNAs. These "CpG island small RNAs" extend outside of the boundaries of annotated rRNA

and piRNA, which account for only a portion of the observations, and do not abide by their

respective canonical length characteristics. It is also possible that these observations could be the

19mer byproduct of piRNA amplification observed elsewhere [81–83] but there are 3 reasons we

do not think this is likely. First, we observe these reads in extracellular vesicles while the

byproducts of piRNA amplification are known only to exist in pre-pachytene spermatocytes and

should be derived from both the sense and anti-sense strands, a characteristic we do not see. If this

observation is due to 19mer piRNA amplification byproducts generated in the epididymis, then

our data provide functional significance to their existence not previously reported. Second, the

reads we observe are expressly within the boundaries of CpG islands with a GC content not

reported elsewhere. The specificity of the expression loci, and the lack of high GC content in other

reports make their identity as byproducts unlikely. Finally, we observe expression from multiple

CpG islands not linked with known piRNA loci. If these are indeed the byproduct of piRNA

amplification, our data then represent the identification of novel piRNA loci that would seem to

be important for the final steps of sperm maturation. Mature sperm transiting the epididymis should

be transcriptionally quiescent [84–87], and therefore transposons should not be expressed.

Ascertaining a functional role of these small RNAs requires further investigation.

Conclusions

EVs from the rat caput epididymis carry a complex repertoire of small RNAs and their

contents are substantially different from the mouse. The dominant features we identify are tRNA

fragments and piRNAs derived from tRNA loci. MicroRNAs are poorly represented in stark

contrast to mice. We also identify two types of small RNA not previously seen in caput EVs, rRNA

fragments and Y RNA fragments, and described a potentially novel small RNA we have termed

CpG island (CpGi) sRNAs. These data represent an exciting collection of possibilities for future

researchers studying basic reproductive biology and the complexities of epigenetic

transgenerational inheritance.

Conflict of Interest: The authors declare no conflict of interest.

Grant funding: Supported by and RO1 ES029464 to A.C.G. and PhRMA Foundation

Postdoctoral Fellowship to R.G.

Acknowledgement: The authors recognize Mandee Bell and Lindsay M. Thompson for their

assistance with animal husbandry, treatment, and sample collection. We would also like to

acknowledge the excellent work (small RNA library preparation and sequencing) performed by

the Genomic Sequencing and Analysis Facility at UT Austin, Center for Biomedical Research

Support. RRID#: SCR_021713.

REFERENCES

1. Hermo, L.; Pelletier, R. ‐ Marc; Cyr, D.G.; Smith, C.E. Surfing the Wave, Cycle, Life
History, and Genes/Proteins Expressed by Testicular Germ Cells. Part 1: Background to
Spermatogenesis, Spermatogonia, and Spermatocytes. *Microsc Res Techniq* 2010, *73*, 241–278,
doi:10.1002/jemt.20783.

2. Cornwall, G.A. New Insights into Epididymal Biology and Function. *Hum Reprod Update*
2009, *15*, 213–227, doi:10.1093/humupd/dmn055.

3. Sullivan, R.; Saez, F. Epididymosomes, Prostasomes, and Liposomes: Their Roles in
Mammalian Male Reproductive Physiology. *Reproduction* 2001, *146*, R21–R35,
doi:10.1530/rep-13-0058.

4. Sullivan, R. Epididymosomes: A Heterogeneous Population of Microvesicles with Multiple
Functions in Sperm Maturation and Storage. *Asian J Androl* 2015, *0*, 0, doi:10.4103/1008-
682x.155255.

5. Martin-DeLeon, P.A. Epididymosomes: Transfer of Fertility-Modulating Proteins to the
Sperm Surface. *Asian J Androl* 2015, *17*, 720–725, doi:10.4103/1008-682x.155538.

6. Martin-DeLeon, P.A. Epididymal SPAM1 and Its Impact on Sperm Function. *Mol Cell*
Endocrinol 2006, *250*, 114–121, doi:10.1016/j.mce.2005.12.033.

7. Fornés, M.W.; Barbieri, A.; Cavicchia, J.C. Morphological and Enzymatic Study of
Membrane ‐ bound Vesicles from the Lumen of the Rat Epididymis. *Andrologia* 1995, *27*, 1–5,
doi:10.1111/j.1439-0272.1995.tb02087.x.

8. Rejraji, H.; Sion, B.; Prensier, G.; Carreras, M.; Motta, C.; Frenoux, J.-M.; Vericel, E.;
Grizard, G.; Vernet, P.; Drevet, J.R. Lipid Remodeling of Murine Epididymosomes and
Spermatozoa During Epididymal Maturation. *Biol Reprod* 2006, *74*, 1104–1113,
doi:10.1095/biolreprod.105.049304.

9. Grimalt, P.; Bertini, F.; Fornes, M.W. HIGH-AFFINITY SITES FOR Beta-D-
GALACTOSIDASE ON MEMBRANE-BOUND VESICLES ISOLATED FROM RAT
EPIDIDYMAL FLUID. *Arch Andrology* 2009, *44*, 85–91, doi:10.1080/014850100262245.

10. Hermo, L.; Jacks, D. Nature's Ingenuity: Bypassing the Classical Secretory Route via
Apocrine Secretion. *Mol Reprod Dev* 2002, *63*, 394–410, doi:10.1002/mrd.90023.

11. Oh, J.S.; Han, C.; Cho, C. ADAM7 Is Associated with Epididymosomes and Integrated into
Sperm Plasma Membrane. *Mol Cells* 2009, *28*, 441–446, doi:10.1007/s10059-009-0140-x.

12. Zhou, W.; Iuliis, G.N.D.; Dun, M.D.; Nixon, B. Characteristics of the Epididymal Luminal
Environment Responsible for Sperm Maturation and Storage. *Front Endocrinol* 2018, *9*, 59,
doi:10.3389/fendo.2018.00059.

13. Frenette, G.; Lessard, C.; Sullivan, R. Polyol Pathway along the Bovine Epididymis. *Mol*
Reprod Dev 2004, *69*, 448–456, doi:10.1002/mrd.20170.

14. Sutovsky, P.; Moreno, R.; Ramalho-Santos, J.; Dominko, T.; Thompson, W.E.; Schatten, G.
A Putative, Ubiquitin-Dependent Mechanism for the Recognition and Elimination of Defective
Spermatozoa in the Mammalian Epididymis. *J Cell Sci* 2001, *114*, 1665–1675,
doi:10.1242/jcs.114.9.1665.

15. D'Amours, O.; Frenette, G.; Bordeleau, L.-J.; Allard, N.; Leclerc, P.; Blondin, P.; Sullivan,
R. Epididymosomes Transfer Epididymal Sperm Binding Protein 1 (ELSPBP1) to Dead
Spermatozoa During Epididymal Transit in Bovine1. *Biol Reprod* 2012, *87*, Article 94, 1-11,
doi:10.1095/biolreprod.112.100990.

16. Belleanne, C.; Calvo, zequiel; Caballero, J.; Sullivan, R. Epididymosomes Convey Different
Repertoires of MicroRNAs Throughout the Bovine Epididymis1. *Biol Reprod* 2013, *89*, Article
30, 1-11, doi:10.1095/biolreprod.113.110486.

17. Reilly, J.N.; McLaughlin, E.A.; Stanger, S.J.; Anderson, A.L.; Hutcheon, K.; Church, K.;
Mihalas, B.P.; Tyagi, S.; Holt, J.E.; Eamens, A.L.; et al. Characterisation of Mouse
Epididymosomes Reveals a Complex Profile of MicroRNAs and a Potential Mechanism for
Modification of the Sperm Epigenome. *Sci Rep-uk* 2016, *6*, 31794, doi:10.1038/srep31794.

18. Caballero, J.N.; Frenette, G.; Belleannée, C.; Sullivan, R. CD9-Positive Microvesicles
Mediate the Transfer of Molecules to Bovine Spermatozoa during Epididymal Maturation. *Plos*
One 2013, *8*, e65364, doi:10.1371/journal.pone.0065364.

19. Sharma, U.; Conine, C.C.; Shea, J.M.; Boskovic, A.; Derr, A.G.; Bing, X.Y.; Belleannee, C.;
Kucukural, A.; Serra, R.W.; Sun, F.; et al. Biogenesis and Function of TRNA Fragments during
Sperm Maturation and Fertilization in Mammals. *Science* 2016, *351*, 391,
doi:10.1126/science.aad6780.

20. Conine, C.C.; Sun, F.; Song, L.; Rivera-Pérez, J.A.; Rando, O.J. Small RNAs Gained during
Epididymal Transit of Sperm Are Essential for Embryonic Development in Mice. *Dev Cell* 2018,
46, 470-480.e3, doi:10.1016/j.devcel.2018.06.024.

21. Suryawanshi, A.R.; Khan, S.A.; Joshi, C.S.; Khole, V.V. Epididymosome‐Mediated
Acquisition of MMSDH, an Androgen‐Dependent and Developmentally Regulated Epididymal
Sperm Protein. *J Androl* 2012, *33*, 963–974, doi:10.2164/jandrol.111.014753.

22. Rodgers, A.B.; Morgan, C.P.; Bronson, S.L.; Revello, S.; Bale, T.L. Paternal Stress Exposure
Alters Sperm MicroRNA Content and Reprograms Offspring HPA Stress Axis Regulation. *J*
Neurosci 2013, *33*, 9003--9012, doi:10.1523/jneurosci.0914-13.2013.

23. Chen, Q.; Yan, M.; Cao, Z.; Li, X.; Zhang, Y.; Shi, J.; Feng, G.; Peng, H.; Zhang, X.; Zhang,
Y.; et al. Sperm TsRNAs Contribute to Intergenerational Inheritance of an Acquired Metabolic
Disorder. *Science* 2016, *351*, 397, doi:10.1126/science.aad7977.

24. Fullston, T.; Teague, E.M.C.O.; Palmer, N.O.; DeBlasio, M.J.; Mitchell, M.; Corbett, M.;
Print, C.G.; Owens, J.A.; Lane, M. Paternal Obesity Initiates Metabolic Disturbances in Two
Generations of Mice with Incomplete Penetrance to the F2 Generation and Alters the
Transcriptional Profile of Testis and Sperm MicroRNA Content. *Faseb J* 2013, *27*, 4226–4243,
doi:10.1096/fj.12-224048.

25. Short, A.; Fennell, K.; Perreau, V.; Fox, A.; O'Bryan, M.; Kim, J.H.; Bredy, T.; Pang, T.;
Hannan, A. Elevated Paternal Glucocorticoid Exposure Alters the Small Noncoding RNA Profile
in Sperm and Modifies Anxiety and Depressive Phenotypes in the Offspring. *Translational
Psychiatry* 2016, *6*.

26. Barbosa, T. de C.; Ingerslev, L.R.; Alm, P.S.; Versteyhe, S.; Massart, J.; Rasmussen, M.;
Donkin, I.; Sjögren, R.; Mudry, J.M.; Vetterli, L.; et al. High-Fat Diet Reprograms the
Epigenome of Rat Spermatozoa and Transgenerationally Affects Metabolism of the Offspring.
Mol Metab 2015, *5*, 184–197, doi:10.1016/j.molmet.2015.12.002.

27. Rompala, G.R.; Mounier, A.; Wolfe, C.M.; Lin, Q.; Lefterov, I.; Homanics, G.E. Heavy
Chronic Intermittent Ethanol Exposure Alters Small Noncoding RNAs in Mouse Sperm and
Epididymosomes. *Frontiers Genetics* 2018, *9*, 32–32, doi:10.3389/fgene.2018.00032.

28. Gapp, K.; Jawaid, A.; Sarkies, P.; Bohacek, J.; Pelczar, P.; Prados, J.; Farinelli, L.; Miska, E.;
Mansuy, I.M. Implication of Sperm RNAs in Transgenerational Inheritance of the Effects of
Early Trauma in Mice. *Nat Neurosci* 2014, *17*, 667–669, doi:10.1038/nn.3695.

29. Chan, J.C.; Morgan, C.P.; Leu, N.A.; Shetty, A.; Cisse, Y.M.; Nugent, B.M.; Morrison, K.E.;
Jašarević, E.; Huang, W.; Kanyuch, N.; et al. Reproductive Tract Extracellular Vesicles Are
Sufficient to Transmit Intergenerational Stress and Program Neurodevelopment. *Nat Commun*
2020, *11*, 1499, doi:10.1038/s41467-020-15305-w.

30. Trigg, N.A.; Eamens, A.L.; Nixon, B. The Contribution of Epididymosomes to the Sperm
Small RNA Profile. *Reproduction* 2019, *1*, R209–R223, doi:10.1530/rep-18-0480.

31. Hutcheon, K.; McLaughlin, E.A.; Stanger, S.J.; Bernstein, I.R.; Dun, M.D.; Eamens, A.L.;
Nixon, B. Analysis of the Small Non-Protein-Coding RNA Profile of Mouse Spermatozoa
Reveals Specific Enrichment of PiRNAs within Mature Spermatozoa. *Rna Biol* 2017, *14*, 1776–
1790, doi:10.1080/15476286.2017.1356569.

32. Wei, H.; Zhou, B.; Zhang, F.; Tu, Y.; Hu, Y.; Zhang, B.; Zhai, Q. Profiling and Identification
of Small RDNA-Derived RNAs and Their Potential Biological Functions. *Plos One* 2013, *8*,
e56842, doi:10.1371/journal.pone.0056842.

33. Wei, Y.; Schatten, H.; Sun, Q.-Y. Environmental Epigenetic Inheritance through Gametes
and Implications for Human Reproduction. *Hum Reprod Update* 2015, *21*, 194–208,
doi:10.1093/humupd/dmu061.

34. Ellenbroek, B.; Youn, J. Rodent Models in Neuroscience Research: Is It a Rat Race? *Dis*
Model Mech 2016, *9*, 1079–1087, doi:10.1242/dmm.026120.

35. Burns, S.B.; Szyszkowicz, J.K.; Luheshi, G.N.; Lutz, P.-E.; Turecki, G. Plasticity of the
Epigenome during Early-Life Stress. *Semin Cell Dev Biol* 2018, *77*, 115–132,
doi:10.1016/j.semcdb.2017.09.033.

36. Honda, S.; Kawamura, T.; Loher, P.; Morichika, K.; Rigoutsos, I.; Kirino, Y. The Biogenesis
Pathway of TRNA-Derived PiRNAs in Bombyx Germ Cells. *Nucleic Acids Res* 2017, *45*, 9108–
9120, doi:10.1093/nar/gkx537.

37. Krahn, N.; Fischer, J.T.; Söll, D. Naturally Occurring TRNAs With Non-Canonical
Structures. *Front Microbiol* 2020, *11*, 596914, doi:10.3389/fmicb.2020.596914.

38. Shen, Y.; Yu, X.; Zhu, L.; Li, T.; Yan, Z.; Guo, J. Transfer RNA-Derived Fragments and
TRNA Halves: Biogenesis, Biological Functions and Their Roles in Diseases. *J Mol Med* 2018,
96, 1167–1176, doi:10.1007/s00109-018-1693-y.

39. Kumar, P.; Anaya, J.; Mudunuri, S.B.; Dutta, A. Meta-Analysis of TRNA Derived RNA
Fragments Reveals That They Are Evolutionarily Conserved and Associate with AGO Proteins
to Recognize Specific RNA Targets. *Bmc Biol* 2014, *12*, 78, doi:10.1186/s12915-014-0078-0.

40. Gray, M.W.; Burger, G.; Lang, B.F. Mitochondrial Evolution. *Science* 1999, *283*, 1476–
1481, doi:10.1126/science.283.5407.1476.

41. Walker, T.A.; Pace, N.R. 5.8S Ribosomal RNA. *Cell* 1983, *33*, 320–322, doi:10.1016/0092-
8674(83)90413-0.

42. Chan, Y.L.; Gutell, R.; Noller, H.F.; Wool, I.G. The Nucleotide Sequence of a Rat 18 S
Ribosomal Ribonucleic Acid Gene and a Proposal for the Secondary Structure of 18 S
Ribosomal Ribonucleic Acid. *J Biol Chem* 1984, *259*, 224–230, doi:10.1016/s0021-
9258(17)43645-3.

43. Hadjiolov, A.A.; Georgiev, O.I.; Nosikov, V.V.; Yavachev, L.P. Primary and Secondary
Structure of Rat 28 S Ribosomal RNA. *Nucleic Acids Res* 1984, *12*, 3677–3693,
doi:10.1093/nar/12.8.3677.

44. Sasaki, M.; Nishida, C.; Kodama, Y. Characterization of Silver-Stained Nucleolus Organizer
Regions (Ag-NORs) in 16 Inbred Strains of the Norway Rat, Rattus Norvegicus. *Cytogenet*
Genome Res 1986, *41*, 83–88, doi:10.1159/000132208.

45. Cherlin, T.; Magee, R.; Jing, Y.; Pliatsika, V.; Loher, P.; Rigoutsos, I. Ribosomal RNA
Fragmentation into Short RNAs (RRFs) Is Modulated in a Sex- and Population of Origin-
Specific Manner. *Bmc Biol* 2020, *18*, 38, doi:10.1186/s12915-020-0763-0.

46. Wolin, S.L.; Steitz, J.A. Genes for Two Small Cytoplasmic Ro RNAs Are Adjacent and
Appear to Be Single-Copy in the Human Genome. *Cell* 1983, *32*, 735–744, doi:10.1016/0092-
8674(83)90059-4.

47. Beyret, E.; Liu, N.; Lin, H. PiRNA Biogenesis during Adult Spermatogenesis in Mice Is
Independent of the Ping-Pong Mechanism. *Cell Res* 2012, *22*, 1429–1439,
doi:10.1038/cr.2012.120.

48. Sullivan, R.; Frenette, G.; Girouard, J. Epididymosomes Are Involved in the Acquisition of
New Sperm Proteins during Epididymal Transit. *Asian J Androl* 2007, *9*, 483–491,
doi:10.1111/j.1745-7262.2007.00281.x.

49. Caballero, J.; Frenette, G.; D'Amours, O.; Belleannée, C.; Lacroix‐Pepin, N.; Robert, C.;
Sullivan, R. Bovine Sperm Raft Membrane Associated Glioma Pathogenesis‐Related 1‐like
Protein 1 (GliPr1L1) Is Modified during the Epididymal Transit and Is Potentially Involved in
Sperm Binding to the Zona Pellucida. *J Cell Physiol* 2012, *227*, 3876–3886,
doi:10.1002/jcp.24099.

50. Fullston, T.; Teague, E.M.C.O.; Palmer, N.O.; DeBlasio, M.J.; Mitchell, M.; Corbett, M.;
Print, C.G.; Owens, J.A.; Lane, M. Paternal Obesity Initiates Metabolic Disturbances in Two
Generations of Mice with Incomplete Penetrance to the F2 Generation and Alters the
Transcriptional Profile of Testis and Sperm MicroRNA Content. *The FASEB Journal* 2013, *27*,
4226–4243.

51. Belleannée, C.; Calvo, E.; Thimon, V.; Cyr, D.G.; Légaré, C.; Garneau, L.; Sullivan, R. Role
of MicroRNAs in Controlling Gene Expression in Different Segments of the Human Epididymis.
Plos One 2012, *7*, e34996, doi:10.1371/journal.pone.0034996.

52. Kowalski, M.P.; Krude, T. Functional Roles of Non-Coding Y RNAs. *Int J Biochem Cell*
Biology 2015, *66*, 20–29, doi:10.1016/j.biocel.2015.07.003.

53. Driedonks, T.A.P.; Hoen, E.N.M.N.-'t Circulating Y-RNAs in Extracellular Vesicles and
Ribonucleoprotein Complexes; Implications for the Immune System. *Front Immunol* 2019, *9*,
3164, doi:10.3389/fimmu.2018.03164.

54. Driedonks, T.A.P.; Mol, S.; Bruin, S. de; Peters, A.-L.; Zhang, X.; Lindenbergh, M.F.S.;
Beuger, B.M.; Stalborch, A.-M.D. van; Spaan, T.; Jong, E.C. de; et al. Y-RNA Subtype Ratios in
Plasma Extracellular Vesicles Are Cell Type- Specific and Are Candidate Biomarkers for
Inflammatory Diseases. *J Extracell Vesicles* 2020, *9*, 1764213,
doi:10.1080/20013078.2020.1764213.

55. Stanger, S.J.; Bernstein, I.R.; Anderson, A.L.; Hutcheon, K.; Dun, M.D.; Eamens, A.L.;
Nixon, B. The Abundance of a Transfer RNA-Derived RNA Fragment Small RNA
Subpopulation Is Enriched in Cauda Spermatozoa. *Exrna* 2020, *2*, 17, doi:10.1186/s41544-020-
00058-x.

56. Gagnon, K.T.; Corey, D.R. Argonaute and the Nuclear RNAs: New Pathways for RNA-
Mediated Control of Gene Expression. *Nucleic Acid Ther* 2012, *22*, 3–16,
doi:10.1089/nat.2011.0330.

57. Grelet, S.; McShane, A.; Hok, E.; Tomberlin, J.; Howe, P.H.; Geslain, R. SPOt: A Novel and
Streamlined Microarray Platform for Observing Cellular TRNA Levels. *Plos One* 2017, *12*,
e0177939, doi:10.1371/journal.pone.0177939.

58. Stermer, A.R.; Reyes, G.; Hall, S.J.; Boekelheide, K. Small RNAs in Rat Sperm Are a
Predictive and Sensitive Biomarker of Exposure to the Testicular Toxicant Ethylene Glycol
Monomethyl Ether. *Toxicol Sci* 2019, *169*, 399–408, doi:10.1093/toxsci/kfz041.

59. Aravin, A.A.; Sachidanandam, R.; Girard, A.; Fejes-Toth, K.; Hannon, G.J. Developmentally
Regulated PiRNA Clusters Implicate MILI in Transposon Control. *Science* 2007, *316*, 744–747,
doi:10.1126/science.1142612.

60. Smorag, L.; Zheng, Y.; Nolte, J.; Zechner, U.; Engel, W.; Pantakani, D.V.K. MicroRNA
Signature in Various Cell Types of Mouse Spermatogenesis: Evidence for Stage‐specifically
Expressed MiRNA‐221, ‐203 and ‐34b‐5p Mediated Spermatogenesis Regulation. *Biol*
Cell 2012, *104*, 677–692, doi:10.1111/boc.201200014.

61. Newkirk, S.J.; Lee, S.; Grandi, F.C.; Gaysinskaya, V.; Rosser, J.M.; Berg, N.V.; Hogarth,
C.A.; Marchetto, M.C.N.; Muotri, A.R.; Griswold, M.D.; et al. Intact PiRNA Pathway Prevents
L1 Mobilization in Male Meiosis. *Proc National Acad Sci* 2017, *114*, E5635–E5644,
doi:10.1073/pnas.1701069114.

62. Peng, J.C.; Lin, H. Beyond Transposons: The Epigenetic and Somatic Functions of the Piwi-
PiRNA Mechanism. *Curr Opin Cell Biol* 2013, *25*, 190–194, doi:10.1016/j.ceb.2013.01.010.

63. Hirano, T.; Iwasaki, Y.W.; Lin, Z.Y.-C.; Imamura, M.; Seki, N.M.; Sasaki, E.; Saito, K.;
Okano, H.; Siomi, M.C.; Siomi, H. Small RNA Profiling and Characterization of PiRNA
Clusters in the Adult Testes of the Common Marmoset, a Model Primate. *Rna* 2014, *20*, 1223–
1237, doi:10.1261/rna.045310.114.

64. Guan, L.; Grigoriev, A. Computational Meta-Analysis of Ribosomal RNA Fragments:
Potential Targets and Interaction Mechanisms. *Nucleic Acids Res* 2021, *49*, gkab190-,
doi:10.1093/nar/gkab190.

65. Goodrich, R.J.; Anton, E.; Krawetz, S.A. Spermatogenesis, Methods and Protocols. *Methods*
Mol Biology 2012, 385–396, doi:10.1007/978-1-62703-038-0_33.

66. Johnson, G.D.; Sendler, E.; Lalancette, C.; Hauser, R.; Diamond, M.P.; Krawetz, S.A.
Cleavage of RRNA Ensures Translational Cessation in Sperm at Fertilization. *Mhr Basic Sci*
Reproductive Medicine 2011, *17*, 721–726, doi:10.1093/molehr/gar054.

67. Almiñana, C.; Tsikis, G.; Labas, V.; Uzbekov, R.; Silveira, J.C. da; Bauersachs, S.;
Mermillod, P. Deciphering the Oviductal Extracellular Vesicles Content across the Estrous
Cycle: Implications for the Gametes-Oviduct Interactions and the Environment of the Potential
Embryo. *Bmc Genomics* 2018, *19*, 622, doi:10.1186/s12864-018-4982-5.

68. Gardiner-Garden, M.; Frommer, M. CpG Islands in Vertebrate Genomes. *J Mol Biol* 1987,
196, 261–282, doi:10.1016/0022-2836(87)90689-9.

69. Cooper, D.N.; Gerber-Huber, S. DNA Methylation and CpG Suppression. *Cell Differ Dev*
1985, *17*, 199–205, doi:10.1016/0045-6039(85)90488-9.

70. Bird, A.P. CpG-Rich Islands and the Function of DNA Methylation. *Nature* 1986, *321*, 209–
213, doi:10.1038/321209a0.

71. Medvedeva, Y.A.; Fridman, M.V.; Oparina, N.J.; Malko, D.B.; Ermakova, E.O.;
Kulakovskiy, I.V.; Heinzel, A.; Makeev, V.J. Intergenic, Gene Terminal, and Intragenic CpG
Islands in the Human Genome. *Bmc Genomics* 2010, *11*, 48–48, doi:10.1186/1471-2164-11-48.

72. Hackenberg, M.; Previti, C.; Luque-Escamilla, P.L.; Carpena, P.; Martínez-Aroza, J.; Oliver,
J.L. CpGcluster: A Distance-Based Algorithm for CpG-Island Detection. *Bmc Bioinformatics*
2006, *7*, 446–446, doi:10.1186/1471-2105-7-446.

73. Aravin, A.; Gaidatzis, D.; Pfeffer, S.; Lagos-Quintana, M.; Landgraf, P.; Iovino, N.; Morris,
P.; Brownstein, M.J.; Kuramochi-Miyagawa, S.; Nakano, T.; et al. A Novel Class of Small
RNAs Bind to MILI Protein in Mouse Testes. *Nature* 2006, *442*, 203–207,
doi:10.1038/nature04916.

74. Girard, A.; Sachidanandam, R.; Hannon, G.J.; Carmell, M.A. A Germline-Specific Class of
Small RNAs Binds Mammalian Piwi Proteins. *Nature* 2006, *442*, 199–202,
doi:10.1038/nature04917.

75. Carmell, M.A.; Girard, A.; Kant, H.J.G. van de; Bourc'his, D.; Bestor, T.H.; Rooij, D.G. de;
Hannon, G.J. MIWI2 Is Essential for Spermatogenesis and Repression of Transposons in the
Mouse Male Germline. *Dev Cell* 2007, *12*, 503–514, doi:10.1016/j.devcel.2007.03.001.

76. Aravin, A.A.; Sachidanandam, R.; Bourc'his, D.; Schaefer, C.; Pezic, D.; Toth, K.F.; Bestor,
T.; Hannon, G.J. A PiRNA Pathway Primed by Individual Transposons Is Linked to De Novo
DNA Methylation in Mice. *Mol Cell* 2008, *31*, 785–799, doi:10.1016/j.molcel.2008.09.003.

77. Brennecke, J.; Aravin, A.A.; Stark, A.; Dus, M.; Kellis, M.; Sachidanandam, R.; Hannon,
G.J. Discrete Small RNA-Generating Loci as Master Regulators of Transposon Activity in
Drosophila. *Cell* 2007, *128*, 1089–1103, doi:10.1016/j.cell.2007.01.043.

78. Gunawardane, L.S.; Saito, K.; Nishida, K.M.; Miyoshi, K.; Kawamura, Y.; Nagami, T.;
Siomi, H.; Siomi, M.C. A Slicer-Mediated Mechanism for Repeat-Associated SiRNA 5' End
Formation in Drosophila. *Science* 2007, *315*, 1587–1590, doi:10.1126/science.1140494.

79. Betel, D.; Sheridan, R.; Marks, D.S.; Sander, C. Computational Analysis of Mouse PiRNA
Sequence and Biogenesis. *Plos Comput Biol* 2007, *3*, e222, doi:10.1371/journal.pcbi.0030222.

80. Marzec, M. Size Does Matter: PiRNA and MiRNA Targeting. *Trends Biochem Sci* 2021, *47*,
287–288, doi:10.1016/j.tibs.2021.12.001.

81. Berninger, P.; Jaskiewicz, L.; Khorshid, M.; Zavolan, M. Conserved Generation of Short
Products at PiRNA Loci. *Bmc Genomics* 2011, *12*, 46–46, doi:10.1186/1471-2164-12-46.

82. Oey, H.M.; Youngson, N.A.; Whitelaw, E. The Characterisation of PiRNA-Related 19mers
in the Mouse. *Bmc Genomics* 2011, *12*, 315–315, doi:10.1186/1471-2164-12-315.

83. Ichiyanagi, T.; Ichiyanagi, K.; Ogawa, A.; Kuramochi-Miyagawa, S.; Nakano, T.; Chuma,
S.; Sasaki, H.; Udono, H. HSP90α Plays an Important Role in PiRNA Biogenesis and
Retrotransposon Repression in Mouse. *Nucleic Acids Res* 2014, *42*, 11903–11911,
doi:10.1093/nar/gku881.

84. Kierszenbaum, A.L.; Tres, L.L. Structural and Transcriptional Features of the Mouse
Spermatid Genome. *J Cell Biology* 1975, *65*, 258–270, doi:10.1083/jcb.65.2.258.

85. Hecht, N.B. Molecular Mechanisms of Male Germ Cell Differentiation. *Bioessays* 1998, *20*,
555–561, doi:10.1002/(sici)1521-1878(199807)20:7<555::aid-bies6>3.0.co;2-j.

86. Steger, K. Transcriptional and Translational Regulation of Gene Expression in Haploid
Spermatids. *Anat Embryol* 1999, *199*, 471–487, doi:10.1007/s004290050245.

87. Fischer, B.E.; Wasbrough, E.; Meadows, L.A.; Randlet, O.; Dorus, S.; Karr, T.L.; Russell, S.
Conserved Properties of Drosophila and Human Spermatozoal MRNA Repertoires. *Proc Royal*
Soc B Biological Sci 2012, *279*, 2636–2644, doi:10.1098/rspb.2012.0153.

88. Corral-Vazquez, C.; Salas-Huetos, A.; Blanco, J.; Vidal, F.; Sarrate, Z.; Anton, E. Sperm
MicroRNA Pairs: New Perspectives in the Search for Male Fertility Biomarkers. *Fertil Steril*
2019, *112*, 831–841, doi:10.1016/j.fertnstert.2019.07.006.

89. Sun, J.; Zhao, Y.; He, J.; Zhou, Q.; El-Ashram, S.; Yuan, S.; Chi, S.; Qin, J.; Huang, Z.; Ye,
M.; et al. Small RNA Expression Patterns in Seminal Plasma Exosomes Isolated from Semen
Containing Spermatozoa with Cytoplasmic Droplets versus Regular Exosomes in Boar Semen.
Theriogenology 2021, *176*, 233–243, doi:10.1016/j.theriogenology.2021.09.031.

90. Alves, M.B.R.; Celeghini, E.C.C.; Belleannée, C. From Sperm Motility to Sperm-Borne
MicroRNA Signatures: New Approaches to Predict Male Fertility Potential. *Frontiers Cell Dev*
Biology 2020, *8*, 791, doi:10.3389/fcell.2020.00791.

91. Liu, W.-M.; Pang, R.T.K.; Chiu, P.C.N.; Wong, B.P.C.; Lao, K.; Lee, K.-F.; Yeung, W.S.B.
Sperm-Borne MicroRNA-34c Is Required for the First Cleavage Division in Mouse. *Proc*
National Acad Sci 2012, *109*, 490–494, doi:10.1073/pnas.1110368109.

740
741

742   **FIGURES**

743   **Figure 1.**

744
745   The size distribution and density of the extracellular vesicles from two separate samples are
746   shown as analyzed by nano-particle tracking (NanoSight 300). The red dashed lines indicate the
747   boundaries of size for canonical epididymosomes (50 – 250 nm [4]).
748

749 **Figure 2.**
750

Results from sequencing small RNA derived from EVs in the rat caput are shown. **A)** The
proportion of small RNA reads assigned to each of 10 annotations in our analysis; tRNA and
piRNA account for the majority of small RNA in caput EVs while miRNAs are notably absent.
B) The remaining small RNA reads are shown as a proportion of the residual 2.8% not assigned
to tRNA or piRNA loci. CpG islands, which are not a known unit of small RNA expression,
account for the majority of the remaining reads. **C)** The size distribution of 4 well-defined
sncRNAs are shown and the one undefined category (CpG islands) demonstrates a unique size
profile (19 nt). The size distribution for piRNA (29-31 nt) and miRNA (22 nt) are exactly as
expected. Reads aligned to tRNA (30 - 31 nt) and rRNA (primary peak at 22 nt and small peaks
at 18 nt and 30 nt) both demonstrate size specific fragmentation.

762 **Figure 3.**

Detailed analysis of tRNA reads is shown. **A)** Reads align to the 5' end of tRNA loci almost
exclusively. **B)** 5'-tRNA^{Gly} is the primary tRNA fragment found in rat caput extracellular
vesicles, followed by a substantially smaller proportion of 5'-tRNA^{Glu} and 5'-tRNA^{His}. **C-D)** The
size distribution of 3' and 5' tRNA reads are shown, respectively. Reads from the 3' end of
tRNA are slightly longer with a broader distribution.

769 **Figure 4.**

770
Reads that aligned to annotated rRNA loci are shown by subcategory. The rat genome is poorly
annotated for rRNA, so the available 5s and 5.8s loci were used for categorization. **A)** The
majority of rRNA reads aligned to 5.8s rRNA. **B-C)** The size distribution for reads aligned to 5s
and 5.8s rRNA show profiles indicative of rRFs that are unique from one another. **B)** 5s rRFs
show a bimodal distribution with peaks at 18 and 30 nt. **C)** 5.8s rRFs show a single peak at 22 nt.

Figure 5

The log transformed density of reads is shown across three CpG islands with different
characteristics. Overlying feature annotations are shown indicated by color; Light blue = CpG
island, Green = piRNA, Black = predicted transcriptional start sites, red = 5.8s rRNA, and
orange = tRNA. **A)** Reads are shown expressed from within the boundaries of a CpG island on
chromosome 6. The reads were not associated with the piRNA (green) within the 3' CpG island
while the 5' CpG island had no other overlying features. **B)** Reads were identified within a CpG
island on chromosome 7 but the majority of them aligned to a piRNA (green) and rRNA (red)
feature. **C)** Reads were identified within a CpG island on chromosome 13 but they aligned
exclusively to piRNA and tRNA loci and are demonstrative of their typical expression profiles.
The CpG islands from **B & C** (and all other CpG islands like these) were removed from the
analysis of small RNAs expressed from CpG islands. The two in **A** are representative of those
that were used for further analysis.

Figure 6

The log transformed reads from a cluster of 8 CpG islands on chromosome 14 are shown with
the primary overlying features indicated by color and label. Light blue = CpG islands, Red = 5.8s
rRNA, Black = predicted transcriptional start sites, Green = piRNA. **A)** The boundaries of the
reads shown are clearly demarcated by CpG islands and not the overlying features such as rRNA
or piRNA. **B)** A detailed view of the largest CpG island in the chromosome 14 cluster. The reads
are not associated with either the rRNA or piRNA features within the CpG island and are not
typical of those features found elsewhere (Figure 5B & C).

TABLES

Table 1

Class	Mode	Median	Mean	SD	90% Range
CpG Islands	19	19	19.89	3.75	15 - 28
lncRNA	22	22	22.27	4.98	15 - 33
miRNA	22	22	22.04	2.34	19 - 27
piRNA	30	30	29.60	2.50	26 - 32
rRNA	22	22	22.65	4.06	17 - 31
snoRNA	27	24	23.44	5.12	15 - 32
snRNA	16	16	18.05	3.85	15 - 26
tRNAs	30	30	30.19	1.18	28 - 31
vault RNA	18	19	22.99	8.02	14 - 41
Y RNA	31	31	28.33	3.89	22 - 31

801
802 Descriptive statistics of the read length for each annotated small RNA category are shown with
803 the mode, median, mean, standard deviation, and length range from which 90 percent of the
804 reads are found.

805 **Table 2**

Name	Reads per Million	Found in Sperm	Found in EVs
miR-1b	16.2	--	--
miR-7a-1	19.7	[88]	[89]
miR-7a-2	19.9	[88]	[89]
miR-let-7c	10.0	[88]	[89]
miR-let-7i	11.7	[88]	[89]
miR-26a	11.7	[90]	--
miR-99a	87.8	[91]	--
miR-143	29.5	[19]	[19]
miR-148a	93.0	[88]	[89]
miR-184	493.7	[88]	[89]
AABR07005004.2	16.0		
AABR07029907.2	34.7		
AABR07057421.1	11.9		
AABR07064716.2	34.2		
AABR07064724.1	22.1		

806
Fifteen miRNA were identified as having substantial (> 10 RPM) expression in rat caput EVs, 10
of which are annotated and 5 that are putative. This is far fewer than expected based on mice
experiments. The majority of miRNAs identified have previously been reported in either sperm
or EVs (citation indicated), except for miR-1b. A miRNA considered a hallmark of caput EVs
(miR-143) was also identified in our rat data set.

Table 3.

Chromosome	Start Location	End Location	Length (nt)	Read Count	RPKM	piRNA Loci	pRNA Protein Loci	rRNA Binding Protein Loci	rRNA Loci
Chr 1	11,903,555	11,916,838	13,283	112,070	63.16	19	1	0	1
Chr 1	11,961,297	11,975,108	13,811	278,633	151.02	27	1	0	1
Chr 5	91,124,274	91,141,016	16,742	341,585	152.72	26	1	0	1
Chr 6	30,627,068	30,633,859	6,791	177,949	196.15	13	0	0	2
Chr 6	31,047,175	31,047,676	501	10,347	154.59	0	0	0	0
Chr 6	31,048,983	31,050,286	1,303	3,282	18.85	1	0	0	0
Chr 14	46,517,594	46,518,401	807	113,831	1,055.86	1	0	0	0
Chr 14	46,521,126	46,527,599	6,473	170,612	197.30	10	0	1	0
Chr 14	46,588,206	46,589,034	828	39,955	361.21	0	0	1	0
Chr 14	46,590,294	46,591,521	1,227	2,562	15.63	1	0	1	0
Chr 14	46,631,880	46,656,213	24,333	429,235	132.04	44	0	1	1
Chr 14	46,679,234	46,688,547	9,313	160,403	128.93	17	0	0	1

813
814   The genomic coordinates (Rnor v6 – Chromosome, Start Location, End Location) of 12 CpG
815   islands found to have substantial expression (> 10 RPKM) of ~19nt small RNA transcripts are
816   shown along with the length of the CpG island, raw read count, and length-corrected read count
817   (RPKM) found within each. The number of various overlying features found overlapping each
818   CpG island is also shown.