Interoceptive Awareness of the Breath Preserves Dorsal Attention Network Activity amidst

Widespread Cortical Deactivation: A Within-Participant Neuroimaging Study

Norman A. S. Farb[1,2], Zoey Zuo[2], Cynthia J. Price[3]

[1] Department of Psychology, University of Toronto Mississauga, Mississauga, Ontario, Canada,

L5L 1C6

[2] Department of Psychological Clinical Sciences, University of Toronto Scarborough,

Scarborough, Ontario, Canada, M1C 1A4

[3] Department of Biobehavioral Nursing and Health Informatics, University of Washington,

Seattle, Washington, USA, 98195

Corresponding Author:

Norman Farb

Department of Psychology

University of Toronto Mississauga

3359 Mississauga Road

Mississauga, ON L5L 1C6 Canada

Email: norman.farb@utoronto.ca

The authors declare that they have no conflicts of interest.

Abstract

Interoception, the representation of the body's internal state, serves as a foundation for emotion, motivation, and wellbeing. Yet despite its centrality in human experience, the neural mechanisms of interoception are poorly understood. The Interoceptive/Exteroceptive Attention Task (IEAT) is a novel neuroimaging paradigm that compares behavioral tracking of the respiratory cycle (Active Interoception) to tracking of a visual stimulus (Active Exteroception). Twenty-two healthy participants completed the IEAT during two separate scanning sessions (N = 44) as part of a randomized control trial of Mindful Awareness in Body-oriented Therapy (MABT). Compared to Exteroception, Interoception deactivated somatomotor and prefrontal regions. Greater interoceptive sensibility (MAIA scale) predicted sparing from deactivation along the anterior cingulate cortex (ACC) and left-lateralized language regions. The right insula—typically described as a primary interoceptive cortex—was only implicated by its further deactivation during an exogenously paced respiration condition (Active Matching). Psychophysiological interaction analysis characterized Active Interoception as promoting greater ACC connectivity with lateral frontal and parietal regions commonly referred to as the Dorsal Attention Network. Interoception of the breath may therefore involve reduced cortical *activity* but greater *connectivity*, with greater sensibility sparing cortical inhibition within well-characterized attentional networks. In contrast to a literature that relates detection of liminal signals such as the heartbeat to anterior insula activity, attention towards accessible body sensations such as the breath may lead to a low activity, high connectivity state in which sensory signals from the body may be better discerned.

Significance Statement (120 words maximum)

Interoception, the representation of the body's internal state, is poorly understood compared to the external senses, with existing neuroimaging studies failing to match task difficulty between interoceptive and exteroceptive tasks. The present study used a novel fMRI task to compare interoceptive and exteroceptive attention, and how this distinction was moderated by self-reported interoceptive awareness. The results implicate three novel interoceptive mechanisms: interoception reduces cortical *activity* while increasing *connectivity*, wherein awareness is linked to preserved activation of the brain's salience network and left-lateralized language regions. These findings characterize interoception as a lower activity state in which awareness depends upon the ability to notice and report on body signals typically obscured by the processing of exteroceptive information and other forms of cognition.

Introduction

Interoception, the sense of the body's internal state, is central to human experience, providing homeostatic cues (Strigo & Craig, 2016) that inform emotion (Barrett, 2017; Wiens, 2005), motivation (Craig, 2003; Critchley & Garfinkel, 2017), and wellbeing (Chen et al., 2021; Farb et al., 2015; Paulus & Khalsa, 2021). Yet interoception's neural mechanisms are still poorly understood when compared to the five "canonical" human senses (Chen et al., 2021).

Our understanding has been limited by a focus on interoceptive *accuracy* by the research community. For example, heartbeat detection paradigms model the ability to detect a liminal cardiac signal (e.g., Garfinkel et al., 2015). Superior detection accuracy is linked to greater activation of the anterior cingulate (ACC) (Critchley et al., 2004) and anterior insula (Chong et al., 2015), hubs of the brain's "Salience Network" (SLN) (Chand & Dhamala, 2016; Seeley et al., 2007). Yet SLN recruitment may indicate a broader error-monitoring system that does not distinguish interoception from other sensory processes (Baltazar et al., 2021).

Moreover, differences in interoceptive accuracy may not be especially relevant for wellbeing. Clinical populations often show normal interoceptive accuracy (Desmedt et al., 2020), and clinically-efficacious interoception-focused interventions such as mindfulness training do not improve accuracy (Khalsa et al., 2020; Parkin et al., 2014). Patients with anxiety or panic disorders have historically demonstrated *superior* accuracy (Ehlers & Breuer, 1992; Zoellner & Craske, 1999), but tend to catastrophize interoceptive experience (Domschke et al., 2010). The complex relationship between interoceptive accuracy and wellbeing demonstrates the need to expand the scope of interoceptive research to other, more clinically relevant constructs.

The ability to sustain interoceptive *attention* remains a compelling if underexplored determinant of mental health. Improving the capacity to skillfully attend to interoceptive cues

remains a central target of contemplative interventions such as mindfulness training (Gibson, 2019; Price & Weng, 2021). Such interventions address intolerance for interoceptive signals, which otherwise leads vulnerable individuals into patterns of experiential avoidance (Anestis et al., 2007; Hayes et al., 1996). Interoceptive *sensibility* — how one appraises interoceptive signals — shows stronger links to wellbeing than interoceptive accuracy (Ferentzi et al., 2019; Schuette et al., 2021), with a growing literature relating trust in interoceptive signals to greater wellbeing (Calì et al., 2015; Trevisan et al., 2019).

Interoceptive attention seems to recruit a distinct neural network from exteroception that features the middle insula (Haruki & Ogawa, 2021), consistent with its proposed role as a bridge to the prefrontal cortex from primary representation cortices in the posterior insula and somatosensory regions (Craig, 2009; Farb et al., 2013a; Pollatos et al., 2007). Activity in the middle insula characterizes wellbeing (Nord et al., 2021), with hypoactivation linked to depression (Avery et al., 2014; Farb et al., 2010; Farb et al., 2022), and hyperactivation linked to anxiety (Kerr et al., 2016; Tan et al., 2018). Breath monitoring in particular appears to strengthen coherence across a frontotemporal-insular network (Herrero et al., 2018).

Yet neuroimaging studies characterizing interoceptive attention to the breath have been confounded by non-equivalent task demands. Farb et al. (2013) contrasted exteroceptive and interoceptive attention by comparing a visual condition requiring behavioral responses against passive breath monitoring; Wang et al. (2019) compared a more difficult interoception task against an easier exteroception task. In both cases, the more demanding task recruited the Dorsal Attention Network (DAN), while the easier/passive task implicated the Default Mode Network (DMN), confounding conclusions on interoceptive representation.

We therefore developed a novel fMRI paradigm for characterizing the neural dynamics of breath-focused interoception vs. a more closely balanced exteroception task. We hypothesized (H1) that interoception would result in greater somato-insular recruitment than exteroception, but reduced DAN activation. We further hypothesized (H2) that greater self-reported interoceptive sensibility would correlate with greater SLN activity during interoception, consistent with a greater capacity for interoceptive monitoring and integration. We also explored: (H3) a paced-breathing condition to understand impact of endogenous vs. exogenous respiratory control, and (H4) functional connectivity differences between Interoception and Exteroception.

Materials and Methods

Experimental Design

This fully within-participant study was conducted to validate a novel interoceptive attention task as part of an NIH-funded pilot study, a two-group randomized control trial to examine the neural correlates of interoceptive awareness in the context of Mindful Awareness in Body-oriented Therapy (MABT) training (www.ClinicalTrials.gov identifier: NCT03583060). MABT is a well-validated, clinical intervention the focuses on developing the interoceptive capacities of identifying, accessing, and appraising internal bodily signal to support adaptive emotion regulation (Price & Hooven, 2018). To maximize power, data from the study's two assessment timepoints were combined into a single dataset, with group and time included in all statistical models as nuisance covariates.

Participants

Twenty-two right-handed study participants (11 male and 11 female), of adult age (mean: 36.1 years, range: 18-62) completed both baseline and post-intervention assessments. Eleven participants (50% of the sample) were randomly allocated to receive eight MABT sessions,

delivered individually once per week for eight weeks. Twenty participants self-identified as

Caucasian, one as African American, and two as Hispanic. Their highest education levels were

high school (n = 5), 2 years of college (n = 2), Bachelor's degree (n = 8), and Master's degree or

higher (n = 7).

Sample size for the study was determined by simulation-based power analysis. Given the

difficulties in *a priori* registration of all fMRI contrasts, power analysis was conducted to

determine minimum sample size for up to 10 focal contrasts while maintaining familywise power

$\geq$ .90; this required a per-test power of $\sqrt[10]{.90} = .9895$. The alpha level was determined by

taking the typical peak Z required for peak voxel familywise error correction (Z = 4.53),

computing its equivalent *p*-value from the normal distribution ($p = 2.95 \times 10^{-6}$), and then

Bonferroni correcting this value for 10 comparisons ($p = 2.95 \times 10^{-7}$). The simulated effect size

was determined by computing the average of the peak Z scores from all significant clusters

revealed by the [Interoception – Exteroception] in Farb et al. (2013), as were the variances of the

contrast main effect and the participant random effects (Z = 5.87, $s^2_{contrasteffect} = .20$, $s^2_{randomeffect} = $

.22). Monte Carlo simulation conducted in the R statistical environment suggested that up to 10

contrasts would be sufficiently powered at N $\geq$ 9 (Figure S1), so the parent trial sample size of N

= 44 scans adequately powered the study design.

Twenty-five healthy individuals with self-reported elevated stress were initially recruited

through advertisements in a local newspaper and through the University of Washington (UW)

research volunteer website and flyers posted on the UW campus. Inclusion criteria were: 1)

being over 18 years of age; 2) Perceived Stress Scale (Taylor, 2015) scores indicating moderate

stress levels; 3) naive to mindfulness-based approaches (no prior experience), 4) agrees to forgo

(non-study) manual therapies (e.g., massage) and mind-body therapies (e.g., mindfulness

meditation) for 12 weeks (baseline to post-test); 5) fluent in English; 6) can attend MABT and assessment sessions; and 7) right-handed (for uniformity of neuroimaging results). Exclusion criteria were: 1) lifetime diagnosis of mental health disorder; 2) unable to complete study participation (including planned relocation, pending inpatient treatment, planned extensive surgical procedures, etc.); 3) cognitive impairment, assessed by the Mini-Mental Status Exam (MMSE) (Folstein et al., 1975) if demonstrated difficulty comprehending the consent; 4) use of medications in the past 30 days that affect hemodynamic response; 5) lifetime head injuries or loss of consciousness longer than 5 minutes; 6) currently pregnant; or 7) contraindications for MRI, e.g., claustrophobia, metal objects in body, etc. The full CONSORT diagram is available in the supplementary materials (Figure S2).

Ethics Statement

All participants provided informed consent. The study procedures were reviewed and approved by the institutional review board at the University of Washington in accord with the World Medical Association Declaration of Helsinki.

The Interoceptive/Exteroceptive Attention Task (IEAT)

The Interoceptive/Exteroceptive Attention Task (IEAT) is a novel paradigm for exploring the neural dynamics of respiratory attention and awareness. The IEAT consisted of five conditions: Passive Exteroception, Passive Interoception, Active Interoception, Active Exteroception, and Active Matching (a paced breathing condition), as shown in Figure 1.

Figure 1. Schematics for the Interoceptive/Exteroceptive Attention Task (IEAT). In Exteroceptive conditions, participants attended to a circle expanding and contracting; in Interoceptive conditions, participants attended to their respiratory inhalation and exhalation. In Passive conditions, participants simply observed the circle or their breath; in Active conditions, participants pressed buttons to track the circle or breath. In the Matching condition, participants tracked the circle's movements while synchronizing their respiration cycle with the circle.

Passive Conditions. During Passive Exteroception, participants were asked to visually monitor a circle as it expanded and contracted periodically on the MRI-compatible visual display without making any behavioral responses. The circle's pulse frequency was set to match participants' in-scanner breathing frequency, obtained from a respiration belt worn by participants in the scanner (usually around 12 Hz). During Passive Interoception, participants viewed a stationary circle on the screen while attending to sensations of the breath.

Active Conditions. During Active Exteroception, participants reported on the expansion and contraction of the circle on the screen, which again was set to pulse at participants' in-scanner respiratory frequency. During Active Interoception, participants reported on inhalations and exhalations by making button box key presses with their right-hand index and middle fingers respectively. The circle on the screen expanded and contracted with these key presses,

approximating the frequency of circle movement during Passive Exteroception. During Active

Matching, participants reported on the expansion and contraction of the circle as in Active

Exteroception, while synchronizing their inhalation to the circle's expansion and their exhalation

to the circle's contraction.

Self-Reported Interoceptive Awareness

Perhaps the most common and general (i.e., not task-specific) index of subjective

interoceptive engagement is the Multidimensional Assessment of Interoceptive Awareness

(MAIA), which in validation has demonstrated strong associations to subjective wellbeing

(Mehling et al., 2012, 2018). The MAIA is a broad and general measure that examines multiple

indicators of subjective interoceptive attention including awareness, comfort, skill at sustaining

attention, and trust. The MAIA is sensitive to treatment effects from clinically efficacious

interoceptive-focused interventions, with studies demonstrating a positive relationship between

improved interoceptive awareness on the MAIA and treatment health outcomes (Fissler et al.,

2016; Price et al., 2019; Roberts et al., 2022).

Here, the MAIA was used to provide a subjective report of the ability to adaptively engage

interoceptive attention. The MAIA is a 32-item scale designed to measure multidimensional

facets of self-reported interoceptive awareness relevant for wellbeing. The eight subscales canvas

experiential domains such as Noticing, Listening, and Emotional Awareness, as well as

regulatory domains such as Attention Regulation, Self-Regulation, and Trusting. As noted above,

the MAIA has an updated version to improve the reliability of the two reverse-coded subscales

(Mehling et al., 2018), but as the new version was not available at study outset, the total score

from the original version was employed, which yielded very good reliability (Cronbach's α =

.85) in the present sample.

Procedures

Participants completed the IEAT during fMRI acquisition at two timepoints, baseline and 3-month follow-up. The MAIA self-report questionnaire was administered to assess interoceptive awareness at both timepoints. Training effects are the subject of separate reports.

Data Analysis

Respiration Confounds. Changes in breathing depth and rate modulate CO_2 concentration in the bloodstream, with faster, shallower breathing increasing blood oxygen level dependent (BOLD) signals throughout the brain (Birn et al., 2006). Failure to correct for task-related variation in breathing rate and depth can therefore lead to spurious associations with experimental conditions (Weiss et al., 2020). Consensus on correction for such effects is an area of ongoing investigation, but fMRI analysis should at a minimum correct for respiratory frequency and volume / time (Power et al., 2020).

Respiration Frequency. Respiration data was acquired using a MR-compatible respiration belt sampling at 500 Hz. Respiration data was first smoothed using a 1-sec zero phase low-pass filter window and then mean-corrected. Breath frequency was then estimated using a Fast Fourier Transform (FFT) of the respiration period. As a first level covariate, frequency was estimated using a 10-sec sliding window across the time series, generating a frequency value for each volume acquired in the timeseries. Trial-specific frequencies were also estimated across each task period to serve as covariates at the second (group) level of analysis.

Respiration Volume / Time. The respiratory signal is influenced by changes in respiratory volume in addition to frequency, with Respiratory Volume per Time (RVT) predicting widespread changes in BOLD activity (Birn et al., 2006). To correct for this influence, RVT was calculated as recently recommend (Power et al., 2020) by first finding peaks and troughs of the

normalized (z-scored) respiratory signal, setting a minimum distance between peak and trough of

0.5 standard deviations, but a minimum distance of 2 seconds between consecutive peaks or

troughs. Linear interpolation of peaks and troughs generated an 'envelope' around the respiratory

signal, and the difference between peak and trough values represented the RVT score across the

timeseries. Trial-averaged RVTs were also estimated for each task period for use as covariates at

the second (group) level of analysis.

Stimulus and Behavior Timeseries. The three active localizer conditions, Active

Exteroception, Active Interoception, and Active Matching required button-presses to track the

sensory target, i.e., inhalation/exhalation during the respiratory cycle, or expansion/contraction

during the visual circle cycle. Each active condition therefore produced three periodic timeseries:

(i) the circle radius, as measured by timestamps in the participant log files, (ii) respiration

volume, as measured by pressure transduction on an MR-compatible respiration belt, and (iii) a

waveform generated from participant button presses, with button presses indicating inflection

points (peaks and troughs) of the response waveform.

The respiration belt waveform was obtained directly from a physiological logfile for each

fMRI run. Respiration data was sampled at 500 Hz and included a scanner-generated timestamp

to precisely indicate the beginning and end of the functional run. Respiration data was segmented

into task trials using timestamps from participant logfiles. The 'pracma' library (Borchers, 2021)

in the R statistical programming environment (R Core Team, 2017) was used to find peaks and

troughs within the timeseries (Figure 2A below). The minimum peak difference was set to the

sampling rate, as individual breaths were unlikely to have a period shorter than 1 second.

Behavioral keypress waveforms were generated using linear interpolation of behavioral

logfile timestamps for button presses using the *approx* function in R. The inhalation/expansion

button "1" was coded as −1 to indicate beginning from the trough of the period, and the

exhalation/contraction button "2" was coded as 1 to indicate peak of the period. As the

waveforms were intended to be compared to the respiratory signal, the waveforms were

interpolated at a sampling rate of 500 Hz to match the data acquired by the respiration belt.

The circle stimulus waveform was obtained from the stimulus presentation software. The

circle period was matched to participant-specific breathing frequencies obtained via respiration

belt during structural MRI acquisition. For each trial, the timeseries began at an arbitrary unit of

0. The expansion/contraction period was symmetrical and defined as half the participant

respiratory period. The circle value was then extrapolated as increasing from −1 for the

expansion period to reach a value of 1 by the end of the period; the circle then paused for 0.2 *

the expansion period; the circle then decreased at the same rate for the contraction period to

reach a value of −1 by the end of the period, and then paused again for 0.2 * the contraction

period, repeating the cycle the end of the task block.

Tracking Accuracy. To calculate trial-specific tracking accuracy in the active tracking

conditions (Active Interoception, Active Exteroception, and Active Matching), the appropriate

sensory stimulus waveform (respiration in Active Interoception, the visual circle in Active

Exteroception) was first phase-corrected to maximally align with the keypress waveform; this

correction served to compensate for systematic signal transduction lags between the three types

of signals. Phase correction was performed using a normalized cross-correlation (NCC) analysis

using the 'dtwclust' library in R to estimate the maximum correlation between behavior and

respiration/circle timeseries (Sardá-Espinosa, 2019). The NCC analysis tested correlation as

function of lag between two timeseries, up to a full second (500 samples) of lag.

Following phase correction, tracking accuracy was assessed by computing the deviation from the respiration/circle inflection points to the nearest keypress, as tracking keypresses were cued by changes in circle/respiration phase (the switch from expansion/inhalation to contraction/exhalation). The average deviation in ms was then calculated for each trial and used as a measure of tracking accuracy, with lower scores indicating more accurate tracking (Figure 2D below). The complete algorithm for generating timeseries waveforms and calculating tracking accuracy is available on the Open Science Framework (https://osf.io/cuz6s).

Neuroimaging Data Acquisition. Neuroimaging was performed using a 3T Philips Achieva scanner (Philips Inc., Amsterdam, Netherlands) at the Diagnostic Imaging Sciences Center, University of Washington. Imaging began with the acquisition of a T1-weighted anatomical scan (MPRAGE) to guide normalization of functional images (~ 6 minutes) with TR = 7.60 ms, TE = 3.52 ms, TI = 1100 ms, acquisition matrix = 256 × 256, flip angle = 7°, shot interval = 2530 ms, and 1mm isotropic voxel size. Functional data were acquired using a T2∗-weighted echo-planar-imaging (EPI) sequence with TR = 2000, TE = 25 ms, flip angle α = 79°, field of view = 240 × 240 × 129 mm, 33 slices, and a voxel size of 3 × 3 × 3.3 mm with 3.3 mm gap. Button presses were registered using a 2-button MR-compatible response pad.

Preprocessing. A set of preprocessing steps was performed using the consortium-developed fMRIprep robust preprocessing pipeline for fMRI data (https://fmriprep.readthedocs.io/en/stable/). Briefly, preprocessing consisted of realignment and unwarping of functional images, slice timing correction, and motion correction. The functional images were resliced using a voxel size of 2 × 2 × 2 mm and smoothed using a 6-mm FWHM isotropic Gaussian kernel.

First Level Analysis. Within subject statistical models were used to characterize the interoceptive networks at the participant level. Participant time series from the interoceptive tasks were submitted to separate first-level general linear statistical models using Statistical Parametric Mapping software (v12). Task-specific boxcar stimulus functions were convolved with the canonical hemodynamic response function, separately modeling the onsets of the interoceptive and visual control conditions for each participant. To control for motion and physiological confounds, six standard movement parameters, the root mean square of temporal change (DVARS), framewise displacement, respiration rate, and RVT were all included as nuisance covariates.

Second Level Analysis. Participant first-level maps for each experimental condition [Passive Interoception, Passive Exteroception, Active Interoception, Active Exteroception, Active Matching] were analyzed at the second level using a full-factorial mixed-model ANOVA in SPM12 (Friston, 2007). The second level contrasts first subdivided the tasks in two main effects and their interaction term: Reporting Demand [Active vs. Passive] × Target [Interoception vs. Exteroception]. Follow up comparisons within Active [Active Interoception vs. Active Exteroception vs. Active Matching] and within Passive [Passive Interoception vs. Passive Exteroception] were also modelled.

Familywise control for multiple comparisons (corrected $p < 0.05$) in whole-brain analyses was achieved through threshold-free cluster enhancement (TFCE), which controls familywise error rate based on a permutation testing approach and determines optimal voxel-wise cluster-forming thresholds using an automated algorithmic method (Smith & Nichols, 2009). The algorithm eliminates the need to choose between arbitrary correction thresholds by sampling

across statistically equivalent peak and cluster size thresholds. Trial-averaged respiration

frequency and RVT were modelled at the second level as nuisance covariates.

Trial-level Confounds. The current study comes from an exploratory clinical trial, the

results of which are the subject of a separate report. We combined data across the trial to power

the comparison of IEAT task conditions and modelled any effects of trial Group (MABT vs.

Control), Time (Baseline vs. Post-Intervention) and their interactions as nuisance covariates. All

models also contained condition-averaged respiration rate as a covariate, to further control for

variation in respiration between experimental conditions. Post hoc analyses that did not include

the nuisance covariates of Group and Time did not meaningfully change the reported results.

Region of Interest (ROI) Analysis. For region of interest (ROI) analysis, all signal

extractions were taken from models containing the nuisance covariates. Using the built-in

SPM12 function, the median value of the raw, unwhitened signal was extracted from all voxels

within the ROI, yielding one value per participant at each time point. These values were entered

into a linear mixed-effects model with restricted likelihood estimation was applied using the

'lme4' library (Bates et al., 2015) in the R statistical programming environment.

Hypothesis Testing. Hypothesis 1 aimed to compare interoceptive and exteroceptive

attention. To this end we first evaluated a whole-brain interaction between reporting demand

[active vs. passive] and attentional target [interoception vs. exteroception] to evaluate whether

the effects of attentional target should be evaluated separately for the active and passive

reporting conditions. Subsequent analyses compared the simple effects of attentional target

within each reporting demand condition, i.e. [Passive Interoception vs. Passive Exteroception]

and [Active Exteroception vs. Active Interoception]. To perform these contrasts, all 5 task

conditions were estimated separately at the first (individual session) level of analysis and entered into a full factorial design in SPM12.

Hypothesis 2 aimed to investigate whether the differences between exteroception and interoception were moderated by individual differences subjective interoceptive awareness (MAIA scores). Focusing on the contrast of [Active Interoception vs. Active Exteroception], we first created contrast maps at the first (within session) level of analysis. These first level maps were then entered into a second (group) level analysis that included normalized (z-scored) MAIA scale total scores as a covariate of interest. The MAIA covariate was subjected to TFCE correction in the same fashion as other whole brain analyses, and respiration rate change between the two conditions was also included in the factorial model as a nuisance covariate to control for variation associated with changes in respiratory rate.

Hypothesis 3 tested the potential moderating factor of endogenous vs. exogenous control of the respiratory cycle by contrasting each of Active Exteroception and Active Interoception against Active Matching using the same five condition model from testing Hypothesis 1.

Hypothesis 4 aimed to understand how engaging in Active Interoception changes brain connectivity relative to Active Exteroception. To accomplish this aim, a psychophysiological interaction analysis (PPI) was conducted in SPM12 using the Generalized PPI Toolbox (v. 13.1), which improves upon standard PPI analyses by estimating the effect each task condition has upon connectivity independently (McLaren et al., 2012). Here, we used the model employed in Hypotheses 2, a second (group) level full factorial model that contained first level contrasts of [Active Interoception – Active Exteroception] and the MAIA covariate term.

To define a region of interest (ROI), the conjunction of the [Active Exteroception - Active Interoception] contrast and the positive MAIA contrast was evaluated, using a threshold of $p <$

.001 for each contrast, resulting in a conjoint probability comparable to conservative FWE

correction of $p < 1 \times 10^{-6}$. The largest cluster observed was used as a seed region, K = 794

voxels, peak Z = 4.06, x = −4, y = 58, z = 14, consistent with dorsal anterior cingulate cortex

(Brodmann area 24). At the first level of analysis, mean timecourse activity extracted from this

seed region was convolved with separate boxcar regressors indicating the onsets and durations of

the Active Exteroception and Active Interoception conditions, and the ensuing whole brain maps

were then contrasted to model the PPI effect for each participant session. These first level PPI

maps were then collected and analyzed using the same full factorial modelling approach

described above.

Test-Retest Reliability. The fact that each participant was scanned twice offered a unique

opportunity to test the reliability of study effects across two independent scanning sessions. A

conjunction analysis for the four main contrasts reported in this paper were conducted. As the

TFCE algorithm does not currently perform conjunction analysis, and halving the sample size

reduces experimental power, the baseline and post-intervention sessions were each analyzed

separately at $p < .001^{1/2} = p < .0316$, so that the resulting overlap would yield a conjunction $p <$

.001; a minimum cluster size of k ≥ 500 was also applied to each estimate as this cluster size

resulted in a FWE corrected p value < .05.

Data and Code Availability

The full study protocol was pre-registered with the Open Science Framework

(https://osf.io/y34ja). All study materials, including behavioral data and fMRI signal extractions,

the code to run the experiment and subsequent fMRI analysis, power analysis, statistics, and

graphing scripts, are freely available on the Open Science Framework (https://osf.io/ctqrh/).

Results

Control Analyses / Manipulation Checks

Figure 2. Respiration and Behavior Tracking. **(A)** Respiratory peak and trough detection was successful for each of the experimental task conditions, allowing for calculation of respiratory frequency (Hz) and respiratory volume over time (RVT). **(B)** Respiration was slower during Active Interoception than in other conditions. **(C)** Respiration was deeper (greater volume / time) during Active Interoception and Active Matching than in other conditions. **(D)** To assess tracking accuracy, the respiration waveform (blue) was first temporally adjusted (black) to match phase with keypress signals (orange). The average error (red) between key press timing and respiratory peaks/troughs was calculated; the same procedure was applied to a waveform describing circle radius in the exteroceptive tracking conditions. **(E)** No differences in tracking accuracy were observed between Active Interoception of the Breath and Active Exteroception of the Circle, nor with but Active Matching to the Circle; however, breath tracking was superior during the Active Matching condition.

Figure 2 summarizes the analysis of the control variables: respiratory frequency, volume/time

(RVT), and tracking error. The peak and trough detection algorithm (Figure 2A) allowed for

RVT estimation (see Supplementary Materials for similar images of every run in the study).

Effects of Attention on Respiration Rate and Respiratory Volume / Time (RVT)

The study-wide average respiration frequency was .21 Hz (Table S1). A main effect of task

condition was observed, $F(4,854) = 9.36$, $p < .001$, with follow-up pairwise comparisons

suggesting that respiration rate was slower during Interoception conditions than Exteroception

conditions, with an average reduction of .02 Hz, 95% CI [.01, .03] (Figure 2B). However,

respiration rate was not slower during than Exteroception conditions in the Active Matching

condition (Table S2).

Condition had a large impact on RVT, $F(4,854) = 109.7$, $p < .001$ with equivalent

respiratory volumes only between the two exteroception conditions (Table S3). Follow-up

analyses confirmed that participants breathed more deeply during Passive Interoception than

Exteroception, even more deeply during Active Matching, and deepest during Active

Interoception (Figure 2C, Table S4).

A frequency / volume tradeoff was also evident, with frequency and RVT negatively

correlated within-participant, $r(878) = -.452$, 95% CI [-.535, -.434], $p < .001$, which may offset

the impact of changing respiration rates on BOLD activity. Nevertheless, to control for the

influence of condition on respiration, block-specific respiration frequency and RVT were

included as covariates in all second level analyses. Controlling for frequency made had little

impact on BOLD activation in subsequent analysis, but greater RVT was associated with

widespread cortical deactivation (Figure S3), reinforcing the importance of controlling for RVT

during data analysis.

Interoceptive and Exteroceptive Tracking Accuracy

Participant tracking of the sensory stimuli (respiration and visual circle) was analyzed to in

terms of error (in milliseconds) between key presses and the peaks and troughs of stimulus

waveforms within each of the active task blocks. Stimulus waveforms were first phase-corrected

to maximally match peaks and troughs with keypresses. Error was then measured as the average

time difference between stimulus peaks/troughs and key presses (Figure 2D).

Participants accurately and reliably tracked both the respiratory and visual stimuli (Figure

2E, Table S5) with no error differences between breath tracking in Active Interoception, circle

tracking in Active Exteroception, or circle tracking in Active Matching (Table S6). Respiration

alignment with keypad responses was superior to all other conditions in Active Matching. These

results suggest the IEAT successfully matched tracking difficulty between the Active conditions.

H1: Effects of Attentional and Reporting Demand

A whole-brain interaction analysis between Reporting Demand [Active vs. Passive] and Attentional Target [Exteroception vs. Interoception] implicated the sensorimotor, temporoparietal, and prefrontal cortex (Figure 3A); follow-up analysis of the median signal across all significant voxels revealed deactivation in the Active Interoception and Active Matching conditions relative to the other task conditions (Figure 3B; Table S7).

Figure 3. Interactions and Simple Effects of Task **(A)** Interoceptive Reporting Demand [Active vs. Passive] x Attentional Target [Exteroception vs. Interoception] implicated the medial and lateral prefrontal cortices (MPFC and LPFC), somatomotor cortices, striatum, and temporal/occipital cortices. **(B)** Median % signal change across the regions revealed highest activation in the Active Exteroception condition and the two Passive conditions relative to Active Interoception and Act Match. **(C)** Simple effects comparisons between Active Exteroception and Interoception conditions largely replicated the interaction effect, suggesting that the Active conditions were driving the interaction. **(D)** Conversely, only area V5/MT distinguished Passive Exteroception from Passive Interoception.

A follow-up simple effects contrast of [Active Exteroception > Active Interoception] largely replicated the interaction effect (Figure 3C); conversely, the contrast of [Passive Exteroception > Passive Interoception] revealed only a small cluster of activation in motion-related lateral occipital area V5/MT (Figure 3D). The V5/MT finding is consistent with a failure to fully match task features, as Passive Interoception was the only condition in which the circle stimulus remained stationary rather than pulsing. Given a lack of other distinctions between the passive monitoring conditions, subsequent analyses focused upon the active tracking conditions.

H2: Covariates of Self-Reported Interoceptive Awareness

To better understand the nature of the deactivation observed during Active Interoception relative to Active Exteroception, a focal analysis was conducted to investigate the potential moderating role of self-reported interoceptive awareness (MAIA scores). Interoceptive awareness was significantly related to the level of deactivation observed across a subset of the cortical regions implicated in interoception-related deactivation (Figure 4). Specifically, activity in the anterior cingulate cortex (ACC), dorsomedial prefrontal cortex, and left lateralized language-related regions demonstrated significant associations with interoceptive awareness, such that greater MAIA scores were associated with *reduced deactivation* during Active Interoception relative to Active Exteroception (Table S8). This pattern was also more generally replicated across both passive and Active Matching conditions (Figure S4).

Figure 4. Covariates of Self-Reported Interoceptive Awareness. **(A)** Orange regions denote positive correlates of MAIA scores within the [ActExt – ActInt] contrast. **(B)** Scatterplot of the relationship between MAIA score and median neural activity during the [ActExt – ActInt] contrast across significant covariate regions.

As the largest and most powerfully moderated cluster, the ACC region, k = 1205, x = -4; y = 12; z = 36, was retained as a seed region of interest (ROI) in the PPI analysis (H4) described below.

H3: Endogenous vs. Exogenous Sources of Interoceptive Control

The next planned comparison contrasted Active Matching against Active Interoception to explore endogenous and exogenous sources of respiratory control. To contextualize Active Matching, we first compared it to Active Exteroception. Active Matching showed even more pronounced and widespread patterns of deactivation than Active Interoception (Figure 3C vs. Figure 5A). Compared to the endogenously paced Active Interoception condition, the exogenously-paced Active Matching condition led to greater deactivation bilaterally along an insula/operculum pathway, somatomotor regions, and within the ventral occipital cortex and cerebellum (Figure 5, Table S9).

Figure 5. Endogenous vs. Exogenous Sources of Interoceptive Control. **(A)** Active Exteroception > Active Matching replicated the deactivation effect observed for Active Interoception. **(B)** Median % signal change across the regions identified via the [Active Exteroception > Active Matching] contrast. **(C)** Active Interoception > Active Matching linked endogenous respiration to greater activation along an insula-operculum pathway relative to the exogenous control condition. **(D)** Median % signal change across the regions identified via the [Active Interoception > Active Matching] contrast.

H4: Psychophysiological Interaction (PPI) Analysis

The final contrast examined changes in functional connectivity between Active

Exteroception and Active Interoception. The anterior cingulate cortex (ACC) region of interest

(Figures 4A & 6A) was entered as a seed region in a psychophysiological interaction (PPI)

analysis. The PPI analysis explored changes in functional connectivity with the ACC (Figure 7C)

as a function of the two experimental conditions (Active Interoception vs Active Exteroception).

The analysis revealed a strong integration of the ACC into regions consistent with the DAN

during Active Interoception (Figure 6B; Table S10).

Figure 6. Psychophysiological Interaction (PPI) Analysis. **(A)** The ACC seed region obtained
from the MAIA covariate analysis. **(B)** PPI between ACC regional activity and task conditions
[Active Interoception > Active Exteroception], demonstrating enhanced ACC connectivity with a
frontoparietal network during Active Interoception relative to Active Exteroception.

Test-Retest Reliability

A conjunction analysis of the main H1-H4 contrasts estimated separately at each of the two

timepoints successfully reproduced the results described above (Figure S5).

Discussion

Relative to visual Exteroception, respiratory Interoception was characterized by pervasive neural deactivation in prefrontal, somatomotor, and temporoparietal regions. This finding deviates from previous respiratory interoception studies (Farb et al., 2013b; Wang et al., 2019), where more difficult sensory tasks yielded greater DAN activity relative to greater DMN activity for less difficult tasks. The finding also differs from liminal heartbeat detection paradigms (c.f., Ring & Brener, 2018), where prior research has related interoceptive accuracy to greater activity and connectivity within the SLN (Chong et al., 2017; Critchley et al., 2004; Tan et al., 2018). The lack of SLN involvement in closely matched, supraliminal tasks supports recent arguments that SLN engagement generally indicates perceptual decision-making rather interoception specifically (Baltazar et al., 2021; Koeppel et al., 2020).

Slower and deeper breathing during the Active Interoception condition could suggest that neural activation differences may be predicated on gross physiological changes driven by different respiration rates, volumes, and spontaneous end-tidal CO_2 (PETCO$_2$; Golestani et al., 2015). However, modelling RVT successfully accounted for pervasive deactivation effects (Figure S3), and controlling for respiration rate and RVT at both the first (acquisition volume) and second (task block) levels of analysis failed to eliminate the Interoception deactivation effect. Furthermore, the rate of respiratory slowing (~0.5 Hz) was too small to provoke hypocapnia, and while CO_2 was not measured in this study, convergent findings suggest that controlling for PETCO$_2$ reduces functional connectivity but not task amplitude (Golestani & Chen, 2020; Madjar et al., 2012). Finally, the Active Matching task, in which respiration frequency and yoking of behavior were both matched to Active Exteroception, still resulted in this deactivation pattern. While there must certainly be interactions between attention and

physiology driving the deactivation effect, they are unlikely to be explained by gross changes in

blood oxygenation or CO_2.

The right anterior insula, commonly implicated in investigations of interoceptive accuracy

(Critchley et al., 2004; Haruki & Ogawa, 2021; Wang et al., 2019), was insensitive to contrasts

between actively-reported Interoception and Exteroception. However, the insula was implicated

in contrasts between Active Interoception and Active Matching, where moving from endogenous

to exogenous respiratory pacing resulted in greater insula deactivation. This effect is not simply

due to the integration of external cues, as this deactivation effect was absent in Active

Exteroception. Endogenous respiratory rhythms may be continuously processed by this

insula/operculum pathway, revealed only when such processing is disrupted by task demands.

Here, exogenous control over the respiratory cycle may have disrupted sensory integration or

signals such as the body's internal respiratory 'clock', a rhythm commonly attributed to the pre-

Bötzinger complex (Smith et al., 1991), but which may communicate with the insula to support

homeostasis (Yang & Feldman, 2018). By this logic, heartbeat detection accuracy paradigms

implicate the insula precisely because they are comparing participants with robust heartbeat

representations to those where such processing is compromised, disrupted, or absent.

Conversely, attention to salient signals such as the breath are unlikely to implicate the insula

unless compared to situations where breath awareness is somehow disrupted or compromised.

The Moderating Influence of Interoceptive Sensibility

A second aim of the study was to explore the potential moderating effects of interoceptive

sensibility (MAIA scores) on interoceptive network engagement, with the hypothesis that greater

subjective interoceptive awareness would be supported by increased SLN activity in the ACC

and anterior insula, commensurate with their established role in supporting interoceptive

accuracy (Chong et al., 2017; Critchley et al., 2004; Harrison, Köchli, et al., 2021).

Greater activity within the SLN was observed in participants who reported greater

interoceptive sensibility. Exactly how the allocation of attention serves to reduce brain activity

without impairing performance remains an intriguing question. Performance may be sustained

because of noise inhibition along interoceptive pathways (c.f., Kuehn et al., 2016), which could

offset the disadvantage seemingly implied by widespread cortical inhibition. This moderating

effect is consistent with the 'low-energy state' theory proposed above, which characterizes

interoceptive attention as a state of preserved attentional monitoring in a less noisy neural

environment, especially given findings of increased SLN connectivity during interoceptive

attention relative to exteroceptive attention.

Characterizing the Network Supporting Interoceptive Awareness

A final aim of the study was to characterize an interoceptive attention network supporting

adaptive interoceptive sensibility. The rostral ACC was identified as a seed region for

psychophysiological interaction (PPI) analysis, convolving the activity in the region implicated

as a MAIA covariate with the Active Interoception vs. Active Exteroception contrast.

Interoception led to increased connectivity in the frontoparietal dorsal attention network

(DAN; Spreng et al., 2013; Szczepanski et al., 2013). While DAN *activity* was inhibited during

interoception, those with greater self-reported interoceptive awareness were likely to offset such

deactivation by greater *connectivity* between the ACC and DAN, a potential biomarker of

interoceptive engagement.

This finding helps to explain how interoceptive tracking accuracy was equivalent to

exteroceptive tracking accuracy despite reduced cortical activity. Interoceptive processing may

be continuous and automatic, but largely obscured by a combination of external sensory signals and internal cognitive processing. Interoceptive attention may therefore employ 'addition by subtraction', a reduction of competing neural representations rather than the activation of a dormant interoceptive pathway.

The idea of an 'always on' interoceptive state is consistent with contemporary theories that place interoception as the background of consciousness, such as Damasio's "Proto Self" (Bosse et al., 2008) or "Core Consciousness" (Parvizi, 2001), and the distinction between a subcortical "Core Self" and higher order representations in the cerebral cortex (Northoff & Panksepp, 2008). Interoception is likely the first sense represented in developing brains because of the critical roles homeostasis plays in ensuring survival (Filippetti, 2021; Fotopoulou & Tsakiris, 2017). The consistent presence of periodic interoceptive signals may lead to neural habituation, in keeping with neural repetition suppression effects for familiar and repeatedly represented sensory signals (Barron et al., 2016; Summerfield et al., 2008). Relative to more varied exteroceptive signals, lifelong habituation to omnipresent interoceptive signals may present as widespread cortical deactivation.

Limitations and Constraints on Generalizability

The IEAT should be improved by introducing a circle motion to the Passive Interoception condition, as the stationary target described here led to a motion confound between passive interoception and the other four experimental conditions. In its current form, Passive Interoception showed relative deactivation in the middle temporal (MT or V5) region of the visual cortex, which is well-established for its sensitivity to motion (Albright & Stoner, 1995).

Several constraints on generalizability are also apparent. We present unexpected and exploratory findings from a relatively small sample; replication is therefore needed, especially

given the paucity of studies in this area. The community sample may also not generalize to

people with clinical conditions associated with interoceptive dysfunction, nor advanced

contemplative practitioners with extensive interoceptive training. Respiration is also only one of

many interoceptive signals that each vary in their reportability and controllability, and so may

possess distinct neural dynamics. Multimodal interoceptive research is needed to determine the

generalizability of the patterns discussed here, to clarify the impact of active reporting vs.

passive engagement, the role sensory signal salience, and the influence of participant

expertise/dysfunction.

A further limitation to the present findings is that all practices were conducted with eyes-

open, which may be dissimilar to many meditation practices. Following repeated evidence that

opening one's eyes decreases the functional connectivity between the SLN and the DMN, it was

theorized that reducing the connectivity between these networks reflects an orientation to

external rather than internal events (Han et al., 2020). However, here interoceptive attention with

eyes open led to greater connectivity between the ACC seed region (an efferent hub of the SLN)

and both the DAN and more posterior aspects of the insula rather than the DMN. It remains

possible that dynamics for interoceptive awareness may be altered in eyes-closed paradigms.

Beyond simply eyes being closed, increased DMN coupling may represent a distinct form of

internal awareness at the elaborative, semantic level of processing distinct from interoceptive

awareness, as has been previously proposed (Andrews-Hanna et al., 2014; Gusnard et al., 2001).

Concluding Remarks

Relative to Exteroception, Interoception reduced somatomotor and prefrontal *activity*,

offset by enhanced *connectivity* between the SLN and the DAN. Greater interoceptive sensibility

was linked to sparing of the SLN and language processing regions from this deactivation pattern.

Together, these processes may allow sensory signals from the body to be better discerned. The

ACC deactivation observed in the IEAT paradigm may serve as a candidate biomarker of

individual differences in interoceptive processing, one that distinguishes adaptive representations

engendered by contemplative training from the avoidance and catastrophizing observed in

anxiety and somatic disorders. Larger, more diverse samples and replication of effects are

needed to test these emerging ideas.

Acknowledgements

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. We extend our gratitude to Natalie Koh and Sophie Xie for their coordination and data collection efforts, and to Dr. Thomas Grabowski, Dr. Chris Gatenby, and Ms. Liza Young at the Integrated Brain Research Center at the University of Washington for their collaboration. This work was supported by a RIFP award from the School of Nursing at the University of Washington and a grant from the National Center for Advancing Translational Sciences of the National Institutes of Health [UL1 TR002319]. Data analysis infrastructure and trainee salary was supported by a Canadian Natural Sciences and Engineering Council Discovery grant [RGPIN-2015-05901].

References

Albright, T. D., & Stoner, G. R. (1995). Visual motion perception. *Proceedings of the National Academy of Sciences*, *92*(7), 2433–2440. https://doi.org/10.1073/pnas.92.7.2433

Andrews-Hanna, J. R., Smallwood, J., & Spreng, R. N. (2014). The default network and self-generated thought: Component processes, dynamic control, and clinical relevance. *Annals of the New York Academy of Sciences*, *1316*, 29–52. https://doi.org/10.1111/nyas.12360

Anestis, M. D., Selby, E. A., Fink, E. L., & Joiner, T. E. (2007). The multifaceted role of distress tolerance in dysregulated eating behaviors. *International Journal of Eating Disorders*, *40*(8), 718–726. https://doi.org/10.1002/eat.20471

Avery, J. A., Drevets, W. C., Moseman, S. E., Bodurka, J., Barcalow, J. C., & Simmons, W. K. (2014). Major Depressive Disorder Is Associated With Abnormal Interoceptive Activity and Functional Connectivity in the Insula. *Biological Psychiatry*, *76*(3), 258–266. https://doi.org/10.1016/j.biopsych.2013.11.027

Baltazar, M., Grezes, J., Geoffray, M., Picq, J., & Conty, L. (2021). Neural correlates of interoceptive accuracy: Beyond cardioception. *European Journal of Neuroscience*, *54*(10), 7642–7653. https://doi.org/10.1111/ejn.15510

Barrett, L. F. (2017). *How emotions are made: The secret life of the brain*. Houghton Mifflin Harcourt.

Barron, H. C., Garvert, M. M., & Behrens, T. E. J. (2016). Repetition suppression: A means to index neural representations using BOLD? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1705), 20150355. https://doi.org/10.1098/rstb.2015.0355

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

    using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

    https://doi.org/10.18637/jss.v067.i01

Birn, R. M., Diamond, J. B., Smith, M. A., & Bandettini, P. A. (2006). Separating respiratory-

    variation-related fluctuations from neuronal-activity-related fluctuations in fMRI.

    Neuroimage, *31*(4), 1536–1548. https://doi.org/10.1016/j.neuroimage.2006.02.048

Borchers, H. W. (2021). *pracma: Practical Numerical Math Functions* (R package   version

    2.3.6). https://CRAN.R-project.org/package=pracma

Bosse, T., Jonker, C. M., & Treur, J. (2008). Formalization of Damasio's theory of emotion,

    feeling and core consciousness. *Consciousness and Cognition*, *17*(1), 94–113.

    https://doi.org/10.1016/j.concog.2007.06.006

Calì, G., Ambrosini, E., Picconi, L., Mehling, W. E., & Committeri, G. (2015). Investigating the

    relationship between interoceptive accuracy, interoceptive awareness, and emotional

    susceptibility. *Frontiers in Psychology*, *6*. https://doi.org/10.3389/fpsyg.2015.01202

Chand, G. B., & Dhamala, M. (2016). The salience network dynamics in perceptual decision-

    making. *NeuroImage*, *134*, 85–93. https://doi.org/10.1016/j.neuroimage.2016.04.018

Chen, W. G., Schloesser, D., Arensdorf, A. M., Simmons, J. M., Cui, C., Valentino, R., Gnadt, J.

    W., Nielsen, L., Hillaire-Clarke, C. St., Spruance, V., Horowitz, T. S., Vallejo, Y. F., &

    Langevin, H. M. (2021). The Emerging Science of Interoception: Sensing, Integrating,

    Interpreting, and Regulating Signals within the Self. *Trends in Neurosciences*, *44*(1), 3–

    16. https://doi.org/10.1016/j.tins.2020.10.007

Chong, J. S. X., Ng, G. J. P., Lee, S. C., & Zhou, J. (2017). Salience network connectivity in the insula is associated with individual differences in interoceptive accuracy. *Brain Structure & Function*, *222*(4), 1635–1644. https://doi.org/10.1007/s00429-016-1297-7

Chong, Y. W., Kee, Y. H., & Chaturvedi, I. (2015). Effects of Brief Mindfulness Induction on Weakening Habits: Evidence from a Computer Mouse Control Task. *Mindfulness*, *6*(3), 582–588. https://doi.org/10.1007/s12671-014-0293-3

Craig, A. D. (2003). Interoception: The sense of the physiological condition of the body. *Curr Opin Neurobiol*, *13*(4), 500–505. https://doi.org/S0959438803000904 [pii]

Craig, A. D. (2009). How do you feel—Now? The anterior insula and human awareness. *Nat Rev Neurosci*, *10*(1), 59–70. https://doi.org/10.1038/nrn2555

Critchley, H. D., & Garfinkel, S. N. (2017). Interoception and emotion. *Current Opinion in Psychology*, *17*, 7–14. https://doi.org/10.1016/j.copsyc.2017.04.020

Critchley, H. D., Wiens, S., Rotshtein, P., Ohman, A., & Dolan, R. J. (2004). Neural systems supporting interoceptive awareness. *Nat Neurosci*, *7*(2), 189–195. https://doi.org/10.1038/nn1176

Desmedt, O., Van Den Houte, M., Walentynowicz, M., Dekeyser, S., Luminet, O., & Corneille, O. (2020). *A Systematic Review and Meta-analysis on the Association Between Heartbeat Counting Task Performance and Mental Disorders and Their Risk Factors Among Adults* [Preprint]. Open Science Framework. https://doi.org/10.31219/osf.io/h3by9

Domschke, K., Stevens, S., Pfleiderer, B., & Gerlach, A. L. (2010). Interoceptive sensitivity in anxiety and anxiety disorders: An overview and integration of neurobiological findings. *Clinical Psychology Review*, *30*(1), 1–11.

Ehlers, A., & Breuer, P. (1992). Increased cardiac awareness in panic disorder. *Journal of Abnormal Psychology*, *101*(3), 371.

Farb, N. A., Anderson, A. K., Mayberg, H., Bean, J., McKeon, D., & Segal, Z. V. (2010). Minding one's emotions: Mindfulness training alters the neural expression of sadness. *Emotion*, *10*(1), 25–33. https://doi.org/10.1037/a0017151

Farb, N. A. S., Daubenmier, J., Price, C. J., Gard, T., Kerr, C., Dunn, B. D., Klein, A. C., Paulus, M. P., & Mehling, W. E. (2015). Interoception, contemplative practice, and health. *Frontiers in Psychology*, *6*. https://doi.org/10.3389/fpsyg.2015.00763

Farb, N. A. S., Desormeau, P., Anderson, A. K., & Segal, Z. V. (2022). Static and treatment-responsive brain biomarkers of depression relapse vulnerability following prophylactic psychotherapy: Evidence from a randomized control trial. *NeuroImage: Clinical*, *34*, 102969. https://doi.org/10.1016/j.nicl.2022.102969

Farb, N. A., Segal, Z. V., & Anderson, A. K. (2013a). Mindfulness meditation training alters cortical representations of interoceptive attention. *Social Cognitive and Affective Neuroscience*, *8*(1), 15–26.

Farb, N. A., Segal, Z. V., & Anderson, A. K. (2013b). Attentional modulation of primary interoceptive and exteroceptive cortices. *Cereb Cortex*, *23*(1), 114–126. https://doi.org/10.1093/cercor/bhr385

Ferentzi, E., Horváth, Á., & Köteles, F. (2019). Do body-related sensations make feel us better? Subjective well-being is associated only with the subjective aspect of interoception. *Psychophysiology*, *56*(4), e13319. https://doi.org/10.1111/psyp.13319

Filippetti, M. L. (2021). Being in Tune With Your Body: The Emergence of Interoceptive

Processing Through Caregiver–Infant Feeding Interactions. *Child Development

Perspectives*, *15*(3), 182–188. https://doi.org/10.1111/cdep.12420

Fissler, M., Winnebeck, E., Schroeter, T., Gummersbach, M., Huntenburg, J. M., Gaertner, M.,

& Barnhofer, T. (2016). An Investigation of the Effects of Brief Mindfulness Training on

Self-Reported Interoceptive Awareness, the Ability to Decenter, and Their Role in the

Reduction of Depressive Symptoms. *Mindfulness*, *7*(5), 1170–1181.

https://doi.org/10.1007/s12671-016-0559-z

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state". A practical

method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*, *12*(3),

189–198. https://doi.org/0022-3956(75)90026-6 [pii]

Fotopoulou, A., & Tsakiris, M. (2017). Mentalizing homeostasis: The social origins of

interoceptive inference. *Neuropsychoanalysis*, *19*(1), 3–28.

https://doi.org/10.1080/15294145.2017.1294031

Friston, K. J. (Ed.). (2007). *Statistical parametric mapping: The analysis of funtional brain

images* (1st ed). Elsevier/Academic Press.

Garfinkel, S. N., Seth, A. K., Barrett, A. B., Suzuki, K., & Critchley, H. D. (2015). Knowing

your own heart: Distinguishing interoceptive accuracy from interoceptive awareness.

Biological Psychology, *104*, 65–74.

Gibson, J. (2019). Mindfulness, Interoception, and the Body: A Contemporary Perspective.

Frontiers in Psychology, *10*. https://doi.org/10.3389/fpsyg.2019.02012

Golestani, A. M., Chang, C., Kwinta, J. B., Khatamian, Y. B., & Jean Chen, J. (2015). Mapping

the end-tidal CO2 response function in the resting-state BOLD fMRI signal: Spatial

specificity, test-retest reliability and effect of fMRI sampling rate. *NeuroImage*, *104*, 266–277. https://doi.org/10.1016/j.neuroimage.2014.10.031

Golestani, A. M., & Chen, J. J. (2020). Controlling for the effect of arterial-CO2 fluctuations in resting-state fMRI: Comparing end-tidal CO2 clamping and retroactive CO2 correction. *NeuroImage*, *216*, 116874. https://doi.org/10.1016/j.neuroimage.2020.116874

Gusnard, D. A., Akbudak, E., Shulman, G. L., & Raichle, M. E. (2001). Medial prefrontal cortex and self-referential mental activity: Relation to a default mode of brain function. *Proc Natl Acad Sci U S A*, *98*(7), 4259–4264. https://doi.org/10.1073/pnas.071043098

Han, J., Wu, X., Wu, H., Wang, D., She, X., Xie, M., Zhang, F., Zhang, D., Zhang, X., & Qin, P. (2020). Eye-Opening Alters the Interaction Between the Salience Network and the Default-Mode Network. *Neuroscience Bulletin*, *36*(12), 1547–1551. https://doi.org/10.1007/s12264-020-00546-y

Harrison, O. K., Köchli, L., Marino, S., Luechinger, R., Hennel, F., Brand, K., Hess, A. J., Frässle, S., Iglesias, S., Vinckier, F., Petzschner, F. H., Harrison, S. J., & Stephan, K. E. (2021). Interoception of breathing and its relationship with anxiety. *Neuron*, *109*(24), 4080-4093.e8. https://doi.org/10.1016/j.neuron.2021.09.045

Haruki, Y., & Ogawa, K. (2021). Role of anatomical insular subdivisions in interoception: Interoceptive attention and accuracy have dissociable substrates. *European Journal of Neuroscience*, *53*(8), 2669–2680. https://doi.org/10.1111/ejn.15157

Hayes, S. C., Wilson, K. G., Gifford, E. V., Follette, V. M., & Strosahl, K. (1996). Experiential avoidance and behavioral disorders: A functional dimensional approach to diagnosis and treatment. *Journal of Consulting and Clinical Psychology*, *64*(6), 1152–1168. https://doi.org/10.1037/0022-006X.64.6.1152

Herrero, J. L., Khuvis, S., Yeagle, E., Cerf, M., & Mehta, A. D. (2018). Breathing above the

    brain stem: Volitional control and attentional modulation in humans. *Journal of*

    Neurophysiology, *119*(1), 145–159. https://doi.org/10.1152/jn.00551.2017

Kerr, K. L., Moseman, S. E., Avery, J. A., Bodurka, J., Zucker, N. L., & Simmons, W. K.

    (2016). Altered Insula Activity during Visceral Interoception in Weight-Restored Patients

    with Anorexia Nervosa. *Neuropsychopharmacology*, *41*(2), 521–528.

    https://doi.org/10.1038/npp.2015.174

Khalsa, S. S., Rudrauf, D., Hassanpour, M. S., Davidson, R. J., & Tranel, D. (2020). The practice

    of meditation is not associated with improved interoceptive awareness of the heartbeat.

    Psychophysiology, *57*(2). https://doi.org/10.1111/psyp.13479

Koeppel, C. J., Ruser, P., Kitzler, H., Hummel, T., & Croy, I. (2020). Interoceptive accuracy and

    its impact on neuronal responses to olfactory stimulation in the insular cortex. *Human*

    Brain Mapping, *41*(11), 2898–2908. https://doi.org/10.1002/hbm.24985

Kuehn, E., Mueller, K., Lohmann, G., & Schuetz-Bosbach, S. (2016). Interoceptive awareness

    changes the posterior insula functional connectivity profile. *Brain Structure and*

    Function, *221*(3), 1555–1571. https://doi.org/10.1007/s00429-015-0989-8

Madjar, C., Gauthier, C. J., Bellec, P., Birn, R. M., Brooks, J. C. W., & Hoge, R. D. (2012).

    Task-related BOLD responses and resting-state functional connectivity during

    physiological clamping of end-tidal CO2. *NeuroImage*, *61*(1), 41–49.

    https://doi.org/10.1016/j.neuroimage.2012.02.080

McLaren, D. G., Ries, M. L., Xu, G., & Johnson, S. C. (2012). A generalized form of context-

    dependent psychophysiological interactions (gPPI): A comparison to standard

approaches. *NeuroImage*, *61*(4), 1277–1286.

https://doi.org/10.1016/j.neuroimage.2012.03.068

Mehling, W. E., Acree, M., Stewart, A., Silas, J., & Jones, A. (2018). The Multidimensional

Assessment of Interoceptive Awareness, Version 2 (MAIA-2). *PLOS ONE*, *13*(12),

e0208034. https://doi.org/10.1371/journal.pone.0208034

Mehling, W. E., Price, C., Daubenmier, J. J., Acree, M., Bartmess, E., & Stewart, A. (2012). The

Multidimensional Assessment of Interoceptive Awareness (MAIA). *PLoS ONE*, *7*(11),

e48230. https://doi.org/10.1371/journal.pone.0048230

Nord, C. L., Lawson, R. P., & Dalgleish, T. (2021). Disrupted Dorsal Mid-Insula Activation

During Interoception Across Psychiatric Disorders. *American Journal of Psychiatry*,

178(8), 761–770. https://doi.org/10.1176/appi.ajp.2020.20091340

Northoff, G., & Panksepp, J. (2008). The trans-species concept of self and the subcortical–

cortical midline system. *Trends in Cognitive Sciences*, *12*(7), 259–264.

https://doi.org/10.1016/j.tics.2008.04.007

Parkin, L., Morgan, R., Rosselli, A., Howard, M., Sheppard, A., Evans, D., Hawkins, A.,

Martinelli, M., Golden, A.-M., Dalgleish, T., & Dunn, B. (2014). Exploring the

Relationship Between Mindfulness and Cardiac Perception. *Mindfulness*, *5*(3), 298–313.

https://doi.org/10.1007/s12671-012-0181-7

Parvizi, J. (2001). Consciousness and the brainstem. *Cognition*, *79*(1–2), 135–160.

https://doi.org/10.1016/S0010-0277(00)00127-X

Paulus, M. P., & Khalsa, S. S. (2021). When You Don't Feel Right Inside: Homeostatic

Dysregulation and the Mid-Insular Cortex in Psychiatric Disorders. *American Journal of*

Psychiatry, *178*(8), 683–685. https://doi.org/10.1176/appi.ajp.2021.21060622

Pollatos, O., Gramann, K., & Schandry, R. (2007). Neural systems connecting interoceptive

awareness and feelings. *Human Brain Mapping*, *28*(1), 9–18.

https://doi.org/10.1002/hbm.20258

Power, J. D., Lynch, C. J., Dubin, M. J., Silver, B. M., Martin, A., & Jones, R. M. (2020).

Characteristics of respiratory measures in young adults scanned at rest, including

systematic changes and "missed" deep breaths. *NeuroImage*, *204*, 116234.

https://doi.org/10.1016/j.neuroimage.2019.116234

Price, C. J., & Hooven, C. (2018). Interoceptive Awareness Skills for Emotion Regulation:

Theory and Approach of Mindful Awareness in Body-Oriented Therapy (MABT).

Frontiers in Psychology, *9*. https://doi.org/10.3389/fpsyg.2018.00798

Price, C. J., Thompson, E. A., Crowell, S., & Pike, K. (2019). Longitudinal effects of

interoceptive awareness training through mindful awareness in body-oriented therapy

(MABT) as an adjunct to women's substance use disorder treatment: A randomized

controlled trial. *Drug and Alcohol Dependence*, *198*, 140–149.

https://doi.org/10.1016/j.drugalcdep.2019.02.012

Price, C. J., & Weng, H. Y. (2021). Facilitating Adaptive Emotion Processing and Somatic

Reappraisal via Sustained Mindful Interoceptive Attention. *Frontiers in Psychology*, *12*,

578827. https://doi.org/10.3389/fpsyg.2021.578827

R Core Team. (2017). *R: A language and environment for statistical computing. R Foundation

for Statistical Computing*. https://www.R-project.org/

Ring, C., & Brener, J. (2018). Heartbeat counting is unrelated to heartbeat detection: A

comparison of methods to quantify interoception. *Psychophysiology*, *55*(9), e13084.

https://doi.org/10.1111/psyp.13084

Roberts, R. L., Ledermann, K., & Garland, E. L. (2022). Mindfulness-oriented recovery

    enhancement improves negative emotion regulation among opioid-treated chronic pain

    patients by increasing interoceptive awareness. *Journal of Psychosomatic Research*, *152*,

    110677. https://doi.org/10.1016/j.jpsychores.2021.110677

Sardá-Espinosa, A. (2019). Time-Series Clustering in R Using the dtwclust Package. *The R

    Journal*, *11*(1), 22. https://doi.org/10.32614/RJ-2019-023

Schuette, S. A., Zucker, N. L., & Smoski, M. J. (2021). Do interoceptive accuracy and

    interoceptive sensibility predict emotion regulation? *Psychological Research*, *85*(5),

    1894–1908. https://doi.org/10.1007/s00426-020-01369-2

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., Reiss, A. L.,

    & Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience

    processing and executive control. *J Neurosci*, *27*(9), 2349–2356.

    https://doi.org/10.1523/JNEUROSCI.5587-06.2007

Smith, J. C., Ellenberger, H. H., Ballanyi, K., Richter, D. W., & Feldman, J. L. (1991). Pre-

    Bötzinger Complex: A Brainstem Region that May Generate Respiratory Rhythm in

    Mammals. *Science*, *254*(5032), 726–729. https://doi.org/10.1126/science.1683005

Smith, S., & Nichols, T. (2009). Threshold-free cluster enhancement: Addressing problems of

    smoothing, threshold dependence and localisation in cluster inference. *NeuroImage*,

    44(1), 83–98. https://doi.org/10.1016/j.neuroimage.2008.03.061

Spreng, R. N., Sepulcre, J., Turner, G. R., Stevens, W. D., & Schacter, D. L. (2013). Intrinsic

    architecture underlying the relations among the default, dorsal attention, and

    frontoparietal control networks of the human brain. *Journal of Cognitive Neuroscience*,

    25(1), 74–86. https://doi.org/10.1162/jocn_a_00281

Strigo, I. A., & Craig, A. D. (Bud). (2016). Interoception, homeostatic emotions and

    sympathovagal balance. *Philosophical Transactions of the Royal Society B: Biological*

    Sciences, *371*(1708), 20160010. https://doi.org/10.1098/rstb.2016.0010

Summerfield, C., Trittschuh, E. H., Monti, J. M., Mesulam, M.-M., & Egner, T. (2008). Neural

    repetition suppression reflects fulfilled perceptual expectations. *Nature Neuroscience*,

    11(9), 1004–1006. https://doi.org/10.1038/nn.2163

Szczepanski, S. M., Pinsk, M. A., Douglas, M. M., Kastner, S., & Saalmann, Y. B. (2013).

    Functional and structural architecture of the human dorsal frontoparietal attention

    network. *Proceedings of the National Academy of Sciences*, *110*(39), 15806–15811.

    https://doi.org/10.1073/pnas.1313903110

Tan, Y., Wei, D., Zhang, M., Yang, J., Jelinčić, V., & Qiu, J. (2018). The role of mid-insula in

    the relationship between cardiac interoceptive attention and anxiety: Evidence from an

    fMRI study. *Scientific Reports*, *8*(1), 17280. https://doi.org/10.1038/s41598-018-35635-6

Taylor, J. M. (2015). Psychometric analysis of the Ten-Item Perceived Stress Scale.

    Psychological Assessment, *27*(1), 90–101.

    http://dx.doi.org.myaccess.library.utoronto.ca/10.1037/a0038100

Trevisan, D. A., Altschuler, M. R., Bagdasarov, A., Carlos, C., Duan, S., Hamo, E., Kala, S.,

    McNair, M. L., Parker, T., Stahl, D., Winkelman, T., Zhou, M., & McPartland, J. C.

    (2019). A meta-analysis on the relationship between interoceptive awareness and

    alexithymia: Distinguishing interoceptive accuracy and sensibility. *Journal of Abnormal*

    Psychology, *128*(8), 765–776. https://doi.org/10.1037/abn0000454

Wang, X., Wu, Q., Egan, L., Gu, X., Liu, P., Gu, H., Yang, Y., Luo, J., Wu, Y., Gao, Z., & Fan,

    J. (2019). Anterior insular cortex plays a critical role in interoceptive attention. *ELife*, *8*,

    e42265. https://doi.org/10.7554/eLife.42265

Weiss, F., Zamoscik, V., Schmidt, S. N. L., Halli, P., Kirsch, P., & Gerchen, M. F. (2020). Just a

    very expensive breathing training? Risk of respiratory artefacts in functional

    connectivity-based real-time fMRI neurofeedback. *NeuroImage*, *210*, 116580.

    https://doi.org/10.1016/j.neuroimage.2020.116580

Wiens, S. (2005). Interoception in emotional experience. *Curr Opin Neurol*, *18*(4), 442–447.

Yang, C. F., & Feldman, J. L. (2018). Efferent projections of excitatory and inhibitory

    preBötzinger Complex neurons. *Journal of Comparative Neurology*, *526*(8), 1389–1402.

    https://doi.org/10.1002/cne.24415

Zoellner, L. A., & Craske, M. G. (1999). Interoceptive Accuracy and Panic. *Behaviour Research*

    and Therapy, *37*(12), 1141–1158. https://doi.org/10.1016/S0005-7967(98)00202-2