Huntingtin S421 phosphorylation increases kinesin and dynein engagement on early endosomes and lysosomes

Emily N. P. Prowse[1], Abdullah R. Chaudhary[1], David Sharon[1], Adam G. Hendricks[1]*

1. Bioengineering Department, McGill University, Montreal, QC, Canada

*adam.hendricks@mcgill.ca

Summary Statement
Huntingtin phosphorylation at S421 increases the outward motility of early endosomes and lysosomes by enhancing the microtubule binding and force generation of kinesin and dynein.

ABSTRACT

Huntingtin (htt) acts as a scaffolding protein that recruits motor proteins to vesicular cargoes, enabling it to regulate kinesin- and dynein-dependent transport. To maintain the native stoichiometry of huntingtin with its interacting partners, we used CRISPR/Cas9 to induce a phosphomimetic mutation of the endogenous htt at S421 (S421Dhtt). Using single-particle tracking, optical tweezers, and immunofluorescence, we examined the effects of this mutation on the motility of early endosomes and lysosomes. In S421Dhtt cells, early endosomes and lysosomes exhibited more outward motility towards the cell periphery compared to wild-type (WT) cells. In contrast, overexpression of htt had variable effects on the processivity, run length, and directional bias of both early endosomes and lysosomes. Kinesins and dyneins exerted greater forces on early endosomes and lysosomes in cells expressing S421Dhtt. Additionally, endosomes had higher binding rates, increased resistance to detachment under load, and enhanced recruitment of kinesins and dyneins in S421Dhtt cells. These data indicate that phosphorylation of the endogenous huntingtin causes early endosomes and lysosomes to move towards the cell periphery by activating both kinesins and dyneins.

List of Symbols and Abbreviations

α	Processivity parameter alpha
BDNF	Brain derived neurotrophic factor
DIC	Dynein intermediate chain
Early	Early endosome
EGF-qdot	Epidermal growth factor coated quantum dot
HD	Huntington's Disease
Htt	Huntingtin
KHC	Kinesin heavy chain
ks	Kolmogorov-Smirnov statistical test
latA	Latrunculin A
NZ	Nocodazole
R_g	Radius of gyration
S421Ahtt	Phosphoresistive huntingtin with the mutation S421A
S421Dhtt	Phosphomimetic huntingtin with the mutation S421D
WT+S421Dhtt	WT cells overexpressing mCherry-S421Dhtt
WT	Wild-type
WT+WThtt	WT cells overexpressing mCherry-WThtt

INTRODUCTION

Huntingtin (htt) is an essential, multifunctional, and ubiquitous protein. It consists of several disordered regions and 34 α-helical HEAT repeats (Fig. 1A) (Guo et al., 2018). Htt is essential for embryonic development (Duyao et al., 1995; Nasir et al., 1995; Schmitt et al., 1995; Zeitlin et al., 1995), and is ubiquitously expressed in tissues including the brain, ovaries, and testis into adulthood (Li et al., 1993; Sharp et al., 1995). It has diverse roles within the cell, including regulation of gene expression (Dunah et al., 2002; Futter et al., 2009; Holbert et al., 2001; Li et al., 2002; Marcora et al., 2003; Zuccato et al., 2003), cell morphology (Lo Sardo et al., 2012; Thion et al., 2015), cell survival (Leavitt et al., 2006), and intracellular transport (Cason et al., 2021; Gauthier et al., 2004; Liot et al., 2013; Twelvetrees et al., 2010; Wong and Holzbaur, 2014). Htt enhances transport of brain derived neurotrophic factor (BDNF) vesicles (Gauthier et al., 2004; Zala et al., 2008), autophagosomes in the mid-axon (Cason et al., 2021; Wong and Holzbaur, 2014), TrkB signalling vesicles (Liot et al., 2013), and synaptic vesicles through GABA_R signalling (Twelvetrees et al., 2010). The remarkable ubiquity and variety of htt's roles underscore the importance of understanding its function in health and disease.

Teams of kinesins and dyneins transport signalling and degradative cargoes. There are multiple kinesins that drive outward transport towards microtubule plus ends at the cell periphery. Different ratios of kinesin-1, kinesin-2, and kinesin-3 associate with distinct cargo populations, allowing different motility characteristics for each population of vesicles (Bentley et al., 2015). Each type of kinesin has a unique binding rate and velocity, which leads combinations of motors to create a motility barcode for cargoes. Unlike kinesins, there is only one form of cytoplasmic dynein that drives organelle transport. Dynein forms a complex with dynactin and activating adaptors such as BicD, Hook1, and others to direct inward transport towards microtubule minus ends at the cell center (Cason et al., 2021; McKenney et al., 2014; Schlager et al., 2014). Scaffolding and adaptor proteins work together to regulate the recruitment and activity of both kinesin and dynein motors on each cargo based on its identity.

Huntingtin recruits teams of kinesin, dynein, and myosin VI motors to cargoes, allowing it
to control the cargo's direction of movement along microtubules and switching between
actin and microtubules (Caviston et al., 2011). Htt interacts with molecular motors
indirectly through adaptors; kinesin-1 and the dynein adaptor dynactin interact with htt
through HAP1 (Engelender, 1997; Li et al., 1998; McGuire et al., 2006; Twelvetrees et
al., 2010), and myosin VI interacts through optineurin (Sahlender et al., 2005). In addition,
dynein interacts directly with htt through its intermediate chain, suggesting a prominent
role for htt in regulating inward transport (Caviston et al., 2007). Htt's interaction with
myosin VI suggests htt regulates transitions to the actin cytoskeleton for short range
transport or tethering. Upon htt deletion or mutation, intracellular transport of many
vesicular cargoes is severely inhibited (Caviston and Holzbaur, 2009; Hilditch-Maguire,
2000).
44
Polyglutamine expansion mutations in htt lead to Huntington's disease (HD), disrupting
multiple cellular functions and leading to neurodegeneration in medium spiny neurons of
the striatum (Gil and Rego, 2008). HD mutations cause htt to form novel interactions.
Notably, through increased self-association mutant htt forms aggregates throughout the
cytoplasm and nucleus (Davies et al., 1997). These interactions cause aberrant initiation
of autophagy, mTOR signalling, mitochondrial fission, and palmitoylation (reviewed in
Wanker et al., 2019). Additional defects occur in neuronal transcription, endoplasmic
reticulum homeostasis, mitochondrial function, calcium signalling, synaptic activity, and
intracellular transport (reviewed in Saudou and Humbert, 2016), demonstrating the
importance of htt's native interaction network in maintaining homeostasis (Colin et al.,
2008; Lemarié et al., 2021; Zala et al., 2008).
56
Huntingtin is regulated by multiple post-translational modifications, including
palmitoylation, phosphorylation, sumoylation, ubiquitination, and acetylation. The
functions and mechanisms of many of these modifications remain unknown. Its
palmitoylation site at C214, regulated by HIP14 in the Golgi, facilitates interactions with
vesicular membranes (Yanai et al., 2006). Palmitoylation is reduced in cells expressing
the polyglutamine-expanded htt protein that leads to HD, while restoring palmitoylation

reduces aggregation and cytotoxicity in vitro (Lemarié et al., 2021). Sumoylation sites

may contribute to nuclear import and transcriptional regulation (Steffan et al., 2004).

Ubiquitination and acetylation regulate huntingtin-mediated autophagy (Wong and

Holzbaur, 2014). Htt has 40 known phosphosites, many of which have unknown biological

functions (reviewed by Saudou and Humbert, 2016). The phosphorylation state of serine

421 affects axonal transport direction and velocity of BDNF vesicles and

autophagosomes (Colin et al., 2008; Wong and Holzbaur, 2014), while mitochondrial

transport is unaffected by this phosphosite (Xu et al., 2020). Each of htt's post-

translational modifications contribute to its role in transport of both signaling and

degradative cargoes. Further, post-translational modifications are reduced in HD,

disrupting the balance of htt's interactions with its binding partners (Colin et al., 2008;

Lemarié et al., 2021; Zala et al., 2008).

Post-translational modifications of scaffolding proteins can influence transport direction.

The phosphomimetic mutation (S421Dhtt) mimics a constitutively phosphorylated amino

acid and biased BDNF transport towards the cell periphery, while phosphoresistive

mutations (S421Ahtt) led to increased transport towards the cell center (Colin et al.,

2008). The S421Dhtt mutation is neuroprotective in cells with the mutation that leads to

HD (Zala et al., 2008), which motivates our attempt to understand the native role of this

phosphosite in cells with normal (WT) htt. It has also been generalized for various

scaffolding proteins including RILP, Miro/TRAK1/TRAK2, JIP1, JIP3/4 that post-

translational modifications can modify transport direction (Fu and Holzbaur, 2014). These

studies suggest that huntingtin phosphorylation directs transport towards the cell

periphery.

To elucidate the effects of S421 phosphorylation at native expression levels and

determine the mechanisms of regulation for both early endosomes and lysosomes, we

applied CRISPR-based gene editing and single-molecule biophysics. Previous studies

analyzed the motility of cargoes overexpressing S421Dhtt or S421Ahtt constructs and

performed Western blotting to determine S421's role in transport regulation (Colin et al.,

2008; Zala et al., 2008). Western blot and immunofluorescence in overexpressing COS7

cells suggests that S421 phosphorylation increases the recruitment of kinesin to cargoes
(Colin et al., 2008; Zala et al., 2008). However, only a subset of the motors bound to a
cargo might be active (Encalada et al., 2011). In addition, the critical stoichiometry of
interactions between scaffolding proteins and motors is disrupted in overexpression
systems. We generated an endogenous mutation to replicate htt's native interactions and
expression levels and observed its effect on organelle transport. We measured the forces
generated by the motors driving early endosome and lysosome transport with optical
tweezers to determine whether phosphorylation of huntingtin modifies motor activity.
Single particle tracking allowed us to obtain nanometer-resolution tracking of cargo
motility. Combining single-molecule biophysical approaches with gene editing provided
mechanistic insight into htt's regulation of intracellular transport by S421 phosphorylation.
105

To understand how S421 phosphorylation of htt regulates intracellular transport, we
examined early endosome and lysosome transport in CRISPR-generated S421Dhtt
HEK293T cells. Motility assays demonstrated an increase in outward (microtubule plus-
ended) motility for both early endosomes and lysosomes in cells expressing S421Dhtt.
Additionally, lysosome run length increased upon S421 phosphorylation. Examining the
forces generated by microtubule motors on early endosomes and lysosomes, we
observed both kinesin and dynein motors had higher binding rates and increased
resistance to unbinding under load with S421Dhtt for both cargoes. Cells with S421Dhtt
showed higher forces in both outward and inward directions on both early endosomes
and lysosomes. A higher fraction of kinesins and dyneins associated to microtubules upon
S421 phosphorylation. The phosphorylation-induced change in directional bias and force
generation for lysosomes was more subtle than those for early endosomes. We
hypothesize that the native motility characteristics of the cargoes dictate the effect of
S421 phosphorylation on htt: inward-directed early endosomes become bidirectional
while bidirectional lysosomes increase their displacement. Our data indicates
phosphorylation of htt at S421 increases motor recruitment and activity of early
endosomes and lysosomes. Additionally, overexpression of WT and S421D htt suggests
overexpressed htt disrupts the transport complex of htt, motors, and adaptors required
for transport, affecting cargo run length, processivity, and direction.

RESULTS

Endogenous S421D mutation does not affect cytoskeletal organization or huntingtin localization

Introducing a phosphomimetic S421D mutation in the endogenous huntingtin did not alter huntingtin expression levels and localization or the organization of the cytoskeleton and golgi. In generating the S421D HEK293T cell line, we aimed to preserve the native expression levels and stoichiometry of interactions between htt and its many interactors. We used a lentiviral CRISPR cassette to generate this S421D point mutant cell line (Fig. 1B) and identified a single positive clone with the desired mutation using next-generation sequencing (Fig. 1C). S421Dhtt did not result in appreciable changes to the actin nor microtubule morphology (Fig. 1D). To determine whether S421Dhtt caused major defects in dynein function, we probed the organization of the Golgi apparatus (Burkhardt et al., 1997; Caviston et al., 2007; Corthésy-Theulaz et al., 1992). We observed no changes in the percentage of cells with organized versus dispersed Golgi, indicating dynein function was not severely disrupted (Fig. 1E). Importantly, the S421D mutation did not affect huntingtin's localization nor expression level (Fig. 1F, G). Overexpression of mCherry-WThtt or mCherry-S421Dhtt leads to elevated expression of both constructs at similar levels (Fig. 1G). In iPSC-derived neurons expressing polyglutamine-expanded huntingtin (180Q), S421Dhtt led to increased mitochondrial surface area while the number of mitochondria remained similar (Xu et al., 2020). Immunofluorescence of the mitochondria in HEK293T cells showed a decrease in the mitochondrial number and area upon S421D mutation (Fig. 1H-J), suggesting that S421Dhtt mitochondrial regulation may be cell-type dependent. Together, immunofluorescence and huntingtin expression level characterization demonstrated that the S421D mutation did not substantially alter the cytoskeleton, Golgi apparatus, nor localization of huntingtin while mitochondrial morphology was affected.

Huntingtin phosphorylation directs early endosomes towards the cell periphery

Introducing the S421D phosphomimetic mutation in the endogenous huntingtin gene
induced an outward directional bias in early endosomes but did not alter run length nor
processivity. We hypothesized that the processivity, run length, and fraction of outwardly-
directed early endosomes would increase upon S421 phosphorylation based on previous
work with BDNF vesicles in neurons (Colin et al., 2008). We quantified the processivity of
organelle trajectories using the slope of the log-log plot of mean-squared displacement
(α). For cargoes transported by microtubule-based motor proteins, we expect that $\alpha > 1$ is
indicative of directed transport, while $\alpha = 1$ indicates diffusive motion. The radius of
gyration (R_g) is an indicator of run length that determines the average position of a
trajectory and measures a radius containing half of the points in the trajectory from its
average position. We transfected cells with Rab5-eGFP to label early endosomes for
particle tracking (Fig. 2A, C). The endogenous S421D mutation increased the outward
motility of early endosomes towards the cell periphery without changing their processivity
or run length (Fig. 2E, F, I). The outward directional bias was previously described when
truncated WThtt and S421Dhtt were overexpressed in primary neurons (Colin et al., 2008;
Zala et al., 2008). To dissect the effects of htt phosphorylation and expression levels, we
transiently transfected full length mCherry-WThtt (WT+WThtt) or mCherry-S421Dhtt
(WT+S421Dhtt). These transfections lead to a ~4-10-fold increase in htt expression (Fig.
1G). Overexpression of S421Dhtt increased the processivity, run length, and
unexpectedly induced inward motility of early endosomes (Fig. 2E, F, I). Interestingly,
overexpression of WThtt increased the outward motility of early endosomes (Fig. 2I).
These unexpected results in overexpression conditions may be due to the change in
stoichiometry of htt's interactions with its native binding partners. The gene edited S421D
HEK293T cells demonstrate an outward bias of early endosomes, while htt
overexpression alters motility. This indicates that S421 phosphorylation is either recruiting
additional kinesins on early endosomes or increasing the activity of the kinesins present
on the early endosome.
183
184 **S421D increases run length and outward bias of lysosomes**
185

S421 phosphorylation increased lysosome run lengths and directed them toward the cell periphery. We hypothesized S421Dhtt's effects to be universal for all htt-associated cargoes, and therefore expected increased outward bias, processivity, and run length in lysosomes. This hypothesis is supported by previous work showing that htt promotes outward motility in the mid-axon of neurons for autophagosomes, which eventually mature into lysosomes (Cason et al., 2021; Wong and Holzbaur, 2014). In both WT and S421D experiments, the raw trajectories demonstrated the expected bidirectional motility (Fig. 2B, D). S421Dhtt induced an increase in run length and outward motility of lysosomes, without affecting their processivity (Fig. 2G, H, J). When we overexpressed htt, we observed a minor increase in processivity for WThtt and a substantial increase in run length regardless of the phosphorylation state (Fig. 2G, H, J). Unexpectedly, WThtt overexpression led to an inward directional bias (Fig. 2J), while overexpression of S421Dhtt reversed the effect by increasing outward motility. Taken together, our results indicate that htt overexpression modifies processivity, run length, and directional bias while phosphorylation shifts lysosome motility towards the cell periphery and increases their run length. This suggests S421 phosphorylation at native expression levels is increasing the number of active kinesin motors, as was observed for early endosomes.

Tethering on actin contributes to the directional bias of early endosomes

Our results suggest that huntingtin's phosphorylation state modulates transient actin tethering and contributes to the transport direction of early endosomes. Huntingtin interacts with optineurin, which recruits myosin VI and could regulate cargo association to actin (Sahlender et al., 2005). We used latrunculin A (latA) to depolymerize actin filaments and reduce cargo association with actin. We expected an increase in run length and processivity upon latA treatment since cargo tethering and confinement by actin are reduced, allowing long range, processive motility by microtubule motors to dominate. We determined the optimal concentration for motility assays via imaging cells labelled with SiR-actin. We determined 250 nM latA was optimal as it resulted in significant depolymerization of actin filaments without loss of native cellular morphology (Fig. S1A). As expected, treatment with latA increased run length and processivity of Rab5 early

endosomes (Fig. S1B, C). Additionally, the phosphorylation-induced directional bias
became less prominent with latA, suggesting that short-range actin-based motility
contributes to the observed directional bias of early endosomes (Fig. S1D). WT early
endosomes have increased outward motility upon latA treatment, indicating that allowing
kinesin and dynein to dominate transport biases it outwards (Fig. S1D). The S421D
mutation did not further shift the directional bias in actin-depleted cells, suggesting that
both loss of actin tethering and htt phosphorylation act through similar mechanisms.

Lysosomes have a higher dependence on microtubule-based transport

Depolymerization of microtubules demonstrated no significant changes in processivity or
run length for early endosomes, while lysosomes showed significant decreases in both
run length and processivity. We used 10 µM nocodazole (NZ) to depolymerize
microtubules and observe how strongly the motility parameters were affected. We
expected that upon NZ treatment, both processivity and run length would decrease
because microtubule-based motors drive the long range processive transport of both
early endosomes and lysosomes. We observed this expected decrease in run length and
processivity for lysosomes, suggesting that they rely heavily on microtubule-based
transport to drive their motility (Fig. S1F, I, J). The large change in processivity and run
length for lysosomes demonstrates the magnitude of changes in these parameters upon
large scale disruption of the cytoskeleton. Early endosomes demonstrated a surprising
lack of effect of nocodazole treatment on their processivity and run length, perhaps due
to their inherently low processivity and run lengths compared to lysosomes (Fig. S1E, G,
H). The lysosomes in NZ-treated cells reached similar values for processivity and run
length to the early endosomes in the control condition, suggesting that the values for early
endosome motility were close to their minimum. Combining this result with our
conclusions from the latA treatment demonstrates that early endosome transport may be
more dependent on switching between actin-based and microtubule-based transport (Fig.
S1). These results indicate early endosome transport has a lower dependence on
microtubules compared to lysosomes, perhaps due to its inherently short run lengths and
lower processivity.

S421 huntingtin phosphorylation activates kinesins and dyneins on early endosomes

To determine how changes in motor activity contribute to the motility we observed, we
measured the force generation of motors on early endosomes with optical tweezers. We
incubated cells with fibronectin-coated 500 nm polystyrene beads for 10-50 mins (Blocker
et al., 1997) and used optical tweezers to measure the forces that motors exerted in living
cells (Fig. 3A, B). At this stage of maturation, the early phagosomes (termed early
endosomes) are positive for Rab5 and other early endosome markers and represent a
subpopulation of the vesicles labeled by Rab5-eGFP. Early endosome force traces
demonstrated the expected bidirectional motility biased inwards (Fig. 3B). We predicted
kinesins would specifically be activated, or dyneins inactivated to generate the directional
bias we observed for early endosomes (Fig. 2I). Unexpectedly, the stall force distributions
for early endosomes shifted towards higher forces in both outward and inward directions
with S421Dhtt (Fig. 3C), suggesting both kinesins and dyneins were activated. In cells,
each kinesin motor exerts forces of 2-7 pN, and each dynein exerts approximately 2-3 pN
(Blehm et al., 2013). Looking closely at the distributions, we observe frequent kinesin
forces around 3 pN and 6 pN for cells with WThtt (Fig. 3C). Both of those populations are
observed in S421Dhtt cells, in addition to high force events near 9 pN and 12 pN.
Observing the inward directed forces, WT cells show frequent dynein forces near 1.5 pN
and 5 pN, while forces shift to ~ 3.5 pN, 9 pN, and 13 pN in S421Dhtt cells. Together,
these forces likely correspond to ~1-2 active kinesins and 1-3 active dyneins in WT cells,
and 1-4 active kinesins and 2-6 active dyneins in S421D cells, which is slightly lower than
previous estimates for late phagosomes (Chaudhary et al., 2018; Hendricks et al., 2010).
We then sought to determine how the force-dependent unbinding rates of each motor
were affected by S421Dhtt, predicting that the kinesin motors would have slower
unbinding and therefore induce the kinesin-based directional bias (Fig. 2I). The unbinding
rates of kinesin and dynein teams on early phagosomes in WT cells is similar (Fig. 3D,
Table S1). We observed slower unbinding under load of both kinesin and dynein teams
with S421Dhtt, consistent with more kinesin and dynein motors engaged with the

microtubule (Fig. 3D, Table S1). Next, we measured the binding rate of early endosomes

moving inwards and outwards and found the binding rate for early endosomes was

increased for both directions in S421Dhtt cells (Fig. 3G, H). The binding rate was more

strongly enhanced for outward forces, suggesting an increased number of active kinesins

enhances binding compared to WThtt (Fig. 3G). Observing the cumulative distribution

functions, we note that outward and inward motor binding rates in S421Dhtt cells were

very similar (Fig. 3H). This suggests that the kinesins are being activated to the same

level as the dyneins, reducing the native inward bias for these cargoes. In summary, the

force data indicate that huntingtin phosphorylation at site S421 increases the net force

generation, affinity, and binding strength of both kinesin and dynein motors in early

endosomes.

S421Dhtt increases lysosome microtubule binding and resistance to unbinding

under load

Like for early endosomes, we hypothesized that kinesins would have increased activity

on lysosomes as they also demonstrated an increased run length and outward directional

bias (Fig. 2H, J). We measured the force characteristics of phagolysosomes by modifying

the optical tweezers assay described for early endosomes with a longer incubation time

of 1-2 hrs. The phagolysosomes measured in the optical tweezer experiments, termed

lysosomes hereafter, represent a subpopulation of lysotracker-positive lysosomes,

compared to our earlier analysis tracking all lysotracker-positive vesicles (Fig. 2). Sample

force traces demonstrated typical bidirectional motility of lysosomes (Fig. 3B). We

observed an increase in forces in both directions for lysosomes as for the early

endosomes, although to a lesser degree (Fig. 3E). The balance of kinesin to dynein forces

remained similar in both control and S421Dhtt conditions, as expected for these

bidirectional cargoes. Predicting the number of motors as described for early endosomes,

the force distributions suggest that 1-2 additional kinesins and 1-2 additional dyneins are

engaged in S421Dhtt cells. The unbinding rates indicate that kinesin teams are more

resistant to unbinding than dyneins for lysosomes (Fig. 3F). In S421Dhtt cells, we

observed a slightly lower resistance to unbinding for lysosomes moving outward, while

the unbinding resistance was higher than WT for inward-moving cargoes (Fig. 3F, Table

S1). Finally, we examined the binding rates for lysosome motors to determine how

S421Dhtt affected microtubule affinity. S421Dhtt increased the binding rate for both

inward- and outward- directed cargoes compared to WT (Fig. 3I, J). The increased forces

and enhanced binding we observed in S421Dhtt cells suggests htt phosphorylation

recruits and/or activates both kinesins and dyneins.

S421Dhtt increases kinesin and dynein association to microtubules

As we did not observe increased kinesin activity compared to dynein upon S421

phosphorylation, we asked if more kinesins were engaged with microtubules to generate

the observed directional bias. We used immunofluorescence imaging to approximate the

differences in the number of motors associated with microtubules in each cell type. We

fixed the cells with methanol to remove cytosolic components and determine the relative

number of motors that were bound to microtubules in each condition at the time of fixation

(Fig. 4A, B). We then created intensity heatmaps of all images from the microtubules and

kinesin-1 heavy chain (KHC) or dynein intermediate chain (DIC) and plotted them on the

same axes (Fig. 4C-F). We detected higher intensities in the motors channel for both KHC

and DIC in the S421Dhtt cells (Fig. 4C-F). The higher intensity indicates a higher

concentration of motors on extracted microtubules in the S421Dhtt cells. Upon calculating

the correlation coefficients between the microtubule and motor channels for each

individual image, we observed that the dynein and microtubule intensities were more

highly correlated in S421Dhtt cells than WT cells. This assay demonstrates that S421Dhtt

increases the fraction of kinesin and dynein motors associated to microtubules. This

supports our data from the optical tweezers analysis, which showed a higher binding rate

and lower unbinding rate for motors in the S421Dhtt cells in both early endosomes and

lysosomes. Further, the signal for microtubule-bound kinesin and dynein motors is higher

in S421Dhtt cells, suggesting htt S421 phosphorylation increases the engagement of both

kinesins and dyneins with microtubules.

DISCUSSION

Our results show that huntingtin phosphorylation at S421 increases the number of active

kinesin and dynein motors on endosomes. Surprisingly, huntingtin activates microtubule

motors on endosomes at varying stages of maturation, from early endosomes to

lysosomes. Huntingtin also regulates the motility of neuronal signalling vesicles including

BDNF (Colin et al., 2008; Zala et al., 2008). In contrast, most cargo adaptors are recruited

to specific cargoes at specific stages of maturation (Cason et al., 2021; Mogre et al.,

2021). The ubiquitous function of huntingtin on a wide array of cargoes illustrates its role

as a central regulator of vesicular transport.

While kinesin and dynein teams exert more force and have enhanced microtubule binding

on both early endosomes and lysosomes in S421D huntingtin cells (Fig. 3,4), its effect on

the motility of different endocytic cargoes varies. For early endosomes, huntingtin

phosphorylation results in a pronounced shift in motility from primarily inward towards the

cell center to bidirectional motility (Fig. 2I). The processivity and displacement of early

endosomes are only weakly affected (Fig. 2E, F). In comparison, lysosomes display a

small but significant shift in motility towards the cell periphery in phosphomimetic htt cells

compared to control cells (Fig. 2J) while their run length is strongly enhanced (Fig. 2H).

Taken together, these results suggest that huntingtin phosphorylation regulates transport

by enhancing the activity of both kinesin and dynein (Fig. 5A, B). However, its effects on

the motility of different cargoes depend on their basal motility characteristics. In the basal

condition, early endosome motility is strongly biased towards the cell center. Activating

both kinesin and dynein results in a loss of the inward bias. Lysosome motility is

approximately equally distributed towards the cell center and periphery, such that

activating kinesin and dynein does not alter directionality, but rather increases the total

run length.

Disrupting actin filaments resulted in an increased outward motility similar to the effect of

S421Dhtt (Fig. S1A-D), suggesting a potential role for myosin VI-mediated tethering to

actin filaments (Sahlender et al., 2005) in generating the directional bias (Al-Haddad et

al., 2001). Yet, optical trapping and immunofluorescence show that S421Dhtt results in

enhanced forces and recruitment to microtubules for both kinesin and dynein (Figs. 3, 4),
suggesting S421D phosphorylation instead acts through activating microtubule motors.
Thus, we hypothesize that outward directional bias occurs in situations where microtubule
motors dominate transport. That could be achieved either by reducing tethering on actin
or by engaging more kinesins and dyneins.
377
Our findings are in broad agreement with previous studies showing that overexpressing
S421Dhtt resulted in increased outward motility of BDNF and TI-VAMP vesicles
compared to cells expressing phosphoresistive S421A htt constructs (Colin et al., 2008;
Zala et al., 2008). Further, Colin et al. observed increased kinesin-1 recruitment to
microtubules and enhanced interactions between the p150glued subunit of dynactin and
KHC upon S421Dhtt expression (Colin et al., 2008). Interestingly, we did not see KHC-
specific recruitment to microtubules with endogenous S421Dhtt compared to wild-type.
Instead, we observed an increase in recruitment of both motors, with slightly higher
dynein recruitment. These differences between previous results and our observations
may reflect effects of htt overexpression. Alternatively, they may be due to differences in
htt function in the embryonic kidney cells used here compared to previous studies in
neurons, which are highly polarized and have higher endogenous htt expression.
390
Overexpression of htt alters the processivity, run length, and directional bias of early
endosomes and lysosomes. The effects of overexpression are highly variable and often
do not reflect the trends observed when inducing mutations in the endogenous htt (Fig.
2), suggesting that maintaining the stoichiometry of htt to its many interacting proteins is
critical. We expect that excess htt in the cytoplasm is interacting with and sequestering
the motors and adaptors required to generate the effect that we observe in transport upon
overexpression (Fig. 5C). This result supports the known consequences of
overexpression experiments, which cause adverse effects by changing the stoichiometry
of interactions with other proteins in the cell. For example, overexpressing the p50
dynamitin subunit of dynactin causes a dominant negative effect in which early
endosomes and lysosomes become peripherally localized due to dynein sequestration
(Burkhardt et al., 1997). Regardless of the phosphorylation state, overexpression of htt

affected the directional bias of early endosomes and lysosomes. Processivity and run
lengths were increased for lysosomes in cells overexpressing either WT or S421Dhtt,
however, early endosomes only demonstrated this effect overexpressing S421Dhtt. This
work, supported by many previous studies discussing expression levels (e.g. (Burkhardt
et al., 1997; Dambournet et al., 2018)) highlights the importance of examining protein
function at endogenous expression levels to ensure biological relevance of the effects.
This is particularly important for scaffolding proteins like huntingtin, which mediates
interactions among many adaptors, effectors, motors, and membranes.
411
Transport is tightly regulated on multiple levels within the cell to ensure specific cargoes
arrive at their intracellular destination. Scaffolding proteins such as htt are a primary
mechanism of regulation that mediate interactions between cargoes and motor proteins,
and are the only known level of regulation that is cargo specific (Cason et al., 2021; Colin
et al., 2008; Fu and Holzbaur, 2014). Many scaffolds specifically interact with a limited
set of cargoes and are regulated by post-translational modifications (Cason et al., 2021;
Fu and Holzbaur, 2013; Fu and Holzbaur, 2014; Reed et al., 2006; Sun et al., 2017).
Here, we illustrate the role of htt in regulating early endosome and lysosome transport.
Htt S421 phosphorylation activates kinesin and dynein on both early endosomes and
lysosomes. However, enhanced kinesin and dynein activity differentially affects early
endosomes and lysosomes, influencing their direction, motor activity, and affinity for
microtubules. Our results underscore the central role of huntingtin in directing endosome
transport and suggest that huntingtin mutations likely result in cargo-specific disruptions
to both signalling and degradative pathways.
426
427 **MATERIALS AND METHODS**
428
Cell Culture
We passaged HEK293T cells (donated by Dr. Kamen's lab) at 80% or higher confluency
in a biosafety cabinet starting by rinsing with 2.5 mL 37 °C 1X phosphate buffered saline
(PBS, Wisent, St-Jean-Baptiste, QC, Canada), followed by adding 400 µL of 4 °C 0.25%
trypsin (Wisent) and incubating for 5 mins at 37 °C. Next, we used 5 mL of 37 °C complete
medium (Dulbecco's Modified Eagle Medium (DMEM), 10% fetal bovine serum (FBS), 2

mM Glutamax, 60 µg/mL penicillin 100 µg/mL streptomycin, all components from Gibco, Waltham, MA, United States) to neutralize the trypsin activity. Finally, we placed a 1/5-1/20 dilution of the resuspended cells in 37 °C fresh complete medium and transferred to the 37 °C, 5% CO_2 incubator. We tested all stocks of cells for mycoplasma.

CRISPR Design and Molecular cloning

We transfected HEK293T cells with 1.25 µg of lentiCRISPR v2 cassette and 1.25 µg of the single-stranded oligodeoxynucleotice (ssODN) for 24-36 hrs. lentiCRISPR v2 was a gift from Feng Zhang (Addgene plasmid # 52961 ; http://n2t.net/addgene:52961 ; RRID:Addgene_52961, Watertown, MA, United States) (Sanjana et al., 2014). Subsequently, we selected expressing cells using 2 µg puromycin for 48-72 hrs. Finally, we isolated single cells using fluorescence activated single cell sorting and grew them for 2-3 weeks in complete media containing 50% FBS. We verified clones containing the S421D sequence with next-generation sequencing.

gRNA sequence:

GTTCCACAATACTCCCACTACGG

ssODN sequence:

ACGCCTCCACCCGAGCTTCTGCAAACCCTGACCGCAGTCGGGGGCATTGGGCAG

CTCACCGCTGCTAAGGAGGAGTCTGGTGGCCGATCTCGTAGTGGGGATATTGTGG

AACTTATAGGCAAGTTATTAGCAAGGTCTACTCTTACAATTAACTTTGCAGTAATAC

TAGTTACACTCTATTGATTATGGGCCTGCCCTGT

Sequencing primers:

Forward: CACCGTTCCACAATACTCCCACTA

Reverse: AAACTAGTGGGAGTATTGTGGAAC

Plasmids and Transfection

We transfected cells with 0.8 µg of eGFP-Rab5 or mCherry-S421D-htt or mCherry-WT-htt huntingtin constructs 24-48 hrs before imaging. mCherry-S421D-htt and mCherry-WT-htt constructs were generously donated from the Saudou lab, generated from the full length pARIS-htt (Pardo et al., 2010) using methods from their previous work (Humbert

et al., 2002). We replaced cell culture medium in the 35 mm round glass bottom dishes
(Mattek, Ashland, MA, United States) containing the cells with 0.2 mL serum-free DMEM.
In separate 1.5 mL tubes, we added 2 µL/dish Lipofectamine 2000 (ThermoFisher
Scientific, Waltham, MA, United States) to 50 µL/dish serum-free DMEM and added 0.8
µg/dish of DNA to 50 µL/dish of serum-free DMEM. We then mixed the DNA and
lipofectamine, incubated for 20 mins at room temperature, and added 100 µL/dish to the
cells. Four hours later, we replaced the serum-free medium with complete media.
473

Western Blotting

We lysed cells by manually twisting while lifting and descending a Dounce homogenizer
20-30 times in 400 µL PBS containing 0.01 M dithiothreitol (DTT, R0681, ThermoFisher
Scientific) and 1X protease inhibitor cocktail (100X stock from PIC002, Bioshop,
Burlington, ON, Canada) on ice. Subsequently, we centrifuged the cell solution at 1500 g
for 10 mins at 4 °C, then centrifuged the supernatant at 16000 g for 10 mins at 4 °C. We
added 20 µL of lysate for each sample to the SDS-PAGE gel, diluted in 2x Laemmli buffer
(161-0737, Biorad, Hercules, CA, USA) with 2.5% 2-mercaptoethanol (M6250, Sigma,
Burlington, MA, United States) after heating to 100 °C for at least 10 mins. To mark
molecular weights, we loaded 5 µL of protein ladder (PI26634, ThermoFisher Scientific).
We used a 25 mM Tris (TRS001.1, Bioshop), 190 mM glycine (800-045-LG, Wisent), and
0.1% SDS (SDS999.100, Bioshop) running buffer, and ran the gel at 80 mA for 60-90
mins. To transfer, we used 25 mM Tris, 190 mM glycine, and 30% methanol (34860-4L-
R, Sigma) transfer buffer for wet transfer at 100 V for 90 mins with a PVDF membrane.
We blocked the membrane for 1 hour with blocking buffer: 5% bovine serum albumin
(BSA, Bioshop) in tris-buffered saline with Tween 20 (TBST, 20 mM Tris, 150 mM NaCl
(SOD004.1, Bioshop), 0.1% Tween 20 (P2287, Sigma). We diluted primary antibodies in
the blocking buffer and incubated at 4 °C overnight. Next, we washed the membrane with
blocking buffer 3 times for 5 mins and incubated with the secondary antibody diluted in
blocking buffer for 90 mins with constant agitation. We washed the membrane 3 times
with TBST before incubating with a 1:1 ratio of horseradish peroxidase substrate and
enzyme mixture (P90720, EMD Millipore, Burlington, MA, USA). We imaged using a
Biorad Chemidoc camera set to detect chemiluminescence. We subsequently stained the

membrane with Coomassie G-250 (CBB555.25, Bioshop), using a previously established
protocol from Goldman et al. titled Coomassie blue R-250 staining (Goldman et al., 2016).
We used the following primary antibodies: 1/1000 dilution of T9026 ms α-tubulin, Sigma
lot: 000089497, and 1/500 dilution of MAB2166 ms-huntingtin, Abcam, Cambridge,
England lot: 3703568. The secondary antibody was rabbit anti-mouse ab6728, Abcam.
502

Motility Assay
In a biological safety cabinet, we mixed 200 µL of 37 °C complete medium with
LysoTracker™ Deep Red (L12492, ThermoFisher Scientific) to a concentration of 70 nM
and added it to HEK293T cells in MatTek 35 mm glass bottom dishes. Cells were
incubated for 10 mins at 37 °C, and subsequently replaced the medium with 1 mL of 37
°C Lebovitz (11415064, Gibco) with 10 % FBS for imaging. We imaged cells in a 37°C
chamber of a Nikon Eclipse Ti-E microscope in near-total internal reflection (TIRF)
fluorescence settings using an EMCCD camera with an exposure time of 120 ms for 3
mins. We repeated the process for several cells within each dish and all cells were imaged
within 1 hour of adding the Lysotracker. For Rab5 transfected cells, we replaced the
medium with 0.5-1 mL of of 37 °C Lebovitz with 10 % FBS and imaged them as the
lysosomes. For experiments using 250 nM latrunculin A (S428021, Sigma) or 10 µM
nocodazole (M1404, Sigma), we added the drug to the Lebovitz medium a minimum of
15 mins prior to imaging.
517

Immunofluorescence: Cytoskeleton and Golgi Apparatus
This protocol was adapted from Jimenez et al. (Jimenez et al., 2020). We passaged
HEK293T cells onto rectangular coverslips (12544-A ThermoFisher Scientific) in 100 mm
petri dishes (83.3902, Sarstedt, Saint-Leonard, QC, Canada) ~24 hrs prior to fixation. We
performed incubations for extraction, fixing and NaBH$_4$ (452882, Sigma) solutions on flat
pieces of parafilm, with droplets containing 100 µL of the described solution. Using fine-
tipped tweezers, we manipulated the coverslips and placed them cell-side down on the
droplets. Firstly, we incubated the cell-adhered coverslips with 37 °C extraction solution
(0.25% Triton-X-100 (X100, Sigma), 0.1% glutaraldehyde (G5882, Sigma)) in PEM
(80mM PIPES (P6757, Sigma), 5mM EGTA (EGT101, Bioshop), 2mM MgCl$_2$ (M2866,

Sigma) at pH=6.8) for 15-45 s. We next fixed the cells with 37 °C 0.25% Triton-X-100 and

0.5% glutaraldehyde in PEM and incubated for 10 mins. We subsequently transferred the

coverslips to 0.1% NaBH$_4$ in PBS (Wisent) and incubated for 7 mins, then rinsed quickly

with PBS twice before blocking. Next, we incubated the fixed cells with blocking buffer

(2% BSA (ALB001, Bioshop) in PBS, (Wisent)) inside a glass petri dish for 30 mins with

gentle rocking on an orbital shaker. Subsequently, we added a 1:400 solution of primary

antibody in blocking buffer (anti-GM130 ab52649 Abcam, anti-tubulin Sigma T6199) in

100 µL droplets on a parafilm-coated humid chamber. We incubated coverslips with the

primary antibody droplets overnight at 4 °C. We then performed 3x 10-minute washes

with ~10 mL blocking buffer in glass petri dishes. After washing, we added a 1:400 dilution

of fluorescent secondary antibody, goat anti-mouse 594 (A11032, ThermoFisher) and

goat anti-rabbit (18772, Sigma), to the humid chamber with new parafilm and incubated

it for 1 hr at room temperature protected from light. We then washed as described after

primary antibody incubation. To label actin, we added phalloidin (P1951, Sigma) labeled

with Alexa-647 (0.094 µM in PBS) and incubated for 20 mins on 100 µL droplets in the

humid chamber. We transferred coverslips to PBS, then set up for imaging by sealing

them to a glass slide with double sided tape and vacuum grease to a glass slide and

adding PBS to the channel. Finally, we imaged the cells in ~1.5-2 µm thick z-stacks using

near-TIRF microscopy, with 1 mW laser power at 640 nm, 561 nm, and 488 nm excitation,

600 ms exposure time, EM gain of 200 using an EMCCD camera of a Nikon Eclipse Ti-E

microscope.

Immunofluorescence: Motor Colocalization

This protocol differs from cytoskeletal imaging in the fixation step; we incubated cells at -

20 °C with 1 mM EGTA (EGT101, Bioshop) in -20 °C methanol, for 8 mins. Subsequent

steps are identical to the cytoskeletal immunofluorescence protocol, except that the

blocking buffer contained 0.2% saponin (4521, Sigma). We first imaged cells for the motor

(kinesin or dynein) and saved their positions on the microscope. Following imaging, we

washed the sample and probed for tubulin while maintaining the slide in the same position

on the microscope, using another colour and shorter incubation times to allow higher

throughput. The protocol was as follows: 1 hr incubation with primary antibody, wash 3x

slide volume with blocking buffer, 45 mins incubation with secondary antibody, wash 3x slide volume with PBS, this sequential antibody staining was done due to the redundancy in antibody species available. The primary antibodies used for experiments were: 1/400 T9026 ms alpha-tubulin, Sigma lot: 000089497, 1/400 ms-kinesin heavy chain MAB1614 lot: 3593684, 1/400 MAB1618 ms-dynein, cytoplasmic lot: 3089121. Secondary antibodies used for experiments were goat anti-mouse 488 (A21236, ThermoFisher) for motors followed by Goat anti-mouse 647 (A21236, Sigma) for microtubules.

Immunoflurescence: Huntingtin Localization

We fixed the cells with prewarmed 4% paraformaldehyde at 37 °C for 10 mins, and subsequently followed the protocol as outlined for cytoskeletal imaging. The huntingtin antibody used was MAB2166 ms-huntingtin, Abcam lot: 3703568.The blocking buffer contained 0.2% saponin (4521, Sigma).

Immunofluorescence: Mitochondria

We fixed cells using prewarmed 4% paraformaldehyde at 37 °C for 20 mins, followed by 3 washes with PBS. Using 50 mM NH_4Cl in PBS, we quenched fixation for 10 mins, and washed again 3 times with PBS. We permeabilized cells using 0.1% TritonX-100 in PBS for 10 mins with gentle rocking, followed by blocking with 10% BSA (ALB001, Bioshop) for 30-45 mins with gentle rocking. We diluted the primary antibody 1/500 (TOMM40 18409-1-AP, Proteintech, Rosemont, IL, United States) with 5% BSA in PBS and incubated overnight at 4 °C. We washed coverslips 3 times with gentle rocking using the 5% BSA in PBS. We diluted the secondary antibody 1/400 in the same solution and incubated for 1 hr at room temperature. We repeated the washing described after the primary antibody incubation, then mounted the slide using PBS as the imaging medium. We imaged samples using an Olympus IX83 with a 100x objective using a disk scanning unit to obtain high resolution z stacking in a 25 µm range using 0.5 µm slices.

Intracellular Optical Tweezers

We prepared a fibronectin (356008 Corning, Bedford, MA, United States) stock at a concentration of 1.4 mg/mL in water and stored it at -20 °C until use. We coated

fluorescent yellow/green 500 nm beads (F8813, ThermoFisher Scientific) by centrifuging

20 µL of stock bead solution at 20000 rcf for 5 mins, then resuspended the beads in 25

µL of the fibronectin solution. We incubated the beads overnight at 4 °C, then washed

with 40 µL PBS 3 times, centrifuging again at 20000 rcf for 5 mins each. We diluted this

bead solution 1/20 in DMEM complete media, sonicated for 5 mins, and incubated with

the cells for 10 mins at 37 °C. For imaging, we replaced the medium with Lebovitz

supplemented with 10% FBS. Early endosome data was acquired from 15 mins–50 mins

after bead internalization, while lysosomes were from 1 - 2 hours post-internalization. We

identified cargoes moving processively within the cell, positioned the location of the center

of the optical tweezer laser above the motile cargo, and opened the shutter to capture the

bead. We acquired 60 s force traces holding the optical tweezer laser (12 W, 1064 nm)

fixed in the original capture position, while monitoring the position signal and live image

to ensure that the bead remained within one bead radius of the center of the tweezer.

Immediately after this measurement, we calibrated the optical tweezers by applying a 275

s multifrequency sinusoid to calculate the position sensitivity (β), trap stiffness (k_{trap}), and

viscoelastic properties (G', G'', Fig. S2E, F) of the cytoplasm as previously described

(Chaubet et al., 2020).

Immunofluorescence Analysis: Actin, Microtubules, Golgi, Huntingtin

For each z-stack, we chose the sharpest image for each channel for qualitative analysis

and observation as overlays.

Mitochondrial Immunofluorescence Analysis

We determined the range of z-values with samples in focus, and only used those slices

for image analysis. We processed the images in Fiji with the white top hat 3D

morphological filter, using a radius of 5 in x and y, and 2 in z. We then set an automatic

threshold and converted the thresholded image to a mask. Next, we ran the analyze

particles function to determine the shape, area, and number of mitochondria in each

plane, repeated for all in focus z planes, and added all the values for each z plane

together. We plotted the results from all cells using MATLAB.

Golgi Dispersal Analysis

Using ImageJ, we observed each of the Golgi images using the same linear LUT (minimum = 145 a.u. and maximum = 1885 a.u.) and then categorized the Golgi into organized, dispersed, or undetermined. Comparing the results from multiple experiments using this binary categorization and excluding the undetermined data, we determined the percentage of organized and dispersed Golgi observed in each cell type.

Motility Assay Analysis- Tracking

We used the Fiji plugin TrackMate (Tinevez et al., 2017) to generate tracks from the images for subsequent analysis. The settings used in TrackMate highly influence the results and are therefore shown in the tables below (Table 1 and 2). We chose the parameters by comparing the maximum intensity projection of all frames with the generated tracks and trying multiple values that would be considered logical for these cargos.

Table 1. TrackMate analysis settings used in the analysis of lysosomal cargoes using LysoTracker™.

Parameter	Value
Estimated blob diameter	1 µm
Intensity threshold	400
Linking maximum distance	0.5 µm
Gap-closing maximum distance	0.5 µm
Gap-closing maximum frame gap	2

Table 2. TrackMate analysis settings used in the analysis of early endosomal cargoes using Rab5-eGFP transfected cells.

Parameter	Value
Estimated blob diameter	0.8 µm
Intensity threshold	300
Linking maximum distance	0.5 µm
Gap-closing maximum distance	0.5 µm
Gap-closing maximum frame gap	2

Motility Assay Analysis-Quantitation

To analyze the raw TrackMate data, we wrote custom MATLAB codes, now available on GitHub: https://github.com/hendricks-lab.

We calculated time averaged mean squared displacement according to the following formula:

$$MSD = \frac{1}{\tau - t} \sum_{t=0}^{\tau-t} (x_{t+1} - x_t)^2$$

We calculated α values by fitting a line to the log-log plot of the mean squared displacement for time delays from 120 ms-5 s. We obtained radius of gyration values using the following formula:

$$R_g^2 = \frac{1}{N} \sum_{k=1}^{N} (r_k - r_{mean})^2$$

Optical Tweezers: Force Trace Analysis

We calculated the trap stiffness and position sensitivity from the calibration data as previously described (Chaubet et al., 2020). Using the calibration data, we converted 60 s position traces to force traces (Fig. S2D) and calculated the viscoelastic moduli in each condition (Fig. S2B, C). We observed brightfield images of the position of the trap and the cell's nucleus to define the directionality of the cargo's movement. Any force event that had a maximum off-axis force, a force in the direction perpendicular to the defined microtubule axis, above 75% of the median on-axis forces, forces along the direction of the microtubule, were excluded. Stall forces were binned into 1 pN size bins and counted as stalls if the duration that they were above 0.5 pN duration was longer than 0.1 s. We calculated binding rates from the force traces by identifying the times that the cargoes unbound from the microtubules after a stall event and determining the time between unbinding and rebinding to the microtubule. We calculated force-dependent unbinding rates from the stall force event before a cargo became detached and categorized them by the direction the cargo was travelling immediately prior to detachment, as previously described (Berger et al., 2019). We performed an exponential fit to the data and presented

670   it in Table S1 following the equation below with U as the unbinding rate, U_0 as the force-
671   independent unbinding rate, F as the force, and F_d as the detachment force.
672

$$U = U_0 e^{\frac{F}{F_d}}$$

Statistical Analysis

We analysed all pairwise histogram comparisons and cumulative distribution functions
using the Kolmogorov-Smirnov (ks) test. We used Student's two-tailed t-test for all
pairwise analyses of violin plots. For optical tweezers stall force distribution analyses, in
addition to the ks test we performed bootstrapping. We took 1000 bootstrap means of
samples equivalent to the number of datapoints for the condition being tested. We then
calculated 95% confidence intervals from the bootstrap mean distribution using the
quantile function in MATLAB. To calculate p-values for bootstrap samples (not reported
in Fig. 3), we subtracted the control from the S421D condition for each of the 1000
bootstrap means for the conditions to be compared and plotted the distribution (Fig. S2E).
The p-value was determined by the fraction of data that overlapped with the 0 value, if
this fraction was lower than 0.05, the difference between the values was considered
statistically significant.

ACKNOWLEDGEMENTS

689   We thank Frederic Saudou (U. Grenoble) for generously sharing huntingtin constructs,
690   Amine Kamen (McGill U.) for guidance on CRISPR/Cas9, and the members of the
691   Hendricks lab for thoughtful comments on the manuscript.

COMPETING INTERESTS

693   The authors report no competing interests.

FUNDING

695   ENPP is supported by a CGS-M fellowship from the Natural Sciences and Engineering
696   Council of Canada. AGH is supported by the Canadian Institutes for Health Research
697   (PJT-159490).

REFERENCES

Al-Haddad, A., Shonn, M. A., Redlich, B., Blocker, A., Burkhardt, J. K., Yu, H.,
Hammer, J. A., Weiss, D. G., Steffen, W., Griffiths, G., et al. (2001). Myosin Va
bound to phagosomes binds to F-actin and delays microtubule-dependent motility.
Mol. Biol. Cell **12**, 2742–2755.

Bentley, M., Decker, H., Luisi, J. and Banker, G. (2015). A novel assay reveals
preferential binding between rabs, kinesins, and specific endosomal
subpopulations. *J. Cell Biol.* **208**, 273–281.

Berger, F., Klumpp, S. and Lipowsky, R. (2019). Force-Dependent Unbinding Rate of
Molecular Motors from Stationary Optical Trap Data. *Nano Lett.* **19**, 2598–2602.

Blehm, B. H., Schroer, T. A., Trybus, K. M., Chemla, Y. R. and Selvin, P. R. (2013).
In vivo optical trapping indicates kinesin's stall force is reduced by dynein during
intracellular transport. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 3381–3386.

Blocker, A., Severin, F. F., Burkhardt, J. K., Bingham, J. B., Yu, H., Olivo, J. C.,
Schroer, T. A., Hyman, A. A. and Griffiths, G. (1997). Molecular requirements for
bi-directional movement of phagosomes along microtubules. *J. Cell Biol.* **137**, 113–
129.

Burkhardt, J. K., Echeverri, C. J., Nilsson, T. and Vallee, R. B. (1997).
Overexpression of the dynamitin (p50) subunit of the dynactin complex disrupts
dynein-dependent maintenance of membrane organelle distribution. *J. Cell Biol.*

Cason, S. E., Carman, P., Van Duyne, C., Goldsmith, J., Dominguez, R. and
Holzbaur, E. L. F. (2021). Dynein effectors regulate autophagosome motility
Sequential dynein effectors regulate axonal autophagosome motility in a
maturation-dependent pathway. *J. Chem. Biol.*

Caviston, J. P. and Holzbaur, E. L. F. (2009). Huntingtin as an essential integrator of
intracellular vesicular trafficking. *Trends Cell Biol.*

Caviston, J. P., Ross, J. L., Antony, S. M., Tokito, M. and Holzbaur, E. L. F. (2007).
Huntingtin facilitates dynein/dynactin-mediated vesicle transport. *Proc. Natl. Acad.*
Sci. U. S. A. **104**, 10045–10050.

Caviston, J. P., Zajac, A. L., Tokito, M. and Holzbaur, E. L. F. (2011). Huntingtin
coordinates the dynein-mediated dynamic positioning of endosomes and
lysosomes. *Mol. Biol. Cell.*

Chaubet, L., Chaudhary, A. R., Heris, H. K., Ehrlicher, A. J. and Hendricks, A. G.
(2020). Dynamic actin cross-linking governs the cytoplasm's transition to fluid-like
behavior. *Mol. Biol. Cell* **31**, 1744–1752.

Chaudhary, A. R., Berger, F., Berger, C. L. and Hendricks, A. G. (2018). Tau directs
intracellular trafficking by regulating the forces exerted by kinesin and dynein
teams. *Traffic* **19**, 111–121.

Chung, P. J., Song, C., Deek, J., Miller, H. P., Li, Y., Choi, M. C., Wilson, L.,
Feinstein, S. C. and Safinya, C. R. (2016). Tau mediates microtubule bundle
architectures mimicking fascicles of microtubules found in the axon initial segment.
Nat. Commun. **7**, 1–9.

Colin, E., Zala, D., Liot, G., Rangone, H., Borrell-Pagès, M., Li, X. J., Saudou, F.
and Humbert, S. (2008). Huntingtin phosphorylation acts as a molecular switch for
anterograde/retrograde transport in neurons. *EMBO J.*

Corthésy-Theulaz, I., Pauloin, A. and Pfeffer, S. R. (1992). Cytoplasmic dynein
participates in the centrosomal localization of the Golgi complex. *J. Cell Biol.* **118**,
1333–1345.

Dambournet, D., Sochacki, K. A., Cheng, A. T., Akamatsu, M., Taraska, J. W.,
Hockemeyer, D. and Drubin, D. G. (2018). Genome-edited human stem cells
expressing fluorescently labeled endocytic markers allow quantitative analysis of
clathrin-mediated endocytosis during differentiation. *J. Cell Biol.* **217**, 3301–3311.

Davies, S. W., Turmaine, M., Cozens, B. A., DiFiglia, M., Sharp, A. H., Ross, C. A.,
Scherzinger, E., Wanker, E. E., Mangiarini, L. and Bates, G. P. (1997).
Formation of neuronal intranuclear inclusions underlies the neurological dysfunction
in mice transgenic for the HD mutation. *Cell*.

Dunah, A. W., Jeong, H., Griffin, A., Kim, Y. M., Standaert, D. G., Hersch, S. M.,
Mouradian, M. M., Young, A. B., Tanese, N. and Krainc, D. (2002). Sp1 and
TAFII130 transcriptional activity disrupted in early Huntington's disease. *Science*
(80-.). **296**, 2238–2243.

Duyao, M. P., Auerbach, A. B., Ryan, A., Persichetti, F., Barnes, G. T., McNeil, S.
M., Ge, P., Vonsattel, J. P., Gusella, J. F., Joyner, A. L., et al. (1995).
Inactivation of the mouse huntington's disease gene homolog Hdh. *Science (80-.).*

269, 407–410.

Ebneth, A., Godemann, R., Stamer, K., Illenberger, S., Trinczek, B., Mandelkow, E. M. and Mandelkow, E. (1998). Overexpression of tau protein inhibits kinesin-dependent trafficking of vesicles, mitochondria, and endoplasmic reticulum: Implications for Alzheimer's disease. *J. Cell Biol.* **143**, 777–794.

Encalada, S. E., Szpankowski, L., Xia, C. H. and Goldstein, L. S. B. (2011). Stable kinesin and dynein assemblies drive the axonal transport of mammalian prion protein vesicles. *Cell* **144**, 551–565.

Engelender, S. (1997). Huntingtin-associated protein 1 (HAP1) interacts with the p150Glued subunit of dynactin. *Hum. Mol. Genet.*

Fu, M. M. and Holzbaur, E. L. F. (2013). JIP1 regulates the directionality of APP axonal transport by coordinating kinesin and dynein motors. *J. Cell Biol.*

Fu, M. meng and Holzbaur, E. L. F. (2014). Integrated regulation of motor-driven organelle transport by scaffolding proteins. *Trends Cell Biol.*

Futter, M., Diekmann, H., Schoenmakers, E., Sadiq, O., Chatterjee, K. and Rubinsztein, D. C. (2009). Wild-type but not mutant huntingtin modulates the transcriptional activity of liver X receptors. *J. Med. Genet.* **46**, 438–446.

Gauthier, L. R., Charrin, B. C., Borrell-Pagès, M., Dompierre, J. P., Rangone, H., Cordelières, F. P., De Mey, J., MacDonald, M. E., Leßmann, V., Humbert, S., et al. (2004). Huntingtin controls neurotrophic support and survival of neurons by enhancing BDNF vesicular transport along microtubules. *Cell* **118**, 127–138.

Gil, J. M. and Rego, A. C. (2008). Mechanisms of neurodegeneration in Huntington's disease. *Eur. J. Neurosci.*

Goldman, A., Harper, S. and Speicher, D. W. (2016). Detection of proteins on blot membranes - Current Protocols in Protein Science. *Curr. Protoc. Protein Sci.* 1–6.

Guo, Q., Huang, B., Cheng, J., Seefelder, M., Engler, T., Pfeifer, G., Oeckl, P., Otto, M., Moser, F., Maurer, M., et al. (2018). The cryo-electron microscopy structure of huntingtin. **555**, 117–120.

Hendricks, A. G., Perlson, E., Ross, J. L., Schroeder, H. W., Tokito, M. and Holzbaur, E. L. F. (2010). Motor Coordination via a Tug-of-War Mechanism Drives Bidirectional Vesicle Transport. *Curr. Biol.*

Hilditch-Maguire, P. (2000). Huntingtin: an iron-regulated protein essential for normal
nuclear and perinuclear organelles. *Hum. Mol. Genet.*

Holbert, S., Denghien, I., Kiechle, T., Rosenblatt, A., Wellington, C., Hayden, M. R.,
Margolis, R. L., Ross, C. A., Dausset, J., Ferrante, R. J., et al. (2001). The Gln-
Ala repeat transcriptional activator CA150 interacts with huntingtin:
Neuropathologic and genetic evidence for a role in Huntington's disease
pathogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 1811–1816.

Humbert, S., Bryson, E. A., Cordelières, F. P., Connors, N. C., Datta, S. R.,
Finkbeiner, S., Greenberg, M. E. and Saudou, F. (2002). The IGF-1/Akt pathway
is neuroprotective in Huntington's disease and involves huntingtin phosphorylation
by Akt. *Dev. Cell.*

Jimenez, A., Friedl, K. and Leterrier, C. (2020). About samples, giving examples:
Optimized Single Molecule Localization Microscopy. *Methods.*

Leavitt, B. R., Van Raamsdonk, J. M., Shehadeh, J., Fernandes, H., Murphy, Z.,
Graham, R. K., Wellington, C. L., Raymond, L. A. and Hayden, M. R. (2006).
Wild-type huntingtin protects neurons from excitotoxicity. *J. Neurochem.* **96**, 1121–
1129.

Lemarié, F. L., Caron, N. S., Sanders, S. S., Schmidt, M. E., Nguyen, Y. T. N., Ko,
S., Xu, X., Pouladi, M. A., Martin, D. D. O. and Hayden, M. R. (2021). Rescue of
aberrant huntingtin palmitoylation ameliorates mutant huntingtin-induced toxicity.
Neurobiol. Dis. **158**,.

Li, S. H., Schilling, G., Young, W. S., Li, X., Margolis, R. L., Stine, O. C., Wagster,
M. V., Abbott, M. H., Franz, M. L., Ranen, N. G., et al. (1993). Huntington's
disease gene (IT15) is widely expressed in human and rat tissues. *Neuron* **11**,
985–993.

Li, S. H., Gutekunst, C. A., Hersch, S. M. and Li, X. J. (1998). Interaction of
Huntingtin-associated protein with dynactin P150(Glued). *J. Neurosci.*

Li, S.-H., Cheng, A. L., Zhou, H., Lam, S., Rao, M., Li, H. and Li, X.-J. (2002).
Interaction of Huntington Disease Protein with Transcriptional Activator Sp1. *Mol.*
Cell. Biol. **22**, 1277–1287.

Liot, G., Zala, D., Pla, P., Mottet, G., Piel, M. and Saudou, F. (2013). Mutant

huntingtin alters retrograde transport of TrkB receptors in striatal dendrites. *J.*

Neurosci. **33**, 6298–6309.

Lo Sardo, V., Zuccato, C., Gaudenzi, G., Vitali, B., Ramos, C., Tartari, M., Myre, M.

A., Walker, J. A., Pistocchi, A., Conti, L., et al. (2012). An evolutionary recent

neuroepithelial cell adhesion function of huntingtin implicates ADAM10-Ncadherin.

Nat. Neurosci. **15**, 713–721.

Marcora, E., Gowan, K. and Lee, J. E. (2003). Stimulation of NeuroD activity by

huntingtin and huntingtin-associated proteins HAP1 and MLK2. *Proc. Natl. Acad.*

Sci. U. S. A. **100**, 9578–9583.

McGuire, J. R., Rong, J., Li, S. H. and Li, X. J. (2006). Interaction of Huntingtin-

associated protein-1 with kinesin light chain: Implications in intracellular trafficking

in neurons. *J. Biol. Chem.* **281**, 3552–3559.

McKenney, R. J., Huynh, W., Tanenbaum, M. E., Bhabha, G. and Vale, R. D. (2014).

Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes.

Science (80-.).

Mogre, S. S., Christensen, J. R., Reck-Peterson, S. L. and Koslover, E. F. (2021).

Optimizing microtubule arrangements for rapid cargo capture. *Biophys. J.* **120**,

4918–4931.

Nasir, J., Floresco, S. B., O'Kusky, J. R., Diewert, V. M., Richman, J. M., Zeisler, J.,

Borowski, A., Marth, J. D., Phillips, A. G. and Hayden, M. R. (1995). Targeted

disruption of the Huntington's disease gene results in embryonic lethality and

behavioral and morphological changes in heterozygotes. *Cell* **81**, 811–823.

Pardo, R., Molina-Calavita, M., Poizat, G., Keryer, G., Humbert, S. and Saudou, F.

(2010). PARIS-htt: An optimised expression platform to study huntingtin reveals

functional domains required for vesicular trafficking. *Mol. Brain* **3**, 1–17.

Reed, N. A., Cai, D., Blasius, T. L., Jih, G. T., Meyhofer, E., Gaertig, J. and Verhey,

K. J. (2006). Microtubule Acetylation Promotes Kinesin-1 Binding and Transport.

Curr. Biol.

Sahlender, D. A., Roberts, R. C., Arden, S. D., Spudich, G., Taylor, M. J., Luzio, J.

P., Kendrick-Jones, J. and Buss, F. (2005). Optineurin links myosin VI to the

Golgi complex and is involved in Golgi organization and exocytosis. *J. Cell Biol.*

Sanjana, N. E., Shalem, O. and Zhang, F. (2014). Improved vectors and genome-wide

libraries for CRISPR screening HHS Public Access Supplementary Material. *Nat*

Methods **11**, 783–784.

Saudou, F. and Humbert, S. (2016). The Biology of Huntingtin. *Neuron*.

Schlager, M. A., Hoang, H. T., Urnavicius, L., Bullock, S. L. and Carter, A. P.

(2014). In vitro reconstitution of a highly processive recombinant human dynein

complex . *EMBO J.* **33**, 1855–1868.

Schmitt, I., Bächner, D., Megow, D., Henkiein, P., Hamelster, H., Epplen, J. T. and

Riess, O. (1995). Expression of the huntington disease gene in rodents: Cloning

the rat homologue and evidence for downregulation in non-neuronal tissues during

development. *Hum. Mol. Genet.* **4**, 1173–1182.

Sharp, A. H., Loev, S. J., Schilling, G., Li, S. H., Li, X. J., Bao, J., Wagster, M. V.,

Kotzuk, J. A., Steiner, J. P., Lo, A., et al. (1995). Widespread expression of

Huntington's disease gene (IT15) protein product. *Neuron* **14**, 1065–1074.

Steffan, J. S., Agrawal, N., Pallos, J., Rockabrand, E., Trotman, L. C., Slepko, N.,

Illes, K., Lukacsovich, T., Zhu, Y. Z., Cattaneo, E., et al. (2004). SUMO

Modification of Huntingtin and Huntington's Disease Pathology. *Science (80-.).*

304, 100–104.

Sun, T., Li, Y., Li, T., Ma, H., Guo, Y., Jiang, X., Hou, M., Huang, S. and Chen, Z.

(2017). JIP1 and JIP3 cooperate to mediate TrkB anterograde axonal transport by

activating kinesin-1. *Cell. Mol. Life Sci.*

Thion, M. S., McGuire, J. R., Sousa, C. M., Fuhrmann, L., Fitamant, J., Leboucher,

S., Vacher, S., Du Montcel, S. T., Bièche, I., Bernet, A., et al. (2015). Unraveling

the Role of Huntingtin in Breast Cancer Metastasis. *J. Natl. Cancer Inst.* **107**, 1–13.

Tinevez, J. Y., Perry, N., Schindelin, J., Hoopes, G. M., Reynolds, G. D., Laplantine,

E., Bednarek, S. Y., Shorte, S. L. and Eliceiri, K. W. (2017). TrackMate: An open

and extensible platform for single-particle tracking. *Methods* **115**, 80–90.

Twelvetrees, A. E., Yuen, E. Y., Arancibia-Carcamo, I. L., MacAskill, A. F.,

Rostaing, P., Lumb, M. J., Humbert, S., Triller, A., Saudou, F., Yan, Z., et al.

(2010). Delivery of GABAARs to Synapses Is Mediated by HAP1-KIF5 and

Disrupted by Mutant Huntingtin. *Neuron* **65**, 53–65.

Wanker, E. E., Ast, A., Schindler, F., Trepte, P. and Schnoegl, S. (2019). The

pathobiology of perturbed mutant huntingtin protein–protein interactions in

Huntington's disease. *J. Neurochem.* **151**, 507–519.

Wong, Y. C. and Holzbaur, E. L. F. (2014). The regulation of autophagosome

dynamics by huntingtin and HAP1 is disrupted by expression of mutant huntingtin,

leading to defective cargo degradation. *J. Neurosci.*

Xu, X., Ng, B., Sim, B., Radulescu, C. I., Yusof, N. A. B. M., Goh, W. I., Lin, S., Lim,

J. S. Y., Cha, Y., Kusko, R., et al. (2020). pS421 huntingtin modulates

mitochondrial phenotypes and confers neuroprotection in an HD hiPSC model. *Cell*

Death Dis.

Yanai, A., Huang, K., Kang, R., Singaraja, R. R., Arstikaitis, P., Gan, L., Orban, P.

C., Mullard, A., Cowan, C. M., Raymond, L. A., et al. (2006). Palmitoylation of

huntingtin by HIP14is essential for its trafficking and function. *Nat. Neurosci.* **9**,

824–831.

Zala, D., Colin, E., Rangone, H., Liot, G., Humbert, S. and Saudou, F. (2008).

Phosphorylation of mutant huntingtin at S421 restores anterograde and retrograde

transport in neurons. *Hum. Mol. Genet.*

Zeitlin, S., Liu, J., Chapman, D. L. and Papaioannou, V. E. (1995). article nullizygous

for the Huntington ' s disease gene homo logue. **11**, 155–163.

Zuccato, C., Tartari, M., Crotti, A., Goffredo, D., Valenza, M., Conti, L., Cataudella,

T., Leavitt, B. R., Hayden, M. R., Timmusk, T., et al. (2003). Huntingtin interacts

with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes.

Nat. Genet. **35**, 76–83.

FIGURE LEGENDS

Fig. 1. S421D HEK293T cell line does not affect cytoskeletal organization, huntingtin localization, or dynein function. (A) Huntingtin protein sequence schematic demonstrating the structured domains (HEAT repeats and other α-helical regions, orange), unstructured domains (blue), S421 phosphosite, polyglutamine repeat region (PolyQ), and proline rich domain (PrD). (B) CRISPR phosphomimetic mutant design guide RNA (gRNA), and ssODN homologous recombination template. (C) Chromatogram of the positive clone containing the phosphomimetic mutation S421D. (D) Immunostaining of actin (top) and microtubules (bottom) indicating no difference between WT (left) and S421D (right) HEK293T cells. Note immunofluorescence images are inverted such that the protein of interest is dark, and the background is white, and all scale bars in the figure are 20 μm. (E) In both WT and S421D HEK293T cells, the Golgi apparatus appeared compact (organized) in ~70% of cells and dispersed in ~30% of cells. WT data for actin, microtubules, and Golgi apparatus staining was from 14 experiments, 84 cells, and S421D data from 11 experiments, 69 cells. (F) Immunofluorescence of huntingtin protein localization in WT (left) compared to S421D (right) HEK293T cells. Immunofluorescence of huntingtin localization for WT cells from 3 experiments, 43 cells and for S421D from 2 experiments, 22 cells. (G) WT and S421D huntingtin endogenous expression levels (top left) detected by Western blot with the raw intensity data from three independent samples (bottom left), compared to levels when overexpressing full length mCherry-WT or mCherry S421D huntingtin (top right) and the fold change relative to native expression levels for three independent samples (bottom right). Samples collected on the same date are connected by a line. (H) Mitochondrial morphology of WT (left) and S421D (right) HEK293T cells. (I) Mitochondrial area of WT (grey) compared to S421D (blue) cells, $p=6.59 \times 10^{-8}$ by Kolmogorov-Smirnov (ks) test. (J) Number of mitochondria per cell in WT (grey) compared to S421D (blue), $p=6.14 \times 10^{-7}$ by ks test. Mitochondrial analysis for WT from 7 experiments, 88 cells, and S421D from 7 experiments, 94 cells.

Fig. 2. Huntingtin S421D mutation increases outward motility of early endosomes and lysosomes while overexpression affects all parameters. (A, B) Maximum intensity projection of early endosomes, labelled with Rab5-eGFP, and lysosomes, labelled with Lysotracker, for endogenous WT and S421D huntingtin expressing cells. All scale bars are 20 µm. (C, D) Sample trajectories for early endosomes (C) and lysosomes (D) in both WT and S421D huntingtin endogenously expressing cells, with their inward (in) or outward (out) directionality indicated by the arrows in the center. Different shades of colours were assigned randomly to facilitate visualization of overlapping segments. (E, G) Early endosome (E) and lysosome (G) processivity per cell, measured by α, the slope of the log-log plot of their mean-squared displacement. Means are represented with a filled line, while medians are dashed lines. (F. H) Radius of gyration (R_g) per cell of early endosomes (F) and lysosomes (H). R_g is a measure of the radius that contains half of the datapoints in the trajectory centered around its average position. (I, J) Early endosome (I) and

lysosome (J) directional bias per cell of processive trajectories toward plus ends of microtubules (outwards),
the remaining fraction of processive trajectories move inwards. Overexpression conditions represented by
+WThtt or +S421Dhtt. Directional bias is determined by fraction of time particles are moving outwards
compared to inwards and averaged for each cell. The number of experiments, cells, and trajectories for
each condition are the following; WT early endosomes: 65 cells, 41994 trajectories, 6 experiments; S421D
early endosomes: 28 cells, 25476 trajectories, 3 experiments; WT+WThtt early endosomes: 28 cells, 17424
trajectories, 5 experiments, WT+S421Dhtt early endosomes: 39 cells, 22854 trajectories, 6 experiments;
WT lysosomes: 55 cells, 25247 trajectories, 5 experiments; S421D lysosomes: 38 cells, 12670 trajectories,
4 experiments; WT+WThtt lysosomes: 29 cells, 7558 trajectories, 5 experiments; WT+S421Dhtt lysosomes:
28 cells, 7031 trajectories, 6 experiments. Statistical significance was determined via Student's two-tailed
t-test and p-values are indicated on each plot where applicable.
959

Fig. 3. Force trace analysis demonstrates S421D huntingtin induces higher binding rates, stronger binding, and higher forces in both directions for early endosomes and lysosomes. (A) Schematic of the optical tweezers assay including the endocytosed bead (green) inside the optical trap (red) along with a brightfield image of a cell during the experiment (top right). The scale bar is 20 μm and the arrowhead indicates the position of the optical trap during the experiment. (B) Sample force traces for early endosomes (early) and lysosomes in WT and S421Dhtt cells. The horizontal line indicates the zero axis, while positive values indicate outward forces and negative values indicate inward forces. (C, E) The fraction of early endosome or lysosome stall events at a given force with a minimum stall duration of 0.1 s, separated into 1 pN bins with the mean and 95% confidence intervals of bootstrap sampled means and p-values for each condition are shown above (for outward) or below (for inward) the distribution. The colours in (C-J) follow the colour legend from (B). (D, F) Force-dependent unbinding rates in both outward and inward directions (filled circles) for early endosomes or lysosomes fit with a single exponential (solid line). (G-J) Binding rates

for each stall event in early endosomes and lysosomes of WT and S421D in both directions in (G, I)
including the cumulative distribution function (CDF) for each condition in (H, J). Statistical significance by
ks test is indicated by the p-value in the figure panel where applicable. All results from optical tweezers
data for WT early endosomes from 10 cells, 5 experiments; S421D early endosomes: 16 cells, 8
experiments; WT lysosomes: 12 cells, 5 experiments, S421D lysosomes: 11 cells, 4 experiments. Statistical
significance was determined by ks test for binding rates and force distributions and p-values are indicated
where applicable.
981

982
Fig. 4. S421D huntingtin increases kinesin and dynein association to extracted microtubules. (A, B)
Inverted immunofluorescence images demonstrating colocalization (bottom) kinesin heavy chain (kinesin-
1, (A), top) or dynein intermediate chain (dynein, (B), top) on extracted microtubules (middle) in WT and
S421D huntingtin cells. In the merge channel, motors (kinesin-1 or dynein) are indicated in magenta and
microtubules in black. All scale bars are 20 µm. (C-F) Colocalization plots were created from plotting
intensity values of each pixel of the microtubule and motor (kinesin or dynein) channels of all images.

Differences between kinesin-1 WT and S421D correlation coefficients resulted in p=0.4748, while dynein
WT compared to S421D correlation gave p=0.0533 by a two-tailed Student's t-test. Data for WT kinesin-1
from 17 cells, 3 experiments, S421D kinesin-1: 15 cells, 2 experiment, WT dynein: 32 cells, 4 experiments,
and S421D dynein: 32 cells, 4 experiments.
993

994
Fig. 5. Model of huntingtin S421 phosphorylation and expression level mediating regulation of early
endosome and lysosome transport. (A) Huntingtin S421 phosphorylation (S421-phos) increases activity
of kinesin and dynein motor proteins on early endosomes, such that the predominantly inward motility shifts
to being more outward. (B) Lysosomes move bidirectionally. When kinesin and dynein activity are enhanced
by huntingtin phosphorylation, the movement remains bidirectional, but the displacement of trajectories
increases. (C) Overexpression sequesters motors and adaptors or disrupts vesicle-bound huntingtin-
adaptor-motor complexes regardless of cargo identity or huntingtin phosphorylation state.
1002

SUPPLEMENTARY MATERIALS

Table S1. Calculated parameters from exponential fits for unbinding rates of early endosomes and lysosomes.

Condition	U_0	F_d (pN)
Early endosome WT outward	2.1×10^{-7}	0.42
Early endosome S421D outward	0.31	3.2
Early endosome WT inward	7.2×10^{-7}	0.43
Early endosome S421D inward	0.31	6.3
Lysosome WT outward	0.49	12.1
Lysosome S421D outward	0.24	6.6
Lysosome WT inward	1.9	14.6
Lysosome S421D inward	0.77	10.8

Fig. S1. Cytoskeletal disruption affects cargo motility. (A) Actin cytoskeleton labelled with SiR actin in S421D HEK293T cells treated with 100 nM-1 μM concentrations of latrunculin A (latA). Scale bars for all images are 20 μm, all images inverted. (B) Per cell α values of Rab5-eGFP labelled early endosomes. Means are represented with a filled line, while medians are dashed lines. (C) Radius of gyration (R_g) per cell is presented for early endosomes (C). (D) Directional bias per cell shown by fraction of time processive

runs moving outward for Rab5-eGFP early endosomes. Note: control WT and S421D data for latrunculin A
experiments is the same data reported in Fig. 2. (E, F) Maximum projections of the same cell with and
without nocodazole (NZ) for early endosome and lysosome analysis. (G-J), Radius of gyration and alpha
values for early endosomes and lysosomes before and after nocodazole treatment. (B-D) WT control data
from 7 independent experiments (41994 trajectories, 64 cells), S421D data from 3 independent experiments
(25476 trajectories, 28 cells), WT latA data from 3 independent experiments (17461 trajectories, 22 cells);
S421D latA data from 5 independent experiments (19879 trajectories, 30 cells). (G, H) WT control data
from 4 experiments, 22 cells, 6109 trajectories and WT+nocodazole data is from 4 experiments, 23 cells,
and 7094 trajectories. (I, J) Contains control data from 4 experiments, 15 cells, 17073 trajectories, and
nocodazole treated data from 4 experiments, 23 cells, and 6639 trajectories. Statistical significance of
$p < 0.05$ by Student's two sample t-test indicated in the figure where applicable.
1025

Fig. S2. Force measurement analysis approach, additional force traces, and viscoelastic properties
of WT and S421D huntingtin cells. (A) Brightfield image of a WT and S421D cell containing a trapped
phagocytosed 500 nm bead (red arrow) during measurement of early endosome (early) or lysosome forces.
(B, C) Viscous (G'') and elastic moduli (G') for WT (black) and S421D (teal) early endosomes and WT (blue)
and S421D (green) lysosomes. (D) Additional sample force traces for each condition in optical tweezers
assays, the horizontal line indicates the zero axis. Positive value forces are directed outward and negative
forces are directed inward. (E) Distributions of bootstrap mean differences evaluated for additional statistical
analysis of force distributions for the differences shown on the x axis of each plot in both early (left) and
late (right) endosomes.
1036

1037
1038 **Fig. S3. Western blot transparency for huntingtin blot in Fig. 1.** (A) Full western blot (above 100 kDa)
1039 of the blot presented in Fig. 1G using the anti-huntingtin antibody. Ladder is indicated by L with molecular
1040 weights to the right of the blot. (B) Coomassie staining of the entire membrane post-antibody staining.
1041
1042