Sucrose metabolism and candidate genes during catkin fibers development in poplar

Running title: Sucrose metabolism in poplar catkin fiber.

Xiong Yang[1, 2 †], Tianyun Zhao[1, 2 †], Pian Rao[1, 2], Ning Yang[1], Guolei Li[1], Liming Jia[1], Xinmin An[2, 3], Zhong Chen[1, 2*]

[1]Key Laboratory of Silviculture and Conservation of the Ministry of Education, National Energy R&D Center for Non-food Biomass, College of Forestry, Beijing Forestry University, Beijing 100083, China;

[2]Beijing Advanced Innovation Center for Tree Breeding by Molecular Design, Beijing Forestry University, Beijing 100083, China;

[3]National Engineering Research Center of Tree Breeding and Ecological Restoration, College of Biological Sciences and Biotechnology, Beijing Forestry University, Beijing 100083, China

[†] X. Y. and T. Z. contributed equally to this work.

E-mail: xiongyang@bjfu.edu.cn; tianyunzhao@126.com; cicipian@gmail.com; ningyang1124@163.com; glli226@163.com; jlm@bjfu.edu.cn; anxinmin@bjfu.edu.cn; zhongchen@bjfu.edu.cn

*Author for Correspondence

Zhong Chen; E-mail: zhongchen@bjfu.edu.cn; Tel.: +86-13020086124

Address: No.35 Qinghua East Road, Haidian District, P. O. Box 407, Beijing Forestry University, Beijing 100083, China

Conflicts of Interests: The authors declare no conflict of interest.

Abstract

Poplar is an important tree species for ecological protection, wood production, landscaping and urban greening; it has been widely planted worldwide. However, the catkin fibers produced by female poplars can cause environmental pollution, human health risks and safety hazards during spring. This study focused on *Populus tomentosa* Carr., and reveals the sucrose metabolism regulatory mechanism of catkin fibers development from morphological, physiological and molecular aspects. And two key genes were selected to demonstrate the exact role. Sucrose degradation plays the dominant role during poplar catkin fibers development. Thereinto, invertase has a function during the early stage and sucrose synthase has a close correlation with the elongation of catkin fibers. The expression patterns revealed that sucrose metabolism-related and *cellulose synthase* (*CesA*) genes play important roles during catkin fiber development. Expression profiles and co-expression networks indicated that the mechanism underlying poplar catkin fiber development is similar, but not identical, to that of cotton fibers. Finally, two key genes, *sucrose synthase 2 (PtoSUS2)* and *vacuolar invertase 3 (PtoVIN3)*, had roles in *Arabidopsis thaliana* trichome density, indicating that sucrose metabolism is important in poplar catkin fibers development. This study is not only helpful for clarifying the mechanism of sucrose regulation during trichome development in perennial woody plants, but also establishes a foundation to solve poplar catkin fiber pollution through biotechnology and genetic engineering methods.

Introduction

Poplar is an ecologically important species and one of the three main tree species used for afforestation in China. The poplar plantation area has reached 7,570,700 hm^2, occupying 13.25% of the total plantation area in 2019. Poplar is not only major contributor to wood production, but also provides resource for eco-environment construction, biomass energy, landscape and urban greening (Sannigrahi et al., 2010). Furthermore, poplar is a crucial model system for research on perennial woody plants due to its small genome, easy propagation and rapid growth (An et al., 2022). In female poplars, the capsules contain seeds and catkin fibers (Fig. 1a), and begin to open as the seeds become mature, usually in early April in northern China (Fig. 1b). Finally, catkin fibers cover the whole catkins (Fig. 1c, d), which is a common phenomenon in poplar (Fig. 1e, f), and the fibers can be carried over long distances with seeds on the wind (Fig. 1g, h). Poplar catkin fibers can survive for > 40 days in the environment depending on the various poplar species and temperatures. The long period during which poplar fibers are suspended in the air can cause serious environmental pollution (Ye et al., 2014), although it can be used as raw materials for bioethanol (Zhang et al., 2018), oil super absorbents (Likon et al., 2013) and textile bulk insulators (Chen and Cluver, 2010). Almost 185,000 female poplar trees that can produce catkins are located in the urban areas of Beijing, the capital of China, and the catkin fibers produced by these trees have extremely harmful effects on human health through inhalation exposure, and they may also affect metal/metalloid concentrations

Fig. 1 Morphological characteristics of capsules and catkin fibers in poplar. a-d

Characteristics of *Populus tomentosa.*. **e, f** Other poplars. **g-h** Environmental pollution

caused by poplar catkin fibers. Scale bar: **a** = 1 cm, **b** = 2 cm.

in the environment (Wan et al., 2020). Ădditionally, the catkin fibers may pose a

safety hazard due to their inflammability. In recent years, studies of sex differentiation

and female flower development in poplar have revealed some key genes such as

type-A cytokinin response regulator (Yang et al., 2021), *FERR-R* and *MSL* (Xue et al.,

2020), *ARABIDOPSIS RESPONSE REGULATOR 17* (Müller et al., 2020),

AGAMOUS and *SEEDSTICK* (Lu et al., 2019), and *LEAFY* (Klocko et al., 2016).

However, the molecular basis of catkin fibers formation and development in poplar is

still unclear. Therefore, analyzing the development process of catkin fibers is helpful

not only for understanding this progress in Salicaceae plants, but also for improving

the urban environment.

Trichomes, which originate from epidermal cells, are distributed on the surfaces

of plants and protect them from biotic and abiotic stresses (Pattanaik et al., 2014).

Many previous studies of trichomes have been conducted in *Arabidopsis thaliana* and

cotton, focusing on epidermal hairs and cotton fibers, respectively (Grebe, 2012;

Naoumkina, 2018). According to morphological characters, trichome development

can be divided into three stages: initiation, extension and maturation. An additional

stage, secondary cell wall (SCW) synthesis, is presumed to occur between extension

and maturation in cotton fibers (Grebe, 2012; Naoumkina, 2018). Similar regulatory

mechanisms were observed during the initiation stage in *A. thaliana* and cotton (Bui

et al., 2019). *GLABROUS1* (*GL1*, a MYB transcription factor [TF] gene),

GLABROUS3 (*GL3*, a bHLH TF gene), *ENHANCER OF GLABRA3* (*EGL3*, a bHLH

TF gene) and *TRANSPARENT TESTA GLABRA1* (*TTG1*, a WD40 TF gene) have been

suggested to play regulatory roles in the trichome development through the induction

of *GLABRA2* (*GL2*, a HD-ZIP IV TF gene) (Rerie et al., 1994; Walker et al., 1999;

Bernhardt et al., 2005; Kirik et al., 2005; Bloomer et al., 2012). In addition,

GhMYB25, another type of *MYB* gene, was found to play a major role in cotton fiber

initiation and extension, and similar functions were found for its homologues,

GhMYB25-LIKE, GhMML3_A12 and *GhMML4_D12* (Wu et al., 2018). *GbML1*,

another HD-ZIP IV TF, was found to have a similar expression pattern to *GhMYB25*

and to interact with *GhMYB25* to control fiber development (Zhang et al., 2010). As a

model of rapid single-cell growth, sucrose metabolism (SM)-related proteins also play

important roles in the process (Ruan, 2014).

Sucrose is an important form of transportable sugar in plants. Plants convert CO_2

into organic carbon through photosynthesis in leaves, and then transport it to

heterotrophic sinks, such as the stem, root and fruit, for energy storage and utilization

(Ruan, 2014). Thus, SM plays a critical role in plant growth and development,

including trichome development. Sucrose can be synthesized by sucrose phosphate

synthase (SPS) and sucrose phosphate phosphatase (SPP), and degraded by sucrose

synthase (SUS) or invertase (INV) in plants (Ruan, 2014). Sugar signals produced by

INV likely influence the cotton fiber initiation through MYB TFs and auxin signals

(Wang et al., 2014). High turgor pressure induced by sucrose degradation via INV and

SUS promotes the rapid elongation of cotton fibers (Wang et al., 2010; Ahmed et al.,

2018). In contrast to INV, uridine diphosphate (UDP)-glucose produced by SUS is a

vital substrate for cellulose synthesis, and it influences cell wall synthesis, especially

of the SCW (Fujii et al., 2010; Ahmed et al., 2018). Additionally, two overexpression

studies showed that SPS has an effect on fiber development (Haigler et al., 2007; Park

et al., 2008),which presumably because SPS can recycle fructose released by SUS to

obtain more sucrose (Martin and Haigler, 2004). Similar to cotton fibers and

Arabidopsis epidermal hairs, in essence, poplar catkin fibers also belong to plant

trichomes. However, compared with the numerous important advances in *Arabidopsis*

and cotton, the research foundation of catkin fibers development in poplar is quite

weak, the relationship between SM and poplar catkin development, and the real role

of many key homologues in poplar are almost unknown.

Chinese white poplar (*Populus tomentosa* Carr.) is a representative species of

poplar, widely planted in northern China. We previously conducted transcriptome

analysis of catkin fibers development in this plant, and found that many Gene

Ontology (GO) terms related to sucrose metabolism were significantly enriched

during this process (Ye et al., 2014). Therefore, in this study, we made histological

observations of catkin fibers development in *P. tomentosa*, and measured the sugar

content and activities of key SM enzymes at various developmental stages. Using the

genomic data of *P. tomentosa* compiled by our group (An et al., 2022), we searched

for SM-related genes and performed detailed fluorescence-based real-time

quantitative polymerase chain reaction (RT-qPCR) analysis. Additionally, based on a

thorough review of the literatures, we summarized the key genes involved in trichome

development in cotton and *A. thaliana*, and then identified their homologues in the

poplar genome database (Table S1). The expression patterns of these potentially

SM-related genes were also analyzed, and a gene co-expression network was

136   constructed. The relationships between SM-related genes and TFs were also analyzed.

137   Finally, two key genes, *PtoSUS2* and *PtoVIN3*, were selected to investigate the

138   function of SM in catkin fibers formation using the model plant *A. thaliana*.

Results

Morphological characteristics of poplar catkin fibers

The morphology of catkin fibers changed rapidly during hydroponic culture (Fig. 2). Histological analysis revealed that no catkin fibers appeared when the branches had been cultured for 0-48 h (Fig. 2a-d). The epidermal cells of funicles initially showed larger nuclei (Fig. 2e, f), in accordance with the endoreduplication stage in *A. thaliana*, indicating the initial stage of development of catkin fibers. Subsequently, the epidermal cells exhibited protrusion and extension from 58 to 120 h, until the catkin fibers finally full filled the ovary (Fig. 2g-j). Thus, according to the morphological characteristics of catkin fibers, we divided the development process into five stages: stage 1 (preparatory stage, cultured for 48 h, Fig. 2d), stage 2 (initial stage, cultured for 58 h, Fig. 2e-f), stage 3 (apophysis stage, cultured for 72 h, Fig. 2g-h), stage 4 (extension stage, cultured for 96 h, Fig. 2i), stage 5 (maturation stage, cultured for 120 h, Fig. 2j). Furthermore, as shown in Fig. 2, catkin fibers were produced from the epidermal cells of funicle and placenta, independent from seeds. Samples from the five stages were used for analyses of enzyme activities and gene expression patterns.

Changes in sugar content and key enzyme activities in poplar catkin fibers

The sucrose content of the catkin of *P. tomentosa* showed a downward trend, indicating that sucrose is gradually consumed during the development of catkin (Fig. 3a). In particular, during the stage 3-4, sucrose was consumed at a dramatic rate when the catkin fibers extended rapidly. Glucose and fructose showed similar trends, which both had higher contents during stage 2-4. The ratio of hexose to sucrose was also

Fig. 2 Histological characteristics of female buds in *P. tomentosa*. a-d Female buds on branches that were hydroponically cultured for 0, 24, 34, 48 h, respectively. No catkin fibers were observed. **e, f** Female buds on branches that were hydroponically cultures for 58 h. **g, h** Female buds on branches that were hydroponically cultured for 72 h. **i** Female buds on branches that were hydroponically cultured for 96 h. **j** Female buds on branches that were hydroponically cultured for 120 h. OW, ovary wall, OV, ovule, FU, funicle, BR, bract, CF, catkin fibers. Scale bar: **a-d, g, i, j** = 500 μm, **e, h** = 200 μm, **f** = 100 μm.

162   analyzed. As shown in Fig. 3b, the ratio increased sharply between the apophysis

Fig. 3 Changes in sugar contents and key enzymes involved in SM during poplar catkin fibers development. Bars represent the standard deviation (SD, n = 3). * P ≤ 0.05.

stage (stage 3) and extension stage (stage 4), and decreased between the extension

stage (stage 4) and maturation stage (stage 5). Changes in the ratio indicated that the

high hexose concentration may play a role in catkin fibers extension. When the catkin

fibers began to extend, most of the sucrose present was degraded into glucose and

fructose, which likely provided greater turgor pressure, thus promoting catkin fibers

elongation. To further explore the role of sucrose in catkin fibers development, we

also determined the activities of key enzymes in SM. Consistent with the results of

sugar content analysis, sucrose degradation was the dominant process.

The activity of CWIN showed no significant fluctuations except at stage 5, when

it decreased as the catkin fibers matured (Fig. 3c). The activities of NIN and VIN

were higher than that of CWIN. VIN activity increased during catkin fibers initiation

and decreased during the catkin fibers maturation, indicating that VIN may play a role

in catkin fibers development (Fig. 3c). Meanwhile, the activity of NIN was high only

during the initial (stage 2) and apophysis stages (stage 3), and appeared to play a role

in the early development of catkin fibers (Fig. 3c). Compared with INV, SUS showed

higher activity during the development of catkin fibers, up to the extension stage

(stage 4); it then decreased during the maturation stage (stage 5), showing an inverse

correlation with sucrose content (Fig. 3e). Similar results were obtained though fitting

analysis. SUS activity showed a significant correlation with the ratio of hexose to

sucrose, indicating that SUS plays a major role in sucrose degradation during the

development of catkin fibers (Fig. 3f). Furthermore, SPS activity was analyzed and

used to represent the sucrose synthesis pathway. As shown in Fig. 3g, SPS exhibited

steady activity throughout catkin fibers development, except during the apophysis

stage (stage 3).

Gene expression patterns related to SM in poplar catkin fibers

The expression of *CWIN* showed a similar trend to enzyme activity. We analyzed

three members of the gene family, *PtoCWIN3*, *PtoCWIN4* and *PtoCWIN5*, during

catkin fibers development. *PtoCWIN3* appeared to play a major role in catkin fibers

development based on its high and variable expression level (Fig. 4a). However,

Pearson correlation analysis showed that *PtoCWIN4* rather than *PtoCWIN3* was

positively correlated with sucrose content, which may be due to the dramatic

decreased in *PtoCWIN3* at the initial stage (Table S4). Two members of the *VIN* gene

family were observed, and both had elevated expression at stage 2-4, coinciding with

the enzyme activity results (Fig. 4b). And both members of the *VIN* gene family had

positive correlations with the concentrations of fructose and glucose (Table S4). For

NIN, nine members were detected during catkin fibers development, including two

low-expression members, *PtoNIN2* and *PtoNIN11* (Fig. 4d, f). According to their

expression patterns, the other members can be divided into two groups. *PtoNIN3* and

PtoNIN4 had higher expression at the apophysis stage (Fig. 4e), while the other five

members peaked at the initial stage (Fig. 4c-d, f-g).

There are seven members in the *SUS* gene family, all of which were detected

during poplar catkin fibers development. Their expression pattern showed a clear

correlation with the phylogenetic tree (Fig. S1c). The first cluster, comprising

PtoSUS1 and *PtoSUS2*, had high expression during catkin fibers development.

Compared with *PtoSUS1*, *PtoSUS2* had higher expression, and both had their highest

Fig. 4 Analysis of SM related gene families in the context of poplar catkin fibers development. a Expression analysis obtained by RT-qPCR for *CWIN* gene family. **b** Expression analysis obtained by RT-qPCR for *VIN* gene family **c-g** Expression analysis obtained by RT-qPCR for the *NIN* gene family. **h-j** Expression analysis obtained by RT-qPCR for the *SUS* gene family. **k, l** Expression analysis obtained by RT-qPCR for *SPS* and *SPP* gene family, respectively. Bars represent the ± SD.

levels during the extension stage (Fig. 4h). The second cluster only contains one

member, *PtoSUS3*, which had low expression but a similar pattern to the first cluster

(Fig. 4i). The other four members belonged to the third cluster, and had high

expression at the apophysis stage (Fig. 4j). The expression levels of most *SUS* gene

members had significant positive correlations with the concentration of hexose (Table

S4).

The expression patterns of *SPS* and *SPP* gene families were analyzed. Five of the

six members of the *SPS* gene family were analyzed, and they all had high expression

at the apophysis stage, in accordance with the results for SPS activity (Fig. 3g and 4k).

All three *SPP* members (*PtoSPP1*, *PtoSPP2* and *PtoSPP3*) were detected, and all had

elevated expression during the apophysis stage (Fig. 4l). The results of both the

enzyme activity and gene expression analyses showed that the sucrose synthesis

pathway might only have a slight effect at the apophysis stage.

***CesA* gene family and other potential genes in poplar catkin fibers development**

Poplar catkin fibers follows a similar development process to cotton fiber, and

cellulose is the main component of cotton fiber (Kim and Triplett, 2007). Thus, we

also checked the expression pattern of the *cellulose synthase* (*CesA*) gene family in

poplar catkin fibers. Thirteen members of the *CesA* gene family were analyzed in this

study, which belonged to five of six groups on the phylogenetic tree, i.e., all except

cluster P2 (Fig. S1d). Among them, *PtoCesA17*, belonging to clade P1, had high

expression at the apophysis stage, while its homologue *PtoCesA7* had low expression

(Fig. 5d). Homologous pairs in clade P3 had similar expression patterns, except

PtoCesA2 and *PtoCesA5* (Fig. 5e-g). *PtoCesA6* and *PtoCesA9* had high expression

Fig. 5 Analysis of the *CesA* gene family and other potential genes in the context of poplar catkin fibers development. a-g Expression analysis based on RT-qPCR for *CesA* gene family. **h, i** Heat map of gene expression obtained by RT-qPCR for TFs-related and other genes, respectively. Bars represent the ± SD.

throughout the process of catkin fibers growth (Fig. 5g). *PtoCesA15* appeared to play

a more important role than *PtoCesA16*, as it had dramatically higher expression

during catkin fibers development (Fig. 5f). Other genes belonging to clades S1, S2

and S3, had low expression in catkin fibers, although their expression increased at the

initial and apophysis stages with the exception of *PtoCesA18* (Fig. 5a-c).

TFs, such as *MYB*, *WD40* and *bHLH*, have been shown to play important roles in

the development of cotton fibers (Bui et al., 2019). Thus, homologues of the factors

that have been identified in cotton were examined in poplar catkin fibers (Table S1).

According to their expression patterns, these homologues can be divided into three

subgroups (Fig. 5h). Among bHLH TFs, *Pto(GL3)-L1* and *2* had highest expression at

the initiation stage. In the *WD40* gene family, *Pto(TTG1)-L1* had high expression at

the initiation stage, while the another WD40 TF *Pto(TTG1)-L2* had high expression at

the extension stage. Members of the MYB gene family may have divergent functions

during catkin fibers development. *Pto(GL1)-L1* and *2* had high expression during

early development of catkin fibers, and *Pto(GL1)-L2* had an additional role in the

extension stage. In contrast, *Pto(GL1)-L3* showed increased expression at the

maturation stage. The homologues of *GhMYB25* all had increased expression at the

maturation stage, except *Pto(GhMYB25)-L1*. *Pto(GhMYB7)-LIKE*, the cotton

homologue of which was speculated to be a lipid transfer protein-related gene, also

had high expression at the maturation stage. Another important TF gene family,

HD-ZIP IV, was also analyzed during poplar catkin fibers development. *Pto(GL2)-L1*

and *2* had high expression at the preparatory and extension stages, and the

homologues of *GhHox3* had functions at the maturation stage. Intriguingly, opposing

results were found between *Pto(GhHD1)-L1* and *Pto(GhHD1)-L2*.

We also examined the expression of other poplar homologues of genes that are

important to the composition of cotton fibers (Table S1). As shown in Fig. 5i, three

subgroups played different roles in the early, middle and late development stages. In

cluster I, *PtoPME-related* genes played an important role in catkin fibers maturation,

which indicted dramatic changes in the composition of cell wall. In cluster II,

PtoCAD6-related and *PtoGH3-related* appeared to be important to the early

development of catkin fibers. During the middle development stage (cluster III), many

genes were involved. Thereinto, *PtoAct1-L1* and *2* and *Pto1,3-β-G-L1* and *5* had high

expression at the initial stage. *PtoGA20x-LIKE, PtoKIS13A1-LIKE, PtoKOR1-LIKE*

and *PtoXTH1-L2* functioned mainly at the apophysis stage, while *PtoHUB2-LIKE,*

Pto1,3-β-G-L3, Pto4CL1-L2, Pto4CL1-L3 and *PtoKT-LIKE* had high expression at

the extension stage.

Co-expression network of genes involved in poplar catkin fibers development

To further explore the relationship between SM and poplar catkin fibers

development, co-expression network analysis was performed using genes related to

SM and homologues of genes that have effects on cotton fiber development. As

shown in Fig. 6a, there were four divergent gene clusters, and SM related genes

mainly occurred in cluster I-III. Thereinto, cluster II was mainly composed of sucrose

synthesis-related genes, with *PtoSPS2* playing a leading role. Notably, *PtoSUS5-7*,

which is a homologue according to the phylogenetic tree, and *PtoGA20x-LIKE* had

close relationships with *SPS* and *SPP*. Different members of the *CesA* gene family

Fig. 6 Co-expression network analysis. a Co-expression network of 108 candidate genes associated with catkin fibers development in *P. tomentosa*. The threshold for significance of the Pearson correlation coefficient was set to ≥ 0.8 (or ≤ − 0.8), p-value ≤ 0.05. Larger and redder the circles, indicate higher connectivity. **b** Co-expression network of SM-related genes and TFs, the threshold for significance of the Pearson correlation coefficient was set to ≥ 0.8 (or ≤ − 0.8), p-value ≤ 0.01.

play specific roles in clusters I and III, respectively. Cluster IV represents the group of

genes that function in the late stages of catkin fibers development. Additionally, we

performed network analysis between SM-related genes and TFs using data from a

previous study (Ye et al., 2014). As shown in Fig. 6b, many TF gene families were

involved in poplar catkin fibers development, including *bHLH*, *ERF*, *WRKY*, *MYB*

and *HD-ZIP*. *Cis*-acting elements were also predicted for genes related to SM in this

study. As shown in Fig. S2, light-responsive elements were the most important

environmental factors, while abscisic acid-, methyl jasmonate (MeJA)- and salicylic

acid- responsive elements were the most common motifs in plant hormone factors.

Overexpression of *PtoSUS2* and *PtoVIN3* increases trichome density

in wild-type *A. thaliana* and *tl1/2* mutant

The results for SUS and VIN indicate roles for SM in poplar catkin fibers

development; two key genes, *PtoSUS2* and *PtoVIN3*, were selected to demonstrate the

exact role. Thereinto, *PtoSUS2* had a stable expression level except in root, while

PtoVIN3 had higher expression levels in stem and root, and the two genes all played

roles during the early development of female and male floral bud (Fig. S3). Because

of the feasibility and long period involved in generating transgenic poplars and

investigating the phenotype of poplar catkins, the *A. thaliana* trichome system was

used to analyze gene function. We introduced *PtoSUS2* and *PtoVIN3* constructs driven

by the 35S promoter into wild-type *A. thaliana* and used T3-generation plants to

analyze the phenotype. Thirteen and twenty-two independent transgenic lines were

finally obtained for *PtoSUS2* and *PtoVIN3*, respectively. Three lines of each

transgenic type were extensively evaluated. In wild-type *A. thaliana*, trichomes are

Fig. 7 Leaf trichome morphology and number, rosette leaf number, fresh weight, and height in col and its transgenic lines. a-f Upper panel, trichome morphology under a stereomicroscope; scale bars = 1 mm. Lower panel, trichome morphology under a scanning electron microscope. **g-j** Phenotypic characteristics of wild-type and transgenic plants; bars represent ± SD.

widely distributed on the adaxial and abaxial surfaces of rosette leaves and cauline

leaves, as well as stems. Compared with the wild-type, the *PtoSUS2* and *PtoVIN3*

overexpression lines had higher leaf trichome densities on the adaxial surface of

rosette leaves and stems at 30 DAG, and trichome shape was unaffected (Fig. 7a-f;

Fig. S4). The number of sixth-rosette leaf trichome in the wild-type was 67 ± 8.80,

significantly fewer than in the transgenic plants, *Pro35S:PtoSUS2/col-2* (87 ± 6.89)

and *Pro35S:PtoVIN3/col-24* (94 ± 8.41) (Fig. 7g). However, the rosette leaf number,

fresh weight and height did not significantly differ between wild-type and

overexpression plants (Fig. 7h-j). Similar results were obtained in terms of root length

(10 DAG), rosette leaf shape (20 DAG) and whole plants (40 DAG) (Fig. S5).

Additionally, the trichome density of transgenic plants on the abaxial surface of

rosette leaves slightly increased, but the numbers of trichomes on the adaxial and

abaxial surfaces of cauline leaves had no significant difference, and these results were

not affected by incubation time (Fig. S6).

Overexpression of *PtoSUS2* and *PtoVIN3* also led to greater trichome density in

mutants *tl1* and *tl2*. Three transgenic lines of each transgenic type were selected for

detailed comparison. In *tl1*, eleven and thirteen independent transgenic plants were

obtained for *PtoSUS2* and *PtoVIN3*, respectively. Compared with the glabrous

phenotype (lacking trichomes) in *tl1*, overexpression of *PtoSUS2* or *PtoVIN3* resulted

in normal trichomes on the adaxial surface of rosette leaves; trichomes were observed

in the stems of some transgenic plants (Fig. 8a-c; Fig. S7). In *tl2*, twelve and eleven

independent transgenic plants were obtained for *PtoSUS2* and *PtoVIN3*, respectively.

In *tl2*, trichomes on the adaxial surface of rosette leaves were malformed; few

Fig. 8 Morphological characteristics of leaf trichome in *tl1/2* and their transgenic lines. Upper panel, trichome morphology under a stereomicroscope; scale bars = 1 mm. Lower panel, trichome morphology under a scanning electron microscope.

trichomes could be distinguished on the leaf edges. Overexpression of *PtoSUS2* or

PtoVIN3 led to greater malformed trichome density in rosette leaves, especially along

the leaf margins. However, the glabrous phenotype of stems was unchanged in *tl2*

transgenic plants (Fig. 8d-f; Fig. S8).

Changes in expression of trichome-related genes in *A. thaliana*

overexpressing *PtoSUS2* and *PtoVIN3*

RT-qPCR was performed to evaluate the expression levels of the exogenous

PtoSUS2 and *PtoVIN3* genes. The two exogenous genes were strongly expressed in

the transgenic lines but were undetectable in wild-type *A. thaliana* (Fig. 9).

Endogenous genes associated with trichome development were also influenced by

PtoSUS2 and *PtoVIN3*. Overexpression of *PtoSUS2* or *PtoVIN3* in the wild-type

increased the expression levels of *AtGL1*, *AtGL2*, *AtGL3*, *AtTTG1*, and *AtMYB23*. The

expression of *AtGL1* had a trend similar to the trends of exogenous *PtoSUS2* and

PtoVIN3; it played a major role in the trichome phenotype of transgenic lines. Similar

results were obtained in mutants *tl1* and *tl2*. The phenotype of transgenic lines was

closely related to gene expression. For individuals with the most pronounced

phenotype in *tl1*, the expression levels of *AtGL2*, *AtGL3*, and *AtMYB23* were

increased in *Pro35S:PtoSUS2/tl1-11* and *Pro35S:PtoVIN3/tl1-14*. In *tl2*, elevated

expression levels of *PtoSUS2* and *PtoVIN3* led to increased expression of *ATGL1*,

AtGL3, *AtMYB23*, and *AtTTG1*, especially the results in *Pro35S:PtoSUS2/tl2-22* and

Pro35S:PtoVIN3/tl2-29. In different transgenic lines, the expression patterns of *AtGL1*,

AtGL3, and *AtMYB23* showed trends similar to *PtoSUS2* and *PtoVIN3;* they were

significantly increased, compared with *tl2*.

Fig. 9 Gene expression patterns in col, tl1/2, and their transgenic lines, as determined RT-qPCR. Bars represent ± SD, ** significant difference at $P < 0.01$, and * significant difference at $P < 0.05$.

Discussion

The catkin fibers are widely presented in poplar, and are similar to *A. thaliana* trichomes and cotton fibers. Numerous studies have indicated that sucrose is important for cotton fiber development (described in detailed in a follow-up), but similar research has not been carried out on poplar catkin fibers. Thus, in this study, SM was thoroughly analyzed during poplar catkin fibers development. And as we performed, sucrose degradation was the primary metabolic factor in catkin fibers.

Invertase has a function during the early development of poplar catkin fibers

According to the pH and subcellular localization, INVs can be divided into CWIN, VIN and NIN, all of which have different functions in plants (Ruan, 2014). Although CWIN and VIN both belong to the acid INV, they can be divided into two subgroups based on evolutionary relationships (Fig. S1a). In cotton fibers, *CWIN* appears to have no regulatory effect in cotton fibers, based on its low and steady expression level (Ruan and Chourey, 1998; Wang et al., 2010). Similarly, low CWIN activity was observed in poplar catkin fibers (Fig. 3c). In contrast with CWIN, VIN has been proven to participate in both initiation and elongation process in cotton fibers. The hexose produced by *GhVIN1* can be sensed by hexokinase, and then influences cotton fiber initiation though MYB TFs and auxin signaling (Wang et al., 2014). Meanwhile, hexose production can promote fiber elongation by increasing osmotic potential (Wang et al., 2010). Two other members of *VIN* were shown to be associated with cotton fiber development (Taliercio et al., 2010). Among them, *GhVIN2*, probably an allele of *GhVIN1*, was expressed primarily in elongating fibers, and

decreased during fiber maturation. Compared with results in cotton fibers, VIN

activity had a similar pattern during poplar catkin fibers development, indicating that

it probably plays the similar role in poplar catkin fibers. Thereinto, *PtoVIN3* has a

closest relationship with *GhVIN1* and *GhVIN2*. Surprisingly, NIN appeared to have a

function during the early development of poplar catkin fibers, differing from cotton

fibers. In contrast with their low expression levels in cotton fiber (Wang et al., 2010),

the NIN high activity in poplar catkin fibers led us to infer that NIN function differs

between poplar and cotton.

376 Sucrose synthase has a close correlation with the elongation of poplar catkin

377 fibers

In this study, SUS showed greater activity than INV (Fig. 3e, f), indicating a

potential role in poplar catkin fibers development. In cotton, SUS has been proposed

to play direct role in fiber yield and quality (Ahmed et al., 2018). *GhSUS3* was firstly

found to have high expression during cotton fiber development, and was thus thought

to have a role in that process (Ruan et al., 1997). Subsequently, fiberless mutant and

SUS-suppression transgenic plant confirmed the function of *GhSUS3* (Ruan and

Chourey, 1998; Ruan et al., 2003). In the fiberless mutant and transgenic plant, the

activity of SUS had a close relationship with fiber quality. Furthermore, Elizabeth et

al. (Elizabeth et al., 2011) found four isoforms of SUS that are expressed during

cotton fiber development. Three of these four members were similar to *GhSUS3*, with

high expression in fibers at the extension stage (at 8 days post-anthesis, dpa). Another

isoform, *GhSUSC*, was highly expressed in fibers during SCW synthesis (at 20 dpa)

(Elizabeth et al., 2011). Different results were obtained in *Gossypium arboretum*

(Chen et al., 2012), with four homologous genes having high expression during fiber

development (at 15 dpa), similar to our results in poplar. Two homologous members,

PtoSUS1 and *PtoSUS2*, had high expression during catkin fibers development, while

the five members in other two clades had consistently low expression, similar to the

pattern in xylem (Zhang et al., 2011). High SUS activity was accompanied by a high

hexose to sucrose ratio, which was also observed in cotton fibers (Ruan et al., 1997).

High turgor pressure, caused by high hexose content, is required for rapid cell

extension (Ruan and Chourey, 1998). Therefore, SUS probably promotes catkin fibers

elongation by breaking down sucrose to increase turgor pressure. Although VIN

activity was also high during catkin fibers extension, fitting analysis suggested that

SUS plays the dominant role (Fig. 3).

The cellulose synthase gene family and other important genes play roles during

poplar catkin fibers development

The *CesA* gene family has been found to play roles during cotton fiber

development (Haigler et al., 2009; Kim et al., 2011; Kim et al., 2012). To elaborate,

the global genome analysis of the *CesA* gene family in cotton was performed (Li et al.,

2013). Members of the S1, S2 and S3 clades showed preferential expression during

SCW synthesis, while other members appeared to play a role in cotton fiber initiation

and extension. However, in poplar catkin fibers, members of S1-S3, which are

involved in SCW synthesis, all had low expression levels, although their levels

increased sharply during early development of catkin fibers except for *PtoCesA18*

(Fig. 5a-c). Members of P1-P3, which are involved in primary cell wall (PCW)

synthesis, played a more critical role in catkin fibers development as their high

expression. Thereinto, *PtoCesA17*, *PtoCesA2* and *PtoCesA15* are notable due to their

unusual expression pattern. Compared with cotton, less changes of *CesAs* expression

were observed in poplar catkin fibers, and we speculate that this difference underlies

the different qualities of catkin fibers in the two plants.

To further elucidate the role of sucrose in poplar catkin fibers, we investigated

TFs and other genes that may be involved in the development process (Fig. 5h, i,

Table S1). Similar expression patterns of *TF* gene family members were observed in

poplar catkin fiber and cotton fibers. *GhMYB109* (Suo et al., 2003), *GhDEL65*

(Shangguan et al., 2016) and *GhTTG1/3* (Humphries et al., 2005) are homologues of

GL1 (MYB), *GL3/EGL3* (bHLH) and *TTG1* (WD40), respectively, in cotton and are

thought to exert effects on fiber initiation. *Pto(GL1)-L1*, *Pto(GL3)-L1* and *2* and

Pto(TTG1)-L1, which are the representatives in poplar, all showed increased

expression at the initiation stage of catkin fibers development. *GhMYB25* (MYB) and

GhHox3 (HD-ZIP IV) were found to play regulatory roles in fiber extension

(Machado et al., 2009; Shan et al., 2014), however their homologues had delayed

expression in poplar catkin fibers, aside from *Pto(GhMYB25)-L1* (Fig. 5h). Notably,

MYB186, the homologue of *Pto(GhMYB25)-L4*, is vital in poplar leaf trichome

development (Plett et al., 2010; Patrick Bewg et al., 2022).

PtoSUS2* and *PtoVIN3* increased trichome density in *A. thaliana

In poplar, *PtoSUS2* is widely expressed in roots, vegetative buds, and floral

catkins; its homologue *PtrSUS2* is highly expressed in xylem tissues and moderately

expressed in leaves and roots (Zhang et al., 2011; An et al., 2014). In this study,

PtoSUS2 had higher expression than other members of the sucrose synthase gene

family during poplar catkin fibers development; it was significantly correlated with

SUS activity (Fig. 4, Table S4). Therefore, *PtoSUS2* may play a major role in poplar

catkin fibers development; this hypothesis was supported by the results of *PtoSUS2*

overexpression in *A. thaliana*. Overexpression wild-type plants had increased

trichome density on rosette leaves and stems, and transgenic plants of the glabrous

mutants produced more trichomes on rosette leaves (Fig. 7, Fig. 8). Another candidate

gene, *PtoVIN3*, was constitutively expressed in roots and leaves; it is also regulated by

abiotic and biotic stresses (Chen et al., 2015). In this study, overexpression of

PtoVIN3 increased trichome density on rosette leaves and stems, indicating that it

functions with *PtoSUS2* to control catkin fibers development in poplar. Notably, the

increased density of trichomes is not due to enhanced vegetative growth in *A. thaliana*

transgenic lines (Fig. 7 h-j, Fig. S5). Trichome development in *A. thaliana* is

controlled by the *AtGL1-AtGL3-AtTTG1* complex (Walker et al., 1999; Bernhardt et

al., 2005; Bloomer et al., 2012; Pesch et al., 2015); *AtGL2* acts downstream of this

complex to control normal trichome development (Rerie et al., 1994). *AtMYB23*

controls trichome branching and initiation at leaf edges; it is functionally redundant

with *AtGL1* (Kirik et al., 2005). In wild-type plants overexpressing *PtoSUS2* and

PtoVIN3, the expression levels of all above-mentioned genes were increased. This

demonstrates the effect of *PtoSUS2* and *PtoVIN3* on trichome development in *A.*

thaliana at the molecular level. In different transgenic lines, the expression patterns of

AtGL1 are closely like the expression of *PtoSUS2* or *PtoVIN3*. *GhSUS3*, which has a

close relationship with *PtoSUS2* in cotton, is implicated in cotton fiber cell initiation

and elongation (Ruan et al., 2003). Overexpression of *SUS* increases cotton fiber

length, while decreased expression of *GhSUS3* induces the fiberless mutation (Ruan

and Chourey, 1998; Ahmed et al., 2018). Additionally, high vacuolar invertase activity

is important for cotton fiber development; the homologue of *PtoVIN3*, *GhVIN1*, is a

key factor in that process (Wang et al., 2010). *GhVIN1* regulates fiber initiation from

the ovule epidermis, probably via sugar signaling with *cis*-acting elements of the

sugar-response element in *GhMYB25* and *GhMYB25like* (Wang et al., 2014), which is

similar to poplar catkin fibers development. The promoter regions of *Pto(GL1)-L1/2/3*,

the homologues of *AtGL1*, and *Pto(GhMYB25)-L1/2/3/4*, the homologues of

GhMYB25, have also been analyzed in poplar. SURE motifs (i.e., sugar-responsive

elements) (Grierson et al., 1994), were found in the promoter region of

Pto(GhMYB25)-L1/2/4 (Fig. S9). Additionally, the SP8B-like motif (TACT(X)TT),

which is similar to another sugar-responsive element (Ishiguro and Nakamura, 1992),

was found in the promoter regions of *Pto(GL1)-L2* and *Pto(GhMYB25)-L3/4*.

Therefore, we speculate that *PtoSUS2* and *PtoVIN3* probably regulate catkin fibers

development via sugar signaling. This result was confirmed by wild-type and *tl2*

transgenic plants because *AtGL1* showed similar trends to *PtoSUS2* and *PtoVIN3*.

In this study, SM was analyzed in detail in the context of poplar catkin fibers

development, and sucrose degradation was found to play the dominant role in the

Fig. 10 Potential regulatory model of poplar catkin fibers development

process. As shown in Fig. 10, sucrose is consumed rapidly as poplar catkin fibers

mature, especially at the extension stage. VIN and SUS, probably derived by *PtoVIN3*

and *PtoSUS2*, were thought to perform the function. Furthermore, the PCW

synthesis-related *PtoCesAs* may play a major role in cellulose synthesis. As an

important metabolic process in poplar catkin fibers, analyzing the regulatory

mechanism of SM is not only helpful for us to understand the regulatory mechanism

in woody plants, and for identifying key candidate genes with high breeding value in

P. tomentosa, but also provides a foundation for solving environmental problems

caused by poplar catkins though genetic engineering.

Materials and Methods

Plant materials and histological analysis

Live branches of adult female *P. tomentosa* were collected in February 2018, at

which time the flower buds were undergoing megasporogenesis and embryo sac

formation (An, 2010). The branches were placed in a greenhouse (25 ± 2°C, 70%

humidity) for hydroponic culture, and samples were collected as described previously

(Ye et al., 2014). Capsules were collected after the branches had been cultured for 0,

24, 34, 48, 58, 72, 96 and 120 h. Samples collected at these eight time points were

subjected to histological analysis; according to the analysis results, the samples from

the last five time points, which were related to catkin fibers development, were

subjected to RNA extraction as well as sugar content and enzyme activity analyses.

RNA extraction and histological analysis were also performed as described previously

(Ye et al., 2014).

The Columbia ecotype (Col) and two trichome mutants, *trichome less 1* (*tl1*,

SALK_133117C) and *trichome less 2* (*tl2*, SALK_039825C), were used in this study.

The seeds were sterilized in 5% sodium hypochlorite solution for 15 min, and then

placed in half-strength Murashige-Skoog (MS) medium at 4°C for 72 h. Subsequently,

the seeds were cultivated in an incubator at 22°C under LD (16 h light/8 h dark)

condition for 7 days, then transplanted into nutrient soil with added perlite.

Quantitation of sugar content and enzyme activities

Samples of 1 g (dry weight, DW) were added to 5 mL distilled water or 90%

alcohol, and the mixture was extracted through ultrasonication for 10 min.

Subsequently the mixture was centrifuged at 4°C and 12,000 rpm for 10 min. The

mixture was filtered through a 0.45-μm hydrophilic filter prior to analysis. The

concentrations of sucrose, fructose and glucose were analyzed by high-performance

liquid chromatography (HPLC) using an Agilent 1260 HPLC system (Agilent

Technologies, Santa Clara, CA, USA), and a carbohydrate column (4.6 × 150 mm, 5

μm, Agilent Technologies) was maintained at 35 °C. The sugar concentrations were

quantified according to a standard solution (Sigma, St. Louis, MO, USA), and defined

in terms of mg•g^{-1} DW.

The activities of SUS and SPS were analyzed according to the method of Yang et

al. (Yang et al., 2013) with some modifications. Sample of 5 g (fresh weight, FW) was

weighed and ground in liquid nitrogen for 5 min, and then extraction buffer (see

Methods S1) was added, followed by mixing and centrifuging at 20,000 rpm for 30

min. Subsequently, the supernatant (1 mL) was desalted using a Sephadex G25 PD-10

column with desalination buffer (see Methods S1). For SUS analysis, 1 mL of reaction

mixture I (see Methods S1) was incubated at 37℃ for 30 min. Subsequently, the

mixture was placing in boiling water for 5 min to stop the reaction. The optical

density (OD) of the mixture was measured at 540 nm using an ultraviolet

spectrophotometer (Cary 5000, Agilent). For SPS activity, 200 μL reaction mixture II

(see Methods S1) was incubated at 30℃ for 30 min. Subsequently, 200 μL NaOH (5

mol/L) was added, and the mixture was placing in boiling water for 10 min to stop the

reaction. After cooling, 3.5 mL of anthrone (0.15%, dissolved in 80% H_2SO_4) was

added, and the mixture was placed in a 40 ℃ water bath for 20 min. The OD of the

mixture was measured at 620 nm using the Cary 5000 instrument. The activities of

neutral/alkaline, vacuolar, and cell wall INVs (NIN, VIN and CWIN, respectively)

were analyzed according to previously described methods (Chen et al., 2015), and the

enzyme activities were recorded as $\mu mol \cdot g^{-1}$ FW$\cdot h^{-1}$.

Candidate gene identification and RT-qPCR

To determine the expression levels of key genes during the development of

catkin fibers, primers were designed to amplify fragments specific to each gene.

Genes from *A. thaliana* and cotton were used as queries for a Basic Local Alignment

Search Tool for Proteins (BLASTP) search against the protein sequences from the *P.*

tomentosa genomic data (An et al., 2022). The sequences of the primer pairs used for

each gene are listed in Table S2.

Total RNA was used for first-strand cDNA synthesis with the Reverse

Transcription System I (Promega, Madison, WI, USA), then diluted 1:50 in water. The

diluted cDNA was used as a template, and PCR amplification was performed with the

7500 Fast Real-Time PCR System (Applied Biosystems, Foster City, CA, USA) using

the following program: 94℃ for 5 min (one cycle); 94℃ for 30 s, 60℃ for 20 s, 72℃

for 30 s (35 cycles); 72℃ for 5 min. The *PtActin* gene was used as an internal

reference gene, as described previously (Chen et al., 2018), and the SYBR Premix Ex

Taq kit (Takara, Shiga, Japan) was used as the reaction solution. All analyses were

performed with three technical and three biological replicates.

Phylogenetic tree and co-expression network

To confirm the evolutionary relationships in SM related-gene families, all genes

were compared with their homologues from poplar*, A. thaliana* and cotton through

multiple sequence alignment (Table S3) using the ClustalW program within MEGA

7.0 software. Phylogenetic trees were draw with MEGA 7.0 using the

neighbor-joining method with complete deletion, and 1,000 replications were used for

bootstrap analysis. Gene co-expression network analysis was performed to identify

correlated genes using R software (v3.6.3), and the networks were visualized using

Cytoscape (v.3.7.1).

Transformation and gene expression analysis in *A. thaliana*

The open reading frame sequences of *PtoSUS2* and *PtoVIN3* were amplified and

cloned into the binary vector pCAMBIA 1301 to generate overexpression vectors.

Transgenic lines were obtained by the *Aarobacterium tumefaciens*-mediated method

established by (Zhang et al., 2006). Transgenic plants were selected on half-strength

MS medium containing 25 mg/L hygromycin. Gene expression analysis was

performed as above, except that relative expression levels were calculated using the

$2^{-\Delta\Delta Ct}$ method. The sequences of the primer pairs used for trichome-related genes in *A.*

thaliana and exogenous genes are listed in Table S2.

Morphological analysis and scanning electron microscopy (SEM)

The sixth rosette leaves, stems (morphological bottom), and bottom-most cauline

leaves were used for trichome comparison with a stereomicroscope (Stemi 305,

ZEISS). The number of trichomes in rosette leaves was determined in plants that had

been cultured for 30 days after germination (DAG). Five biological replicates were

performed. For SEM, the samples were cut into 1×1-cm pieces, and dehydrated in a

vacuum freeze dryer (FreeZone 4.5L, Labconco) for 48 h. Images were obtained using

a scanning electron microscope (S-3400N, Hitachi, Japan).

Statistical Analysis

578 Sugar content, enzyme activity and gene expression data were analyzed through

579 one-way analysis of variance (ANOVA) using SPSS software (version 22.0 for

580 Windows; SPSS Inc., Chicago, IL, USA). The correlations between genes expression

581 patterns in SM and sugar contents was analyzed by Pearson correlation using the same

582 software. Two-tail test was used to test for significance level as $P < 0.05$ and $P < 0.01$.

Accession numbers

584 The gene sequences can be found under the accession numbers: *PtoSUS2*

585 (ON534005), *PtoVIN3* (ON534006).

Acknowledgements

This work was funded by the National Natural Science Foundation of China

(31800555 and 31570661) and the Fundamental Research Funds for the Central

Universities (BLX201701). We gratefully acknowledge the families who participated

by giving research assistance. We also thank the families who participated by giving

research assistance.

Authors contributions

Z.C. and X.Y. designed the work; X.Y., T.Z. and N.Y. performed experiments and

analyzed data; Co-expression network analysis was carried out by P.R.; X.Y. and T. Z.

drafted the manuscript; and Z.C., G.L., L. J. and X.A. revised the manuscript. X. Y.

and T. Z. contributed equally to this work.

Supporting information

Fig. S1 Phylogenetic tree of the SM-related gene families.

Fig. S2 Analysis of cis-acting elements for SM-related genes in *P. tomentosa*.

Fig. S3 Tissue-specific expression analysis of *PtoSUS2* and *PtoVIN3*.

Fig. S4 Morphological characteristics of trichome on rosette leaves and stems in

***A. thaliana* wild-type and transgenic lines.**

Fig. S5 Characteristics of root, rosette leaves, and whole plant in *A. thaliana*

wild-type and transgenic lines.

Fig. S6 Changed in trichome density on leaves during leaf development in *A.*

***thaliana* wild-type and transgenic lines.**

Fig. S7 Morphological characteristics of trichome on leaves and stems in

transgenic lines of *tl1* and *tl1*.

Fig. S8 Morphological characteristics of trichome on leaves and stems in

transgenic lines of *tl2* and *tl2*.

Fig. S9 Analysis of sugar signal-related cis-acting elements for *Pto(GL1)-L1/2/3*

and *Pto(GhMYB25)-L1/2/3/4*.

Table S1 Comparison of potential TFs and other genes in poplar and cotton

Table S2 Primers used for RT-qPCR

Table S3 Accession numbers of genes used for phylogenetic tree construction

Table S4 Correlations between the expression patterns of genes involved in SM

and sugar contents in poplar catkin fibers development

Methods S1: reagent formulation for enzyme activities quantitation

Parsed Citations

Ahmed M, Shahid AA, Akhtar S, Latif A, ud Din S, Fanglu M, Rao AQ, Sarwar MB, Husnain T, Xuede W (2018) Sucrose synthase genes: a way forward for cotton fiber improvement. Biologia 73: 703-713
Google Scholar: Author Only Title Only Author and Title

An X, Chen Z, Wang J, Ye M, Ji L (2014) Identification and characterization of the Populus sucrose synthase gene family. Gene Amsterdam
Google Scholar: Author Only Title Only Author and Title

An X, Gao K, Chen Z, Li J, Yang X, Yang X, Zhou J, Guo T, Zhao T, Huang S, Miao D, Ullah Khan W, Rao P, Ye M, Lei B, Liao W, Wang J, Ji L, Li Y, Guo B, Siddig Mustafa N, Li S, Yun Q, Keller SR, Mao JF, Zhang RG, Strauss SH (2022) High quality haplotype-resolved genome assemblies of Populus tomentosa Carr., a stabilized interspecific hybrid species widespread in Asia. Mol Ecol Resour 22: 786-802
Google Scholar: Author Only Title Only Author and Title

An XW, Dongmei; Wang, Zeliang; Wang, Jingcheng; Cao, Guanlin; Bo, Wenhao; Zhang, Zhiyi (2010) Expression profile of PtLFY in floral bud development associated with floral bud morphological differentiation in Populus tomentosa. Sci Silvae Sinicae 46: 32-38
Google Scholar: Author Only Title Only Author and Title

Bernhardt C, Zhao M, Gonzalez A, Lloyd A, Schiefelbein J (2005) The bHLH genes GL3 and EGL3 participate in an intercellular regulatory circuit that controls cell patterning in the Arabidopsis root epidermis. Development 132: 291-298
Google Scholar: Author Only Title Only Author and Title

Bloomer R, Juenger T, Symonds V (2012) Natural variation in GL1 and its effects on trichome density in Arabidopsis thaliana. Molecular Ecology 21: 3501-3515
Google Scholar: Author Only Title Only Author and Title

Bui APN, Ong B-N, Huu T-AN, Tran H-D (2019) Similarities between Initiation Mechanism between Cotton Fiber and Arabidopsis Trichome: Prospects in Improving Cotton Fiber Yield. annual research & review in biology: 1-9
Google Scholar: Author Only Title Only Author and Title

Chen A, He S, Li F, Li Z, Ding M, Liu Q, Rong J (2012) Analyses of the sucrose synthase gene family in cotton: structure, phylogeny and expression patterns. BMC plant biology 12: 85
Google Scholar: Author Only Title Only Author and Title

Chen HL, Cluver r (2010) Assessment of Poplar Seed Hair Fibers as a Potential Bulk Textile Thermal Insulation Material. Clothing & Textiles Research Journal 28: p.255-262
Google Scholar: Author Only Title Only Author and Title

Chen Z, Gao K, Su X, Rao P, An X (2015) Genome-Wide Identification of the Invertase Gene Family in Populus. plos one 10
Google Scholar: Author Only Title Only Author and Title

Chen Z, Rao P, Yang X, Su X, Zhao T, Gao K, Yang X, An X (2018) A Global View of Transcriptome Dynamics During Male Floral Bud Development in Populus tomentosa. Scientific Reports 8: 722
Google Scholar: Author Only Title Only Author and Title

Elizabeth B, Michel VT, G WR, Danny L, M CP, Steven E, Yong-Ling R, Tony A, T FR (2011) A novel isoform of sucrose synthase is targeted to the cell wall during secondary cell wall synthesis in cotton fiber. Plant Physiology 157: 40-54
Google Scholar: Author Only Title Only Author and Title

Fujii S, Hayashi T, Mizuno K (2010) Sucrose synthase is an integral component of the cellulose synthesis machinery. Plant and cell physiology 51: 294-301
Google Scholar: Author Only Title Only Author and Title

Grebe M (2012) The patterning of epidermal hairs in Arabidopsis : updated. current opinion in plant biology 15: 31-37
Google Scholar: Author Only Title Only Author and Title

Grierson CS, Du J, Torreszabala MD, Beggs K, Holdsworth M, Bevan M (1994) Separate cis sequences and trans factors interact to direct metabolic and developmental regulation of a potato tuber storage protein gene.

Haigler CH, Singh B, Wang G, Zhang D (2009) Genomics of cotton fiber secondary wall deposition and cellulose biogenesis. In AH Paterson, ed, Genetics and genomics of cotton. Springer, New York, NY, pp 385-417
Google Scholar: Author Only Title Only Author and Title

Haigler CH, Singh B, Zhang D, Hwang S, Wu C, Cai WX, Hozain M, Kang W, Kiedaisch B, Strauss RE (2007) Transgenic cotton over-producing spinach sucrose phosphate synthase showed enhanced leaf sucrose synthesis and improved fiber quality under controlled environmental conditions. Plant Molecular Biology 63: 815-832
Google Scholar: Author Only Title Only Author and Title

Humphries JA, Walker AR, Timmis JN, Orford SJ (2005) Two WD-repeat genes from cotton are functional homologues of the Arabidopsis thalianaTRANSPARENT TESTA GLABRA1 (TTG1) gene. Plant molecular biology 57: 67-81
Google Scholar: Author Only Title Only Author and Title

Ishiguro S, Nakamura K (1992) The nuclear factor SP8BF binds to the 5'-upstream regions of three different genes coding for major proteins of sweet potato tuberous roots. Plant Molecular Biology 18: 97-108
Google Scholar: Author Only Title Only Author and Title

Kim H, Triplett BA (2007) Cellulose synthase catalytic subunit (Cesa) genes associated with primary or secondary wall biosynthesis in developing cotton fibers (Gossypium hirsutum). In World Cotton Research Conference-4. ICAC, Lubbock, Texas, USA
Google Scholar: Author Only Title Only Author and Title

Kim HJ, Murai N, Fang DD, Triplett BA (2011) Functional analysis of Gossypium hirsutum cellulose synthase catalytic subunit 4 promoter in transgenic Arabidopsis and cotton tissues. Plant science 180: 323-332
Google Scholar: Author Only Title Only Author and Title

Kim HJ, Triplett BA, Zhang H-B, Lee M-K, Hinchliffe DJ, Li P, Fang DD (2012) Cloning and characterization of homeologous cellulose synthase catalytic subunit 2 genes from allotetraploid cotton (Gossypium hirsutum L.). Gene 494: 181-189
Google Scholar: Author Only Title Only Author and Title

Kirik V, Lee MM, Wester K, Herrmann U, Zheng Z, Oppenheimer D, Schiefelbein J, Hulskamp M (2005) Functional diversification of MYB23 and GL1 genes in trichome morphogenesis and initiation. Development 132: 1477-1485
Google Scholar: Author Only Title Only Author and Title

Klocko AL, Brunner AM, Huang J, Meilan R, Lu H, Ma C, Morel A, Zhao D, Ault K, Dow M, Howe G, Shevchenko O, Strauss SH (2016) Containment of transgenic trees by suppression of LEAFY. Nat Biotechnol 34: 918-922
Google Scholar: Author Only Title Only Author and Title

Li A, Xia T, Xu W, Chen T, Li X, Fan J, Wang R, Feng S, Wang Y, Wang B (2013) An integrative analysis of four CESA isoforms specific for fiber cellulose production between Gossypium hirsutum and Gossypium barbadense. Planta 237: 1585-1597
Google Scholar: Author Only Title Only Author and Title

Likon M, Remškar M, Ducman V, Švegl F (2013) Populus seed fibers as a natural source for production of oil super absorbents. Journal of Environmental Management 114: 158-167
Google Scholar: Author Only Title Only Author and Title

Lu H, Klocko AL, Brunner AM, Ma C, Magnuson AC, Howe GT, An X, Strauss SH (2019) RNA interference suppression of AGAMOUS and SEEDSTICK alters floral organ identity and impairs floral organ determinacy, ovule differentiation, and seed-hair development in Populus. New Phytol 222: 923-937
Google Scholar: Author Only Title Only Author and Title

Machado A, Wu Y, Yang Y, Llewellyn DJ, Dennis ES (2009) The MYB transcription factor GhMYB25 regulates early fibre and trichome development. The Plant Journal 59: 52-62
Google Scholar: Author Only Title Only Author and Title

Martin LK, Haigler CH (2004) Cool temperature hinders flux from glucose to sucrose during cellulose synthesis in secondary wall stage cotton fibers. Cellulose 11: 339-349
Google Scholar: Author Only Title Only Author and Title

Müller NA, Kersten B, Leite Montalvão AP, Mähler N, Bernhardsson C, Bräutigam K, Carracedo Lorenzo Z, Hoenicka H, Kumar V, Mader M, Pakull B, Robinson KM, Sabatti M, Vettori C, Ingvarsson PK, Cronk Q, Street NR, Fladung M (2020) A single gene underlies the dynamic evolution of poplar sex determination. Nat Plants 6: 630-637
Google Scholar: Author Only Title Only Author and Title

Naoumkina M (2018) Advances in Understanding of Cotton Fiber Cell Differentiation and Elongation. In DD Fang, ed, Cotton Fiber: Physics, Chemistry and Biology. Springer International Publishing, Cham, pp 179-191
Google Scholar: Author Only Title Only Author and Title

Park JY, Canam T, Kang KY, Ellis DD, Mansfield SD (2008) Over-expression of an arabidopsis family A sucrose phosphate synthase (SPS) gene alters plant growth and fibre development. Transgenic Research 17: 181-192
Google Scholar: Author Only Title Only Author and Title

Patrick Bewg W, Harding SA, Engle NL, Vaidya BN, Zhou R, Reeves J, Horn TW, Joshee N, Jenkins JW, Shu S, Barry KW, Yoshinaga Y, Grimwood J, Schmitz RJ, Schmutz J, Tschaplinski TJ, Tsai CJ (2022) Multiplex knockout of trichome-regulating MYB duplicates in hybrid poplar using a single gRNA. Plant Physiol
Google Scholar: Author Only Title Only Author and Title

Pattanaik S, Patra B, Singh SK, Yuan L (2014) An overview of the gene regulatory network controlling trichome development in the model plant, Arabidopsis. frontiers in plant science 5: 259-259

Google Scholar: Author Only Title Only Author and Title

Pesch M, Schultheiß I, Klopffleisch K, Uhrig JF, Koegl M, Clemen CS, Simon R, Weidtkamp-Peters S, Hülskamp M (2015) TRANSPARENT TESTA GLABRA1 and GLABRA1 Compete for Binding to GLABRA3 in Arabidopsis. Plant Physiol 168: 584-597
  Google Scholar: Author Only Title Only Author and Title

Plett JM, Wilkins O, Campbell MM, Ralph SG, Regan S (2010) Endogenous overexpression of Populus MYB186 increases trichome density, improves insect pest resistance, and impacts plant growth. Plant J 64: 419-432
  Google Scholar: Author Only Title Only Author and Title

Rerie WG, Feldmann KA, Marks MD (1994) The GLABRA2 gene encodes a homeo domain protein required for normal trichome development in Arabidopsis. Genes & development 8: 1388-1399
  Google Scholar: Author Only Title Only Author and Title

Ruan Y-L (2014) Sucrose metabolism: gateway to diverse carbon use and sugar signaling. Annual review of plant biology 65: 33-67
  Google Scholar: Author Only Title Only Author and Title

Ruan Y-L, Chourey PS (1998) A fiberless seed mutation in cotton is associated with lack of fiber cell initiation in ovule epidermis and alterations in sucrose synthase expression and carbon partitioning in developing seeds. Plant physiology 118: 399-406
  Google Scholar: Author Only Title Only Author and Title

Ruan Y-L, Chourey PS, Delmer DP, Perez-Grau L (1997) The differential expression of sucrose synthase in relation to diverse patterns of carbon partitioning in developing cotton seed. Plant Physiology 115: 375-385
  Google Scholar: Author Only Title Only Author and Title

Ruan Y-L, Llewellyn DJ, Furbank RT (2003) Suppression of sucrose synthase gene expression represses cotton fiber cell initiation, elongation, and seed development. The Plant Cell 15: 952-964
  Google Scholar: Author Only Title Only Author and Title

Sannigrahi P, Ragauskas AJ, Tuskan GA (2010) Poplar as a feedstock for biofuels: A review of compositional characteristics. biofuels bioproducts and biorefining 4: 209-226
  Google Scholar: Author Only Title Only Author and Title

Shan C-M, Shangguan X-X, Zhao B, Zhang X-F, Chao L-m, Yang C-Q, Wang L-J, Zhu H-Y, Zeng Y-D, Guo W-Z (2014) Control of cotton fibre elongation by a homeodomain transcription factor GhHOX3. Nature communications 5: 5519
  Google Scholar: Author Only Title Only Author and Title

Shangguan XX, Yang CQ, Zhang XF, Wang LJ (2016) Functional characterization of a basic Helix-Loop-Helix (bHLH) transcription factor GhDEL65 from cotton (Gossypium hirsutum). Physiologia plantarum 158: 200-212
  Google Scholar: Author Only Title Only Author and Title

Suo J, Liang X, Pu L, Zhang Y, Xue Y (2003) Identification of GhMYB109 encoding a R2R3 MYB transcription factor that expressed specifically in fiber initials and elongating fibers of cotton (Gossypium hirsutum L.). Biochimica et Biophysica Acta (BBA)-Gene Structure and Expression 1630: 25-34
  Google Scholar: Author Only Title Only Author and Title

Taliercio E, Scheffler J, Scheffler B (2010) Characterization of two cotton (Gossypium hirsutum L) invertase genes. Molecular biology reports 37: 3915-3920
  Google Scholar: Author Only Title Only Author and Title

Walker AR, Davison PA, Bolognesi-Winfield AC, James CM, Srinivasan N, Blundell TL, Esch JJ, Marks MD, Gray JC (1999) The TRANSPARENT TESTA GLABRA1 locus, which regulates trichome differentiation and anthocyanin biosynthesis in Arabidopsis, encodes a WD40 repeat protein. The Plant Cell 11: 1337-1349
  Google Scholar: Author Only Title Only Author and Title

Wan X, Gu G, Lei M, Zeng W (2020) Bioaccessibility of metals/metalloids in willow catkins collected in urban parks of Beijing and their health risks to human beings. science of the total environment 717
  Google Scholar: Author Only Title Only Author and Title

Wang L, Cook A, Patrick JW, Chen XY, Ruan YL (2014) Silencing the vacuolar invertase gene GhVIN 1 blocks cotton fiber initiation from the ovule epidermis, probably by suppressing a cohort of regulatory genes via sugar signaling. The Plant Journal 78: 686-696
  Google Scholar: Author Only Title Only Author and Title

Wang L, Li X-R, Lian H, Ni D-A, He Y-k, Chen X-Y, Ruan Y-L (2010) Evidence that high activity of vacuolar invertase is required for cotton fiber and Arabidopsis root elongation through osmotic dependent and independent pathways, respectively. Plant Physiology 154: 744-756
  Google Scholar: Author Only Title Only Author and Title

Wu H, Tian Y, Wan Q, Fang L, Guan X, Chen J, Hu Y, Ye W, Zhang H, Guo W (2018) Genetics and evolution of MIXTA genes regulating cotton lint fiber development. new phytologist 217: 883-895

Google Scholar: Author Only Title Only Author and Title

Xue L, Wu H, Chen Y, Li X, Hou J, Lu J, Wei S, Dai X, Olson MS, Liu J, Wang M, Charlesworth D, Yin T (2020) Evidences for a role of two Y-specific genes in sex determination in Populus deltoides. Nat Commun 11: 5893
  Google Scholar: Author Only Title Only Author and Title

Yang W, Wang D, Li Y, Zhang Z, Tong S, Li M, Zhang X, Zhang L, Ren L, Ma X, Zhou R, Sanderson BJ, Keefover-Ring K, Yin T, Smart LB, Liu J, DiFazio SP, Olson M, Ma T (2021) A General Model to Explain Repeated Turnovers of Sex Determination in the Salicaceae. Mol Biol Evol 38: 968-980
  Google Scholar: Author Only Title Only Author and Title

Yang Z, Wang T, Wang H, Huang X, Qin Y, Hu G (2013) Patterns of enzyme activities and gene expressions in sucrose metabolism in relation to sugar accumulation and composition in the aril of Litchi chinensis Sonn. Journal of Plant Physiology 170: 731-740
  Google Scholar: Author Only Title Only Author and Title

Ye M, Zhong C, Su X, Ji L, Jia W, Liao... W (2014) Study of seed hair growth in Populus tomentosa, an important character of female floral bud development. BMC Genomics 15
  Google Scholar: Author Only Title Only Author and Title

Zhang D, Xu B, Yang X, Zhang Z, Li B (2011) The sucrose synthase gene family in Populus: structure, expression, and evolution. Tree genetics & genomes 7: 443-456
  Google Scholar: Author Only Title Only Author and Title

Zhang F, Zuo K, Zhang J, Liu X, Zhang L, Sun X, Tang K (2010) An L1 box binding protein, GbML1, interacts with GbMYB25 to control cotton fibre development. Journal of experimental botany 61: 3599-3613
  Google Scholar: Author Only Title Only Author and Title

Zhang X, Henriques R, Lin SS, Niu QW, Chua NH (2006) Agrobacterium-mediated transformation of Arabidopsis thaliana using the floral dip method. Nature Protocols 1: 641
  Google Scholar: Author Only Title Only Author and Title

Zhang X, Li Z, Yu Y, Wang H (2018) Characterizations of poplar catkin fibers and their potential for enzymatic hydrolysis. JOURNAL OF WOOD SCIENCE 64: 458-462
  Google Scholar: Author Only Title Only Author and Title