Genetic architectures of postmating isolation and morphology of two highly

diverged rockfishes (genus *Sebastes*)

Nozomu Muto[1], Takuma Kawasaki[2], Ryo Kakioka[3], Atsushi J. Nagano[4,5], Yuta Shimizu[1],

Shu Inose[1], Yohei Shimizu[2], Hiroshi Takahashi[6]

[1]School of Biological Sciences, Tokai University, Hokkaido, Japan

[2]Mariculture Fisheries Research Institute, Fisheries Research Department, Hokkaido

Research Organization, Hokkaido, Japan

[3]Tropical Biosphere Research Center, University of the Ryukyus, Okinawa, Japan

[4]Faculty of Agriculture, Ryukoku University, Shiga, Japan

[5]Institute for Advanced Biosciences, Keio University, Yamagata, Japan

[6]Department of Applied Aquabiology, National Fisheries University, Yamaguchi, Japan

Correspondence

Nozomu Muto, School of Biological Sciences, Tokai University, 5-1-1-1 Minamisawa,

Minami-ku, Sapporo, Hokkaido 005-8601, Japan.

Email: nzmutou@gmail.com

Funding information

JSPS KAKENHI (Grant No. 19K16217); The Yanmar Environmental Sustainability

Support Association (KI019078); The Hokusui Society Foundation.

Abstract

Speciation is a continuous process, and postmating isolation is thought to be an

important driver of the late stages of speciation. However, relatively little is empirically

known about the process compared to other isolating mechanisms that drive the early

stages of speciation, especially in non-model organisms. In this study, we

characterized the genetic architecture of postmating isolation between two rockfishes,

Sebastes schlegelii and *S. trivittatus*, whose reproductive isolation is complete. We

examined transmission ratio distortion (TRD) patterns of genetic markers in two

reciprocal backcross populations. Markers showing either of the two types of TRD was

widespread across the genome, with some of the distorted markers forming extensive

clusters around the recombination coldspots. These suggest that the postmating

isolation effectively prevents gene flow across the genome and the recombination

landscape contributes to the genetic architecture. Comparisons between two backcross

families and two developmental stages showed little similarity in the distorted markers,

suggesting asymmetry and stage-specificity of the isolation. This may be due to hybrid

incompatibility involving maternal factors or extrinsic selection. The lack of sex-ratio

distortion in the mapping families suggested that Haldane's rule in terms of hybrid

inviability does not hold. Additionally, QTL mapping detected significant QTLs for sex

and the morphological traits relevant to divergence and convergence of rockfishes,

including the body colour patterns and the overall intensities of pigmentations. These

findings, coupled with the first linkage map for rockfishes presented herein, constitute

an essential step towards a comprehensive understanding of speciation and

morphological diversification of rockfishes.

KEYWORDS

speciation, postzygotic isolation, hybrid inviability, extrinsic selection, recombination,

convergence, body colouration, sex, linkage map, QTL

1. INTRODUCTION

Speciation is a continuous process whereby gene flow between diverging species

diminishes with reproductive isolation. Existing species pairs can be ordered in a

continuous sequence—a speciation continuum—according to the degree of divergence

(Coyne & Orr, 1989a). Studying the species pairs at different points of the continuum

enables the reconstruction of the order and the rate of evolution of different forms of

reproductive barriers in a given system (Coyne & Orr, 1989a). A widely accepted view

is that premating isolation due to local adaptation or sexual isolation tends to evolve

early in speciation, whereas postmating isolation evolves later, thereby completing

speciation or maintaining separate species (Coyne & Orr, 2004; Schemske, 2010).

However, the latter process is relatively unclear, reflecting lesser attention being paid to

highly diverged species pairs whose reproductive isolation is strong or complete

(Matute & Cooper, 2021).

Postmating isolation may result from reduced fertilization success of heterospecific

gametes (gametic isolation), intrinsic hybrid sterility and inviability (intrinsic postzygotic

isolation), and environmentally-dependent hybrid sterility and inviability (extrinsic

postzygotic isolation) (Coyne & Orr, 2004). Among them, intrinsic postzygotic isolation

is the most intensively studied in terms of both its genetic model and contribution to

speciation; little is known about the other two mechanisms (Rometsch et al., 2020).

Intrinsic postzygotic isolation is caused by negative epistatic interactions involving two

or more genes in hybrids, i.e., Dobzhansky-Muller Incompatibility (DMI) (Maheshwari &

Barbash, 2011). Previous empirical studies have shown that the strength of intrinsic

postzygotic isolation, as well as the number of loci involved in DMI, often evolve

exponentially with divergence ('snowball effect') (Matute et al., 2010; Stelkens et al.,

2010), consistent with theoretical predictions (Orr & Turelli, 2001). Similarly, intrinsic

postzygotic isolation characterises the isolation of most 'good species' examined

(Coyne & Orr, 2004). However, it is still largely unknown what factors account for the

variations in tempo and mode of evolution of intrinsic postzygotic isolation across

different systems and how its genetic architecture relates to its efficacy (Coughlan &

Matute, 2020). Therefore, understanding how postmating isolation contributes to the

later stages of speciation requires more studies on various groups of organisms with

diverse life histories and degrees of divergence, especially the understudied non-model

organisms, and on gametic and extrinsic postzygotic isolation.

Postmating isolation between a pair of species can be detected by analysing the

deviation of marker genotype frequencies from the Mendelian expectations

(transmission ratio distortion, TRD) in hybrid mapping populations, such as backcross

and F_2 generations (Fishman & McIntosh, 2019). Gametic isolation and hybrid

inviability between parental species may cause variations in the fertilization success of

gametes and mortality of progeny in hybrid crosses, resulting in TRD at genetic

markers linked to the underlying loci of these barriers. Importantly, hybrid inviability due

to DMI is largely masked in F_1, but manifests in later generation hybrids (Orr, 1995; Orr

& Turelli, 2001). Extrinsic hybrid inviability can also be inferred from the patterns of

TRD relative to the rearing conditions. Moreover, mapping the distorted markers on a

reference genome or a linkage map can reveal the genetic architecture of the isolating

barriers (e.g., Brennan et al., 2014). Using TRD as a proxy of postmating isolation is

particularly useful when direct fitness measurement is difficult. For example, gametic

isolation and prenatal hybrid inviability in internally fertilising animals could be inferred

from TRD in the progeny immediately after parturition, which is otherwise, highly

challenging.

Speciation and associated phenotypic diversification is sometimes accompanied

by recurrent evolution of convergent traits (Schluter, 2000). While convergent traits

between distant lineages are often interpreted as independent adaptations to similar

ecological challenges, whether and to what extent they share similar genetic

mechanisms varies across systems (Rosenblum et al., 2014). The similarity in genetic

mechanisms of convergence between distant lineages ranges from independent

fixations or introgression of identical mutations to shared molecular pathways with

different nucleotides and genes (Chen et al., 1997; Evans et al., 2012; Stern, 2013).

The first critical step toward understanding the convergence at the genetic level is to

characterize the genetic architecture of the convergent trait in each lineage.

Rockfishes of the genus *Sebastes* are an ideal system to study speciation and the

associated phenotypic diversification in marine environments. *Sebastes* is a genus

comprising more than 100 species of viviparous marine fishes, most of which diverged

in the last 8 million years (Hyde & Vetter, 2007; Kolora et al., 2021). These species

have colonised diverse habitats from the near-shore rocky reefs of less than 1 m depth

to the deep-sea continental slopes of up to 1,500 m depth (Love et al., 2002; Nakabo &

Kai, 2013). Numerous incipient species pairs hybridising in nature are scattered across

the phylogeny of the genus, enabling multiple independent studies on the earliest stage

of speciation (Buonaccorsi et al., 2011; Muto et al., 2013; Saha et al., 2017; Schwenke

et al., 2018). Convergence in some morphological traits among independent lineages

has also been suggested either verbally or through phylogenetic comparative methods

(Ingram & Kai, 2014; Kai et al., 2003), providing opportunities to study how those traits

evolved along with the radiation in diverse environments. Furthermore, artificial crosses

can produce fertile F_1 hybrids even between species pairs with no record of natural

hybrids (T. Kawasaki, unpublished data). This feature enables genetic dissections of

postmating isolation and quantitative trait loci (QTL) analysis of adaptive traits using

later generation hybrids.

Nevertheless, little is known about the speciation and morphological diversification

of rockfishes. Incipient species often occupy habitats of different depths and have

distinct body colourations, implying the role of depth segregation or colour-related

assortative mating in the early stages of speciation (Ingram, 2011; Longo et al., 2022;

Muto et al., 2019; Shum et al., 2014; Venerus et al., 2013). Genes presumably related

to these isolating barriers have also been identified (Behrens et al., 2021; Heras &

Aguilar, 2019). However, the influence of other isolating barriers has not been tested,

and studies on isolating barriers between highly diverged species with strong

reproductive isolation is lacking. Consequently, how reproductive barriers develop

towards the completion of speciation in *Sebastes* remains elusive. As for morphological

convergence, the genetic bases of the convergent traits or any other morphological

traits have not been examined.

In this study, we characterise genetic architectures of postmating isolation and the

morphological difference between the two rockfishes *S. trivittatus* and *S. schlegelii*.

They are estimated to have diverged 6–7 million years ago (mya) (Hyde & Vetter, 2007;

Kolora et al., 2021), with no documented record of natural hybridisation (Figure S1).

They have partially overlapping geographical distributions in the western Pacific: *S.*

trivittatus is distributed in higher latitudes, from the coasts of Sakhalin Island and the

southern Kamchatka Peninsula to the Yellow Sea; whereas *S. schlegelii* is found from

the coasts of Sakhalin and the southern Kuril Islands to the Bohai Sea (Nakabo & Kai,

2013; Parin et al., 2014). Their morphological difference involves several traits

important in the contexts of speciation and convergence of rockfishes. Using two

reciprocal backcross families (single F_1 male × females of *S. trivittatus* and *S.*

schlegelii), we first constructed a hybrid linkage map, which is the first linkage map for

rockfishes. We then explored the TRD of genetic markers as an approximation of

genetic architectures of postmating isolation. Comparisons of the patterns of TRD

between two developmental stages and between the families revealed stage-specificity

and asymmetry of the isolation. We also performed a QTL analysis for the

morphological traits and sex. This study signifies the first critical step toward a better

understanding of speciation and morphological diversification of *Sebastes*, which in

turn will provide a valuable contrast to those in well-studied model organisms.

2 MATERIALS AND METHODS

2.1 Experimental crosses and sampling

Rockfishes are viviparous and have prolonged storage periods of sperm and zygotes in

a female ovary (each from one to several months) (Love et al., 2002). A female

releases hundreds of thousands to millions of hatched larvae per spawning. Nutrients

are supplied to the progeny both maternally (matrotrophy) and via yolk (Macfarlane &

Bowers, 1995; Tengfei et al., 2021). We artificially inseminated a wild female *S.*

trivittatus in November 2012 using sperm obtained from a wild male *S. schlegelii*, both

collected off Hokkaido, Japan, according to Kawasaki et al. (2017). Briefly, a sperm

suspension was prepared by mincing testis and sperm reservoir in urine obtained from

the male fish euthanized with tricaine methanesulfonate (MS222), then deposited into

the external genital pore of the female fish. Parturition of the F_1 yolk-sac larvae

occurred ca. seven months from the insemination. Subsequently, a single male of the

F_1 progeny was crossed with a wild female *S. trivittatus* (different individual from the

founder female) and a wild female *S. schlegelii* in December 2015 according to

Kawasaki et al. (2017). The parturitions occurred ca. six months later. The resulting

backcross families are hereafter referred to as BC_Str and BC_Ssc, respectively.

Rearing conditions for the fishes and a schematic diagram of the experimental design

are provided in Appendix S1 and Figure S2. We sampled approximately 100 progeny of

each family at each of the two stages: yolk-sac larvae (0 d after parturition) for

genotyping and immature subadult (ca. 270 d: see Figure S2) for both phenotyping and

genotyping. All the females used for the crosses had been separated from males at

least for one year to prevent unintended mating. We could not use F_1 females for the

cross because female rockfishes naturally require a few years more than males for

maturation (Love et al., 2002).

2.2 Phenotyping

Independent lineages of *Sebastes* sometimes have strikingly similar colour patterns,

including the presence of vertical bands on the dorsum, a horizontal stripe along the

lateral line, and a horizontal stripe along the dorsal-fin base, but not necessarily similar

colours. Besides, both colour and colour patterns are often the most important trait to

distinguish sister species of the genus (Ingram, 2011; Longo et al., 2022; Muto et al.,

2019; Shum et al., 2014; Venerus et al., 2013). Convergence in meristic and

morphometric traits involves gill-raker number and length, eye size, body depth and

width, pectoral-fin length and width, upper- and lower-jaw lengths, and head spination

(Ingram & Kai, 2014; Kai et al., 2003). We phenotyped these traits and other

taxonomically important traits in subadults of the mapping families (Tables S1 and S2).

As for sex determination, the XY system controlled by the male master sex-determining

gene *amhy* was demonstrated for *S. schlegelii*, *S. koreanus*, and *S. pachycephalus*.

The latter two species belong to the same subclade of *Sebastes* with *S. schlegelii* and

S. trivittatus (Song et al., 2021) (Figure S1). The gene *amhy* is present in a ca. 5 kb

male-specific region of the chromosome (CHR) 12 in the chromosome-level genome

assembly of *S. schlegelii* (He et al., 2019), which we used as a reference genome in

this study.

We measured standard length (SL) and body weight of each subadult fish and took

a digital image of the left side of the body along with a KODAK Colour Separation

Guide using Pentax K200D immediately after euthanasia. We kept the distance

between the camera and the fish constant across individuals. The sex of subadults was

determined through visual inspections of the overall appearance of the fresh gonads:

males have a pair of elongated, almost thread-like white gonads, whereas females

have partly rounded orange gonads. The visual inspection was validated for a subset of

individuals via histological observation. The fish was then fixed in 10% formalin and

preserved in 70% ethanol.

The body colour and the colour patterns of each fish were characterised in three

ways using the digital image (Figure S3). First, we coded the presence/absence of the

vertical bands on the dorsum and the horizontal stripes along the lateral line and along

the dorsal-fin base in each fish as binary traits based on visual inspection. The coding

was done twice by the same person for consistency. Second, we quantified the overall

intensities of black pigmentation (supposedly melanophore-based) and yellow

pigmentation (xanthophore-based) on the body by counting the number of

corresponding pixels. Each image was colour-corrected by setting the grey patch of the

colour separation guide as the neutral grey using the eyedropper tool in the Curves

adjustment implemented in Adobe Photoshop CS6. We then cropped the image by a

polygon defined by 11 landmarks on the body to restrict the analysis to an

unambiguously definable area, excluding the fins, which were sometimes damaged.

We counted the number of pixels on the cropped image that fell within the ranges of

the hue/saturation/brightness (HSB) colour space using the Color Threshold function of

ImageJ 1.53c (Schneider et al., 2012); black: H = 0–255, S = 0–255, B = 0–100;

yellow: H = 10–50, S = 0–255, B = 100–200. These ranges were determined by

preliminary visual inspections of the colour of the extracted pixels, of a subset of

individuals, under variable combinations of HSB values. Third, we quantified the colour

patterns of the fish using the 'patternize' v0.0.2 package (Van Belleghem et al., 2018)

on R v3.6.3 (R Core Team, 2020). We cropped the colour-corrected digital image by a

polygon defined by six landmarks on the body to restrict the analysis to the upper half

of the flank, where the focal colour patterns are expressed. We used the *patRegK*

function to automatically align all images, assign pixel colours to a predefined number

of clusters (K), and extract the colour patterns from the image based on the distribution

of each cluster. We analysed with K = 4–6 (Figure S4). The variation in the extracted

pattern was summarised by a Principal Component Analysis (PCA), performed for each

colour cluster of each K using the *patPCA* function. For QTL mapping, we used the PC

scores derived from three colour clusters from K = 5, which captured the colour pattern

variation most successfully (see Appendix S1 for more detail).

We quantified the overall body shape from the lateral view using geometric

morphometrics. We used tpsDig2 (Rohlf, 2017) to digitize 16 landmarks on the digital

image of each fish (Figure S3). Generalized Procrustes Analysis was performed on

these landmarks using the *gpagen* function implemented in the R package 'geomorph'

v4.0.0 (Adams et al., 2021). The variation in the obtained Procrustes shape

coordinates among individuals was summarized by a PCA using the *gm.prcomp*

function. Linear measurements and counts were made on 34 and 13 traits,

respectively, on the left side of the body of the fixed samples. Methods for measuring

and counting followed Muto et al. (2011, 2019) with the following additions: head

spines, the number of spines on the dorsal aspect of the cranium; lacrimal spines, the

number of spines on the lacrimal.

We $\log_{10}$ transformed SL, weight, all linear measurements, and intensities of

pigmentations. We tested the dependence of each trait on size (SL) and sex in each

family, and the difference between the two families, using either linear regression, the

Wilcoxon's rank-sum test, or the Fisher's exact test, depending on the distribution of

the trait (Tables S1, S2). The traits used for QTL mapping were tested for pairwise

correlations using the *corr.test* function implemented in the R package 'psych' v2.1.6

(Revelle, 2021), with the false discover rate (FDR) correction. We used the residuals of

linear regression of each trait on size or sex for the correlation test, if the trait was

dependent on these variables.

2.3 Genotyping

All parental individuals, 137 BC_Str individuals (48 larvae and 89 subadults) and 136

BC_Ssc individuals (48 larvae and 88 subadults) were subjected to genotyping by

double-digest restriction-site-associated DNA sequencing (ddRAD-seq), according to

Peterson et al. (2012) with minor modifications (see Appendix S1 for details). For

subadults of BC_Str and their parents, a single library was prepared and sequenced

with 50-bp single-end reads on an Illumina HiSeq 2500 system. A single library was

prepared and sequenced with 150-bp paired-end reads on an illumina HiSeq X system

for the remaining individuals. In either case, the sequencing was performed by

Macrogen Japan Corporation (Kyoto, Japan). We used two different sequencing

platforms because the availability of related facilities changed during the study. Only

the forward sequences from the paired-end reads were used for consistency across the

platforms. Parental individuals were sequenced twice or thrice for stringent genotyping.

We processed the ddRAD-seq reads using fastp v0.23.1 (Chen et al., 2018) to

remove adapter sequences and trim the lengths to 50 bp. Reads with one or more

bases with q-values < 20 were discarded. Two individuals of BC_Str subadults and an

individual of BC_Ssc subadults were removed because of the few reads. The trimmed

sequences were mapped to the chromosome-level reference genome assembly of a

male _S. schlegelii_ (He et al., 2019) (China National GeneBank Sequence Archive

accession No. CNA0000824), using BWA-mem v0.7.17 (Li, 2013) with a mismatch

penalty of 5. Unmapped reads and reads with a mapping quality < 5 were removed by

Samtools v1.12 (Danecek et al., 2021). The resulting BAM files were used for

identifying SNPs using 'gstacks' implemented in Stacks v2.5.4 (Catchen et al., 2013)

with the default settings. The subsequent filtering steps were performed separately for

each family to maximize the number of usable SNPs in the downstream analyses. We

created a vcf file using 'populations' implemented in Stacks, retaining only the first SNP

of a locus if two or more were detected. Genotypes with read depths < 10 or > 300

were removed, and SNPs were discarded if the resulting genotyping rate per SNP was

< 0.6, using VCFtools v0.1.16 (Danecek et al., 2011). We then used SnpSift v4.3

(Cingolani et al., 2012) to retain only the SNPs for which the alleles observed in

backcross progeny could be unambiguously determined to be inherited from either _S._

trivittatus or _S. schlegelii_. This was done by keeping only SNPs for which genotypes of

grandparents and parents conformed to all the following criteria: 1) either homozygous

in both grandparents or heterozygous in only one of the grandparents, 2) heterozygous

in the F_1 male parent, and 3) homozygous in the pure female parent. We then removed

the SNPs genotyped in < 60 BC individuals.

2.4 Linkage map and QTL

We first constructed a hybrid linkage map for each family using the R package 'R/qtl'

(Arends et al., 2010) and then a consensus map of the two families using the R

package 'LPmerge' v1.7 (Endelman & Plomion, 2014). The map for each family was

constructed as follows: We removed markers showing deviation of genotype frequency

from the expected even ratio of homozygotes and heterozygotes (hereafter referred to

as single-locus TRD; chi-square test, $p < 0.001$ after FDR correction). Linkage groups

(LGs) were initially formed with a minimum LOD score of 6 and a maximum

recombination fraction (RF) of 0.35. The initial linkage grouping was highly consistent

with the reference genome assembly of *S. schlegelii*, most markers on each LG being

mapped to one of the 24 CHRs of the reference genome. However, deviations from this

pattern were observed; a few markers on each LG mapped to an alternative (non-

syntenic) CHR, and some markers formed an extra 25th LG. Such markers were either

discarded, manually reassigned to another LG, or left unmodified based on

examinations of individual LOD scores and RFs (Appendix S1; Figure S5; Table S3).

Next, we ordered markers in each LG using the *orderMarkers* function. We checked

the quality of the initial order by inspecting the plots of LOD score and RF between

pairs of markers on each LG and corrected putative errors using either the *switch.order*

or the *orderMarkers* function. We also sought better orders for LOD score and

chromosome length using the *ripple* function. To identify problematic markers, we

investigated the change in the length of each LG by eliminating one marker at a time

using the *droponemarker* function. We removed the interior marker whose omission

decreased the length by > 10 centimorgans (cM). We inferred genotyping errors using

the *calc.errorlod* function and removed the genotypes with error LOD scores > 6. We

also removed one outlier individual of each of BC_Str larvae and subadults concerning

the numbers of crossovers. We then used the *ripple* function to search for marker

orders with higher LOD scores. Finally, we compared the LOD score of the resulting

marker order of each LG with that of the alternative order based on the physical

position of markers on the corresponding CHR of the reference genome, and retained

the order with a higher LOD score. If the order based on the physical position was

chosen, we applied the *ripple* function again to search for orders with further higher

LOD scores.

We conducted QTL mapping separately for each family based on the consensus

map using the R package 'R/qtl2' v0.24 (Broman et al., 2019). Among the phenotyped

traits, colour-related traits, head spination, numbers of gill-rakers, lateral-line pores,

and fin rays have been shown to differ between *S. trivittatus* and *S. schlegelii*; however,

the difference in the other traits is unknown (Nakabo & Kai, 2013). Therefore, for QTL

mapping, we used only the traits for which significant differences were observed

between the two backcross families, assuming that such differences represent the

difference between the parental species. We used family-specific markers as well as

markers shared by the two families. We first inserted pseudomarkers in the linkage

map at intervals of 1 cM. We conducted a genome scan for each trait using the

scanone function based on the Haley-Knott regression (Haley & Knott, 1992). We

accounted for the effects of size (SL) or sex by using these variables as additive

covariates if the focal trait was significantly associated with them. Genome scans for

body weight were performed with or without size as a covariate to differentiate potential

QTLs of growth and condition. We conducted 10,000 permutations to estimate the

genome-wide significance threshold of LOD score at alpha = 0.05 using the *scan1perm*

function. We estimated a 95% confidence interval (CI) for each QTL using the

bayes_int function. We estimated the effect of a QTL by calculating the proportion of

phenotypic variance explained (PVE): PVE = (sum of squares between genotypes) /

(sum of squares total) (e.g., Takahashi et al., 2021). For the calculation of PVE and the

visualisation of phenotypic variation by genotype, we used the residuals of the

regression of each trait on size or sex, if the trait was dependent on these variables;

otherwise, we used raw phenotype values.

2.5 TRD

We tested two classes of TRD. First, we tested the single-locus TRD of each marker

and assessed the distribution of the distorted markers on the reference genome. We

used both the markers that were or were not filtered out during the construction of the

linkage map for this analysis. Second, we tested non-random associations of

genotypes between pairs of markers. In the present mapping families, the following

two-marker genotypes are possible for a given pair of markers: StSt/StSt, StSt/StSc,

StSc/StSt, and StSc/StSc for BC_Str; ScSc/ScSc, ScSc/StSc, StSc/ScSc, and

StSc/StSc for BC_Ssc, with St and Sc representing alleles from *S. trivittatus* and *S.*

schlegelii, respectively, and genotypes of two markers separated by a slash. Negative

epistatic interaction (DMI) between alleles at loci near the markers results in

underrepresenting one or more of the two-marker genotypes, which is detectable as

non-random associations of genotypes (Berdan et al., 2021; Pritchard et al., 2011). We

calculated the expected two-marker genotype frequency from the observed genotype

frequency of each marker. We then compared the observed and the expected two-

marker genotype frequencies to detect non-random associations of genotypes using

the chi-square test. For this analysis, we reduced the markers so that the minimum

distance between adjacent markers became minimum 1 cM using the

pickMarkerSubset function in R/qtl. Consequently, a total of 47,694 tests among 316

markers and 59,580 tests among 353 markers were performed for BC_Str and

BC_Ssc, respectively. Distributions of the pairs of markers showing significant non-

random associations of genotypes were assessed on the linkage map rather than the

reference genome since the analysed markers had been filtered based on their

positions on the linkage map. Both types of TRD were analysed separately for larvae

and subadults of each family and corrected for FDR using the *p.adjust* function in R.

Results

3.1 Linkage map

Genotype data after filtering consisted of 642 and 823 markers for BC_Str and

BC_Ssc, respectively, with 545 markers common between them, and both families

consisted of 135 individuals (87 subadults; 47 larvae). No marker was mapped to the

unanchored sequences of the reference genome assembly. The hybrid linkage maps

constructed separately for each family consisted of 24 LGs, consistent with karyotypes

of both parental species as well as the reference genome (Figure 1) (Arai, 2011). The

consensus map consisted of 898 markers, including 636 markers for BC_Str, 799

markers for BC_Ssc, and 537 shared markers, with a total length of 1413.5 cM and an

average marker interval of 1.6 cM (Table S4). Each LG was syntenic to one of the 24

CHRs of the reference genome, showing high collinearity (Figure S6). Thus, the LGs

were numbered according to the corresponding CHRs. Exceptions to the syntenic

relationships were one or a few markers on LGs 5, 12, 15, 19, 20, and 22,

unequivocally assigned to these LGs in both BC_Str and BC_Ssc maps while being

mapped to CHRs 7, 7, 11, 21, 22, 18, and 16, respectively (Table S3). Each LG had a

'recombination coldspot', where little or no recombination was observed across a wide

range of the corresponding CHR of the reference genome (Figure S6). These regions

were positioned distally on each LG except for that observed in the middle of LG 4,

consistent with 23 acrocentric and one metacentric chromosome in both parental

species (Ida et al., 1982).

3.2 TRD

Distributions of genotype frequencies and single-locus TRD across the reference

genome are shown in Figures 2, S7, and S8. Significant TRD at $p < 0.05$ after FDR

correction was observed for a total of 131 markers, most of which showing TRD in only

the subadults of either family (Figure 3; Table S5). In BC_Str, none of the distorted

markers was common in both stages. In BC_Ssc, six distorted markers were shared

between the stages, and one of them ('scaffold6_37022538') additionally showed TRD

in the subadults of BC_Str. This marker, positioned terminally on CHR 6, showed an

interesting pattern of distortion in which heterozygotes and homozygotes were

overrepresented in BC_Str and BC_Ssc, respectively. Only 34.1% of the distorted

markers of BC_Str subadults showed excesses of homozygotes. This largely reflects

the presence of a few CHRs harbouring extended regions with excesses of

heterozygotes (see below). In contrast, 100%, 92.6%, and 83.3% of the distorted

markers showed excesses of homozygotes in BC_Str larvae, BC_Ssc larvae, and

BC_Ssc subadults, respectively.

Several extended genomic regions, harbouring more than ca. 20 markers with

single-locus TRD in the same direction, were recorded (Figure 2). In BC_Str subadults,

CHR 2 exhibited an excess of heterozygotes across its entire length. Approximately 27

Mb of CHR 6 showed an excess of heterozygotes in BC_Str and, in turn, an excess of

homozygotes in BC_Ssc. CHRs 10 and 14 in BC_Ssc and BC_Str subadults,

respectively, showed excesses of homozygotes across more than ca. 20 Mb. These

regions on CHRs 6, 10, and 14 roughly corresponded to the recombination coldspots

(Figure S6). Smaller clusters of 2–5 distorted markers spanning ca. 1.2–3.8 Mb were

also observed in CHRs 7, 21, and 23 in BC_Str subadults and CHRs 4 and 19 in

BC_Ssc subadults (Figures S7, S8). The remaining markers showing single-locus TRD

were interspersed across the genome without forming a distinct cluster.

Tests for non-random associations of genotypes between pairs of markers on

separate LGs yielded no significant result after FDR correction (all $p > 0.42$), partly

because of numerous tests, making the tests highly conservative. We, therefore,

focused on the marker pairs showing associations at nominal $p < 0.01$ and additional

supports for the associations. The additional supports were the clustering of multiple

marker pairs whereby two or more adjacent markers on a LG (hereafter referred to as a

marker block) showed associations with another marker block on a different LG. Fifty

five such associations (pairs of marker blocks) were found (Figure 4; Table S6);

however, none of these was common across developmental stages or families, i.e.,

none of the associations observed in different stages/families involved marker blocks

with overlapping ranges on both LGs. The numbers of the observed associations were

comparable between the developmental stages in each family (Figure 3).

Sex ratio was slightly male-biased in subadults of BC_Str but did not differ

significantly from parity (female:male = 0.40:0.60, n = 80; $\chi^2 = 0.89$, df = 1, $p = 0.35$),

and was more even in BC_Ssc (female:male = 0.48:0.52, n = 95; $\chi^2 = 0.09$, df = 1, $p = $

0.75).

3.3 QTL mapping

Thirty-nine morphological traits were subjected to QTL mapping, showing significant

differences between the two families (Figure S9; Tables S1, S2). Significant QTLs with

LOD scores greater than the genome-wide thresholds were detected for 11 and 9 traits

in BC_Str and BC_Ssc, respectively (Figure 1; Table 1; Figures S10, S11). Single

significant QTL for each trait was detected, except for yellow pigmentations of BC_Str,

where two QTLs were detected on different LGs. As for colour-related traits, QTLs for

the following traits were detected: yellow pigmentations for both families (on different

LGs); black pigmentations and PC1 scores based on the 1st and the 3rd colour

clusters of the patternize analysis for BC_Str; vertical bands on dorsum and horizontal

stripe along the lateral line for BC_Ssc. The 1st and the 3rd colour clusters of the

patternize analysis represented the variations in a stripe along the lateral line and

vertical bands on the dorsum, respectively. As for body shape and counts, QTLs for the

following traits were detected: lower caudal-peduncle length in both families (on

different LGs); PC2 scores based on the geometric morphometrics, predorsal length,

anal-fin base length, 2nd anal-fin spine length, and number of head spines in BC_Str;

orbit length, interorbital width, and numbers of pectoral-fin rays and gill-rakers on lower

gill arch in BC_Ssc. PC2 scores based on the geometric morphometrics represented

variations in the relative positions of the head and body. Directions of the QTLs were
consistent with the differences between the parental species except for the following
traits: predorsal length and the 3rd colour cluster of the patternize analysis in BC_Str;
lower caudal-peduncle length and yellow pigmentations in BC_Ssc. The QTL for a
horizontal stripe along the lateral line exhibited the highest PVE among all
morphological traits. In addition, QTL for sex with a high LOD score was detected on
LG 12 in each family, with the position roughly corresponding to that of the *amhy* gene:
26.4 Mb on the reference genome (Song et al., 2021), ca. 25 cM on the consensus
map (Figure S6).
The QTLs for the following traits had overlapping CIs: PC2 scores of geometric
morphometrics and 2nd anal fin length on LG 1, yellow and black pigmentations on LG
5, sex, yellow pigmentations, and the 1st colour cluster of the patternize analysis on LG
12 of BC_Str; orbit and lower caudal-peduncle lengths on LG 15 of BC_Ssc.
489
490

491 **4. Discussion**
492 **4.1 TRD and postmating isolation**
493 **4.1.1 General pattern**
494
We observed tens of markers showing two classes of TRD: single-locus TRD and non-
random associations of genotypes between pairs of markers. These markers,
presumably linked to the loci that conferred survival variation among hybrid progeny
and hence the postmating isolation between *S. trivittatus* and *S. schlegelii* (barrier loci),
were widely distributed across the genome and some of them encompassed large
proportions of particular chromosomes. Such a genetic architecture suggests genetic
redundancy and the multifarious nature of the postmating isolation. These features
would strongly reduce gene flow across the genome if the two species did mate,
thereby enabling the postmating isolation to play a critical role in completing speciation
or maintaining distinct gene pools (Coughlan & Matute, 2020; Nosil et al., 2009). Three
of the four extensive genomic regions with single-locus TRD in subadults of one or both
families coincided with the recombination coldspots. This suggests that each of these
three regions represents the isolating effects of one or a few barrier loci affecting broad
surrounding regions due to the tight linkage, rather than the effects of more numerous
and widespread barrier loci. Overall, the combined effects of the recombination

landscape and the effects of individual barrier loci created the observed complex

genetic architecture.

TRD was observed in each of the four samples differing in the developmental

stages and the directions of the crosses, with only a few distorted loci shared between

them (Figure 3). TRD observed in larvae can be ascribed to postmating isolating

barriers acting at the prenatal stage, from the artificial insemination until the parturition,

including gametic isolation and hybrid inviability. TRD in subadults can be based on

hybrid inviability at the postnatal stage from the parturition until sacrifice. In fact, the

distinct genetic architectures observed for larvae and subadults suggest that largely

different isolating mechanisms have been acting at the pre and postnatal stages. On

the other hand, asymmetric isolation has been frequently reported in previous studies

employing reciprocal interspecific crosses, and its primary cause is thought to be hybrid

incompatibility involving uniparentally inherited genetic factors or maternal effects

(Turelli & Moyle, 2007). More discussion on specific mechanisms of TRD in different

samples is provided in section 4.1.2.

We found no significant bias in the sex ratio in either family, suggesting little or no

difference in the fitness between sexes. We also observed no tendency for markers on

the sex chromosome CHR 12 to be more frequently involved in either type of TRD.

These results suggest that, regarding hybrid inviability, the present system does not

obey the two 'rules' concerning reproductive isolation: disproportionate contribution of X

or Z chromosomes to postmating isolation (large X effect) and more severe fitness

reduction of heterogametic sex in hybrid crosses (Haldane's rule) (Coyne & Orr,

1989b). This observation is consistent with homomorphic, recombining sex

chromosomes in *Sebastes* (Arai, 2011; Song et al., 2021; this study) since most

explanations for the two rules require heteromorphic sex chromosomes (Coyne, 2018).

Teleosts generally do not obey Haldane's rule regarding hybrid inviability (reviewed in

Schilthuizen et al., 2011); the present system is considered one of such cases.

We found discordance between the current linkage map and the reference

genome of *S. schlegelii* regarding the assignments of some markers to LGs/CHRs

(Figure S6). Some of the markers also showed non-random associations of genotypes

with markers on another LG (Figure 4). These results are reminiscent of reciprocal

translocations between the parental species. However, reciprocal translocation in

parental species generally has more extensive effects on a hybrid linkage map

whereby LGs involved in the translocation are connected into a single LG via markers

on the translocated region (Farré et al., 2011; Fishman et al., 2013), which was not

observed herein (Figure S5). We suspect that this discordance resulted from slight

errors in our linkage map or the genome assembly. Further studies will be necessary to

clarify whether the discordance represents some form of chromosomal

rearrangements, an important mechanism of reproductive isolation and adaptive

divergence between taxa (e.g., Matschiner et al., 2022).

A previous genomic study by Behrens et al. (2021) has identified several genomic

regions showing elevated values of both F_{ST} and D_{XY} between each of the two pairs of

incipient species of *Sebastes* (*S. carnatus*–*S. chrysomelas* and *S. miniatus*–"*S.

crocotulus*"), both pairs belonging to different subclades from the present parental

species. These regions are supposedly involved in adaptive divergence and

reproductive isolation in the presence of gene flow in each pair. The regions were

located at 38.5–39.0 Mb on CHR 3, 18.5–20.0 Mb on CHR 4, 0.5–1.5 Mb on CHR 7,

10.5–11.0 Mb on CHR 9, and 30.0–32.5 Mb on CHR 13 on the same reference

genome as that used in the present study. None of these regions overlaps with the

extended TRD possibly associated with extrinsic selection in the present system (see

below). Instead, these regions except that on CHR 3 coincided with the recombination

coldspots (Figure S6), implicating conserved recombination landscapes across the

phylogeny of rockfishes and their role in determining the genetic architecture of

reproductive isolation (Burri et al., 2015; Nachman & Payseur, 2012).

4.1.2 Possible mechanisms of isolation

Owing to the present cross design and the distinct (asymmetric) genetic architectures

of the two families, we suggest that he most likely isolating mechanisms causing TRD

at the prenatal stage are gametic isolation and hybrid inviability involving maternal

factors. Under the present cross design, both F_1 sperm and embryos of the two families

have been stored in ovaries of reciprocal maternal species during sperm storage and

gestation, with the embryos possessing maternally inherited genetic factors from their

respective maternal species (Figure S2). DMI involving maternally inherited factors

such as mito-nuclear incompatibility is a predominant mechanism of asymmetric

reproductive isolation in reciprocal hybrid crosses (Turelli & Moyle, 2007). Empirical

evidence that mitochondrial genomes from reciprocal parental populations cause TRD

in distinct sets of chromosomes in reciprocal F_2 crosses has been reported for marine

copepod (Lima et al., 2019). Besides, interactions between maternally expressed

factors and sperm or zygotes within the mother's body can result in asymmetric

reproductive isolation (e.g., Devigili et al., 2018). Notably, such interactions are

predicted to be driven by mother-offspring conflict in viviparous, matrotrophic animals,

including rockfishes (viviparity-driven conflict hypothesis; Zeh & Zeh, 2000). In any

case, the mechanisms of the TRD at the prenatal stage might be genetically complex,

involving epistatic interactions among multiple nuclear loci, as well as the interactions

with maternally inherited or expressed factors.

The TRD at the postnatal stage may also be explained by multiple factors,

including intrinsic hybrid inviability due to DMI involving maternally inherited factors and

environmentally-dependent (extrinsic) inviability. The former can have a common

mechanistic and genetic basis with that acting at the prenatal stage. For instance, mito-

nuclear incompatibility may disrupt the metabolic function of mitochondria essential to

all developmental stages (Burton & Barreto, 2012). In fact, five loci showing significant

excess of homozygotes on CHRs 1, 4, 7, and 13 were shared between the two stages

in BC_Ssc. These may represent loci that confer fitness effects across the

developmental stages via intrinsic asymmetric hybrid inviability. The remaining distorted

loci that were not shared with the prenatal stage are considered to have been caused

by stage-specific intrinsic hybrid inviability or extrinsic inviability.

Extrinsic hybrid inviability occurs when ecologically divergent species produce

hybrids with intermediate phenotypes, and the hybrid suffers reduced fitness in the

parental environments (Nosil, 2012). The present study was not specifically designed

to examine this process. Nonetheless, ca. 20 Mb region of CHR 6 showing

heterozygote excess in BC_Str and homozygote excess in BC_Ssc suggest additive

selection against *S. trivittatus* alleles, implying extrinsic hybrid inviability under an

environment where *S. trivittatus* is less fit than *S. schlegelii* (Rundle & Whitlock, 2001).

The pattern also implicates immigrant inviability, a class of premating isolation whereby

immigrants of parental species suffer reduced fitness in their non-native habitats (Nosil,

2012). Additionally, regions with homozygote excess in BC_Ssc subadults (e.g., CHR

10) or heterozygote excess in BC_Str subadults (e.g., CHR 2) with no concurrent TRD

in the other family could also be associated with extrinsic selection against *S. trivittatus*

alleles with dominance effects. A possible environmental factor underlying the extrinsic

selection is water temperature. Species distribution models for *S. schlegelii* have

suggested that water temperature is the most important environmental variable

determining the natural distribution of the species, and it prefers water temperatures of

3–20 ºC (Chen et al., 2021). Although the thermal tolerance of *S. trivittatus* is unknown,

it is reasonable to assume that this species is adapted to colder waters than *S.*

schlegelii, given the more northern distributional of the former. On the other hand, the

water temperatures of the tanks were 3.1–21.3 ºC in the present study (see Appendix

S1). These temperatures are within the range tolerated by *S. schlegelii* but may have

deviated from that by *S. trivittatus*, thus, selecting against *S. trivittatus* alleles at

relevant loci.

The terminal locus ('scaffold6_37022538') of the distorted region of CHR 6

additionally showed TRD in larvae, although the distortion in BC_Str larvae was not

statistically significant (Figure 2; Table S5). The directions of the distortion were the

same between stages in each family (heterozygote excess in BC_Str; homozygote

excess in BC_Ssc), suggesting that the extrinsic selection may have acted at the

prenatal stage as well. Effects of ambient thermal conditions on prenatal viability of

offspring via mother's body temperature of viviparous snake have been previously

documented (Lourdais et al., 2004).

4.2 QTLs for morphological traits and sex determination

Among the morphological traits with significant QTLs detected, the following traits have

been previously suggested to be related to convergence or speciation of rockfishes:

orbit diameter, number of gill-rakers on the lower arch, head spination, intensities of

pigmentations, stripe along the lateral line, and dark bands on the dorsum. The colour

clusters based on the patternize analysis are also considered among such traits since

these are the quantitative representations of the colour patterns. We do not nominate

candidate genes underlying these traits, considering the broad CIs for each QTL,

presumably due to the limited sample sizes. Nonetheless, the present study constitutes

the first step toward understanding the genetic bases of the traits at the individual gene

level, which will, in turn, provide insights into the traits' adaptive significance and the

evolutionary trajectories within and across lineages (e.g., Kokita et al., 2021; Kratochwil

et al., 2018).

The variations in the body colour (overall intensities of pigmentations) and the

colour patterns in the present system were suggested to have partially independent

genetic bases. More specifically, genomic regions containing the QTLs for colour

related traits were divided into three classes: containing the QTLs for the intensities of

pigmentations, the colour patterns, and both (Figure 1). We suspect that these classes

reflect the functions of genes that confer the association of each genomic region with

the variations in the traits. For example, in the melanocortin system of teleosts,

tyrosinase-related protein 1 is involved in pigment formation and therefore can affect

the overall body colour, leaving the patterns unaffected, whereas Melanocortin 1

receptor is involved in the proliferation of chromatophores and therefore can affect both

the colour and the colour patterns (Cal et al., 2017; Singh & Nüsslein-Volhard, 2015).

Mutations in genes such as *leopard* in zebrafish are known to affect interactions among

chromatophores, affecting only the colour patterns (Singh & Nüsslein-Volhard, 2015).

The supposed partially independent genetic bases for the colour and the colour-

patterns in the present system is consistent with the colour pattern convergence in

rockfishes, where independent lineages have similar colour patterns but not

necessarily similar colours.

The present result implicated a common sex-determination system for *S. schlegelii*

and *S. trivittatus*, i.e., male-determination by *amhy*. The role of *amhy* has been

originally demonstrated for *S. schlegelii* and additionally confirmed in *S.*

pachycephalus, and *S. koreanus* (Song et al., 2021). Under the present cross design,

amhy inherited from the male F_0 *S. schlegelii* segregated in each backcross family.

Besides, a highly significant QTL for sex was detected on LG 12 in each family, with

CIs encompassing the presumed position of *amhy*. Taken together, *amhy* from F_0 *S.*

schlegelii successfully determined sex in both families despite the contrasting degrees

of genetic contributions from *S. schlegelii* and *S. trivittatus*, implying that the sex of *S.*

trivittatus is also controlled by this gene. The sex-determination role of *amhy* might be

conserved across the members of the northwest Pacific subclade of the genus, to

which the aforementioned four species belong (Figure S1).

4.3 Concluding remarks

We provided a snapshot of postmating isolation in the speciation continuum of

rockfishes at the point where reproductive isolation is complete in nature. It is most

likely characterised by 1) the involvement of tens of loci across the genome, 2)

asymmetry of the underlying loci concerning the direction of the crosses, and 3)

multifarious mechanisms including asymmetric gametic isolation and hybrid inviability

due to interaction with maternal factors, and extrinsic selection. Future studies

employing direct observations of the isolating mechanisms at the phenotypic level, and

comparisons with other pairs of species at various points in the speciation continuum,

will contribute to a more comprehensive understanding of the speciation in rockfishes.

Besides, several limitations of the present study should also be noted. First, some of

the observed TRD may reflect isolating mechanisms that have evolved after the

completion of reproductive isolation. Second, the slight difference in rearing conditions

for the two families could have affected the results. Third, some of the TRD that were

unique to larvae of each family might be false positives. This is because markers

showing TRD in larvae should also do so in subadults unless the TRD in larvae is

counterbalanced by selective mortality at the postnatal stage. Finally, TRD may occur

due to mechanisms other than reproductive isolation between parental species, such

as inbreeding depression and DMI involving alleles segregating within species

(Corbett-Detig et al., 2013; Plough & Hedgecock, 2011). These issues can be partly

addressed by careful control of rearing conditions and appropriate cross designs (e.g.,

reciprocal F_2 crosses; Lima et al., 2019).

ACKNOWLEDGMENTS

We thank K. Tsuboi (Tokai University) and the staff of Mariculture Fisheries Research

Institute, Hokkaido Research Organization for their help in phenotyping and fish

husbandry, S. Kondo (Ryukoku University) for help in genotyping. Editors and

anonymous reviewers provided valuable comments on the manuscript. This study was

funded by The Yanmar Environmental Sustainability Support Association (grant No.

KI019078), The Hokusui Society Foundation, and JSPS KAKENHI (19K16217) to NM.

This study was done in compliance with the guidelines for the care and use of animals

for scientific purposes at Tokai University.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

NM, TK, YoS, and HT conceived the study. TK and YoS performed the crosses and

oversaw the fish husbandry. NM, TK, YuS, SI, YoS, and HT performed the phenotyping.

NM, RK, and AJN performed the genotyping. NM and RK performed the analyses. NM

wrote the manuscript. All authors read and approved the manuscript.

DATA AVAILABILITY STATEMENT

Sequence data have been deposited in Sequence Read Archive in DDBJ/EMBL-

EBI/NCBI under accession number DRA014198. Genotype and phenotype data and

linkage maps are available from Dryad.

REFERENCES

Adams, D. C., Collyer, M., Kaliontzopoulou, A., & Baken, E. (2021). *Geomorph:*

Software for geometric morphometric analyses. R package version 4.0.

https://cran.r-project.org/package=geomorph.

Arai, R. (2011). *Fish Karyotypes: A check list*. Springer.

Arends, D., Prins, P., Jansen, R. C., & Broman, K. W. (2010). R/qtl: High-throughput

multiple QTL mapping. *Bioinformatics*, *26*(23), 2990–2992.

https://doi.org/10.1093/bioinformatics/btq565

Behrens, K. A., Girasek, Q. L., Sickler, A., Hyde, J., & Buonaccorsi, V. P. (2021).

Regions of genetic divergence in depth-separated *Sebastes* rockfish species pairs:

Depth as a potential driver of speciation. *Molecular Ecology*, *30*(17), 4259–4275.

https://doi.org/10.1111/mec.16046

Berdan, E. L., Fuller, R. C., & Kozak, G. M. (2021). Genomic landscape of reproductive

isolation in *Lucania* killifish: The role of sex loci and salinity. *Journal of Evolutionary*

Biology, *34*(1), 157–174. https://doi.org/10.1111/jeb.13725

Brennan, A. C., Hiscock, S. J., & Abbott, R. J. (2014). Interspecific crossing and

genetic mapping reveal intrinsic genomic incompatibility between two *Senecio*

species that form a hybrid zone on Mount Etna, Sicily. *Heredity*, *113*(3), 195–204.

https://doi.org/10.1038/hdy.2014.14

Broman, K. W., Gatti, D. M., Simecek, P., Furlotte, N. A., Prins, P., Sen, Ś., Yandell, B.

S., & Churchill, G. A. (2019). R/qtl2: Software for mapping quantitative trait loci

with high-dimensional data and multiparent populations. *Genetics*, *211*(2), 495–

502. https://doi.org/10.1534/genetics.118.301595

Buonaccorsi, V. P., Narum, S. R., Karkoska, K. A., Gregory, S., Deptola, T., & Weimer,

A. B. (2011). Characterization of a genomic divergence island between black-and-

yellow and gopher *Sebastes* rockfishes. *Molecular Ecology*, *20*(12), 2603–2618.

https://doi.org/10.1111/j.1365-294X.2011.05119.x

Burri, R., Nater, A., Kawakami, T., Mugal, C. F., Olason, P. I., Smeds, L., Suh, A.,

Dutoit, L., Bureš, S., Garamszegi, L. Z., Hogner, S., Moreno, J., Qvarnström, A.,

Ružić, M., Sæther, S.-A., Sætre, G.-P., Török, J., & Ellegren, H. (2015). Linked

selection and recombination rate variation drive the evolution of the genomic

landscape of differentiation across the speciation continuum of *Ficedula*

flycatchers. *Genome Research*, *25*(11), 1656–1665.

https://doi.org/10.1101/gr.196485.115

Burton, R. S., & Barreto, F. S. (2012). A disproportionate role for mtDNA in

Dobzhansky–Muller incompatibilities? *Molecular Ecology*, *21*(20), 4942–4957.

https://doi.org/10.1111/mec.12006

Cal, L., Suarez-Bregua, P., Cerdá-Reverter, J. M., Braasch, I., & Rotllant, J. (2017).

Fish pigmentation and the melanocortin system. *Comparative Biochemistry and*

Physiology Part A: Molecular & Integrative Physiology, *211*, 26–33.

https://doi.org/10.1016/j.cbpa.2017.06.001

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013).

Stacks: An analysis tool set for population genomics. *Molecular Ecology*, *22*(11),

3124–3140. https://doi.org/10.1111/mec.12354

Chen, L., DeVries, A. L., & Cheng, C.-H. C. (1997). Convergent evolution of antifreeze

glycoproteins in Antarctic notothenioid fish and Arctic cod. *Proceedings of the*

National Academy of Sciences, *94*(8), 3817–3822.

https://doi.org/10.1073/pnas.94.8.3817

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics*, *34*(17), i884–i890.

https://doi.org/10.1093/bioinformatics/bty560

Chen, Y., Shan, X., Ovando, D., Yang, T., Dai, F., & Jin, X. (2021). Predicting current

and future global distribution of black rockfish (*Sebastes schlegelii*) under changing

climate. *Ecological Indicators*, *128*, 107799.

https://doi.org/10.1016/j.ecolind.2021.107799

Cingolani, P., Patel, V. M., Coon, M., Nguyen, T., Land, S. J., Ruden, D. M., & Lu, X.

(2012). Using *Drosophila melanogaster* as a Model for Genotoxic Chemical

Mutational Studies with a New Program, SnpSift. *Frontiers in Genetics*, *3*.

https://doi.org/10.3389/fgene.2012.00035

Corbett-Detig, R. B., Zhou, J., Clark, A. G., Hartl, D. L., & Ayroles, J. F. (2013). Genetic

incompatibilities are widespread within species. *Nature*, *504*(7478), 135–137.

https://doi.org/10.1038/nature12678

Coughlan, J. M., & Matute, D. R. (2020). The importance of intrinsic postzygotic
barriers throughout the speciation process. *Philosophical Transactions of the*
Royal Society B: Biological Sciences, *375*(1806), 20190533.
https://doi.org/10.1098/rstb.2019.0533

Coyne, J. A. (2018). "Two Rules of Speciation" revisited. *Molecular Ecology*, *27*(19),
3749–3752. https://doi.org/10.1111/mec.14790

Coyne, J. A., & Orr, H. A. (1989a). Patterns of speciation in *Drosophila*. *Evolution*,
43(2), 362–381. https://doi.org/10.1111/j.1558-5646.1989.tb04233.x

Coyne, J. A., & Orr, H. A. (1989b). Two rules of speciation. In *Speciation and Its*
Consequences. Sinauer Associates.

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sinauer Associates.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A.,
Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R.,
& 1000 Genomes Project Analysis Group. (2011). The variant call format and
VCFtools. *Bioinformatics*, *27*(15), 2156–2158.
https://doi.org/10.1093/bioinformatics/btr330

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,
Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve
years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008.
https://doi.org/10.1093/gigascience/giab008

Devigili, A., Fitzpatrick, J. L., Gasparini, C., Ramnarine, I. W., Pilastro, A., & Evans, J.
P. (2018). Possible glimpses into early speciation: The effect of ovarian fluid on
sperm velocity accords with post-copulatory isolation between two guppy
populations. *Journal of Evolutionary Biology*, *31*(1), 66–74.
https://doi.org/10.1111/jeb.13194

Endelman, J. B., & Plomion, C. (2014). LPmerge: An R package for merging genetic
maps by linear programming. *Bioinformatics*, *30*(11), 1623–1624.
https://doi.org/10.1093/bioinformatics/btu091

Evans, C. W., Hellman, L., Middleditch, M., Wojnar, J. M., Brimble, M. A., & Devries, A.
L. (2012). Synthesis and recycling of antifreeze glycoproteins in polar fishes.
Antarctic Science, *24*(3), 259–268. https://doi.org/10.1017/S0954102012000119

Farré, A., Lacasa Benito, I., Cistué, L., de Jong, J. H., Romagosa, I., & Jansen, J.
(2011). Linkage map construction involving a reciprocal translocation. *Theoretical*
and Applied Genetics, *122*(5), 1029–1037. https://doi.org/10.1007/s00122-010-
1507-2

Fishman, L., & McIntosh, M. (2019). Standard Deviations: The biological bases of

transmission ratio distortion. *Annual Review of Genetics*, *53*(1), 347–372.

https://doi.org/10.1146/annurev-genet-112618-043905

Fishman, L., Stathos, A., Beardsley, P. M., Williams, C. F., & Hill, J. P. (2013).

Chromosomal rearrangements and the genetics of reproductive barriers in

Mimulus (monkey flowers). *Evolution*, *67*(9), 2547–2560.

https://doi.org/10.1111/evo.12154

Haley, C. S., & Knott, S. A. (1992). A simple regression method for mapping

quantitative trait loci in line crosses using flanking markers. *Heredity*, *69*(4), 315–

324. https://doi.org/10.1038/hdy.1992.131

He, Y., Chang, Y., Bao, L., Yu, M., Li, R., Niu, J., Fan, G., Song, W., Seim, I., Qin, Y.,

Li, X., Liu, J., Kong, X., Peng, M., Sun, M., Wang, M., Qu, J., Wang, X., Liu, X., …

Qi, J. (2019). A chromosome-level genome of black rockfish, *Sebastes schlegelii*,

provides insights into the evolution of live birth. *Molecular Ecology Resources*,

19(5), 1309–1321. https://doi.org/10.1111/1755-0998.13034

Heras, J., & Aguilar, A. (2019). Comparative transcriptomics reveals patterns of

adaptive evolution associated with depth and age within marine rockfishes

(*Sebastes*). *Journal of Heredity*, *110*(3), 340–350.

https://doi.org/10.1093/jhered/esy070

Hyde, J. R., & Vetter, R. D. (2007). The origin, evolution, and diversification of

rockfishes of the genus *Sebastes* (Cuvier). *Molecular Phylogenetics and Evolution*,

44(2), 790–811. https://doi.org/10.1016/j.ympev.2006.12.026

Ida, H., Iwasawa, T., & Kamitori, M. (1982). Karyotypes in eight species of *Sebastes*

from Japan. *Japanese Journal of Ichthyology*, *29*, 162–168.

https://doi.org/10.11369/jji1950.29.162

Ingram, T. (2011). Speciation along a depth gradient in a marine adaptive radiation.

Proceedings of the Royal Society B: Biological Sciences, *278*(1705), 613–618.

https://doi.org/10.1098/rspb.2010.1127

Ingram, T., & Kai, Y. (2014). The geography of morphological convergence in the

radiations of Pacific *Sebastes* rockfishes. *The American Naturalist*, *184*(5), E115–

E131. https://doi.org/10.1086/678053

Kai, Y., Nakayama, K., & Nakabo, T. (2003). Molecular phylogenetic perspective on

speciation in the genus *Sebastes* (Scorpaenidae) from the Northwest Pacific and

the position of *Sebastes* within the subfamily Sebastinae. *Ichthyological Research*,

50(3), 239–244. https://doi.org/10.1007/s10228-003-0163-9

Kawasaki, T., Yohei, S., Tatsunari, M., Hiramatsu, N., & Todo, T. (2017). Development
of artificial insemination techniques for viviparous black rockfish (*Sebastes*
schlegelii). *Aquaculture Science*, *65*(1), 73–82.

Kokita, T., Ueno, K., Yamasaki, Y. Y., Matsuda, M., Tabata, R., Nagano, A. J., Mishina,
T., & Watanabe, K. (2021). Gudgeon fish with and without genetically determined
countershading coexist in heterogeneous littoral environments of an ancient lake.
Ecology and Evolution, *11*(19), 13283–13294. https://doi.org/10.1002/ece3.8050

Kolora, S. R. R., Owens, G. L., Vazquez, J. M., Stubbs, A., Chatla, K., Jainese, C.,
Seeto, K., McCrea, M., Sandel, M. W., Vianna, J. A., Maslenikov, K., Bachtrog, D.,
Orr, J. W., Love, M., & Sudmant, P. H. (2021). Origins and evolution of extreme life
span in Pacific Ocean rockfishes. *Science*, *374*(6569), 842–847.
https://doi.org/10.1126/science.abg5332

Kratochwil, C. F., Gerwin, J., Woltering, J. M., Meyer, A., Hulsey, C. D., Liang, Y.,
Urban, S., Machado-Schiaffino, G., & Henning, F. (2018). Agouti-related peptide 2
facilitates convergent evolution of stripe patterns across cichlid fish radiations.
Science, *362*(6413), 457–460. https://doi.org/10.1126/science.aao6809

Kulmuni, J., Butlin, R. K., Lucek, K., Savolainen, V., & Westram, A. M. (2020). Towards
the completion of speciation: The evolution of reproductive isolation beyond the
first barriers. *Philosophical Transactions of the Royal Society B: Biological*
Sciences, *375*(1806), 20190528. https://doi.org/10.1098/rstb.2019.0528

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. *ArXiv*, *1303.3997v2*, 1–3. http://arxiv.org/abs/1303.3997

Lima, T. G., Burton, R. S., & Willett, C. S. (2019). Genomic scans reveal multiple
mito‐nuclear incompatibilities in population crosses of the copepod *Tigriopus*
californicus. *Evolution*, *73*(3), 609–620. https://doi.org/10.1111/evo.13690

Longo, G. C., Harms, J., Hyde, J. R., Craig, M. T., Ramón-Laca, A., & Nichols, K. M.
(2022). Genome-wide markers reveal differentiation between and within the cryptic
sister species, sunset and vermilion rockfish. *Conservation Genetics*, *23*(1), 75–89.
https://doi.org/10.1007/s10592-021-01397-4

Lourdais, O., Shine, R., Bonnet, X., Guillon, M., & Naulleau, G. (2004). Climate affects
embryonic development in a viviparous snake, *Vipera aspis*. *Oikos*, *104*(3), 551–
560. https://doi.org/10.1111/j.0030-1299.2004.12961.x

Love, M. S., Yoklavich, M. M., & Thorsteinson, L. K. (2002). *The rockfishes of the*
northeast Pacific. University of California Press.

Macfarlane, R., & Bowers, M. (1995). Matrotrophic viviparity in the yellowtail rockfish

889    *Sebastes flavidus*. *Journal of Experimental Biology*, *198*(5), 1197–1206.

890    https://doi.org/10.1242/jeb.198.5.1197

891 Maheshwari, S., & Barbash, D. A. (2011). The Genetics of Hybrid Incompatibilities.

892    *Annual Review of Genetics*, *45*(1), 331–355. https://doi.org/10.1146/annurev-

893    genet-110410-132514

894 Matschiner, M., Barth, J. M. I., Tørresen, O. K., Star, B., Baalsrud, H. T., Brieuc, M. S.

895    O., Pampoulie, C., Bradbury, I., Jakobsen, K. S., & Jentoft, S. (2022). Supergene

896    origin and maintenance in Atlantic cod. *Nature Ecology & Evolution*, *6*(4), 469–481.

897    https://doi.org/10.1038/s41559-022-01661-x

898 Matute, D. R., Butler, I. A., Turissini, D. A., & Coyne, J. A. (2010). A test of the snowball

899    theory for the rate of evolution of hybrid incompatibilities. *Science*.

900    https://doi.org/10.1126/science.1193440

901 Matute, D. R., & Cooper, B. S. (2021). Comparative studies on speciation: 30 years

902    since Coyne and Orr. *Evolution*, *75*(4), 764–778. https://doi.org/10.1111/evo.14181

903 Muto, N., Kai, Y., & Nakabo, T. (2011). Genetic and morphological differences between

904    *Sebastes vulpes* and *S. zonatus* (Teleostei: Scorpaeniformes: Scorpaenidae).

905    *Fishery Bulletin*, *109*, 429–439. https://spo.nmfs.noaa.gov/sites/default/files/pdf-

906    content/2011/1094/1094muto.pdf

907 Muto, N., Kai, Y., & Nakabo, T. (2019). Taxonomic review of the *Sebastes vulpes*

908    complex (Scorpaenoidei: Sebastidae). *Ichthyological Research*, *66*(1), 9–29.

909    https://doi.org/10.1007/s10228-018-0641-8

910 Muto, N., Kai, Y., Noda, T., & Nakabo, T. (2013). Extensive hybridization and

911    associated geographic trends between two rockfishes *Sebastes vulpes* and *S.*

912    *zonatus* (Teleostei: Scorpaeniformes: Sebastidae). *Journal of Evolutionary*

913    *Biology*, *26*(8), 1750–1762. https://doi.org/10.1111/jeb.12175

914 Nachman, M. W., & Payseur, B. A. (2012). Recombination rate variation and

915    speciation: Theoretical predictions and empirical results from rabbits and mice.

916    *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1587),

917    409–421. https://doi.org/10.1098/rstb.2011.0249

918 Nakabo, T., & Kai, Y. (2013). Sebastidae. In *Fishes of Japan, with pictorial keys to*

919    *species.* (3rd edition). Tokai University Press.

920 Nosil, P. (2012). *Ecological Speciation.* Oxford Univ. Press.

921 Nosil, P., Harmon, L. J., & Seehausen, O. (2009). Ecological explanations for

922    (incomplete) speciation. *Trends in Ecology & Evolution, 24*(3)*,* 145–156.

923    https://doi.org/10.1016/j.tree.2008.10.011

Orr, H. A. (1995). The population genetics of speciation: The evolution of hybrid
incompatibilities. *Genetics*, *139*(4), 1805–1813.
https://doi.org/10.1093/genetics/139.4.1805

Orr, H. A., & Turelli, M. (2001). The evolution of postzygotic isolation: accumulating
Dobzhansky-Muller incompatibilities. *Evolution*, *55*(6), 1085–1094.
https://doi.org/10.1111/j.0014-3820.2001.tb00628.x

Parin, N. V., Evseenko, S. A., & Vasil'eva, E. D. (2014). *Fishes of Russian Seas:*
Annotated Catalogue. KMK.

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012).
Double Digest RADseq: An inexpensive method for de novo SNP discovery and
genotyping in model and non-model species. *PLOS ONE*, *7*(5), e37135.
https://doi.org/10.1371/journal.pone.0037135

Plough, L. V., & Hedgecock, D. (2011). Quantitative trait locus analysis of stage-
specific inbreeding depression in the Pacific oyster *Crassostrea gigas*. *Genetics*,
189(4), 1473–1486. https://doi.org/10.1534/genetics.111.131854

Pritchard, V. L., Dimond, L., Harrison, J., S Velázquez, C., Zieba, J. T., Burton, R. S., &
Edmands, S. (2011). Interpopulation hybridization results in widespread viability
selection across the genome in *Tigriopus californicus*. *BMC Genetics*, *12*(1), 54.
https://doi.org/10.1186/1471-2156-12-54

R Core Team. (2020). *R: A language and environment for statistical computing*. R
Foundation for Statistical Computing.

Revelle, W. (2021). *psych: Procedures for Psychological, Psychometric, and*
Personality Research. (2.1.6). Northwestern University. https://CRAN.R-
project.org/package=psych

Rohlf, F. J. (2017). *tpsDig, digitize landmarks and outlines, version 2.31*. Department of
Ecology and Evolution, State University of New York at Stony Brook.
http://www.sbmorphometrics.org/soft-dataacq.html

Rometsch, S. J., Torres-Dowdall, J., & Meyer, A. (2020). Evolutionary dynamics of pre-
and postzygotic reproductive isolation in cichlid fishes. *Philosophical Transactions*
of the Royal Society B: Biological Sciences, *375*(1806), 20190535.
https://doi.org/10.1098/rstb.2019.0535

Rosenblum, E. B., Parent, C. E., & Brandt, E. E. (2014). The Molecular basis of
phenotypic convergence. *Annual Review of Ecology, Evolution, and Systematics*,
45(1), 203–226. https://doi.org/10.1146/annurev-ecolsys-120213-091851

Rundle, H. D., & Whitlock, M. C. (2001). A genetic interpretation of ecologically
dependent isolation. *Evolution*, *55*(1), 198–201. https://doi.org/10.1111/j.0014-
3820.2001.tb01284.x

Saha, A., Johansen, T., Hedeholm, R., Nielsen, E. E., Westgaard, J.-I., Hauser, L.,
Planque, B., Cadrin, S. X., & Boje, J. (2017). Geographic extent of introgression in
Sebastes mentella and its effect on genetic population structure. *Evolutionary*
Applications, *10*(1), 77–90. https://doi.org/10.1111/eva.12429

Schemske, D. W. (2010). Adaptation and the origin of species. *The American*
Naturalist, *176*(S1), S4–S25. https://doi.org/10.1086/657060

Schilthuizen, M., Giesbers, M. C. W. G., & Beukeboom, L. W. (2011). Haldane's rule in
the 21st century. *Heredity*, *107*(2), 95–102. https://doi.org/10.1038/hdy.2010.170

Schluter, D. (2000). *The ecology of adaptive radiation*. Oxford Univ. Press.

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25
years of image analysis. *Nature Methods*, *9*(7), 671–675.
https://doi.org/10.1038/nmeth.2089

Schrader, M., & Travis, J. (2008). Testing the viviparity-driven-conflict hypothesis:
parent-offspring conflict and the evolution of reproductive isolation in a Poeciliid
fish. *The American Naturalist*, *172*(6), 806–817. https://doi.org/10.1086/592999

Schwenke, P. L., Park, L. K., & Hauser, L. (2018). Introgression among three rockfish
species (*Sebastes* spp.) in the Salish Sea, northeast Pacific Ocean. *PLOS ONE*,
13(3), e0194068. https://doi.org/10.1371/journal.pone.0194068

Shum, P., Pampoulie, C., Sacchi, C., & Mariani, S. (2014). Divergence by depth in an
oceanic fish. *PeerJ*, *2*, e525. https://doi.org/10.7717/peerj.525

Singh, A. P., & Nüsslein-Volhard, C. (2015). Zebrafish stripes as a model for vertebrate
colour pattern formation. *Current Biology*, *25*(2), R81–R92.
https://doi.org/10.1016/j.cub.2014.11.013

Song, W., Xie, Y., Sun, M., Li, X., Fitzpatrick, C. K., Vaux, F., O'Malley, K. G., Zhang,
Q., Qi, J., & He, Y. (2021). A duplicated *amh* is the master sex-determining gene
for *Sebastes* rockfish in the Northwest Pacific. *Open Biology*, *11*(7), 210063.
https://doi.org/10.1098/rsob.210063

Stelkens, R. B., Young, K. A., & Seehausen, O. (2010). The accumulation of
reproductive incompatibilities in African cichlid fish. *Evolution*, *64*(3), 617–633.
https://doi.org/10.1111/j.1558-5646.2009.00849.x

Stern, D. L. (2013). The genetic causes of convergent evolution. *Nature Reviews*
Genetics, *14*(11), 751–764. https://doi.org/10.1038/nrg3483

Takahashi, T., Nagano, A. J., & Sota, T. (2021). Mapping of quantitative trait loci

underlying a magic trait in ongoing ecological speciation. *BMC Genomics*, *22*(1),

615. https://doi.org/10.1186/s12864-021-07908-4

Tengfei, D., Yongshuang, X., Haixia, Z., Li, Z., Qinghua, L., Xueying, W., Jun, L.,

Shihong, X., Yanfeng, W., Jiachen, Y., Lele, W., Yunong, W., & Guang, G. (2021).

Multiple fetal nutritional patterns before parturition in viviparous fish *Sebastes*

schlegelii (Hilgendorf, 1880). *Frontiers in Marine Science*, *7*.

https://www.frontiersin.org/article/10.3389/fmars.2020.571946

Turelli, M., & Moyle, L. C. (2007). Asymmetric postmating isolation: Darwin's corollary

to Haldane's rule. *Genetics*, *176*(2), 1059–1088.

https://doi.org/10.1534/genetics.106.065979

Van Belleghem, S. M., Papa, R., Ortiz‐Zuazaga, H., Hendrickx, F., Jiggins, C. D.,

Owen McMillan, W., & Counterman, B. A. (2018). patternize: An R package for

quantifying colour pattern variation. *Methods in Ecology and Evolution*, *9*(2), 390–

398. https://doi.org/10.1111/2041-210X.12853

Venerus, L. A., Ciancio, J. E., Riva-Rossi, C., Gilbert-Horvath, E. A., Gosztonyi, A. E.,

& Garza, J. C. (2013). Genetic structure and different color morphotypes suggest

the occurrence and bathymetric segregation of two incipient species of *Sebastes*

off Argentina. *Naturwissenschaften*, *100*(7), 645–658.

https://doi.org/10.1007/s00114-013-1061-2

Zeh, D. W., & Zeh, J. A. (2000). Reproductive mode and speciation: The viviparity-

driven conflict hypothesis. *BioEssays*, *22*(10), 938–946.

https://doi.org/10.1002/1521-1878(200010)22:10<938::AID-BIES9>3.0.CO;2-9

TABLES

TABLE 1 Significant QTLs for body colouration, shape, counts, and sex. The 'Direction' column indicates whether *S.*

trivittatus (Str) or *S. schlegelii* (Ssc) alleles cause higher trait values (for continuous traits) or the presence of the focal trait

(for binary traits). Asterisks indicate allelic effects in the direction expected from the divergence between the parental species.

Trait	Family	LG	Pos (cM)	Pos (Mb)[a]	LOD	CI lower	CI upper	PVE	Direction
Colouration									
Colour cluster 1: PC1	BC_Str	12	0.0	0.7	3.00	0.0	26.2	14.6	Ssc*
Colour cluster 3: PC1	BC_Str	16	60.8	25.8	3.36	51.6	60.8	16.6	Str
Black pigmentation	BC_Str	5	11.0	1.3	2.77	0.0	45.9	6.2	Ssc*
Yellow pigmentation	BC_Str	5	34.1	11.4	3.21	0.0	47.4	12.2	Str*
Yellow pigmentation	BC_Str	12	15.0	16.7	3.31	0.0	18.5	11.8	Str*
Yellow pigmentation	BC_Ssc	22	58.0	23.4	5.86	43.9	60.5	22.9	Ssc
Stripe along lateral line	BC_Ssc	1	51.0	29.6	9.27	40.8	51.5	46.9	Str*
Dark bands on dorsum	BC_Ssc	2	35.3	23.0	3.41	17.5	48.7	19.0	Str*
Shape									
GM: PC2	BC_Str	1	40.8	27.2	3.05	37.8	60.2	15.8	Str*
Orbit diameter	BC_Ssc	15	6.0	12.9	3.25	0.0	54.0	6.4	Ssc*
Interorbital width	BC_Ssc	13	2.0	2.7	3.75	0.0	51.6	4.2	Ssc*
Lower peduncle length	BC_Str	12	35.6	32.8	3.19	29.2	42.5	5.7	Ssc*
Lower peduncle length	BC_Ssc	15	24.9	23.9	3.04	0.0	54.0	5.0	Str
Predorsal length	BC_Str	20	37.0	6.8	2.70	11.5	51.3	2.2	Ssc
Anal-fin base length	BC_Str	24	24.0	12.1	2.51	0.8	39.6	17.4	Str*

Anal-fin 2nd spine length	BC_Str	1	51.0	29.6	4.84	43.9	60.2	10.5	Str*
Counts									
Gill rakers on lower arch	BC_Ssc	3	23.0	32.3	2.78	0.0	60.5	15.5	Str*
Pectoral-fin rays	BC_Ssc	11	37.0	17.5	2.79	0.0	58.3	14.7	Ssc*
Spines on head	BC_Str	11	54.0	21.5	3.42	0.0	60.2	16.7	Str*
Sex	BC_Str	12	25.2	26.1	7.71	15.4	29.2	49.6	NA
Sex	BC_Ssc	12	21.0	27.6	20.80	18.5	23.2	94.9	NA

1021 [a] Based on the marker position on the reference genome assembly of *S. schlegelii*. For the QTL on pseudomarker, the

1022 physical position of the nearest marker is provided.

1023 Abbreviations: CI, confidence interval; cM: centimorgan; GM, geometric morphometrics; LG, linkage group; Mb, megabase;

1024 PVE, per cent variance explained.

 Figures

 FIGURE 1 Consensus linkage maps and 95% confidence intervals of positions of QTLs

 for colouration (blue bars), shape (orange), counts (red), and sex (grey): a, PC2 scores

 based on geometric morphometrics; b, Anal-fin 2nd spine length; c, black pigmentation;

 d, yellow pigmentation; e, head spination; f, PC1 scores based on colour cluster 1

 derived from patternize analysis; g, lower peduncle length; h, sex; i, PC1 scores based

 on colour cluster 3; j, predorsal length; k, anal-fin base length; l, stripe along lateral line;

 m, dark bands on dorsum; n, gill-rakers on lower arch; o, pectoral-fin rays; p,

 interorbital width; q, orbit diameter. The difference in the length of each linkage group

 between families is due to the presence of markers unique to either family (see also

 Table S4).

FIGURE 2 Genotype frequencies and transmission ratio distortion of markers on selected chromosomes, shown on the reference genome assembly of *S. schlegelii*. For each marker, frequencies of homozygotes (left column) and *p* values as $-\log_{10}p$ from the chi-square tests for deviations of genotype frequencies from Mendelian expectations corrected for false discovery rates (FDR) (right column) are given. Dotted line indicates FDR-corrected *p* = 0.05.

FIGURE 3 Venn diagrams for the number of markers showing significant single-locus
transmission ratio distortion (TRD) and the number of pairs of marker blocks (groups of
two or more adjacent markers) on separate linkage groups (LGs) showing non-random
associations of genotypes. The non-random associations of genotypes between pairs of
marker blocks were regarded as common across samples of different developmental
stages/families if the associations in different samples involved marker blocks with
overlapping ranges on both LGs (none observed in the present study).
1052

FIGURE 4 Possible epistatic interactions between pairs of markers on separate linkage groups (LGs). Each link represents a pair of marker blocks showing a non-random association of genotypes (chi-square test; nominal *p* < 0.01). Blue and yellow links correspond to the associations observed in larvae and subadults, respectively. The outer and inner circles represent each LG of the consensus linkage map and marker positions used for this analysis, respectively. Red ticks indicate the markers assigned to different (non-syntenic) LGs/chromosomes on the consensus linkage map and on the reference genome assembly of *S. schlegelii*.