SNV-FEAST: microbial source tracking with single nucleotide variants

Leah Briscoe[1]*, Eran Halperin[2,3,4,5,6], Nandita R. Garud[3,7]*

1. Bioinformatics Interdepartmental Program, University of California Los Angeles, Los Angeles, Los Angeles, CA, United States of America
2. Department of Computer Science, University of California Los Angeles, Los Angeles, CA, United States of America
3. Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, United States of America
4. Department of Computational Medicine, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, United States of America
5. Department of Anesthesiology and Perioperative Medicine, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, United States of America
6. Institute of Precision Health, University of California Los Angeles, CA, United States of America
7. Department of Ecology and Evolutionary Biology, University of California Los Angeles, CA, United States of America

*Correspondence to leahpbriscoe@gmail.com and ngarud@ucla.edu

ABSTRACT

Elucidating the sources of a microbiome can provide insight into the ecological dynamics responsible for the formation of these communities. "Source tracking" approaches to date leverage species abundance information, however, single nucleotide variants (SNVs) may be more informative because of their high specificity to certain sources. To overcome the computational burden of utilizing all SNVs for a given sample, we introduce a novel method to identify signature SNVs for source tracking. We show that signature SNVs used as input into a previously designed source tracking algorithm, FEAST, can more accurately estimate contributions than species and provide novel insights, demonstrated in three case studies.

Keywords: Source tracking, microbiome, single nucleotide variants, transmission, strains

BACKGROUND

Understanding the sources that could contribute to the formation of a given microbiome

is of great interest in elucidating the ecological processes that give rise to these complex

communities and the impact of these communities on human and environmental health. For

example, a hospital environment may introduce antibiotic resistance genes to an infant gut

microbiome, and local selective pressures may result in vastly different microbial compositions

in different parts of the ocean. Approaches for determining the proportion of a microbiome of

interest (the "sink") that is attributed to different microbiomes (the "sources") is known as

"source tracking" (Knights et al., 2011; Shenhav et al., 2019). Source tracking is useful for

forensics, categorization of samples, detecting contamination, and tracing transmissions between

different hosts or environments. While source tracking was developed as a way to quantitatively

characterize a sample based on a set of samples with known origin, in most studies, the true

source of samples may never be collected. In these cases, source tracking approaches are useful

in identifying similarities between microbiome samples even if they cannot be used to

definitively identify the true source of origin.

Current approaches for source tracking include the Bayesian approach, SourceTracker

(Knights et al., 2011) and more recently the expectation-maximization approach, FEAST

(Shenhav et al., 2019). These source tracking methods use species abundance profiles of the

sample of interest (the sink) and of potential sources and compute percentages of sinks that are

attributable to each potential source. However, species abundance profiles miss important sub-

species single nucleotide variants (SNVs), which may provide higher resolution information than

species about transmission patterns. For example, (Nayfach et al., 2016) found that the sharing of

microbiome SNVs private to mothers and their infants decreases over the first year of the

infant's life while species sharing increases. This suggests that while the infant microbiome

increasingly resembles the adult microbiome ecologically, sources other than the mother also

colonize the infant. Thus, species-level resolution may obscure true sources of microbes while

SNVs can reveal actual transmissions to the infant.

While tracking strain transmissions with SNVs has been highly successful in a number of

studies (Asnicar et al., 2017a; Ferretti et al., 2018; Korpela et al., 2018; Li et al., 2016; Nayfach

et al., 2016; Olm et al., 2021; Schmidt et al., 2019) current approaches to strain tracking are

limited. These methods provide binary information by inferring whether or not a strain

transmission has occurred per species but they do not shed light on the relative proportions of

microbiomes that are similar. A specific example of this is inStrain (Olm et al., 2021) which

computes a pairwise population-level average nucleotide identity (popANI) between two

samples. If an infant harbors several strains derived from the mother at low frequency, these

shared strains will have high popANI values, but they will represent a relatively small proportion

of the infant's microbiome. By contrast, source tracking allows us to simultaneously infer the

putative proportions for multiple sources contributing to a given sink, integrated over all

community members in the sink. As shown in **Figure 1**, one may be able to estimate that an

infant microbiome is explained 25% by the mother, 10% by the dog, and 30% by unknown

sources (Knights et al., 2011; Shenhav et al., 2019). In other words, source tracking with SNVs

leverages not only the genetic variants within species, but also the relative abundances of the

species that carry the SNVs.

Here, we evaluate whether source contributions estimated with SNVs are more accurate

than with only species when provided as inputs to FEAST (Shenhav et al., 2019) (hereafter

referred to as SNV-FEAST and species-FEAST, respectively). FEAST (Shenhav et al., 2019) is

faster and more accurate than previous source tracking tools (Knights et al., 2011), and therefore,

is ideal for adaptation to SNV source tracking since it can accept larger numbers of features and

input sources. Despite this improved computational efficiency, the potentially millions of single

nucleotide variants across all microbiome species in a given host still can computationally

overwhelm FEAST. To address this, we introduce a novel approach to determine signature SNVs

that can be used as input to FEAST. This both reduces memory requirements and computation

time in the FEAST estimation, allowing us to optimally estimate the source contribution of a

sink. We find that SNV-FEAST and species-FEAST yield different outcomes when applied to

simulated data, with SNV-FEAST frequently out-performing species-FEAST. We apply SNV-

FEAST to three real-world case studies, including source tracking between infants and their

mothers in the first year of life, between infants and the neonatal intensive care unit (NICU), and

between oceans around the world. We confirm the ability of SNV-FEAST by recapitulating

several previously published findings in our case studies, as well as discover new source tracking

patterns across oceans. In sum, we show that SNVs can be used to estimate potential

transmissions across hosts and across environments.

RESULTS

SNV-FEAST algorithm

Here we adapt FEAST to accept SNV abundance instead of species abundance as input.

A computational challenge in using SNVs instead of species as input to FEAST is that SNVs

contribute a significantly larger feature space. The number of different species comprising a

microbiome can range from a few hundred to a few thousand, while the number of possible

SNVs for a given species alone can be in the thousands (Schloissnig et al., 2013). This difference

in number of input features can result in FEAST runtimes that last several hours instead of a few

minutes and memory intensive storage of read counts at all sites of variation.

We devised a likelihood-based approach for selecting a set of informative or "signature"

SNVs for a given source tracking analysis, allowing us to overcome the time and memory

intensive challenges of utilizing SNV-level data. We identify these informative SNVs by

computing a signature score (**Figure 1A**) (see **Methods)** that quantifies the extent to which

SNVs in the sink are most likely derived from one of the potential sources. This is analogous to

identifying SNVs private to sources and their sinks, but more generalized to include SNVs that

may be found in multiple sources, albeit at higher frequency in one of the potential sources (see

Methods).

To compute a signature score for a given SNV, two hypotheses are compared for each

potential source: (1) that one source solely explains the observed allele counts in the sink and (2)

all sources except that one source collectively explain the observed allele counts in the sink. For

each hypothesis, we calculate the binomial log-likelihood for the estimate of the allele frequency

in the sink, θ.

Hypothesis 1: Source i with allele frequency p_i explains the allele counts in the sink.

$$\hat{\theta} = p_i$$

Hypothesis 2: A combination of all other sources except i (sources $j \neq i$) explain the observed

allele count distribution in the sink. The estimate of the sink allele frequency is computed using a

mixture of the allele frequencies p_j from those sources. The mixing parameter α_j is learned using

Sequential Least Squares Programming with the constraint that $\sum_{j \neq i} \alpha_j = 1$.

The binomial log-likelihood is calculated as follows, where there are *n* reads with the reference

allele and *m* reads with the alternative allele in the sink.

A log likelihood ratio representing the support for hypothesis 1 relative to hypothesis 2 is

calculated per site per potential source. The maximum log likelihood ratio per site is the

signature score for that SNV, representing how favorably one of the sources explains the sink

over all other sources. Signature SNVs are those with scores greater than two standard deviations

over the mean signature score computed for all SNVs (**Methods**).

Figure 1: Signature SNV selection and SNV-FEAST. (A) A signature SNV is present in one or

few but not all sources. By contrast, a non-signature SNV is generically present in multiple

sources and thus provides little discriminating information. (B) SNV-FEAST estimates the

proportion a given sink derived from various sources using the read counts for each allele in

sinks and sources.

Evaluation of SNV-FEAST in simulations

To compare the accuracy of species-FEAST and SNV-FEAST, we performed simulations mimicking mother-infant transmissions with the goal of estimating contributions of different sources to an infant sink. Our simulations tested the ability of SNVs and species to recapitulate the true source composition in synthetic samples comprised of a mixture of reads drawn from multiple real fecal adult samples. To construct these synthetic infant microbiomes, we mixed metagenomic data from mothers sampled in a mother-infant dataset (Bäckhed et al., 2015) at various proportions as described below (**Methods**).

The difficulty of source tracking increases with the number of contributing sources (Shenhav et al., 2019). Thus, we simulate infants that have a small (<=5) versus large (6 – 10) number of contributing sources (**Supplementary Table 1**), including an unknown source (e.g. a randomly selected unrelated mother). Known source contributions to the simulated gut microbiome sample of the infant were varied between 1 and 90% while the unknown contribution varied between 10 and 90%. The unknown source was not presented to FEAST as a potential known source.

Additionally, not all species in a mother are transmitted to the infant (Asnicar et al., 2017b; Ferretti et al., 2018; Korpela et al., 2018; Sprockett et al., 2020; Yassour et al., 2018). Thus, in our simulations, species transmission rates were determined using a beta distribution, which is a natural model for values between (0,1) and often proposed for microbial abundance data (E. Z. Chen & Li, 2016; Martin et al., 2020; Sloan et al., 2006, 2007) (see **Methods**). We therefore consider four simulated scenarios: a combination of low versus high number of sources and low versus high transmission rates (see **Methods**).

Figure 2 compares the performance of SNV-FEAST and species-FEAST in estimating the true contribution of sources. FEAST using SNVs has equal if not better performance than species in most scenarios, and performs especially well when transmission rates are low and unknown source proportions are high. SNVs have a lower root mean squared error (RMSE) compared to species in three of the four scenarios and higher Pearson correlation between true and estimated contributions in all four scenarios. The difference in these correlations for SNVs versus species is significant in all four cases when using a paired Wilcoxon signed rank test (high transmission: p-value = 0.00560, 0.00251 for small and large number of sources, low transmission: p-value = 0.00024, 0.002340 for small and large number of sources). These results suggest that SNVs may offer useful signatures of transmission.

Figure 2: Ability of SNV and species-FEAST to recapitulate true contributions in simulations. Estimated known and unknown source proportions for infant microbiomes simulated with in silico mixtures of real maternal fecal microbiomes under different scenarios: either low number of contributing sources (<=5) or high number of sources (6-11), and high transmission rate of species or low transmission rate. Transmission rate is the probability of an infant being colonized by a given species, simulated using a beta distribution centered on the relative abundance of species in sources (**Methods**). 23 infants were simulated with five or fewer sources and 19 infants were simulated with a large number of sources (**Table S1**). The black line indicates the ground truth for proportions. For each simulated infant, there are 11 points plotted,

whereby 10 correspond to known sources (some of which have zero contribution), and one

corresponds to an unknown source which are indicated by a hollow circles in the plot.

To assess whether all species and all signatures SNVs in the sink are needed for accurate

source tracking, we varied the proportion of species (from 10%, 50% or 100%) and SNVs (from

10%, 50% or 100%) included as inputs to the algorithm (**Figure S1**). We used Pearson

correlation between the true and estimated proportions to represent accuracy of SNV-FEAST.

When decreasing the percentage of SNVs used, there is no statistically significant change in the

performance. However, when decreasing the percentage of species used, there are statistically

significant decreases in the performance (**Figure S1**).

To illustrate the advantage of SNV-FEAST over traditional strain tracking approaches

such as inStrain (Olm et al., 2021), we used the same synthetic communities produced in the

above simulation for inStrain profiling between each infant and each of their potential

contributing sources (**Figure S2**). InStrain computes a popANI score, which represents the

average nucleotide identity between two different metagenomic samples for a given species. As

per the inStrain paper, popANI values > 99.999% represent the same strain for that species being

shared between samples (**Methods**). However, this approach provides a binarization as to

whether or not a strain was transmitted, and does not account for the relative abundance of the

strain in the sink. Thus, we computed the fraction of each infant's species that have popANI

$\geq$99.999%, with each potential source.

As expected, both SNV-FEAST and inStrain produce estimates of sharing that correlate

positively with the ground truth mixture proportions of the contributing source samples in each

infant (**Figure S2**). We found inStrain results yielded a 0.742 Pearson correlation ($p < 1 \times 10^{-12}$)

with the true mixture proportions, whereas SNV-FEAST has a 0.866 Pearson correlation ($p < 1 \times$

10^{-12}) with the true proportions. The higher correlation values for SNV-FEAST likely reflect that

relative abundances of strains and their genomic identities are simultaneously taken into account

for source tracking, whereas inStrain only accounts for genomic identities. Finally, several of the

shared species in the simulations had popANI values < 99.999%, reflecting the complex

mixtures from multiple sources.

We next compared SNV-FEAST with the strain tracking procedure in Nayfach et al.

2016. Again, we used the same synthetic communities produced in the simulation to determine

marker alleles as defined in Nayfach et al. 2016 (**Methods**). Here a marker allele is determined

to be a SNV that is private to mother, infant, or the mother-infant dyad, and absent from the

background population, which consisted of other samples in the dataset as well as samples from

United States adults in the Human Microbiome Project (**Methods**). Species with $\geq$ 5% marker

allele sharing between mother and infant were deemed to share a strain (**Methods**). We found a

high correlation between the true mixture proportions (on x-axis) and the percentage of species

with transmission events (y-axis) (Pearson correlation 0.915 , $p < 1 \times 10^{-16}$). The higher

correlation for the Nayfach et al. 2016 approach compared to the inStrain approach possibly

reflects horizontal gene transfers between lineages residing in infants and mothers. By contrast,

there was a lower correlation between the true mixture proportions (x-axis) and the sharing for

all marker alleles across species present in the infant (y-axis) and (0.575 Pearson correlation, $p <$

1×10^{-16}) (**Figure S3B**).

Source tracking in infants over the first year of life

Having assessed the abilities of SNV-FEAST in synthetic data, we next estimated the

contribution from the true mother over time to the true infant with SNV and species-FEAST in

the Backhed et al. 2015 dataset. This dataset is composed of metagenomic samples from infants

collected at four days, four months, and 12 months after birth, as well as their mothers at the time

of delivery. Previous analyses on this data have shown that even while species similarity

increases, infants and their mothers share fewer proportions of strains over time as revealed by

sharing of SNVs private to mother-infant dyads (Nayfach et al., 2016). Thus, SNVs belonging to

strains shared only by the infant and their mother may be more informative of the true source

compared to species. Here we sought to test whether SNV and species-FEAST recapitulate these

results (**Methods**).

In applying FEAST to the Backhed et al. 2015 dataset, we estimated the proportion of

infant at birth attributable to mother. For 4 month infants, we estimated the proportion

attributable to the mother and itself at birth. For 12 month infants, we estimated the proportion

attributable to the mother and itself at birth and four months (Shenhav et al. 2019). This allowed

"unknown" to be more strictly defined as the component of the infant microbiome that could not

be explained by the mother. It also allowed us to better discern if completely new strains were

acquired at the 4th and 12th months of life (that were not already acquired during previous life

stages).

First, consistent with previous findings made with species and SNVs (Nayfach et al.,

2016), species-FEAST estimates an increasing contribution of the mother over time (t-test p-

value = 5.1 x 10^{-4}), but SNV-FEAST estimates a decrease over time (p-value = 0.063) (**Figure**

3).

Second, we assessed the ability of species and SNV-FEAST to distinguish the true

mother from three randomly selected unrelated mothers. Species-FEAST estimates an increasing

contribution of unrelated mothers over time (t-test p-value = 0.014) while SNV-FEAST

estimates no significant change over time (t-test p-value = 0.59) (**Figure 3**). The increase in

contribution from unrelated mothers with species-FEAST does not suggest that these particular

unrelated mothers are seeding the infant. Rather, the opposing trend observed with SNVs

suggests that similarity at the species level is consistent with the maturation of the infant

microbiome over time.

Finally, we estimated contributions from unknown sources, i.e. the proportion of the

infant microbiome not explainable by the true mother, the three randomly selected unrelated

mothers, or any previous time point. Species-FEAST estimates a sharp decline in contribution of

unknown sources over the first year of life (t-test p-value =7.1 x 10^{-12}) (**Figure 3**). This

significant decrease in unknown at the species level reflects the infant microbiome maturation

over the first year of life. By contrast, SNV-FEAST estimates little change in the contribution of

unknown sources (t-test p-value = 0.49) (**Figure 3**). Note that this unknown component reflects

what was gained since a previous time point. In other words, at 12 months, the infant on average

acquired the same fraction of unknown as it did at 4 months and birth. When source tracking is

run without including previous time points as sources, the unknown component increases over

the first year of life for SNVs only (**Figure S5**).

Next, we sought to understand the effect of swapping sink and source in the re-analysis of

Backhed et al. 2015 data. In **Figure 3G and H**, the infant at birth is the potential source and

mother is the sink. The estimated contribution from baby to mother is significantly smaller

(species-FEAST: 11.9 difference, Wilcoxon rank sum test p-value = 0.013; SNV-FEAST: 16.0

difference, p-value = 2.2 x 10^{-5}) compared to that of mother to baby. This trend may be

 suggestive, but is not conclusive, of directionality, whereby a less diverse source is seeded by a

 more diverse source.

Figure 3. Source tracking in the infant gut microbiome over the first year of life. Species- and SNV-FEAST were applied to Backhed et al. 2019 data to estimate the contribution of (A, B) mother, (C, D) unrelated mothers and (E, F) unknown sources to infants sampled at birth, four months, and twelve months. The black line and inset statistics pertain to the linear regression fit for the source estimates as a function of age of the infant. (G, H) are flipped source tracking analyses with mother and infant swapped when using species-FEAST and SNV-FEAST,

289 respectively. **Figure S4** shows the species that were included in species-FEAST and species that

290 had SNVs included in SNV-FEAST. **Figure S5** shows the estimate of the unknown component

291 when previous time points of the infant are excluded from the sources.

Contribution of the NICU built environment to infant microbiomes

Next, we re-analyzed a metagenomic dataset studying the contribution of the hospital

environment to the infant gut microbiome in the neonatal intensive care unit (NICU) (Brooks et

al. 2017). This dataset is composed of microbiomes of infant stool, as well as the NICU rooms of

the same infants at frequently touched surfaces, sink basins, the floor, and isolette-top sampled

over an 11-month period (Brooks et al., 2017). We applied SNV and species-FEAST to assess

the contribution of the infant's own NICU room as well as a different NICU room in the vicinity

of the infant's gut microbiome (**see Methods**).

Concordant with the findings of Brooks et al., both SNV and species-FEAST detected

that the most common source contributing to the infant microbiome was the floor and isolette-top

from the infant's own room (**Figures 4A and B**). SNV-FEAST found Infant 18 also had large

contributions from their own room's touched surfaces at multiple time points (**Figure 4B**), which

is consistent with a finding by Brooks et al. that three strains found in Infant 18 perfectly

matched ($> 99.999\%$ average nucleotide identity) strains found in the touched surfaces samples

of Infant 18's own room. Lastly, both species-FEAST and SNV-FEAST found Infant 6's

microbiome was explained almost entirely by samples from a different room with SNV-FEAST

finding a sizeable contribution from both the floor and isolette top and the sink basin in this

different room. This is concordant with Brooks et al.'s finding of multiple cases of strain sharing

across rooms of Infant 6 and 12 for the different surfaces. FEAST with both data types is able to

quantify the extent to which Infant 6's microbiome was influenced by strains present in the built

environment.

Through application of SNV and species-FEAST, we are able to quantify any trends over

time the influence of the built environment on the infant microbiome (**Figures 4A and B**). SNV-

FEAST more consistently finds that contribution from the infant's own room exceeds

contributions from a different room over time (paired Wilcoxon signed rank test for same room $>$

different room: Infant 3: p-value = 1.95×10^{-9}, Infant 6: 1, Infant 12: 3.05×10^{-5}, Infant 18: 3.81

$\times 10^{-6}$) as compared to species-FEAST (Infant 3: p-value = 0.41, Infant 6: 1, Infant 12: 5.8×10^{-4},

Infant 18: 3.81 x 10^{-6}). Interestingly, species-FEAST assigns one dominant source primarily,

whereas SNV-FEAST more often finds a combination of sources for a given sample.

Additionally, both SNV and species-FEAST estimated a large unknown component for

all four infants, with Infant 18 showing the largest mean unknown component across the NICU

stay based on SNVs (**Figure S6**). This unknown component is important because it signifies the

extent to which other sources such as the mother and diet impact infant gut colonization.

We then asked the question: is the infant more explained by the built environment rather

than vice-versa, the built environment is more explained by the infant. We tested this by

swapping the infant and each of the three built environment sources (**Figure 4C and D**). The

estimated contribution of room to infant is significantly higher than the estimated contribution of

infant to room, but this asymmetry is more pronounced with SNV-FEAST. SNV-FEAST showed

significantly higher contribution of room to infant for two of the three surface types (floor and

isolette top: Wilcoxon rank sum test p-value = 7.00x 10^{-9}, touched surface: p-value = 0.0058,

sink basin: p-value = 0.274) while species-FEAST found this to be true for one of the three

surface types (floor and isolette top: Wilcoxon rank sum test p-value = 7.1x 10^{-5}, touched

surface: p-value = 0.968, sink basin: p-value = 0.998). Interestingly, the built environments of

different rooms highly resemble each other. This is especially apparent with species-FEAST,

suggestive of similar ecological forces operating in similar built environments. By contrast,

SNV-FEAST reveals a higher diversity of contributing sources of the built environment samples

to other NICU built environments, once again highlighting the utility of performing source

tracking with SNVs.

Figure 4: Source tracking of infant gut microbiome in the NICU. (A) species-FEAST and

(B) SNV-FEAST applied to infants in the NICU. Each bar represents one sampling day in the

NICU stay of an infant. Infants 3 and 6 stayed in the same room, but at different times. The same

applies to Infants 12 and 18. The contribution of a different room was determined by using

samples from Infant 12's room for Infants 3 and 6, and samples from Infants 6's room for Infants

12 and 18 for each of the categories of surfaces per infant: touched surface, sink basin, or floor

and isolette top surface. The asterisks represent the result of a paired Wilcoxon signed rank test

indicating whether the total contribution of surfaces from the infant's own room were higher than

contributions from the other room: **** for p-value < 0.0001, *** for p-value < 0.001, ** for p-

value < 0.001, * for p-value < 0.05, and n.s. for p-value > 0.05. Iterative swapping of the infant

sink and each potential source for source tracking with (C) species-FEAST and (D) SNV-

FEAST. The first column shows source tracking results in which the infant was treated as the

sink. In each column after the first column, a different environmental source was swapped with

the infant and considered as a sink.

Global source tracking of ocean microbiomes

The ocean microbiome is a complex community that displays biogeography at the species

and functional levels (Nayfach et al., 2016; Sunagawa et al., 2015). To further understand global

patterns of ocean microbiomes, we applied SNV and species-FEAST to the Tara Oceans

microbiome dataset (Sunagawa et al., 2015). In the source tracking context, rather than defining

sharing as evidence of a transmission event (which is more likely in mother-infant data),

estimated source contributions at best explain the extent to which a given ocean sample

resembles other ocean samples. On one extreme, an ocean sample might be entirely explainable

by a single ocean's samples, and at the other extreme, an ocean sample might be explainable by

multiple oceans at the same time. Another alternative is for an ocean sample to not be

explainable by any of the provided sources, resulting in a high unknown component and

potentially suggesting high endemism. These source tracking estimates could be indicative of the

extent to which oceans mix or may be reflective of similar niches.

Tara Oceans is composed of 182 whole metagenomic sequencing samples derived from

64 stations at multiple depths. Previous research indicates that temperature is one of the highest

drivers of variability in microbial composition in the ocean (Ladau et al., 2013; Sunagawa et al.,

2015). For this reason, we restricted the source tracking analysis to sinks and sources from the

same temperature and depth range: above 20 degrees Celsius and within an average of 5 meters

below the surface.

Figure 5. Microbial source tracking in the Tara Oceans dataset with SNV and species-

FEAST. World map indicating the location of sampling sites (A). Source tracking estimates for

the contribution of different oceans to the South Pacific (n=16) (B) and Indian Oceans (n=16)

(C) are depicted with vertical bars. In each experiment, all stations around the world excluding

those from the "sink" ocean are considered potential sources. Light blue, for example, represents

the total contribution of the four stations from the Mediterranean Sea that had samples in the

surface layer that were also greater than 20°C in temperature.

Figure 6. Source tracking with ocean samples. Distance decay in contribution of a "source" ocean to a "sink" ocean when using (A) species-FEAST and (B) SNV-FEAST. In each experiment, only stations from one ocean were considered as sources for a given sink station. For example, when performing source tracking between the mediterranean and north atlantic, for each mediterranean station, the 10 available north atlantic stations were considered as potential sources. Thus, plotted are 10 points for a given mediterranean sink, where each point represents the contribution of a source station from the North Atlantic to the Mediterranean sink station in question. Shown in inset text are the slope and t-test p-value for the slope. (C) and (D) are flipped source tracking analysis with the Red Sea and Mediterranean, as well as the South Pacific Ocean and North Atlantic Ocean using species-FEAST and SNV-FEAST, respectively.

First, we performed source tracking between oceans using SNV and species-FEAST. We

treated each station around the world as a sink and estimated the contribution of different oceans

around the world to that sink (**Methods**). Unknown represents any portion of the microbiome in

these sink samples that cannot be explained by any of the provided source samples. We found

that species and SNV-FEAST estimate different amounts of sharing between oceans, where

SNVs estimate a higher unknown on average, potentially indicative of endemism. The finding

that SNV-FEAST estimates a higher unknown contribution on average is most evident in the

North Pacific, North Atlantic, South Atlantic, and Mediterranean oceans (**Figure S7**).

Additionally, in some oceans, SNVs identify contributions from oceans that species-FEAST does

not detect (**Figure 5, Figure S7**). For example, in applying FEAST to Indian Ocean samples we

find that there is measurable sharing of microbes with the Mediterranean Sea, but this is not

detected with species (**Figure 5C**). Such differences were found in samples from other oceans as

well (**Figure S7**).

Next, we assessed whether source tracking estimates display a distance-decay

relationship. Previous studies found that genetic distance, such as that represented by fixation

index F_{ST}, increases with geographic distance between populations (Cavalli-Sforza & Feldman,

2003; DeGiorgio & Rosenberg, 2013). Based on these findings, our expectation was that samples

that are further away from a given station will have reduced resemblance to that station. To

assess this distance-decay relationship, we plotted pairwiseh source tracking results across all

possible pairs of ocean samples (**Figure 6A and B**). We found that indeed as the distance

increases, the % explainability of a given source ocean decreases -0.23 % per thousand km

according to species-FEAST (t-test p-value $< 1 \times 10^{-16}$), and -0.5% per thousand km according to

SNV-FEAST (t-test p-value = 0.0018). The steeper slope for SNV-FEAST suggests that SNVs

may be more sensitive to distance decay signals on a global level.

Finally, we investigated whether some oceans have higher estimated contributions to

other oceans than vice versa, potentially indicative of the directionality of transmissions (though

see Discussion). Specifically, we investigated the relationship between the Red Sea to the

Mediterranean Sea (**Figure 6C and D**). Migration from the Red Sea to the Mediterranean,

known as Lessepsian migration, is well-documented for not only microorganisms but also

macroorganisms like fish (Bentur et al., 2008; Bianchi & Morri, 2003; Golani, 2009). Anti-

Lessepsian migration (Red Sea to Mediterranean), on the other hand, has been primarily thought

to be rare due to the Additionally, recent studies suggest that there is also evidence for anti-

Lessepsian migration of bacteria (Mediterranean to Red Sea) may be more common than

Lessepsian migration (Elsaeed et al., 2021). Research studies find that Mediterranean has brine

pools that produce similar a similar environment to the Red Sea's (Antunes et al., 2011),

allowing for bacteria from the MS to potentially thrive in the RS.

By swapping the Red Sea and Mediterranean as source and sink, we found that there was

indeed a significant difference in the estimated contribution from one direction to another with

SNVs but not species (**Figure 6C and D**). SNV-FEAST found the Mediterranean explained an

average of 15% of the Red Sea, while the Red Sea explained an average of 1.8% of the

Mediterranean (Wilcoxon rank sum test, p-value =0.02), consistent with anti-Lessepsian

migration. Meanwhile, a similar analysis with species-FEAST found the Mediterranean

explained 2.5% of the Red Sea and the Red Sea explained 4.9% of the Mediterranean (Wilcoxon

rank sum test, p-value = 0.25). In a similar analysis between North Atlantic and South Pacific we

found that both species and SNVs supported significantly greater contributions from the North

Atlantic to the South Pacific, with SNV-FEAST estimating a greater contribution (17%,

Wilcoxon rank sum test p-value = 5.1×10^{-11}) than species-FEAST (10%, Wilcoxon rank sum

test p-value =1.8×10^{-4}).

Together, these results suggest that on average, SNV and species FEAST generate similar

source tracking results in the Tara Oceans dataset, with SNVs displaying stronger signals of

endemism, distance-decay relationships, and potential directionality of transmission.

DISCUSSION

Source tracking provides insight into potential source contributions to a metagenomic

sample as well as similarities between metagenomic samples. While species abundances have

been informative in source tracking in several studies (Flores et al., 2011; Knights et al., 2011;

McGhee et al., 2020; Shenhav et al., 2019), they may be limited in their resolution. SNVs

provide a potential alternative because of their ability to distinguish sources of strain

transmissions. Here we compared the ability of a previously published source tracking algorithm

FEAST using species versus SNVs as input data. In application of species and SNV-FEAST to

simulations as well as three case studies, we demonstrate that the two input types can provide

distinct insights into microbial sharing and similarities across different environments. As a

hypothetical example, two unrelated samples may have very similar species composition due to

similar colonization processes and similar environmental influences without any actual microbial

sharing. It would be unlikely for these two unrelated samples to share rare SNVs, however. This

distinction suggests that SNVs indeed can provide insight into the ecological processes shaping

microbial communities that species information alone cannot, and our three case studies are able

to demonstrate this.

In the first case study, we confirmed previous findings that SNV sharing between

mothers and infants decreases over the first year of life while species sharing increases (Nayfach

et al., 2016), suggesting that while the infant microbiome matures to resemble adults at the

species level, sources other than the mother may seed the infant over time. In the second case

study, we confirmed source contributions from the NICU built environment to the infant

microbiome (Brooks et al., 2017), and found that SNVs detect a more consistent estimate in

source contributions overtime compared to species as well as detecting contribution from sources

not detectable by species-FEAST.

In the third case study, we perform source tracking in the Tara oceans dataset and found

SNVs display a stronger distance decay relationship. These distance-decay results parallel recent

findings made with gene content (Dlugosch et al., 2022). While previous studies have examined

the biogeography of the ocean using species profiles, genes (Dlugosch et al., 2022; Nayfach et

al., 2016) or amino acid variants from a single species (SAR11) (Delmont et al., 2019), for the

first time, we leverage the use of SNVs across all detected prevalent species in the ocean

microbiome to identify proportions of sharing across oceans. A benefit of using SNVs in the

ocean microbiome is that SNVs can track fragments of DNA that have moved due to horizontal

gene transfer in the distant past rather than relying on inference of whole genomes or presence of

private SNVs that may been transmitted in the recent past. This global-level source tracking is

analogous to admixture estimation with human genotypes (Alexander et al., 2009; Chiu et al.,

2022).

We note that source tracking provides insights into similarities between microbiomes and

potential transmissions, though the directionality is less conclusive. It is possible that increased

contributions in one direction but not the other is suggestive of directionality of transmission. For

example, in the case of the mother-infant data from Backhed et al. 2015, FEAST predicted

higher contribution from mother to baby than vice versa. This is consistent with work done on

crAss-like phage transmissions between mother and infant in the same dataset that showed

evidence of directionality by tracking the accumulation of mutations over time that are private to

the infant and absent from the mother (Siranosian et al., 2020). But in the case of the ocean, it is

possible that over longer time periods, differences in relative contributions from one part of the

world to another (e.g. Mediterranean to Red Sea) are more reflective of local selection pressures

that permit certain species and genotypes (Delmont et al., 2019). Thus, source tracking in certain

instances, such as the ocean microbiome, at best reflects the extent of similarity between samples

and is less conclusive about directionality.

A popular approach used to track strain transmissions is by detecting high average

nucleotide identity (ANI) for species shared between source and sink. For example, inStrain

(Olm et al., 2021) identifies a match between samples for a given species when ANI exceeds

99.999%. However, it is to be noted that inStrain provides distinct and complementary

information from FEAST given its binarization of whether or not a strain is shared. For

illustration purposes, if an infant harbors 100 species, of which only 1 came from their mother,

but that species' strain's relative abundance is 50% of the infant's microbiome, SNV-FEAST

would infer that the mother's contribution is 50%, while inStrain would infer that only $1/100^{th}$ of

the species are derived from the mother.

Other studies rely on tracking transmissions of strains with private SNVs shared only

between the sink and putative source (Bäckhed et al., 2015; Korpela et al., 2018; Nayfach et al.,

2016; Schmidt et al., 2019). The private marker allele tracking approach in Nayfach et al. 2016

provides an improved estimate of true percentage of species that share some portion of their

genome with putative sources compared to inStrain (**Figure S2, S3**). It is possible that requiring

only 5% of marker alleles to be shared rather than a 99.999% ANI permits detection of

horizontal gene transfers between lineages residing in mothers and infants (D. W. Chen &

Garud, 2022; Vatanen et al., 2022). However, in FEAST, by using any SNV with an informative

distribution across sources as determined by our signature scoring method, we are able to

quantify the relative contribution of all the sampled environments and assign a proportion to

these putative sources. Another advantage to FEAST is that the contribution of unknown sources

can be quantified. For example, the significant fraction of marine biodiversity estimated to be

'unknown' may be endemic, as previously noted in the Mediterranean (Katsanevakis et al.,

2014).

A drawback, however, with using SNVs over species is deeper, whole genome

sequencing is required to accurately call SNVs. Moreover, even when there is sufficient

coverage, there is still the challenge of a large number of SNVs. We demonstrate one way to

subset SNVs that uses a scoring method for informativeness, but there may yet be other methods

for filtering SNVs to the most informative set. Another potential caveat of SNV filtering is that

not all species present will be represented in the final signature SNV set (**Figure S4**). Species

with higher abundance are more likely to be represented in the signature SNV set. However, we

show that not all species need to contribute signature SNVs in order to make accurate inferences,

and likewise, not all SNVs are needed to make accurate inferences (**Figure S1**).

Ascertainment of SNVs from metagenomic data in a high-throughput manner, especially

common SNVs with microbiome genotyping technology (Shi et al., 2021), is becoming an

increasing priority for the field as metagenomic datasets become more abundant. A genotyper for

prokaryotes has already been developed and tested on a catalog of over 100 million SNVs in

order to characterize population structure (Shi et al., 2021). Such a catalog of informative SNVs

could be invaluable for source tracking. With source tracking enabling us to characterize samples

by their relationship to known samples, we have a powerful tool to explore samples in new

contexts we have yet to discover.

METHODS

Data

For simulations and analyses of infant microbiomes in the first year of life (Bäckhed et

al., 2015), we downloaded the raw shotgun metagenomic sequencing reads from public read

archives under accession number PRJEB6456. We downloaded the raw sequence reads for the

NICU analysis (Brooks et al., 2017) from accession number PRJEB323631, and the equivalent

for the Tara Oceans analyses (Sunagawa et al., 2015) were downloaded from accession number

PRJEB402. Data from the HMP Consortium (Methé et al., 2012) and Lloyd-Price et al (Lloyd-

Price et al., 2017) was downloaded from the following

URL: https://aws.amazon.com/datasets/human-microbiome-project/.

Estimation of species and SNV content of metagenomic samples

We used MIDAS (Metagenomic Intra-Species Diversity Analysis System, version 1.2,

downloaded on November 21, 2016 (Nayfach et al., 2016) to estimate species abundance and SNV

content per species in each metagenomic shotgun sequencing sample. The database we used to apply

MIDAS consisted of 31,007 bacterial genomes that are clustered into 5,952 species. The parameters we

used to estimate species abundances and SNVs were described in (Garud et al., 2019). A species was

considered present if there are at least 3 reads mapping to a set of single copy marker genes on average.

To call SNVs, we used the default MIDAS settings in order to map reads to a single representative

reference genome. The mapping was done with Bowtie 2 (Langmead & Salzberg, 2012): global

alignment, MAPID$\geq$94.0%, READQ$\geq$20, ALN_COV$\geq$0.75, and MAPQ$\geq$20, where species with reads

mapped to less than 40% of the genome were excluded from the SNV calls. We excluded samples with

depth lower than 5 reads, and excluded genetic sites using the default site filters of MIDAS (e.g.

ALLELE_FREQ$\geq$0.01, with the exception of SITE_DEPTH which was set to 3.

Application of FEAST algorithm

FEAST, originally introduced by Shenhav et al., is an R-based method that models the

mixture proportions for various "source" microbial samples for a given "sink" (Shenhav et al.,

2019). This method utilizes expectation maximization to estimate the proportions when given

any sort of count-based feature matrix representing the potential sources and sinks. The intuition

behind the estimation process is that a source with a similar species distribution to the sink would

have a higher contribution estimate to the sink. A species with non-zero counts only in source *j*

and the sink would increase the estimated contribution of source *j*. However, in many cases, the

same species are found in multiple sources simultaneous. The algorithm does not uniquely assign

a species to a source but rather simultaneously utilizes all species information to infer the source

contributions. The method was originally tested and evaluated on species and not on more fine

scale genetic data such as SNVs. The number of different species, on average, range in number

from a few hundred to a few thousand, while the number of possible nucleotide sites that vary

across different sources can number in millions. For this reason, a SNV-filtering process is

necessary so that the algorithm can run within a reasonable time and with reasonable memory

requirements.

Application of FEAST to the Backhed et al. 2015 dataset:

For both species and SNV-FEAST, the same set of sources and sinks were fed into the

FEAST algorithm. In the case study of infants in the first year of life (Bäckhed et al., 2015), the

sink consisted of the infant fecal sample at either four days, four months, or 12 months and the

potential sources consisted of fecal samples from the true mother, three randomly selected

mothers from the same dataset, and also any previous time points for the infant.

Species-FEAST utilized all species present in the infant whereas SNV-FEAST used

signature SNVs from the subset of species that had signature SNVs. Shown in **Figure S3** are the

distribution of species included in species and SNV-FEAST.

Application of FEAST to the Brooks et al. 2017 dataset:

For the case study of infants in the NICU (Brooks et al., 2017), the sink consisted of the

fecal sample of the infant at a given time point and the potential sources consisted of pooled

reads from the touched surfaces, the sink basin and the floor and isolette top from both the

infant's own room as well as a different room. The different room was Infant 12's room for

Infants 3 and 6, Infants 6's room for Infants 12 and 18.

Application of FEAST to the Sunagawa et al. 2015 dataset: Determining the signature SNV set

Signature SNVs were identified as described in the main text. We provide specific steps for

determining signature SNVs:

(1) Filter sites: only sites of the genome with at least the required number of reads mapping

to the site are considered. In the case study of infants in the first year of life (Bäckhed et

al., 2015) and infants in the NICU (Brooks et al., 2017), the minimum coverage

requirement is 10 across the sink and *J* sources. For the Tara Ocean (Sunagawa et al.,

2015) samples, the minimum coverage is five reads (Sunagawa et al., 2015).

Additionally, sites that are biallelic must have more than one read mapped to each allele

to be considered.

(2) Perform per site per source parameter estimates: for each potential source compute the

estimated allele frequency in the sink θ under two different hypotheses:

Hypothesis 1: Source *i* with allele frequency p_i explains the allele counts in the sink.

$$\hat{\theta} = p_i$$

614   **Hypothesis 2:** A combination of all other sources except i (sources $j \neq i$) explain the

615   observed allele count distribution in the sink. The estimate of the sink allele frequency is

616   computed using a mixture of the allele frequencies p_j from those sources. The mixing

617   parameter α_j is learned using Sequential Least Squares Programming (scipy.minimize())

618   with the constraint of summing to 1 with bounds of 0 to 1 inclusive: $\sum_{j \neq i} \alpha_j = 1$.

$$\hat{\theta} = \sum_{j \neq i} \alpha_j p_i$$

620   **(3)** Compute per site per source log likelihoods: Compute the binomial log-likelihood under

621   hypotheses 1 and 2, given n reads with the reference allele and m reads with the

622   alternative allele in the sink:

$$l(\hat{\theta}) = n \log \hat{\theta} + m \log(1 - \hat{\theta})$$

624   (4) Compute per site per source log likelihood ratio:

$$l_1(\theta) - l_2(\theta)$$

(5) Compute per site summary signature score: The maximum log likelihood ratio per site is

the signature score for that SNV, representing how favorably one of the sources explains

the sink over all other sources

(6) Filtering of SNVs using signature score: One signature score for that SNV represents

how favorably one source explains the sink better than all other sources. All the scores

are ranked across SNVs and SNVs with scores that are greater than two standard

deviations over the mean signature score within each 200 kbp window of the genome are

retained as signature SNVs. This window size was chosen for to optimize run time and

memory requirements.

Note, if only one source passes minimum coverage filtering, $l_2(\theta) = 0$ resulting in a

very high likelihood ratio as represented by $l_1(\theta)$ for the one source. These SNVs are

more likely to pass the signature score filtering. One exception for SNVs that are

included in the signature SNV set without passing signature score filtering are SNVs with

an allele that is completely unique to the infant, as these represent SNVs that are

641    potentially derived from an unknown source. Signature SNVs are obtained from the SNV

642    profile of every species for which there is MIDAS output.

644  *Simulating mother to infant transmission*

645    The mixture proportions for 28 simulated infants is shown in **Table S1**. Four possible

646  scenarios are simulated using a combination of either low or high number of sources and low or

647  high transmission probabilities of species. High transmission of species was simulated by

648  drawing separate transmission probabilities for each species in each contributing source based on

649  a beta distribution with a mean equal to the species relative abundance and variance equal to 0.1,

650  a value selected to emulate Backhed et al.'s mean relative abundance and variance. For the low

651  transmission scenario, transmission probabilities were drawn from a beta distribution with mean

652  0.1 times the relative abundance and variance at 0.1. To determine if a species from each source

653  was transmitted to a given infant, a binomial draw was performed J times, where J = number of

654  sources, and the probability of a mother transmitting the species is p_j based on the beta-drawn

655  transmission probability. If any of the draws yields a one, that species is transmitted to the infant

656  from all sources. The same simulated data under these scenarios is used for both SNV and

657  species source tracking.

658    The source tracking estimates are compared to the true mixing proportions using

659  Spearman correlation. The significance of correlation is calculated using the stat_cor function in

660  the 'ggpubr' package (*CRAN - Package Ggpubr*, n.d.).

662  *Comparison to inStrain*

663    We ran inStrain (Olm et al., 2021) on the same synthetic samples as described above.

664  InStrain "profile" (Olm et al., 2021) and inStrain "compare" (Olm et al., 2021) were run for

665  every possible infant-source pair. For example, for simulated infant 1 there were 10 putative

666  sources, therefore inStrain compare was run 10 times for each putative source. InStrain reports

667  popANI calculated per scaffold for a given species. To compute a single statistic per species, we

668  computed the average popANI across scaffolds for a given species. The percent infant

669  microbiome species that had strains shared with mother was computed as the number of species

670  in which popANI was >= 99.999% divided by the total number of species with coverage >= 5.

671  PopANI was only calculated in scaffolds that had >=5 coverage in both samples of the pair.

Comparison with strain tracking approach in Nayfach et al. 2016

We applied the strain tracking approach in Nayfach et al. 2016 on the same synthetic communities described above. In Nayfach et al. 2016, strain transmissions are tracked by identifying 'marker alleles' which are private to the infant, mother, or infant-mother dyad, and absent from the broader population. A strain is considered to be shared if at least 5% of all marker alleles for a mother-infant dyad are shared. Note that the approach for strain tracking proposed in Nayfach et al. 2016 utilizes SNV information outputted by MIDAS, but is not a part of MIDAS.

Each simulated infant had up to 10 sources that were real maternal samples from Backhed et al. 2015 For each possible pair of infants and maternal sources (10 pairings per infant, with 48 infants), we found the set of infant-only marker alleles, mother-only marker alleles, and mother-infant dyad marker aleles. As described in Nayfach et al, 2016, only sites with minimum 30 reads and only alleles that were supported by at least 10% of the total reads aligned to that site were considered. The infant marker allele and mother marker allele were defined as alleles that were present only in the focal sample and absent from the background samples (or below 3 reads = 10% * 30 reads). For the infant, the background consisted of all mothers (including mothers that were used to simulate the infant), real infant samples (excluding infants of mothers used to simulate the infant), and 337 samples of adults from the United States in the HMP (which includes 180 unique adults) that were obtained from the metagenomics repository of HMP under project ID SRP002163 and SRP056641 (Lloyd-Price et al., 2017; Methé et al., 2012). For the mother, the background consisted of all mother and infant samples in addition to the HMP samples. For computing shared marker alleles, an allele must be present in both the mother and infant but absent from the background, which consisted of all mothers and the HMP samples.

To compute sharing, two quantities were considered: "total sharing", defined as % shared marker alleles/ (infant marker alleles + mother marker alleles + shared marker alleles) and proportion of infant marker alleles that are shared: % shared marker alleles/ (infant marker alleles + shared marker alleles). The first quantity compared to FEAST estimates was the percentage of infant species in which the "total sharing" was at least 5%. The second quantity

compared to FEAST was the pooled proportion of infant marker alleles that are shared across all
species.

Distance Decay Analysis

To study the relationship between source tracking estimates and geographic distance, we
analyzed all oceans as either a sink or source against all other possible oceans. To compute
geographic distance between stations, we applied the Haversine distance to the longitude and
latitude of the sampling sites provided by (Sunagawa et al., 2015) using the package "geosphere"
(Hijmans et al., 2021). Source tracking estimates were computed as described above using either
SNV-FEAST or Species FEAST. The regression line for the distance decay analysis was
computed using a linear mixed model "contribution ~ distance + (1| sink_ocean)".

Ethics declarations
The authors declare that they have no competing interests.

Acknowledgements
We thank Nicole Zeltser for processing the Tara Oceans data through the MIDAS pipeline. We
thank Richard Wolff for his advice on simulations. We thank Michael Wasney for testing the
associated software. We thank members of the Garud lab for their feedback on the manuscript.
LB was supported by the NSF Graduate Research Fellowship Program grant numbers DGE-
1650604 and DGE-2034835. N.R.G. received support from the Paul Allen Frontiers Group and
the Research Corporation for Science Advancement.

Availability of Data and Materials
All metagenomic data was obtained from public repositories. The applicable accessions numbers
are PRJEB6456 for Backhed et al. 2015 (mother-infant), PRJEB323631 for Brooks et al. 2017
(NICU), PRJEB402 for Sunagawa et al. 2015 (Tara Oceans), and SRP002163 and SRP056641
for HMP.

Source code for SNV-FEAST signature SNV selection as well as analyses in this paper are
available at GitHub (https://github.com/garudlab/Signature-SNVs), Zenodo (DOI

10.5281/zenodo.7515044), and PyPi for pip installation (https://pypi.org/project/Signature-

SNVs/0.0.1/).

References

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in

unrelated individuals. *Genome Research*, *19*(9), 1655–1664.

https://doi.org/10.1101/GR.094052.109

Antunes, A., Ngugi, D. K., & Stingl, U. (2011). Microbiology of the Red Sea (and other) deep-

sea anoxic brine lakes. *Environmental Microbiology Reports*, *3*(4), 416–433.

https://doi.org/10.1111/J.1758-2229.2011.00264.X

Asnicar, F., Manara, S., Zolfo, M., Truong, D. T., Scholz, M., Armanini, F., Ferretti, P., Gorfer,

V., Pedrotti, A., Tett, A., & Segata, N. (2017a). Studying Vertical Microbiome

Transmission from Mothers to Infants by Strain-Level Metagenomic Profiling. *MSystems*,

2(1). https://doi.org/10.1128/msystems.00164-16

Asnicar, F., Manara, S., Zolfo, M., Truong, D. T., Scholz, M., Armanini, F., Ferretti, P., Gorfer,

V., Pedrotti, A., Tett, A., & Segata, N. (2017b). Studying Vertical Microbiome

Transmission from Mothers to Infants by Strain-Level Metagenomic Profiling. *MSystems*,

2(1). https://doi.org/10.1128/MSYSTEMS.00164-16/ASSET/54C4C531-C6DB-421B-

8C8A-10C0ECFE3BE9/ASSETS/GRAPHIC/SYS0011720800004.JPEG

Bäckhed, F., Roswall, J., Peng, Y., Feng, Q., Jia, H., Kovatcheva-Datchary, P., Li, Y., Xia, Y.,

Xie, H., Zhong, H., Khan, M. T., Zhang, J., Li, J., Xiao, L., Al-Aama, J., Zhang, D., Lee, Y.

S., Kotowska, D., Colding, C., … Jun, W. (2015). Dynamics and stabilization of the human

gut microbiome during the first year of life. *Cell Host and Microbe*, *17*(5), 690–703.

https://doi.org/10.1016/j.chom.2015.04.004

Bentur, Y., Ashkar, J., Lurie, Y., Levy, Y., Azzam, Z. S., Litmanovich, M., Golik, M., Gurevych,

B., Golani, D., & Eisenman, A. (2008). Lessepsian migration and tetrodotoxin poisoning

due to Lagocephalus sceleratus in the eastern Mediterranean. *Toxicon*, *52*(8), 964–968.

https://doi.org/10.1016/J.TOXICON.2008.10.001

Bianchi, C. N., & Morri, C. (2003). Global sea warming and "tropicalization" of the

Mediterranean Sea: biogeographic and ecological aspects. *Biogeographia – The Journal of*

Integrative Biogeography, *24*(1). https://doi.org/10.21426/B6110129

Brooks, B., Olm, M. R., Firek, B. A., Baker, R., Thomas, B. C., Morowitz, M. J., & Banfield, J.

F. (2017). Strain-resolved analysis of hospital rooms and infants reveals overlap between

the human and room microbiome. *Nature Communications*, *8*(1), 1–7.

https://doi.org/10.1038/s41467-017-02018-w

Cavalli-Sforza, L. L., & Feldman, M. W. (2003). The application of molecular genetic

approaches to the study of human evolution. *Nature Genetics 2003 33:3*, *33*(3), 266–275.

https://doi.org/10.1038/ng1113

Chen, D. W., & Garud, N. R. (2022). Rapid evolution and strain turnover in the infant gut

microbiome. *Genome Research*, *32*(6), 1124–1136.

https://doi.org/10.1101/GR.276306.121/-/DC1

Chen, E. Z., & Li, H. (2016). A two-part mixed-effects model for analyzing longitudinal

microbiome compositional data. *Bioinformatics*, *32*(17), 2611–2617.

https://doi.org/10.1093/BIOINFORMATICS/BTW308

Chiu, A. M., Molloy, E. K., Tan, Z., Talwalkar, A., & Sankararaman, S. (2022). Inferring

population structure in biobank-scale genomic data. *The American Journal of Human*

Genetics. https://doi.org/10.1016/J.AJHG.2022.02.015

CRAN - Package ggpubr. (n.d.). Retrieved March 6, 2022, from https://cran.r-

project.org/web/packages/ggpubr/index.html

DeGiorgio, M., & Rosenberg, N. A. (2013). Geographic Sampling Scheme as a Determinant of

the Major Axis of Genetic Variation in Principal Components Analysis. *Molecular Biology*

and Evolution, *30*(2), 480–488. https://doi.org/10.1093/MOLBEV/MSS233

Delmont, T. O., Kiefl, E., Kilinc, O., Esen, O. C., Uysal, I., Rappé, M. S., Giovannoni, S., &

Eren, A. M. (2019). Single-amino acid variants reveal evolutionary processes that shape the

biogeography of a global SAR11 subclade. *ELife*, *8*. https://doi.org/10.7554/ELIFE.46497

Dlugosch, L., Poehlein, A., Wemheuer, B., Pfeiffer, B., Badewien, T. H., Daniel, R., & Simon,

M. (2022). Significance of gene variants for the functional biogeography of the near-surface

Atlantic Ocean microbiome. *Nature Communications 2022 13:1*, *13*(1), 1–11.

https://doi.org/10.1038/s41467-022-28128-8

Elsaeed, E., Fahmy, N., Hanora, A., & Enany, S. (2021). Bacterial Taxa Migrating from the

Mediterranean Sea into the Red Sea Revealed a Higher Prevalence of Anti-Lessepsian

Migrations. *OMICS A Journal of Integrative Biology*, *25*(1), 60–71.

https://doi.org/10.1089/omi.2020.0140

Ferretti, P., Pasolli, E., Tett, A., Asnicar, F., Gorfer, V., Fedi, S., Armanini, F., Truong, D. T.,

Manara, S., Zolfo, M., Beghini, F., Bertorelli, R., De Sanctis, V., Bariletti, I., Canto, R.,

Clementi, R., Cologna, M., Crifò, T., Cusumano, G., … Segata, N. (2018). Mother-to-Infant

Microbial Transmission from Different Body Sites Shapes the Developing Infant Gut

Microbiome. *Cell Host and Microbe*, *24*(1), 133-145.e5.

https://doi.org/10.1016/j.chom.2018.06.005

Flores, G. E., Bates, S. T., Knights, D., Lauber, C. L., Stombaugh, J., Knight, R., & Fierer, N.

(2011). Microbial Biogeography of Public Restroom Surfaces. *PLoS ONE*, *6*(11), e28132.

https://doi.org/10.1371/journal.pone.0028132

Garud, N. R., Good, B. H., Hallatschek, O., & Pollard, K. S. (2019). Evolutionary dynamics of

bacteria in the gut microbiome within and across hosts. *PLoS Biology*, *17*(1), e3000102.

https://doi.org/10.1371/JOURNAL.PBIO.3000102

Golani, D. (2009). Distribution of Lessepsian migrant fish in the Mediterranean.

Http://Dx.Doi.Org/10.1080/11250009809386801, *65*(S1), 95–99.

https://doi.org/10.1080/11250009809386801

Hijmans, R. J., Karney, C., Geographiclib,] (, Williams, E., Vennes, C., & Maintainer,]. (2021).

Package "geosphere." https://doi.org/10.1007/s00190012

Katsanevakis, S., Coll, M., Piroddi, C., Steenbeek, J., Lasram, F. B. R., Zenetos, A., & Cardoso,

A. C. (2014). Invading the Mediterranean Sea: Biodiversity patterns shaped by human

activities. *Frontiers in Marine Science*, *1*(SEP), 32.

https://doi.org/10.3389/FMARS.2014.00032/ABSTRACT

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G.,

Bushman, F. D., Knight, R., & Kelley, S. T. (2011). Bayesian community-wide culture-

independent microbial source tracking. *Nature Methods*, *8*(9), 761–765.

https://doi.org/10.1038/nmeth.1650

Korpela, K., Costea, P., Coelho, L. P., Kandels-Lewis, S., Willemsen, G., Boomsma, D. I.,

Segata, N., & Bork, P. (2018). Selective maternal seeding and environment shape the

human gut microbiome. *Genome Research*, *28*(4), 561–568.

https://doi.org/10.1101/GR.233940.117/-/DC1

Ladau, J., Sharpton, T. J., Finucane, M. M., Jospin, G., Kembel, S. W., O'Dwyer, J., Koeppel, A.

F., Green, J. L., & Pollard, K. S. (2013). Global marine bacterial diversity peaks at high

latitudes in winter. *The ISME Journal 2013 7:9*, *7*(9), 1669–1677.

https://doi.org/10.1038/ismej.2013.37

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods 2012 9:4, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Li, S. S., Zhu, A., Benes, V., Costea, P. I., Hercog, R., Hildebrand, F., Huerta-Cepas, J.,

Nieuwdorp, M., Salojärvi, J., Voigt, A. Y., Zeller, G., Sunagawa, S., De Vos, W. M., &

Bork, P. (2016). Durable coexistence of donor and recipient strains after fecal microbiota

transplantation. *Science*, *352*(6285), 586–589.

https://doi.org/10.1126/SCIENCE.AAD8852/SUPPL_FILE/LI-SM.PDF

Lloyd-Price, J., Mahurkar, A., Rahnavard, G., Crabtree, J., Orvis, J., Hall, A. B., Brady, A.,

Creasy, H. H., McCracken, C., Giglio, M. G., McDonald, D., Franzosa, E. A., Knight, R.,

White, O., & Huttenhower, C. (2017). Strains, functions and dynamics in the expanded

Human Microbiome Project. *Nature 2017 550:7674*, *550*(7674), 61–66.

https://doi.org/10.1038/nature23889

Martin, B. D., Witten, D., & Willis, A. D. (2020). MODELING MICROBIAL ABUNDANCES

AND DYSBIOSIS WITH BETA-BINOMIAL REGRESSION. *The Annals of Applied*

Statistics, *14*(1), 94. https://doi.org/10.1214/19-AOAS1283

McGhee, J. J., Rawson, N., Bailey, B. A., Fernandez-Guerra, A., Sisk-Hackworth, L., & Kelley,

S. T. (2020). Meta-SourceTracker: application of Bayesian source tracking to shotgun

metagenomics. *PeerJ*, *8*, e8783. https://doi.org/10.7717/peerj.8783

Methé, B. A., Nelson, K. E., Pop, M., Creasy, H. H., Giglio, M. G., Huttenhower, C., Gevers, D.,

Petrosino, J. F., Abubucker, S., Badger, J. H., Chinwalla, A. T., Earl, A. M., Fitzgerald, M.

G., Fulton, R. S., Hallsworth-Pepin, K., Lobos, E. A., Madupu, R., Magrini, V., Martin, J.

C., … White, O. (2012). A framework for human microbiome research. *Nature 2012*

486:7402, *486*(7402), 215–221. https://doi.org/10.1038/nature11209

Nayfach, S., Rodriguez-Mueller, B., Garud, N., & Pollard, K. S. (2016). An integrated

metagenomics pipeline for strain profiling reveals novel patterns of bacterial transmission

and biogeography. *Genome Research*, *26*(11), 1612–1625.

https://doi.org/10.1101/gr.201863.115

Olm, M. R., Crits-Christoph, A., Bouma-Gregson, K., Firek, B. A., Morowitz, M. J., & Banfield,
J. F. (2021). inStrain profiles population microdiversity from metagenomic data and
sensitively detects shared microbial strains. *Nature Biotechnology 2021 39:6*, *39*(6), 727–
736. https://doi.org/10.1038/s41587-020-00797-0

Schloissnig, S., Arumugam, M., Sunagawa, S., Mitreva, M., Tap, J., Zhu, A., Waller, A., Mende,
D. R., Kultima, J. R., Martin, J., Kota, K., Sunyaev, S. R., & Weinstock, G. M. (2013).
Genomic variation landscape of the human gut microbiome. *Nature*.
https://doi.org/10.1038/nature11711

Schmidt, T. S. B., Hayward, M. R., Coelho, L. P., Li, S. S., Costea, P. I., Voigt, A. Y., Wirbel, J.,
Maistrenko, O. M., Alves, R. J. C., Bergsten, E., de Beaufort, C., Sobhani, I., Heintz-
Buschart, A., Sunagawa, S., Zeller, G., Wilmes, P., & Bork, P. (2019). Extensive
transmission of microbes along the gastrointestinal tract. *ELife*, *8*.
https://doi.org/10.7554/ELIFE.42693

Shenhav, L., Thompson, M., Joseph, T. A., Briscoe, L., Furman, O., Bogumil, D., Mizrahi, I.,
Pe'er, I., & Halperin, E. (2019). FEAST: fast expectation-maximization for microbial
source tracking. *Nature Methods*, *16*(7). https://doi.org/10.1038/s41592-019-0431-x

Shi, Z. J., Dimitrov, B., Zhao, C., Nayfach, S., & Pollard, K. S. (2021). Fast and accurate
metagenotyping of the human gut microbiome with GT-Pro. *Nature Biotechnology 2021*
40:4, *40*(4), 507–516. https://doi.org/10.1038/s41587-021-01102-3

Siranosian, B. A., Tamburini, F. B., Sherlock, G., & Bhatt, A. S. (2020). Acquisition,
transmission and strain diversity of human gut-colonizing crAss-like phages. *Nature*
Communications 2020 11:1, *11*(1), 1–11. https://doi.org/10.1038/s41467-019-14103-3

Sloan, W. T., Lunn, M., Woodcock, S., Head, I. M., Nee, S., & Curtis, T. P. (2006). Quantifying
the roles of immigration and chance in shaping prokaryote community structure.
Environmental Microbiology, *8*(4), 732–740. https://doi.org/10.1111/J.1462-
2920.2005.00956.X

Sloan, W. T., Woodcock, S., Lunn, M., Head, I. M., & Curtis, T. P. (2007). Modeling taxa-
abundance distributions in microbial communities using environmental sequence data.
Microbial Ecology, *53*(3), 443–455. https://doi.org/10.1007/S00248-006-9141-
X/FIGURES/4

Sprockett, D. D., Martin, M., Costello, E. K., Burns, A. R., Holmes, S. P., Gurven, M. D., &

Relman, D. A. (2020). Microbiota assembly, structure, and dynamics among Tsimane

horticulturalists of the Bolivian Amazon. *Nature Communications 2020 11:1*, *11*(1), 1–14.

https://doi.org/10.1038/s41467-020-17541-6

Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G., Djahanschiri,

B., Zeller, G., Mende, D. R., Alberti, A., Cornejo-Castillo, F. M., Costea, P. I., Cruaud, C.,

d'Ovidio, F., Engelen, S., Ferrera, I., Gasol, J. M., Guidi, L., Hildebrand, F., … Bork, P.

(2015). Structure and function of the global ocean microbiome. *Science*, *348*(6237).

https://doi.org/10.1126/SCIENCE.1261359

Vatanen, T., Jabbar, K. S., Vlamakis, H., Knip, M., & Correspondence, R. J. X. (2022). Mobile

genetic elements from the maternal microbiome shape infant gut microbial assembly and

metabolism. *Cell*, *185*, 4921-4936.e15. https://doi.org/10.1016/j.cell.2022.11.023

Yassour, M., Jason, E., Hogstrom, L. J., Arthur, T. D., Tripathi, S., Siljander, H., Selvenius, J.,

Oikarinen, S., Hyöty, H., Virtanen, S. M., Ilonen, J., Ferretti, P., Pasolli, E., Tett, A.,

Asnicar, F., Segata, N., Vlamakis, H., Lander, E. S., Huttenhower, C., … Xavier, R. J.

(2018). Strain-Level Analysis of Mother-to-Child Bacterial Transmission during the First

Few Months of Life. *Cell Host & Microbe*, *24*(1), 146-154.e4.

https://doi.org/10.1016/J.CHOM.2018.06.007