Elevated temperatures reduce population-specific transcriptional plasticity in developing lake
sturgeon (*Acipenser fulvescens*)

William S. Bugg*[1], Matt J. Thorstensen[1], Katie E. Marshall[2], W. Gary Anderson[1], and Ken M.
Jeffries[1]

*Corresponding Author: wsbuggfiles@gmail.com, Department of Biological Sciences, 50 Sifton

Road, University of Manitoba, Winnipeg, Manitoba, Canada, R3T 2N2

1. Department of Biological Sciences, University of Manitoba, 50 Sifton Road, Winnipeg,
Manitoba, Canada, R3T 2N2
2. Department of Zoology, University of British Columbia, Vancouver, BC, Canada V6T
1Z4

Abstract:

Rising mean and variance in temperatures elevate threats to endangered freshwater

species such as lake sturgeon, *Acipenser fulvescens*. Previous research demonstrated that higher

temperatures during development result in physiological consequences for lake sturgeon

populations throughout Manitoba, Canada, with alteration of metabolic rate, thermal tolerance,

transcriptional responses, growth, and mortality. We acclimated lake sturgeon (30 – 60 days post

fertilization, a period of high mortality) from northern and southern populations (56° 02′ 46.5″ N,

96° 54′ 18.6″ W and 50° 17′ 52″ N, 95° 32′ 51″ W respectively, separated by approximately 650

km) within Manitoba to current (summer highs of 20-23°C) and future projected (+2-3°C)

environmental temperatures of 16, 20, and 24°C for 30 days, and measured gill transcriptional

responses using RNAseq. Transcripts revealed SNPs consistent with genetically distinct

populations and transcriptional responses altered by acclimation temperature. There were a

higher number of differentially expressed transcripts observed in the southern, compared to the

northern, population as temperatures increased, indicating enhanced transcriptional plasticity.

Both lake sturgeon populations responded to elevated acclimation temperatures by

downregulating the transcription of genes involved in protein synthesis and energy production.

Further, there were population-specific thresholds for the downregulation of processes promoting

transcriptional plasticity as well as mitochondrial function as the northern population showed

decreases at 20°C, while this capacity was not diminished until 24°C in the southern population.

These transcriptional responses highlight the molecular impacts of increasing temperatures for

divergent lake sturgeon populations during vulnerable developmental periods and the critical

influence of transcriptome plasticity on acclimation capacity.

1. Introduction

Freshwater fishes have experienced the highest rates of extinction among all vertebrates throughout the 20[th] century, influenced by the compounding and dynamic climate change-related stressors of elevated temperatures, hypoxic conditions, and pathogens which continue to imperil freshwater fishes and limit suitable habitat (Marcos-López et al., 2010; Burkhead, 2012; Comte et al., 2012; Schade et al., 2014; Krabbenhoft et al., 2020; Vollset et al., 2020; Jane et al., 2021). Climate change is increasing both the mean and variance of environmental temperature (Thornton et al., 2014; Dillon et al., 2016). Environmental changes are often occurring faster than organisms can adapt to increasing temperatures, especially in species with long generation times (Crozier and Hutchings, 2014; Morgan et al., 2020; Logan and Cox, 2020). Therefore, phenotypic plasticity will be critical for maintaining fitness in response to climate change and promoting survival in these changing environments (Burggren, 2018; Scheiner et al., 2019; Earhart et al., 2022).

Genomic and epigenetic variation can contribute to transgenerational plasticity, while intragenerational plasticity induced to alter an organism's phenotype at the transcriptomic level may act as an adaptive trait (Muschick et al., 2011; Jonsson and Jonsson, 2019). Together these transgenerational and intragenerational mechanisms provide adaptive plasticity which may enable molecular physiological changes, promoting survival (Lafuente and Beldade, 2019; Oomen and Hutchings, 2017; Wellband and Heath, 2017). Further, polyploid organisms with larger genomes may have increased acclimatory capacity at both the transgenerational and intragenerational timescales through transcriptome plasticity (Ellis et al., 2014; Trifonov et al., 2016; Van de Peer, 2017). In populations of polyploid species with divergent genetic histories, which may be under genetic constraints preventing performance traits from evolving rapidly (Logan and Cox, 2020), genome level changes may be anticipated but alteration of phenotypes through transcriptomic plasticity may be necessary to promote survival as environmental changes intensify.

During early development, change in temperature is a critical abiotic factor necessitating prompt phenotypic alteration. These plastic, individual-level phenotypic changes can alter organismal physiology, buffer against acute environmental changes, lead to long-term phenotypic impacts, and be selected upon across generations (McKenzie et al., 2021). Thus,

plasticity and physiological flexibility can be an adaptive response stimulated in stressful

environments and is a reversible phenotypic change that may be persistent for days to months,

associated with the process of acclimation (Hockachka and Somero, 2002; Crozier and

Hutchings 2014; Mackey et al., 2021). Underlying these phenotypic modifications are

transcriptional mechanisms that impact the establishment of long-term phenotypes (e.g.,

developmental plasticity; Jonsson and Jonsson, 2019). Exposing developing fish to increasing

environmental temperatures will thus alter traits that may have transient as well as long-lasting

physiological consequences. The basis for plasticity is variation in phenotypes produced by the

same genotype under different environmental conditions (Lafuente and Beldade, 2019; Earhart et

al., 2022). Acclimatory plasticity is especially important in early life to ensure sufficient

phenotypic flexibility, enabling fishes to survive in variable environmental conditions until

adulthood when their reproductive potential can be reached (Burggren, 2018). In early

developmental stages, different phenotypes and developmental trajectories can emerge from

genetically distinct populations exposed to similar environmental change based, in part, on

differences in transcriptome plasticity (Logan and Cox, 2020). Induced transcriptional responses

to stressors often differ between populations and may be indicative of long-term fitness and

survival consequences if phenotypes cannot be adequately altered in early life to allow

individuals to persist in changing environments (Whitehead et al., 2012; Bugg et al., 2020;

Mundy et al., 2020; Bugg et al., 2021).

A primary mechanism underlying physiological responses in the face of environmental

change are modifications to the abundance of messenger RNA (mRNA) through transcriptional

processes (Smith et al., 2013; Connon et al., 2018; Jeffries et al., 2019; Jeffries et al., 2021).

Through mRNA sequencing (i.e., RNAseq) and *de novo* transcriptome assembly, transcriptomic

approaches can be used when few genomic resources are available for a species (Alvarez et al.,

2015; Connon et al., 2018; Oomen & Hutchings 2017; Thorstensen et al., 2020; Komoroske et

al., 2020). Examining genomic sequence differences (SNPs) and differential expression genes

through transcriptome-wide approaches can illuminate an organism's physiologically- and

ecologically-relevant responses to environmental change, aid in the identification of tolerance

thresholds, and inform conservation management practices (Connon et al., 2018). As

environmental extremes in freshwater systems escalate, transcriptomic techniques provide

powerful tools to investigate adaptive capacities in species of conservation concern, highlighting

the genetic mechanisms that promote physiological acclimation to environmental change. While

acclimation may promote thermal resilience, the potential for decreased transcriptional plasticity

as well as limitations to other physiological responses in the face of multiple stressors may

suggest increased vulnerability, the possibility of severe short-term outcomes, and the potential

for impacts on long-term fitness (Smith et al., 2013; Jeffries et al., 2018).

Greater than 50% of the world's river basins face widespread anthropogenic impacts

which limit productivity, species richness, and biodiversity, threatening approximately a quarter

of freshwater fish species globally (Palmer et al., 2008; IUCN, 2018; Grill et al., 2019; Su et al.,

2021; Warkentin et al., 2022). Habitat loss, commercial exploitation, and changing climates,

have pushed many species to the brink of extirpation (Arthington et al., 2016). Sturgeons are

some of the most endangered groups of vertebrates on the planet (WWF, 2022), have long

generation times (often > 15 years) and have persisted relatively unchanged for hundreds of

millions of years. However, because of a rapidly changing environment, sturgeon populations are

in widespread decline. Lake sturgeon, *Acipenser fulvescens* are widely distributed throughout

North America, and within Canada have experienced population declines >90% in the northern

extent of their range since the 1960's (COSEWIC, 2006). Historically, lake sturgeon population

declines in the 1800s were caused by commercial exploitation for caviar. Since the 1950's

however, habitat loss and degradation associated with dams and other barriers have become the

most prominent threats to populations and the species' persistence (Cleator et al., 2010;

COSEWIC 2017; Van der Lee and Koops, 2021).

Further, with the intensification of their environments due to changing climates, lake

sturgeon now likely face additional stressors, especially so for northern populations. Summer

temperatures for Canadian populations can reach upwards of 23°C, exceeding temperatures of

20°C for approximately 50 days per year, imposing thermal stress and whole-organism

physiological impacts for developing lake sturgeon (Bugg et al., 2020). Temperatures in the

northern range for lake sturgeon are also increasing faster than the global average (projected 2.1-

3.4°C increase by 2050; Manitoba Hydro, 2015), which may further limit suitable habitat, and

imperil these endangered sub-Arctic populations as water temperatures exceed their thermal

limits (Vincent et al., 2015; Zhang et al., 2019; Bugg et al., 2021).

Increasing environmental temperatures directly influence the development, metabolism,

transcriptomic responses, and survival of northern populations of lake sturgeon under laboratory

conditions and may contribute to observed declines in wild stocks with high mortality (> 90%)

observed in early development (Caroffino et al., 2011; Bugg et al., 2020; Bugg et al., 2021). In

fishes like lake sturgeon, the gill lamellae is a multi-purpose tissue comprised of multiple cell

types reactive to the external environment with roles in gas exchange, osmoregulation, pathogen

defense, acid-base balance, and responses to thermal stress (Lazado and Caipang, 2014;

Komoroske et al., 2015; Akbarzadeh et al., 2018; Gilmour and Perry, 2018). Indirectly, elevating

temperatures may impact many of these processes in the gill, resulting in increased vulnerability

with additional environmental stressors like increase hypoxia or pathogen exposure (LaPatra et

al., 1994; Georgiadis et al., 2001; Fujimoto, 2018; Clouthier et al., 2020). Sturgeons may be

especially susceptible to the combined effects of increasing temperature, hypoxic environments,

and pathogens in early development, when mortality is often high (> 90%; Caroffino et al., 2010)

and sturgeon have limited immune responses and ability to escape stressful environments (Webb,

1986; Peake et al., 1997; Schindler, 2001; Deslauriers and Kieffer, 2012; Gradil et al., 2014 a,b;

Verhille et al., 2014). However, despite these vulnerabilities in early development, sturgeon have

successfully persisted for hundreds of millions of years in ever-changing environments, evolving

elaborate polyploid genetic complexity, and demonstrating high levels of phenotypic plasticity in

early development, highlighting the potential for inducible transcriptional plasticity (Braasch and

Postlethwait, 2012; Trifonov et al., 2016; Bugg et al., 2020; Bugg et al., 2021; Penman, 2021;

Yoon et al., 2021; Brandt et al., 2021, Brandt et al., 2022). Therefore, early development

represents a crucial window to examine lake sturgeon responses to thermal acclimation,

especially within the gill where key homeostatic functions are critical to physiological

performance (Burggren, 2018; Barley et al., 2021).

Here we used RNAseq to investigate differential transcriptomic responses to increased

environmental temperatures between two latitudinally distinct populations of developing lake

sturgeon (McDougall et al., 2017; Bugg et al., 2020). Genomic differences were used to provide

context for potential genetic differentiation between the two populations using SNPs in the RNA

sequence reads. As sturgeons are ancestral polyploid species, with highly variable genomic

architecture and phenotypic plasticity (Braasch and Postlethwait, 2012; Trifonov et al., 2016;

Bugg et al., 2021), we expected population-specific responses to temperature treatments in lake

sturgeon. We examined the gill mRNA transcript abundance in $n = 36$ individuals from three

thermal acclimation treatments in the northern and southern population of lake sturgeon within

Manitoba. We targeted early development, 60 days post fertilization, for this analysis to highlight

key vulnerabilities to thermal stress for these populations during a critical period in life when

high mortality (> 90%; Caroffino et al., 2011) often occurs in the wild. We then investigated

phenotypic plasticity through exploration of 9 total contrasts spanning the molecular mechanisms

shared between populations as acclimation temperatures increased, as well as population and

acclimation-specific responses to increased environmental temperatures. Previous research

demonstrated that acclimating lake sturgeon for 30 days to thermally stressful, but ecologically-

relevant summer temperatures, of $20°C$ and $24°C$, resulted in population and acclimation-specific

changes in mortality, metabolism, thermal tolerance, and transcriptional responses (Bugg et al.

2020). Therefore, we hypothesized that acclimation to these same elevated temperatures during

early life history would elicit transcriptome-wide responses associated with energy mobilization,

immune response, cellular repair, and heat shock proteins indicative of chronic thermal stress.

Given the higher mortality and lower thermal tolerance observed in the northern population post-

temperature acclimation in a previous study (Bugg et al., 2020), we predicted that the northern

population of lake sturgeon would show greater transcriptional signatures of thermal stress as

temperatures increased, relative to the southern population.

2. Materials and Methods:

2.1 Lake Sturgeon Husbandry and Acclimation

Lake sturgeon used in the current study were from the same populations and acclimation

treatments described in Bugg et al. (2020). Briefly, gametes were collected from wild male and

female lake sturgeon downstream from the Pointe du Boise Generating Station on the southern

Winnipeg River ($50°17'52''$N, $95°32'51''$W) and first rapids on the northern Burntwood River

($56°02'46.5''$ N, $96°54'18.6''$W; separated by approximately 650 km) (Figure 1). Eggs and

sperm from the Winnipeg River were fertilized at the University of Manitoba animal holding

facility while those from the Burntwood River were fertilized at the Grand Rapids Fish Hatchery,

and subsequently transferred to the University of Manitoba. Progeny from the Winnipeg River

population were the product of the fertilization of the eggs from two females with the sperm

from two males, while individuals from the Burntwood River were the product of the

fertilization of eggs from one female with the sperm from six males. Immediately post-

fertilization, embryos from each population were submerged in a clay substrate to de-adhease

them and then were gently stirred by hand for 1 h. Next, embryos were rinsed with dechlorinated

freshwater and kept in tumbling jars to incubate at $12°C$ until hatch. Water temperature was

maintained at $12°C$ until 13 days post fertilization after which it was increased by $1°C$ day^{-1} until

$16°C$, matching typical hatchery rearing conditions in Manitoba. Starting at 19 days post

fertilization, and continuing throughout the length of the experiment, developing lake sturgeon

were fed freshly hatched artemia (Artemia International LLC; Texas, USA) three times daily.

Acclimation began at 30 days post fertilization, by increasing treatments to $20°C$ and $24°C$ at a

rate of $1°C$ day^{-1} and keeping one treatment from each population at $16°C$. Lake sturgeon from

both populations were acclimated to these temperatures for 30 days with water temperatures

recorded by HOBO Water Temperature PROv2 Data Loggers (Onset Computer Corporation;

Bourne, MA, USA) and checked by thermometer at least three times daily. All animals in this

study were reared and sampled under guidelines established by the Canadian Council for Animal

Care and approved by the Animal Care Committee at the University of Manitoba under Protocol

#F15-007.

2.2 Sampling and RNA extraction

At the end of the 30 day acclimation, six larval lake sturgeon per treatment were netted

and euthanized by immersion in an overdose of tricaine methanesulfonate solution (250 mg L^{-1};

MS-222, Syndel Laboratory, Vancouver, Canada) with an equal volume of sodium bicarbonate

to buffer against pH change. Next, gill tissue was extracted from each larval fish and preserved

in RNA*later* (Invitrogen; Carlsbad, CA, USA), held at 4°C for 24 h and stored at −80°C prior to

RNA extraction. Total RNA was extracted from the gill tissue by homogenization in 500 µl of

lysis buffer (PureLink RNA Mini Kit; Invitrogen; Ambion Life Technologies; Waltham, MA,

USA) for 10 min at 50 Hz using a TissueLyser II (Qiagen; Germantown, MD, USA) and then

using a PureLink RNA Mini Kit (Invitrogen; Ambion Life Technologies) following

manufacturer instructions. Total RNA purity and concentration was evaluated for all samples

using a NanoDrop One (Thermo Fisher Scientific) as well as gel electrophoresis to assess RNA

integrity. Samples were then stored at -80$°C$ prior to sequencing.

2.3 *De novo* Transcriptome Assembly

Prior to the present study, a lake sturgeon gill reference transcriptome was created by

sequencing mRNA from the pooled RNA of 3 individuals. We used a mRNA isolation, stranded

library preparation approach to produce 50 million reads of 100 base pair paired-end reads. The

transcriptome was *de novo* assembled using Trinity (Grabherr et al., 2011) through the Canadian

Centre for Computational Genomics (C3G; https://computationalgenomics.ca/). Assembled

transcripts were then annotated with Trinotate (http://trinotate.sourceforge.net/) using a modified

approach. First assembled transcripts were blasted against a fish-specific database with

annotation hits returned. Transcripts that were not annotated from the fish specific database were

then blasted against the entire SwissProt library.

2.4 Sequencing

Total RNA for each sample was normalized to approximately 15 ng μL^{-1} at a total

volume of 10 μL and shipped to the McGill Applied Genomics Innovation Core sequencing

facility (https://www.mcgillgenomecentre.ca/) for cDNA library preparation and sequencing.

Samples were run on a Bioanalyzer to confirm RNA quality and all samples had a RNA Integrity

Number (RIN) between 8.7-10. A mRNA isolation step was performed using NEBNext Poly(A)

Magnetic Isolation Modules (New England Biolabs) followed by using NEBNext Ultra II

Directional RNA Library Prep Kit for Illumina (New England Biolabs) to produce stranded

cDNA libraries. NEBNext dual adaptors (New England Biolabs) were then individually applied

to barcode each library prior to sequencing of 100 base pair paired-end reads. All 36 sturgeon

gill samples were sequenced on a single NovaSeq 6000 (Illumina) lane producing 1.53 billion

total reads with an average of 42,528,434 (± 17,812,372 SD) reads per sample.

2.5 Quality control and Trimming

Raw read files were uploaded to Galaxy servers (Giardine et al., 2005) and paired reads

were joined and checked for quality using FastQC version 0.11.8 (Andrews, 2010). To filter low-

quality reads, Trimmomatic version 0.38.0 was applied with the Illuminaclip option set to

standard for TruSeq3 adapter sequences, a maximum mismatch count of 2, and a palindrome

read alignment threshold of 30. Additionally, a sliding window trimming step was used with a

size of 4 base pairs as well as a minimum read length filter set at 20 bases. Ultimately this

trimming removed approximately 2.8% (± 0.26% SD) of reads from each file. Full details for

quality control and trimming procedures are available in Supplementary File 1, Sheets A-C.

2.6 Read Mapping and Differential Expression

After filtering, HISAT2 version 2.1 and StringTie version 2.1.1 (Kim et al., 2015; Pertea

et al., 2015) were used to align paired read libraries to the *de novo* assembled gill transcriptome

and produce a set of assembled transcripts for each sample, respectively. HISAT2 was run with

default parameters and produced an overall alignment rate of approximately 83% (± 0.7% SD)

per sample. HISAT2 aligned bam files were then input into StringTie to assemble transcripts,

using default settings, producing an average of 94,992 (± 11,075 SD) transcripts per sample.

StringTie files for each sample were then merged producing a master list of 139,456 transcripts.

To assess differential transcript expression each of these transcripts were first counted in

HISAT2 produced aligned read files using featureCounts version 1.6.2 with default settings

(Liao et al., 2014; Available in Supplementary File 2). Total counts were then assessed for

differential expression using the R v4.0.0 package EdgeR v3.32.1 (Robinson et al., 2010; R Core

Team, 2022). Prior to analysis the "filterByExpr" function with default arguments was used to

retain transcripts expressed among any individual, resulting in a final count of 95,593 transcripts.

Full details for read mapping of each individual sample are available in Supplementary File 1,

Sheet D.

2.7 Principal Component Analysis, Differentially Expressed Transcripts, and Aggregate

Heatmapping

A principal component analysis (PCA) was conducted on featureCounts from the above

95,693 transcripts after normalizing the data to the library size using the plotMDS function in the

limma package (Ritchie et al., 2015). Next, numbers of differentially expressed genes between

the Burntwood River and Winnipeg River populations of lake sturgeon acclimated to 16°C, 20°C,

and 24°C for 30 days in early development were tabulated for upregulated and downregulated

genes independently, representing genes with a $\log_2$ fold change > 1 and a *q*-value (FDR adjusted

p value) < 0.05 for each of the 9 individual contrast (Details for all differentially expressed

transcripts as outlined in Figure 3B can be found in Supplementary File 3, Sheets A-C). The 9

total contrasts compared the populations at each acclimation temperature as well as across

acclimation temperatures for each population. Log$_2$ counts per million (CPM) data for all 95,693

transcripts were then generated and the top 500 genes from each contrast with the highest

absolute value log$_2$ CPM fold change were included in an aggregate list of the most highly

differentially expressed genes across all contrasts, resulting in a total of 4,500 transcripts, 2,285

uniquely included (full CPM data are available for all 95,693 transcripts in Supplementary File

4). A heatmap with hierarchical clustering was then produced using the log$_2$ CPM expression of

these aggregate most highly differentially expressed 2,800 unique transcripts using the pheatmap

package (Kolde, 2015).

2.8 Functional Term Enrichment

Transcripts annotated to the *de novo* lake sturgeon transcriptome were split into groups of

upregulated and downregulated genes by treatment based on their log$_2$ CPM fold change (< -1 or

> 1) and were counted for distinct gene names (hereafter referred to as split-contrast). The

positively and negatively regulated annotated genes were then analyzed for functional

enrichment using gene ontology (GO) terms through enrichR (Chen et al., 2013; Kuleshov et al.,

2016; Xie et al., 2021) with positive hits returned from the "GO_Biological_Processes_2018",

"GO_Molecular_Function_2018", and "GO_Cellular_Component_2018 databases and filtered

for an adjusted *p*-value < 0.05. The resulting 213 enrichment terms across the upregulated and

downregulated spit-contrast were then tabulated into a master file detailing their presence or

absence in each split-contrast (0 or 1; Full presence and absence data across each split-contrast is

available in Supplementary File 5). This master file was then used to create UpSet plots of the

shared and population-specific terms both upregulated and downregulated across all split-

contrasts, upregulated terms as temperatures increased from 16°C to 20°C, and terms which

decreased as temperatures increased from 20°C to 24°C using the 'UpsetR' package v1.4.0

(Conway et al., 2017). As the split-contrasts with the most GO terms were largely

downregulated, split-contrasts with downregulated functional enrichment terms as temperatures

increased were then focused on for interpretation of results.

Next, non-redundant enrichment terms in the GO Biological Process category were

identified with REVIGO for summary and visualization (Supek et al., 2011). Adjusted *p* values

were supplied for all REVIGO analyses to guide the clustering of redundant GO terms by the

REVIGO algorithm (Supek et al., 2011). Where GO terms were shared across populations or

acclimation treatments, the geometric mean of the adjusted *p* values from all shared groups was

input into REVIGO. All REVIGO output is available in Supplementary File 6, Sheets A-I.

2.9 Aggregate Gene Network Analysis and Enrichment

An aggregate of the most highly differentially expressed transcripts was then investigated,

producing a total of 503 uniquely annotated most differentially expressed transcripts, from the

aggregate list of 4,500 most differentially expressed transcripts across contrasts used for the

heatmap (the full aggregate list of 503 annotated most differentially expressed transcripts is

available in Supplementary File 7). These 503 unique gene names were uploaded to OmicsNet

(Zhou and Xia, 2018) and input under the genes category using the "official gene symbol"

classification and zebrafish *Danio rerio* as the study organism, as this was the most closely

related model species option (used for OmicsNet aggregate network analysis only). Next,

protein-protein interactions were selected in the network building panel, using STRING the

database (a comprehensive collection of physical and functional protein-protein interactions

covering 5090 organism across a broad and diverse selection of benchmarked data sources;

Szklarczyk et al., 2019), to assembly a network of the unique genes and their potential

interactions. The network was then changed to a 2-D model, using a force atlas layout, with seed

nodes highlighted with a blue ring around the node, and edge bundling applied. Functional

analysis of the produced network was then categorized using the Function Explorer, querying all

nodes against the GO: Biological Process database. Processes with a *p* value < 0.05 are reported

in Supplementary File 8. Additionally, degree of connectivity and node betweenness scores were

tabulated for each transcript in the network and are included in Supplementary File 8 as well.

2.10   SNP Calling and Analyses

Population differentiation was assessed with a combination of RNA-seq and reduced

representation-oriented programs to mitigate issues with population genetic analyses of polyploid

mRNA data (Thorstensen et al., 2021; Thorstensen et al., 2022a). The program fastp v0.20.1 was

used for adapter trimming and read filtering, with a minimum phred quality of 15, a maximum of

40% of unqualified bases allowed in a read before read filtering, a minimum length of 100 base

pairs, and polyG read tails were trimmed (Chen et al., 2018). STAR v2.7.9a was used as a splice-

aware aligner in two-pass mode to map reads to the lake sturgeon gill transcriptome used for

differential expression analyses in this study (Dobin et al., 2013; Thorstensen et al., 2022a).

Picard v2.26.3 was used to clean the resulting bam files, add read groups, and mark duplicate

reads. GATK v4.2.4.0 was used to split reads that contained Ns in cigar reads (Van der Auwera

et al., 2013). Next, STACKS v2.60 was used to create loci for SNP calling (gstacks) and call

SNPs (populations) with a minimum minor allele count of 2, at least 60% of the data present

within each of the Winnipeg and Burntwood River groups, and 90% present data overall

(Catchen et al., 2013). Because paralogous loci may lead to biased population genetics measures,

HDplot was used to filter out loci with heterozygosity ≤ 0.7 and read ratio deviations ≤ 7

(McKinney et al., 2017). SNPRelate v1.30.1 was used to filter out SNPs with $R^2 > 0.20$ along 500

kilobase segments, treating each transcript as a chromosome (Zheng et al., 2012).

Hierfstat v0.5-11 was used to assess Weir & Cockerham's F_{ST} between the samples from the

Burntwood and Winnipeg Rivers, with 95% confidence intervals generated by 1000 bootstrap

iterations (Goudet, 2005). Admixture v1.3.0 was used to estimate ancestry coefficients for each

group between K values of 2, and pophelper v2.3.1 was used to visualize the results from

Admixture (Alexander et al., 2009; Francis et al., 2017). While further population genetic and

association-based analyses were possible (e.g., tests for signatures of selection), extensive family

structure remained in the data because genotyped samples were *not* random representations of

their populations of origin. Therefore, the present population genetic analyses were used to

assess genetic dissimilarities between groups in the context of other research and to demonstrate

how transcriptional differences between groups may be consistent with underlying genetic

differences.

3. Results

3.1 Principal Component analysis, Differentially Expressed Transcripts and Aggregate

Heatmapping

Examination of the PCA (Figure 3A), number of upregulated and downregulated genes

(Figure 3B), as well as the heatmap displaying the most highly differentially expressed genes

across contrasts (Figure 3C), demonstrated that the Burntwood River and Winnipeg River

populations of lake sturgeon responded differently to elevated temperatures both in the

magnitude and trajectory of transcriptional change. The PCA showed divergent response patterns

with Winnipeg River sturgeon in treatments with populations grouping out separately along a

linear trajectory as temperatures increased. Acclimation treatments primarily separated out

linearly across principal component 1 (16.9% contribution), while populations were dispersed

vertically across principal component 2 (12.4% contribution). As temperatures increased from

16°C to 20°C both higher temperature treatments separated out cleanly from their 16°C

counterparts. However, as temperatures increased from 20°C to 24°C, these two temperature

treatments remained with some overlap for the northern Burntwood River population, while the

24°C treatment for the southern Winnipeg River population continued on its responsive

trajectory, separating out from the 20°C treatment.

Patterns of upregulation and downregulation between the two populations at a given

acclimation temperature, and within populations across acclimation temperatures, demonstrated

population-specific and acclimation-specific responses to elevated temperatures in developing

lake sturgeon. When comparing Burntwood River and Winnipeg River sturgeon at each

acclimation temperature there are similar numbers of up and downregulated transcripts.

However, within population comparisons reveal a much higher level of transcriptional plasticity

in the southern Winnipeg River population. The largest changes in transcript regulation in the

Burntwood River and Winnipeg River sturgeon occurred as acclimation temperatures increased

from 16°C to 24°C, where the populations differentially expressed a total of 3,442 and 8,792

transcripts, respectively. These changes in transcript expression between 16°C and 24°C represent

the differential expression of approximately 3.6% and 9.2% of all transcripts for the Burntwood

River and Winnipeg River populations of sturgeon included in the analysis, respectively.

Between 16°C and 20°C these numbers of differentially expressed transcripts were less than half

when compared to the 16 to 24°C transition, with just 1,398 and 4,022 transcripts differentially

expressed in the Burntwood River and Winnipeg River, respectively. While producing similar

qualitative patterns of expression to the Burntwood River, the number of differentially expressed

transcripts was higher in the southern Winnipeg River across acclimation treatments, with lower

values in the northern latitude Burntwood River population.

The heatmap outlined specific patterns of transcript expression of the most highly

differentially expressed transcripts, with divergent $\log_2$ CPM expression values as temperatures

increased above 16°C for both populations but with differences in these patterns across

populations (Figure 3C). Differential expression patterns of these transcripts are largely

qualitatively similar across the Burntwood River 20°C and 24°C treatments, with those from the

Winnipeg River 20°C and 24°C treatments. However, there is evidence of a much larger change

in transcriptional regulation observed between the Winnipeg River 16°C and higher temperature

treatments (20°C and 24°C), especially when compared to patterns of expression observed in the

Burntwood River sturgeon as temperatures increase. Overall, the patterns in the most highly

differentially expressed genes align with the PCA in the increased separation between Winnipeg

River 16 and 20°C acclimation treatments, and the lack of separation between Burntwood River

20 and 24°C acclimation treatments compared to Winnipeg River sturgeon in the same

temperatures. Overall, these patterns suggest that the most highly differentially expressed

transcripts reflect similar transcriptional trajectories as shown in the PCA of all transcripts.

3.2 Functional Term Enrichment

Term enrichment demonstrated 213 unique GO pathway terms distributed across the 17

split-contrasts (Supplemental Figure 2A). Overall, the split-contrasts with the largest number of

assigned terms were those underlying processes being upregulated as temperatures increased

from 16 to 20°C (WR16v20_neg and BWR16v20_neg; 75 and 85 terms respectively) and those

processes that decreased in expression as temperatures continued to increase from 20°C to 24°C

(WR20v24_pos and BWR20v24_pos; 62 and 97 terms, respectively). The split-contrasts

between the Burntwood River and Winnipeg River populations as temperatures increase from

16°C to 20°C and again from 20°C to 24°C specifically revealed conserved and population-

specific biological processes upregulated as temperatures increase from 16°C to 20°C and those

that are decreased with further temperature increase to 24°C. As temperatures increased from

16°C to 20°C there were 50 upregulated functional term responses shared between populations,

as well as 39 responses specific to the northern Burntwood River population and 25 distinct to

the southern Winnipeg River population (Supplemental Figure 2B). As acclimation temperatures

increased further from 20°C to 24°C there were 26 functional term responses shared between

populations, with 71 terms specific to the Burntwood River population and 38 distinct to the

Winnipeg River population (Supplemental Figure 2B).

3.3 GO Term and REVIGO Analysis

Overwhelmingly, the majority of all 213 terms were categorized as GO Biological

Processes, encompassing 77% of terms identified and highlighting transcriptional mechanisms

altered across lake sturgeon populations as temperatures increase, therefore these terms were

focused on for further analysis.

Conserved responses to increasing temperature

Across both temperature increases (16 to 20°C and 16 to 24°C), there were 8 GO

Biological Process terms identified as downregulated and conserved across both populations.

These mechanisms were involved in RNA maturation, mRNA splicing, mitochondrial protein

insertion, and ribosome biogenesis (Figure 4; Supplementary File 6, Sheet A).

As the acclimation temperature increased from 16°C to 20°C there were 29 conserved GO

Biological Process terms identified as downregulated and shared between the northern

Burntwood River and southern Winnipeg River populations of lake sturgeon. These shared

processes were involved in chromosomal structure (telomere maintenance, histone methylation),

mRNA processing and maturation (maturation of rRNA, mRNA splicing, mRNA processing,

miRNA-mediated gene silencing), ribosomal biogenesis and protein modification (ribosome

biogenesis, protein methylation), as well as mitochondrial function (protein targeting to

mitochondrion, protein insertion into mitochondrial inner membrane; Supplementary File 6,

Sheet B).

As acclimation temperature further increased from 16°C to 24°C there were 22 conserved

GO Biological Process terms that decreased in expression in both the Burntwood River and

Winnipeg River lake sturgeon populations. These shared processes were involved nuclear

regulation and DNA repair (double-strand break repair, mitotic nuclear membrane reassembly,

mitotic nuclear membrane organization, DNA duplex unwinding), further mitochondrial

dysregulation (mitochondrial translation, elongation, organization, protein localization, in

addition to terms above), mRNA and RNA splicing (via spliceosome, and transesterification

reactions respectively), ribosomal biogenesis, and protein insertion into the ER membrane

(Supplementary File 6, Sheet C), demonstrating further disruption at the DNA, mRNA,

mitochondrial, and protein synthesis levels as temperatures increased.

Population-specific responses to increasing temperature

As temperatures increased from 16°C to 20°C, there were also population-specific

increases in transcripts involved in various biological processes, fairly evenly split across both

populations. Specific to the Winnipeg River population, there were 38 GO Biological Process

terms identified as downregulated (Supplementary Figure 3A) largely characterized by changes

in mRNA modification and transcription (gene expression, mRNA processing, non-sense

mediated decay, tRNA export, mRNA stability regulation, RNA polymerase I transcription,

mRNA cleavage, RNA mediated gene silencing), protein production and modification

(ribosomal disassembly, translation, regulation of translation, translation termination, peptide

biosynthetic process, amino acid modification), signaling (negative regulation of Wnt signaling

pathway, tyrosine phosphate signaling pathway, G1 and p53 DNA damage signaling), and

mitochondrial function (mitochondrial gene expression, protein localization, as well as

translational elongation and termination; Supplementary File 6, Sheet D). In the Burntwood

River population, there were an additional 20 population-specific GO Biological Process terms

downregulated (Supplementary Figure 3B). These Burntwood River specific upregulated terms

involved further impacts on mRNA processes and alternative splicing mechanisms (histone

arginine methylation, peptidyl-lysine monomethylation, regulation of alternative splicing via

spliceosome, mRNA 3'end processing, RNA nuclear export, miRNA metabolism, and different

signaling mechanism from the Winnipeg River population (positive regulation of Wnt signaling,

canonical Wnt signaling pathway, smoothened signaling pathway, regulation of p38MAPK

cascade; Supplementary File 6, Sheet E). Additionally, the Burntwood River population

exhibited population-specific mitochondrial processes terms including lipid metabolism and

mitochondrial respiratory chain complex assembly, key for energy production.

As temperatures increased further from 16°C to 24°C there were more population-specific

GO Biological Process terms identified as downregulated, however these changes were almost

solely in the Winnipeg River population, with 56 population-specific process to 3 in the

Burntwood River population (Supplementary Figure 3C). These terms involved increased signs

of cellular disfunction with downregulation of processes involving translational disruption,

telomere maintenance (via telomere lengthening and telomerases) DNA damage (cellular

response to DNA damage, nucleotide excision repair, DNA repair), mitochondrial and oxidative

stress (respiratory chain complex III assembly), ubiquitin and protein cycling (ubiquitin protein

transferase and ligase activity), and immune function (regulation of type I interferon signaling;

Supplementary File 6, Sheet F). In contrast, there were only 3 population specific processes for

this temperature contrast in the Burntwood River involving the downregulation of endosomal

transport via the multivesicular body sorting pathway, neuromuscular processes, and ureter

development.

3.4 Aggregate Transcript Network Analysis and Functional Enrichment

The aggregate most highly differentially expressed transcript network demonstrated

connections across many different functional groups of transcripts that were altered in

developing lake sturgeon during acclimation to elevated temperatures. Transcripts identified as

hubs (orange circles, highlighted in blue; Figure 5) form many responsive network connections

and are active in DNA regulation and repair processes (*cdk2, ddit4, trip12, exo1, mcm9, chd1l,*

smarcad1a), mRNA splicing (*zmat5, cwc27, prpf31*), protein turnover and synthesis (*rps27a,*

cops5, hap1, rnf8, usp13, fbxo45, smurf2, rplp0, bop1, eif3d), chaperone activity and protein

folding (*hsc70, hspa8, hyou1, dnajc2, cct4, unc45b, naca*), as well as immune and inflammatory

activation (*traf6, b2m, banf1, rsad2, egr1, jak1, prkcbb*). Additional hub genes were involved in

more general processes like calcium regulation (*fyna, cam2kd1*), cell cycling (*plk4, spc24,*

cdc14ab, pds5a), autophagy (*atg4b, trim13*), development (*kif7*), osmotic balance (*cftr*) and

mitochondrial function (*gfm1*). Many of these processes were also present in the network GO

Biological Process term enrichment with enriched pathways including protein ubiquitination,

catabolism, folding and stability, RNA splicing (including spliceosomal assembly), immune

function (inflammation, virus, immune system process, adaptive immune response, cytokine-

mediated signaling pathway, lymphocyte differentiation), responses to DNA damage and repair,

heart development, and responses to stress (including heat; Supplementary File 8). The hub

transcript with the highest degree of connectivity and network betweenness score

(Supplementary File 8) was *rps27a* with dual roles in ribosome and ubiquitin structure.

3.5 SNP Calling and Analyses

An initial 32,244 SNPs were called with STACKS from the mapped reads, of which

24,396 were accepted as non-paralogous, and 14,517 were accepted after pruning for linkage

disequilibrium. This dataset of 14,517 SNPs was used for subsequent analyses.

A principal component analysis for SNPs between lake sturgeon samples in the present

data demonstrate that the northern Burntwood River and southern Winnipeg River populations

are genetically distinct (Figure 2) with an F_{ST} of 0.086 (0.082 – 0.090 95% CI). The admixture

plot (Supplemental Figure 1) demonstrates distinct populations with ancestry from the southern

Winnipeg River population in the downstream northern Burntwood River population attributable

to introgression or uncertainty in ancestry coefficient estimates.

4. Discussion

In the current study, we demonstrated conserved and population-specific transcriptional

responses as environmental temperatures increased in two geographically and genetically distinct

endangered populations of lake sturgeon. These responses spanned wide ranging cellular

protective processes induced during acclimation to thermally stressful conditions. Specifically,

these processes related to the cellular machinery involved in transcriptional and translational

regulation, mitochondrial function, oxidative damage, and immunocompetence. Overarching

transcript expression patterns indicated population-specific capacity for gene transcription as

environmental temperatures increased, with divergent transcript response trajectories, different

temperature thresholds to induce transcriptional change, and a higher magnitude of

transcriptional plasticity apparent in the southern Winnipeg River population. Conserved and

population-specific biological process responses illustrated that this transcriptional plasticity

promoted whole-organism acclimation to elevated temperatures in developing lake sturgeon with

potential for adaptive responses at the population scale. Ultimately, these processes contribute to

the ability of lake sturgeon to acclimate to environments with increasing temperature and

highlight the physiological consequences once acclimatory capacity has been exceeded.

Plasticity as an Adaptive Trait

Thermally responsive molecular pathways first rely on the ability of lake sturgeon to

transcriptionally respond to environmental change. Without the necessary translational

machinery in place, mRNA abundance changes cannot effectively modify the proteome or the

physiology of the organism (Vornanen et al., 2005; Smith et al., 2013; Akbarzadeh et al., 2018).

Thus, as thermal thresholds are reached and lake sturgeon face a reduction in the ability to

transcribe mRNA's, produce ribosomes, and maintain mitochondria, those consequences likely

negatively impact the organism's ability to respond to changes in their environment with

impaired and diminished mRNA, protein, and energy production.

Increasing temperatures induced alteration in the abundance of transcripts associated with

post-transcriptional modifications in both populations of lake sturgeon, influencing cellular

plasticity. Conserved responses of alternative splicing and gene silencing via miRNA suggest

that these mechanisms are part of the inducible stress response of sturgeons and likely play a role

in regulating phenotypic plasticity, as observed in other fishes (Somero, 2018; Healy & Schulte,

2019; Tan et al., 2019; Shiina and Shimizu, 2020; Verta and Jacobs, 2021; Thorstensen et al.,

2022b; Steward et al., 2022). However, these processes were ultimately downregulated as

temperatures increased to 24°C, indicating thermal thresholds on post-translational as well as

transcriptional responsiveness. The conservation of these mechanisms across both populations of

lake sturgeon, as temperatures increase and under stressful conditions in various other species of

fish, highlight the importance of post-transcriptional modifications in response to environmental

stressors across the evolution of fish lineages.

As temperatures surpass thermal thresholds, and the effectiveness of some transcriptional

and translational mechanisms are diminished, overall physiological plasticity is likely reduced.

Fishes' susceptibility to the effects of additional stressors are thereby likely increased by

compromising their ability to respond at multiple levels of biological organization (Smith et al.,

2013; Brennan et al., 2022). As temperatures increased, transcription modifying processes were

more highly preserved in Winnipeg River sturgeon that historically experience higher

temperature thermal environments in the wild. As the fish used from each wild population are

genetically distinct (F_{ST} of 0.086), experience different thermal environments, and have been

geographically separated throughout recent history (McDougall et al., 2017; Bugg et al., 2020),

the transgenerational preservation of these transcriptional mechanisms may itself be an adaptive

mechanism, enhancing the southern Winnipeg River population's plasticity and ability to

respond to environmental change (Fox et al., 2019). Enhanced transcriptional plasticity likely

results in the observed increases in growth, metabolism, thermal tolerance, and survival of the

southern Winnipeg River population under thermally stressful conditions, when compared to

their Burntwood River counterparts (Bugg et al., 2020).

Transcriptomic Responses to Increasing Temperatures

As temperatures increased first from 16°C to 20°C and then to 24°C, analysis of

downregulated biological processes revealed 8 shared transcriptional responses conserved

between the lake sturgeon populations. Together, these processes illustrated the effects of

increased temperatures with impacts on key cellular mechanisms involved in transcription,

translation, and energy production, processes fundamental to cellular function. Without the

necessary cellular machinery in place, the capacity for protein level changes to plastically alter

the phenotype of developing lake sturgeon and promote environmental acclimation at the whole

organism level is diminished (Buccitelli and Selbach, 2020). Further, acclimation to 20°C was

associated with decreases in processes involved in telomere organization in both populations,

which have the potential to disrupt normal cell cycling and proliferation (Barrington et al., 2017).

In both populations, temperature increase to 24°C highlighted decreases in DNA repair

mechanisms as thermal stress mounts, which can result in a wide array of cellular consequences

such as genome instability, modulation of disease-states, impacts to cellular health and ultimately

senescence (Chatterjee and Walker, 2017; Barnes et al., 2019). These impacts of temperature

increase likely elevate the lake sturgeon's susceptibility to the effects of compounding

environmental stressors, which may be expected as environmental conditions intensify (Smith et

al., 2013; Todgham and Stillman, 2013; Brennan et al., 2022).

Population-specific responses to increased temperature highlight transcriptional

mechanisms specific to the northern or southern lake sturgeon. As temperatures increased to

20°C, the northern Burntwood River population decreased processes promoting transcriptional

plasticity including alternative splicing and methylation (similarly as observed in Atlantic

Sturgeon, *A. oxyrinchus;* Mastrangelo et al., 2012; Metzger and Schulte, 2016; Steward et al.,

2022; Thorstensen et al., 2022b; Penny et al., 2023), as well as mitochondrial processes involved

in energy production such as lipid metabolism and respiratory chain complex assembly (Greene

and Selivonchick 1987, Chung and Schulte, 2015). In contrast, the southern Winnipeg River

population demonstrated decreases in cellular processes dominated by mRNA modification and

transcriptional mechanisms. For the Winnipeg River population, downregulation of genes

involved in alternative splicing and methylation, and energy producing mechanisms did not

occur until 24°C was reached, highlighting potential mechanisms that underly observed

population-specific thresholds in transcriptional plasticity as well as mitochondrial function and

energy production (Bugg et al., 2020). Overall, as temperatures reached 24°C population-specific

processes primarily occurred only in the southern Winnipeg River population, further

demonstrating the preservation of transcriptional plasticity in this population. Specifically, we

observed downregulation of mechanisms involved in response to DNA damage, oxidative stress,

protein turnover, and immune function. As the southern Winnipeg River population is exposed to

higher yearly temperatures and demonstrates population-specific transcriptional processes when

exposed to increased temperatures, as compared to their more northerly Burntwood River

counterparts (Bugg et al., 2020; Bugg et al., 2021), these unique responses are likely indicative

of adaptive mechanisms downregulated to preserve energy and promote survival as temperatures

increase (Fox et al., 2019).

Network Analysis

Predictive transcript network analysis, of the aggregate most highly differentially

expressed transcripts, revealed functional processes possibly significant for thermal acclimation.

Key hub transcripts and enriched pathways in these regulatory networks involved impacts on

DNA integrity, mRNA splicing, protein turnover and synthesis, chaperone activity and protein

folding as well as immune responsive mechanisms, with within-network GO analysis indicating

the recruitment of similar biological processes. Together, these highlighted transcripts and

pathways indicate the importance of these systems in the transcriptomic response to thermal

acclimation across different populations of developing lake sturgeon.

The effects of thermal stress had direct impacts on genes involved in the integrity of

DNA, altering damage signaling, repair processes, and cellular cycling (pathways: DNA repair,

response to DNA damage stimulus, DNA damage checkpoint; hub transcripts: *cdk2*, *ddit4,*

trip12, exo1, mcm9, chd1l, smarcad1a). These transcripts are largely involved in damage

detection during the G1/S phase transition, and associated DNA replication (Bacevic, et al.,

2017) suggesting key cellular processes disrupted by thermal stress in lake sturgeon. This G1

checkpoint specifically detects DNA damage prior to entering the S phase of DNA synthesis,

preventing proliferation of irreversible DNA damage in future cells (Bertoli et al., 2013).

Differential expression of transcripts involved in mRNA splicing (pathways:

spliceosomal complex assembly, RNA splicing; hub transcripts: *zmat5, cwc27, prpf31*) highlight

modified post-transcriptional RNA processes occurring as temperatures increased. Post-

transcriptional modifications by pre-mRNA splicing can provide increased transcriptional

plasticity, with the potential for enhanced acclimatory capacity when faced with environmental

change (Mastrangelo et al., 2012; Steward et al., 2022; Thorstensen et al., 2022b). However,

ultimately as temperatures reached $24°C$, these processes were downregulated across

populations, demonstrating the impacts that increasing temperatures have on transcriptional

plasticity, reducing the capacity for physiological plasticity through alternative splicing

mechanisms as stressors mount.

As temperatures increased, there were modifications to protein synthesis and turnover

processes (pathway: protein catabolic process, protein ubiquitination, protein polyubiquitination;

hub transcripts: *rps27a, cops5, hap1, rnf8, usp13, fbxo45, smurf2, rplp0, bop1, eif3d*) key to

translation, ubiquitination, and amino acid recycling. Interestingly, *rps27a,* with dual roles

providing both ubiquitin and ribosome structural components (Klinge et al., 2012; Luo, et al.,

2022), was at the center of the network with the largest number of protein-protein interactions

and betweenness score, emphasizing its role within the regulatory network in response to thermal

stress. These protein synthesis and recycling mechanisms are key to modifying the organism's

existing phenotype as environments shift, and without their proper function may leave the

organism vulnerable to the effects of compounding stressors (Davies et al., 2016).

Increased temperatures also modulated chaperone activity and protein folding processes

(pathways: protein folding, regulation of protein stability, response to heat; hub transcripts:

hsc70, hspa8, hyou1, dnajc2, cct4, unc45b, naca) in the face of thermal stress and reactive

oxygen species formation, consistent with a broad literature of molecular responses to thermal

stress (Place and Hoffman, 2001; Jackson and Hewitt 2016; Vallin and Grantham, 2019; Penny

et al, 2023). Modulation of these chaperone proteins, paired with an observed decrease in

metabolic rate for lake sturgeon in temperatures of 24°C from previous research (Bugg et al.,

2020), suggest that there was a transcriptional modification of heat shock protein production

under thermal stress. Transcriptionally modification of heat shock protein production and

decreased mitochondrial oxygen consumption (Bugg et al., 2020) could aid to suppress the

formation of reactive oxygen species as a mechanism to stabilize the existing cellular proteome

to enhance long-term fitness and survival (Storey and Storey, 2011).

Finally, elevated temperatures altered immune system processes (pathways:

inflammation, response to virus, immune system process, adaptive immune response, cytokine-

mediated signaling pathway, lymphocyte differentiation; hub transcripts: *traf6, b2m, banf1,*

rsad2, egr1, jak1, prkcbb). Pathways and related hub genes indicate that there were altered innate

and adaptive immune markers associated with viral infection suggesting immune compromise

and increased susceptibility to pathogens as temperatures increased, with a potential threshold of

20°C for immunocompetence (Choi, 2005; Li et al., 2016; Wei et al., 2017; Ebrahimi, 2018).

Thus, future increases in temperature will not only have direct effects on the lake sturgeon's

thermally responsive mechanisms, but likely indirect effects on pathogen tolerance and survival

as combining stressors mount (Dittmar et al., 2014; Alfonso et al., 2020).

Mitochondrial Limits Under Thermal Stress

Based on transcriptional evidence, mitochondrial function is likely inhibited across both

populations of lake sturgeon as temperatures increased to 24°C. Many mitochondrial processes

were downregulated in response to initial thermal stress at 20°C and continued to decline as

temperatures increased. Transcriptional markers of dysfunction and resulting damages were

apparent at 20°C in northern Burntwood River sturgeon and were more extreme when compared

to southern Winnipeg River sturgeon, highlighting potential population-specific mitochondrial

capacity. Additionally, a highly differentially expressed hub transcript (*gfm1*) key to

mitochondrial translation and oxidative phosphorylation (Berenguer et al., 2021) was altered as

temperatures increased. Aside from the direct effects of decreased ATP production and

increased reactive oxygen species damage, a decline in mitochondrial function also has

implications for stress signaling, viral immune response, and formation of heme as well as iron

cluster assemblies required for oxygen transport and numerous cellular processes (Tovar et al.,

2003; Ajioka et al., 2006; Lin et al., 2006; Qin et al., 2010; Karnkowska et al., 2016;

Hernansanz-Agustin et al., 2020; Braymer et al., 2021). The impairment of these widespread

mitochondrial processes in developing lake sturgeon may inhibit the function of the

mitochondria itself, related cellular systems, and ultimately gill function, with the potential for

population-specific thresholds to impact whole-organism fitness during changes in

environmental temperatures (Iftikar and Hickey, 2013; Jeffries et al., 2018).

Study Limitations

As lake sturgeon are an octoploid species, transcriptomic analysis can be vulnerable to

the effects of gene duplication, resulting homeologs, and their effect on the interpretation of

resulting sequence data (Ogden et al., 2013; Limborg et al., 2016). To mitigate this effect, we

selected a tissue-specific, *de novo*, sequencing approach to help isolate different gene isoforms.

Additionally, we selected a transcriptome assembly program, Trinity, that has been shown to

have success with the accurate assembly of polyploid transcriptomes (Grabherr et al., 2011;

Voshall & Moriyama, 2020). With respect to SNP calling using the transcriptome, we used a

splice-aware read mapping program (STAR; Dobin et al., 2013) and addressed paralogous loci

with heterozygosity and read depth ratio deviations (McKinney et al., 2017), in addition to

standard quality control and linkage disequilibrium pruning steps. Nevertheless, the genetic

differentiation presented in the present study is biased with family structure by our experimental

design (studied individuals represent F_1 crosses of wild-caught adults), and therefore are not

representative of differentiation between the two studied populations overall (Kjartanson et al.,

2023). For instance, the PCA with SNPs may not accurately reflect patterns of genetic variation

in F0 wild individuals. Nor are these genetic data appropriate for studying heritability or additive

genetic effects because of the difficulties of ploidy and SNPs with mRNA data. Instead, the

genetic data gathered from fish in each population provides context for the observed

transcriptional plasticity, which is at least partially attributable to genetic differences between the

individuals used in the study.

While the octoploid lake sturgeon genome can complicate transcriptomic analysis (Ogden

et al., 2013; Limborg et al., 2016), the consistency in responses to temperature across the

populations, and with other literature, lends credence to the reliability of the interpretation of

sequencing data. Moreover, the thermal resolution of our study (4°C windows) may make the

exact detection of thermal thresholds difficult, especially if thresholds are non-linear across

populations. Further, this work only represents the findings from two latitudinally distributed

populations. Future research should expand these findings across the full latitudinal distribution

of lake sturgeon, or across other latitudinally distributed sturgeon species.

Ecological Relevance

In the current study, temperatures of 20°C and above limited the transcriptional plasticity

of developing lake sturgeon, which would ultimately diminish their plasticity to other stressors.

As temperatures continue to increase in the wild, the combined effects of thermal, oxidative, and

pathogen related stressors will likely limit the survival of lake sturgeon populations within

Manitoba, Canada, with increased severity for northern sub-Arctic populations throughout the

province (Bugg et al., 2020). The population designation units now group lake sturgeon

inhabiting northern regions with those from several other populations in Manitoba to form a

larger designation unit (COSEWIC 2006; COSEWIC 2017). However, new evidence suggesting

large discrepancies in transcriptomic, physiological, and genetic differences between

populations, as well as their natural thermal environments and tolerance thresholds (McDougall

et al., 2017; Bugg et al., 2020; Bugg et al., 2021; Kjartanson et al., 2023) indicate that they are

differentially susceptible to increasing temperatures and require distinct management strategies

for conservation. While the experimental design used precludes analyses of heritability and

additive effects of genetic variation underlying the observed transcriptional plasticity, these

results are nevertheless valuable for their physiological and population-level implications.

Currently, temperatures in the Winnipeg River exceed 20°C for approximately 50 days

per year, while this temperature threshold is projected to be exceeded during June, July, and

August in the Burntwood River in the coming decades (Manitoba Hydro, 2015; Vincent et al.,

2015; Zhang et al., 2019; Bugg et al., 2020). These seasonal increases in temperature occur when

newly hatched sturgeon are in their first few months of life, which are the same life stages

examined in the current study, and could exacerbate already high rates of mortality during this

period (Bugg et al., 2020; Bugg et al., 2021). As temperatures increase transcriptional

modification occurs, but many key mechanisms across both populations are downregulated by

24°C. Additionally, endemic viral pathogens are persistent in both northern and southern

populations of lake sturgeon throughout the province, with higher abundance and probability of

detection in younger fish (Clouthier et al., 2013; Clouthier et al., 2020), and as temperatures

increase. With the present experimental evidence, we suggest that if river temperatures continue

to increase as projected (Manitoba Hydro, 2015) and there are not sufficient thermal refugia,

these physiological limitations will likely impede future lake sturgeon recruitment, especially in

the most northern populations. Therefore, increasing temperatures may influence recruitment,

movement patterns, and distributions of lake sturgeon within Manitoba, as sturgeon shift to areas

of thermal refuge to preserve energy and persist in the face of environmental change (Moore et

al., 2020; Robertson et al., 2021). However, even with sufficient refugia, fish do not always

travel to habitat with increased suitability due to life history constraints (Sutton et al., 2007;

White et al., 2019). If lake sturgeon are unable to redistribute themselves in areas of thermal

refuge, evidence suggests that the impacts of increasing environmental temperature may be more

severe, with impacts to population-level processes and survival based on observed transcriptional

plasticity and physiological responses when exposed to elevated temperatures.

Data Accessibility:

The data used in this study is available in the included supplementary files (1-8) and the raw

sequencing reads are available at the National Center for Biotechnology Information Sequence

Read Archive with the accession number PRJNA815828. The Trinity assembly and Trinotate

annotation file are available via figshare (https://doi.org/10.6084/m9.figshare.19209753.v1).

Author Contributions:

WSB, WGA, and KMJ designed the research. WSB analyzed the count data with assistance from

KM and MJT. MJT analyzed the single nucleotide polymorphism data. WSB wrote the first draft

of the manuscript and all authors provided comments on the final version.

Acknowledgements:

The authors thank North/South Consultants Inc. and Manitoba Hydro for their assistance in the

capture of spawning adults and collection of gametes from these wild sturgeon populations,

Evelien de Greef for her support with the map (Figure 1), as well as Madison Earhart for

valuable discussions and sturgeon illustrations (Figure 2 and Figure 3A). We would also like to

thank the staff at the University of Manitoba Animal Holding Facility and numerous Anderson

and Jeffries Laboratory members for assistance in the care and maintenance of fish. This work

was conducted at the University of Manitoba located on original lands of the Anishinaabeg,

Cree, Oji-Cree, Dakota and Dene peoples, and on the homeland of the Métis Nation. We

recognize that water supplied for our fish at the University of Manitoba was sourced from the

Shoal Lake 40 First Nation. Many bioinformatics steps were enabled by our chance to use

Digital Research Alliance of Canada computing clusters. Funding for this study was provided by

NSERC/Manitoba Hydro Industrial Research Chair awarded to W.G.A. and NSERC Discovery

Grants (grant numbers 05348 and 05479) awarded to W.G.A. and K.M.J., respectively. W.S.B.

was supported by a University of Manitoba Graduate Fellowship as well as the International

Graduate Student Scholarship.

Figures:

Figure 1. Geographic localities of two lake sturgeon populations (*Acipenser fulvescens*) within

Manitoba, Canada, highlighting the northern Burntwood River (BWR) and more southernly

(WR) sturgeon populations, separated by approximately 650 km, as well as the direction of flow

throughout the watershed. This figure was assembled with the assistance of Evelien de Greef.

Figure 2. Principal components analysis of 14,517 mRNA SNPs called in lake sturgeon

(*Acipenser fulvescens*) from the northern Burntwood River and southern Winnipeg River

populations within Manitoba, Canada.

Figure 3. A) Principal component analysis and B) Numbers of differentially expressed transcripts across all transcripts, as well as C) Aggregate heat map of the top 500 differentially expressed transcripts between each contrast of northern Burntwood River and southern Winnipeg River populations of lake sturgeon within Manitoba, Canada, acclimated to $16°$C, $20°$C, and $24°$C for 30 days in early development, measured using RNAseq. Transcript numbers represent upregulation and downregulation with a $\log_2$ fold change > 1, respectively, and with a q-value (FDR adjusted p value) < 0.05. Contrasts are oriented Burntwood River-Winnipeg River at the interpopulation level and lower temperature-higher temperature across acclimation treatments (e.g. up means increased with temperature, and in the Burntwood River population, relative to Winnipeg River). Each population and acclimation treatment is represented by 6 individual gill samples (n = 6). For the aggregate heat map, only unique transcripts were included for a total of 2,800 transcripts represented. Data are expressed as $\log_2$ counts per million (CPM).

Figure 4. REVIGO (Reduce + Visualize Gene Ontology) plots of GO (gene ontology) biological processes terms that decreased during a 30-day thermal acclimation from both 16 to 20°C and 16 to 24°C for developing northern and southern lake sturgeon (*Acipenser fulvescens*) populations within Manitoba, Canada, measured using RNAseq. These terms were conserved across both the southern Winnipeg River (WR) and northern Burntwood River (BWR) lake sturgeon populations as temperatures increased. Bubble color indicates the significance of the adjusted *p* value at after applying a log$_{10}$ scale (darker color is a more significant term), while bubble size indicates the frequency of a GO term GO annotation database (larger bubble is a more common GO term in the data set).

Figure 5. Aggregate most differentially expressed transcript network demonstrating predicted

protein-protein interactions across the most upregulated and downregulated transcripts of

northern and southern lake sturgeon, *Acipenser fulvescens*, populations, within Manitoba,

Canada, acclimated to 16°C, 20°C, and 24°C for 30 days in early development, measured using

RNAseq. Orange transcript circles outlined in blue with bolded names indicate hub transcripts

within the network. Hub size and naming is scaled according to their degree of interaction

(number of connections) with surrounding hubs (Zhou and Xia, 2019).

Supplemental Figures:

Supplemental Figure 1. Admixture plot illustration the distribution of SNPs between northern

Burntwood River and Southern Winnipeg River populations of lake sturgeon, *Acipenser*

fulvescens, within Manitoba, Canada.

Supplementary Figure 2. GO (gene ontology) term distribution illustrated using UpSet plots

demonstrating shared and population-specific terms across A) all split-contrasts and B) split

contrasts with terms decreasing as environmental temperatures increase. *'s on split-contrast in

A indicate the contrasts which were included in B. These data represent change in the

transcriptional mechanisms in the gill of developing lake sturgeon from both a northern

Burntwood River (BWR) and southern Winnipeg River (WR) population of lake sturgeon

(*Acipenser fulvescens*), within Manitoba, Canada, acclimated to temperatures 16°C, 20°C, and

24°C for 30 days in early development, measured using RNAseq. Individual dots indicate

processes unique to that contrast, while a line connecting multiple dots indicate processes that

are shared between the contrast connected by the line. Split-contrasts are oriented BWR-WR at

the interpopulation level and lower temperature-higher temperature across acclimation

treatments so that "neg" indicates upregulation at higher temperatures, and "pos" indicates

downregulation at higher temperatures (e.g. WR16v20_pos includes terms that were

downregulated in the WR as temperatures increased). Each population and acclimation treatment

is represented by 6 individual gill samples (n = 6).

Supplementary Figure 3. REVIGO (Reduce + Visualize Gene Ontology) plots of GO (gene

ontology) population-specific biological processes terms for the Winnipeg River and Burntwood

River populations of developing lake sturgeon (*Acipenser fulvescens*) within Manitoba, Canada,

as acclimation temperatures increase during a 30-day acclimation, measured using RNAseq.

Plots represent A) Winnipeg River population-specific responses downregulated from 16 to 20°C

B) Burntwood River population-specific responses downregulated from 16 to 20°C) Winnipeg

River population-specific responses decreased from 16 to 24°C. Bubble color indicates the

significance of the adjusted p value after applying a $\log_{10}$ scale (darker color is a more significant

term), while bubble size indicates the frequency of a GO term GO annotation database (larger

bubble is a more common GO term in the data set).

References:

Aijoka, R.S., Philips, J.D., Kushner, J.P. 2006. Biosynthesis of heme in mammals. *Biochim Biophys Acta*. 1763, 723-736. https://doi.org/10.1016/j.bbamcr.2006.05.005

Akbarzadeh, A., Günther, O.P., Houde, A.L., Li, S., Ming, T.J., Jeffries, K.M., Hinch, S.G., Miller, K.M. 2018. Developing specific molecular biomarkers for thermal stress in salmonids. *BMC Genom*. 19, 749. https://doi.org/10.1186/s12864-018-5108-9

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. Genome research, 19(9), 1655-1664.

Alvarez, M., Schrey, A., Richards, C. 2015. Ten years of transcriptomics in wild populations: What have we learned about their ecology and evolution? *Mol Ecol*. 24, 710–725. https://doi.org/10.1111/mec.13055

Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data.524https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Arthington, A.H., Dulvy, N.K., Gladstone, W., Winfield, I.J. 2016. Fish conservation in freshwater and marine realms: status, threats and management. *Aquatic Conservation*. 26, 838-857. https://doi.org/10.1002/aqc.2712

Bacevic, K., Lossaint, G., Achour, T.N., Georget, V., Fisher, D., Dulic, V. 2017. Cdk2 strengthens the intra-S checkpoint and counteracts cell cycle exit induced DNA damage. *Sci Rep*. 7, 13429. https://doi.org/10.1038/s41598-017-12868-5

Baker, D.W., Matey, V., Huynh, K.T., Wilson, J.M., Morgan, J.D., Brauner, C.J. 2009. Complete intracellular pH protection during extracellular pH depression is associated with hypercarbia tolerance in white sturgeon, *Acipenser transmontanus*. *Am J Physiol Regul Integr Comp Physiol*. 296, R1868-R1880. https://doi.org/10.1152/ajpregu.90767.2008

Barley, J.M., Cheng, B.S., Sasaki, M., Gignoux-Wolfsohn, S., Hays, C.G., Putnam, A.B., Sheth, S., Villeneuve, A.R., Kelly, M. 2021. Limited plasticity in thermally tolerant ectotherm populations: evidence for a trade-off. *Proc Royal Soc B*. 288, 21210765. https://doi.org/10.1098/rspb.2021.0765

Barnes, R.P., Fouquerel, E., Opresko, P.L. 2019. The impact of oxidative DNA damage and stress on telomere homeostasis. *Mechanisms of Ageing and Development*. 177, 37-45. https://doi.org/10.1016/j.mad.2018.03.013

Barrington, C., Pezic, D., Hadjur, S. 2017. Chromosome structure dynamics during the cell cycle: a structure to fit every phase. *The EMBO Journal*. 36, 2661-2663. https://doi.org/10.15252/embj.201798014

Berenguer, M.M., Julia, F.V., Ramos, S.P., Allende, M.T.S., Camara, Y., Torronteras, J.V., Marti, R. 2021. Dysfunctional mitochondrial translation and combined oxidative

phosphorylation deficiency in a mouse model of hepatoencephalopathy due to *Gfm1* mutations. *FASEB*. 36, e22091.

Bertoli, C., Skotheim, J.M., de Bruin, R.A.M. 2015. Control of cell cycle transcription during G1 and S phases. *Nat Rev Mol Cell Biol*. 14, 518-528. https://doi.org/10.1038/nrm3629

Birnie-Gauvin, K., Costantini, D., Cooke, S.J., Willmore, W.G. 2017. A comparative and evolutionary approach to oxidative stress in fish: A review. *Fish Fish*. 18, 928-942. https://doi.org/10.1111/faf.12215

Bolger, A.M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics*, btu170. https://doi.org/10.1093/bioinformatics/btu170

Braasch, I., Postlethwait, J.H. 2012. Polyploidy in Fish and the Teleost Genome Duplication. In: Soltis P., Soltis D. (eds) Polyploidy and Genome Evolution. Springer, Berlin, Heidelberg. pp. 341-383.

Brandt, C., Bugg, W., Groening, L., Klassen, C., Anderson, W.G. 2021. Effects of rearing temperature on volitional and escape response swimming performance in lake sturgeon, *Acipenser fulvescens*, from hatch to age 1. *Environ Biol Fishes*. 104, 737-750. https://doi.org/10.1007/s10641-021-01112-9.

Brandt, C., Groening, L., Klassen, C., Anderson, W.G. Effects of rearing temperature on yolksac volume and growth rate in Lake Sturgeon, *A. fulvescens*, from hatch to age-1. *Aquaculture*. 546, 737352. https://doi.org/10.1016/j.aquaculture.2021.737352

Braymer, J.J., Freibert, S.A., Rakwalska-Bange, M., Lill. 2021. Mechanistic concepts of iron-sulfur protein biogenesis in biology. *Biochim Biophys Acta Mol Cell Res*. 1868, 118863. https://doi.org/10.1016/j.bbamcr.2020.118863

Brennan, B.S., deMayo, J.A., Dam, H.G., Finiguerra, M.B., Baumann, H., Pespeni, M.H. 2022. Loss of transcriptional plasticity but sustained adaptive capacity after adaptation to global change conditions in a marine copepod. *Nat Commun*. 13, 1147. https://doi.org/10.1038/s41467-022-28742-6

Buccitelli, C., Selbach, M. 2020. mRNAs, proteins and the emerging principles of gene expression control. *Nat Rev Genet*. 21, 630-644. https://doi.org/10.1038/s41576-020-0258-4

Bugg, W.S., Yoon, G.R., Brandt, C., Earhart, M., Anderson, W.G., Jeffries, K.M. 2021. The effects of population and thermal acclimation on the growth, condition, and cold responsive mRNA expression of age-0 lake sturgeon (*Acipenser fulvescens*). *J Fish Biol*. 99, 1-16. https://doi.org/10.1111/jfb.14897

Bugg, W.S., Yoon, G.R., Schoen, A., Laluk, A., Brandt, C., Anderson, W.G., Jeffries, K.M. 2020. Effects of Acclimation Temperature on the Thermal Physiology in two geographically distinct Populations of Age-0 Lake Sturgeon (*Acipenser fulvescens*). Conservation Physiology. 8, coaa087. https://doi.org/10.1093/conphys/coaa087

Burggren, W. 2018. Developmental phenotypic plasticity helps bridge stochastic weather events
associated with climate change. *J Exp Biol*. 221: jeb161984.
https://doi.org/10.1242/jeb.161984

Burggren, W., Crews, D. 2014. Epigenetics in comparative biology: why we should pay
attention. *Integr Comp Biol*. 54, 7-20. https://doi.org/10.1093/icb/icu013

Burkhead, N. M. 2012. Extinction rates in North American freshwater fishes, 1900–2010.
BioScience. 62, 798–808. https://doi.org/10.1525/bio.2012.62.9.5

Caroffino, D.C., Sutton, T.M.S., Elliot, R.F., Donofrio, M.C. 2010. Early Life Stage Mortality
Rates of Lake Sturgeon in the Peshtigo River, Wisconsin. *N Am J Fish Manag*. 30, 295-
304. https://doi.org/10.1577/M09-082.1

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an
analysis tool set for population genomics. Molecular ecology, 22(11), 3124-3140.

Chandel, N.S. 2010. Mitochondrial Complex III: An essential component of universal oxygen
sensing machinery? *Respir Physiol Nuerobiol*. 174, 175-181.
https://doi.org/10.1016/j.resp.2010.08.004

Chatterjee, N., Walker, G.C. 2017. Mechanisms of DNA damage, repair and mutagenesis.
Environ Mol Mutagen. 58, 235-263.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., Ma'ayan, A.
2013. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool.
BMC Bioinform. 14, 128. https://doi.org/10.1186/1471-2105-14-128

Chen, S., Zhou, Y., Chen, Y., Gu, J. 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics, 34(17), i884-i890.

Choi, Y., 2005. Role of TRAF6 in the immune system. *Mechanisms of Lymphocyte Activation*
and Immune Regulation X, pp.77-82.
1046
1047 Chung, D.J., Schulte, P.M. 2015. Mechanisms and costs of mitochondrial thermal acclimation in
1048   a eurythermal killifish (*Fundulus heteroclitus*). *Journal of Experimental Biology*. 218,
1049   1621-1631.
1050
Cioce, M., Lamond, A.I. 2005. CAJAL BODIES: A Long History of Discovery. *Annu Rev Cell*
Dev Biol. 21, 105-131. https://doi.org/10.1146/annurev.cellbio.20.010403.103738

Cleator, H., Martin, K.A., Pratt, T.C., Macdonald, D. 2010. Information relevant to a recovery
potential assessment of Lake Sturgeon: Nelson River population (DU3). DFO Canadian
Science Advisory Secretariat Document 2010/082. vi + 33p

Clouthier, S., VanWalleghem, E., Copeland, S., Klassen, C., Hobbs, G., Nielsen, O., Anderson,
E.D. 2013. A new species of nucleo-cytoplasmic large DNA virus (NCLDV) associated
with mortalities in Manitoba Lake sturgeon *Acipenser fulvescens*. *Dis Aquat Org*. 102,
195-209.

Clouthier, S., Caskenette, A., Walleghem, E.V., Schroeder, T., Macdonald, D., Anderson, E.D.
2020. Molecular phylogeny of sturgeon mimiviruses and Bayesian hierarchical modeling
of their effect on wild Lake Sturgeon (*Acipenser fulvescens*) in Central Canada. *Infect*
Genet Evol. 84, 104491. https://doi.org/10.1016/j.meegid.2020.104491

Comte, L., Buisson, L., Daufresne, M., Grenouillet, G. 2012. Climate-induced changes in the
distribution of freshwater fish: observed and predicted trends. *Freshw Biol*. 58, 625-639.
https://doi.org/10.1111/fwb.12081

Connon, R.E., Jeffries, K.M., Komoroske, L.M., Todgham, A.E., Fangue, N.A. 2018. The utility
of transcriptomics in fish conservation. *J Exp Biol*. 221, jeb148833.
https://doi.org/10.1242/jeb.148833

Conway, J.R., Lex, A., Gehlenborg, N. 2017. UpSetR: an R package for visualization of
intersecting sets and their properties. *Bioinformatics*. 33, 2938-2940.
https://doi.org/10.1093/bioinformatics/btx364

COSEWIC. 2006. COSEWIC assessment and update status report on the lake sturgeon
Acipenser fulvescens in Canada. COSEWIC, 107.

COSEWIC. 2017. COSEWIC assessment and update status report on the Lake Sturgeon
Acipenser fulvescens, Western Hudson Bay populations, Saskatchewan-Nelson River
populations, Southern Hudson Bay – James Bay populations and Great Lakes – Upper St.
Lawrence populations in Canada. Committee on the Status of Endangered Wildlife in
Canada, Ottawa, ON. xxx + 153 p.

Crozier, L.G., Hutchings, J.A. 2014. Plastic and evolutionary responses to climate change in fish.
Evolutionary applications. 7, 68-87. https://doi.org/10.1111/eva.12135

Davies, S.W., Marchetti, A., Ries, J.B., Castillo, K.D. 2016. Thermal and pCO2 Stress Elicit
Divergent Transcriptomic Responses in a Resilient Coral. *Frontiers in Marine Science*.
https://doi.org/10.3389/fmars.2016.00112

Deslauriers, D., Kieffer, J.D. 2012. Swimming performance and behaviour of young-of-the-year
shortnose sturgeon (*Acipenser brevirostrum*) under fixed and increased velocity
swimming tests. *Can J Zool*. 351, 345–351. https://doi.org/10.1139/z2012-004

Dillon, M.E., Woods, H.A., Wang, G., Fey, S.B., Vasseur, D.A., Telemeco, R.S., Marshall, K.,
Pincebourde, S. 2016. Life in the Frequency Domain: the Biological Impacts of Changes
in Climate Variability at Multiple Time Scales. *ICB*.56, 14-30.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., ... & Gingeras, T. R.
(2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics, 29(1), 15-21.

Earhart, M.L., Blanchard, T.S., Harman, A.A., Schulte, P.M. 2022. Hypoxia and High
Temperature as Interacting Stressors: Will Plasticity Promote Resilience of Fishes in a
Changing World? *The Biological Bulletin*. 243, 85-103. https://doi.org/10.1086/722115

Ebrahimi, K.H. 2018. A unifying view of the broad-spectrum antiviral activity of RSAD2
(viperin) based on its radical-SAM chemistry. *Metallomics*. 10, 539-552.

Ellis, L.L., Huang, W., Quinn, A.M., Ahuja, A., Alfrejd, B., Gomez, F.E., Hjelmen, C.E., Moore,
K.L., Mackay, T.F.C., Johnston, J.P., Tarone, A.M. 2014. Intrapopulation Genome Size
Variation in *D. melanogaster* Reflects Life History Variation and Plasticity. *PLoS Genet*.
10, e1004522. https://doi.org/10.1371/journal.pgen.1004522

Filho, D.W. 2007. Reactive oxygen species, antioxidants and fish mitochondria. *Front Biosci*.
12, 1229-1237. https://doi.org/10.2741/2141

Fox, R.J., Donelson, J.M., Schunter, C., Ravasi, T., Espitia, J.D.G. 2019. Beyond buying time:
the role of plasticity in phenotypic adaptation to rapid environmental change.
Philosophical Transactions of the Royal Society B. 374, 20180174.
https://doi.org/10.1098/rstb.2018.0174

Francis, R. M. (2017). pophelper: an R package and web app to analyse and visualize population
structure. Molecular ecology resources, 17(1), 27-32.

Fujimoto, M., Lovett, B., Angoshtari, R., Nirenberg, P., Loch, T.P., Scribner, K.T., Marsh, T.L.
2018. Antagonistic interactions and biofilm forming capabilities among bacterial strains
isolated from egg surfaces of lake sturgeon (*Acipenser fulvescens*). *Microb Ecol*. 75, 22-
37. https://doi.org/10.1007/s00248-017-1013-z

Gardil, A.M., Wright, G.M., Wadowska, D.W., Fast, M.D. 2014b. Ontogeny of the immune
system in acipenserid juveniles. *Dev Comp Immunol*. 44, 303-314.
https://doi.org/10.1016/j.dci.2014.01.006

Georgiadis, M.P., Hedrick, R.P., Carpenter, T.E., Gardner, I.A., 2001. Factors influencing
transmission, onset and severity of outbreaks due to white sturgeon iridovirus in a
commercial hatchery. *Aquaculture*. 194, 21–35. https://doi.org/10.1016/S0044-
8486(00)00503-2

Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y.,
Blankenberg, D., Albert, I., Taylor, J., et al. 2005. Galaxy: A platform for interactive
large-scale genome analysis. *Genome Res*. 15, 1451-1455.
https://doi.org/10.1101/gr.4086505

Gilmour, K.M., Perry, S.F. 2018. Conflict and Compromise: Using Reversible Remodeling to
Manage Competing Physiological Demands at the Fish Gill. *Am Physio. Soc.* 33, 412-
422. https://doi.org/10.1152/physiol.00031.2018

Gomez-Gallardo, M., Sanchez-Briones, L.A., Diaz-Perez, A.L., Lopez-Alvarez, A., Cortes-Rojo,
C., Aguilar-Mendez., Saavedra-Molina, A., Campos-Garcia, J. 2016. The Function of the
Iron-Sulfur Cluster Assembly Machinery is involved in Electron Trasport Chain Activity
and Oxidative Stress in *Saccharomyces cerevisiae*. *FASEB J*. 30, 863.1-863.1.
https://doi.org/10.1096/fasebj.30.1_supplement.863.1

Goudet, J. (2005). Hierfstat, a package for R to compute and test hierarchical F-statistics. Molecular ecology notes, 5(1), 184-186.

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regevm A. 2011. Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq data. *Nat Biotechnol*. 29, 644-652. https://doi.org/10.1038/nbt.1883

Gradil, A.M., Wright, G.M., Speare, D.J., Wadowksa, D.W., Purcell, S., Fast, M.D. 2014a. The effects of temperature and body size on immunological development and responsiveness in juvenile shortnose sturgeon (*Acipenser brevirostrum*). *Fish Shellfish Immunol*. 40, 545-555. https://doi.org/10.1016/j.fsi.2014.07.036

Greene, D.H.S., Selivonchick, D.P. 1987. Lipid Metabolism in fish. *Progress in Lipid Research*. 26, 53-86.

Grill, G., et al. "Mapping the world's free-flowing rivers." Nature, vol. 569, no. 7755, 2019, p. 215. Gale OneFile: CPI.Q, link.gale.com/apps/doc/A593382054/CPI?u=winn62981&sid=CPI&xid=ff4c09e3. Accessed 25 May 2021. https://doi.org/10.1038/s41586-019-1111-9

Guo, C., Sun, L., Chen, X., Zhang, D. 2013. Oxidative stress, mitochondrial damage and neurodegenerative diseases. *Nueral Regen Res*. 25, 2003-2014. https://doi.org/10.3969/j.issn.1673-5374.2013.21.009

Guo, K., Zhao, C., Lang, B., Wang, H., Zheng, H., and Zhang, F. 2021. Regulator of Chromosome Condensation 2 Modulates Cell Cycle Progression, Tumorigenesis, and Therapeutic Resistance. *Front mol biosci*. 7:620973. https://doi.org/10.3389/fmolb.2020.620973

Healy, T. M., Schulte, P. M. 2019. Patterns of alternative splicing in response to cold acclimation in fish. *J Exp Biol*. 222, jeb193516. https://doi.org/10.1242/jeb.193516.

Hernansanz-Agustin, P., et al., 2020. Na+ controls hypoxic signaling by the mitochondrial respiratory chain. 2020. *Nature*. 586, 287-291. https://doi.org/10.1038/s41586-020-2551-y

Hwang, P.P., Lee, T.H., Lin, L.Y. 2011. Ion regulation in fish gills: recent progress in the cellular and molecular mechanisms. *Am J Physiol Regul Integr Comp Physiol*. 301, R28-47. https://doi.org/10.1152/ajpregu.00047.2011

Iftikar, F.I., Hickey, A.J.R. 2013. Do Mitochondria Limit Hot Fish Hearts? Understanding the Role of Mitochondrial Function with Heat Stress in *Notolabrus celidotus*. *PLOS ONE*. 8, 64120. https://doi.org/10.1371/journal.pone.0064120

IUCN, *The IUCN Red List of Threatened Species* Version 2017-3 (2018); www.iucnredlist.org.

Jackson, M.P., Hewitt, E.W. 2016. Cellular proteostasis: degradation of misfolded proteins by
lysosomes. *Essays Biochem*. 60, 173-180. https://doi.org/10.1042/EBC20160005

Jane, S.F., Gretchen, J.A., et al. 2021. Widespread deoxygenation of temperate lakes. *Nature*.
594, 60-70. https://doi.org/10.1038/s41586-021-03550-y

Jeffries, K. M., Connon, R. E., Verhille, C. E., Dabruzzi, T. F., Britton, M. T., Durbin-Johnson,
B. P., Fangue, N. A. 2019. Divergent transcriptomic signatures in response to salinity
exposure in two populations of an estuarine fish. *Evol App*. eva.12799.
https://doi.org/10.1111/eva.12799

Jeffries, K. M., Teffer, A., Michaleski, S., Bernier, N. J., Heath, D. D., & Miller, K. M. 2021.
The use of non-lethal sampling for transcriptomics to assess the physiological status of
wild fishes. *Comp Biochem Physiol B, Biochem.* 256, 110629.
https://doi.org/10.1016/j.cbpb.2021.110629

Jeffries, K.M. Fangue, N.A., Connon, R.E. 2018. Multiple sub-lethal thresholds for cellular
responses to thermal stressors in an estuarine fish. *Comp Bio Phys A*. 255, 33-45.
https://doi.org/10.1016/j.cbpa.2018.06.020

Jeffries, K.M., Komoroske, L.M., Truong, J., Werner, I., Matthias, H., Hasenbein, S., Fangue,
N.A., Connon, R.E. 2015. The transcriptome-wide effects of exposure to a pyrethroid
pesticide on the critically endangered delta smelt *Hypomesus transpacificus*. *Endanger
Species Res*. 28, 43-60. https://doi.org/10.3354/esr00679

Jonsson, B., Jonsson, N. 2019. Phenotypic plasticity and epigenetics of fish: embryo temperature
affects later-developing life history traits. *Aquat Biol*. 28, 21-32.
https://doi.org/10.3354/ab00707

Karnkokowska, A., Vacet, V., et al. 2016. A Eukaryote without a Mitochondrial Organelle. *Curr
Biol*. 26, 1274-1284. https://doi.org/10.1016/j.cub.2016.03.053

Kim, D., Langmead, B., Salzberg, S.L. 2015. HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods.* 12, 357-360. https://doi:10.1038/nmeth.3317

Kjartanson, S.L., Haxton, T., Wozney, K., Lovejoy, N., Wilson, C.C. 2023. Conservation
Genetics of Lake Sturgeon (*Acipenser fulvescens*): Nuclear Phylogeography Drives
Contemporary Patterns of Genetic Structure and Diversity. *Diversity*. 15, 385.
https://doi.org/10.3390/d15030385

Klinge, S., Hoffmann, F.V., Leibundgut, M., Ban, N. 2012. Atomic structures of the eukaryotic
ribosome. *Trends in Biochemical Sciences*. 37, 189-198.

Kolde, R. 2015. pheatmap: Pretty heatmaps. https://CRAN. R-project. org/package= pheatmap

Komoroske, L.M., Connon, R.E., Jeffries, K.M., Fangue, N.A. 2015. Linking transcriptional
responses to organismal tolerance reveals mechanisms of thermal sensitivity in a
mesothermal endangered fish. *Mol Ecol*. 24, 4960-4981. https://doi:10.1111/mec.13373

Komoroske, L.M., Jeffries, K.M., Whitehead, A., Roach, J.L., Britton, M., Connon, R.E.,
Verhille, C., Brander, S.M., Fangue, N.A. 2020. Transcriptional flexibility during thermal
challenge corresponds with expanded thermal tolerance in an invasive compared to native
fish. *Evol Appl.* 14, 931-949. https://doi.org/10.1111/eva.13172

Krabbenhoft, T.J., Myers, B.J.E., Wong, J.P., Chu, C., Tingley, R.W., Falke, J.A., Kwak, T.J.,
Paukert, C.P., Lynch, A.J. 2020. FiCli, the Fish and Climate Change Database, informs
climate adaptation and management for freshwater fishes. *Sci Data.* 7, 1–6.
https://doi.org/10.1038/s41597-020-0465-z

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S.,
Jenkins, S.L., Jagodnik, K.M., Lachmann, A., McDermott, M.G., Monteiro, C.D.,
Gundersen, G.W., Ma'ayan, A. 2016. Enrichr: a comprehensive gene set enrichment
analysis web server 2016 update. *Nucleic Acids Res.* https://doi.org/gkw377.
10.1093/nar/gkw377

Lafuente, E., Beldade, P. 2019. Genomics of Developmental Plasticity in Animals. *Front Genet.*
10:720. https://doi.org/10.3389/fgene.2019.00720

LaPatra, S.E., Groff, J.M., Jones, G.R., Munn, B., Patterson, T.L., Holt, R.A., Hauck, A.K.,
Hedrick, R.P., 1994. Occurrence of white sturgeon iridovirus infections among cultured
white sturgeon in the Pacific Northwest. *Aquaculture.* 126, 201–210.
https://doi.org/10.1016/0044-8486(94)90036-1

Lazado, C.C., Caipang, C.M.A. 2014. Mucosal immunity and probiotics in fish. *Fish Shellfish
Immunol.* 39, 78-89. https://doi.org/10.1016/j.fsi.2014.04.015

Liao, Y., Smyth, G.K., Shi, W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics.* 30, 923-930.
https://doi.org/10.1093/bioinformatics/btt656

Limborg, Sweeb, L.W., Seeb, J.E. 2016. Sorting duplicated loci disentangles complexities of
polyploid genomes masked by genotyping by sequencing. *Mol Ecol.* 25, 2117-29.
https://doi.org/10.1111/mec.13601

Li, L., Dong, M., Wang, X.G. 2016. The Implication and Significance of Beta 2 Microglobulin:
A conservative Multifunctional Regulator. *Chin Med J.* 129, 448-455.

Lin, R., Lacoste, J., Nakhaei, P., Sun, Q., Yang, L., Paz, S., Wilkinson, P., Julkunen, I., Vitour,
D., Meurs, E., Hiscott, J. 2006. Dissociation of MAVS/IPS-1/VISA/Cardif-IKKe
Molecular Complex from the Mitochondrial Outer Membrane by Hepatitis C Virus NS3-
4A Proteolytic Cleavage. *J Virol.* 80, 6072-6083. https://doi.org/10.1128/JVI.02495-05

Logan, M.L., Cox, C.L. 2020. Genetic Constraints, Transcriptome Plasticity, and the
Evolutionary Response to Climate Change. *Front Genet.* 11, 538226.
https://doi.org/10.3389/fgene.2020.538226

Lou, J., Zhao, H., Chen, L., Liu, M. 2022. Multifaceted functions of RPS27a: An unconventional
ribosomal protein. *Journal of Cellular Physiology.* https://doi.org/10.1002/jcp.30941

Manitoba Hydro .2015. Manitoba Hydro Climate Change Report Fiscal Year 2014–2015.
https://www.hydro.mb.ca/environment/pdf/climate_change_report_2014_15.pdf. (last
accessed 12 May 2020).

Marcos-López, M., Gale, P., Oidtmann, B.C., Peeler, E.J. 2010. Assessing the Impact of Climate
Change on Disease Emergence in Freshwater Fish in the United Kingdom. *Transbound*
Emerg Dis. 57, 293-304. https://doi/10.1111/j.1865-1682.2010.01150.x

Mastrangelo, A.M., Marone, D., Laido, G., Leonardis, A.M., Vita, P.D. 2012. Alternative
splicing: Enhanced ability to cope with stress via transcriptome plasticity. *Plant Science*.
185, 40-49.

McDougall, C.A., Welsh, A.B., Gosselin, T., Anders, W.G., Nelson, P.A. 2017. Rethinking the
influence of hydroelectric development on gene flow in long-lived fish, the lake sturgeon
Acipenser fulvescens. *PLoS One*. 12: e0174269.
https://doi/10.1371/journal.pone.0174269.

McKinney, G. J., Waples, R. K., Seeb, L. W., & Seeb, J. E. (2017). Paralogs are revealed by
proportion of heterozygotes and deviations in read ratios in genotyping-by-sequencing
data from natural populations. Molecular Ecology Resources, 17(4), 656-669.

Metzger, D.C.H., Schulte, P.M. 2016. Epigenomics in marine fishes. *Marine Genomics*. 30, 43-
54.

Miao, T., Symonds, A.L.J., Singh, R., Symonds, J.D., Ogbe, A., Omodho, B., Zhu, B., Li, S.,
Wang, P. 2017. Egr2 and 3 control adaptive immune responses by temporally uncoupling
expansion from T cell differentiation. *J Exp Med*. 214, 1787- 1808.
https://doi/10.1084/jem.20160553

Moore, M.J., Paukert, C.P., Moore, T.L. 2021. Effects of Latitude, Season, and Temperature on
Lake Sturgeon Movement. *N Am J Fish Manag*. 41, 916-928.
https://doi.org/10.1002/nafm.10416.

Morgan, R., Finnoen, M.H., Jensen, H., Pelabon, C., Jutfelt, F. 2020. Low potential for
evolutionary rescue from climate change in a tropical fish. *PNAS*. 117, 33365-33372.
https://doi.org/10.1073/pnas.2011419117.

Mundy, P.C., Jeffries, K.M., Fangue, N.A., Connon, R.E. 2020. Differential regulation of select
osmoregulatory genes and Na+/K+-ATPase paralogs may contribute to population
differences in salinity tolerance in semi-anadromous fish. *Comp Biochem Physiol A*.
110584. https://doi.org/10.1016/j.cbpa.2019.110584

Muschick, M., Barluenga, M., Salzburger, W., Meyer, A. 2011. Adaptive phenotypic plasticity in
the Midas cichlid fish pharyngeal jaw and its relevance in adaptive radiation. *BMC Evol*
Biol. 11, 116. https://doi.org/10.1186/1471-2148-11-116

Ogden, R., Gharbi, K., Mugue, N., Martinsohn, J., Senn, H., Davey, J.W., Pourkazemi, M.,
McEwing, R., Eland, C., Vidotto, M., Sergeev, A., Congiu, L. 2013. Sturgeon

conservation genomics: SNP discovery and validation using RAD sequencing. *Mol Ecol*. 22, 3112-23. https://doi.org/10.1111/mec.12234.

Oomen, R. A., Hutchings, J. A. 2017. Transcriptomic responses to environmental change in fishes: insights from RNA sequencing. *FACETS* 2, 610–641. https://doi.org/10.1139/facets-2017-0015.

Paital, B., Panda, S.K., Hati, A.K., Mohanty, B., Mohapatra, M.K., Kanungo, S., Chainy, G.B.N. 2016. Longevity of animals under reactive oxygen species stress and disease susceptibility due to global warming. *World J Biol Chem*. 7, 110-127. https://doi.org/10.4331/wjbc.v7.i1.110

Palmer, M.A., Liermann, C.A.R., Nilsson, C., Flörke, M., Alcamo, J., Lake, P.S., Bond, N. 2008. Climate change and the world's river basins: anticipating management options. *Front Ecol Environ*. 6, 81-89. https://doi.org/10.1890/060148

Peake, S., Beamish, F.W.H., McKinley, R.S., Scruton, D.A., Katopodis, C. 1997. Relating swimming performance of lake sturgeon, Acipenser fulvescens, to fishway design. *Can J Fish Aquat Sci*. 1366, 1361–1366. https://doi.org/10.1139/f97-039

Pelseter, B., Egg, M. 2018. Hypoxia-inducible transcription factors in fish: expression, function and interconnection with the circadian clock. *J Exp Biol*. 4, 13. https://doi.org/10.1242/jeb.163709

Penny, F.M., Bugg, W.S., Kieffer, J.D., Jeffries, K.M., Pavey, S.A. 2023. Atlantic sturgeon and shortnose sturgeon exhibit highly divergent transcriptomic responses to acute heat stress. *Comp Bio Phys D*. 45, 101058.

Penman, R. 2021. The effects of temperature acclimation and heating rate on the thermal tolerance of juvenile white sturgeon (*Acipenser transmontanus*). *Masters thesis*. University of British Columbia.

Pertea, M., Pertea, G., Antonescu, C. et al. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat Biotechnol*. 33, 290–295. https://doi.org/10.1038/nbt.3122.

Portner, H.O., Bock, C., Mark, F.C. 2017. Oxygen- and capacity-limited thermal tolerance: bridging ecology and physiology.*J Exp Biol*. 220, 2685-2696. https://doi.org/10.1242/jeb.134585

Portner, H.O., Lanning, G. 2009. Chapter 4 Oxygen and Capacity Limited Thermal Tolerance. Fish Physiology. 27, 143-191.

Qian, X., Ba, Y., Zhuang, Q., Zhong, G. 2014. RNA-Seq technology and its application in fish transcriptomics. *OMICS J Integr Biol*. 18, 98-110. https://doi.org/10.1089/omi.2013.0110

Qin, Z., Raoof, M., Chen, Y., Sumi, Y., Sursal, T., Junger, W., Brohi, W., Itagaki, K., Hauser, C.J. 2010. Circulating mitochondrial DAMPS cause inflammatory responses to injury. *Nature*. 464, 104-107. https://doi.org/10.1038/nature08780

R Core Team. 2022. **R**: A language and environment for statistical computing. **R** Foundation for
Statistical Computing, Vienna, Austria. URL http://www.**R**-project.org/

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law., C.W., Shi, W., Smyth, G.K. 2015. limma
powers differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res. 43, e47. https://doi.org/10.1093/nar/gkv007

Rivera-Ingraham, G.A., Lignot, J.H. 2017. Osmoregulation, bioenergetics and oxidative stress in
coastal marine invertebgrates: raising the questions for future research. *J Exp Biol*. 220,
1749-1760. https://doi.org/10.1242/jeb.135624

Robertson, M.D., Gao, J., Regular, P.M., Morgan, M.J., Zhang, F. 2021. Lagged recovery of fish
spatial distributions following a cold-water perturbation. *Sci Rep*. 11, 9513.
https://doi.org/10.1038/s41598-021-89066-x

Robinson, M.D., McCarthy, D.J., Smyth, G.K. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*. 26, 139-
140. https://doi.org/10.1093/bioinformatics/btp616

Saari, G.N., Wang, Z., Brooks, B.W. 2018. Revisiting inland hypoxia: diverse exceedances of
dissolved oxygen thresholds for freshwater aquatic life. *Environ Sci Pollut Res*. 25, 3139-
3150. https://doi.org/10.1007/s11356-017-8908-6

Schade, F.M., Shama, L.N.S., Wegner, K.M. 2014. Impact of thermal stress on evolutionary
trajectories of pathogen resistance in three-spined stickleback. *BMC Evol Biol*. 14, 164.
https://doi.org/10.1186/s12862-014-0164-5

Scheiner, S.M., Barfield, M., Holt, R.D. 2019. The genetics of phenotypic plasticity. XVII.
Response to climate change. *Evol Appl*. 13, 388-399.

Schindler, D.W. 2001. The cumulative effect of climate change and multiple stressors. *Can J
Fish Aquatic Sci*. 2958, 18–29. https://doi.org/10.1139/f00-179

Schlichting, C.D., Smith, H. 2002. Phenotypic plasticity: linking molecular mechanisms with
evolutionary outcomes. *Evol Ecol*. 16, 189-211.
https://doi.org/10.1023/A:1019624425971

Seebacher, F., White, C.R., Franklin, C.E. 2015. Physiological plasticity increases resilience of
ectothermic animals to climate change. *Nat Clim Change*. 5, 61-66.
https://doi.org/10.1038/nclimate2457

Shiina, T., Shimizu, Y. 2020. Temperature-Dependent Alternative Splicing of Precursor mRNAs
and Its Biological Significance: A Review Focused on Post-Transcriptional Regulation of
a Cold Shock Protein Gene in Hibernating Mammals. *Int J Mol Sci*. 14, 7599.
https://doi.org/ 10.3390/ijms21207599

Smith, S., Bernatchez, L., Beheregaray, L.B. 2013. RNA-seq analysis reveals extensive
transcriptional plasticity to temperature stress in a freshwater fish species. *BMC Genom*.
14, 1-12. https://doi.org/10.1186/1471-2164-14-375

Somero, G.N. 2010. The physiology of climate change: how potentials for acclimatization and
genetic adaption will determine 'winners' and 'losers'. *J Exp Biol*. 213, 912-920.
https://doi.org/10.1242/jeb.037473

Somero, G.N. 2018. RNA thermosensors: how might animals exploit their regulatory potential? *J*
Exp Biol. 22, 221. https://doi.org/10.1242/jeb.162842

Steward, R.A., de Jong, M.A., Oostra, V., Wheat, C.W. 2022. Alternative splicing in seasonal
plasticity and the potential for adaptation to environmental change. *Nat Commun*. 13,
755. https://doi.org/10.1038/s41467-022-28306-8

Storey, K.B., Storey, J.M. 2011. Heat shock proteins and hypometabolism: adaptive strategy for
proteome preservation. *Res Rep Biol*. 2, 57-68. https://doi.org./10.2147/RRB.S13351

Su, G., Logez, M., Xu, J., Tao, S., Villéger, S., Brosse, S. 2021. Human impacts on global
freshwater fish biodiversity. *Science*. 371, 835-838.
https://doi.org/10.1126/science.abd3369

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists of
Gene Ontology terms. *PLoS One*. 6, e21800.
https://doi.org/10.1371/journal.pone.0021800

Sutton, R.J., Deas, M.L., Tanaka, S.K., Soto, T., Corum, A. 2007. Salmonid observations at the
Klamath River thermal refuge under various hydrological and meteorological conditions.
River Res Appl. 23, 775-785. https://doi.org/10.1002/rra.1026

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
Doncheva, N.T., Morris, J.H., Bork, P., Jensen, L.J., von Mering, C. 2018. STRING v11:
protein-protein association networks with increased coverage, supporting functional
discovery in genome-wide experimental datasets. *Nucleic Acids Res*. 8, D607-D613.
https://doi.org/10.1093/nar/gky1131

Tan, S., Wang, W., Tian, C., Niu, D., Zhou, T., Jin, Y., Yang, Y., Gao, D., Dunham, R., Liu, Z.
2019a. Heat stress induced alternative splicing in catfish as determined by transcriptome
analysis. *Comp Biochem Physiol*. 29. 166–172. https://doi.org/10.1016/j.cbd.2018.11.008

Tan, S., Wang, W., Tian, C., Niu, D., Zhou, T., Yang, Y., Gao, D., Liu, Z. 2019b. Post-
transcriptional regulation through alternative splicing after infection with *Flavobacterium*
columnare in channel catfish (*Ictaluru punctatus*). *Fish Shellfish Immunol*. 91, 188-193.
https://doi.org/10.1016/j.fsi.2019.05.008

Tan, S., Wang, W., Zhong, X., Tian, C., Niu, D., Bao, L., Zhou, T., Jin, Y., Yang, Y., Yuan, Z.,
Gao, D., Dunham, R., Liu, Z. 2018. Increased alternative splicing as a host response to
edwardsiella ictalurid infection in catfish. *Mar Biotechnol*.
https://doi.org/10.1007/s10126-018-9844-2

Taylor, S.J., Shalloway, D. 1996. Src and the control of cell division. *Bioessays*. 18, 9-11.
https://doi.org/10.1002/bies.950180105

Thornton, P. K., Ericksen, P. J., Herrero, M., Challinor, A.J. 2014. Climate variability and vulnerability to climate change: a review. *Glob Chang Biol*. 20, 3313-3328. https://doi.org/10.1111/gcb.12581

Thorstensen, M.J., Baerwald, M.R., Jeffries, K.M. 2021. RNA sequencing describes both population structure and plasticity-selection dynamics in a non-model fish. *BMC Genom*. 22, 1-12. https://doi.org/10.1186/s12864-021-07592-4

Thorstensen, M.J., Jeffrey, J.D., Treberg, J.R., Watkinson, D.A., Enders, E.C., Jeffries, K.M. 2020. Genomic signals found using RNA sequencing show signatures of selection and subtle population differentiation in walleye (*Sander vitreus*) in a large freshwater ecosystem. *Ecol Evol*. 10, 7173-7188. https://doi.org/10.1002/ece3.6418

Thorstensen, M. J., Weinrauch, A. M., Bugg, W. S., Jeffries, K. M., Anderson, W.G. 2022a. Tissue-specific transcriptomes reveal mechanisms of microbiome regulation in an ancient fish. bioRxiv, 2022-10. https://doi.org/10.1101/2022.10.12.511976

Thorstensen, M.J., Turko, A.J., Heath, D.D., Jeffries, K.M., Pitcher, T.E. 2022b. Acute thermal stress elicits interactions between gene expression and alternative splicing in a fish of conservation concern. *J Exp Biol*. 15, jeb244162.

Todgham, A.E., Stillman, J.H. 2013. Physiological Responses to Shifts in Multiple Environmental Stressors: Relevance in a Changing World. *Integrative and Comparative Biology*. 53, 539-544.

Tovar, J., Leon-Avila, G., Sanchez, L.B., et al., 2003. Mitochondrial remnant organelles of Giardia function in iron-sulphur protein maturation. *Nature*. 13, 172-176. https://doi.org/10.1038/nature01945

Trifonov, V.A., Romanenko, S.S., Beklemisheva, V.R., Biltueva, L.S., Makunin, A.I., Lemskaya, N.A., Kulemzina, A.I., Stanyon, R., Graphodatsky, A.S. 2016. Evolutionary plasticity of acipenseriform genomes. *Chromosoma*. 125, 661-668. https://doi.org/10.1007/s00412-016-0609-2

Vallin, J., Grantham, J. 2019. The role of the molecular chaperone CCT in protein folding and mediation of cytoskeleton-associated processes: implications for cancer cell biology. *Cell Stress Chaperones*. 24, 17-27. https://doi.org/10.1007/s12192-018-0949-3

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine, A., ... & DePristo, M. A. (2013). From FastQ data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. Current protocols in bioinformatics, 43(1), 11-10.

Van de Peer, Y., Mizrachi, E. & Marchal, K. 2017. The evolutionary significance of polyploidy. *Nat Rev Genet*. 18, 411–424. https://doi.org/10.1038/nrg.2017.26

Van der Lee, A.S., Koops, M.A. 2021. Recovery Potential Modelling of Lake Sturgeon (*Acipenser fulvescens*) in Canada. DFO Can. Sci. Advis. Sec. Res. Doc. 2021/025. iv + 48 p.

Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D.G., McCann, K.S., Savage,
V., Tunney, T.D., O'Connor, M.I. 2014. Increased temperature variation poses a greater
risk to species than climate warming. *Proc Royal Soc B*. 281, 20132612.
https://doi.org/10.1098/rspb.2013.2612

Verhille, C.E., Poletto, J.B., Cocherell, D.E., DeCourten, B., Baird, S., Cech, Jr, J.J., Fangue,
N.A. 2014. Larval green and white sturgeon swimming performance in relation to water-
diversion flows. *Conserv Physiol*. 2, cou031. https://doi.org/10.1093/conphys/cou031

Verta, J.P., Jacobs, A. 2021. The role of alternative splicing in adaptation and evolution. *Trends
Ecol Evol*. 37, 4. https://doi.org/10.1016/j.tree.2021.11.010

Vincent, L.A., Zhang, X., Brown, R.D., Feng, Y., Mekis, E., Milewska, E.J., Wan, H., Wang,
X.L. 2015. Observed trends in Canada's climate and influence of low-frequency
variability modes. *J Climate* 28: 4545-4560. https://doi.org/10.1175/JCLI-D-14-00697.1

Vollset, K.W., Lennox, R.J., Davidsen, J.G., Eldøy, Isaksen, T.E., Madhun, A., Karlsson, S.,
Miller, K.M. 2020. Wild salmonids are running the gauntlet of pathogens and climate as
fish farms expand northwards. *ICES J Mar Sci*. https://doi.org/10.1093/icesjms/fsaa138

Vornanen M, Hassinen M, Koskinen H, Krasnov A. 2005. Steady-state effects of temperature
acclimation on the transcriptome of the rainbow trout heart. *Am J Physiol Regul Integr
Comp Physiol*. 289, R1177–R1184. https://doi.org/10.1152/ajpregu.00157.2005

Warkentin, L., Parken, C.K., Bailey, R., Moore, J.W. 2022. Low summer river flows associated
with low productivity of Chinook salmon in watershed with shifting hydrology. *Ecolog
Solut Evid*. 3, e12124. https://doi.org/10.1002/2688-8319.12124.

Webb, W. 1986. Kinematics of lake sturgeon, *Acipenser fulvescens*, at cruising speeds. *Can J
Zool*. 64, 2137–2141. https://doi.org/10.1139/z86-328

Wei, J., Zang, S., Xu, M., Zheng, Q., Chen, X. and Qin, Q., 2017. TRAF6 is a critical factor in
fish immune response to virus infection. *Fish Shellfish Immunol*. 60, pp.6-12.
https://doi.org/10.1016/j.fsi.2016.11.008

Wellband, K. W., Heath, D. D. 2017. Plasticity in gene transcription explains the differential
performance of two invasive fish species. *Evol Appl*. 10, 563–576. https://doi.org/
10.1111/eva.12463

White, S.L., Kline, B.C., Hitt, N.P., Wagner, T. 2019. Individual behaviour and resource use of
thermally stress brook trout *Salveliunus fontinalis* portend the conservation potential of
thermal refugia. *J Fish Biol*. 95, 1061-1071. https://doi.org/10.1111/jfb.14099

Whitehead, A., Roach, J.L., Zhang, S., Galvez, F. 2012. Salinity-and population-dependent
genome regulatory response during osmotic acclimation in the killifish (*Fundulus
heteroclitus*) gill. *J Exp Biol*. 215, 1293-1305. https://doi.org/10.1242/jeb.062075

WWF. 2022. Sturgeon slipping towards extinction: Chances of survival dimming for world's
most threatened group of species. See https://wwfcee.org/news/sturgeon-slipping-
towards-extinction-chances-of-survival-dimming-for-worlds-most-threatened-group-of-
species.

Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann
A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., Jeon, M., Ma'ayan, A. 2021.
Gene set knowledge discovery with Enrichr. *Curr Protoc*. 1, e90.
https://doi.org/10.1002/cpz1.90

Yoon, G.R., Earhart, M., Wang, Y., Suh, M., Anderson, G.W. 2021. Effects of temperature and
food availability on liver fatty acid composition and plasma cortisol concentration in age-
0 lake sturgeon: Support for homeoviscous adaptation. *Comp Biochem and Phys.* 261,
111056. https://doi.org/10.1016/j.cbpa.2021.111056

Zhang X, Flato G, Kirchmeier-Young M, Vincent L, Wan H, Wang X, Rong R, Fyfe J, Li G,
Kharin W. 2019. Changes in Temperatures and Precipitation Across Canada; Chapter 4 *in*
Bush E and Lemmens DS (Eds.). *Canada's Changing Climate Report*. *Government of*
Canada, Ottawa, Ontario, p. 112-193.

Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-
performance computing toolset for relatedness and principal component analysis of SNP
data. Bioinformatics, 28(24), 3326-3328.

Zhou, G., Xia, J. 2018. OmicsNet: a web-based tool for creation and visual analysis of biological
networks in 3D space. *Nucleic Acids Res*. 46, W514-W522.
https://doi.org/10.1093/nar/gky510

Zhou, G., Xia, J. 2019. Using OmicsNet for networking integration and 3D visualization. *Curr*
Protoc Bioinform. 65, e69. https://doi.org/10.1002/cpbi.69

1489