1 **Title:**
2 M-Risk: A framework for assessing global fisheries management efficacy of sharks, rays, and
3 chimaeras
4
5 **Alternate titles:**
6   1. M-Risk: An updated framework to assess efficacy of fisheries management of sharks,
7   rays, and chimaeras based on intrinsic sensitivity
8   2. Developing a framework for rapid assessment of global fisheries management of
9   sharks, rays, and chimaeras
10
11 **Running title:**
12 M-Risk shark fisheries assessments
13

Authors:
C. Samantha Sherman* [1,2]
Glenn Sant [2,3]
Colin A. Simpfendorfer [4]
Eric D. Digel [1]
Patrick Zubick [1]
Grant Johnson [5]
Michael Usher [5]
Nicholas K. Dulvy [1]
23
24 **Corresponding author:**
25 C. Samantha Sherman
26 E-mail: sammsherman27@gmail.com
27
Affiliations:
1. Earth to Oceans Research Group, Department of Biological Sciences, Simon Fraser
University, Burnaby, British Columbia, Canada
2. TRAFFIC International, Cambridge, U.K.
3. Australian National Centre for Ocean Resources and Security, University of
Wollongong, NSW, Australia
4. Institute of Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania,
Australia
5. Department of Industry, Tourism and Trade, Fisheries Branch, Northern Territory
Government, Darwin, Northern Territory, Australia

38
39
40
41

42
43
44

ABSTRACT

Fisheries management is essential to guarantee sustainable capture of target species and

avoid undesirable declines of incidentally captured species. A key challenge is halting and

reversing declines of shark and ray species, and specifically assessing the degree to which

management is sufficient to avoid declines in relatively data-poor fisheries. While ecological

risk analyses focus on intrinsic 'productivity' and extrinsic 'susceptibility', one would ideally

consider the influence of 'fisheries management'. Currently, there is no single management

evaluation that can be applied to a combination of fishery types at the scale of individual

country or Regional Fisheries Management Organisations (RFMOs). Here, we outline a

management risk (M-Risk) framework for sharks, rays, and chimaeras used to evaluate

species' risk to overfishing resulting from ineffective management. We illustrate our

approach with application to one country (Ecuador) and RFMO (Inter-American Tropical

Tuna Commission) and illustrate the variation in scores among species. We found that while

both management units assessed had similar overall scores, the scores for individual

attributes varied. Ecuador scored higher in reporting-related attributes, while the IATTC

scored higher in attributes related to data collection and use. We evaluated whether

management of individual species was sufficient for their relative sensitivity by combining

the management risk score for each species with their intrinsic sensitivity to determine a

final M-Risk score. This framework can be applied to determine which species face the

greatest risk of overfishing and be used by fisheries managers to identify effective

management policies by replicating regulations from countries with lower risk scores.

KEYWORDS

Data-poor fisheries, ecological risk assessment, elasmobranch, marine conservation,

resource management, socio-ecological resilience

1. INTRODUCTION

Catch in marine fisheries globally has decreased since the mid-1990s despite increasing effort (FAO, 2020; Pauly, Zeller, & Palomares, 2021; Rousseau, Watson, Blanchard, & Fulton, 2019). Ensuring long-term sustainable fisheries requires management not only of target species but also of all other species affected by the fishery (Hilborn et al., 2003). However, halting biodiversity loss and ecosystem management ideally requires knowledge of the status and fishing mortality associated with all species taken in the fishery. Globally, at least 13,060 different species are caught (FAO, 2021), leaving over 97% of species without a stock assessment – there are only 957 stock assessments for 360 unique species (RAM Legacy Stock Assessment Database, 2021). For such data-rich stock assessed populations and species, we know their levels of exploitation and their sustainable fishing mortality levels (RAM Legacy Stock Assessment Database, 2021; Ricard, Minto, Jensen, & Baum, 2012). But for data-poor species, how do we determine their status? Much of our understanding on data-poor species' catch is based on landings data reported to the Food and Agriculture Organization of the United Nations (FAO)(Froese, Zeller, Kleisner, & Pauly, 2012). Data-poor species comprise >80% of global catch and almost two-thirds of species are overexploited (Costello et al., 2012; Guan, Chen, Boenish, Jin, & Shan, 2020). However, landings data are not consistent or reliable globally, particularly in countries with many landing sites and/or high levels of artisanal catch, particularly for sharks and their relatives (Khan et al., 2020; Okes & Sant, 2022; Ruano-Chamorro, Subida, & Fernández, 2017). Additionally, the catches of data-poor species, if characterised, are often reported as aggregates; grouped together by genus, family, or higher, giving little information on the catch of individual species (Cashion, Bailly, & Pauly, 2019; FAO, 2019). The portfolio effects of these aggregate catch statistics mask serial depletions, cryptic declines, and local extinctions (Dulvy, Metcalfe, Glanville, Pawson, & Reynolds, 2000; Lawson et al., 2020; Schindler et al., 2010). Many methods have been developed to assess data-poor species and stocks, which require, *inter alia*, different data inputs such as catches (Anderson, Branch, Ricard, & Lotze, 2012), length compositions (Cope & Punt, 2009; Froese, 2004), age-at-maturity (Brooks, Powers, & Cortés, 2010; Cope, 2013), gear selectivity (Le Quesne & Jennings, 2012). However, all have major assumptions and biases associated with the outputs (Chrysafi & Kuparinen, 2015). There remains a need for a rapid risk assessment technique, when detailed stock assessments are not possible, at the scale of countries and RFMOs.

When assessing the risk of a species within a fishery, the terminology used throughout the

risk assessment literature is inconsistent. Therefore, we have used the most common terms

and defined them where appropriate. Here, we use the framing of Vulnerability as derived

from the social science, hazard assessment, and climate risk literature (Turner II et al.,

2003), and increasingly used in biological risk assessment (Allison et al., 2009; Williams,

Shoo, Isaac, Hoffmann, & Langham, 2008). Vulnerability is typically considered to be the

interaction of intrinsic sensitivity (biological traits that inform extinction risk) with exposure

(the overlap between a threat and the species' range) to a threatening process (i.e., fishing

or climate change). Broadly, vulnerability can be considered to be the inverse of ecological

resilience (Allison et al., 2009). Taken together, the Potential Impact (sensitivity x exposure)

can be offset either by phenotypical plasticity or genotypic evolution when considering a

species, or through building Adaptive Capacity of the human system, which can be

strengthened through management or disaster planning and preparedness (Allen Consulting

Group, 2005; Allison et al., 2009; Dulvy et al., 2011). For example, fishers in Kenya increased

their Adaptive Capacity to climate change by increasing community infrastructure and

access to credit (Cinner et al., 2015). Ecological risk assessments can take several different

forms, however, all measure similar aspects of risk such that a species' vulnerability (V) is a

function of their intrinsic life history sensitivity (Intrinsic Sensitivity), Exposure to fishing

pressure, say as inferred from the spatial and depth overlap with the species' distribution,

and the species' catchability in the fishing gears (Hobday et al., 2011; Walker et al., 2021).

All this Potential Impact can be offset by Adaptive Capacity:

$$\text{Equation 1:} \quad Vulnerability = f(\frac{Intrinsic\ Sensitivity * Exposure}{Adaptive\ Capacity})$$

This generic risk assessment framework has been pruned down to focus only on the

Potential Impact, specifically the interplay of sensitivity and exposure. This is most

commonly treated by Productivity Susceptibility Assessments (PSA), a type of ecological risk

analysis which has been typically applied to single fishery to compare the relative risk of the

full range incidentally captured species (Fletcher, 2005; Hobday et al., 2011; Micheli, De Leo,

Butner, Martone, & Shester, 2014) (**Figure 1**). This approach can be applied to data-poor

fisheries and in highly diverse systems with taxonomically diverse species in the incidental

catch, such as in an Australian Prawn Trawl Fishery (Astles et al., 2006; Stobutzki, Miller, &

Brewer, 2001).

As with all risk assessments, there are trade-offs to achieve a result that best informs the

goals of the assessment. Other risk frameworks, like PSAs, tend to focus on a specific

assessment unit comprised of a single fishery, set of species, or gear type, or combination

(Astles et al., 2006; Cortés, Brooks, & Shertzer, 2015; Zhou, Milton, & Fry, 2012). PSA

customization allows for a deeper, more nuanced assessment within a single assessment

unit. However, specificity of a PSA makes comparisons difficult to other PSAs with different

criteria or scoring. Thus, the need for a method that can be used across all fishery types

including different gear types, target species, and data availability at the scale of countries

and RFMOs. Similarly, while a risk assessment can be used to rank the species at risk within

a fishery, it does not consider the degree to which the Potential Impact is or can be offset by

the existence of some form of management (Turner II et al., 2003) (**Figure 1**).

To incorporate management into risk assessments, a rapid management risk assessment

(M-Risk) was proposed for shark species and tested for species with high intrinsic sensitivity,

many of which are CITES listed (Lack, Sant, Burgener, & Okes, 2014). The results showed

variation in management efficacy despite protections that should have resulted from

compliance with CITES listings. While the original risk framework was fit for the purpose of

threatened species, it was not necessarily optimised for species caught and traded in high

volumes. M-Risk considers that vulnerability (V) is a function of a species' intrinsic sensitivity

(IS), exposure to fishing pressure (E), and Management, substituted for Adaptive Capacity

from equation 1:

$$\text{Equation 2: } Vulnerability = f(\frac{Intrinsic\ Sensitivity * Exposure}{Management})$$

We note that while management does not influence intrinsic susceptibility *per se* it does

influence the interaction of intrinsic sensitivity and exposure *inter alia* by managing the

availability, encounterability, selectivity, and post-release mortality of species through

various technical approaches (Hobday et al., 2011).

Sharks, rays, and chimaeras (class: Chondrichthyes; >1,199 species, hereafter "sharks and

rays") present a challenge for fisheries managers as they are frequently retained as

secondary catch or discarded as incidental catch (Stevens, Bonfil, Dulvy, & Walker, 2000).

Overfishing, both targeted and incidental, has caused populations of sharks and rays to

decline dramatically over the past 50 years, leading to a high rate of elevated extinction risk

(Dulvy et al., 2021; Pacoureau et al., 2021). As they are commonly considered unavoidable

incidental or secondary catch, sharks and rays are frequently treated as unimportant or

unavoidable in a management context. Therefore, sharks and rays are an ideal candidate

group for an M-Risk assessment. Many sharks and rays are longer-lived species, with low

intrinsic rates of population increase, and cannot be fished at the same rate as most

teleosts, despite often being caught alongside them (Brander, 1981; Musick, Burgess,

Cailliet, Camhi, & Fordham, 2000; Myers & Worm, 2005; Pardo, Cooper, Reynolds, & Dulvy,

2018; Stevens, Walker, & Simpfendorfer, 1997). Fisheries management of sharks and rays is

further complicated because of the diversity of species, gears used, jurisdictions in which

they are caught (Dulvy et al., 2017; Simpfendorfer & Dulvy, 2017), and their complex

migration patterns that transit the waters of multiple countries, which increases their

exposure in multiple fisheries and cumulative impacts of being caught throughout their

migration routes (Dragičević, Dulčić, & Capapé, 2009; Heupel et al., 2015; Sellas et al., 2015).

Despite the high intrinsic sensitivity of many species, sustainable shark and ray fisheries are

possible if assessment and management is adequate. There are 39 sustainably fished

populations of 33 species of sharks and rays around the globe (Simpfendorfer & Dulvy,

2017) and signs of recovery in US and EU managed populations (Amelot et al., 2021;

Peterson et al., 2017). Sustainable shark and ray fisheries all have common characteristics

that distinguish them from unsustainable shark and ray fisheries. These centre on fishing

mortality to sustainable levels through limits on catch and/ or effort, supported by robust

legislation, well-enforced regulations, and science-based advisory processes (Simpfendorfer

& Dulvy, 2017; Woodhams, Peddemors, Braccini, Victorian Fisheries Authority, & Lyle,

2021).

Here, we provide a revised set of attributes based on the original M-Risk framework (Lack et

al. 2014), for application to any shark or ray species in any fishery to rapidly and objectively

assess management efficacy in a comparable consistent manner at country and RFMO

scales (**Figure 1**). First, we illustrate the revised M-Risk framework with two case studies:

one for all species in a country (Ecuador, 29 species) and one for 24 species in a Regional

Fisheries Management Organisation (RFMO) – the Inter-American Tropical Tuna

Commission (IATTC). Second, once assessments were completed, management scores for

each species were combined with their intrinsic risk scores to attain a final M-Risk score per

species in each management unit. Third, we discuss the methodology and initial

assessments with the goal of completing assessments for a larger suite of management

units globally, hence species mentioned includes some not assessed in this paper. These

scores are intended for two anticipated uses: (1) by fisheries managers so that

improvements can be made specifically for species that are undermanaged relative to their

intrinsic sensitivity or for overall fisheries improvements (risk management) and (2) for

comparative analysis of the state of the world's shark and ray fisheries.

2. DEVELOPING THE NEW M-RISK FRAMEWORK

2.1 Unit of Assessment – country and RFMO

When considering exploited marine species and their management, there are two levels at

which an assessment can be completed: stock or fishery. First, a stock is usually a political

construct that consists of a demographically isolated portion of the global population that is

often managed as a single unit (Begg & Waldman, 1999). Where a single stock is

widespread, the fisheries may be managed separately by different jurisdictions. This can

complicate the risk assessment of the stock, as it may not be clear which management unit

is responsible for the species' status if it is not uniformly co-managed across jurisdictions,

and the assessment must factor in the effects of multiple sectors and the cumulative

impacts on the status of the stock (Urquhart, Acott, Symes, & Zhao, 2014). For example, the

Winter Skate (*Leucoraja ocellata*) occurs in both Canadian and USA waters; however, the

population trend in the USA is increasing, while the population trend in Canadian waters is

decreasing (Kulka et al., 2020). The discrepancy may be due to differing management focus

and effectiveness of the fisheries in each country (Kulka et al., 2020). Second, in our case, a

fishery is defined by its management such that each separate fishery is managed by a single

jurisdiction that regulates the laws of operation within the spatial extent of the fishery.

We are applying M-Risk at two scales (country and RFMO), hereafter referred to as

"Management Units". Within a single country (Ecuador, here), we selected a representative

fishery based on which one had the highest level of catch (retained and discarded) with the

species under assessment. Each fishery will be assessed as a single unit, not considering if

the species is caught in other fisheries within or outside of the management unit, or the

cumulative impacts of multiple fisheries contributing to fishing mortality of the species. High

seas fishing was not considered when assessing countries, only fishing occurring within their

respective Exclusive Economic Zones (EEZs), as high seas management should be picked up

in RFMO assessments. Vessels from member countries of each RFMO are obligated to

adhere to the RFMO regulations when operating in those fishery grounds. Countries may

apply stricter regulations to their flagged vessels fishing within RFMO grounds. However, we

are applying our criteria only to legislation that is applicable for **all** member countries of the

RFMO, as this is the minimum standard in those fishing grounds.

2.2 Species Selection

Sharks and rays are commonly caught and traded globally (Dent & Clarke, 2015). As there

are currently over 1,200 described species (Ebert, Dando, & Fowler, 2021; Last et al., 2016),

we curated a list consisting of the most frequently traded species worldwide, based on four

sources: (i) catches reported to the Food and Agriculture Organization of the United Nations

(FAO) (FAO, 2019), (ii) reconstructed catches from the Sea Around Us Project (Pauly & Zeller,

2016), (iii) species listed on the Appendices of the Convention on International Trade in

Endangered Species of Wild Fauna and Flora (CITES) (https://cites.org), and (iv) species

listed on the Appendices of the Convention on the Conservation of Migratory Species of

Wild Animals (CMS) (https://www.cms.int). Additionally, we included species or groups that

are caught in high abundances but usually identified at a higher taxonomic level, such as

cowtail rays (*Pastinachus* spp.) in the Indo-Pacific. The final list for the full M-Risk project

included 69 sharks identified to species level apart from one genus (*Etmopterus*; n = 38), 27

rays identified to either species level or a ray group (e.g., eagle rays) (16 species; 11 groups,

n = 80), and 6 ghost shark species for a total of 102 species or groups (**Data S1**). The ray

groups included different levels of taxonomic resolution for species that were difficult to

distinguish, have similar life history characteristics, distribution, and human use, or are

taxonomically unresolved or may comprise a species complex (e.g., maskrays – *Neotrygon*

spp.). The number of species assessed in this paper was lower, at 36 species (30 sharks, 2

ray species, and 4 ray groups). However, the larger suite of 102 species is referred to here

because intrinsic risk scores of the 36 species considered here are expressed relative to the

entire 102 species of the project. Assessments were completed for each species separately,

except for the groups, for which a single assessment was completed for the whole group.

The combined assessments were necessary, as there are very limited species-specific data

or management system details within the groups. All species that have a distribution

overlapping with the management unit's spatial grounds are assessed for that management

unit. Final M-Risk scores were calculated at the species level, as intrinsic sensitivity of

species within each group differed.

2.3 Management Assessment Framework

We devised 21 measurable attributes to assess the degree to which management is

adequate to prevent sharks and rays from being overexploited. These attributes were

chosen based on regulations that (1) enabled understanding, and (2) curbed fishing

mortality on the focal shark and ray species and was informed by previous work on

sustainable shark fisheries (Davidson, Krawchuk, & Dulvy, 2015; FAO, 1999; Melnychuk,

Peterson, Elliott, & Hilborn, 2017). Some attributes were considered but could not be

included due to data paucity. For example, total catch of each species is not available in

most fisheries and, therefore, was not included as an attribute at present but could be

considered as more of these data become available.

Twenty-one attributes were created comprised of three common classes and two classes

related to the spatial unit of analysis (**Table 1**):

(i) the management system (5 attributes),

(ii) fishing practices and catch (5 attributes),

(iii) compliance and enforcement (5 attributes),

and an additional category considered the management unit and how it relates to other

management units which consisted of attributes either specific to:

(iv) country (4 attributes), OR

(v) RFMO (2 attributes).

Therefore, countries were scored for 19 of the attributes and RFMOs are scored for 17 of

the attributes. These numbers were different as there were more aspects of a country that

could be assessed. Next, we unpack each of the five attribute classes.

2.3.1 *Management System*

The management system consists of a regulatory body, fishery permits, and assessments to

understand the potential risk the fishery poses. Without such foundational management in

place, it is unlikely the fishery will have any more sophisticated regulations or any capacity

to enforce regulations in place. These attributes are intended to increase our understanding

of the capability of management for improvement. Having basic management, such as a

permitting system, enables fisheries managers to have records of all vessels operating in the

fishery, which can enable the setting of limits on catches or effort. For management units

that receive low scores in the 'management system' attributes, we would recommend the

government allocates resources to strengthen the capacity of the fishery to manage the

resource as a first step to introducing intricate regulations that are more likely to curb

fishing mortality and increase sustainability.

2.3.2 *Fishing Practices and Catch*

Fishing practices and catch include landing limits, shark finning, post-release survival, and

closures. Management of at-sea fishing operations is difficult due to the size of the area in

which fishing occurs and the costs associated with patrolling large spaces (Rowlands, Brown,

Soule, Boluda, & Rogers, 2019). Ideally, fisheries managers would know exactly the number

of each species caught and their fate post-release (if released), in addition to having spatial

and/or seasonal closures in place for at-risk species at sensitive times or locations. Simply,

having regulations in place that affect at-sea operations but can be measured or enforced

upon landing are the most effective, as these do not require boarding of vessels at-sea. For

example, the requirement that sharks are landed with fins naturally attached is something

that can be checked upon landing the catch (Fowler & Séret, 2010). Similarly, the existence

of spatial and temporal closures can be monitored through use of Vessel Monitoring

Systems (VMS) that can be checked from land (Enguehard, Devillers, & Hoeber, 2013). For

management units with low scores in the 'fishing practices and catch' attributes, we would

recommend the inclusion of regulations that decrease fishing mortality on intrinsically

sensitive species, like sharks and rays.

2.3.3 Compliance, Monitoring, and Enforcement

Compliance, monitoring, and enforcement consists of reporting catch, ensuring reports are

valid, having enforcement in place for violations, and monitoring of illegal, unregulated, and

unreported fishing (IUU). Compliance, meaning operating within the established legislation,

often requires fisher agreement. Fishers are more likely to comply with legislation (1) when

they understand regulations are in place for their own interest by increasing sustainability,

(2) through feeling obligated to comply, and (3) through enforcement measures that may

decrease their profitability if non-compliance is discovered (Hønneland, 1999). Additionally,

fishers with a better relationship and trust with the management authority are more likely

to comply with regulations (Hauck, 2008). In an ideal world, fishers would have 100%

compliance with the regulations, however, this is not realistic, and therefore, must be

monitored and enforced (Price et al., 2016). The attributes within this class were designed

to assess how the quantity and composition of the catch is validated and what measures are

in place to ensure there is high compliance within the fishery. Similarly, we include an

attribute considering how IUU fishing is accounted for through management measures. As

IUU fishing has been estimated to be up to 26 million tonnes per year of the global catch,

this can have a significant impact on fishing mortality and must be accounted for in

assessing and managing the fishery (Agnew et al., 2009). The attributes within this group

require more nuanced regulations, therefore, management units with higher scores in this

category are expected to have higher scores in the previous categories as well. For

management units with lower scores in the 'compliance, monitoring, and enforcement'

attributes, we would recommend allocating funding to increase fishery officer coverage,

including at landing ports to observe unloading and in office to monitor vessel operations.

2.3.4   Country Attributes

Country attributes include whether the country is involved in international agreements, how

effective their National Plan of Action for Sharks (NPOA-Sharks) is, if one exists, and the

amount given in subsidies. These attributes deal with the relationship of the country to the

rest of the world through the engagement with international treaties and agreements like

CITES and CMS. The scores in this group are indicative of how managers in each country

consider preservation of at-risk species. For countries with lower scores in the country

attributes, we would recommend improving or drafting an NPOA-Sharks, and reallocating

subsidy funding to more beneficial outcomes like marine protected area implementation

and management as opposed to tax exemptions and fuel subsidies.

2.3.5   RFMO Attributes

RFMO attributes include whether membership parties are involved in international treaties

and agreements like CITES and CMS, and their progress on ecosystem-based fisheries

management (EBFM). Similar to the country attribute, membership in CITES and CMS shows

willingness to protect species-at-risk and EBFM progress shows the care of the RFMO for the

overall ecosystem in which they are fishing. For RFMOs with lower scores in these

attributes, we would recommend improving their progress towards EBFM and encouraging

member countries to participate in CITES and CMS, without species reservations.

2.4   Scoring of Attributes

Each attribute was scored individually based on detailed value statements (e.g., **Table 2**),

such that a higher score indicated a higher likelihood of achieving a sustainable outcome for

the species (full value statements available in **Supplementary Information 1**). Our scoring of

attributes assigned the highest scores to the ideal 'counterfactual' situation (Juan-Jordá,

Murua, Arrizabalaga, Dulvy, & Restrepo, 2018). We chose narrow ranges for ordinal scoring,

(either 0-3, 0-4, or 0-5, similar to a Likert scale) and scored attributes against the value

statements found in **supplementary information 1**. In cases where information was absent,

precautionary scores of zero were given (Hobday et al., 2011). In some cases, an attribute

was not applicable to the species being assessed (i.e., an attribute determining post-release

survival is not relevant to a species that is always retained). When this occurred, the

attribute would simply not be scored and left out of the final calculations. The narrow range

of scores ensured consistency of scoring over time, across jurisdictions, and across

assessors. As such the range of uncertainty in scoring can only be low. We did consider

scoring over a wider scale, say 0-9, which would have allowed assigning a range instead of a

single score, and, therefore, allow incorporation and propagation of uncertainty, say with a

Fuzzy Logic or Bayesian approach. However, we deemed that it would be much more

difficult to ensure consistent scoring across assessors and jurisdictions. For countries, nine

of the attributes were universal (U) for the entire fishery and the remaining ten attributes

were specific to each species being assessed (SP). For RFMOs, seven were universal and ten

were species-specific (**Table 1**). The final management score, when all attributes were

assessed, was calculated based on points scored out of potential points available and

converted to a percentage towards ideal management.

For RFMOs, scores were assigned based on legislation agreed to by all member countries.

Some countries may have additional requirements, however, these were not considered for

the RFMO score because we took a precautionary approach and assessed based on the

minimum standards for all vessels operating within a fishery. Here, we use the term 'score'

rather than the term 'index'. While the academic literature uses the term 'index' to refer to

any mathematical formula by which different variables are combined, the policy world tends

to use the word 'index' only when a metric or score has been adopted for use. The reality is

that there are many candidate indicators, but very few are fit-for-purpose following

extensive performance testing {Newson, 2009 #8482; Rice, 2005 #8483}. The highest

possible score (100%) indicates ideal management for sharks and rays; therefore, final

management scores for each management unit indicated their progress towards the ideal.

Management scores are intended to be used for comparisons between management units

and species, and are of little value alone as no fisheries are expected to achieve a score of

100%. Comparisons can be made (1) between different species within a single management

unit, (2) the average score between different management units, (3) between a single

species in each management unit that it has been assessed, and (4) between different

taxonomic groups within or across different management units.

2.5 Representing Intrinsic Sensitivity

To understand Vulnerability (equation 2) we calculate the intrinsic sensitivity of a species to

overexploitation in addition to the management risk assessment. Factors that contribute to

a greater likelihood of population decline or higher intrinsic sensitivity for marine species

include large body size and a slow speed of life (e.g. slow somatic growth rate, late age-at-

maturity, or long generation length) (Juan-Jordá, Mosqueira, Freire, & Dulvy, 2015; Lee &

Jetz, 2011; Reynolds, Dulvy, Goodwin, & Hutchings, 2005).

Elasmobranchs as a group are characterised as having long life spans, late age-at-maturity,

and low fecundity (Cortés, 2000; Field, Meekan, Buckworth, & Bradshaw, 2009), and

consequently they have a range of intrinsic sensitivities (Pardo, Kindsvater, Reynolds, &

Dulvy, 2016). However, within the group there are extremes for all characteristics. For

example, the Whale Shark (*Rhincodon typus*) is the largest fish species in the world,

attaining a maximum body length of up to 20 m (Chen, Liu, & Joung, 1997). On the other

end of the scale, the Dwarf Lanternshark (*Etmopterus perryi*) attains a maximum body

length of just 21 cm, approximately one hundredth the maximum body length of Whale

Shark (Ebert et al., 2021). Relating to the speed of life, Rigby and Simpfendorfer (2015)

discuss the high intrinsic sensitivity of deepwater sharks due to their late age-at-maturity

despite their relatively small body size, and thus, the consequences related to their capture

in developing deepwater fisheries.

Intrinsic sensitivity can be categorized as low, medium, or high, based on a number of traits

that can include age-at-maturity, length-at-maturity, longevity, maximum body length,

fecundity, reproductive strategy, and trophic level (Cheung, Pitcher, & Pauly, 2005;

Georgeson et al., 2020; Musick, 1999). Data are not available for most shark and ray species

for all these traits, therefore we can only use a few reliably available across all species.

Oldfield et al. (2012) suggested that minimum age-at-maturity, reproductive strategy, and

maximum body length were the three most important factors for sharks and rays,

respectively. However, there have been considerable advances in comparative life history

theory and it is clear that there are three dimensions to consider in this order of

importance: maximum body length, speed of life traits (growth rate, age at maturity,

maximum age), and reproductive output (Cortés, 2000; Juan-Jordá, Mosqueira, Freire, &

Dulvy, 2013). Many of these traits are not commonly known for shark and ray species and

surprisingly, reproductive output contributes little to the maximum intrinsic rate of

population increase, except in sharks and rays with very low fecundity of typically fewer

than five pups per year (Forrest & Walters, 2009; Pardo et al., 2016), therefore, fecundity is

least useful in determining intrinsic sensitivity. Similarly, reproductive strategy would be a

binary option, either egg-laying (oviparity) or giving birth to live young (viviparity), and

without corresponding information on fecundity or the relationship to maximum intrinsic

rate of population increase, would be difficult to directly compare. Finally, trophic level is

widely available based on dietary analyses and was considered, however, due to limited

species-specific data and the change in trophic level with ontogenetic dietary shifts, we did

not include this as a variable when calculating intrinsic sensitivity (Bethea et al., 2007;

Lucifora, García, Menni, Escalante, & Hozbor, 2009).

For our analyses, we used two traits to represent intrinsic sensitivity – generation length

(the midpoint between age-at-maturity and maximum age) and maximum body length.

Many of the shark and ray generation lengths reported in IUCN Red List Assessments are

inferred or suspected from closely related species (65% of species considered by M-Risk;

Supplementary Information 2). Thus, we categorised generation lengths into 5-year bins,

such that species with longer generation lengths have higher intrinsic sensitivities. We

scored our confidence in the generation length for each species based on whether it was

species-specific and the range was within a 5-year bin (high confidence), range spanned

multiple 5-year bins or was based on estimated life-history parameters (medium), or was

inferred from a congener (low)(**Data S1**). There was a significant positive linear relationship

between generation length and relative maximum body length (**Supplementary Information

2**). However, when considering only species with high confidence generation lengths, there

was no significant relationship (**Supplementary Information 2 - Figure 1**). This is likely a

result of estimated generation lengths being scaled to maximum body length. We then

included maximum body size as a second measure of intrinsic sensitivity, of which we could

be confident in the species-specific values (**Figure 2**). Ideally, we would use maximum

weight but there are few such measures for all species and we encourage their collection.

Therefore, we used maximum linear dimension, derived from either maximum length or

maximum disc width for some rays, and scored it such that the largest species (*Maximum

size$_L$*) of those we assessed was assigned the reference highest intrinsic sensitivity (i.e.,

Relative Size = 100%) and the *Maximum size$_x$* for each other species (*x*) was scaled as a

percentage of the largest species (**Figure 2**):

$$\text{Equation 3:} \quad Relative\ Size = \left(\frac{Maximum\ size_x}{Maximum\ size_L}\right) * 100$$

We considered four groups of body morphologies because they are measured in different

ways (sharks and shark-like rays, pelagic rays, classic rays, and ghost sharks), and took the

largest species for each body morphology and assigned them at 100% *Relative Size*. Whale

Shark and Basking Shark (*Cetorhinus maximus*) are outliers, measuring over 1,200 cm longer

than the next largest species, therefore, we considered the White Shark (*Carcharodon*

carcharias) for these calculations, which attains a maximum size of 640 cm. All these three

shark species were assigned a *Relative Size* score of 100% for their maximum size. Intrinsic

sensitivity values in this paper are based on the larger project species list (**Data S1**), rather

than just those species within this paper to ensure that *Relative Size* is conserved. Both

intrinsic sensitivity scores (*Relative GL* and *Relative Size*) were averaged for an overall

intrinsic sensitivity (*IS*) score (**Figure 2**):

$$\text{Equation 4:}\quad Intrinsic\ Sensitivity = \frac{(Relative\ Size\ +\ Relative\ GL)}{2}$$

526   It is not possible for a species to receive an intrinsic sensitivity score of 0% because all

527   species are intrinsically at risk, even if the risk is small.

529   The intrinsic sensitivity (IS) was then divided by the management assessment score to attain

530   a final M-Risk vulnerability score for each species within each management unit they are

531   assessed:

$$\text{Equation 5:}\quad Vulnerability = (\frac{Intrinsic\ Sensitivity * Exposure}{Management})$$

As species exposure to each fishery is difficult to measure, we assume that Exposure = 1 as

assessments are only performed on species that are caught or directly affected by the

fishery in question. This is the maximum possible exposure and is consistent with the

precautionary principle. For countries, there are 19 attribute scores (*A*) with a maximum

possible score for each attribute of *Max.A*, and weighting (*W*), therefore, the formula to

calculate the M-Risk score is:

$$\text{Equation 6: } Vulnerability = \frac{\left(\frac{\left(\frac{Maximum\ size_x}{Maximum\ size_L} * 100\right) + Relative\ GL)}{2}\right)}{\sum_{n=1}^{19}\left(W_n\left[\frac{A_n}{Max.A_n}\right] * 100\right)}$$

538   We have weighted all attributes equally (*W* = 1) so the bottom of equation 6 simplifies to

539   the total points scored over the total possible points. The final M-Risk formula simplifies to:

$$\text{Equation 7:}\quad Vulnerability = (\frac{Intrinsic\ Sensitivity}{Management})$$

3. APPLYING THE M-RISK FRAMEWORK

To explore the application of the M-Risk, we applied our criteria to two case studies, one country (Ecuador), and one RFMO (Inter-American Tropical Tuna Commission, IATTC).

3.1 Case Studies: Country and RFMO for Management Assessments

We completed assessments for all species that occur within two management units in order to determine the efficacy of our management-risk assessment framework; one country (Ecuador) and one RFMO (Inter-American Tropical Tuna Commission, IATTC). These assessments were completed for 35 species in Ecuador (23 sharks, 12 rays) and 27 species in the IATTC (20 sharks, 7 rays).

3.1.1 Case Study 1: Ecuador

3.1.1.1 Information Sources

We searched Google and Google Scholar in both English and Spanish for the following four elements in sequence: (i) Ecuadorian fisheries regulations, (ii) specific attribute keywords (i.e., "Ecuador shark finning regulations"), (iii) species name (English, Ecuadorian, Spanish, and Latin binomial) and "Ecuador fishery", and (iv) species name in all relevant languages and specific attribute keywords (i.e., "Ecuador Whale Shark catch limits") as has been done previously in ecological risk assessments (Cortés et al., 2010). Ten of the attributes were fishery-specific and did not require steps iii and iv. All attributes were scored on the most current publicly available information. We acknowledge, however, that in some cases this information may be out-of-date, and the most recent regulations are not available online. In cases where no information was available, the precautionary approach was applied, and it was concluded that there were no regulations related to the species in question and a score of zero was given. Again, we acknowledge that this information may exist, but we were unable to access due to it not being publicly available or readily accessible. For this reason, we do not consider uncertainty surrounding scores as exhaustive searches were completed and our framework was developed with the intent to reward transparency, i.e., if we couldn't find the information in a reasonable amount of time then this was regarded as problematic reflecting lower levels of transparency and availability and thus scored as a

zero. The point is that knowledge that exists somewhere, but is not available, is not likely

helpful to management or transparency.

Four primary resources were used to complete the Ecuador management risk assessments:

(1) Ecuador's National Fisheries Institute website (www.institutopesca.gob.ec), (2) Ministry

of Aquaculture and Fisheries website (acuaculturaypesca.gob.ec), (3) the Ecuador Law

website (www.derechoecuador.com), and (4) a paper by Martínez-Ortiz et al. (2015)

describing enforcement measures and species-specific catch data. Although multiple

fisheries exist in Ecuador, two were chosen to be representative of their management. The

Large Pelagic Artisanal Fishery was selected as it accounted for ~93% of shark and ray catch

in Ecuador (Alava, Lindop, & Jacquet, 2015). The only exception was for the Whale Shark as

there is common interaction between this species and purse-seine vessels that does not

exist with other species, thus the Purse Seine Fishery was chosen to assess Whale Shark

management in Ecuador (Dagorn, Holland, Restrepo, & Moreno, 2013; Rowat & Brooks,

2012).

3.1.1.2 Management Assessment Results

Management assessment scores from all 35 species ranged from 53 to 68% of the ideal

score. One species (Whale Shark) received the highest score of 68%, five species (3 sharks, 2

rays) scored 67%, and the final 29 species (19 sharks, 10 rays) scored 53 to 62% (**Figure 3a;**

Table S1). The differences in management scores were mainly due to three attributes: (1)

landing limits, (2) post-release survival, and (3) catch reporting. Scores for the landing limits

attribute ranged from 0 to 3 (the highest score possible), while post-release survival had

range of 0 to 2 (out of a possible 3) and catch reporting ranged from 1 to 4 (out of 4). The

scores for the remaining 16 attributes were the same across all species in the Large Pelagic

Artisanal Fishery.

3.1.1.3 Management Assessment Discussion

There was little connection between the IUCN status and M-Risk scores with threatened

species included in both higher and lower scoring groups. Almost all species with higher

scores are listed on both CITES and CMS appendices, indicating clear progress for listed

species. All species listed on CITES prior to 2014 were included in the higher scoring group.

Eight CITES and nine CMS-listed species were in the lower scoring group, indicating it takes

some time to implement regulations relating to these agreements. This has been seen in

other CITES-listed species, like seahorses, which were listed on CITES in 2002, however,

Thailand continued exporting high numbers without a positive Non-Detriment Finding until

2016 (Kuo, Laksanawimol, Aylesworth, Foster, & Vincent, 2018). What was more likely to

determine a higher score was species charisma. Charismatic species likely to be of high

tourism interest are often subject to increased conservation efforts (Albert, Luque, &

Courchamp, 2018; Hausmann, Slotow, Fraser, & Di Minin, 2016; McClenachan, Cooper,

Carpenter, & Dulvy, 2011), whereas a very large fraction of highly threatened species can go

unstudied and unmanaged (Guy et al., 2021). This was apparent in the scores in Ecuador, as

all species in the higher scoring group had some type of retention ban or catch limit (**Table**

S1). These limits, in addition to CITES and CMS compliance, were likely placed to protect

tourism value and public perception.

3.1.2 Case Study 2: IATTC

3.1.2.1 Information Sources

Like assessments for Ecuador, we searched Google and Google Scholar for the same four

elements: (i) IATTC fisheries regulations, (ii) specific attribute keywords, (iii) species name

(English and Latin binomial), and (iv) species name and specific attribute keywords. For

these assessments, we scored both the purse seine and the longline regulations, and the

higher score was used. We used one primary and three secondary resources to complete

the IATTC management risk assessments. Primarily, we used the IATTC website

(www.iattc.org), which provided in-depth documents related to management regulations

and fishery knowledge. Additionally, we found information from: (1) the Food and

Agriculture Organization of the United Nations website (ww.fao.org), (2) the National

Oceanic and Atmospheric Agency website (www.noaa.gov), and (3) the International

Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean website

(isc.fra.go.jp).

3.1.2.2 Management Assessment Results

Scores from the 27 species assessed ranged from 46–71% of an ideal score in the IATTC,

with an average score of 58.3% (**Figure 3b; Table S1**). A single species, the Silky Shark

(*Carcharhinus falciformis*), stood out with the highest score (71%). Three species, the Pelagic

Stingray (*Pteroplatytrygon violacea*) and two eagle ray species (*Aetobatus laticeps* and *A.*

ocellatus) both had scores of 50% or lower (50, 46 and 46%, respectively) and 14 of the 27

species scored between 50-59% (**Figure 3b; Table S1**). Catch reporting was the attribute

with the lowest scores for the IATTC because half of the species were only reported to

broad taxonomic categories (i.e., "sharks" or "rays"). Only six species assessed had any form

of landing limits in place (Silky Shark, Oceanic Whitetip Shark (*Carcharhinus longimanus*),

Giant Manta Ray (*Mobula birostris*), and devil rays (*Mobula mobular, M. munkiana,* and *M.*

thurstoni), while the rest were only *recommended* to be released once caught. These results

show that overall, the IATTC has species-specific regulations in place for only a select few

species and the others are subject to marginal management applicable to elasmobranchs. It

remains unclear whether these measures can significantly reduce fishing mortality and if the

management focus is on the species that are most at-risk to overfishing within the fishery.

3.1.2.3 Management Assessment Discussion

Within the IATTC, there does not appear to be any relationship between global IUCN status

and the management assessment scores. Of the 27 species assessed, only five are not

currently listed in a threatened category, yet these are not the five with the lowest scores.

Similarly, CITES and CMS listings appear to have not had any bearing on the management

assessment scores. The Whale Shark has been listed on both CITES (2003) and CMS (App I in

2017, App II in 1999) for much longer than many other sharks and rays and is

unquestionably one of the most charismatic species, which we would expect to lead to a

higher management score. However, it received a score of only 55% in the IATTC. The score

may be lower than expected as there is little interaction between the fishery and whale

sharks, thus lesser concern to provide specific legislation for this species. Additionally,

where interactions do occur, there is a prohibition on setting nets when a whale shark is

sighted (Inter-American Tropical Tuna Commission, 2019). The Silky Shark, which has been

more recently listed on both CITES (2017) and CMS (2014) appendices, received the highest

score of all species assessed. This surprisingly rapid management response may be due to

the large amount of recent attention and market pressure to reduce the plight of Silky Shark

caught in fish aggregating devices (FADs) and their susceptibility to purse seines (Duffy et

al., 2015; Filmalter, Capello, Deneubourg, Cowley, & Dagorn, 2013; Hutchinson, Itano, Muir,

& Holland, 2015). Although the purse-seine tuna fishery catches mostly shark 'bycatch',

there are several pelagic ray species caught within the fishery. Although CITES listing may

not necessarily be considered by RFMOs when discussing regulations, it should be. CITES is a

legally binding agreement which covers relevant issues like 'Introduction From the Sea',

including the high seas that constitute the fishing grounds plied by vessels operating under

RFMOs, and countries' responsibilities for legal chain of custody when CITES-listed species

are landed from high seas vessels (Pavitt et al., 2021). With the exception of manta and

devil rays, which are no-retention species (Inter-American Tropical Tuna Commission, 2015),

there are no ray-specific regulations to reduce their catch. While shark species benefit from

more generic legislation including the prohibition of shark lines (which intentionally catch

sharks as retained bycatch), a code of practice to increase post-release survival, and fin-to-

meat landing ratios to reduce finning, rays are not afforded any additional regulations

(Inter-American Tropical Tuna Commission, 2005, 2016). In order to decrease management

risk for elasmobranch species in the IATTC, implementing science-based landing limits for

the most commonly caught species and including more legislation to evaluate and reduce

ray fishing and post-release morality are recommended.

3.2 Final M-Risk Scores

The final M-Risk score for a species shows whether the current management is appropriate

based on its intrinsic sensitivity (IS). Species with high intrinsic sensitivity will need higher

management assessment scores to achieve a similar M-Risk score to lower IS species. As an

example, the Great Hammerhead (*Sphyrna mokarran*) has a generation length of 24.8 years

for a *Relative GL* score of 80% and reaches a maximum size of 610 cm total length, which is

95.3% of the maximum sized shark (White Shark – 640 cm), therefore, the combined

intrinsic sensitivity score for Great Hammerhead is 87.7% (**Figure 2; Table S1**). Compared to

the Pelagic Stingray (*Pteroplatytrygon violacea*), which has a generation length of 6.5 years

(*Relative GL* of 20%) and reaches a maximum size of 90 cm disc width, which is 34.6% of the

maximum sized classic ray (Spiny Butterfly Ray, *Gymnura altavela* – 260 cm), therefore, the

combined intrinsic sensitivity score for Pelagic Stingray is 27.3%. Although both the Great

Hammerhead and Pelagic Stingray received similar scores in their management assessments

for both Ecuador and the IATTC, when combined with their intrinsic sensitivity scores, the

final Vulnerability is vastly different. The Great Hammerhead is vastly undermanaged for its

life history compared to the Pelagic Stingray (**Figure 3; Table S1**). Although the Pelagic

Stingray scored lower in the management assessment for the IATTC than the Great

Hammerhead (50.0% and 58.9%, respectively), the final M-Risk scores show it is almost one-

third at risk due to undermanagement in the IATTC than the Great Hammerhead due to its

lower intrinsic sensitivity (M-Risk scores of 0.51 and 1.50, respectively). In Ecuador, the

Pelagic Stingray is similarly approximately one-third as at-risk compared to Great

Hammerhead (0.55 and 1.49, respectively). These scores describe the true risk to

undermanagement by each of these species based on our current understanding of these

species and their populations.

4. DISCUSSION

Here, we presented a framework for a management risk assessment (M-Risk) that is

designed: (1) for assessing different management approaches for individual species *within* a

management unit (of country or RFMO), (2) for comparing the efficacy of shark and ray

management *across* different management units globally, and (3) for comparing

management efficacy of a single species in all management units it is found. We also

summarise the two main findings from this proof-of-concept. Firstly, different shark and ray

species within a single management unit have management assessment scores that range

from ~50-70% of a score that would be consistent with the 'ideal' management in the two

fisheries assessed (Ecuador and the IATTC). Second, when accounting for species' intrinsic

sensitivities, the final M-Risk scores (Vulnerability in Equation 5) best show which species

are most at risk due to management deficiencies.

Intrinsic sensitivity of each species is important and particularly beneficial to our

understanding of risk differences for species caught in poorly managed fisheries that have

low intrinsic sensitivity, compared to high-risk species in adequately managed fisheries.

Incorporating species-specific sensitivities to this analysis shows the nuance of managing

sharks and rays, which are frequently viewed as a large group with similar attributes. With

this approach, the difficulty of having good management for higher risk species is

demonstrated as those with high intrinsic sensitivity scores will always have a higher final

M-Risk score than species with lower intrinsic sensitivities. Upon completion of a wider

array of countries with varying management regimes, we will be able to better assign final

M-Risk scores to either low, medium, or high-risk groupings. This will provide an

understanding and priority setting for which species, in which management units, are most

at risk of overexploitation and at what M-Risk score intervention should begin. Additionally,

after more management units are scored, we will evaluate the strength of pairwise

correlations between attribute scores and/or country and species traits. This will allow us to

ask questions such as, "do countries with higher Human Development Indices always score

higher in the 'Country Attributes'?" and "in fisheries with a high score in the Catch

Validation attribute, do they always receive a high score for Enforcement Methods?" The

answers to these questions may uncover causal relationships that provide clear paths to

improving fisheries management. Next, we explore the differences in scores *across*

management units, how our M-Risk approach compares to other PSAs, and future directions

for this work.

Two management units (IATTC and Ecuador) were assessed using the M-Risk framework.

Once additional management units are assessed, countries will be compared to other

countries and RFMOs can all be compared to one another. In describing the applications of

this risk assessment framework, this discussion will assume these two management units

are directly comparable. Despite both receiving similar scores for species in their respective

jurisdictions, the scores for specific attribute categories differed. For example, the IATTC

species, on average, scored 30% higher in the "Management System" attributes than

species in Ecuador (73% and 44%, respectively). However, in the "Fishing Practices and

Catch" attributes, species in Ecuador scored, on average, 20% higher than those in the IATTC

(59% and 37%, respectively; **Figure 3; Table S1**). Based on these results, each management

unit could benefit from studying regulations in the other to improve their own shark and ray

management. Comparing management units using the same framework not only enables a

level playing field, but also allows for management units with lower scores to more easily

identify areas to improve their management efficacy. Management units lacking sufficient

funding to complete their own fisheries research into which regulations have the greatest

impact on shark and ray fishing mortality can learn from or borrow legislation from those

with highly effective management strategies. This can also apply to sections of management

that are lacking in sufficient shark and ray management. If adopted, this would increase the

efficacy of global fisheries management for sharks and rays overall. A limitation of this type

of assessment is it does not consider compliance with legislation. This includes compliance

by the fishers, fisheries officers, and managers who are policing the regulations and

assumes countries are meeting their obligations to agreements to which they are

signatories, including CITES and for RFMOs. Without effective compliance and enforcement

of the fisheries regulations, the management scores we have assigned for each fishery are a

best-case scenario and likely over-estimate the management effectiveness. Since these

assessments are intended to be completed by external assessors using publicly available

data, the scores should be considered minimum possible scores for each management unit

if all regulations are not readily available online and consequently, our approach is

precautionary.

Future directions for the M-Risk framework may include a capacity to weight attributes and

include an evaluation of the exposure component. For this global comparison, weighting

attributes is not appropriate as the goal is to compare across all fishery types, gears,

countries, etc. For example, in Bangladesh the commercial and artisanal sectors operate

with different goals and budgets and in different areas (Islam et al., 2017; Kumar et al.,

2019). Therefore, an attribute that may be weighted higher for assessing the commercial

fisheries may not make sense to be weighted that way when assessing subsistence fisheries.

There are methods available to assign post-hoc weightings to attributes that remove the

subjectivity of determining "importance" (Chen, 2019). Post-hoc weighting of attributes may

be applied when using the management assessment framework to suit a particular goal.

However, as the framework is designed for ongoing use, post-hoc weightings may continue

to change as more assessments are completed. In our current assessments, exposure has

been treated as a binary variable, despite normally ranging from 0–1, because we are

assessing species in trade, therefore, they are assumed to be highly exposed. However, if

information about the fishing grounds and depths of species becomes easier to acquire,

there is potential to include exposure in future M-Risk work. Additionally, this framework

does not consider cumulative impacts due to multiple fisheries or threats, including climate

change, environmental modifications and other anthropogenic hazards, that may

significantly increase overall risk to some species (Walker et al., 2021). However, this

framework could be readily adapted to incorporate vulnerability to climate change through

the addition of Equation 3, section 2.2.12 of Walker et al. (2021). We have not explored the

implications of multiple assessments through time, however, anticipate this could provide

valuable indications of management improvement or decline, given the assessments are

completed against the same criteria in the same manner.

The results of this proof of concept provide a basis for further investigation of shark and ray

management globally. With a larger number of management units assessed, we will have a

better understanding of the efficacy of shark and ray management globally and which

management units or species are most at risk due to undermanagement based on their

intrinsic sensitivity. The M-Risk assessment tool may be a step to identify species-at-risk and

appropriate management to implement that would lower risk to those species. Currently,

international agreements like CITES and CMS are in place once species have been depleted.

However, identifying species-at-risk to overexploitation and implementing management to

rebuild and achieve sustainable fisheries and avoid population collapses, may reduce the

need for some species to be included in these international agreements in the first place as

maintenance is simpler than recovery.

ACKNOWLEDGEMENTS

The authors thank Heather Patterson for her input of attributes and their scoring. We thank

Thomasina Oldfield and Dulvy lab members for comments on manuscript drafts.

Additionally, we would like to thank Ian Gregor for help with coding the graphs. We are

grateful to the three reviewers and editor who provided valuable comments that improved

this manuscript. This work was supported by a grant to GS from the Shark Conservation

Fund, a philanthropic collaborative that pools expertise and resources to meet the threats

facing the world's sharks and rays. The Shark Conservation Fund is a project of Rockefeller

Philanthropy Advisors. NKD was supported by the Discovery and Accelerator grants from

Natural Science and Engineering Research Council and the Canada Research Chair program.

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

Individual attribute scores for each species in each management unit, and the management

documents used for scoring are available in the Supplementary Information (**Data S2** and

Data S3, respectively).

REFERENCES

Agnew, D. J., Pearce, J., Pramod, G., Peatman, T., Watson, R., Beddington, J. R., & Pitcher, T.
J. (2009). Estimating the worldwide extent of illegal fishing. *PLoS ONE, 4*(2), e4570.
doi:10.1371/journal.pone.0004570

Alava, J. J., Lindop, A., & Jacquet, J. (2015). *Marine fisheries catch reconstructions for*
continental Ecuador: 1950-2010. Retrieved from Vancouver, British Columbia,
Canada: doi:10.13140/2.1.1150.5447

Albert, C., Luque, G. M., & Courchamp, F. (2018). The twenty most charismatic species. *PLoS*
ONE, 13(7), e0199149. doi:10.1371/journal.pone.0199149

Allen Consulting Group. (2005). *Climate Change Risk and Vulnerability: Promoting an*
Efficient Adaptation Response in Australia. Retrieved from Canberra, ACT:
http://www.sfrpc.com/Climate%20Change/4.pdf

Allison, E. H., Perry, A. L., Badjeck, M.-C., Adger, W. N., Brown, K., Conway, D., . . . Dulvy, N.
K. (2009). Vulnerability of national economies to the impacts of climate change on
fisheries. *Fish and Fisheries, 10*, 173-196. doi:10.1111/j.1467-2979.2008.00310.x

Amelot, M., Batsleer, J., Foucher, E., Girardin, R., Marchal, P., Poos, J. J., & Sys, K. (2021).
Evidence of difference in landings and discards patterns in the English Channel and
North Sea Rajidae complex fishery. *Fisheries Research, 242*, 106028.
doi:10.1016/j.fishres.2021.106028

Anderson, S. C., Branch, T. A., Ricard, D., & Lotze, H. K. (2012). Assessing global marine
fishery status with a revised dynamic catch-based method and stock-assessment
reference points. *ICES Journal of Marine Science, 69*(8), 1491-1500.
doi:10.1093/icesjms/fss105

Astles, K. L., Holloway, M. G., Steffe, A., Green, M., Ganassin, C., & Gibbs, P. J. (2006). An
ecological method for qualitative risk assessment and its use in the management of
fisheries in New South Wales, Australia. *Fisheries Research, 82*, 290-303.
doi:10.1016/j.fishres.2006.05.013

Begg, G. A., & Waldman, J. R. (1999). An holistic approach to fish stock identification.
Fisheries Research, 43(1–3), 35-44. doi:10.1016/S0165-7836(99)00065-X

Bethea, D. M., Hale, L., Carlson, J. K., Cortes, E., Manire, C. A., & Gelsleichter, J. (2007).
Geographic and ontogenetic variation in the diet and daily ration of the bonnethead
shark, Sphyrna tiburo, from the eastern Gulf of Mexico. *Marine Biology, 152*(5),
1009-1020.

Brander, K. (1981). Disappearance of common skate *Raja batis* from Irish Sea. *Nature, 290*,
48-49. doi:10.1038/290048a0

Brooks, E. N., Powers, J. E., & Cortés, E. (2010). Analytical reference points for age-
structured models: application to data-poor fisheries. *ICES Journal of Marine Science,*
67(1), 165-175. doi:10.1093/icesjms/fsp225

Cashion, M. S., Bailly, N., & Pauly, D. (2019). Official catch data underrepresent shark and
ray taxa caught in Mediterranean and Black Sea fisheries. *Marine Policy, 105*, 1-9.
doi:10.1016/j.marpol.2019.02.041
Chen, C. T., Liu, K. M., & Joung, S. J. (1997). Preliminary report on Taiwan's whale shark
fishery. *TRAFFIC Bulletin, 17*(1), 53-57.
Chen, P. (2019). On the diversity-based weighting method for risk assessment and decision-
making about natural hazards. *Entropy, 21*, 269. doi:10.3390/e21030269
Cheung, W. W. L., Pitcher, T. J., & Pauly, D. (2005). A fuzzy logic expert system to estimate
intrinsic extinction vulnerabilities of marine fishes to fishing. *Biological Conservation,*
124(1), 97-111.
Chrysafi, A., & Kuparinen, A. (2015). Assessing abundance of populations with limited data:
lessons learned from data-poor fisheries stock assessments. *Environmental Reviews,*
24, 25-38. doi:dx.doi.org/10.1139/er-2015-0044
Cinner, J. E., Huchery, C., Hicks, C. C., Daw, T. M., Marshall, N., Wamukota, A., & Allison, E. H.
(2015). Changes in adaptive capacity of Kenyan fishing community. *Nature Climate*
Change, 5, 872-877. doi:10.1038/NCLIMATE2690
Cope, J. M. (2013). Implementing a statistical catch-at-age model (Stock Synthesis) as a tool
for deriving overfishing limits in data-limited situations. *Fisheries Research, 142*, 3-
14. doi:10.1016/j.fishres.2012.03.006
Cope, J. M., & Punt, A. E. (2009). Length-based reference points for data-limited situations:
applications and restrictions. *Marine and Coastal Fisheries: Dynamics, Management,*
and Ecosystem Science, 1, 169-186. doi:10.1577/C08-025.1
Cortés, E. (2000). Life history patterns and correlations in sharks. *Reviews in Fisheries*
Science, 8(4), 299-344. doi:10.1080/10408340308951115
Cortés, E., Arocha, F., Beerkircher, L., Carvalho, F., Domingo, A., Heupel, M., . . .
Simpfendorfer, C. (2010). Ecological risk assessment of pelagic sharks caught in
Atlantic pelagic longline fisheries. *Aquatic Living Resources, 23*, 25-34.
doi:10.1051/alr/2009044
Cortés, E., Brooks, E. N., & Shertzer, K. W. (2015). Risk assessment of cartilaginous fish
populations. *ICES Journal of Marine Science, 72*(3), 1057-1068.
doi:10.1093/icesjms/fsu157
Costello, C., Ovando, D., Hilborn, R., Gaines, S. D., Deschenes, O., & Lester, S. E. (2012).
Status and solutions for the world's unassessed fisheries. *Science, 338*, 517-520.
doi:10.1126/science.1223389
Dagorn, L., Holland, K. N., Restrepo, V., & Moreno, G. (2013). Is it good or bad to fish with
FADs? What are the real impacts of the use of drifting FADs on pelagic marine
ecosystems? *Fish and Fisheries, 14*(3), 391-415. doi:10.1111/j.1467-
2979.2012.00478.x
Davidson, L. N. K., Krawchuk, M. A., & Dulvy, N. K. (2015). Why have global shark and ray
landings declines: improved management or overfishing? *Fish and Fisheries, 17*(2),
438-458. doi:10.1111/faf.12119
Dent, F., & Clarke, S. (2015). *State of the global market for shark products* (590). Retrieved
from https://www.fao.org/in-action/globefish/publications/details-
publication/en/c/338282/
Dragičević, B., Dulčić, J., & Capapé, C. (2009). Capture of a rare shark, *Oxynotus centrina*
(Chondrichthyes: Oxynotidae) in the eastern Adriatic Sea. *Journal of Applied*
Ichthyology, 25(1), 56-59. doi:10.1111/j.1439-0426.2009.01265.x

Duffy, L. M., Olson, R. J., Lennert-Cody, C. E., Galvan-Magana, F., Bocanegra-Castillo, N., &
Kuhnert, P. M. (2015). Foraging ecology of silky sharks, *Carcharhinus falciformis*,
captured by the tuna purse-seine fishery in the eastern Pacific Ocean. *Marine*
Biology, 162(3), 571-593. doi:10.1007/s00227-014-2606-4
Dulvy, N., Reynolds, J. D., Pilling, G. M., Pinnegar, J. K., Scutt Phillips, J., Allison, E. H., &
Badjeck, M.-C. (2011). Fisheries management and governance challenges in a climate
change. In *The Economics of Adapting Fisheries to Climate Change* (pp. 31-88). Paris,
FR.: OECD Publishing.
Dulvy, N. K., Metcalfe, J. D., Glanville, J., Pawson, M. G., & Reynolds, J. D. (2000). Fishery
stability, local extinctions, and shifts in community structure in skates. *Conservation*
Biology, 14(1), 283-293.
Dulvy, N. K., Pacoureau, N., Rigby, C. L., Pollom, R. A., Jabado, R. W., Ebert, D. A., . . .
Simpfendorfer, C. A. (2021). Overfishing drives over one-third of all sharks and rays
toward a global extinction crisis. *Current Biology, 31*, 1-15.
doi:10.1016/j.cub.2021.08.062
Dulvy, N. K., Simpfendorfer, C. A., Davidson, L. N. K., Fordham, S. V., Bräutigam, A., Sant, G.,
& Welch, D. J. (2017). Challenges and priorities in shark and ray conservation.
Current Biology, 27, R565-R572. doi:10.1016/j.cub.2017.04.038
Ebert, D. A., Dando, M., & Fowler, S. (2021). *Sharks of the World* (2nd ed.). Princeton, New
Jersey: Princeton University Press.
Enguehard, R. A., Devillers, R., & Hoeber, O. (2013). Comparing interactive and automated
mapping systems for supporting fisheries enforement activities - a case study on
vessel monitoring systems (VMS). *Journal of Coastal Conservation, 17*, 105-119.
doi:10.1007/s11852-012-0222-3
FAO. (1999). *International Plan of Action for reducing incidental catch of seabirds in longline*
fisheries. International Plan of Action for the conservation and management of
sharks. International Plan of Action for the management of fishing capacity.
Retrieved from Rome: https://www.fao.org/3/x3170e/x3170e.pdf
FAO. (2019). FishStatJ (Version 4.00.0). Rome: FAO.
FAO. (2020). *The State of World Fisheries and Aquaculture 2020. Sustainability in action*.
Retrieved from Rome, Italy: https://www.fao.org/documents/card/en/c/ca9229en/
FAO. (2021). *ASFIS List of Species for Fisheries Statistics Purposes*. Retrieved from:
https://www.fao.org/fishery/collection/asfis/en
Field, I. C., Meekan, M. G., Buckworth, R. C., & Bradshaw, C. J. A. (2009). Susceptibility of
sharks, rays and chimaeras to global extinction. *Advances in Marine Biology, 56*, 275-
363. doi:10.1016/S0065-2881(09)56004-X
Filmalter, J. D., Capello, M., Deneubourg, J.-L., Cowley, P. D., & Dagorn, L. (2013). Looking
behind the curtain: quantifying massive shark mortality in fish aggregating devices.
Frontiers in Ecology and the Environment. doi:10.1890/130045
Fletcher, W. J. (2005). The application of qualitative risk assessment methodology to
prioritize issues for fisheries management. *ICES Journal of Marine Science, 62*, 1576-
1587. doi:10.1016/j.icesjms.2005.06.005
Forrest, R. E., & Walters, C. J. (2009). Estimating thresholds to optimal harvest rate for long-
lived, low-fecundity sharks accounting for selectivity and density dependence in
recruitment. *Canadian Journal of Fishery and Aquatic Sciences, 66*, 2062-2080.
doi:10.1139/F09-137

Fowler, S., & Séret, B. (2010). *Shark fins in Europe: Implications for reforming the EU finning*
ban. Retrieved from http://eulasmo.org/blog/shark-fins-europe-ban/
Froese, R. (2004). Keep it simple: three indicators to deal with overfishing. *Fish and*
Fisheries, 5(1), 86-91. doi:10.1111/j.1467-2979.2004.00144.x
Froese, R., Zeller, D., Kleisner, K., & Pauly, D. (2012). What catch data can tell us about the
status of global fisheries. *Marine Biology, 159*, 1283-1292. doi:10.1007/s00227-012-
1909-6
Georgeson, L., Rigby, C. L., Emery, T. J., Fuller, M., Hartog, H., Williams, A. J., . . . Nicol, S. J.
(2020). Ecological risks of demersal fishing on deepwater chondrichthyan
populations in the Southern Indian and South Pacific Oceans. *ICES Journal of Marine*
Science, 77(5), 1711-1727. doi:10.1093/icesjms/fsaa019
Guan, L., Chen, Y., Boenish, R., Jin, X., & Shan, X. (2020). Improving data-limited stock
assessment with sporadic stock index information in stock reduction analysis.
Canadian Journal of Fisheries and Aquatic Sciences, 77, 857-868. doi:10.1139/cjfas-
2018-0500
Guy, C. S., Cox, T. L., Williams, J. R., Brown, C. D., Eckelbecker, R. W., Glassic, H. C., . . .
Siemiantkowski, M. J. (2021). A paradoxical knowledge gap in science for critically
endangered fishes and game fishes during the sixth mass extinction. *Scientific*
Reports, 11, 8447. doi:10.1038/s41598-021-87871-y
Hauck, M. (2008). Rethinking small-scale fisheries compliance. *Marine Policy, 32*, 635-642.
doi:10.1016/j.marpol.2007.11.004
Hausmann, A., Slotow, R., Fraser, I., & Di Minin, E. (2016). Ecotourism marketing alternative
to charismatic megafauna can also support biodiversity conservation. *Animal*
Conservation, 20, 91-100. doi:10.1111/acv.12292
Heupel, M. R., Simpfendorfer, C. A., Espinoza, M., Smoothey, A. F., Tobin, A., & Peddemors,
V. (2015). Conservation challenges of sharks with continental scale migrations.
Frontiers in Marine Science, 2, 1-7. doi:10.3389/fmars.2015.00012
Hilborn, R., Branch, T. A., Ernst, B., Magnusson, A., Minte-Vera, C. V., Scheuerell, M. D., &
Valero, J. L. (2003). State of the world's fisheries. *Annual Review of Environment and*
Resources, 28, 359-399. doi:10.1146/annurev.energy.28.050302.105509
Hobday, A. J., Smith, A. D. M., Stobutzki, I. C., Bulman, C., Daley, R., Dambacher, J. M., . . .
Zhou, S. (2011). Ecological risk assessment for the effects of fishing. *Fisheries*
Research, 108(2-3), 372-384. doi:10.1016/j.fishres.2011.01.013
Hønneland, G. (1999). A model of compliance in fisheries: theoretical foundations and
practical application. *Ocean & Coastal Management, 42*(8), 699-716.
doi:10.1016/S0964-5691(99)00041-1
Hutchinson, M. R., Itano, D. G., Muir, J. A., & Holland, K. N. (2015). Post-release survival of
juvenile silky sharks captured in a tropical tuna purse seine fishery. *Marine Ecology*
Progress Series, 521, 143-154. doi:10.3354/meps11073
Inter-American Tropical Tuna Commission. (2005). *Resolution on the Conservation of Sharks*
Caught in Association with Fisheries in the Eastern Pacific Ocean. (Resolution C-05-
03). Lanzarote, Spain: Inter-American Tropical Tuna Commission 73rd Meeting.
Inter-American Tropical Tuna Commission. (2015). *Resolution on the Conservation of*
Mobulid Rays Caught in Association With Fisheries in the IATTC Convention Area.
(Regulation C-15-04). Guayaquil, Ecuador: Inter-American Tropical Tuna Commission
89th Meeting.

Inter-American Tropical Tuna Commission. (2016). *Resolution on the Management of Shark*
Species. (Resolution C-16-05). La Jolla, California, USA: Inter-American Tropical Tuna
Commission 90th Meeting.
Inter-American Tropical Tuna Commission. (2019). *Conservation of Whale Sharks*.
(Resolution C-19-06). Bilbao, Spain: Inter-American Tropical Tuna Commission 94th
Meeting.
Islam, M. M., Shamsuzzaman, M. M., Mozumder, M. M. H., Xiangmin, X., Ming, Y., & Jewel,
M. A. S. (2017). Exploitation and conservation of coastal and marine fisheries in
Bangladesh: do the fishery laws matter? *Marine Policy, 76*, 143-151.
doi:10.1016/j.marpol.2016.11.026
Juan-Jordá, M. J., Mosqueira, I., Freire, J., & Dulvy, N. K. (2013). Life in 3-D: life history
strategies in tunas, mackerels and bonitos. *Reviews in Fish Biology & Fisheries, 23*,
135-155. doi:doi.org/10.1007/s11160-012-9284-4
Juan-Jordá, M. J., Mosqueira, I., Freire, J., & Dulvy, N. K. (2015). Population declines of tuna
and relatives depend on their speed of life. *Proceedings of the Royal Society B, 282*,
20150322. doi:10.1098/rspb.2015.0322
Juan-Jordá, M. J., Murua, H., Arrizabalaga, H., Dulvy, N. K., & Restrepo, V. (2018). Report
card on ecosystem-based fisheries management in tuna regional fisheries
management organizations. *Fish and Fisheries, 19*, 321-339. doi:10.1111/faf.12256
Khan, A. M. A., Mill, A. C., Gray, T. S., Jiang, M., Arief, H., Brown, A., . . . Polunin, N. V. C.
(2020). Reliability of the data on tuna catches obtained from the dockside in
Indonesia: a study of stakeholders' perceptions. *Marine Policy, 122*, 104242.
doi:10.1016/j.marpol.2020.104242
Kulka, D. W., Anderson, B., Cotton, C. F., Derrick, D., Pacoureau, N., & Dulvy, N. K. (2020).
Leucoraja ocellata. The IUCN Red List of Threatened Species, 2020,
eT161631A124518400. doi:10.2305/IUCN.UK.2020-3.RLTS.T161631A124518400.en
Kumar, U., Helen, A. M., Das, J., Parvez, M. S., Biswas, S. K., & Ray, S. (2019). Unraveling the
hidden truth in a poorly managed ecosystem: the case of discarded species of
conservation interest in Bangladesh industrial marine fisheries. *Regional Studies in*
Marine Science, 32, 100813. doi:10.1016/j.rsma.2019.100813
Kuo, T.-C., Laksanawimol, P., Aylesworth, L., Foster, S. J., & Vincent, A. C. J. (2018). Changes
in the trade of bycatch species corresponding to CITES regulations: the case of dried
seahorse trade in Thailand. *Biodiversity and Conservation, 27*, 3447-3468.
doi:doi.org/10.1007/s10531-018-1610-2
Lack, M., Sant, G., Burgener, M., & Okes, N. (2014). *Development of a rapid management-*
risk assessment method for fish species through its application to sharks: framework
and results. Retrieved from https://www.cms.int/en/document/development-rapid-
management-risk-assessment-method-fish-species-through-its-application-0
Last, P. R., White, W. T., Carvalho, M. R. d., Séret, B., Stehmann, M. F. W., & Naylor, G. J. P.
(2016). *Rays of the World*. Australia: CSIRO Publishing.
Lawson, J. M., Pollom, R. A., Gordon, C. A., Barker, J., Meyers, E. K. M., Zidowitz, H., . . .
Dulvy, N. K. (2020). Extinction risk and conservation of critically endangered angel
sharks in the Eastern Atlantic and Mediterranean Sea. *ICES Journal of Marine*
Science, 77(1), 12-29. doi:10.1093/icesjms/fsz222
Le Quesne, W. J. F., & Jennings, S. (2012). Predicting species vulnerability with minimal data
to support rapid risk assessment of fishing impacts on biodiversity. *Journal of Applied*
Ecology, 49, 20-28. doi:10.1111/j.1365-2664.2011.02087.x

Lee, T. M., & Jetz, W. (2011). Unravelling the structure of species extinction risk for
predictive conservation science. *Proceedings of the Royal Society B, 278*, 1329-1338.
doi:10.1098/rspb.2010.1877
Lucifora, L. O., García, V. B., Menni, R. C., Escalante, A. H., & Hozbor, N. M. (2009). Effects of
body size, age and maturity stage on diet in a large shark: ecological and applied
implications. *Ecological Research, 24*, 109-118. doi:10.1007/s11284-008-0487-z
Martínez-Ortiz, J., Aires-da-Silva, A. M., Lennert-Cody, C. E., & Maunder, M. N. (2015). The
Ecuadorian artisanal fishery for large pelagics: species composition and spatio-
temporal dynamics. *PLoS ONE, 10*(8), e0135136. doi:10.1371/journal.pone.0135136
McClenachan, L., Cooper, A. B., Carpenter, K. E., & Dulvy, N. K. (2011). Extinction risk and
bottlenecks in the conservation of charismatic marine species. *Conservation Letters,
5*, 73-80. doi:10.1111/j.1755-263x.2011.00206.x
Melnychuk, M. C., Peterson, E., Elliott, M., & Hilborn, R. (2017). Fisheries management
impacts on target species status. *Proceedings of the National Academy of Science,
114*(1), 178-183. doi:10.1073/pnas.1609915114
Micheli, F., De Leo, G., Butner, C., Martone, R. G., & Shester, G. (2014). A risk-based
framework for assessing the cumulative impact of multiple fisheries. *Biological
Conservation, 176*, 224-235. doi:10.1016/j.biocon.2014.05.031
Musick, J. A. (1999). Criteria to define extinction risk in marine fishes - The American
Fisheries Society initiative. *Fisheries, 24*(12), 6-14. doi:10.1577/1548-
8446(1999)024<0006:CTDERI>2.0.CO;2
Musick, J. A., Burgess, G., Cailliet, G., Camhi, M., & Fordham, S. (2000). Management of
sharks and their relatives (Elasmobranchii). *Fisheries, 25*(3), 9-13. doi:10.1577/1548-
8446(2000)025<0009:MOSATR>2.0.CO;2
Myers, R. A., & Worm, B. (2005). Extinction, survival or recovery of large predatory fishes.
Philosophical Transactions of the Royal Society B-Biological Sciences, 360(1453), 13-
20. doi:10.1098/rstb.2004.1573
Okes, N., & Sant, G. (2022). Missing sharks: A country review of catch, trade and
management recommendations for CITES-listed shark species. In *Seventy-fourth
meeting of the Standing Committee* (pp. 39). Lyon, France: TRAFFIC.
Oldfield, T. E. E., Outhwaite, W., Goodman, G., & Sant, G. (2012). *Report on assessing the
intrinsic vulnerability of harvested sharks*. Retrieved from Bonn, Germany:
https://www.cms.int/sharks/en/document/assessing-intrinsic-vulnerability-
harvested-sharks-submitted-uk-english-only
Pacoureau, N., Rigby, C. L., Kyne, P. M., Sherley, R. B., Winker, H., Carlson, J. K., . . . Dulvy, N.
K. (2021). Half a century of global decline in oceanic sharks and rays. *Nature, 589*,
567-571. doi:10.1038/s41586-020-03173-9
Pardo, S. A., Cooper, A. B., Reynolds, J. D., & Dulvy, N. K. (2018). Quantifying the known
unknowns: estimating maximum intrinsic rate of population increase in the face of
uncertainty. *ICES Journal of Marine Science, 75*(3), 953-963.
doi:10.1093/icesjms/fsx220
Pardo, S. A., Kindsvater, H. K., Reynolds, J. D., & Dulvy, N. K. (2016). Maximum intrinsic rate
of population increase in sharks, rays, and chimaeras: the importance of survival to
maturity. *Canadian Journal of Fisheries and Aquatic Sciences, 73*(8), 1159-1163.
doi:10.1139/cjfas-2016-0069

Pauly, D., & Zeller, D. (2016). Catch reconstructions reveal that global marine fisheries
1101   catches are higher than reported and declining. *Nature Communications, 7*, 10244.
1102   doi:10.1038/ncomms10244
1103   Pauly, D., Zeller, D., & Palomares, M. L. D. (2021). Sea Around Us Concepts, Design and Data.
1104   Retrieved from http://seaaroundus.org
1105   Pavitt, A., Malsch, K., King, E., Chevalier, A., Kachelriess, D., Vannuccini, S., & Friedman, K.
1106   (2021). *CITES and the sea: Trade in commercially exploited CITES-listed marine
1107   species*. Retrieved from Rome:
1108   Peterson, C. D., Belcher, C. N., Bethea, D. M., Driggers III, W. B., Frazier, B. S., & Latour, R. J.
1109   (2017). Preliminary recovery of coastal sharks in the south-east United States. *Fish
1110   and Fisheries, 18*, 845-859. doi:10.1111/faf.12210
1111   Price, E., Melville-Smith, R., King, D., Green, T., Dixon, W., Lambert, S., & Spencer, T. (2016).
1112   *Measurement of Fisheries Compliance Outcomes: A Preliminary National Study*.
1113   Perth, Australia: Fisheries Research and Development Corporation Retrieved from
1114   http://www.fish.wa.gov.au/Documents/research_reports/frr275.pdf.
1115   RAM Legacy Stock Assessment Database. (2021). *Extended RAM Legacy Stock Assessment
1116   Database version 4.493 (v4.493) [Data set]*. Retrieved from:
1117   https://doi.org/10.5281/zenodo.4458275
1118   Reynolds, J. D., Dulvy, N. K., Goodwin, N. B., & Hutchings, J. A. (2005). Biology of extinction
1119   risk in marine fishes. *Proceedings of the Royal Society B-Biological Sciences,
1120   272*(1579), 2337-2344. doi:10.1098/rspb.2005.3281
1121   Ricard, D., Minto, C., Jensen, O. P., & Baum, J. K. (2012). Examining the knowledge base and
1122   status of commercially exploited marine species with the RAM Legacy Stock
1123   Assessment Database. *Fish and Fisheries, 13*, 380-398. doi:10.1111/j.1467-
1124   2979.2011.00435.x
1125   Rigby, C., & Simpfendorfer, C. A. (2015). Patterns in life history traits of deep-water
1126   chondrichthyans. *Deep Sea Research Part II: Topical Studies in Oceanography, 115*,
1127   30-40. doi:10.1016/j.dsr2.2013.09.004
1128   Rousseau, Y., Watson, R. A., Blanchard, J. L., & Fulton, E. A. (2019). Evolution of global
1129   marine fishing fleets and the response of fished resources. *Proceedings of the
1130   National Academy of Science, 116*(25), 12238-12243. doi:10.1073/pnas.1820344116
1131   Rowat, D., & Brooks, K. S. (2012). A review of the biology, fisheries and conservation of the
1132   whale shark *Rhincodon typus. Journal of Fish Biology, 80*(5), 1019-1056.
1133   doi:10.1111/j.1095-8649.2012.03252.x
1134   Rowlands, G., Brown, J., Soule, B., Boluda, P. T., & Rogers, A. D. (2019). Satellite surveillance
1135   of fishing vessel activity in the Ascension Island Exclusive Economic Zone and Marine
1136   Protected Area. *Marine Policy, 101*, 39-50. doi:10.1016/j.marpol.2018.11.006
1137   Ruano-Chamorro, C., Subida, M. D., & Fernández, M. (2017). Fishers' perception: an
1138   alternative source of information to assess the data-poor benthic small-scale
1139   artisanal fisheries of central Chile. *Ocean and Coastal Management, 146*, 67-76.
1140   doi:10.1016/j.ocecoaman.2017.06.007
1141   Schindler, D. E., Hilborn, R., Chasco, B., Boatright, C. P., Quinn, T. P., Rogers, L. A., &
1142   Webster, M. S. (2010). Population diversity and the portfolio effect in an exploited
1143   species. *Nature, 465*, 609-612. doi:https://doi.org/10.1038/nature09060
1144   Sellas, A. B., Bassos-Hull, K., Perez-Jimenez, J. C., Angulo-Valdes, J. A., Bernal, M. A., &
1145   Hueter, R. E. (2015). Population structure and seasonal migration of the spotted

eagle ray, *Aetobatus narinari. Journal of Heredity, 106*(3), 266-275.
doi:10.1093/jhered/esv011
Simpfendorfer, C. A., & Dulvy, N. K. (2017). Bright spots of sustainable shark fishing. *Current*
Biology, 27, R83-R102. doi:10.1016/j.cub.2016.12.017
Stevens, J. D., Bonfil, R., Dulvy, N. K., & Walker, P. A. (2000). The effects of fishing on sharks,
rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems.
ICES Journal of Marine Science, 57, 476-494. doi:10.1006/jmsc.2000.0724
Stevens, J. D., Walker, T. I., & Simpfendorfer, C. A. (1997). Are southern Australian shark
fisheries sustainable? In D. A. Hancock, D. C. Smith, A. Grant, & J. P. Beumer (Eds.),
Developing and Sustaining World Fisheries Resources. The State of Science and
Management. Melbourne: CSIRO Publishing.
Stobutzki, I., Miller, M., & Brewer, D. (2001). Sustainability of fishery bycatch: a process for
assessing highly diverse and numerous bycatch. *Environmental Conservation, 28*(2),
167-181. doi:10.1017/S0376892901000170
Turner II, B. L., Kasperson, R. E., Matson, P. A., McCarthy, J. J., Corell, R. W., Christensen, L., .
. . Schiller, A. (2003). A framework for vulnerability analysis in sustainability science.
Proceedings of the National Academy of Science, 100(14), 8074-8079.
doi:10.1073/pnas.1231335100
Urquhart, J., Acott, T. G., Symes, D., & Zhao, M. (2014). Introduction: Social Issues in
Sustainable Fisheries Management. In J. Urquhart, T. G. Acott, D. Symes, & M. Zhao
(Eds.), *Social Issues in Sustainable Fisheries Management*. London, U.K.: Springer.
Walker, T. I., Day, R. W., Awruch, C. A., Bell, J. D., Braccini, J. M., Dapp, D. R., . . . Reina, R. D.
(2021). Ecological vulnerability of the chondrichthyan fauna of southern Australia to
the stressors of climate change, fishing and other anthropogenic hazards. *Fish and*
Fisheries, 22, 1105-1135. doi:10.1111/faf.12571
Williams, S. E., Shoo, L. P., Isaac, J. L., Hoffmann, A. A., & Langham, G. (2008). Towards an
integrated framework for assessing the vulnerability of species to climate change.
PLoS Biology, 6(12), e325. doi:10.1371/journal.pbio.0060325
Woodhams, J., Peddemors, V., Braccini, M., Victorian Fisheries Authority, & Lyle, J. (2021).
Gummy shark Mustelus antarcticus (2020). Retrieved from
https://fish.gov.au/report/301-Gummy-Shark-2020
Zhou, S. J., Milton, D. A., & Fry, G. C. (2012). Integrated risk analysis for rare marine species
impacted by fishing: sustainability assessment and population trend modelling. *ICES*
Journal of Marine Science, 69(2), 271-280. doi:10.1093/icesjms/fss009
1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195 **Table 1.** The 21 M-Risk Attributes used for assessments. Universal attributes are indicated
1196 by a (U) and species-specific attributes are indicated by a (SP). For complete explanation and
1197 scoring of each attribute see **supplementary information 1**.

Attribute Class	Question Posed by Attribute
Management System	Is there a regulatory body in place? (U)
	What permits are associated with the fishery? (U)
	Is there a stock status or risk assessment performed for the species being assessed? (SP)
	How is the sustainable fishing level for non-shark and ray target species determined? (U)
	What efforts are in place to reduce incidental catch? (SP)
Fishing Practices and Catch	What is the taxonomic resolution of landing limits? (SP)
	How are finning and removal of other high-value products at sea dealt with? (SP)
	Are there any seasonal closures? (SP)
	Are there any spatial closures? (SP)
	What measures are in place to increase post-release survival? (SP)
Compliance and Enforcement	What is the taxonomic resolution of the catch reported? (SP)
	How is illegal, unregulated, and unreported fishing handled? (U)
	What compliance measures are in place to reduce fishing mortality of non-target species? (SP)
	How is reported catch validated? (U)
	How are regulations enforced? (U)
Country Attributes	Is there a larger amount given for beneficial subsidies than for deleterious subsidies? (U)
	Does the country's NPOA-Sharks address the ten recommended objectives? (U)
	Is the country a member of CITES without reservations on elasmobranch species? (SP)
	Is the country a member of CMS and are they a signatory on the MOU-Sharks? (U)
RFMO Attributes	Are all member countries also members of CITES and CMS with no reservations on elasmobranch species? (SP)
	How well has the RFMO progressed with the implementation of ecosystem-based fisheries management? (U)

1198
1199

1208 **Table 2.** Example of value statements for the attribute that asks, "What is the taxonomic
1209 resolution of the catch reported?"

Score	Value Statement
0	No reporting of elasmobranch catch
1	Catch reporting to broad categories (i.e. "sharks" "elasmobranchs" or "rays") OR list of species listed with no associated numbers of actual catch
2	Catch reporting to narrow categories (i.e. "mackerel sharks," "reef sharks," "deepwater sharks," "whaler sharks," etc.)
3	Catch reporting as similar/related species grouped ("deepwater dogfishes," "gulper sharks," "mako," "hammerhead," etc.)
4	Species-specific catch reporting

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

FIGURE LEGENDS

Figure 1. The goal of M-Risk assessments is to determine relative species risk within and
across fisheries at country and RFMO scales. Whereas Productivity Susceptibility Analysis
(PSA) is applied to a specific assessment unit (i.e., typically the fishery), and hence it is not
typical to make comparisons among PSAs. PSAs identify Vulnerability based on a species'
attributes (Productivity) and its overlap with the fishery being assessed (Susceptibility). M-
Risk also identifies Vulnerability based on a species' attributes (albeit with a different name:
Intrinsic Sensitivity), however, this M-Risk is contingent on its management within the
fishery, not just spatial overlap with the fishery.
1243

1244 **Figure 2.** Intrinsic sensitivity (IS) scores based on generation length and maximum size.
1245 Higher intrinsic sensitivities are assigned to sharks and rays with longer generation lengths
1246 and larger relative sizes. Relative generation length is scored by bins (0-5 years = 0%, 5.1-10
1247 = 20%, 10.1-15 = 40%, 15.1-20 = 60%, 20.1-25 = 80%, >25 = 100%).
1248

Figure 3. All attribute scores, management scores, intrinsic sensitivity scores, and final M-
Risk scores for all species assessed in (a) Ecuador and (b) IATTC. Attribute columns are
arranged from highest average score to lowest average score within the management unit.
Final M-Risk scores are presented as the percentage of the highest score such that the
higher M-Risk scores represent the most at-risk species. No NA values are included in the
figure as only relevant attributes for each management unit are included.
1255

■ White Shark *Carcharodon carcharias*			Intrinsic Sensitivity = 100%	
■ Grey reef shark *Carcharhinus amblyrhynchos*			IS = 39.92%	
■ Scalloped bonnethead *Sphyrna corona*			IS = 17.18%	

0 cm ─── 640 cm

14.4%		39.8%	Relative Max Size	100%

0-5 yrs	5.1-10	10.1-15	15.1-20	20.1-25	> 25
20%	40%				Relative Generation Length 100%

0% ─── 100%

17.2%	39.9%	Intrinsic Sensitivity	100%