Transcriptome profiling of histone acetylation and methylation writers and erasers gene families during mouse spermatogenesis

Candela R. González [1], Gastón Barbero [1], Maximiliano de Sousa Serro[2], Camila Perez Lujan [2], Alfredo D. Vitullo [1], Betina González [2]*

[1] Centro de Estudios Biomédicos Básicos, Aplicados y Desarrollo (CEBBAD), Universidad Maimónides, Ciudad Autónoma de Buenos Aires, Buenos Aires, Argentina.

[2] Instituto de Investigaciones Farmacológicas (Universidad de Buenos Aires–Consejo Nacional de Investigaciones Científicas y Técnicas), Ciudad Autónoma de Buenos Aires, Buenos Aires, Argentina.

* Corresponding author

E-mail: bgonzalez@ffyb.uba.ar

Keywords: **spermatogenesis, epigenetic enzymes, histone post-transcriptional modification**

ABSTRACT

Accumulating evidence points out that sperm carry epigenetic instructions to embryo in the form of

retained histones marks and RNA cargo that can transmit metabolic and behavioral traits to

offspring. However, the mechanisms behind epigenetic inheritance of paternal environment are still

poorly understood. Here, we analyzed the expression profile of all known histone lysine writers and

erasers enzymes across spermatogenesis, unraveling the developmental windows at which they are

upregulated, and the specific activity related to canonical and non-canonical histone marks

deposition and removal. We also characterized the epigenetic enzymes signature in the mature

sperm RNA cargo, showing most of them positive translation at pre-cleavage zygote, suggesting

that paternally-derived enzymes mRNA cooperate with maternal factors to embryo chromatin

assembly. Our study shows important mechanistic aspects behind transgenerational epigenetics,

where epigenetic enzymes not only can respond to environmental stressors, but could function as

vectors of epigenetic information and participate in chromatin organization during maternal-to-

zygote transition.

INTRODUCTION

In the last years, accumulating research has documented that male germ cell maturation presents windows of vulnerability for epigenetic reprogramming by environmental stressors that can affect fertility and even transmit developmental, metabolic and behavioral traits to offspring (Bale, 2015). This type of non-genetic "Lamarckian" transmission has been established for paternal lifestyle and different forms of chronic stress, drug abuse, dietary change, and social defeat, involving changes in non-coding RNAs cargo, DNA methylation, and histone post-translational modifications (PTMs) (Bale, 2105). However, most of the research in epigenetic inheritance has documented the role of non-coding RNAs and DNA methylation patterns, whereas the mechanisms by which the environment can trigger specific changes on histone PTMs are still largely unexplored.

Spermatogenesis is a finely regulated process where unipotent spermatogonia undergo meiosis and, subsequently, spermiogenesis, to become sperm cells. During the developmental stages of spermatogenesis, male germ cells experience dramatic chromatin reorganization, where approximately 90% (human) to 95% (mouse) of histones are evicted and replaced by protamines to compact the paternal genome (Rajender et al., 2011). Importantly, a small fraction of histones in the sperm genome is retained in specific locations and carries several PTMs that play critical roles in epigenetic regulation of spermatogenesis and early embryonic development (Luense et al., 2016). The epigenetic program of histone PTMs that controls spermatogenesis is driven by enzymatic families known as writers and erasers of the so-called "histone code", that modulate chromatin structure and transcription (Kouzarides, 2007). These epigenetic enzymes catalyze the deposition or removal of specific PTMs such as acetylation, methylation, phosphorylation, crotonylation, among others, at specific amino acid residues on the nucleosome's core

histones H2A, H2B, H3, and H4, and also on the linker H1/H5 (Luense et al., 2016). In male germ cells, histone PTMs not only control active vs repressed chromatin states but also a range of processes including DNA replication and repair, chromosome maintenance and histone eviction (Luense et al., 2016). So far, the most studied histone PTMs related to the control of germ cells mitosis, meiosis and spermiogenesis are methylated and acetylated lysines (K) on H2A/B, H3 and H4. Histone K acetylation can be dynamically regulated by the opposing action of acetyltransferases (HATs) and deacetylases (HDACs). Acetylation of K residues neutralizes the positive charge on histones, allowing DNA-binding proteins better access to the DNA and resulting in activation of gene expression as well as histone eviction (Rajender et al., 2011). Unlike acetylation, methylation does not affect histone charge but regulates recognition and interaction with chromatin-binding proteins that control the transcription or respond to DNA damage (Rajender et al., 2011). Histone K methylation is finely regulated by methyltransferases (KMTs) and demethylases (KDMs) that control the mono-, di-, and/or tri-methylation of specific residues, and this can either activate or repress transcription depending on the residue position, the number of methylations and the presence of other methyl or acetyl groups in the vicinity (Mosammaparast and Shi, 2010).

The role of epigenetic enzymes comes to focus, as they are responsible for the histones´ PTMs writing and erasing balance. Moreover, histone-modifying enzymes have been shown to respond to the organism physiology and metabolism, several disease conditions, and environmental stressors (Bale, 2015; Nebbioso et al., 2018). Even though epigenetic enzymes are the bridge linking the environment with epigenetic inheritance through retained histones PTMs, an in-depth analysis of their expression patterns throughout the spermatogenic process is still lacking. Moreover, these enzymes provide

new targets for therapies for numerous diseases (Prachayasittikul et al., 2017) but the

potential effect of these compounds on epigenetic inheritance is generally overlooked. To

further elucidate the mechanisms driving the histone PTMs during spermatogenesis, a

better understanding of the gene expression profiles of epigenetic modifying enzymes in

male germ cells is required. Here, we curated transcriptomic data from spermatogonia to

mature sperm populations and analyzed the expression profile and dynamic changes of all

the known families of histone K acetylation and methylation writers and erasers. We

provide a complete picture of the epigenetic enzymes across spermatogenesis in mice and

the developmental windows when the transcription of these enzymes may be more

susceptible to environmental disruption. Moreover, we analyzed pre-cleavage zygote

translatome, confirming that mRNAs of several histone modifying enzymes carried by

sperm are associated with ribosomes for protein synthesis in 1-cell embryo. Not only do we

expand the knowledge on epigenetic enzymes with recognized roles in spermatogenesis,

but provide evidence of many more whose roles in male germ cell development and zygote

have not been described yet.

RESULTS

Epigenetic enzymes and histone K acetylation and methylation marks distribution during spermatogenic stages

Table 1 shows a curated list of writers and erasers families for histone K acetylation and methylation obtained from the Uniprot database, as proteins with confirmed catalytic activity towards histones K residues, and their most representative K targets reported in mammalian cells. Figure 1A shows the canonical histone methylation sites that are found on H3K4/9/27/36/79 and H4K20, with well-established roles in chromatin functions. For instance, H3K4me1/2/3, H3K36me3, H3K9me1, H3K27me1, H3K79me2/3, and H4K20me1 are found in enhancers, promoters, and gene bodies of active genes, whereas H3K9me2/3, H3K27me2/3 and H4K20me3 are found in—heterochromatin, telomeric regions and inactivated X chromosome (Fig. 1A). Open chromatin is characterized by the presence of acetylated histones at several K residues such as H2AK5/9/13/15ac, H2BK5/12/15/20/23/24ac, H3K4/9/14/18/27/36/56/79/122ac, and H4K5/8/12/16/20/79/91ac. Many of these sites are also targets of mutually exclusive methylation and characteristic of active or repressed chromatin. In addition, there are multiple non-canonical histone methylation sites on H2AZK7, H3K23/63 and H4K5/12 with unknown functions. In Figure 1B we summarized the available information on the most important histone K acetylation and methylation marks distribution during spermatogenic stages, obtained from reported immunohistochemical and/or proteomic studies. We observed two main profiles on the marks distribution across spermatogenesis: H3/4ac, H3K4me2/3 and H3K9me2/3 increase with spermatogonia differentiation and early meiosis, decrease at late meiosis, and regain high levels at spermiogenesis, whereas

H3K36me2/3, H3K27me2/3, H3K79me3 and H4K20me3 show sustained increased

expression, but H4K20me is erased at RStid stage.

Histone K acetylation and methylation enzymes expression during spermatogenesis

To analyze the expression profiles of writers and erasers families for histone K

acetylation and methylation across spermatogenesis, we curated RNA-seq data available in

purified male germ cells from adult mice and selected the dataset GSE162740 (Mayorek et

al., 2022). This dataset contains the mRNA profiles of 6 populations of germ cells isolated

in triplicates: undifferentiated (SGund) and differentiated spermatogonia (SGdiff)

(spermatogonial phase), primary spermatocytes, including pre-leptotene (PreL),

leptotene/zygotene (LZ) and pachytene/diplotene (PD) stages (meiosis), and round

spermatids (RStid) (spermiogenesis phase). We downloaded the FASTQ files and mapped

them to MM10 genome using STAR and featureCounts to obtain the count matrix. We

corroborated the cell stages and purity of the dataset by analyzing spermatogenic and

somatic cell marker genes profiles in each sample (Fig.S1 and Table S1). Figure 2 shows

the hierarchical clustering of z-scores for the 99 epigenetic enzymes listed in Table 1.

Column clustering shows two main profiles between the spermatogonial and early meiotic

vs late meiotic and spermiogenic phases. SGdiff and PreL stages are the most similar,

whereas SGund and LZ show large clusters of enzyme upregulation. The late meiotic and

spermiogenesis genes show two main clusters: genes specifically upregulated at PD and

genes that increase at PD and remain high or even increase expression at RStid stage.

Epigenetic enzymes upregulation across spermatogenesis

To analyze upregulated mRNAs of epigenetic enzymes across male germ cells

differentiation, we performed differential gene expression (DEG) analysis using the

DESeq2 algorithm and compared consecutive stages of cell development: SGdiff vs

SGund, PreL vs SGdiff, LZ vs PreL, PD vs LZ and RStid vs PD. For each analysis, we

selected genes with padj < 0.05 and log2FC > 0.5. Figure 3A shows the number of

upregulated enzymes detected for each comparison, while Figure 3B shows enzymes that

were not found significantly upregulated at any cell stage. These enzymes show a pattern of

high expression at the spermatogonial phase that decreases with cell differentiation. To

evaluate until which stage these enzymes maintain high expression, we analyzed the

downregulated genes at each comparison (padj < 0.05, log2FC < -0.5). The statistical

analysis showed that i) *Kat6a*, *Kat6b*, *Mcm3ap* and *Kmt2b* significantly decreased at PreL

vs SGdiff, ii) *Hdac2*, *Smyd2* and *Kdm8* significantly decreased at LZ vs PreL, iii) *Naa60*,

Ash1l, *Kmt2a*, *Kmt5b* and *Kdm3b* significantly decreased at PD vs LZ, and iv) *Setd2*

significantly decreased at RStid vs PD.

Expression profile of histone K acetylation and methylation enzymes during

spermatogenesis phases

For the analysis of the increased epigenetic enzymes profiles during spermatogonial,

meiosis prophase I and spermiogenesis phases, we constructed heatmaps for the

upregulated enzymes at each developmental cell stage, and dot plots indicating the mRNA

abundance, the writer/eraser nature and target histone mark. During the spermatogonial

phase, the SGund population is characterized by a mitotic activity to maintain testicular

homeostasis through self-renewal. Some of these SGund initiate extensive chromatin

reorganization and their epigenetic landscape shifts dramatically to differentiate to

committed SGdiff (Shirakawa et al., 2013). Figure 4 shows the differential expression of

histone K acetylation and methylation writers and erasers in SGund and SGdiff. Several

histone acetylation writers decrease with the transition of SGund to SGdiff, including

Kat2a, *Kat2b*, *Supt7l* and *Ep300*. The SGund also showed high expression of class I

Hdac8, class II *Hdac4/7/9* and class IV *Hdac11*, whereas SGdiff upregulated class I

Hdac1/3 and class III *Sirt1*. We also found specific expression profiles of enzymes

involved in heterochromatin formation, including increased KDM3/4 family of H3K9me

erasers at SGund, and increased KMT1 H3K9me and KMT5 H4K20me writers families at

SGdiff. Interestingly, the SGund showed high expression of the polycomb H3K27me writer

subunit *Ezh1*, and the transition to SGdiff upregulated the H3K27me eraser *Kdm6b*. The

transition to SGdiff also decreased the levels of several writers and erasers for H3K4me and

H3K36me but specifically upregulated *Prdm9* and *Nsd2* writers. Finally, Figure 2 shows

high expression of *Ehmt2* and *Kdm1a* at the spermatogonial phase, coincident with the high

protein expression described for these enzymes at this stage.

Meiotic entry occurs concomitantly with the pre-meiotic S phase and is followed by

the meiotic prophase that initiates with PreL stage. Here, we found some enzymes that

continue to increase expression from SGdiff to LZ stage, including *Prdm9*, *Setdb2* and

Kmt5a/c, and enzymes that seem to be specific to each meiotic stage, like *Setd6* at PreL

stage, that deposits H2AZK7me1 mark, and *N6amt1* at LZ stage, that deposits H4K12me

mark (Fig. 5). Histone acetylation writers increased from PreL to PD stage, with

upregulated expression of almost all members of GNATs, NATs, SRC, MYST, and

p300/CBP families, and other HATs like *Supt7l*, *Msl3*, *Clock*, *Gtf3c4* and *Brca2*, showing

most of them high RPKM values. During meiotic prophase I, several class I, II and III

HDACs were increased, including *Hdac1/8*, *Hdac4/5/9* and *Sirt2/3/5/6/7*. Additionally,

note that the major enzymatic regulation of active H3K4me and H3K36me marks occurs

during chromosomal synapsis at LZ stage, allowing the crossing over of homologous

chromosomes at hotspots and then at PD stage. Moreover, H3K36me writers *Nsd1/2/3,*

Setd5 and *Prdm9* peak expression from PreL to LZ stage, while PD stage does not show

peak expression of neither H3K36me writers nor erasers. Also, prophase I stages showed

increased expression of KDM5 family and *Kdm1a*, consistent with the decreased

H3K4me2/3 global expression reported at this stage (Fig. 1B). We also detected increased

expression and high RPKM values of H3K27me writers and erasers at PreL and LZ stages,

including polycomb writers *Ezh1/2*, and *Kdm6a*, which form the COMPASS complex with

Kmt2b/c/d and *Ashl2* H3K4me writers increased at these stages. Meiotic prophase I also

showed upregulation of KDM7 family, *Kdm7a* and *Phf2/8*, that mediate gene activation

programs by removal of all histone repressive marks. In addition, there is considerable

enzymatic regulation on H3K9me mark, evidenced in the number and peak expression of

H3K9me writers and erasers during all stages of prophase I, including KDM3/4 families

that would mediate the decreased global H3K9me3 levels detected at this stage (Fig. 1B).

Spermiogenesis is the final phase of spermatogenesis, where the haploid RStids

engage in chromatin remodeling programs involving specific histone variants

incorporation, H3/4 hyperacetylation and subsequent replacement of 90-99% histones by

protamines (PRMs) to compact the nucleus. Figures 6 shows that the RStid stage presents

expression of histone K acetylation writers and erasers that continued to increase

expression from PD stage, including *Kdm4a*, *Msl3*, *Ehmt2*, *Rsbn1* and *Dot1l*, and others

that seemed to reactivate expression from previous spermatogenic and early meiotic stages.

We found several HATs specifically upregulated at RStid, and also *Kat8* that maintained

high levels from PD stage (Fig. 2) consistent with the histone hyperacetylation that takes

place at this stage (Fig. 1B). Also, RStid stage showed high RPKM values for the

H3K79me writer *Dot1l*, the H3K27me writer *Ezh2* and the H3K9me2 writer *Ehmt2*,

consistent with the high global expression of these marks at spermiogenesis stage (Fig. 1B).

Additionally, RStid population showed peak expression of *Rsbn1*, a known specific eraser

involved in decreasing H4K20me levels at spermiogenesis (Fig. 1B).

Epigenetic enzymes detected at mature sperm and 1 cell embryo translatome

To characterize the epigenetic enzymes mRNAs present in the mature sperm, we

analyzed data from three datasets that performed long non-coding and mRNA RNAseq:

GSE81216 (total and head sperm, Schuster et al., 2016), GSE88732 (Zhang et al., 2017),

and E-MTAB-5834 (Gapp et al., 2020). For each study, we obtained the mean RPKM for

each gene and converted it to percent rank to transform values into a 0-1 scale, where 0

means no detection and 1 is the highest detected gene in each study (see Supplemental

Table1). Then, we calculated the mean percent rank for the three studies. Figure 7A shows

a positive correlation between sperm and RStid mean percent rank, which favors the idea

that the sperm enzymes mRNA comes from the spermiogenesis phase. In addition, with the

aim of elucidating if the epigenetic enzymes detected in mature sperm may participate in

embryo chromatin assembly, we analyzed recently reported data that profiled the mRNA

translation landscape in mouse pre-implantation embryos by Ribo-seq (Zhang et al., 2022).

We analyzed the translation efficiency (TE), obtained as mean RPKM detected at

ribosomes/mean RPKM detected at whole RNAseq (RPKM>0.5) (see Supplemental Table

S3). Figure 7B shows a word cloud plot where word size indicates enzyme abundance

(mean percent rank) in mature sperm, and red color indicates positive translation detected at

1 cell embryo. Figure 7C shows that several writers and erasers of all histone acetylation

and methylation marks are detected in the translatome of the 1-cell embryo, where many of

these mRNAs are highly detected in the sperm RNA pool.

DISCUSSION

Accumulating evidence has put on focus the existence of windows of vulnerability

during spermatogenesis, at which environmental stressors can induce epigenetic

reprogramming and the transmission of developmental, metabolic, and behavioral traits to

offspring. To further elucidate the mechanisms driving histone PTMs changes during the

spermatogenic process, we analyzed the epigenetic enzyme expression programs across

germ cell differentiation to mature sperm. We show that cell transition across

spermatogenesis is characterized by the upregulation of specific histone K methylation and

acetylation writers and erasers driving the epigenome changes necessary for spermatogonia

differentiation, meiosis entry and spermiogenesis. Moreover, most of these enzymes are

detected in the mature sperm mRNA pool, and some are also detected in the early zygote

translatome, suggesting they can be delivered and participate in the 1-cell embryo

chromatin assembly. Our study shows important mechanistic aspects behind

transgenerational epigenetics, where epigenetic enzymes not only can respond to

environmental stressors and alter the germ cell epigenome but could function as vectors of

epigenetic information transmission themselves, by participating in chromatin organization

at the maternal to zygote transition.

The spermatogonia stages seem to be vulnerable for stress-induced epigenetic

reprogramming since they reside outside the blood-testis barrier (González et al., 2020).

Furthermore, current data support a dynamic stem cell model in which the fate of SGund

population is context-dependent and plastic, and several epigenetic mechanisms are

implicated in the regulation of spermatogonia maintenance and cell fate (Mäkelä et al.

2019). Here, we found that the transition to the SGdiff population is characterized by

increased expression of writers of activation marks H3K4me and H3K36me and repression

marks H3K9me and H4K20me. At the spermatogonial phase, we found high expression of

H3K4me epigenetic enzymes, including *Kmt2a/b/c* writers and *Kdm1a/b* erasers. In this

context, at spermatogonial stage occurs the deposition of monovalent and bivalent H3K4me

marking at promoters by *Kmt2b* (*Mll2*), preparing germ cells for gene activation at late

spermatogenesis and embryonic development (Tomizawa et al., 2018). Accumulating

evidence also points to H3K4me2 in the maintenance of transcriptional states during cell

development and, its removal by erasers expressed at spermatogonia such as

KDM1A/LSD1 is a key step for epigenetic reprogramming and cell fate (Martinez-Gamero

et al., 2021). KDM1A/LSD1 was found to mediate spermatogonia commitment and

differentiation, as testicular deletion of *Kdm1a* leads to progressive germ cell loss,

downregulation of self-renewal factors and abnormal accumulation of meiotic

spermatocytes (Lambrot et al., 2015; Myrick et al., 2017). Interestingly, overexpression of

Kdm1a altered the H3K4me2/3 levels and the RNA profile of sperm and embryos

transgenerationally (Siklenka et al., 2015; Lismer et al., 2020). Hence, the spermatogonial

phase seems to be especially sensitive to alterations in H3K4me balance linked to

epigenetic inheritance. SGdiff showed a specific increase of *Prdm9*, responsible for

H3K4me3 and H3K36me3 that mediate the double-strand breaks (DSBs) formation at

meiotic hot spots (Sun et al., 2020; Getun et al., 2017). In line with this, it has been shown

that meiosis entry of SGdiff is driven by increased levels of H3K36me3 (Godmann et al.,

2007; Song et al., 2011; Zuo et al., 2018). Here, we found *Setd2* expression until PD stage,

and *Setd5* specifically upregulated at SGund, suggesting their involvement in the specific

H3K36me3 landscape of spermatogonia differentiation. The SGund population also

expressed H3K4me/H3K36me eraser *Kdm2b*, a subunit of non-canonical PRC1.1 complex

that was found to protect the polycomb-silenced promoters against ectopic de novo

H3K36me methylation (Blackledge and Klose, 2021). Furthermore, KDM2B was recently

found to erase H3K79me2/3 and to induce transcriptional repression via SIRT1 chromatin

silencing (Kang et al., 2018).

It is established that SGund typically lacks heterochromatin, which forms as they

differentiate to the spermatogonia population committed to meiosis (Chiarini-Garcia et al.,

2002). The SGund showed increased expression of H3K9me erasers *Kdm3a/b* and *Jmjd1c*

(*Kdm3c*), that were found to counteract the enzymatic activity of EHMT2 (Tachibana et al.,

2007), which is highly expressed in the spermatogonia population. Methylation of H3K9 by

EHMT2 blocks gene expression of the stem cell factors OCT4 and NANOG, therefore,

demethylation of H3K9 by KDM3 family enzymes may be a key step in the maintenance of

self-renewal of spermatogonial stem cells (Kuroki et al., 2020; Chioccarelli et al., 2020). In

agreement, we found increased *Ehmt1*, *Suv39h2* and *Setdb2* expression at SGdiff stage,

consistent with H3K9me2/3 deposition during spermatogonia differentiation. In particular,

EHMT1 was found in a complex that binds E2F- and Myc-Responsive genes to repress the

mitotic program (Ogawa et al., 2002). Moreover, SGdiff population showed decreased

Ezh1 and increased *Kdm6b* compared to SGund, suggesting that spermatogonia

differentiation involves H3K27me removal. We also observed that the transition to the

SGdiff population is characterized by specific HDACs upregulation. In this sense, it was

shown that class IIa HDACs enzymatic activity depends on their recruitment into a

complex containing class I HDAC3 and retinoid acid (RA) receptor (RAR), suggesting that

these HDACs could mediate chromatin silencing induced by RA, essential for the transition

of SGund into SGdiff (Parra, 2015). Moreover, SGdiff population showed increased *Sirt1,*

described as a key regulator of spermatogonia differentiation since SIRT1-deficient mice

show a delay of pre-meiotic differentiation, aberrant expression of spermatogenic genes,

abnormal spermatozoa with elevated DNA damage, and reduced fertility (Coussens et al.,

2008; Bell et al., 2014). Among histone acetylation writers, *Kat2b* expressed in SGund

functions as a co-activator for RAR and can interact with NCOA1 to promote transcription

(Spencer et al., 1997), which is upregulated from SGdiff to LZ stage. Also, we observed

peak expression of *Kat2a* in SGund*,* its germ-cell-specific knockout results in abnormal

chromatin dynamics, leading to increased sperm histone retention and severe reproductive

phenotype (Luense et al., 2019). Therefore, the action of certain epigenetic enzymes in pre-

meiotic stages seems to enable the post-meiotic transcriptome and chromatin remodeling

processes.

Initiation of male meiosis is one of the most important events that coincide with

spermatocyte differentiation, and meiotic prophase is accompanied by several alterations of

epigenetic and gene expression programs for post-meiotic spermiogenesis (Maezawa et al.,

2020). Histone acetylation writers increase during prophase I progression, and

H3K9/18/23ac and H4K5/8/12/16/91ac marks have been associated with open chromatin

and hot spot cores (Getun et al., 2017; Chioccarelli et al., 2020). Accordingly, MYST

enzymes were previously found strongly related to meiosis control, increasing their

expression from early pachytene through diplotene stages (Getun et al., 2017).

Interestingly, p300/CBP family and KAT2B were found to regulate histone eviction by

acetylation of transition nuclear protein (TNP) 2, affecting DNA condensation properties

and interaction with histone chaperones (Pradeepa et al., 2009). Among other acetylation

writers upregulated during prophase I, MSL3 binds to H3K36me3 marked sites and

deposits H4K16ac, controlling meiosis entry and STRA8 (stimulated by retinoic acid 8)

functions (McCarthy et al., 2022). On the other hand, we detected several SIRT enzymes

throughout prophase I. Deacetylation of H3K9 by SIRT6 modulates telomeric chromatin

function (Michishita et al., 2008) while SIRT7 is highly selective toward H3K18ac, and

might play an upstream role in DNA repair and telomere maintenance (Wu et al., 2018).

We also detected peak expression of *Hdac1* at the PD stage that was shown to form a

complex with DNA methyltransferase DNMT3L to regulate X chromosome compaction

(Deplus et al., 2002; Zamudio et al., 2011).

During prophase I, recombination hotspots are mainly marked by H3K4me3 catalyzed

by KMT2 family and PRDM9 methyltransferases (Sollier et al., 2004; Buard et al., 2009).

Moreover, the levels of H3K4me3 and H3K36me3 are highly correlated at hotspots, but

mutually exclusive elsewhere, and PRDM9 is capable of placing both marks on the same

nucleosomes in vivo (Powers et al., 2016). It was shown that *Prdm9* knockout changes the

distribution of DSBs across the genome inducing defective synapses and male infertility

(Paigen and Petkov, 2018; Bhattacharyya et al., 2019). We also detected peak expression of

H3K79me writer *Dot1L* at PD stage*, consistent with previous reports showing increased

levels of DOT1L, and H3K79me2/3 from pachytene onwards (Ontoso et al., 2014). The

heterochromatic centromeric regions and the sex body are enriched for H3K79me3, while

H3K79me2 is present all over the chromatin, but is largely excluded from the sex body

despite the accumulation of DOT1L (Ontoso et al., 2014). Repressive histone methylation

marks H3K27me3, H3K9me3 and H4K20me3 are present in chromatin regions with

reduced recombination or involved in heterochromatin formation. *In vitro* and *in vivo*

models disrupting the action of H3K9me writers *Suv39h1, Setdb1/2 and Ehmt2* detected

during prophase I, showed an abnormal distribution of H3K9me3 at pericentromeric

heterochromatin, compromised meiotic silencing of unsynapsed chromatin (MSUC),

misregulation of meiotic and somatic genes, anomalous synapsis and misssegregation of

chromosomes and apoptosis at pachytene stage (Peters et al., 2001;, Tachibana et al., 2007;

Takada et al., 2011). Interestingly, low protein diet in the father was linked to altered levels

of H3K9me2 through EHMT2, that changed tRNAs in sperm and transmitted metabolic

phenotypes to the offspring (Yoshida et al., 2020). Also, SETDB1 haploinsufficiency in

mice was shown to induce changes in DNA methylation in transposable elements, and to

influence coat color in the offspring, further linking epigenetic inheritance with altered

epigenetic enzymes levels during spermatogenesis (Daxinger et al., 2016). A recent work

by Barral et al., 2022 reported dual regions in mouse embryonic stem cells that rely on the

SETDB1 and NSD proteins to generate H3K9me3 and H3K36me3, respectively. They

found that SETDB1 removal induces loss of both marks in dual regions, gains signatures of

active enhancers, and comes into contact with upregulated genes, providing a mechanistic

insight by which genes are controlled by heterochromatin (Barral et al., 2022). Moreover,

NSD1-mediated H3K36me2 was found to prevent H3K27me3 deposition by PRC2,

modulating PRC2-mediated H3K27me domains demarcation (Blackledge and Klose,

2021). In line with this, we detected the expression of H3K27me writers *Ezh1/2* at PreL and

LZ stages that were found involved in mammalian X chromosome inactivation

(Schuettengruber et al., 2007). Furthermore, *Kdm6a* is part of the COMPASS complex with

Kmt2b/c/d and *Ashl2* that activate genes in response to RA by H3K4me deposition and

H3K27me removal (Lavery et al., 2020). On the contrary, the loss of H3K27me3 from

bivalent signature in the promoter of endonuclease SPO11 induces chromatin activation

that favors meiotic entry (Hammoud et al., 2014). At PreL stage, we detected greater

expression of H3K27me erasers than writers, including the expression of *Kdm6a*, which

alteration in its expression induce defects that persisted transgenerationally (Siklenka et al.,

2015, Lesch et al., 2019). We also detected high expression of H2AZK7me1 writer *Setd6* at

PreL stage, which in ESC was found with H3K27me3 close to differentiation marker genes

and removed upon RA signal (Binda et al., 2013), suggesting that SETD6 may have a role

at early male meiosis. Finally, we observed expression of the H4K12me writer *N6amt1* at

LZ stage, a recently described epigenetic mark found at promoters of genes encoding cell

cycle regulators (Metzger et al., 2019).

Chromatin organization dramatically changes during mid- to late-spermiogenesis, due

to histone eviction and replacing by PRMs to facilitate the condensation and packaging of

the paternal genome. In mice, 1%-10% of histones are retained in sperm chromatin and

form a heterogeneous mixture of nucleo-histones and nucleo-protamines (Rajender et al.,

2011; Luense et al., 2016). At the spermiogenesis phase, histone acetylation is essential in

destabilization and remodeling of nucleosomes. Here, the RStid population showed high

transcriptional levels of H4K16ac writers *Msl3* and *Kat8*, a histone PTM essential for

histone eviction which does not occur in the absence of this specific mark (Lu et al., 2010).

Also, *p300/CBP* and *Kat2b* keep their expression from spermatocyte stage, due to their

crucial role to regulate histone eviction (Pradeepa et al., 2009). Moreover, KAT2B was

found in spermatids and involved in H3K9ac, an epigenetic mark detected at unmethylated

active genes / enhancers (Hammoud et al., 2014) that could influence gene expression

directly after fertilization (Steilmann et al., 2011). Additionally, *Hat1* expression was found

related to the incorporation of H4K5/12ac and H3.3 variant at DSBs sites and the

promotion of DNA repair (Yang et al., 2013). H4K5/8/12 marks have been also observed

just before eviction of histones during spermiogenesis (Meistrich et al, 1992; Hecht et al.,

2009) and moreover, H4K8/12ac was detected prior to full decondensation of the sperm

nucleus, suggesting that these marks are transmitted to the zygote (van der Heijden et al.,

2006).

Multiple histone methylation have been also identified in spermatids, pointing to a

balance of "opened" and "closed" chromatin regions during the histone-to-protamine

transition (Wang et al., 2019). Here, we detected upregulated expression of H3K79me

writer *Dotl1* and H3K9me writer *Ehtm2* at RStid stage. It was found that H3K79me writer

DOTL1 is enriched in post-meiotic stages of mouse germ cells and precedes the histone-to-

protamine transition (Dottermusch-Heidel et al., 2014). During spermiogenesis, there is a

reactivation of transcription from MSUC and MSCI sites that is enabled by the deposition

of histone crotonylation (Kcr). It was proposed that CDYL, an HKcr eraser, prevents post-

meiotic chromatin reactivation by binding to H3K9me3 and H3K27me2/3 marked sites and

facilitating H3K9me2 deposition by EHMT2 (Mulligan et al., 2008). Moreover, there is

specific H3K27me deposition to establish bivalency at developmental genes (Maezawa et

al., 2018), and proteomic studies found substantial H3K27me3/H3K36me2 double marking

in RStid and sperm (Luense et al., 2016).

It is established that mature sperm transport a cargo of miRNAs, tsRNAs, lncRNAs,

circRNAs, and protein-coding mRNAs, that carry an epigenetic blueprint involved in early

embryo development (Guo et al., 2017). Some of these RNAs remain from the last stages of

elongated spermatids, and others are acquired along the passage through the epididymis, as

sperm absorb epididymosomes released by somatic cells (Trigg et al., 2019). Here, we

found that the mature sperm transport a signature of epigenetic enzymes mRNAs similar to

the one detected at spermiogenesis phase, and most of these enzymes show positive

translation at 1 cell embryo. After fertilization, the zygote genome is transcriptionally

silent, and cellular processes are carried on with inherited RNAs and proteins until the

onset of zygotic genome activation (ZGA) around the 2-4 cells stage (Schulz and Harrison,

2019). Also, at 1 cell stage there is significant removal of some maternal RNAs and

specific translation of others, including RNAs specifically delivered by the sperm like the

egg-activating factor PLC-zeta (Sone et al., 2005; Yao et al., 2010). The parental pronuclei

have asymmetric reprogramming capacities and the reprogramming factors reside

predominantly in the male pronucleus. Here, we detected high sperm values and embryo

translation of several enzymes involved in histone acetylation and methylation, including

the H4K20me2/3 writer *Kmt5c*, the H3K9me2 writers *Ehmt1/2,* and the H3K9me3 writers

Suv39h2 and *Setdb1/2,* with established roles in chromatin structure organization. In this

context, it was found that EHMT2 activates soon after fertilization and deposit H3K9me2

patterns in the paternal genome (Ma et al., 2015). Also, previous work suggested that

KMT5C deposits H4K20me3 to allow the timely and coordinated progression of replication

after fertilization (Eid et al., 2016).

In summary, the analysis performed here shows important windows during

spermatogenesis, where interference with epigenetic enzymes gene expression may have

phenotypic consequences in the offspring, even when they are not inherited. Moreover, we

show that epigenetic enzymes mRNA could become functional in pre-cleavage zygotes and

contribute to early chromatin organization with deep implications in future embryo

development. The epigenetic enzyme's paternal contribution could be another mechanism

for epigenetic inheritance that deserves further consideration.

METHODS

Data retrieval

FASTQ files were downloaded from NCBI GEO and ArrayExpress databases.

Germ cells data transcriptomic was obtained from GSE162740, that used crossed Stra8-cre

mice with CAG-LSLtdTomato mice to generate "Stra8-Tom" mice. The gradient

expression of the transgene under Stra8 expression profile, combined with immunostaining

for cKit to further discriminate the SGund and SGdiff spermatogonia, was used to FACS

sort 6 populations of germ cells in triplicates, and sequence in an Illumina HiSeq 2500

platform. Sperm cells data were obtained from: i) GSE81216, that sampled 2 total sperm

and 2 sperm heads from C57BL/6J (JAX) mice using Ion Torrent PGM platform, ii) GSE,

that sampled 4 mature sperm from adult C57BL/6J mice using Illumina HiSeq 4000

platform, and iii) E-MTAB-5834, that sampled 4 control mature sperm from adult

C57BL/6J mice using Illumina HiSeq 2500 platform. One cell embryo, 2 total and 2

ribosome-bound RNA RPKMs were obtained from GSE169632.

Data processing

FASTQ files were processed with the following pipeline: quality control with

FASTQC, trimming with fastp, mapping with STAR and GRCm39 (MM10

gencode.vM29.annotation), and counting with featureCounts. Processed data was

manipulated using Rstudio (v1.2.1033) and Tidyverse packages Dplyr and ggplot2

(Wickham et al., 2019). Counts were converted to RPKM using mean gene length extracted

from the gencode.vM29 file. The count matrix was converted to z score by gene and

performed hierarchical clustering with heatmap package from R base. The sperm RPKMs

obtained for the three selected datasets were converted to percentile rank, and the sperm

mean percentile rank calculated. Word cloud plot was performed with ggwordcloud

package. The embryo traslational eficiency (TE) was calculated as Ribo-seq RPKM / total

RPKM for each gene, with RPKM > 0.5. All the plots shown in this study can be

reproduced by downloading the counts tables, enzymes metadata table and R scripts

available at https://github.com/Gonzalez-Lab/Gonzalez-2022-germ-cells.

Author contributions

B.G and C.R.G conceived the project, analyzed the data and wrote the manuscript. G.B,

M.S.F and C.P.L contributed to data mining and analysis. A.D.V contributed to the critical

reading and editing of the manuscript. All authors read and approved the submitted version.

Declaration of interests

The authors declare no competing interests.

Table 1. Writers and erasers families for histone K acetylation and methylation and the targets described.

Acetylation						Methylation					
WRITERS			**ERASERS**			**WRITERS**			**ERASERS**		
FAMILY	NAME	TARGET	FAMILY	NAME	TARGET	FAMILY	NAME	TARGET	FAMILY	NAME	TARGET
GNAT	KAT2A - GCN5	H3ac	Class I	HDAC1	Hac	KMT1	KMT1A - SUV39H1	H3K9me3	KDM1	KDM1A - LSD1	H3K4me1/2
	KAT2B - PCAF	H3ac		HDAC2	Hac		KMT1B - SUV39H2	H3K9me3			H3K9me1/2
	HAT1	H4ac		HDAC3	Hac		KMT1C - EHMT2 - G9a	H3K9me1/2		KDM1B - LSD2	H3K4me1/2
	ATF2	H4/2Bac		HDAC8	Hac		KMT1D - EHMT1 - GLP	H3K9me1/2	KDM2	KDM2A - JHDM1A	H3K36me2
NAT	NAA40 - NAT11	H4ac	Class IIa	HDAC4	Hac		KMT1E - SETDB1	H3K9me3			H3K4me3
	NAA50 - NAT13	H4ac		HDAC5	Hac		KMT1F - SETDB2	H3K9me3		KDM2B - JHDM1B	H3K36me2
	NAA60 - NAT15	H4K79/91ac		HDAC7	Hac	KMT2	KMT2A - MLL1	H3K4me1/2/3			H3K79me2/3
MYST	KAT5 - TIP60	H4/ H2AZac		HDAC9	Hac		KMT2B - MLL2	H3K4me1/2/3	KDM3	KDM3A - JMJD1A - JHDM2A	H3K9me1/2
	KAT8 - MYST1 - MOF	H4K16ac	Class III	SIRT1	H4K16ac		KMT2C - MLL3	H3K4me1		KDM3B - JMJD1B	H3K9me1/2
	KAT7 - MYST2 - HBO1	H3/4ac		SIRT2	H4K16ac		KMT2D - MLL4	H3K4me1		KDM3C - JMJD1C	H3K9me1/2
	KAT6A - MYST3 - MOZ	H3/4ac		SIRT3	H4K16ac		KMT2F - SETD1A	H3K4me1/2/3	KDM4	KDM4A - JMJD2A - JHDM3A	H3K9me2/3
	KAT6B - MYST4 - MORF	H3ac		SIRT5	Hac		KMT2G - SETD1B	H3K4me1/2/3			H3K36me2/3
p300/CBP	KAT3A - CREBBP	H3ac		SIRT6	H3K9ac		KMT2H - ASH1L	H3K36me2		KDM4B - JMJD2B - JHDM3B	H3K9me3
	KAT3B - EP300	H3ac		SIRT7	H3K18ac	KMT2 core	ASH2L	H3K4me1/2/3			H3K36me3
SRC	NCOA1	H3/4ac	Class IV	HDAC11	Hac	KMT3	KMT3A - SETD2	H3K36me3		KDM4C - JMJD2C - JHDM3C	H3K9me3
	NCOA3	H3ac					KMT3B - NSD1	H3K36me2			H3K36me3
Other	CLOCK	H3ac					KMT3C - SMYD2	H3K4me1/2/3		KDM4D - JMJD2D - JHDM3E	H3K9me2/3
	MSL3	H4K16ac						H3K36me2	KDM5	KDM5A - JARID1A	H3K4me2/3
	GTF3C4	H3ac					KMT3D - SMYD1	H3K4me1/2/3		KDM5B - JARID1B	H3K4me1/2/3
	BRCA2	H3ac					KMT3E - SMYD3	H3K4me2/3		KDM5C - JARID1C	H3K4me2/3
	MCM3AP	H3ac					KMT3G - NSD2 - WHSC1	H3K36me2		KDM5D - JARID1D	H3K4me2/3
	SUPT7L	H3ac					KMT3F - NSD3 - WHSC1L1	H3K36me2	KDM6	KDM6A - UTX	H3K27me2/3
						KMT4	DOT1L	H3K79me1/2/3		KDM6B - JMJD3	H3K27me2/3
						KMT5	KMT5A - SETD8	H4K20me1	KDM7	KDM7A - JHDM1D	H3K9me2
							KMT5B - SUV420H1	H4K20me1/2/3			H3K27me2
							KMT5C - SUV420H2	H4K20me1/2/3			H3K9me1/2
						KMT6	KMT6A - EZH2	H3K27me1/2/3		KDM7B - JHDM1F - PHF8	H3K27me2
							KMT6B - EZH1	H3K27me1/2/3			H4K20me1
						KMT7	KMT7 - SETD7	H3K4me1		KDM7C- JHDM1E - PHF2	H3K9me2
						KMT8	KMT8B - PRDM9	H3K4me1/2/3			H4K20me3
								H3K36me1/2/3	KDM8	KDM8 - JMJD5	H3K9me1
							KMT8A - PRDM2	H3K9me1			H3K36me1/2
							KMT8D - PRDM8	H3K9me2	KDM9	KDM9 - RSBN1	H4K20me2/3
							KMT8C - PRDM6	H4K20me			
						KMT9	KMT9 - N6AMT1- HEMK2	H4K12me1			
						SMYD	SMYD4	H3K4me1/2/3			
							SMYD5	H4K20me3			
						SETD	SETD4	H3K4me1/2			
								H4K20me3			
							SETD5	H3K36me3			
							SETD6	H2AZK7me			
							SETMAR	H3K36me2			
						PRDM	PRDM15	H3K9me1			

The HATs can be divided into six major families: 1) the GNATs (GCN5-related N-acetyltransferases), 2) the NATs (N-terminal acetylases, which also possess a GNAT domain), 3) the MYST, 4) the p300/CBP, 5) the SRC steroid receptors coactivators, and 6) other HATs (Wapenaar and Dekker, 2016). HATs actions are counteracted by the HDACs, which include zinc-dependent class I, class IIa, and class IV HDACs, and NAD-dependent class III Sirtuins (Wapenaar and Dekker, 2016). On the other hand, the KMTs present two domains with annotated lysine methyltransferase activity: 1) SET domains, that catalyze all the canonical methylation sites except for H3K79, and 2) 7βS domains (DOT1L and N6AMT1) (Mosammaparast and Shi, 2010). The KDMs can be divided according to their catalytic activity in: 1) amine oxidases (KDM1), and 2) Jumonji C demethylases (KDM2/3/4/5/6/7/8) (Mosammaparast and Shi, 2010).

501
Figure 1. Canonical histone K marks and writers/erasers families distribution during spermatogenesis. A) Histone K methylation PTMs distribution at DNA elements and functional roles in gene expression for an active gene (pink shading) or inactive gene (gray shading). TSS: transcription start site. Only K modifications with sufficient and consensual information about distribution across genes are shown. B) Main histone K PTMs levels involved in chromatin remodeling during spermatogenesis. The data presented is summarized from the following references: Song et al., 2011; Godmann et al., 2007; Zuo et al., 2018; Luense et al., 2016; Wang et al., 2021). SGund: spermatogonia Thy1+/Kit-, SGdiff: spermatogonia Kit+, PreL: pre-leptotene spermatocytes, LZ: leptotene/zygotene spermatocytes, PD: pachytene/diplotene spermatocytes, RStid: round spermatid.

516 **Figure 2. Unsupervised hierarchical clustering analysis of male germ cells transcriptome**.
517 Heatmap showing z-score values of histone K acetylation and methylation writer/eraser enzymes.
518 SGund: spermatogonia Kit-, SGdiff: spermatogonia Kit+, PreL: pre-leptotene spermatocytes, LZ:
519 leptotene/zygotene spermatocytes, PD: pachytene/diplotene spermatocytes, RStid: round spermatid.
520

521
Figure 3. Histone K acetylation and methylation epigenetic enzymes expression during
spermatogenesis A) Number of upregulated epigenetic enzymes for each germ cell
stage.Differentiall gene expression analysis was conducted with DESeq2 on consecutive
populations across differentiationn (SGdiff vs SGund, PreL vs SGdiff, LZ vs PreL, PD vs LZ and
RStid vs PD), and upregulated genes selected with padj<0.05 and log2FC>0.5. B) Heatmap
showing epigenetic enzymes not significantly upregulated at any germ cell stage across
spermatogenesis. SGund: spermatogonia Kit-, SGdiff: spermatogonia Kit+, PreL: pre-leptotene
spermatocytes, LZ: leptotene/zygotene spermatocytes, PD:pachytenee/diplotene spermatocytes,
RStid: round spermatid.
531
532

Figure 4. Epigenetic histone K writers and erasers upregulation at spermatogonial phase. The left panel indicates the heatmap showing z score values and the right panel indicates the Dot plot showing enzymes RPKM values in SGund (A) and SGdiff (B) population. Blue dots: writers, red dots: erasers. SGund: spermatogonia Kit-, SGdiff: spermatogonia Kit+.

Figure 5. Epigenetic histone K writers and erasers upregulation at meiosis prophase I. The left panel indicates the heatmap showing z score values and the right panel indicates the Dot plot showing enzymes RPKM values, in PreL (A), LZ (B) and PD (C) spermatocytes. Blue dots: writers, red dots: erasers. PreL: pre-leptotene spermatocytes, LZ: leptotene/zygotene spermatocytes, PD:pachytenee/diplotene spermatocytes.

547 **Figure 6. Epigenetic histone K writers and erasers upregulation at spermiogenesis.** The left

548 panel indicates the heatmap showing z score values and the right panel indicates the Dot plot

549 showing enzymes RPKM values in RStid. Blue dots: writers, red dots: erasers. RStid: round

550 spermatid, PD: pachytene/diplotene spermatocytes.

Figure 7. Epigenetic histone K writers and erasers detected at mature sperm RNA cargo and
1-cell embryo translatome. A) Pearson correlation between epigenetic enzymes mean percent
rank detected at RStid and sperm. B) Mature sperm word cloud for epigenetic enzymes
indicating the enzyme abundance (word size = mean percent rank) and translation
efficiency (TE) detected at 1 cell embryo (red: TE > 0, black: TE=0). C) Epigenetic
enzymes TE in mouse pre-implantation embryos (TE = RNA RPKM at ribosome / total
RNA RPKM, RPKM>0.5)

REFERENCES

1. Bale T. L. (2015). Epigenetic and transgenerational reprogramming of brain development. *Nature reviews. Neuroscience*, 16(6), 332–344. https://doi.org/10.1038/nrn3818

2. Barral, A., Pozo, G., Ducrot, L., Papadopoulos, G. L., Sauzet, S., Oldfield, A. J., Cavalli, G., & Déjardin, J. (2022). SETDB1/NSD-dependent H3K9me3/H3K36me3 dual heterochromatin maintains gene expression profiles by bookmarking poised enhancers. Molecular cell, 82(4), 816–832.e12. https://doi.org/10.1016/j.molcel.2021.12.037

3. Bell, E. L., Nagamori, I., Williams, E. O., Del Rosario, A. M., Bryson, B. D., Watson, N., White, F. M., Sassone-Corsi, P., & Guarente, L. (2014). SirT1 is required in the male germ cell for differentiation and fecundity in mice. *Development (Cambridge, England)*, 141(18), 3495–3504. https://doi.org/10.1242/dev.110627

4. Bhattacharyya, T., Walker, M., Powers, N. R., Brunton, C., Fine, A. D., Petkov, P. M., & Handel, M. A. (2019). Prdm9 and Meiotic Cohesin Proteins Cooperatively Promote DNA Double-Strand Break Formation in Mammalian Spermatocytes. Current biology : CB, 29(6), 1002–1018.e7. https://doi.org/10.1016/j.cub.2019.02.007

5. Binda, O., Sevilla, A., LeRoy, G., Lemischka, I. R., Garcia, B. A., & Richard, S. (2013). SETD6 monomethylates H2AZ on lysine 7 and is required for the maintenance of embryonic stem cell self-renewal. *Epigenetics*, 8(2), 177–183. https://doi.org/10.4161/epi.23416

6. Blackledge, N. P., & Klose, R. J. (2021). The molecular principles of gene regulation by Polycomb repressive complexes. Nature reviews. Molecular cell biology, 22(12), 815–833. https://doi.org/10.1038/s41580-021-00398-y

7. Buard, J., Barthès, P., Grey, C., & de Massy, B. (2009). Distinct histone modifications define initiation and repair of meiotic recombination in the mouse. The EMBO journal, 28(17), 2616–2624. https://doi.org/10.1038/emboj.2009.207

8. Chiarini-Garcia, H., & Russell, L. D. (2002). Characterization of mouse spermatogonia by transmission electron microscopy. Reproduction (Cambridge, England), 123(4), 567–577. https://doi.org/10.1530/rep.0.1230567

9. Chioccarelli, T., Pierantoni, R., Manfrevola, F., Porreca, V., Fasano, S., Chianese, R., & Cobellis, G. (2020). Histone Post-Translational Modifications and CircRNAs in Mouse and Human Spermatozoa: Potential Epigenetic Marks to Assess Human Sperm Quality. Journal of clinical medicine, 9(3), 640. https://doi.org/10.3390/jcm9030640

10. Coussens, M., Maresh, J. G., Yanagimachi, R., Maeda, G., & Allsopp, R. (2008). Sirt1 deficiency attenuates spermatogenesis and germ cell function. PloS one, 3(2), e1571. https://doi.org/10.1371/journal.pone.0001571

11. Daxinger, L., Oey, H., Isbel, L., Whitelaw, N. C., Youngson, N. A., Spurling, A., Vonk, K. K., & Whitelaw, E. (2016). Hypomethylation of ERVs in the sperm of mice haploinsufficient for the histone methyltransferase Setdb1 correlates with a paternal effect on phenotype. Scientific reports, 6, 25004. https://doi.org/10.1038/srep25004

12. Deplus, R., Brenner, C., Burgers, W. A., Putmans, P., Kouzarides, T., de Launoit, Y., & Fuks, F. (2002). Dnmt3L is a transcriptional repressor that recruits histone deacetylase. Nucleic acids research, 30(17), 3831–3838. https://doi.org/10.1093/nar/gkf509

13. Dottermusch-Heidel, C., Klaus, E. S., Gonzalez, N. H., Bhushan, S., Meinhardt, A., Bergmann, M., Renkawitz-Pohl, R., Rathke, C., & Steger, K. (2014). H3K79 methylation directly precedes the histone-to-protamine transition in mammalian spermatids and is sensitive to bacterial infections. Andrology, 2(5), 655–665. https://doi.org/10.1111/j.2047-2927.2014.00248.x

14. Eid, A., Rodriguez-Terrones, D., Burton, A., & Torres-Padilla, M. E. (2016). SUV4-20 activity in the preimplantation mouse embryo controls timely replication. Genes & development, 30(22), 2513–2526. https://doi.org/10.1101/gad.288969.116

15. Gapp, K., van Steenwyk, G., Germain, P. L., Matsushima, W., Rudolph, K., Manuella, F., Roszkowski, M., Vernaz, G., Ghosh, T., Pelczar, P., Mansuy, I. M., & Miska, E. A. (2020). Alterations in sperm long RNA contribute to the epigenetic inheritance of the effects of postnatal trauma. Molecular psychiatry, 25(9), 2162–2174. https://doi.org/10.1038/s41380-018-0271-6

16. Getun, I. V., Wu, Z., Fallahi, M., Ouizem, S., Liu, Q., Li, W., Costi, R., Roush, W. R., Cleveland, J. L., & Bois, P. (2017). Functional Roles of Acetylated Histone Marks at Mouse Meiotic Recombination Hot Spots. *Molecular and cellular biology*, *37*(3), e00942-15. https://doi.org/10.1128/MCB.00942-15

17. Godmann, M., Auger, V., Ferraroni-Aguiar, V., Di Sauro, A., Sette, C., Behr, R., & Kimmins, S. (2007). Dynamic regulation of histone H3 methylation at lysine 4 in mammalian spermatogenesis. *Biology of reproduction*, *77*(5), 754–764. https://doi.org/10.1095/biolreprod.107.062265

18. González, B., Gancedo, S. N., Garazatua, S., Roldán, E., Vitullo, A. D., & González, C. R. (2020). Dopamine Receptor D1 Contributes to Cocaine Epigenetic Reprogramming of Histone Modifications in Male Germ Cells. Frontiers in cell and developmental biology, 8, 216. https://doi.org/10.3389/fcell.2020.00216

19. Guo, L., Chao, S. B., Xiao, L., Wang, Z. B., Meng, T. G., Li, Y. Y., Han, Z. M., Ouyang, Y. C., Hou, Y., Sun, Q. Y., & Ou, X. H. (2017). Sperm-carried RNAs play critical roles in mouse embryonic development. Oncotarget, 8(40), 67394–67405. https://doi.org/10.18632/oncotarget.18672

20. Hammoud, S. S., Low, D. H., Yi, C., Carrell, D. T., Guccione, E., & Cairns, B. R. (2014). Chromatin and transcription transitions of mammalian adult germline stem cells and spermatogenesis. Cell stem cell, 15(2), 239–253. https://doi.org/10.1016/j.stem.2014.04.006

21. Hecht, N., Behr, R., Hild, A., Bergmann, M., Weidner, W., & Steger, K. (2009). The common marmoset (Callithrix jacchus) as a model for histone and protamine expression during human spermatogenesis. Human reproduction (Oxford, England), 24(3), 536–545. https://doi.org/10.1093/humrep/den390

22. Kang, J. Y., Kim, J. Y., Kim, K. B., Park, J. W., Cho, H., Hahm, J. Y., Chae, Y. C., Kim, D., Kook, H., Rhee, S., Ha, N. C., & Seo, S. B. (2018). KDM2B is a histone H3K79 demethylase and induces transcriptional repression via sirtuin-1-mediated chromatin silencing. FASEB journal : official publication of the Federation of American Societies for Experimental Biology, 32(10), 5737–5750. https://doi.org/10.1096/fj.201800242R

23. Kouzarides T. (2007). Chromatin modifications and their function. *Cell*, *128*(4), 693–705. https://doi.org/10.1016/j.cell.2007.02.005

24. Kuroki, S., Maeda, R., Yano, M., Kitano, S., Miyachi, H., Fukuda, M., Shinkai, Y., & Tachibana, M. (2020). H3K9 Demethylases JMJD1A and JMJD1B Control Prospermatogonia to Spermatogonia Transition in Mouse Germline. Stem cell reports, 15(2), 424–438. https://doi.org/10.1016/j.stemcr.2020.06.013

25. Lambrot, R., Lafleur, C., & Kimmins, S. (2015). The histone demethylase KDM1A is essential for the maintenance and differentiation of spermatogonial stem cells and progenitors. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*, *29*(11), 4402–4416. https://doi.org/10.1096/fj.14-267328

26. Lavery, W. J., Barski, A., Wiley, S., Schorry, E. K., & Lindsley, A. W. (2020). KMT2C/D COMPASS complex-associated diseases [KCDCOM-ADs]: an emerging class of

congenital regulopathies. Clinical epigenetics, 12(1), 10. https://doi.org/10.1186/s13148-019-0802-2

27. Lesch, B. J., Tothova, Z., Morgan, E. A., Liao, Z., Bronson, R. T., Ebert, B. L., & Page, D. C. (2019). Intergenerational epigenetic inheritance of cancer susceptibility in mammals. eLife, 8, e39380. https://doi.org/10.7554/eLife.39380

28. Lismer, A., Siklenka, K., Lafleur, C., Dumeaux, V., & Kimmins, S. (2020). Sperm histone H3 lysine 4 trimethylation is altered in a genetic mouse model of transgenerational epigenetic inheritance. *Nucleic acids research*, *48*(20), 11380–11393. https://doi.org/10.1093/nar/gkaa712

29. Lu, L. Y., Wu, J., Ye, L., Gavrilina, G. B., Saunders, T. L., & Yu, X. (2010). RNF8-dependent histone modifications regulate nucleosome removal during spermatogenesis. Developmental cell, 18(3), 371–384. https://doi.org/10.1016/j.devcel.2010.01.010

30. Luense, L. J., Donahue, G., Lin-Shiao, E., Rangel, R., Weller, A. H., Bartolomei, M. S., & Berger, S. L. (2019). Gcn5-Mediated Histone Acetylation Governs Nucleosome Dynamics in Spermiogenesis. *Developmental cell*, *51*(6), 745–758.e6. https://doi.org/10.1016/j.devcel.2019.10.024

31. Luense, L. J., Wang, X., Schon, S. B., Weller, A. H., Lin Shiao, E., Bryant, J. M., Bartolomei, M. S., Coutifaris, C., Garcia, B. A., & Berger, S. L. (2016). Comprehensive analysis of histone post-translational modifications in mouse and human male germ cells. *Epigenetics & chromatin*, *9*, 24. https://doi.org/10.1186/s13072-016-0072-6

32. Ma, X. S., Chao, S. B., Huang, X. J., Lin, F., Qin, L., Wang, X. G., Meng, T. G., Zhu, C. C., Schatten, H., Liu, H. L., & Sun, Q. Y. (2015). The Dynamics and Regulatory Mechanism of Pronuclear H3k9me2 Asymmetry in Mouse Zygotes. Scientific reports, 5, 17924. https://doi.org/10.1038/srep17924

33. Maezawa, S., Hasegawa, K., Yukawa, M., Kubo, N., Sakashita, A., Alavattam, K. G., Sin, H. S., Kartashov, A. V., Sasaki, H., Barski, A., & Namekawa, S. H. (2018). Polycomb protein SCML2 facilitates H3K27me3 to establish bivalent domains in the male germline. Proceedings of the National Academy of Sciences of the United States of America, 115(19), 4957–4962. https://doi.org/10.1073/pnas.1804512115

34. Mäkelä, J. A., & Hobbs, R. M. (2019). Molecular regulation of spermatogonial stem cell renewal and differentiation. Reproduction (Cambridge, England), 158(5), R169–R187. https://doi.org/10.1530/REP-18-0476

35. Martinez-Gamero, C., Malla, S., & Aguilo, F. (2021). LSD1: Expanding Functions in Stem Cells and Differentiation. *Cells*, *10*(11), 3252. https://doi.org/10.3390/cells10113252

36. Mayorek, N., Schlossberg, M.,, Mansour, Y., Pillar, N., Stein, I., Mushasha, F., Baziza, G., Medvedev, E., Manevitch, Z., Menzel, J., Aizenshtein, E., Sarvin, B., Sarvin, N., Shlomi, T., Klutstein, M., & Pikarsky. E. (2022). 2-hydroxyglutarate controls centromere and heterochromatin conformation and function in the male germline. bioRxiv 2022.05.29.493890; doi: https://doi.org/10.1101/2022.05.29.493890

37. McCarthy, A., Sarkar, K., Martin, E. T., Upadhyay, M., Jang, S., Williams, N. D., Forni, P. E., Buszczak, M., & Rangan, P. (2022). Msl3 promotes germline stem cell differentiation in female Drosophila. *Development (Cambridge, England)*, *149*(1), dev199625. https://doi.org/10.1242/dev.199625

38. Meistrich, M. L., Trostle-Weige, P. K., Lin, R., Bhatnagar, Y. M., & Allis, C. D. (1992). Highly acetylated H4 is associated with histone displacement in rat spermatids. Molecular reproduction and development, 31(3), 170–181. https://doi.org/10.1002/mrd.1080310303

39. Metzger, E., Wang, S., Urban, S., Willmann, D., Schmidt, A., Offermann, A., Allen, A., Sum, M., Obier, N., Cottard, F., Ulferts, S., Preca, B. T., Hermann, B., Maurer, J.,

Greschik, H., Hornung, V., Einsle, O., Perner, S., Imhof, A., Jung, M., … Schüle, R. (2019). KMT9 monomethylates histone H4 lysine 12 and controls proliferation of prostate cancer cells. Nature structural & molecular biology, 26(5), 361–371. https://doi.org/10.1038/s41594-019-0219-9

40. Michishita, E., McCord, R. A., Berber, E., Kioi, M., Padilla-Nash, H., Damian, M., Cheung, P., Kusumoto, R., Kawahara, T. L., Barrett, J. C., Chang, H. Y., Bohr, V. A., Ried, T., Gozani, O., & Chua, K. F. (2008). SIRT6 is a histone H3 lysine 9 deacetylase that modulates telomeric chromatin. *Nature*, *452*(7186), 492–496. https://doi.org/10.1038/nature06736

41. Mosammaparast, N., & Shi, Y. (2010). Reversal of histone methylation: biochemical and molecular mechanisms of histone demethylases. *Annual review of biochemistry*, 79, 155–179. https://doi.org/10.1146/annurev.biochem.78.070907.103946

42. Mulligan, P., Westbrook, T. F., Ottinger, M., Pavlova, N., Chang, B., Macia, E., Shi, Y. J., Barretina, J., Liu, J., Howley, P. M., Elledge, S. J., & Shi, Y. (2008). CDYL bridges REST and histone methyltransferases for gene repression and suppression of cellular transformation. Molecular cell, 32(5), 718–726. https://doi.org/10.1016/j.molcel.2008.10.025

43. Myrick, D. A., Christopher, M. A., Scott, A. M., Simon, A. K., Donlin-Asp, P. G., Kelly, W. G., & Katz, D. J. (2017). KDM1A/LSD1 regulates the differentiation and maintenance of spermatogonia in mice. PloS one, 12(5), e0177473. https://doi.org/10.1371/journal.pone.0177473

44. Nebbioso, A., Tambaro, F. P., Dell'Aversana, C., & Altucci, L. (2018). Cancer epigenetics: Moving forward. *PLoS genetics*, *14*(6), e1007362. https://doi.org/10.1371/journal.pgen.1007362

45. Ogawa, H., Ishiguro, K., Gaubatz, S., Livingston, D. M., & Nakatani, Y. (2002). A complex with chromatin modifiers that occupies E2F- and Myc-responsive genes in G0 cells. Science (New York, N.Y.), 296(5570), 1132–1136. https://doi.org/10.1126/science.1069861

46. Ontoso, D., Kauppi, L., Keeney, S., & San-Segundo, P. A. (2014). Dynamics of DOT1L localization and H3K79 methylation during meiotic prophase I in mouse spermatocytes. Chromosoma, 123(1-2), 147–164. https://doi.org/10.1007/s00412-013-0438-5

47. Paigen, K., & Petkov, P. M. (2018). PRDM9 and Its Role in Genetic Recombination. Trends in genetics : TIG, 34(4), 291–300. https://doi.org/10.1016/j.tig.2017.12.017

48. Parra M. (2015). Class IIa HDACs - new insights into their functions in physiology and pathology. *The FEBS journal*, *282*(9), 1736–1744. https://doi.org/10.1111/febs.13061

49. Peters, A. H., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer, C., Weipoltshammer, K., Pagani, M., Lachner, M., Kohlmaier, A., Opravil, S., Doyle, M., Sibilia, M., & Jenuwein, T. (2001). Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin and genome stability. Cell, 107(3), 323–337. https://doi.org/10.1016/s0092-8674(01)00542-6

50. Powers, N. R., Parvanov, E. D., Baker, C. L., Walker, M., Petkov, P. M., & Paigen, K. (2016). The Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at Recombination Hotspots In Vivo. PLoS genetics, 12(6), e1006146. https://doi.org/10.1371/journal.pgen.1006146

51. Prachayasittikul, V., Prathipati, P., Pratiwi, R., Phanus-Umporn, C., Malik, A. A., Schaduangrat, N., Seenprachawong, K., Wongchitrat, P., Supokawej, A., Prachayasittikul, V., Wikberg, J. E., & Nantasenamat, C. (2017). Exploring the epigenetic drug discovery

landscape. *Expert opinion on drug discovery*, *12*(4), 345–362. https://doi.org/10.1080/17460441.2017.1295954

52. Pradeepa, M. M., Nikhil, G., Hari Kishore, A., Bharath, G. N., Kundu, T. K., & Rao, M. R. (2009). Acetylation of transition protein 2 (TP2) by KAT3B (p300) alters its DNA condensation property and interaction with putative histone chaperone NPM3. *The Journal of biological chemistry*, *284*(43), 29956–29967. https://doi.org/10.1074/jbc.M109.052043

53. Rajender, S., Avery, K., & Agarwal, A. (2011). Epigenetics, spermatogenesis and male infertility. Mutation research, 727(3), 62–71. https://doi.org/10.1016/j.mrrev.2011.04.002

54. Schuettengruber, B., Chourrout, D., Vervoort, M., Leblanc, B., & Cavalli, G. (2007). Genome regulation by polycomb and trithorax proteins. Cell, 128(4), 735–745. https://doi.org/10.1016/j.cell.2007.02.009

55. Schulz, K. N., & Harrison, M. M. (2019). Mechanisms regulating zygotic genome activation. Nature reviews. Genetics, 20(4), 221–234. https://doi.org/10.1038/s41576-018-0087-x

56. Schuster, A., Tang, C., Xie, Y., Ortogero, N., Yuan, S., & Yan, W. (2016). SpermBase: A Database for Sperm-Borne RNA Contents. Biology of reproduction, 95(5), 99. https://doi.org/10.1095/biolreprod.116.142190

57. Shirakawa, T., Yaman-Deveci, R., Tomizawa, S., Kamizato, Y., Nakajima, K., Sone, H., Sato, Y., Sharif, J., Yamashita, A., Takada-Horisawa, Y., Yoshida, S., Ura, K., Muto, M., Koseki, H., Suda, T., & Ohbo, K. (2013). An epigenetic switch is crucial for spermatogonia to exit the undifferentiated state toward a Kit-positive identity. Development (Cambridge, England), 140(17), 3565–3576. https://doi.org/10.1242/dev.094045

58. Siklenka, K., Erkek, S., Godmann, M., Lambrot, R., McGraw, S., Lafleur, C., Cohen, T., Xia, J., Suderman, M., Hallett, M., Trasler, J., Peters, A. H., & Kimmins, S. (2015). Disruption of histone methylation in developing sperm impairs offspring health transgenerationally. Science (New York, N.Y.), 350(6261), aab2006. https://doi.org/10.1126/science.aab2006

59. Sollier, J., Lin, W., Soustelle, C., Suhre, K., Nicolas, A., Géli, V., & de La Roche Saint-André, C. (2004). Set1 is required for meiotic S-phase onset, double-strand break formation and middle gene expression. The EMBO journal, 23(9), 1957–1967. https://doi.org/10.1038/sj.emboj.7600204

60. Sone, Y., Ito, M., Shirakawa, H., Shikano, T., Takeuchi, H., Kinoshita, K., & Miyazaki, S. (2005). Nuclear translocation of phospholipase C-zeta, an egg-activating factor, during early embryonic development. Biochemical and biophysical research communications, 330(3), 690–694. https://doi.org/10.1016/j.bbrc.2005.03.032

61. Song, N., Liu, J., An, S., Nishino, T., Hishikawa, Y., & Koji, T. (2011). Immunohistochemical Analysis of Histone H3 Modifications in Germ Cells during Mouse Spermatogenesis. *Acta histochemica et cytochemica*, *44*(4), 183–190. https://doi.org/10.1267/ahc.11027

62. Spencer, T. E., Jenster, G., Burcin, M. M., Allis, C. D., Zhou, J., Mizzen, C. A., McKenna, N. J., Onate, S. A., Tsai, S. Y., Tsai, M. J., & O'Malley, B. W. (1997). Steroid receptor coactivator-1 is a histone acetyltransferase. Nature, 389(6647), 194–198. https://doi.org/10.1038/38304

63. Steilmann, C., Paradowska, A., Bartkuhn, M., Vieweg, M., Schuppe, H. C., Bergmann, M., Kliesch, S., Weidner, W., & Steger, K. (2011). Presence of histone H3 acetylated at lysine 9 in male germ cells and its distribution pattern in the genome of human spermatozoa. Reproduction, fertility, and development, 23(8), 997–1011. https://doi.org/10.1071/RD10197

64. Sun, Z., Zhang, Y., Jia, J., Fang, Y., Tang, Y., Wu, H., & Fang, D. (2020). H3K36me3, message from chromatin to DNA damage repair. Cell & bioscience, 10, 9. https://doi.org/10.1186/s13578-020-0374-z

65. Tachibana, M., Nozaki, M., Takeda, N., & Shinkai, Y. (2007). Functional dynamics of H3K9 methylation during meiotic prophase progression. *The EMBO journal*, *26*(14), 3346–3359. https://doi.org/10.1038/sj.emboj.7601767

66. Takada, Y., Naruse, C., Costa, Y., Shirakawa, T., Tachibana, M., Sharif, J., Kezuka-Shiotani, F., Kakiuchi, D., Masumoto, H., Shinkai, Y., Ohbo, K., Peters, A. H., Turner, J. M., Asano, M., & Koseki, H. (2011). HP1γ links histone methylation marks to meiotic synapsis in mice. Development (Cambridge, England), 138(19), 4207–4217. https://doi.org/10.1242/dev.064444

67. Tomizawa, S. I., Kobayashi, Y., Shirakawa, T., Watanabe, K., Mizoguchi, K., Hoshi, I., Nakajima, K., Nakabayashi, J., Singh, S., Dahl, A., Alexopoulou, D., Seki, M., Suzuki, Y., Royo, H., Peters, A., Anastassiadis, K., Stewart, A. F., & Ohbo, K. (2018). Kmt2b conveys monovalent and bivalent H3K4me3 in mouse spermatogonial stem cells at germline and embryonic promoters. Development (Cambridge, England), 145(23), dev169102. https://doi.org/10.1242/dev.169102

68. Trigg, N. A., Eamens, A. L., & Nixon, B. (2019). The contribution of epididymosomes to the sperm small RNA profile. Reproduction (Cambridge, England), 157(6), R209–R223. https://doi.org/10.1530/REP-18-0480

69. van der Heijden, G. W., Derijck, A. A., Ramos, L., Giele, M., van der Vlag, J., & de Boer, P. (2006). Transmission of modified nucleosomes from the mouse male germline to the zygote and subsequent remodeling of paternal chromatin. Developmental biology, 298(2), 458–469. https://doi.org/10.1016/j.ydbio.2006.06.051

70. Wang, T., Gao, H., Li, W., & Liu, C. (2019). Essential Role of Histone Replacement and Modifications in Male Fertility. Frontiers in genetics, 10, 962. https://doi.org/10.3389/fgene.2019.00962

71. Wang, Y., Iwamori, T., Kaneko, T., Iida, H., & Iwamori, N. (2021). Comparative distributions of RSBN1 and methylated histone H4 Lysine 20 in the mouse spermatogenesis. *PloS one*, *16*(6), e0253897. https://doi.org/10.1371/journal.pone.0253897

72. Wapenaar, H., & Dekker, F. J. (2016). Histone acetyltransferases: challenges in targeting bi-substrate enzymes. *Clinical epigenetics*, 8, 59. https://doi.org/10.1186/s13148-016-0225-2

73. Wu, D., Li, Y., Zhu, K. S., Wang, H., & Zhu, W. G. (2018). Advances in Cellular Characterization of the Sirtuin Isoform, SIRT7. Frontiers in endocrinology, 9, 652. https://doi.org/10.3389/fendo.2018.00652

74. Yang, X., Li, L., Liang, J., Shi, L., Yang, J., Yi, X., Zhang, D., Han, X., Yu, N., & Shang, Y. (2013). Histone acetyltransferase 1 promotes homologous recombination in DNA repair by facilitating histone turnover. The Journal of biological chemistry, 288(25), 18271–18282. https://doi.org/10.1074/jbc.M113.473199

75. Yao, C. J., Xu, W. J., Gong, X. L., Zhou, Y., Yan, Z. Q., Zhu, Z. J., Wang, Z. X., Li, Q. L., Guo, X. B., Wang, L. Y., Ma, D., & Qiao, Z. D. (2010). The role of Dby mRNA in early development of male mouse zygotes. Asian journal of andrology, 12(4), 567–577. https://doi.org/10.1038/aja.2010.28

76. Yoshida, K., Maekawa, T., Ly, N. H., Fujita, S. I., Muratani, M., Ando, M., Katou, Y., Araki, H., Miura, F., Shirahige, K., Okada, M., Ito, T., Chatton, B., & Ishii, S. (2020). ATF7-Dependent Epigenetic Changes Are Required for the Intergenerational Effect of a

Paternal Low-Protein Diet. Molecular cell, 78(3), 445–458.e6. https://doi.org/10.1016/j.molcel.2020.02.028

77. Zamudio, N. M., Scott, H. S., Wolski, K., Lo, C. Y., Law, C., Leong, D., Kinkel, S. A., Chong, S., Jolley, D., Smyth, G. K., de Kretser, D., Whitelaw, E., & O'Bryan, M. K. (2011). DNMT3L is a regulator of X chromosome compaction and post-meiotic gene transcription. PloS one, 6(3), e18276. https://doi.org/10.1371/journal.pone.0018276

78. Zhang, C., Wang, M., Li, Y., & Zhang, Y. (2022). Profiling and functional characterization of maternal mRNA translation during mouse maternal-to-zygotic transition. Science advances, 8(5), eabj3967. https://doi.org/10.1126/sciadv.abj3967

79. Zhang, X., Gao, F., Fu, J., Zhang, P., Wang, Y., & Zeng, X. (2017). Systematic identification and characterization of long non-coding RNAs in mouse mature sperm. PloS one, 12(3), e0173402. https://doi.org/10.1371/journal.pone.0173402

80. Zuo, X., Rong, B., Li, L., Lv, R., Lan, F., & Tong, M. H. (2018). The histone methyltransferase SETD2 is required for expression of acrosin-binding protein 1 and protamines and essential for spermiogenesis in mice. *The Journal of biological chemistry*, *293*(24), 9188–9197. https://doi.org/10.1074/jbc.RA118.002851

Table 1. Writers and erasers families for histone K acetylation and methylation and the targets described.

Acetylation — WRITERS

FAMILY	NAME	TARGET
GNAT	KAT2A - GCN5	H3ac
	KAT2B - PCAF	H3ac
	HAT1	H4ac
	ATF2	H4/2Bac
NAT	NAA40 - NAT11	H4ac
	NAA50 - NAT13	H4ac
	NAA60 - NAT15	H4K79/91ac
MYST	KAT5 - TIP60	H4/ H2AZac
	KAT8 - MYST1 - MOF	H4K16ac
	KAT7 - MYST2 - HBO1	H3/4ac
	KAT6A - MYST3 - MOZ	H3/4ac
	KAT6B - MYST4 - MORF	H3ac
p300/CBP	KAT3A - CREBBP	H3ac
	KAT3B - EP300	H3ac
SRC	NCOA1	H3/4ac
	NCOA3	H3ac
Other	CLOCK	H3ac
	MSL3	H4K16ac
	GTF3C4	H3ac
	BRCA2	H3ac
	MCM3AP	H3ac
	SUPT7L	H3ac

Acetylation — ERASERS

FAMILY	NAME	TARGET
Class I	HDAC1	Hac
	HDAC2	Hac
	HDAC3	Hac
	HDAC8	Hac
Class IIa	HDAC4	Hac
	HDAC5	Hac
	HDAC7	Hac
	HDAC9	Hac
Class III	SIRT1	H4K16ac
	SIRT2	H4K16ac
	SIRT3	H4K16ac
	SIRT5	Hac
	SIRT6	H3K9ac
	SIRT7	H3K18ac
Class IV	HDAC11	Hac

Methylation — WRITERS

FAMILY	NAME	TARGET
KMT1	KMT1A - SUV39H1	H3K9me3
	KMT1B - SUV39H2	H3K9me3
	KMT1C - EHMT2 - G9a	H3K9me1/2
	KMT1D - EHMT1 - GLP	H3K9me1/2
	KMT1E - SETDB1	H3K9me3
	KMT1F - SETDB2	H3K9me3
KMT2	KMT2A - MLL1	H3K4me1/2/3
	KMT2B - MLL2	H3K4me1/2/3
	KMT2C - MLL3	H3K4me1
	KMT2D - MLL4	H3K4me1
	KMT2F - SETD1A	H3K4me1/2/3
	KMT2G - SETD1B	H3K4me1/2/3
	KMT2H - ASH1L	H3K36me2
KMT2 core	ASH2L	H3K4me1/2/3
KMT3	KMT3A - SETD2	H3K36me3
	KMT3B - NSD1	H3K36me2
	KMT3C - SMYD2	H3K4me1/2/3
		H3K36me2
	KMT3D - SMYD1	H3K4me1/2/3
	KMT3E - SMYD3	H3K4me2/3
	KMT3G - NSD2 - WHSC1	H3K36me2
	KMT3F - NSD3 - WHSC1L1	H3K36me2
KMT4	DOT1L	H3K79me1/2/3
KMT5	KMT5A - SETD8	H4K20me1
	KMT5B - SUV420H1	H4K20me1/2/3
	KMT5C - SUV420H2	H4K20me1/2/3
KMT6	KMT6A - EZH2	H3K27me1/2/3
	KMT6B - EZH1	H3K27me1/2/3
KMT7	KMT7 - SETD7	H3K4me1
KMT8	KMT8B - PRDM9	H3K4me1/2/3
		H3K36me1/2/3
	KMT8A - PRDM2	H3K9me1
	KMT8D - PRDM8	H3K9me2
	KMT8C - PRDM6	H4K20me
KMT9	KMT9 - N6AMT1- HEMK2	H4K12me1
SMYD	SMYD4	H3K4me1/2/3
	SMYD5	H4K20me3
SETD	SETD4	H3K4me1/2
		H4K20me3
	SETD5	H3K36me3
	SETD6	H2AZK7me
	SETMAR	H3K36me2
PRDM	PRDM15	H3K9me1

Methylation — ERASERS

FAMILY	NAME	TARGET
KDM1	KDM1A - LSD1	H3K4me1/2
		H3K9me1/2
	KDM1B - LSD2	H3K4me1/2
KDM2	KDM2A - JHDM1A	H3K36me2
		H3K4me3
	KDM2B - JHDM1B	H3K36me2
		H3K79me2/3
KDM3	KDM3A - JMJD1A - JHDM2A	H3K9me1/2
	KDM3B - JMJD1B	H3K9me1/2
	KDM3C - JMJD1C	H3K9me1/2
KDM4	KDM4A - JMJD2A - JHDM3A	H3K9me2/3
		H3K36me2/3
	KDM4B - JMJD2B - JHDM3B	H3K9me3
		H3K36me3
	KDM4C - JMJD2C - JHDM3C	H3K9me3
		H3K36me3
	KDM4D - JMJD2D - JHDM3E	H3K9me2/3
KDM5	KDM5A - JARID1A	H3K4me2/3
	KDM5B - JARID1B	H3K4me1/2/3
	KDM5C - JARID1C	H3K4me2/3
	KDM5D - JARID1D	H3K4me2/3
KDM6	KDM6A - UTX	H3K27me2/3
	KDM6B - JMJD3	H3K27me2/3
KDM7	KDM7A - JHDM1D	H3K9me2
		H3K27me2
		H3K9me1/2
	KDM7B - JHDM1F - PHF8	H3K27me2
		H4K20me1
	KDM7C- JHDM1E - PHF2	H3K9me2
		H4K20me3
KDM8	KDM8 - JMJD5	H3K9me1
		H3K36me1/2
KDM9	KDM9 - RSBN1	H4K20me2/3

The HATs can be divided into six major families: 1) the GNATs (GCN5-related N-acetyltransferases), 2) the NATs (N-terminal acetylases, which also possess a GNAT domain), 3) the MYST, 4) the p300/CBP, 5) the SRC steroid receptors coactivators, and 6) other HATs (Wapenaar and Dekker, 2016). HATs actions are counteracted by the HDACs, which include zinc-dependent class I, class IIa, and class IV HDACs, and NAD-dependent class III Sirtuins (Wapenaar and Dekker, 2016). On the other hand, the KMTs present two domains with annotated lysine methyltransferase activity: 1) SET domains, that catalyze all the canonical methylation sites except for H3K79, and 2) 7βS domains (DOT1L and N6AMT1) (Mosammaparast and Shi, 2010). The KDMs can be divided according to their catalytic activity in: 1) amine oxidases (KDM1), and 2) Jumonji C demethylases (KDM2/3/4/5/6/7/8) (Mosammaparast and Shi, 2010).

501
Figure 1. Canonical histone K marks and writers/erasers families distribution during
spermatogenesis. A) Histone K methylation PTMs distribution at DNA elements and functional
roles in gene expression for an active gene (pink shading) or inactive gene (gray shading). TSS:
transcription start site. Only K modifications with sufficient and consensual information about
distribution across genes are shown. B) Main histone K PTMs levels involved in chromatin
remodeling during spermatogenesis. The data presented is summarized from the following
references: Song et al., 2011; Godmann et al., 2007; Zuo et al., 2018; Luense et al., 2016; Wang et
al., 2021). SGund: spermatogonia Thy1+/Kit-, SGdiff: spermatogonia Kit+, PreL: pre-leptotene
spermatocytes, LZ: leptotene/zygotene spermatocytes, PD: pachytene/diplotene spermatocytes,
RStid: round spermatid.

512
513
514

Figure 2. Unsupervised hierarchical clustering analysis of male germ cells transcriptome.

Heatmap showing z-score values of histone K acetylation and methylation writer/eraser enzymes.

SGund: spermatogonia Kit-, SGdiff: spermatogonia Kit+, PreL: pre-leptotene spermatocytes, LZ:

leptotene/zygotene spermatocytes, PD: pachytene/diplotene spermatocytes, RStid: round spermatid.

Figure 3. Histone K acetylation and methylation epigenetic enzymes expression during

spermatogenesis A) Number of upregulated epigenetic enzymes for each germ cell

stage.Differentiall gene expression analysis was conducted with DESeq2 on consecutive

populations across differentiationn (SGdiff vs SGund, PreL vs SGdiff, LZ vs PreL, PD vs LZ and

RStid vs PD), and upregulated genes selected with padj<0.05 and log2FC>0.5. B) Heatmap

showing epigenetic enzymes not significantly upregulated at any germ cell stage across

spermatogenesis. SGund: spermatogonia Kit-, SGdiff: spermatogonia Kit+, PreL: pre-leptotene

spermatocytes, LZ: leptotene/zygotene spermatocytes, PD:pachytenee/diplotene spermatocytes,

RStid: round spermatid.

Figure 4. Epigenetic histone K writers and erasers upregulation at spermatogonial phase. The
left panel indicates the heatmap showing z score values and the right panel indicates the Dot plot
showing enzymes RPKM values in SGund (A) and SGdiff (B) population. Blue dots: writers, red
dots: erasers. SGund: spermatogonia Kit-, SGdiff: spermatogonia Kit+.

Figure 5. Epigenetic histone K writers and erasers upregulation at meiosis prophase I. The left panel indicates the heatmap showing z score values and the right panel indicates the Dot plot showing enzymes RPKM values, in PreL (A), LZ (B) and PD (C) spermatocytes. Blue dots: writers, red dots: erasers. PreL: pre-leptotene spermatocytes, LZ: leptotene/zygotene spermatocytes, PD:pachytenee/diplotene spermatocytes.

547 **Figure 6. Epigenetic histone K writers and erasers upregulation at spermiogenesis.** The left

548 panel indicates the heatmap showing z score values and the right panel indicates the Dot plot

549 showing enzymes RPKM values in RStid. Blue dots: writers, red dots: erasers. RStid: round

550 spermatid, PD: pachytene/diplotene spermatocytes.

Figure 7. Epigenetic histone K writers and erasers detected at mature sperm RNA cargo and
1-cell embryo translatome. A) Pearson correlation between epigenetic enzymes mean percent
rank detected at RStid and sperm. B) Mature sperm word cloud for epigenetic enzymes
indicating the enzyme abundance (word size = mean percent rank) and translation
efficiency (TE) detected at 1 cell embryo (red: TE > 0, black: TE=0). C) Epigenetic
enzymes TE in mouse pre-implantation embryos (TE = RNA RPKM at ribosome / total
RNA RPKM, RPKM>0.5)

560