Transcriptome profiling of histone acetylation and methylation writers and erasers families across male germ cell development and pre-cleavage zygote

Candela R. González [1], Camila Perez Lujan [2], Alfredo D. Vitullo [1], Betina González [2]*

[1] Centro de Estudios Biomédicos Básicos, Aplicados y Desarrollo (CEBBAD), Universidad Maimónides, Ciudad Autónoma de Buenos Aires, Buenos Aires, Argentina.

[2] Instituto de Investigaciones Farmacológicas (Universidad de Buenos Aires–Consejo Nacional de Investigaciones Científicas y Técnicas), Ciudad Autónoma de Buenos Aires, Buenos Aires, Argentina.

* Corresponding author
E-mail: bgonzalez@ffyb.uba.ar

Keywords: **spermatogenesis, epigenetic enzymes, histone post-transcriptional modification**

ABSTRACT

Accumulating evidence put on the focus that male spermatogenesis presents windows of vulnerability for epigenetic reprogramming by environmental stressors that not only can affect fertility but even transmit developmental, metabolic, and behavioral traits to offspring. This work analyzed the transcriptomic programs of epigenetic enzymes involved in histone K acetylation and methylation from undifferentiated spermatogonia to mature sperm and pre-cleavage zygote in the mouse. We dissected the peak expression at each stage for families and individual enzymes, their target histone PTMs functions and reported dynamics during spermatogenesis and early embryogenesis. This approach allowed us to identify epigenetic enzymes with known functions during spermatogenesis and, by the rule of "guilty by association", many others not even described yet in male germ cell development. Moreover, some of these enzymes' mRNA were detected higher in the pre-

cleavage zygote than in the transcriptionally silent MII oocyte, suggesting a paternal

contribution. Our study contributes to the mechanistic aspects behind transgenerational

epigenetics, described for the first time in such a comprehensive approach.

INTRODUCTION

In the last years, accumulating research has documented that male germ cell

maturation presents windows of vulnerability for epigenetic reprogramming by

environmental stressors that not only can affect fertility but even transmit developmental,

metabolic and behavioral traits to offspring (Bale, 2015). This type of non-genetic

"Lamarckian" transmission has been established for paternal lifestyle and different forms of

chronic stress, drug abuse, dietary change, and social defeat, involving changes in non-

coding RNAs cargo, DNA methylation, and histone PTMs (Bale, 2105). However, most of

the research in epigenetic inheritance has documented the role of non-coding RNAs and

DNA methylation patterns, whereas the mechanisms by which the environment can trigger

specific changes on histone PTMs are still largely unexplored (Skvortsova et al., 2018).

Spermatogenesis is a finely regulated process where unipotent spermatogonia

undergo meiosis and, subsequently, spermiogenesis, to become sperm cells. During

developmental stages of spermatogenesis, male germ cells experience dramatic chromatin

reorganization, where approximately 90% (human) to 95% (mouse) of histones are evicted

and replaced by protamines to compact the paternal genome (Rajender et al., 2011, Carrel

2012). Importantly, a small fraction of histones in the sperm genome is retained on specific

locations and carries several post-translational modifications (PTMs) that play critical roles

in epigenetic regulation of spermatogenesis and early embryonic development (Luense et

al., 2016). The epigenetic program of histone PTMs that controls spermatogenesis is driven

by enzymatic families known as writers and erasers of the so-called "histone code", that

modulate chromatin structure and transcription (Kouzarides, 2007). These epigenetic

enzymes catalyze the deposition or removal of specific PTMs such as acetylation,

methylation, phosphorylation, crotonylation, etc, at specific amino acid residues on the

nucleosome's core histones H2A, H2B, H3, and H4, and also on the linker H1/H5 (Luense

et al., 2016). In male germ cells, histone PTMs not only control active vs repressed

chromatin states but also a range of processes including DNA replication and repair, chromosome maintenance and histone eviction (Luense et al., 2016). So far, the most studied histone PTMs related to the control of germ cells mitosis, meiosis, spermiogenesis and early embryo development are methylated and acetylated lysines (K) on H2A/B, H3 and H4. Histone K acetylation can be dynamically regulated by the opposing action of acetyltransferases (HATs) and deacetylases (HDACs). Acetylation of K residues neutralizes the positive charge on histones, allowing DNA-binding proteins better access to the DNA and resulting in activation of gene expression as well as histone eviction (Rajender et al., 2011). Unlike acetylation, methylation does not affect histone charge but regulates recognition and interaction with chromatin-binding proteins that control the transcription or respond to DNA damage (Rajender et al., 2011). Histone K methylation is finely regulated by methyltransferases (KMTs) and demethylases (KDMs) that control the mono-, di-, and/or tri-methylation of specific residues, and this can either activate or repress transcription depending on the residue position, the number of methylations and the presence of other methyl or acetyl groups in the vicinity (Black et al., 2012).

The role of epigenetic enzymes comes to focus, as they are responsible for the histone's PTMs writing and erasing balance. Moreover, histone-modifying enzymes have been shown to respond to the organism physiology and metabolism, several disease conditions, and environmental stressors (Fan et al., 2016; Nebbioso et al., 2018; Gonzalez et al., 2020). Even though epigenetic enzymes are the bridge linking the environment with epigenetic inheritance through retained histones PTMs, an in-depth analysis of their expression patterns throughout the spermatogenic process is still lacking. Moreover, these enzymes provide new targets for therapies of numerous diseases (Prachayasittikul et al., 2017) but the potential effect of these compounds on epigenetic inheritance is generally overlooked. Here, we curated transcriptomic data from spermatogonia to mature sperm populations and pre-cleavage zygote and analyzed the expression profile and dynamic changes of all the known families of histone K acetylation and methylation writers and erasers. We provide a complete picture of the epigenetic enzymes across the spermatogenesis in mice and the developmental windows when the transcription of these enzymes may be more susceptible to environmental disruption. Not only we expand the knowledge on enzymes with recognized roles in spermatogenesis, but provide evidence of

many more whose role in male germ cells development and zygote have not been described

yet.

RESULTS

We curated RNA-seq data available in purified male germ cells from adult mice and

selected the dataset of one of the most referenced studies regarding male germ cells

epigenomic and transcriptomic profiles published by Hammoud et al. (2014). We analyzed

the transcriptome from spermatogonia (SG), including the self-renewing and stem-like

Thy1+/Kit- (SGThy1) and the committed Thy1-/Kit+ (SGKit) population, primary

spermatocyte (SCyte), round spermatid (STid) and mature sperm stages.

1. Epigenetic writers and erasers involved in male germ cells histone K acetylation

and methylation dynamics

Table 1 shows writers and erasers families for histone K acetylation and

methylation, and the most representative targets described in mammalian cells. The HATs

can be divided into six major families: 1) the GNATs (GCN5-related N-acetyltransferases),

2) the NATs (N-terminal acetylases, which also possess a GNAT domain), 3) the MYST, 4)

the p300/CBP, 5) the SRC steroid receptors coactivators, and 6) other HATs (Wapenaar

and Dekker, 2016). HATs actions are counteracted by the HDACs, which include zinc-

dependent class I, class IIa, and class IV HDACs, and NAD-dependent class III Sirtuins

(Wapenaar and Dekker, 2016). On the other hand, the KMTs present two domains with

annotated lysine methyltransferase activity: 1) SET domains, that catalyze all the canonical

methylation sites except for H3K79, and 2) 7βS domains (the H3K79-specific DOT1L)

(Mosammaparast and Shi, 2010). The KDMs can be divided according to their catalytic

activity in: 1) amine oxidases (KDM1), and 2) Jumonji C demethylases

(KDM2/3/4/5/6/7/8) (Mosammaparast and Shi, 2010).

Table 1. Writers and erasers families for histone K acetylation and methylation and

the targets described.

Acetylation

WRITERS			ERASERS		
FAMILY	NAME	TARGET	FAMILY	NAME	TARGET
GNAT	KAT2A - GCN5	H3ac	Class I	HDAC1	Hac
	KAT2B - PCAF	H3ac		HDAC2	Hac
	HAT1	H4ac		HDAC3	Hac
	ATF2	H4/2Bac		HDAC8	Hac
NAT	NAA40 - NAT11	H4ac	Class IIa	HDAC4	Hac
	NAA50 - NAT13	H4ac		HDAC5	Hac
	NAA60 - NAT15	H4K79/91ac		HDAC7	Hac
MYST	KAT5 - TIP60	H4/ H2AZac		HDAC9	Hac
	KAT8 - MYST1 - MOF	H4K16ac	Class III	SIRT1	H4K16ac
	KAT7 - MYST2 - HBO1	H3/4ac		SIRT2	H4K16ac
	KAT6A - MYST3 - MOZ	H3/4ac		SIRT3	H4K16ac
	KAT6B - MYST4 - MORF	H3ac		SIRT5	Hac
p300/CBP	KAT3A - CREBBP	H3ac		SIRT6	H3K9ac
	KAT3B - EP300	H3ac		SIRT7	H3K18ac
SRC	NCOA1	H3/4ac	Class IV	HDAC11	Hac
	NCOA3	H3ac			
Other	CLOCK	H3ac			
	MSL3	H4K16ac			
	GTF3C4	H3ac			
	BRCA2	H3ac			
	MCM3AP	H3ac			
	SUPT7L	H3ac			

Methylation

WRITERS			ERASERS		
FAMILY	NAME	TARGET	FAMILY	NAME	TARGET
KMT1	KMT1A - SUV39H1	H3K9me3	KDM1	KDM1A - LSD1	H3K4me1/2
	KMT1B - SUV39H2	H3K9me3			H3K9me1/2
	KMT1C - EHMT2 - G9a	H3K9me1/2		KDM1B - LSD2	H3K4me1/2
	KMT1D - EHMT1 - GLP	H3K9me1/2	KDM2	KDM2A - JHDM1A	H3K36me2
	KMT1E - SETDB1	H3K9me3			H3K4me3
	KMT1F - SETDB2	H3K9me3		KDM2B - JHDM1B	H3K36me2
KMT2	KMT2A - MLL1	H3K4me1/2/3			H3K79me2/3
	KMT2B - MLL2	H3K4me1/2/3	KDM3	KDM3A - JMJD1A - JHDM2A	H3K9me1/2
	KMT2C - MLL3	H3K4me1/2/3		KDM3B - JMJD1B	H3K9me1/2
	KMT2F - SETD1A	H3K4me1/2/3		KDM3C - JMJD1C	H3K9me1/2
	KMT2H - ASH1L	H3K36me2	KDM4	KDM4A - JMJD2A - JHDM3A	H3K9me2/3
KMT2 core	ASH2L	H3K4me1/2/3			H3K36me2/3
KMT3	KMT3A - SETD2	H3K36me3		KDM4B - JMJD2B - JHDM3B	H3K9me3
	KMT3B - NSD1	H3K36me2			H3K36me3
	KMT3C - SMYD2	H3K4me1/2/3 H3K36me2		KDM4C - JMJD2C - JHDM3C	H3K9me3 H3K36me3
	KMT3D - SMYD1	H3K4me1/2/3		KDM4D - JMJD2D - JHDM3E	H3K9me2/3
	KMT3E - SMYD3	H3K4me2/3	KDM5	KDM5A - JARID1A	H3K4me2/3
	KMT3G - NSD2 - WHSC1	H3K36me2		KDM5B - JARID1B	H3K4me1/2/3
	KMT3F - NSD3 - WHSC1L1	H3K36me2		KDM5C - JARID1C	H3K4me2/3
KMT4	DOT1L	H3K79me1/2/3		KDM5D - JARID1D	H3K4me2/3
KMT5	KMT5A - SETD8	H4K20me1	KDM6	KDM6A - UTX	H3K27me2/3
	KMT5B - SUV420H1	H4K20me1/2/3		KDM6B - JMJD3	H3K27me2/3
	KMT5C - SUV420H2	H4K20me1/2/3	KDM7	KDM7A - JHDM1D	H3K9me2 H3K27me2
KMT6	KMT6A - EZH2	H3K27me1/2/3			H3K9me1/2
	KMT6B - EZH1	H3K27me1/2/3		KDM7B - JHDM1F - PHF8	H3K27me2 H4K20me1
KMT7	KMT7 - SETD7	H3K4me1		KDM7C- JHDM1E - PHF2	H3K9me2 H4K20me3
KMT8	KMT8B - PRDM9	H3K4me1/2/3 H3K36me1/2/3	KDM8	KDM8 - JMJD5	H3K9me1 H3K36me1/2
	KMT8A - PRDM2	H3K9me1	KDM9	KDM9 - RSBN1	H4K20me2/3
	KMT8D - PRDM8	H3K9me2			
	KMT8C - PRDM6	H4K20me			
SMYD	SMYD4	H3K4me1/2/3			
	SMYD5	H4K20me3			
SETD	SETD4	H3K4me1/2 H4K20me3			
	SETD5	H3K36me3			
	SETD6	H2AZK7me			
	SETMAR	H3K36me2			
PRDM	PRDM15	H3K9me1			

Open chromatin is characterized by the presence of acetylated histones at several K residues such as H2AK5/9/13/15ac, H2BK5/12/15/20/23/24ac, H3K4/9/14/18/27/36/56/79/122ac, and H4K5/8/12/16/20/79/91ac. Many of these sites are also targets of mutually exclusive methylation and characteristic of active or repressed chromatin. The canonical histone methylation sites are found on H3K4/9/27/36/79 and H4K20, with well-established roles on chromatin functions (Husmann and Gozani 2019) (Figure 1A). For instance, H3K4me1/2/3, H3K36me3, H3K9me1, H3K27me1, H3K79me2, and H4K20me1 are found in enhancers, promoters, and gene bodies of active genes, whereas H3K9me2/3, H3K27me2/3, H4K20me3, and H3K79me3 are found in repressed chromatin such as heterochromatin, telomeric regions and inactivated X chromosome (Husmann and Gozani 2019) (Figure 1A). In addition, there are multiple non-canonical histone methylation sites on H2AZK7me, H3K23/63 and H4K5/12 with unknown functions. Figure 1B shows the most important histone K acetylation and methylation distribution during spermatogenic stages while Figure 1C the stacked RPKM accumulation

139   of enzymes families' expression associated with each histone mark. The accumulated
140   expression patterns for writers and erasers families are consistent with global dynamics
141   described for main histone K marks levels and functions during spermatogenesis, which
142   will be further discussed below.
143

Figure 1. Canonical histone K marks and writers/erasers families distribution during spermatogenesis. A) Histone K methylation PTMs distribution at DNA elements and functional roles in gene expression for an active gene (pink shading) or inactive gene (gray shading). TSS: transcription start site. Only K modifications with sufficient and consensual information about distribution across genes are shown. B) Main histone K PTMs levels involved in chromatin remodeling during spermatogenesis. The data presented is summarized from the following references: Song et al., 2011; Godmann et al., 2007; Zuo et al., 2018; Luense et al., 2016; Wang et al., 2021, Tatehana et al., 2020. C) Accumulated expression of writers/erasers for main canonical histone marks. Legend indicates the maximum number of methylations catalyzed by each enzyme family. SGThy1: spermatogonia Thy1+/Kit-, SGKit: spermatogonia Kit+, SCyte: primary spermatocyte, STid: round spermatid.

2. Clustering analysis of writers and erasers across male germ cell differentiation

Expression profiling analyses of male germ cells were conducted performing k-means partitioning clustering (k=5) (Figure 2A). Then, we searched in each cluster for epigenetic enzymes and marker genes for cell identity and developmental stage. Figure 2B shows that cluster 1 is enriched at SGThy1 cells, showing peak expression of undifferentiated SG marker genes *Thy1*, *Nanos1*, *Nanos2*, *Gfra1*, *Fgfr3*, *Mlh1* and *Cd9*. Cluster 2 is enriched in SGKit cells, showing differentiated SG marker genes *Kit*, *Stra8*, *Dazl*, *H2afv*, *Aldh2* and *Sycp3*. Cluster 3 is enriched in markers of meiosis characteristic of SCyte cells, such as *Sycp1/2*, *Top2a*, *Piwil1*, *Ddx4*, *Dkkl1*, *Spo11* and *H2Afx*. Clusters 4 and 5 correspond to the spermatic stage, characterized by peak expression of transition proteins (TNP) *Tnp1/2*, protamine (PRM) *Prm1/2*, *Izumo1*, acrosome proteins *Acr*, *Acrbp*, *Acrv1*, and *Pgk2*. Cluster 4 showed peak expression at round STid and cluster 5 at sperm cells.

Figure 2. Clustering analysis (k-means) of male germ cells transcriptome. A) Principal

component analysis (PCA) plot showing partitioning clustering results. B) Heatmap showing z-

score from RPKM values of histone K acetylation (Hac) and methylation (Hme) writer/eraser
enzymes (black), and marker genes for male germ cell development (green). SGThy1:
spermatogonia Thy1+/Kit-, SGKit: spermatogonia Kit+, SCyte: primary spermatocyte, STid: round
spermatid.

2.1 Enzymes with peak expression at SGThy1 population

Spermatogonial stem cells (SSC) are characterized by a mitotic activity to maintain
testicular homeostasis through self-renewal during spermatogenesis. Some of these germ
cells are subject to extensive chromatin reorganization and the epigenetic landscape of SG
shifts dramatically to differentiate to committed SGKit. Few variations were found in DNA
methy1ation at promoter regions between SGThy1 and SGKit, suggesting that histone
PTMs may be responsible for the transition between self-renewal and differentiation
(Hammoud et al., 2014). Accordingly, we observed that SGThy1 population showed
different epigenetic enzymes families expression patterns compared to SGKit, having most
of them a peak expression that seems to decrease after SGKit stage.

SGThy1 cluster (Figure 3A) showed peak expression of HATs *Naa60*, *Clock*, *Ncoa3*
and *Supt7l*, none of which have been described yet in the SG population. CLOCK is a
central component of the circadian pacemaker complex that contributes to chromatin-
remodeling events implicated in circadian control of gene expression (Hirayama et al.,
2007). SUPT7L and NCOA3 were found to interact with transcription factors to allow the
RNA polymerase II preinitiation complexes on protein-encoding genes (Demény et al.,
2007; Percharde et al., 2012). Moreover, NCOA3 was related to stem cell self-renewal,
reprogramming, and pluripotency maintenance (Percharde et al., 2012; Percharde and
Azuara, 2013). *Naa60*, which showed the highest RPKM value, was linked to acetylation of
free histone H4K79/91ac and the promotion of nucleosome assembly (Yang et al., 2011).
Among the HDACs family, we found peak expression and high RPKM of *Sirt1/3/5*, and
also *Hdac8* and *Hdac5/7*. SIRT1 is a key regulator of SG differentiation, as SIRT1-
deficient mice show a delay of pre-meiotic differentiation, aberrant expression of
spermatogenic genes, abnormal spermatozoa with elevated DNA damage, and reduced
fertility (Coussens et al., 2008; Bell et al., 2014). SIRT1 also prevents abnormal DNA
methylation of developmental genes by antagonizing DNA methyltransferase DNMT3L
(Geo et al., 2017). SIRT3/5 are mainly mitochondrial but also found in the nucleus linked

to H4K16ac, a mark involved in the formation of higher-order chromatin (Scher et al.,
2007; Giblin et al., 2021). Finally, class IIa HDACs were found to induce pluripotency by
repressing the function of MEF2 (Myocyte Enhancer Factor 2) (Luo et al., 2019).

It is established that undifferentiated SG typically lacks heterochromatin, which forms
as they differentiate to SGKit population committed to meiosis (Chiarini-Garcia et al.,
2002). Self-renewal and maintenance of the undifferentiated SG pool require the
transcriptional repressor PLZF (Promyelocytic Leukaemia Zinc Finger), which mediates
the deposition of repressive marks H3K9me2/3, H3K27me2/3 and H4K20me3 at specific
perinuclear foci (Costoya et al., 2004; Payne and Braun, 2006). These repressive marks
distribution, reflect the epigenomic landscape associated with chromatin silencing
necessary to maintain the stem state and suppress somatic cell differentiation (Payne and
Braun, 2006; Eun et al., 2010). Here, we show that SGThy1 exhibited high expression of
H3K9me2/3 writers *Suv39h2* and *Prdm8*, and H4K20me3 writers *Suv420h*1, *Suv420h2* and
Smyd5, pointing to these enzymes as important drivers of repressive chromatin. We also
found peak expression of *Kdm8*, which was associated with H3K36me erasing at
centromeric regions and mitosis regulation (He et al., 2016), pointing to a novel role of this
enzyme at SGThy1 self-renewal.

Accumulating evidence points to H3K4me2 in the maintenance of transcriptional states
during cell development and, its removal by erasers such as KDM1A/LSD1 is a key step
for epigenetic reprogramming and cell fate (Martinez-Gamero et al., 2021). This is
consistent with the low expression of H3K4me writers found at the SGThy1 stage, as
opposed to the high levels of *Kdm1a* and *Kdm1b* (Figure 1C and 2). In germ cells, KDM1A
is part of a repressive complex that mediates chromatin compaction and transcriptional
repression (Zhang et al., 2013). It was found that KDM1A mediates SG commitment and
differentiation, as testicular deletion of *Kdm1a* leads to progressive germ cell loss and
abnormal accumulation of meiotic SCytes (Lambrot et al., 2015; Myrick et al., 2017).
SGThy1 cluster also showed peak expression of H3K36me2 writers *Nsd1, Smyd2*, and
Setmar. In embryonic stem cells (ESC), NSD1-mediated H3K36me2 was found to prevent
H3K27me3 deposition by polycomb repressive complex 2 (PRC2), modulating polycomb-
mediated H3K27me domains demarcation (Streubel et al., 2018). Moreover, NSD1 was
found to safeguard a subset of genes against H3K27me3-mediated transcriptional silencing

and to have a critical role in de novo DNA methylation in prospermatogonia (Shirane et al.,

2020). In line with this, adult SSC exhibits low H3K27me2/3 levels that increase with cell

differentiation and meiotic progression (Payne and Braun, 2006). In agreement, we found

the highest total H3K36me and lowest H3K27me writers families expression at SGThy1

population (Figure 1C), suggesting that H3K36me2 landscape could be set prior to

increased polycomb activity from SGKit onward. H3K27me erasers *Kdm6b* and *Kdm7a*

showed peak expression in the SGThy1 population and may participate in maintaining low

H3K27me levels. In this context, KDM6B was found to regulate the fragmentation of

spermatogonial cysts, and its down-regulation promotes an increase in undifferentiated SG

without affecting their differentiation (Iwamori et al., 2013). Interestingly, we also

observed high expression of *Sedt6*, a writer that specifically deposits H2AZK7me1, which

in ESC was found with H3K27me3 close to differentiation marker genes and removed upon

retinoic acid (RA) signal, suggesting that SETD6 may have a role in self-renewal and

differentiation of SG (Binda et al., 2013).

2.2 Enzymes with peak expression at SGKit population

SGKit population is highly committed to gametogenesis and express RA-induced

STRA8 (stimulated by retinoic acid 8), consistent with known roles for RA in committing

Kit+ germ cells to meiosis. SGKit population is characterized by the activation of

transcriptional programs that control meiosis entry and post-meiotic differentiation,

consisting of genes that increase and maintain their expression from committed SG, and

genes that are transiently silenced during meiosis and regain expression at round STids

(Chen et al., 2018).

Germ cell differentiation and meiosis entry is driven by increased levels of H3/4ac,

H3K4me2/3 and H3K36me3 (Song et al., 2011; Godmann et al., 2007; Zuo et al., 2018)

(Figure 2B). Here, SGKit population (Figure 3B) showed peak levels of specific HAT

families including GNATs *Gcn5* and *Atf2*, p300/CBP *Ep300* and *Crebbp*, and *Myst4*, *Msl3*

and *Mcm3ap*. Among them, GCN5, CBP and p300 are co-activators for several

transcription factors and nuclear receptors including AR and ERα (Jin et al., 2011).

Moreover, *Gcn5* removal by CRE under *Stra8* promoter disrupted histone eviction,

promoted nucleosome retention in sperm, and affected fertility (Luense et al., 2019).

P300/CBP was linked to metabolism programming of pre- and post-meiotic male germ
cells (Boussouar et al., 2014) and to chromatin reorganization at STids (Pradeepa et al.,
2009). Importantly, it was shown that MSL3, a member of the NSL complex, binds to
H3K36me3 marked sites and deposits H4K16ac, promoting changes in higher order
chromatin structure that prepares meiosis entry and enables STRA8 functions (McCarthy et
al., 2022). In line with this, we also found peak expression of H3K36me2 writers *Ash1l* and
Nsd3, and a specific increase of H3K36me3 writers, including *Setd2/5* and *Prdm9*. PRDM9
and ASH1L were found strongly linked to meiosis, as both enzymes are responsible for
H3K4me3 and H3K36me3 that mediate the double-strand breaks (DSBs) formation at
meiotic hot spots (discussed below) (Sun et al., 2020; Getun et al., 2017).

SGKit population also expressed peak levels of *Hdac2/3*, *Hdac4* and *Sirt6/7*. It was
shown that class IIa HDACs enzymatic activity depends on their recruitment into a
complex containing HDAC3 and RA nuclear receptor (Parra 2015; Fischle et al., 2002),
suggesting that these HDACs could mediate chromatin silencing induced by RA signal.
HDAC3 was also found increased in the last stages of meiosis, inducing repression of
meiotic/spermatogonial genes and activation of postmeiotic haploid gene programs during
meiotic exit, with associated histone acetylation alterations (Yin et al., 2021). On the other
hand, SIRT6 and SIRT7 were found associated with heterochromatic regions and nucleoli,
respectively (Michishita et al., 2005). Deacetylation of H3K9 by SIRT6 modulates
telomeric chromatin function (Michishita et al., 2008) while SIRT7 is highly selective
toward H3K18ac, and might play an upstream role in DNA repair and telomere
maintenance pathway (Barber et al., 2012; Wu et al., 2018).

SGKit population showed peak expression of the KMT2 members *Setd1a, Mll2* and
Mll3, which are part of the COMPASS complex family that display both redundant and
non-redundant H3K4me deposition. In mammalian cells, SETD1A/B reportedly catalyzes
most of the H3K4me3 (Couture and Skiniotis, 2013) while MLL3 deposits H3K4me1 at
enhancers (Herz et al., 2012) and MLL1/2 H3K4me2/3 at developmental genes (Hu et al.,
2017; Rickels et al., 2016). Importantly, targeted loss of H3K4me methylation in SSC by
reduction of MLL2 activity results in a dramatic reduction in the number of SCytes (Glaser
et al., 2009), and manipulating H3K4me3 patterns can induce transgenerational inheritance
(Lismer et al., 2020). During the SGKit stage, certain genes such as *Stra8*, *Aldh2* and *Spo11*

gain expression by losing H3K27me3 bivalency and transitioning from a bivalent to active

chromatin state (Hammoud et al., 2014). In line with this, we detected peak expression of

H3K27me2/3 erasers *Kdm6a* and *Kdm7b* in the SGKit population. Interestingly, KDM6A is

also a component of the MLL2/MLL3 COMPASS complex that mediates enhancer

activation via simultaneous H3K4me3 deposition and H3K27me2/3 removal (Fagan and

Dingwall, 2019). Moreover, the COMPASS complex promotes H3K27ac by interacting

with p300/CBP (Fagan and Dingwall, 2019). Taken together, these data could reveal the

importance of MSL and COMPASS complexes during meiosis entry, deserving further

future attention.

SGKit stage also showed peak expression of repressive marks writers indicative of

facultative heterochromatin and cell differentiation, including KMT1 H3K9me2 *Ehmt1* and

H3K9me3 *Setdb1/2*, and KMT6 H3K27me2/3 polycomb *Ezh1/2* families. At this stage

there is de novo formation of bivalent domains by H3K27me deposition at developmental

genes (Maezawa et al., 2018). Her, we detected peak expression of the H4K20me1 writer

Setd8, which was found to physically interact with STRA8 and to mutually control each

other transcription (Niu et al., 2020). Moreover, SETD8, H4K20me1 and PCNA co-localize

with STRA8, indicating a role for SETD8 in the cell cycle transition from mitosis to

meiosis (Niu et al., 2020). We also found peak expression of H3K36me *Kdm2a* and

H3K9/K36me *Kdm4a/b/c* erasers. KDM2A was strongly linked to heterochromatin

maintenance, transcriptional repression of repetitive DNA elements and inactive genes, and

the proper segregation of chromosomes during mitosis (Frescas et al., 2008), but the exact

roles during spermatogenesis remain to be further explored. On the other hand, the KDM4

family participates in DNA replication control, where KDM4B is linked to core

transcription factors NANOG and SOX2 and KDM4C to PRC2 and Myc regulatory

modules (Das et al., 2014). Interestingly, in meiotic and post-meiotic SCytes, KDM4A/B

were found to remove H3K36me2/3 after binding to H3K23me3, a poorly understood

histone modification recently shown to be enriched in heterochromatin at meiosis onset (Su

et al., 2016). Here, we also detected high expression of the KDM5 family which removes

H3K4me2/3. KDM5A is an integral, conserved component of the Notch/RBP-J repressor

complex and fundamental for the switch-off of the Notch target genes that control cell fate

transitions (Liefke et al., 2010.). Among the KDM5 family, KDM5A/D were found

strongly related to spermatogenesis progression and fertility (Nishio et al., 2014;
Gegenschatz-Schmid et al, 2019). Importantly, KDM5D forms a complex with the MSH5
DNA repair factor that controls synapsis and crossing-over at meiotic prophase and locates
to condensed DNA during the leptotene/zygotene stage (Akimoto et al., 2008).

A *Cluster 1*: SGThy1 population

B *Cluster 2*: SGKit population

Figure 3. Cluster 1 and 2 epigenetic enzymes showing peak expression at SGThy1 (A) and

SGKit population (B). Left: heatmap showing z score values for each male germ cell population.

Right: Dot plot showing enzymes RPKM values in SGThy1 (A) and SGKit (B) population,

repetively. SGThy1: spermatogonia Thy1+/Kit-, SGKit: spermatogonia Kit+, SCyte: primary

spermatocyte, STid: round spermatid.

2.3 Enzymes with peak expression at primary SCyte population

Meiotic division is associated with specific chromatin changes that enable genetic

exchange at meiotic recombination: i) DNA DSBs deposition and repair at synapsed

chromatin, and ii) epigenetic silencing of unpaired chromatin (MSUC), including meiotic

sex chromosome inactivation (MSCI). Several studies showed that histone PTMs contribute

to synapsis formation, recombination at hot spots and to MSCI and sex body formation

(Buard et al., 2009; Baudat et al., 2010; Parvanov et al., 2010).

Hot spot cores of SG and pre-leptotene-stage cells harbor H3K9/18/23ac and

H4K5/8/12/16/91ac that are associated with open chromatin (Getun et al., 2017;

Chioccarelli et al., 2020). We observed that SCyte cluster (Figure 4A) showed peak

expression of MYST family *Myst1, Myst2* and *Tip60*, and also *Naa40*, *Brca2* and *Gtf3c4,*

showing most of them high RPKM values. Accordingly, MYST enzymes were previously

found strongly related to meiosis control, increasing their expression from early pachytene

through diplotene stages (Getun et al., 2017). Moreover, MYST1/MOF, a specific

H4K16ac writer, was found involved in DSBs repair and crossover numbers (Jiang et al.,

2018). In line with this, H4K91ac was also linked to chromatin assembly of repair

machinery (Ye et al., 2005). In addition, we detected peak expression of the H3K36me3

writer *Setd2*, which has been also associated with DSBs establishment and repair in

transcribed genes (Jha et al., 2014). Importantly, *Setd2* knockout in germ cells results in

complete loss of H3K36me3, decreased expression of genes encoding PRMs and altered

spermiogenesis with acrosomal malformation at the round STid stage (Zuo et al., 2018).

During MSCI, both sex chromosomes undergo extensive H3/4 deacetylation and

H3K9me2/3 marking to create the sex body (Khalil et al., 2004). At the SCyte stage, we

detected peak expression of *Hdac1* that was shown to form a complex with DNA

methyltransferase DNMT3L to regulate X chromosome compaction (Zamudio et al., 2011;

Deplus et al., 2002). Upon meiosis completion, there is an increase in H3K4me3 and a drop

in H3/H4ac and H3K9me3 (Figure 1B). Here, we found high RPKM values for

H3K9me2/3 erasers *Kdm3a* and *Kdm4d* that remain elevated in the STid stage. In line with

this, H3K9me demethylation by KDM3A was found essential for post-meiotic chromatin

organization, enabling the transcriptional activation of TNP1 and PRM1 (Okada et al.,

2007). SCyte stage also showed peak expression and high RPKM values of the H3K4me

writer *Ash2l*, a key core member of the WRAD complex that interacts with KMT2

COMPASS writers to confer methyltransferase activity (Ernst and Vakoc, 2012).

Interestingly, this enzyme also showed high expression at SGThy1, consistent with

previous work showing that *Ash2l* controls ESC maintenance and differentiation (Wan et

al., 2013). Among H3K4me erasers, we found high RPKM values of *Kdm5b*, strongly

linked to spermatogonial differentiation. Moreover, *Kdm5b* is highly expressed in pre-

meiotic SG and pachytene-diplotene SCytes, where it is thought to repress the expression

of genes that would otherwise cause the SG to remain in the mitotic phase (Madsen et al.,

2003).

2.4 Enzymes with peak expression at round STid population

Spermiogenesis is the final phase of spermatogenesis, where haploid round STids

undergo dramatic morphological changes to transform into elongated STids and sperm.

During this phase, extreme chromatin remodeling takes place to facilitate the condensation

and packaging of the paternal genome. Spermiogenesis involves transient DSBs formation

and repair, specific histone variants incorporation and H3/4 hyperacetylation to destabilize

the nucleosome, and subsequent replacement of 95-99% histones by TNPs and PRMs

(Rajender et al., 2011, Carrel 2012).

During spermiogenesis, H4K16ac is essential for histone eviction which does not occur

in the absence of this specific mark (Lu et al., 2010). Here, SCyte cluster (Figure 4A)

showed peak transcriptional levels of MYST enzymes *Tip60* and *Myst1*, which were linked

to H4ac prior to histone eviction (Dong et al., 2017, Lu et al., 2010). In line with this, it was

previously reported that pachytene SCytes show higher expression of genes related to

spermiogenesis than round STids, showing a gap between the time of peak transcription

and the execution of certain programs (Da Cruz et al., 2016). The existence of translational

delays in spermatogenesis is well known and has been associated with a strategy to regulate the timing of protein synthesis in STids, as transcription in these cells is silenced after histone eviction (Kleene et al., 2001; Idler and Yan, 2012; Braun 1998; Lee et al., 1995). A characteristic of STid enzymes is that they also show high expression at SGKit with or without transient silencing at SCyte stage, in line with the notion that many STid genes begin to express at SG population (Jan et al., 2017). In a very interesting study, Bell et al., (2014) found that SIRT1 deletion by CRE under protamine promoter has no effect, but CRE under *Stra8* promoter not only dampens SG differentiation but STids show reduced H4K5/8/12ac and TNP2 do not localize in the nucleus leading to chromatin condensation defects in sperm. Similar results were reported for GCN5 (Luense et al., 2019). Therefore, the action of certain epigenetic enzymes in pre-meiotic stages seems to enable the post-meiotic transcriptome and chromatin remodeling processes.

Histone eviction happens in an initial wave when nucleosomes become accessible at the round STid stage followed by a "major" wave of nucleosome accessibility at elongated STids (Luense et al., 2019). The specific epigenetic programs controlling the histone eviction waves are not fully understood, but different HATs activities are likely to be involved. It is important to note that SGKit and SCyte cluster HATs show moderate/high expression at STid population as well, suggesting they could be involved in the acetylation waves prior to histone eviction, except for p300/CBP that has been already discarded (Boussouar et al., 2014). Here, STid cluster (Figure 4B) showed peak expression of *Pcaf*, *Ncoa1* and *Hat1*. PCAF functions as a co-activator for ERα and RA receptor (RAR) (Blanco et al, 1998), and can interact with NCOA1 to promote transcription (Spencer et al., 1997). Interestingly, p300/CBP and PCAF were found to regulate histone eviction by acetylation of TNP2, affecting DNA condensation properties and interaction with histone chaperones (Pradeepa et al., 2009). Concerning *Hat1* expression, a cytoplasmic HAT responsible for acetylating newly synthesized H4, it was found related to the incorporation of H4K5/12ac and H3.3 variant at DSBs sites and the promotion of DNA repair (Yang et al., 2013).

STid population showed peak expression of H3K9me2/3 writers *Ehmt2* and *Suv39h1*, which have been previously found related to heterochromatin formation during meiosis (Tachibana et al., 2007). Importantly, deficiency of these enzymes totally disrupts SCytes

maturation and induces cell arrest (Tachibana et al., 2007, Peters et al., 2001). Therefore,
given their crucial role in the meiotic stage, the involvement of these KMT1 enzymes in
STids may have been overlooked. At this stage, there is a reactivation of transcription from
MSUC and MSCI sites that is enabled by the deposition of histone crotonylation (Kcr). It
was proposed that CDYL, an HKcr eraser, prevents post-meiotic chromatin reactivation by
binding to H3K9me3 and H3K27me2/3 marked sites and facilitating H3K9me2 deposition
by EHMT2 (Mulligan et al., 2008). We also observed high expression of KMT *Prdm2* that
was found in an activation complex with p300 that drives Hac upon ER binding (Di Zazzo
et al., 2013). *Nsd2* and *Ezh2* presented highly similar RPKM values in SGKit, SCyte and
STid population, and we actually consider that the real peak expression of these enzymes
comprises the three stages. At STid stage there is specific H3K27me deposition to stablish
bivalency at developmental genes (Maezawa et al., 2018), and proteomic studies found
substantial H3K27me3/K36me2 double marking in STid and sperm (Brunner et al., 2014;
Luense et al., 2016). EZH2 and NSD2 were found to form a regulatory axis that promotes
the oncogenic phenotype, where EZH2 acts upstream NSD2 regulating its expression and
H3K36me2-mediated chromatin activation by a microRNA network (Asangani et al., 2013;
Popovic et al., 2014). Hence, through this axis, EZH2 repressive marking leads to
deposition of activating H3K36me2 marking, showing yet another level of complexity in
the epigenetic landscape dynamics of cells. Additionally, the γH2AX-MDC1 signaling
pathway is involved in the recruitment of NSD2 to DSBs, where it catalyzes H4K20me2
methylation involved in DNA damage repair pathway (Pei et al., 2011).

2.5 Enzymes with peak expression at mature sperm population

The mature sperm transports a cargo of RNAs that have been the focus of intense
attention, composed of non-coding RNAs like miRNAs, lncRNAs, tRNAs and circRNAs,
and protein-coding mRNAs (Guo et al., 2017; Jodar et al., 2013). Some of these RNAs are
remnants of the last stages of elongated STids, and others are acquired along the passage
through the epididymis, as sperm absorb epididymosomes released by somatic cells (Trigg
et al., 2019). Sperm RNA sequencing shows a set of mRNAs, from marker genes of the
spermatogenic process (Figure 2B), known as the "memory of spermatogenesis", and many

other RNAs that were found to carry instructions for early embryo development and
metabolic and behavioral traits in the offspring (Guo et al., 2017; Jodar et al., 2013).

We found a very specific pattern in sperm cluster (Figure 4C) of HDACs including
Hdac9, *Sirt2*, and *Hdac1*, and peak RNA values for HATs *Naa50* and *Myst3*. Importantly,
HDACs are also highly detected in human late STids from integrated single-cell RNAseq
data (Figure S1). There is no information concerning possible roles for the HATs and
HDACs of the sperm cluster, but HDAC11 was linked to early embryo development, as
will be discussed below. On the other hand, we observed peak expression of the H3K79me
writer *Dot1l*, which was found expressed throughout embryogenesis and with apparently
higher levels in the ovary and testis (Barry et al., 2010). Moreover, in ESCs, DOT1L may
play a role in telomere maintenance since H3K79me2 is enriched at constitutive telomeric
and centromeric heterochromatin (Jones et al., 2008). The DOT1L binding to the H4 tail is
regulated by H4K16ac, and also deletion or overexpression of DOT1L leads to
mislocalization of the SIRT complex, causing silencing defects (Farooq et al., 2016). Here,
we also detected the H3K9me1/2 eraser *Kdm3b*, which would reflect other contributions
than spermatogenesis, as this enzyme is highly expressed by the epididymal epithelium
(Liu et al., 2015) and therefore, likely enriched in epididymosomes (Trigg et al., 2019).
Finally, we also detected *Kdm2b* expression, which erases H3K4/K36me and was found to
protect the polycomb-silenced promoters against ectopic de novo methylation (Boulard et
al., 2015). Furthermore, KDM2B is identified as one subunit of non-canonical PRC1.1
complex (Wang et al., 2018), consistent with the last stages of spermiogenesis, when
Polycomb KMT6 family and H3K27me2/3 expression are the highest (Figure 1B, C).
Importantly, KDM2B was recently found to erase H3K79me2/3 and to induce
transcriptional repression via SIRT1 chromatin silencing (Kang et al., 2018). Here, we also
found *Setd7*, previously associated with mesoderm lineage differentiation, muscle and
cardiac development (Bultman et al., 2019).

A *Cluster 3*: primary SCyte population

B *Cluster 4*: round STid population

C *Cluster 5*: mature Sperm population

Figure 4. Cluster 3, 4 and 5 epigenetic enzymes showing peak expression at primary SCyte (A), round STid (B) and mature sperm (B) populations. Left: heatmap showing z score values for each male germ cell population. Right: Dot plot showing enzymes RPKM values in primary SCyte (A), round STid (B) and sperm (C) population, respectively. SGThy1: spermatogonia Thy1+/Kit-, SGKit: spermatogonia Kit+, SCyte: primary spermatocyte, STid: round spermatid.

2.6 Sperm enzymes: potential contribution to early embryo development

Recent findings indicate that the sperm delivers RNAs to the oocyte at fertilization which stay stable until the activation of the embryonic genome, having putative roles at zygote formation (Boerke et al., 2007; Smith and Spadafora, 2005). Early molecular processes in the zygote include differential epigenetic reprogramming of the parental genomes, where PRMs are quickly released from the paternal genome to be repackaged by maternal nucleosomal histones (McLay and Clarke, 2003). The zygote genome is translationally silent, and cellular processes are carried on with inherited RNAs and proteins until the onset of zygotic genome activation (ZGA) around the 2-4 cells stage (Schulz and Harrison, 2019). In this sense, Guo et al. (2017) characterized that sperm-carried RNAs are enriched in genes associated with the ontologies of embryonic cleavage and blastocyst formation. For example, it was shown that coding mRNA for the egg-activating factor PLC-zeta, is inherited by the father and participates in early embryo development (Yao et al., 2010; Sone et al., 2005).

Here, we evaluated the possible paternal contribution of epigenetic enzymes to the zygote, by analyzing RNAseq data from meiosis II (MII) oocytes and pre-cleavage zygotes (Wu et al., 2016). We searched for enzymes showing increased RPKM values in zygotes compared to MII-oocytes (Fold > 1, Table 2), indicative of potential paternal contribution to the zygotic mRNA pool. This approach allowed us to detect many marker mRNAs previously described as high in sperm and with roles early in embryo development (Guo et al., 2017), as well as many enzymes with peak expression in the different clusters that were also present in sperm population (Table 2). Remarkably, we detected several enzymes from the sperm cluster, including *Naa50*, *Hdac11*, *Sirt2,* and *Setd7*. In this context, previous work *in silico* identified *Hdac11* as a paternally-inherited mRNA that could potentially interact co-operatively and synergistically with maternal factors during zygote formation (Ntostis et al., 2017). Among enzymes of clusters 1-4, we found *Hdac1* expression, which

was previously linked to cell cycle regulation and ZGA during pre-implantation (Ma et al.,
2008). We also detected key enzymes involved in chromatin structure organization: the
H4K20me2/3 writer *Suv420h2*, the H3K9me3 writers *Suv39h1/2* and *Setdb2*, and
H3K9me2 writer *Ehmt2*. In line with this, the active mark H3K4me3 is detected in the male
pronucleus, but H3K9me2/3, H3K27me3 and H4K20me3, are mostly absent (Lepikhov et
al., 2010). The parental pronuclei have asymmetric reprogramming capacities and the
reprogramming factors reside predominantly in the male pronucleus. In this context, it was
found that EHMT2 activates soon after fertilization and deposit H3K9me2 patterns in the
paternal genome (Ma et al., 2015). Also, previous work suggested that SUV420H2 deposits
H4K20me3 to allow the timely and coordinated progression of replication after fertilization
(Eid et al., 2016). Finally, we also detected several other writers and erasers with unknown
functions at pre-cleavage embryo.

Symbol	Sperm	MII_oocyte	Zygote	Fold
Clock	3.14	11.86	12.37	1.04
Myst1	28.90	9.98	25.49	2.55
Gtf3c4	18.18	0.24	0.27	1.13
Pcaf	6.12	0.03	0.07	2.45
Naa50	14.01	35.67	99.97	2.80
Hdac1	1.91	0	1.73	-
Hdac11	178.31	0	0.04	-
Sirt1	3.63	2.23	3.84	1.73
Sirt2	96.57	0.36	1.96	5.45
Sirt5	1.96	1.82	2.79	1.53
Sirt7	2.64	6.89	10.22	1.48
Ash2l	48.31	9.88	13.33	1.35
Setd7	4.14	0.01	0.01	1.44
Setd4	1.29	7.11	13.96	1.96
Nsd1	10.31	14.68	20.44	1.39
Nsd3	7.16	1.90	2.35	1.24
Prdm9	1.14	0.06	0.25	4.27
Setd5	14.71	0.71	1.05	1.47
Ehmt2	67.83	0	0.03	-
Suv39h1	1.36	1.54	3.61	2.34
Suv39h2	13.08	4.73	9.82	2.08
Setdb2	7.09	6.53	11.54	1.77
Suv420h1	7.64	0.47	1.30	2.78
Kdm5a	11.82	0.19	0.29	1.50
Kdm5b	42.30	0.74	1.30	1.76
Kdm5d	1.06	0	0.02	-
Kdm4c	3.32	0.02	0.10	6.43
Kdm7a	5.53	20.61	21.80	1.06
Aatf	61.530	2.866	4.098	1.430
Pik3ca	4.663	2.301	2.395	1.041
Rpl7l1	7.437	0.631	3.203	5.073
Rad51	1.937	66.832	78.557	1.175
Nek2	64.205	10.929	34.466	3.154
Itch	18.287	5.405	5.926	1.096
Serpine1	0.740	0	0.016	-
Sp1	10.705	0.728	2.144	2.945
Cacna1g	0.579	0.008	0.017	2.157
Pnrc1	8.158	6.300	38.548	6.119
Ltbp1	1.727	0	0.007	-
Map1lc3a	1858.090	0.566	1.415	2.501
Zfp821	495.639	5.905	6.054	1.025

Table 2: RPKM values of writers and erasers in mature sperm, meiosis II oocytes and pre-cleavage zigotes. Fold represent the ratio between zygote and MII oocytes RPKMs. Blue: writers, red: erasers, green: marker genes for paternal contribution to zygote RNA pool.

DISCUSSION

A new trend in research points toward sperm epigenome contribution to embryonic development and offspring health. However, few links were established between alterations in sperm histone PTMs, the specific enzymes responsible, and the developmental window where these marks may be reprogrammed. We show for the first time a comprehensive evaluation of all known families of histone acetylation and methylation writers and erasers from undifferentiated SG to mature sperm. Not only we accurately detected germ cell markers genes, but also by the rule of "guilty by association", enzymes with known functions and many others not even described yet in male germ cells. The analysis performed here shows important windows during spermatogenesis, where interference with epigenetic enzymes gene expression may have phenotypic consequences in the offspring, even when they are not inherited.

Many studies have implicated different histone PTMs in sperm chromatin as substrates for transgenerational inheritance (Skvortsova et al., 2018, Brunner et al., 2014, Hammoud et al., 2009; Brykczynska et al., 2010). Altered H3K4me3 in sperm was associated with diet-induced phenotypes in the offspring and developmental defects (Lismer et al., 2021). In *C elegans*, inheritance of H3K27me3-lacking sperm chromatin resulted in lost repression of neuronal genes from sperm-inherited alleles, which primed germ cells for losing their identity and adopting a neuronal fate (Kaneshiro et al., 2019). In *Drosophila,* paternal restraint stress affected offspring metabolism via ATF2-dependent mechanisms (Seong et al., 2020). H4K8/12ac was found prior to full decondensation of the sperm nucleus, suggesting that these marks are transmitted to the zygote (van der Heijden et al., 2006). Moreover, it was recently reported that alterations in the expression of paternal KDM1A/LSD1 and KDM6A enzymes were able to induced defects that persisted transgenerationally (Siklenka et al., 2015, Lesch et al., 2019).

Finally, we report that mature sperm carries several histone-modifying enzymes

mRNAs with moderate to high RPKM values. Moreover, some of these enzymes' mRNAs

were detected higher in the zygote than in the transcriptionally silent MII oocyte, pointing

to a paternal contribution. Many pieces of evidence showed that sperm RNAs, in particular

miRNAs and tsRNAs, can mediate intergenerational transmission of paternally acquired

phenotypes such as diet-induced metabolic and neurological phenotypes (Chen et al.,

2016). Moreover, it was proposed that their mechanisms involve changes in chromatin

structure and DNA methylation. Dynamic chromatin remodeling by histone PTMs occurs

after fertilization to facilitate the transition from gametic-to-embryonic chromatin, and

inherited mRNAs from the gametes are selectively translated before the ZGA. The data

presented here, show that epigenetic enzymes mRNA could become functional in pre-

cleavage zygotes and contribute to early chromatin organization with deep implications in

future embryo development. The epigenetic enzyme's paternal contribution could be

another mechanism for epigenetic inheritance that deserves further consideration.

METHODS

Data retrieval, cleaning and processing

Data was manipulated using Rstudio (v1.2.1033) and Tidyverse packages Dplyr and

ggplot2 (Wickham et al., 2019). Male germ cell processed RNA-seq data tables (RPKM)

were downloaded from NCBI GEO database, deposited by Hammoud et al., (2014), under

accession number GSE49622. Data contained one replica of SGThy, spermatogonia SGKit

and mature sperm, and 5 replicas of primary SCyte and round STid. We removed 54 extra

genes in SCyte, STid and sperm that were not included in the SG tables. The mean value

for each gene was calculated in SCyte and STid samples, and generated a data matrix by

merging the mean SCyte and mean STid with SGThy1, SGKit and sperm samples (RPKM

matrix). The accumulated RPKM for each enzyme family was calculated and generated

stacked bar plots. To perform clustering analysis, RPKM matrix was converted to z score

by gene and performed k-means clustering from the R Base Stats package, selectig k=5 and

nstart = 25. We generated an enzyme metadata table, and retrieved the genes of interest

RPKM values and cluster assigned. Heatmap plots were plotted using pheatmap package

from R base. The complete samples tables, enzymes metadata table and R scripts used in

this study are available for download and further editing at https://github.com/Gonzalez-
Lab/Gonzalez-2022-germ-cells. Finally, meiosis II oocyte and zygote accessed RNA-seq
data tables (RPKM) were downloaded from NCBI GEO database, deposited by Wu et al.,
(2016), under accession number GSE66582.

Author contributions

597 B.G and C.R.G conceived the project, analyzed the data and wrote the manuscript. C.P.L
598 contributed to data mining and analysis. A.D.V contributed to the critical reading and
599 editing of the manuscript. All authors read and approved the submitted version.

Declaration of interests

602 The authors declare no competing interests.

REFERENCES

Akimoto, C., Kitagawa, H., Matsumoto, T., & Kato, S. (2008). Spermatogenesis-specific
association of SMCY and MSH5. *Genes to cells : devoted to molecular & cellular*
mechanisms, *13*(6), 623–633. https://doi.org/10.1111/j.1365-2443.2008.01193.x

Asangani, I. A., Ateeq, B., Cao, Q., Dodson, L., Pandhi, M., Kunju, L. P., Mehra, R., Lonigro, R. J.,
Siddiqui, J., Palanisamy, N., Wu, Y. M., Cao, X., Kim, J. H., Zhao, M., Qin, Z. S., Iyer, M. K.,
Maher, C. A., Kumar-Sinha, C., Varambally, S., & Chinnaiyan, A. M. (2013). Characterization of
the EZH2-MMSET histone methyltransferase regulatory axis in cancer. Molecular cell, 49(1),
80–93. https://doi.org/10.1016/j.molcel.2012.10.008

Bale T. L. (2015). Epigenetic and transgenerational reprogramming of brain development. *Nature*
reviews. Neuroscience, 16(6), 332–344. https://doi.org/10.1038/nrn3818

Barber, M. F., Michishita-Kioi, E., Xi, Y., Tasselli, L., Kioi, M., Moqtaderi, Z., Tennen, R. I.,
Paredes, S., Young, N. L., Chen, K., Struhl, K., Garcia, B. A., Gozani, O., Li, W., & Chua, K. F.
(2012). SIRT7 links H3K18 deacetylation to maintenance of oncogenic
transformation. *Nature*, *487*(7405), 114–118. https://doi.org/10.1038/nature11043

Barry, E. R., Corry, G. N., & Rasmussen, T. P. (2010). Targeting DOT1L action and interactions in leukemia: the role of DOT1L in transformation and development. Expert opinion on therapeutic targets, 14(4), 405–418. https://doi.org/10.1517/14728221003623241

Baudat, F., Buard, J., Grey, C., Fledel-Alon, A., Ober, C., Przeworski, M., Coop, G., & de Massy, B. (2010). PRDM9 is a major determinant of meiotic recombination hotspots in humans and mice. Science (New York, N.Y.), 327(5967), 836–840. https://doi.org/10.1126/science.1183439

Bell, E. L., Nagamori, I., Williams, E. O., Del Rosario, A. M., Bryson, B. D., Watson, N., White, F. M., Sassone-Corsi, P., & Guarente, L. (2014). SirT1 is required in the male germ cell for differentiation and fecundity in mice. *Development (Cambridge, England)*, 141(18), 3495–3504. https://doi.org/10.1242/dev.110627

Binda, O., Sevilla, A., LeRoy, G., Lemischka, I. R., Garcia, B. A., & Richard, S. (2013). SETD6 monomethylates H2AZ on lysine 7 and is required for the maintenance of embryonic stem cell self-renewal. *Epigenetics*, 8(2), 177–183. https://doi.org/10.4161/epi.23416

Black, J. C., Van Rechem, C., & Whetstine, J. R. (2012). Histone lysine methylation dynamics: establishment, regulation, and biological impact. *Molecular cell*, 48(4), 491–507. https://doi.org/10.1016/j.molcel.2012.11.006

Blanco, J. C., Minucci, S., Lu, J., Yang, X. J., Walker, K. K., Chen, H., Evans, R. M., Nakatani, Y., & Ozato, K. (1998). The histone acetylase PCAF is a nuclear receptor coactivator. Genes & development, 12(11), 1638–1651. https://doi.org/10.1101/gad.12.11.1638

Boerke, A., Dieleman, S. J., & Gadella, B. M. (2007). A possible role for sperm RNA in early embryo development. Theriogenology, 68 Suppl 1, S147–S155. https://doi.org/10.1016/j.theriogenology.2007.05.058

Boulard, M., Edwards, J. R., & Bestor, T. H. (2015). FBXL10 protects Polycomb-bound genes from hypermethylation. Nature genetics, 47(5), 479–485. https://doi.org/10.1038/ng.3272

Boussouar, F., Goudarzi, A., Buchou, T., Shiota, H., Barral, S., Debernardi, A., Guardiola, P., Brindle, P., Martinez, G., Arnoult, C., Khochbin, S., & Rousseaux, S. (2014). A specific CBP/p300-dependent gene expression programme drives the metabolic remodelling in late stages of spermatogenesis. *Andrology*, *2*(3), 351–359. https://doi.org/10.1111/j.2047-2927.2014.00184.x

Braun R. E. (1998). Post-transcriptional control of gene expression during spermatogenesis. Seminars in cell & developmental biology, 9(4), 483–489. https://doi.org/10.1006/scdb.1998.0226

Brunner, A. M., Nanni, P., & Mansuy, I. M. (2014). Epigenetic marking of sperm by post-translational modification of histones and protamines. Epigenetics & chromatin, 7(1), 2. https://doi.org/10.1186/1756-8935-7-2

Brykczynska, U., Hisano, M., Erkek, S., Ramos, L., Oakeley, E. J., Roloff, T. C., Beisel, C., Schübeler, D., Stadler, M. B., & Peters, A. H. (2010). Repressive and active histone methylation mark distinct promoters in human and mouse spermatozoa. Nature structural & molecular biology, 17(6), 679–687. https://doi.org/10.1038/nsmb.1821

Buard, J., Barthès, P., Grey, C., & de Massy, B. (2009). Distinct histone modifications define initiation and repair of meiotic recombination in the mouse. The EMBO journal, 28(17), 2616–2624. https://doi.org/10.1038/emboj.2009.207

Bultman S. J. (2019). SETD7 interacts with other chromatin-modifying factors to regulate cardiac development. *Stem cell investigation*, 6, 14. https://doi.org/10.21037/sci.2019.05.03

Chen, Q., Yan, W., & Duan, E. (2016). Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. Nature reviews. Genetics, 17(12), 733–743. https://doi.org/10.1038/nrg.2016.106

Chen, Y., Zheng, Y., Gao, Y., Lin, Z., Yang, S., Wang, T., Wang, Q., Xie, N., Hua, R., Liu, M., Sha, J., Griswold, M. D., Li, J., Tang, F., & Tong, M. H. (2018). Single-cell RNA-seq uncovers dynamic processes and critical regulators in mouse spermatogenesis. *Cell research*, *28*(9), 879–896. https://doi.org/10.1038/s41422-018-0074-y

Chiarini-Garcia, H., & Russell, L. D. (2002). Characterization of mouse spermatogonia by transmission electron microscopy. *Reproduction (Cambridge, England)*, *123*(4), 567–577. https://doi.org/10.1530/rep.0.1230567

Costoya, J. A., Hobbs, R. M., Barna, M., Cattoretti, G., Manova, K., Sukhwani, M., Orwig, K. E., Wolgemuth, D. J., & Pandolfi, P. P. (2004). Essential role of Plzf in maintenance of spermatogonial stem cells. *Nature genetics*, *36*(6), 653–659. https://doi.org/10.1038/ng1367

Coussens, M., Maresh, J. G., Yanagimachi, R., Maeda, G., & Allsopp, R. (2008). Sirt1 deficiency attenuates spermatogenesis and germ cell function. *PloS one*, *3*(2), e1571. https://doi.org/10.1371/journal.pone.0001571

Couture, J. F., & Skiniotis, G. (2013). Assembling a COMPASS. Epigenetics, 8(4), 349–354. https://doi.org/10.4161/epi.24177

da Cruz, I., Rodríguez-Casuriaga, R., Santiñaque, F. F., Farías, J., Curti, G., Capoano, C. A., Folle, G. A., Benavente, R., Sotelo-Silveira, J. R., & Geisinger, A. (2016). Transcriptome analysis of highly purified mouse spermatogenic cell populations: gene expression signatures switch from

meiotic-to postmeiotic-related processes at pachytene stage. BMC genomics, 17, 294.
https://doi.org/10.1186/s12864-016-2618-1

Das, P. P., Shao, Z., Beyaz, S., Apostolou, E., Pinello, L., De Los Angeles, A., O'Brien, K., Atsma,
J. M., Fujiwara, Y., Nguyen, M., Ljuboja, D., Guo, G., Woo, A., Yuan, G. C., Onder, T., Daley,
G., Hochedlinger, K., Kim, J., & Orkin, S. H. (2014). Distinct and combinatorial functions of
Jmjd2b/Kdm4b and Jmjd2c/Kdm4c in mouse embryonic stem cell identity. *Molecular cell*, *53*(1),
32–48. https://doi.org/10.1016/j.molcel.2013.11.011

Demény, M. A., Soutoglou, E., Nagy, Z., Scheer, E., Jànoshàzi, A., Richardot, M., Argentini, M.,
Kessler, P., & Tora, L. (2007). Identification of a small TAF complex and its role in the assembly
of TAF-containing complexes. *PloS one*, *2*(3), e316.
https://doi.org/10.1371/journal.pone.0000316

Deplus, R., Brenner, C., Burgers, W. A., Putmans, P., Kouzarides, T., de Launoit, Y., & Fuks, F.
(2002). Dnmt3L is a transcriptional repressor that recruits histone deacetylase. Nucleic acids
research, 30(17), 3831–3838. https://doi.org/10.1093/nar/gkf509

Di Zazzo, E., De Rosa, C., Abbondanza, C., & Moncharmont, B. (2013). PRDM Proteins:
Molecular Mechanisms in Signal Transduction and Transcriptional Regulation. Biology, 2(1),
107–141. https://doi.org/10.3390/biology2010107

Dong, Y., Isono, K. I., Ohbo, K., Endo, T. A., Ohara, O., Maekawa, M., Toyama, Y., Ito, C.,
Toshimori, K., Helin, K., Ogonuki, N., Inoue, K., Ogura, A., Yamagata, K., Kitabayashi, I., &
Koseki, H. (2017). EPC1/TIP60-Mediated Histone Acetylation Facilitates Spermiogenesis in
Mice. Molecular and cellular biology, 37(19), e00082-17. https://doi.org/10.1128/MCB.00082-17

Eid, A., Rodriguez-Terrones, D., Burton, A., & Torres-Padilla, M. E. (2016). SUV4-20 activity in
the preimplantation mouse embryo controls timely replication. Genes & development, 30(22),
2513–2526. https://doi.org/10.1101/gad.288969.116

Ernst, P., & Vakoc, C. R. (2012). WRAD: enabler of the SET1-family of H3K4 methyltransferases.
Briefings in functional genomics, 11(3), 217–226. https://doi.org/10.1093/bfgp/els017

Eun, S. H., Gan, Q., & Chen, X. (2010). Epigenetic regulation of germ cell differentiation. *Current
opinion in cell biology*, *22*(6), 737–743. https://doi.org/10.1016/j.ceb.2010.09.004

Fagan, R. J., & Dingwall, A. K. (2019). COMPASS Ascending: Emerging clues regarding the roles
of MLL3/KMT2C and MLL2/KMT2D proteins in cancer. *Cancer letters*, 458, 56–65.
https://doi.org/10.1016/j.canlet.2019.05.024

Fan, J., Krautkramer, K. A., Feldman, J. L., & Denu, J. M. (2015). Metabolic regulation of histone
post-translational modifications. *ACS chemical biology*, 10(1), 95–108.
https://doi.org/10.1021/cb500846u

Farooq, Z., Banday, S., Pandita, T. K., & Altaf, M. (2016). The many faces of histone H3K79
methylation. Mutation research. Reviews in mutation research, 768, 46–52.
https://doi.org/10.1016/j.mrrev.2016.03.005

Fischle, W., Dequiedt, F., Hendzel, M. J., Guenther, M. G., Lazar, M. A., Voelter, W., & Verdin, E.
(2002). Enzymatic activity associated with class II HDACs is dependent on a multiprotein
complex containing HDAC3 and SMRT/N-CoR. *Molecular cell*, 9(1), 45–57.
https://doi.org/10.1016/s1097-2765(01)00429-4

Frescas, D., Guardavaccaro, D., Kuchay, S. M., Kato, H., Poleshko, A., Basrur, V., Elenitoba-
Johnson, K. S., Katz, R. A., & Pagano, M. (2008). KDM2A represses transcription of centromeric
satellite repeats and maintains the heterochromatic state. *Cell cycle (Georgetown, Tex.)*, 7(22),
3539–3547. https://doi.org/10.4161/cc.7.22.7062

Gegenschatz-Schmid, K., Verkauskas, G., Stadler, M. B., & Hadziselimovic, F. (2019). Genes
located in Y-chromosomal regions important for male fertility show altered transcript levels in
cryptorchidism and respond to curative hormone treatment. *Basic and clinical andrology*, 29, 8.
https://doi.org/10.1186/s12610-019-0089-3

Getun, I. V., Wu, Z., Fallahi, M., Ouizem, S., Liu, Q., Li, W., Costi, R., Roush, W. R., Cleveland, J.
L., & Bois, P. (2017). Functional Roles of Acetylated Histone Marks at Mouse Meiotic
Recombination Hot Spots. *Molecular and cellular biology*, 37(3), e00942-15.
https://doi.org/10.1128/MCB.00942-15

Giblin, W., Bringman-Rodenbarger, L., Guo, A. H., Kumar, S., Monovich, A. C., Mostafa, A. M.,
Skinner, M. E., Azar, M., Mady, A. S., Chung, C. H., Kadambi, N., Melong, K. A., Lee, H. J.,
Zhang, L., Sajjakulnukit, P., Trefely, S., Varner, E. L., Iyer, S., Wang, M., Wilmott, J. S., …
Lombard, D. B. (2021). The deacylase SIRT5 supports melanoma viability by influencing
chromatin dynamics. *The Journal of clinical investigation*, 131(12), e138926.
https://doi.org/10.1172/JCI138926

Glaser, S., Lubitz, S., Loveland, K. L., Ohbo, K., Robb, L., Schwenk, F., Seibler, J., Roellig, D.,
Kranz, A., Anastassiadis, K., & Stewart, A. F. (2009). The histone 3 lysine 4 methyltransferase,
Mll2, is only required briefly in development and spermatogenesis. *Epigenetics &*
chromatin, 2(1), 5. https://doi.org/10.1186/1756-8935-2-5

Godmann, M., Auger, V., Ferraroni-Aguiar, V., Di Sauro, A., Sette, C., Behr, R., & Kimmins, S.
(2007). Dynamic regulation of histone H3 methylation at lysine 4 in mammalian
spermatogenesis. *Biology of reproduction*, *77*(5), 754–764.
https://doi.org/10.1095/biolreprod.107.062265

González, B., Gancedo, S. N., Garazatua, S., Roldán, E., Vitullo, A. D., & González, C. R. (2020).
Dopamine Receptor D1 Contributes to Cocaine Epigenetic Reprogramming of Histone
Modifications in Male Germ Cells. Frontiers in cell and developmental biology, 8, 216.
https://doi.org/10.3389/fcell.2020.00216

Guo, L., Chao, S. B., Xiao, L., Wang, Z. B., Meng, T. G., Li, Y. Y., Han, Z. M., Ouyang, Y. C.,
Hou, Y., Sun, Q. Y., & Ou, X. H. (2017). Sperm-carried RNAs play critical roles in mouse
embryonic development. Oncotarget, 8(40), 67394–67405.
https://doi.org/10.18632/oncotarget.18672

Hammoud, S. S., Low, D. H., Yi, C., Carrell, D. T., Guccione, E., & Cairns, B. R. (2014).
Chromatin and transcription transitions of mammalian adult germline stem cells and
spermatogenesis. Cell stem cell, 15(2), 239–253. https://doi.org/10.1016/j.stem.2014.04.006

Hammoud, S. S., Nix, D. A., Zhang, H., Purwar, J., Carrell, D. T., & Cairns, B. R. (2009).
Distinctive chromatin in human sperm packages genes for embryo development. Nature,
460(7254), 473–478. https://doi.org/10.1038/nature08162

He, Z., Wu, J., Su, X., Zhang, Y., Pan, L., Wei, H., Fang, Q., Li, H., Wang, D. L., & Sun, F. L.
(2016). JMJD5 (Jumonji Domain-containing 5) Associates with Spindle Microtubules and Is
Required for Proper Mitosis. *The Journal of biological chemistry*, *291*(9), 4684–4697.
https://doi.org/10.1074/jbc.M115.672642

Herz, H. M., Mohan, M., Garruss, A. S., Liang, K., Takahashi, Y. H., Mickey, K., Voets, O.,
Verrijzer, C. P., & Shilatifard, A. (2012). Enhancer-associated H3K4 monomethylation by
Trithorax-related, the Drosophila homolog of mammalian Mll3/Mll4. *Genes &*
development, *26*(23), 2604–2620. https://doi.org/10.1101/gad.201327.112

Hirayama, J., Sahar, S., Grimaldi, B., Tamaru, T., Takamatsu, K., Nakahata, Y., & Sassone-Corsi,
P. (2007). CLOCK-mediated acetylation of BMAL1 controls circadian
function. *Nature*, *450*(7172), 1086–1090. https://doi.org/10.1038/nature06394

Hu, D., Gao, X., Cao, K., Morgan, M. A., Mas, G., Smith, E. R., Volk, A. G., Bartom, E. T.,
Crispino, J. D., Di Croce, L., & Shilatifard, A. (2017). Not All H3K4 Methylations Are Created
Equal: Mll2/COMPASS Dependency in Primordial Germ Cell Specification. *Molecular*
cell, *65*(3), 460–475.e6. https://doi.org/10.1016/j.molcel.2017.01.013

Husmann, D., & Gozani, O. (2019). Histone lysine methyltransferases in biology and disease.

Nature structural & molecular biology, 26(10), 880–889. https://doi.org/10.1038/s41594-019-

0298-7

Idler, R. K., & Yan, W. (2012). Control of messenger RNA fate by RNA-binding proteins: an

emphasis on mammalian spermatogenesis. Journal of andrology, 33(3), 309–337.

https://doi.org/10.2164/jandrol.111.014167

Iwamori, N., Iwamori, T., & Matzuk, M. M. (2013). H3K27 demethylase, JMJD3, regulates

fragmentation of spermatogonial cysts. *PloS one*, *8*(8), e72689.

https://doi.org/10.1371/journal.pone.0072689

Jan, S. Z., Vormer, T. L., Jongejan, A., Röling, M. D., Silber, S. J., de Rooij, D. G., Hamer, G.,

Repping, S., & van Pelt, A. (2017). Unraveling transcriptome dynamics in human

spermatogenesis. Development (Cambridge, England), 144(20), 3659–3673.

https://doi.org/10.1242/dev.152413

Jha, D. K., & Strahl, B. D. (2014). An RNA polymerase II-coupled function for histone H3K36

methylation in checkpoint activation and DSB repair. Nature communications, 5, 3965.

https://doi.org/10.1038/ncomms4965

Jiang, H., Gao, Q., Zheng, W., Yin, S., Wang, L., Zhong, L., Ali, A., Khan, T., Hao, Q., Fang, H.,

Sun, X., Xu, P., Pandita, T. K., Jiang, X., & Shi, Q. (2018). MOF influences meiotic expansion of

H2AX phosphorylation and spermatogenesis in mice. PLoS genetics, 14(5), e1007300.

https://doi.org/10.1371/journal.pgen.1007300

Jin, Q., Yu, L. R., Wang, L., Zhang, Z., Kasper, L. H., Lee, J. E., Wang, C., Brindle, P. K., Dent, S.

Y., & Ge, K. (2011). Distinct roles of GCN5/PCAF-mediated H3K9ac and CBP/p300-mediated

H3K18/27ac in nuclear receptor transactivation. *The EMBO journal*, *30*(2), 249–262.

https://doi.org/10.1038/emboj.2010.318

Jodar, M., Selvaraju, S., Sendler, E., Diamond, M. P., Krawetz, S. A., & Reproductive Medicine

Network (2013). The presence, role and clinical use of spermatozoal RNAs. Human reproduction

update, 19(6), 604–624. https://doi.org/10.1093/humupd/dmt031

Jones, B., Su, H., Bhat, A., Lei, H., Bajko, J., Hevi, S., Baltus, G. A., Kadam, S., Zhai, H., Valdez,

R., Gonzalo, S., Zhang, Y., Li, E., & Chen, T. (2008). The histone H3K79 methyltransferase

Dot1L is essential for mammalian development and heterochromatin structure. PLoS genetics,

4(9), e1000190. https://doi.org/10.1371/journal.pgen.1000190

Kaneshiro, K. R., Rechtsteiner, A., & Strome, S. (2019). Sperm-inherited H3K27me3 impacts
offspring transcription and development in C. elegans. Nature communications, 10(1), 1271.
https://doi.org/10.1038/s41467-019-09141-w

Kang, J. Y., Kim, J. Y., Kim, K. B., Park, J. W., Cho, H., Hahm, J. Y., Chae, Y. C., Kim, D., Kook,
H., Rhee, S., Ha, N. C., & Seo, S. B. (2018). KDM2B is a histone H3K79 demethylase and
induces transcriptional repression via sirtuin-1-mediated chromatin silencing. FASEB journal :
official publication of the Federation of American Societies for Experimental Biology, 32(10),
5737–5750. https://doi.org/10.1096/fj.201800242R

Khalil, A. M., Boyar, F. Z., & Driscoll, D. J. (2004). Dynamic histone modifications mark sex
chromosome inactivation and reactivation during mammalian spermatogenesis. Proceedings of
the National Academy of Sciences of the United States of America, 101(47), 16583–16587.
https://doi.org/10.1073/pnas.0406325101

Kleene K. C. (2001). A possible meiotic function of the peculiar patterns of gene expression in
mammalian   spermatogenic   cells.   Mechanisms   of   development,   106(1-2),   3–23.
https://doi.org/10.1016/s0925-4773(01)00413-0

Kouzarides T. (2007). Chromatin modifications and their function. *Cell*, *128*(4), 693–705.
https://doi.org/10.1016/j.cell.2007.02.005

Lambrot, R., Lafleur, C., & Kimmins, S. (2015). The histone demethylase KDM1A is essential for
the maintenance and differentiation of spermatogonial stem cells and progenitors. *FASEB journal*
: official publication of the Federation of American Societies for Experimental Biology, *29*(11),
4402–4416. https://doi.org/10.1096/fj.14-267328

Lee, K., Haugen, H. S., Clegg, C. H., & Braun, R. E. (1995). Premature translation of protamine 1
mRNA causes precocious nuclear condensation and arrests spermatid differentiation in mice.
Proceedings of the National Academy of Sciences of the United States of America, 92(26),
12451–12455. https://doi.org/10.1073/pnas.92.26.12451

Lepikhov, K., Wossidlo, M., Arand, J., & Walter, J. (2010). DNA methylation reprogramming and
DNA repair in the mouse zygote. The International journal of developmental biology, 54(11-12),
1565–1574. https://doi.org/10.1387/ijdb.103206kl

Lesch, B. J., Tothova, Z., Morgan, E. A., Liao, Z., Bronson, R. T., Ebert, B. L., & Page, D. C.
(2019). Intergenerational epigenetic inheritance of cancer susceptibility in mammals. eLife, 8,
e39380. https://doi.org/10.7554/eLife.39380

Liefke, R., Oswald, F., Alvarado, C., Ferres-Marco, D., Mittler, G., Rodriguez, P., Dominguez, M.,
& Borggrefe, T. (2010). Histone demethylase KDM5A is an integral part of the core Notch-RBP-
J repressor complex. *Genes & development*, *24*(6), 590–601. https://doi.org/10.1101/gad.563210

Lismer, A., Dumeaux, V., Lafleur, C., Lambrot, R., Brind'Amour, J., Lorincz, M. C., & Kimmins,
S. (2021). Histone H3 lysine 4 trimethylation in sperm is transmitted to the embryo and associated
with diet-induced phenotypes in the offspring. Developmental cell, 56(5), 671–686.e6.
https://doi.org/10.1016/j.devcel.2021.01.014

Lismer, A., Siklenka, K., Lafleur, C., Dumeaux, V., & Kimmins, S. (2020). Sperm histone H3
lysine 4 trimethylation is altered in a genetic mouse model of transgenerational epigenetic
inheritance. *Nucleic acids research*, *48*(20), 11380–11393. https://doi.org/10.1093/nar/gkaa712

Liu, Z., Oyola, M. G., Zhou, S., Chen, X., Liao, L., Tien, J. C., Mani, S. K., & Xu, J. (2015).
Knockout of the Histone Demethylase Kdm3b Decreases Spermatogenesis and Impairs Male
Sexual Behaviors. International journal of biological sciences, 11(12), 1447–1457.
https://doi.org/10.7150/ijbs.13795

Lu, L. Y., Wu, J., Ye, L., Gavrilina, G. B., Saunders, T. L., & Yu, X. (2010). RNF8-dependent
histone modifications regulate nucleosome removal during spermatogenesis. Developmental cell,
18(3), 371–384. https://doi.org/10.1016/j.devcel.2010.01.010

Luense, L. J., Donahue, G., Lin-Shiao, E., Rangel, R., Weller, A. H., Bartolomei, M. S., & Berger,
S. L. (2019). Gcn5-Mediated Histone Acetylation Governs Nucleosome Dynamics in
Spermiogenesis. *Developmental cell*, *51*(6), 745–758.e6.
https://doi.org/10.1016/j.devcel.2019.10.024

Luense, L. J., Wang, X., Schon, S. B., Weller, A. H., Lin Shiao, E., Bryant, J. M., Bartolomei, M.
S., Coutifaris, C., Garcia, B. A., & Berger, S. L. (2016). Comprehensive analysis of histone post-
translational modifications in mouse and human male germ cells. *Epigenetics & chromatin*, *9*, 24.
https://doi.org/10.1186/s13072-016-0072-6

Luo, Z., Qing, X., Benda, C., Huang, Z., Zhang, M., Huang, Y., Zhang, H., Wang, L., Lai, Y.,
Ward, C., Volpe, G., Zhong, X., Qin, B., Zhuang, Q., Esteban, M. A., & Li, W. (2019). Nuclear-
cytoplasmic shuttling of class IIa histone deacetylases regulates somatic cell reprogramming. *Cell*
regeneration (London, England), *8*(1), 21–29. https://doi.org/10.1016/j.cr.2018.11.001

Ma, P., & Schultz, R. M. (2008). Histone deacetylase 1 (HDAC1) regulates histone acetylation,
development, and gene expression in preimplantation mouse embryos. Developmental biology,
319(1), 110–120. https://doi.org/10.1016/j.ydbio.2008.04.011

Ma, X. S., Chao, S. B., Huang, X. J., Lin, F., Qin, L., Wang, X. G., Meng, T. G., Zhu, C. C.,
Schatten, H., Liu, H. L., & Sun, Q. Y. (2015). The Dynamics and Regulatory Mechanism of
Pronuclear H3k9me2 Asymmetry in Mouse Zygotes. Scientific reports, 5, 17924.
https://doi.org/10.1038/srep17924

Madsen, B., Tarsounas, M., Burchell, J. M., Hall, D., Poulsom, R., & Taylor-Papadimitriou, J.
(2003). PLU-1, a transcriptional repressor and putative testis-cancer antigen, has a specific
expression and localisation pattern during meiosis. Chromosoma, 112(3), 124–132.
https://doi.org/10.1007/s00412-003-0252-6

Maezawa, S., Hasegawa, K., Yukawa, M., Kubo, N., Sakashita, A., Alavattam, K. G., Sin, H. S.,
Kartashov, A. V., Sasaki, H., Barski, A., & Namekawa, S. H. (2018). Polycomb protein SCML2
facilitates H3K27me3 to establish bivalent domains in the male germline. Proceedings of the
National Academy of Sciences of the United States of America, 115(19), 4957–4962.
https://doi.org/10.1073/pnas.1804512115

Martinez-Gamero, C., Malla, S., & Aguilo, F. (2021). LSD1: Expanding Functions in Stem Cells
and Differentiation. *Cells*, *10*(11), 3252. https://doi.org/10.3390/cells10113252

McCarthy, A., Sarkar, K., Martin, E. T., Upadhyay, M., Jang, S., Williams, N. D., Forni, P. E.,
Buszczak, M., & Rangan, P. (2022). Msl3 promotes germline stem cell differentiation in female
Drosophila. *Development   (Cambridge,   England)*, *149*(1),   dev199625.
https://doi.org/10.1242/dev.199625

McLay, D. W., & Clarke, H. J. (2003). Remodelling the paternal chromatin at fertilization in
mammals.   Reproduction   (Cambridge,   England),   125(5),   625–633.
https://doi.org/10.1530/rep.0.1250625

Michishita, E., McCord, R. A., Berber, E., Kioi, M., Padilla-Nash, H., Damian, M., Cheung, P.,
Kusumoto, R., Kawahara, T. L., Barrett, J. C., Chang, H. Y., Bohr, V. A., Ried, T., Gozani, O., &
Chua, K. F. (2008). SIRT6 is a histone H3 lysine 9 deacetylase that modulates telomeric
chromatin. *Nature*, *452*(7186), 492–496. https://doi.org/10.1038/nature06736

Michishita, E., Park, J. Y., Burneskis, J. M., Barrett, J. C., & Horikawa, I. (2005). Evolutionarily
conserved and nonconserved cellular localizations and functions of human SIRT
proteins. *Molecular biology of the cell*, *16*(10), 4623–4635. https://doi.org/10.1091/mbc.e05-01-
0033

Mosammaparast, N., & Shi, Y. (2010). Reversal of histone methylation: biochemical and molecular
mechanisms of histone demethylases. *Annual review of biochemistry*, 79, 155–179.
https://doi.org/10.1146/annurev.biochem.78.070907.103946

Mulligan, P., Westbrook, T. F., Ottinger, M., Pavlova, N., Chang, B., Macia, E., Shi, Y. J., Barretina, J., Liu, J., Howley, P. M., Elledge, S. J., & Shi, Y. (2008). CDYL bridges REST and histone methyltransferases for gene repression and suppression of cellular transformation. Molecular cell, 32(5), 718–726. https://doi.org/10.1016/j.molcel.2008.10.025

Myrick, D. A., Christopher, M. A., Scott, A. M., Simon, A. K., Donlin-Asp, P. G., Kelly, W. G., & Katz, D. J. (2017). KDM1A/LSD1 regulates the differentiation and maintenance of spermatogonia in mice. *PloS one*, *12*(5), e0177473. https://doi.org/10.1371/journal.pone.0177473

Nebbioso, A., Tambaro, F. P., Dell'Aversana, C., & Altucci, L. (2018). Cancer epigenetics: Moving forward. *PLoS genetics*, *14*(6), e1007362. https://doi.org/10.1371/journal.pgen.1007362

Nishio, H., Hayashi, Y., Moritoki, Y., Kamisawa, H., Mizuno, K., Kojima, Y., & Kohri, K. (2014). Distinctive changes in histone H3K4 modification mediated via Kdm5a expression in spermatogonial stem cells of cryptorchid testes. *The Journal of urology*, *191*(5 Suppl), 1564–1572. https://doi.org/10.1016/j.juro.2013.10.071

Niu, C., Guo, J., Shen, X., Ma, S., Xia, M., Xia, J., & Zheng, Y. (2020). Meiotic gatekeeper STRA8 regulates cell cycle by interacting with SETD8 during spermatogenesis. *Journal of cellular and molecular medicine*, *24*(7), 4194–4211. https://doi.org/10.1111/jcmm.15080

Ntostis, P., Carter, D., Iles, D., Huntriss, J., Tzetis, M., & Miller, D. (2017). Potential sperm contributions to the murine zygote predicted by in silico analysis. Reproduction (Cambridge, England), 154(6), 777–788. https://doi.org/10.1530/REP-17-0097

Okada, Y., Scott, G., Ray, M. K., Mishina, Y., & Zhang, Y. (2007). Histone demethylase JHDM2A is critical for Tnp1 and Prm1 transcription and spermatogenesis. Nature, 450(7166), 119–123. https://doi.org/10.1038/nature06236

Parra M. (2015). Class IIa HDACs - new insights into their functions in physiology and pathology. *The FEBS journal*, *282*(9), 1736–1744. https://doi.org/10.1111/febs.13061

Parvanov, E. D., Petkov, P. M., & Paigen, K. (2010). Prdm9 controls activation of mammalian recombination hotspots. Science (New York, N.Y.), 327(5967), 835. https://doi.org/10.1126/science.1181495

Payne, C., & Braun, R. E. (2006). Histone lysine trimethylation exhibits a distinct perinuclear distribution in Plzf-expressing spermatogonia. Developmental biology, 293(2), 461–472. https://doi.org/10.1016/j.ydbio.2006.02.013

Pei, H., Zhang, L., Luo, K., Qin, Y., Chesi, M., Fei, F., Bergsagel, P. L., Wang, L., You, Z., & Lou, Z. (2011). MMSET regulates histone H4K20 methylation and 53BP1 accumulation at DNA damage sites. Nature, 470(7332), 124–128. https://doi.org/10.1038/nature09658

Percharde, M., & Azuara, V. (2013). Essential roles for the nuclear receptor coactivator Ncoa3 in
pluripotency. *Cell cycle (Georgetown, Tex.)*, *12*(2), 195–196. https://doi.org/10.4161/cc.23377

Percharde, M., Lavial, F., Ng, J. H., Kumar, V., Tomaz, R. A., Martin, N., Yeo, J. C., Gil, J.,
Prabhakar, S., Ng, H. H., Parker, M. G., & Azuara, V. (2012). Ncoa3 functions as an essential
Esrrb coactivator to sustain embryonic stem cell self-renewal and reprogramming. *Genes &*
development, *26*(20), 2286–2298. https://doi.org/10.1101/gad.195545.112

Peters, A. H., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer, C., Weipoltshammer,
K., Pagani, M., Lachner, M., Kohlmaier, A., Opravil, S., Doyle, M., Sibilia, M., & Jenuwein, T.
(2001). Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin and
genome stability. *Cell*, *107*(3), 323–337. https://doi.org/10.1016/s0092-8674(01)00542-6

Prachayasittikul, V., Prathipati, P., Pratiwi, R., Phanus-Umporn, C., Malik, A. A., Schaduangrat, N.,
Seenprachawong, K., Wongchitrat, P., Supokawej, A., Prachayasittikul, V., Wikberg, J. E., &
Nantasenamat, C. (2017). Exploring the epigenetic drug discovery landscape. *Expert opinion on*
drug discovery, *12*(4), 345–362. https://doi.org/10.1080/17460441.2017.1295954

Pradeepa, M. M., Nikhil, G., Hari Kishore, A., Bharath, G. N., Kundu, T. K., & Rao, M. R. (2009).
Acetylation of transition protein 2 (TP2) by KAT3B (p300) alters its DNA condensation property
and interaction with putative histone chaperone NPM3. *The Journal of biological*
chemistry, *284*(43), 29956–29967. https://doi.org/10.1074/jbc.M109.052043

Rajender, S., Avery, K., & Agarwal, A. (2011). Epigenetics, spermatogenesis and male infertility.
Mutation research, 727(3), 62–71. https://doi.org/10.1016/j.mrrev.2011.04.002

Rickels, R., Hu, D., Collings, C. K., Woodfin, A. R., Piunti, A., Mohan, M., Herz, H. M., Kvon, E.,
& Shilatifard, A. (2016). An Evolutionary Conserved Epigenetic Mark of Polycomb Response
Elements Implemented by Trx/MLL/COMPASS. *Molecular cell*, *63*(2), 318–328.
https://doi.org/10.1016/j.molcel.2016.06.018

Scher, M. B., Vaquero, A., & Reinberg, D. (2007). SirT3 is a nuclear NAD+-dependent histone
deacetylase that translocates to the mitochondria upon cellular stress. *Genes &*
development, *21*(8), 920–928. https://doi.org/10.1101/gad.1527307

Schulz, K. N., & Harrison, M. M. (2019). Mechanisms regulating zygotic genome activation.
Nature reviews. Genetics, 20(4), 221–234. https://doi.org/10.1038/s41576-018-0087-x

Seong, K. H., Ly, N. H., Katou, Y., Yokota, N., Nakato, R., Murakami, S., Hirayama, A., Fukuda,
S., Kang, S., Soga, T., Shirahige, K., & Ishii, S. (2020). Paternal restraint stress affects offspring
metabolism via ATF-2 dependent mechanisms in Drosophila melanogaster germ cells.
Communications biology, 3(1), 208. https://doi.org/10.1038/s42003-020-0935-z

Shirane, K., Miura, F., Ito, T., & Lorincz, M. C. (2020). NSD1-deposited H3K36me2 directs de
novo methylation in the mouse male germline and counteracts Polycomb-associated
silencing. *Nature genetics*, *52*(10), 1088–1098. https://doi.org/10.1038/s41588-020-0689-z

Siklenka, K., Erkek, S., Godmann, M., Lambrot, R., McGraw, S., Lafleur, C., Cohen, T., Xia, J.,
Suderman, M., Hallett, M., Trasler, J., Peters, A. H., & Kimmins, S. (2015). Disruption of histone
methylation in developing sperm impairs offspring health transgenerationally. Science (New
York, N.Y.), 350(6261), aab2006. https://doi.org/10.1126/science.aab2006

Skvortsova, K., Iovino, N., & Bogdanović, O. (2018). Functions and mechanisms of epigenetic
inheritance in animals. *Nature reviews. Molecular cell biology*, *19*(12), 774–790.
https://doi.org/10.1038/s41580-018-0074-2

Smith, K., & Spadafora, C. (2005). Sperm-mediated gene transfer: applications and implications.
BioEssays : news and reviews in molecular, cellular and developmental biology, 27(5), 551–562.
https://doi.org/10.1002/bies.20211

Sone, Y., Ito, M., Shirakawa, H., Shikano, T., Takeuchi, H., Kinoshita, K., & Miyazaki, S. (2005).
Nuclear translocation of phospholipase C-zeta, an egg-activating factor, during early embryonic
development. Biochemical and biophysical research communications, 330(3), 690–694.
https://doi.org/10.1016/j.bbrc.2005.03.032

Song, N., Liu, J., An, S., Nishino, T., Hishikawa, Y., & Koji, T. (2011). Immunohistochemical
Analysis of Histone H3 Modifications in Germ Cells during Mouse Spermatogenesis. *Acta
histochemica et cytochemica*, *44*(4), 183–190. https://doi.org/10.1267/ahc.11027

Spencer, T. E., Jenster, G., Burcin, M. M., Allis, C. D., Zhou, J., Mizzen, C. A., McKenna, N. J.,
Onate, S. A., Tsai, S. Y., Tsai, M. J., & O'Malley, B. W. (1997). Steroid receptor coactivator-1 is
a histone acetyltransferase. Nature, 389(6647), 194–198. https://doi.org/10.1038/38304

Streubel, G., Watson, A., Jammula, S. G., Scelfo, A., Fitzpatrick, D. J., Oliviero, G., McCole, R.,
Conway, E., Glancy, E., Negri, G. L., Dillon, E., Wynne, K., Pasini, D., Krogan, N. J., Bracken,
A. P., & Cagney, G. (2018). The H3K36me2 Methyltransferase Nsd1 Demarcates PRC2-
Mediated H3K27me2 and H3K27me3 Domains in Embryonic Stem Cells. *Molecular cell*, *70*(2),
371–379.e5. https://doi.org/10.1016/j.molcel.2018.02.027

Su, Z., Wang, F., Lee, J. H., Stephens, K. E., Papazyan, R., Voronina, E., Krautkramer, K. A.,
Raman, A., Thorpe, J. J., Boersma, M. D., Kuznetsov, V. I., Miller, M. D., Taverna, S. D.,
Phillips, G. N., Jr, & Denu, J. M. (2016). Reader domain specificity and lysine demethylase-4
family function. *Nature communications*, *7*, 13387. https://doi.org/10.1038/ncomms13387

Sun, Z., Zhang, Y., Jia, J., Fang, Y., Tang, Y., Wu, H., & Fang, D. (2020). H3K36me3, message from chromatin to DNA damage repair. Cell & bioscience, 10, 9. https://doi.org/10.1186/s13578-020-0374-z

Tachibana, M., Nozaki, M., Takeda, N., & Shinkai, Y. (2007). Functional dynamics of H3K9 methylation during meiotic prophase progression. *The EMBO journal*, *26*(14), 3346–3359. https://doi.org/10.1038/sj.emboj.7601767

Trigg, N. A., Eamens, A. L., & Nixon, B. (2019). The contribution of epididymosomes to the sperm small RNA profile. Reproduction (Cambridge, England), 157(6), R209–R223. https://doi.org/10.1530/REP-18-0480

van der Heijden, G. W., Ramos, L., Baart, E. B., van den Berg, I. M., Derijck, A. A., van der Vlag, J., Martini, E., & de Boer, P. (2008). Sperm-derived histones contribute to zygotic chromatin in humans. *BMC developmental biology*, *8*, 34. https://doi.org/10.1186/1471-213X-8-34

Wan, M., Liang, J., Xiong, Y., Shi, F., Zhang, Y., Lu, W., He, Q., Yang, D., Chen, R., Liu, D., Barton, M., & Songyang, Z. (2013). The trithorax group protein Ash2l is essential for pluripotency and maintaining open chromatin in embryonic stem cells. The Journal of biological chemistry, 288(7), 5039–5048. https://doi.org/10.1074/jbc.M112.424515

Wang, Y., Iwamori, T., Kaneko, T., Iida, H., & Iwamori, N. (2021). Comparative distributions of RSBN1 and methylated histone H4 Lysine 20 in the mouse spermatogenesis. *PloS one*, *16*(6), e0253897. https://doi.org/10.1371/journal.pone.0253897

Wapenaar, H., & Dekker, F. J. (2016). Histone acetyltransferases: challenges in targeting bi-substrate enzymes. *Clinical epigenetics*, 8, 59. https://doi.org/10.1186/s13148-016-0225-2

Warren, J. S., Tracy, C. M., Miller, M. R., Makaju, A., Szulik, M. W., Oka, S. I., Yuzyuk, T. N., Cox, J. E., Kumar, A., Lozier, B. K., Wang, L., Llana, J. G., Sabry, A. D., Cawley, K. M., Barton, D. W., Han, Y. H., Boudina, S., Fiehn, O., Tucker, H. O., Zaitsev, A. V., … Franklin, S. (2018). Histone methyltransferase Smyd1 regulates mitochondrial energetics in the heart. Proceedings of the National Academy of Sciences of the United States of America, 115(33), E7871–E7880. https://doi.org/10.1073/pnas.1800680115

Wickham et al., (2019). Welcome to the Tidyverse. Journal of Open Source Software, 4(43), 1686, https://doi.org/10.21105/joss.01686

Wu, D., Li, Y., Zhu, K. S., Wang, H., & Zhu, W. G. (2018). Advances in Cellular Characterization of the Sirtuin Isoform, SIRT7. Frontiers in endocrinology, 9, 652. https://doi.org/10.3389/fendo.2018.00652

Wu, J., Huang, B., Chen, H., Yin, Q., Liu, Y., Xiang, Y., Zhang, B., Liu, B., Wang, Q., Xia, W., Li, W., Li, Y., Ma, J., Peng, X., Zheng, H., Ming, J., Zhang, W., Zhang, J., Tian, G., Xu, F., … Xie, W. (2016). The landscape of accessible chromatin in mammalian preimplantation embryos. Nature, 534(7609), 652–657. https://doi.org/10.1038/nature18606

Yang, X., Li, L., Liang, J., Shi, L., Yang, J., Yi, X., Zhang, D., Han, X., Yu, N., & Shang, Y. (2013). Histone acetyltransferase 1 promotes homologous recombination in DNA repair by facilitating histone turnover. The Journal of biological chemistry, 288(25), 18271–18282. https://doi.org/10.1074/jbc.M113.473199

Yang, X., Yu, W., Shi, L., Sun, L., Liang, J., Yi, X., Li, Q., Zhang, Y., Yang, F., Han, X., Zhang, D., Yang, J., Yao, Z., & Shang, Y. (2011). HAT4, a Golgi apparatus-anchored B-type histone acetyltransferase, acetylates free histone H4 and facilitates chromatin assembly. *Molecular cell*, *44*(1), 39–50. https://doi.org/10.1016/j.molcel.2011.07.032

Yao, C. J., Xu, W. J., Gong, X. L., Zhou, Y., Yan, Z. Q., Zhu, Z. J., Wang, Z. X., Li, Q. L., Guo, X. B., Wang, L. Y., Ma, D., & Qiao, Z. D. (2010). The role of Dby mRNA in early development of male mouse zygotes. Asian journal of andrology, 12(4), 567–577. https://doi.org/10.1038/aja.2010.28

Ye, J., Ai, X., Eugeni, E. E., Zhang, L., Carpenter, L. R., Jelinek, M. A., Freitas, M. A., & Parthun, M. R. (2005). Histone H4 lysine 91 acetylation a core domain modification associated with chromatin assembly. Molecular cell, 18(1), 123–130. https://doi.org/10.1016/j.molcel.2005.02.031

Yin, H., Kang, Z., Zhang, Y., Gong, Y., Liu, M., Xue, Y., He, W., Wang, Y., Zhang, S., Xu, Q., Fu, K., Zheng, B., Xie, J., Zhang, J., Wang, Y., Lin, M., Zhang, Y., Feng, H., Xin, C., Guan, Y., … Ye, L. (2021). HDAC3 controls male fertility through enzyme-independent transcriptional regulation at the meiotic exit of spermatogenesis. *Nucleic acids research*, 49(9), 5106–5123. https://doi.org/10.1093/nar/gkab313

Zamudio, N. M., Scott, H. S., Wolski, K., Lo, C. Y., Law, C., Leong, D., Kinkel, S. A., Chong, S., Jolley, D., Smyth, G. K., de Kretser, D., Whitelaw, E., & O'Bryan, M. K. (2011). DNMT3L is a regulator of X chromosome compaction and post-meiotic gene transcription. PloS one, 6(3), e18276. https://doi.org/10.1371/journal.pone.0018276

Zhang, J., Bonasio, R., Strino, F., Kluger, Y., Holloway, J. K., Modzelewski, A. J., Cohen, P. E., & Reinberg, D. (2013). SFMBT1 functions with LSD1 to regulate expression of canonical histone genes and chromatin-related factors. *Genes & development*, *27*(7), 749–766. https://doi.org/10.1101/gad.210963.112

1071   Zuo, X., Rong, B., Li, L., Lv, R., Lan, F., & Tong, M. H. (2018). The histone methyltransferase
1072   SETD2 is required for expression of acrosin-binding protein 1 and protamines and essential for
1073   spermiogenesis in mice. *The Journal of biological chemistry*, *293*(24), 9188–9197.
1074   https://doi.org/10.1074/jbc.RA118.002851

1075