A Role for Steroid 5 alpha-reductase 1 in Vascular Remodelling During Endometrial Decidualisation

Shaw, I.W[1]., Kirkwood, P.M.[1], Rebourcet, D.[2], Cousins, F.L. [1], Ainslie, R.J. [1], Livingstone, D.E.W.[3], Smith L.B.[2], Saunders P.T.K. [1], Gibson, D.A[1]*.

1. *Centre for Inflammation Research, The University of Edinburgh*
2. *MRC Centre for Reproductive Health, The University of Edinburgh*
3. *BHF Centre for Cardiovascular Sciences, The University of Edinburgh*

Key words: Decidualisation, angiogenesis, vascular remodelling, 5 alpha-reductase, Srd5a1, dihydrotestosterone, androgen, intracrinology, finasteride.

Short title: DHT-dependent decidualisation

*to whom correspondence should be addressed.

D.A.Gibson@ed.ac.uk

Centre for Inflammation Research, The University of Edinburgh, Institute for Regeneration and Repair, Edinburgh Bioquarter, Edinburgh

16   **Abstract**

Decidualisation is the hormone-dependent process of endometrial remodelling that is

18   essential for fertility and reproductive health. It is characterised by dynamic changes in the

endometrial stromal compartment including differentiation of fibroblasts, immune cell

20   trafficking and vascular remodelling. Deficits in decidualisation are implicated in disorders of

pregnancy such as implantation failure, intra-uterine growth restriction, and pre-eclampsia.

22   Androgens are key regulators of decidualisation that promote optimal differentiation of

stromal fibroblasts and activation of downstream signalling pathways required for

24   endometrial remodelling. We have shown that androgen biosynthesis, via 5α-reductase-

dependent production of dihydrotestosterone, is required for optimal decidualisation of

26   human stromal fibroblasts *in vitro*, but whether this is required for decidualisation *in vivo* has

not been tested.

28   In the current study we used steroid 5α-reductase type 1 (SRD5A1) deficient mice (*Srd5a1-/-*

mice) and a validated model of induced decidualisation to investigate the role of SRD5A1

30   and intracrine androgen signalling in endometrial decidualisation. We measured

decidualisation response (weight/proportion), transcriptomic changes, and morphological

32   and functional parameters of vascular development. These investigations revealed a striking

effect of 5α-reductase deficiency on the decidualisation response. Furthermore, vessel

34   permeability and transcriptional regulation of angiogenesis signalling pathways, particularly

those that involved vascular endothelial growth factor (VEGF), were disrupted in the

36   absence of 5α-reductase. In *Srd5a1-/-* mice, injection of dihydrotestosterone co-incident with

decidualisation restored decidualisation responses, vessel permeability, and expression of

38   angiogenesis genes to wild type levels.

Androgen availability declines with age which may contribute to age-related risk of

40   pregnancy disorders. These findings show that intracrine androgen signalling is required for

optimal decidualisation in vivo and confirm a major role for androgens in the development of

42   the vasculature during decidualisation through regulation of the VEGF pathway. These

findings highlight new opportunities for improving age-related deficits in fertility and

44   pregnancy health by targeting androgen-dependent signalling in the endometrium.

46   **Introduction**

Decidualisation is a fundamental step in the establishment of pregnancy that involves
48   coordinated remodelling of the endometrial stroma. It is a hormone-dependent process
characterised by differentiation of endometrial stromal fibroblasts (hESF) (Dunn et al., 2003),
50   immune cell trafficking (Yang et al., 2019) and vascular remodelling (Plaisier, 2011). Deficits
in decidualisation are implicated in disorders of pregnancy such as implantation failure, intra-
52   uterine growth restriction, and pre-eclampsia (Dunk et al., 2008, Gibson et al., 2016b). Risk
of pregnancy disorders increases with age which may be associated with age-related decline
54   in hormone production or availability.

Androgens are key regulators of decidualisation that promote optimal differentiation of
56   stromal fibroblasts and activation of downstream signalling pathways required for
endometrial remodelling. Androgen supplementation enhances secretion of decidualization
58   markers in hESF *in vitro* (Narukawa et al., 1994). Exogenous dihydrotestosterone (DHT)
enhances and maintains decidualisation responses in mice, an effect which is attenuated by
60   co-administration of the AR antagonist flutamide (Zhang and Croy, 1996). Endometrial
fibroblasts in both mouse and human endometrium express androgen receptor (AR) and
62   their function is altered by AR-dependent signalling (Marshall et al., 2011, Simitsidellis et al.,
2019, Simitsidellis et al., 2016). Using assays that combined *in vitro* decidualisation and
64   knockdown of receptors Cloke et al. reported that AR regulated distinct decidual gene
networks involved in cytoskeletal organization, cell motility and regulation of the cell cycle
66   (Cloke et al., 2008). Using a pharmacologic approach, we found that flutamide attenuated
the expression of decidualisation and endometrial receptivity markers in hESF (Gibson et al.,
68   2016a). Together, these studies confirm the importance of androgens and AR-dependent
signalling for optimal decidualisation responses.

70   In women, androgens and their precursors are abundant in the circulation but production
declines with age (Zumoff et al., 1995, Labrie et al., 1997). Circulating concentrations of the
72   most potent androgen dihydrotestosterone (DHT) are low but this is because DHT is

primarily a product of local metabolism within target tissues in women; mediating its affects

74   via intracrine signalling (Rothman et al., 2011, Labrie et al., 2017). We have shown that DHT

is actively produced by endometrial stromal fibroblasts during decidualisation via expression

76   of the enzyme SRD5A1 and that the abundance of DHT is affected by availability of steroid

precursors such as dehydroepiandrosterone (DHEA) (Gibson et al., 2018a, Gibson et al.,

78   2016a, Gibson et al., 2018b, Gibson et al., 2016b). When we increased intracrine DHT

signalling by supplementing with DHEA we found that decidualisation responses were

80   enhanced and expression of implantation markers were also increased. Androgen precursor

availability and intracrine androgen signalling can therefore dictate the extent of

82   decidualisation responses.

To date, a role for intracrine androgen signalling in regulating decidualisation *in vivo* has not

84   been rigorously investigated. Previous studies have been hampered by the developmental

uterine defects reported in global AR knock-out female mice (Walters et al., 2016). In the

86   current study, we have used mice which are homozygous for a recessive knock-out mutation

in the gene encoding the type 1 steroid 5α-reductase (SRD5A1) enzyme (*Srd5a1[-/-]*) which

88   lack the capacity to convert T to DHT (Mahendroo et al., 1996). Previously, Srd5a1[-/-] mice

have been shown to have reduced fertility due to oestrogen-induced foetal death and a

90   parturition defect caused by lack of 5α-androstan-3α,17β-diol, another product of SRD5A1

(Mahendroo et al., 1997). The potential role of this enzyme and the contribution of 5α-

92   reduced androgens to decidualisation has not been tested, but it is plausible that the

reduced fecundity reported in Srd5a1[-/-] mice (Mahendroo and Russell, 1999) may in part be

94   due to decidualisation defects as a result of intracrine androgen deficiency. To investigate

this, we have assessed the impact of 5α-reductase deletion or inhibition in a mouse model of

96   induced decidualisation.

We show that 5α-reductase deficiency leads to impaired decidualisation, structural and

98   functional changes to decidual blood vessels, and transcriptomic changes affecting

angiogenesis signalling pathways. Deficits in decidualisation in Srd5a1[-/-] mice were reversed

100   by restoration of androgen signalling through exogenous DHT administration. We conclude

that intracrine androgens are necessary for optimal decidualisation and vascular remodelling

102   required for establishment of pregnancy.

104 **Materials and Methods**

Mice

106 All mouse work was performed in accordance with the Animals (Scientific Procedures) Act

1986 under UK law and were bred on a C57Bl/6J or C57Bl6/JCrl background. The Srd5a1$^{-/-}$

108 mouse line was obtained from the Jackson Laboratories (https://www.jax.org/strain/002793).

The original paper describing the generation of mice with a null allele was published by

110 Mahendroo, Cala, and Russel (Mahendroo et al., 1996). Due to parturition defects in

homozygous mice heterozygote crosses were used to maintain the colony and produce

112 *Srd5a1$^{-/-}$* progeny. Wildtype (WT) mice were either littermates or purchased from Charles

River Laboratories (Tranent, Scotland).

114 *Mouse model of induced decidualisation*

Mice were ovariectomised to remove endogenous ovarian steroids at day 0 (d0), then

116 administered oestradiol (E2, Sigma-Aldrich, Poole) by subcutaneous injection daily on d7-9

(5ug.kg^{-1} in 200ul sesame oil) and on d13-15 (0.25ug.kg^{-1} in 200ul sesame oil) with a P4

118 pellet inserted on d13. On day 15 a decidualisation stimulus was administered by

intrauterine injection of 20ul sesame oil into the uterus via transvaginal delivery using a non-

120 surgical embryo transfer device (NSET). Animals were sacrificed 4 days later and uterine

tissues collected (Fig. 1A, (Cousins et al., 2014)). The decidualisation response was

122 immediately scored in each horn as outlined in Fig. 1B. Specifically a non-decidualised horn

had no decidualisation reaction whatsoever (Non), whereas a fully decidualised horn had

124 continuous decidualisation along the majority (>50%) of its length (Full). Partial

decidualisation is any level of decidualisation between these two extremes (Fig. 1B).

126 Collectively, samples designated Full or Partial are referred to as decidualised (Dec).

Finasteride (Sigma-Aldrich, Poole) was dissolved in 100% EtOH, then diluted to 5% v/v in

128 sesame oil to a final concentration of 12.5mg.mL^{-1} and 100uL (50 mg.kg^{-1}) was administered

via intraperitoneal injection daily from the point of the decidualisation stimulus onwards, i.e.

130   d15-18, as show in Figure S1A. Finasteride is a synthetic 4-azasteroid that competitively inhibits both type I and II 5α-reductase enzymes in rodents (Livingstone et al., 2015, Thigpen

132   and Russell, 1992).

DHT (Sigma-Aldrich, Poole) was dissolved in 100% ethanol (EtOH) then diluted to 5% v/v in

134   0.4% w/v methylcellulose solution for a final DHT concentration of 2mg.mL^{-1}. 100uL (8mg.kg^{-1}) was injected subcutaneously on d15 at the point of decidualisation stimulus. At the end of

136   experiments mice were terminated by exposure to rising concentrations of CO_2 combined with cervical dislocation.

138   ***Histology and immunolabelling***

Uterine tissue for histology was fixed in 4% paraformaldehyde then dehydrated and

embedded in paraffin blocks and sectioned according to standard protocols. Protocols for 3,3'-diaminobenzidine (DAB) and immunofluorescence staining were performed as

previously reported (Kirkwood et al., 2021, Simitsidellis et al., 2016). Briefly, slides were dewaxed, rehydrated, and subjected to antigen retrieval in pH 6.0 citrate buffer using a

microwave. Following a wash in phosphate buffered saline (PBS) slides were incubated with 3% H_2O_2 solution in PBS for 15min to remove endogenous peroxidase activity. Following

several PBS washes slides were incubated with normal goat serum solution (NGS, PBS containing 20% goat serum and 0.05% (w/v) bovine serum albumin) to block non-specific

interactions. Anti-CD31 primary antibody (Abcam, ab28364, 1:500 dilution) in NGS was added and incubated overnight at 4^{o}C. Following several washes, slides were incubated with

goat-raised peroxidase-conjugated secondary antibody (Abcam, ab7171, 1:500 dilution) for 1hr at room temperature. Following several washes, slides were incubated with either DAB

reagent or Opal red 570 reagent for 10 minutes or until sufficiently developed. After a brief wash, slides were counterstained either with haematoxylin and eosin as per standard

procedures, or with DAPI before being mounted with coverslips.

Slides were imaged using a Zeiss Z1 microscope, or a Zeiss AxioScan Slide Scanner using

156   conventional fluorescent setups.

Vessel quantification

158   To quantify vessels, regions of non-decidualised endometrium (Non) or decidua (Dec samples) were selected digitally on 20X magnification slide scans of CD31-labelled uterus

160   sections and overlaid with a grid with edge length 250 pixels (~ 28μm). The intersections on the grid were sequentially classified as either vessel or not vessel, and the total proportion of

162   vessel to non-vessel intersections was used as a measure of vessel count. Additionally, each time a vessel was identified, it was given an irregularity score equalling the number of

164   intersections on a tree used to trace out the vessel's cross-sectional shape. For example, a circular or or 'L' shaped cross-section would have a score of 0, a 'T' shape a score of 1, an

166   'H' shape a score of 2, etc. Irregularity scores were averaged across all observations.

RNA extraction

168   Uterine samples for RNA analysis were collected in RNALater solution (Qiagen, Hilden, Germany) and stored at $-80^{o}C$ until tissue extraction. For decidualised samples, only regions

170   of uterus containing decidualised endometrium were selected. Samples were thawed and homogenised in 1 mL Trizol reagent using a TissueLyser machine (6min, 25Hz).

172   Homogenate was transferred to MaXtract High Density tubes along with 200ul chloroform and 100ul RNase free water, shaken vigorously and centrifuged. The aqueous layer was

174   added to RNeasy spin columns (Qiagen, Hilden, Germany) and processed as per manufacturer's instructions.

176   ### Quantitative RT-PCR

cDNA was synthesised from $100ng.uL^{-1}$ RNA using Superscript VILO cDNA synthesis kit

178   (Thermo Fisher Scientific Life Sciences, Schwerte, Germany) as per the manufacturer's instructions. Thermal cycler settings were $25^{o}C$ for 10min, $42^{o}C$ for 60min, $85^{o}C$ for 5min.

180   Primers were designed using Universal Probe Library Assay Design Center (Roche Applied Science, Burgess Hill, UK) or, when this was discontinued, using Neoformit

182   (https://primers.neoformit.com/ Accessed: 6/4/22; sequences in Supplementary Data 1).

Primers were synthesised by Eurofins MWG Operon (Ebersberg, Gemany). Reactions were

184   run in duplicate on a Quantstudio 5 384-well PCR machine (Thermo Fisher Scientific Life

Sciences, Schwerte, Germany) using the following settings: 95^{o}C for 10min then 40 cycles of

186   95^{o}C for 15s and 60^{o}C for 1min. A serial dilution of a standard cDNA solution (consisting of a

mix of all samples tested) was used to plot a standard curve and expression levels

188   interpolated from this. Data was processed using QuantStudio Design and Analysis Software.

Nanostring gene expression analysis and data processing

190   Samples were analysed on the NanoString nCounter Analysis System using a NanoString

Mouse Pancancer Pathways Panel kit (Nanostring, Edinburgh, UK). RNA at 20ng.uL^{-1} was

192   used. Processing was performed by the Host and Tumour Profiling Unit Microarray Services,

Institute of Genetics and Cancer, University of Edinburgh, as per the manufacturer's

194   instructions. Briefly, the reporter codeset was mixed with hybridisation buffer and added to

each of the RNA samples, followed by the capture codeset. This mix was hybridised at 65^{o}C

196   for 18hr. Following hybridisation samples were loaded into the provided cartridges on the

nCounter prep station and processed using the High Sensitivity protocol. Cartridges were

198   then sealed and read using the digital analyser on the max setting. No QC flags were

registered for any of the samples.

200   Data was processed using the nSolver analysis software. Data was normalised using a

combination of positive controls and housekeeping genes. Firstly, background was

202   subtracted using a background value of the mean plus two standard deviations of the

negative control values. Normalisation using positive controls and housekeeper genes was

204   performed using the geometric mean to compute the normalisation factor, and the

manufacturer-provided list of housekeeper genes was used excluding the following genes

206   based on the results of the geNorm algorithm in the Nanostring Advanced Analysis modules:

Hprt, *Alas1*, *G6pdx*, *Gusb*, *Ppia*.

208   Normalised counts data was then analysed for differential gene expression using R,

specifically the edgeR analysis workflow. Dispersions were estimated with

210   the *estimateDisp()* command and fit to the design model using *glmQLFit()*, before testing for

differential expression using *glmQLFTest()*. Graphs were produced using the ggplot2

212   package. Finally, to perform gene ontogeny (GO) analysis we utilised the clusterProfiler

analysis workflow. Ensembl IDs were mapped from the Bioconductor org.Mm.eg.db genome

214   annotation and enrichment assessed using the *enrichGO()* command with the 'universe' set

as the genes included in the Nanostring Pancancer Pathways panel. Graphs were produced

216   using the *dotplot ()* and *cnetplot ()* commands from the enrichplot R package.

Comparison of Nanostring gene expression with scRNAseq gene expression

218   Normalised counts and differential gene expression data was further analysed in the context

of publicly available single cell RNA sequencing (scRNAseq) data, accessed from NCBI's

220   Gene Expression Omnibus (Edgar et al., 2002) through GEO Series accession numbers

GSE160772 and GSE198556. These scRNAseq data defined the transcriptomic profile of

222   over 20,000 perivascular, fibroblast and epithelial cell types present in the mouse uterus.

The AverageExpression() command (Seurat) was used to calculate the mean expression of

224   Nanostring DEgenes in perivascular, fibroblast and epithelial cell clusters from scRNAseq

data. Gene expression matrices were normalised and clustered heatmaps produced using

226   the pheatmap package. Expression values shown are scaled but not centred. This

comparison allowed us to attribute the expression of genes that were found to be

228   differentially expressed between Srd5a1$^{-/-}$ and WT genotypes to certain cell fractions in the

mouse endometrium.

230   ***Creation and imaging of resin casts***

Resin casts were prepared using a method previously validated in male mice (Rebourcet et

232   al., 2016). Briefly, female mice (Non and Dec) were culled using a terminal dose of sodium

pentobarbital (150 mg/kg, ip) and perfusion fixation of the vasculature was achieved via the

234   left ventricle. Heparinized PBS (heparin, 20 U/mL) was infused at 6 mL/min for 2 minutes.

Low-viscosity resin (10 mL; Microfil MV-122; Flow Tech Inc) was prepared according to the

236   manufacturer's instructions and then infused via the left ventricle. Uterine tissues were

recovered, trimmed, fixed in paraformaldehyde and embedded in 1.5% low melting point

238 agarose (Invitrogen, UK), dehydrated in methanol (100%; 24 hours) and optically cleared in

benzyl alcohol:benzyl benzoate (1:2 v/v; 24 hours). Imaging was carried out as described in

240 Rebourcet et al. with a positive control tissue being a previously prepared mouse testis

(Rebourcet et al., 2016).

242 ***Evans Blue assay of vascular permeability***

Evans Blue (EB) dye binds to albumin in the blood so its presence in tissues, which can be

244 quantified spectrophotometrically, implies a breakdown in the endothelial barrier. The Evans

Blue dye (Sigma-Aldrich, Poole) was dissolved in sterile saline solution to a concentration of

246 0.5% w/v, and 200uL was injected i.v. (via tail vein) 30-90 minutes prior to cull via cervical

dislocation. The interval between EB injection and cull was recorded and found not to

248 associate with dye retention in tissues.

To quantify the amount of Evans blue in the tissues, following sacrifice a portion of the

250 uterus was taken, precisely weighed, and transferred to 500µl of formamide. The gross

appearance of the tissue was recorded using a digital camera. For decidualised samples,

252 only sections of uterus containing decidualised endometrium were selected. Samples were

incubated for 24h at $55^{o}C$ to extract the dye, then tissue pieces were discarded, the solution

254 centrifuged to remove debris, and absorbance read in triplicate at 610nm.

Data processing and statistical analysis

256 Unless otherwise stated, data was processed using Microsoft Excel and Python. Statistics

were performed using either Python or Graphpad Prism 5.

258 In the analysis of Nanostring data, two-way ANOVAs were performed on all genes found to

be expressed and ANOVA results corrected for multiple comparisons using the Benjamini

260 and Hochberg method (Benjamini and Hochberg, 1995). The false discovery rate (FDR) was

0.05.

262   In the analysis of Evans Blue absorbance data, absorbance readings are shown as a ratio

relative to the mean value of WT or vehicle-treated WT data. In the case of data from Figure

264   2, ratios have been calculated separately for partially and fully decidualised horns and the

results combined for the subsequent statistical analysis.

266   Statistical tests performed for other comparisons are stated in figure legends and the text.

268 **Results**

5α-Reductase deficiency impairs decidualisation in mouse uterus

270 To investigate the functional requirement for 5α-reductase during decidualisation we induced

decidualisation in WT and Srd5a1[-/-] mice using a previously validated protocol (Fig. 1A;

272 (Cousins et al., 2014)). We used a ternary system to classify the extent of decidualisation.

Uterine horns were classified as non-, partially- or fully-decidualised (Non, Partial, Full,

274 respectively; Fig. 1B). For the purposes of analysis individual uterine horns were treated as

separate biological entities. In instances where only the decidualised regions of the uterus

276 are analysed Partial and Full horns were grouped together: referred to as 'decidualised'

(Dec). Decidualisation was observed in both genotypes, however Srd5a1[-/-] uteri had a

278 noticeably impaired decidualisation response compared with those of WT. Specifically uteri

from Srd5a1[-/-] mice had fewer 'Full' horns and decidualised areas in 'Partial' horns were

280 smaller (Fig 1C). When quantified and compared statistically we were able to reject the

hypothesis that the number in each of the decidualisation response groups is the same

282 between genotypes, which supports our interpretation that decidualisation is impaired in

Srd5a1[-/-] mice (Fig 1D).

284 In line with expectations uterine weight increased significantly in Dec horns compared to

non-decidualised horns in both WT and Srd5a1[-/-] mice (Fig 1E). Uterine horn weights of

286 partially decidualised horns from WT mice were significantly heavier than those from Srd5a1[-/-] females (Fig 1F). The expression of mRNAs encoded by *Prl8*, a marker of decidualisation,

288 was significantly increased in Dec compared to Non horns in WT mice. A similar, non-

significant, trend for increased *Prl8* mRNA expression was observed in Srd5a1[-/-] mice (Fig

290 1G).

To complement these findings we tested the impact of transiently blocking 5α-reductase

292 activity in WT mice using the pharmacologic 5α-reductase inhibitor finasteride (for 4 days

from days 15-18; Supplementary Figures 1A). Notably this treatment regime did not

294 significantly affect the decidualisation response, uterine horn weights, nor expression of *Prl8*

(Supplementary Figures 1B,C,D). Thus transient inhibition of 5α-reductase under this

296　paradigm was insufficient to replicate the effects of the global *Srd5a1* knock-out on

decidualisation responses.

298　***5α-reductase is required for normal endothelial barrier function and vascular***

remodelling during decidualisation

300　To investigate if impaired decidualisation in Srd5a1$^{-/-}$ mice was also associated with deficient

vascular morphology and function we stained blood vessels with the endothelial cell marker

302　CD31. Observational analysis of tissue sections suggested blood vessels in mutant mice

appeared less numerous and more dilated compared to WT (Fig 2A). To quantify the vessels,

304　counts were performed on samples fluorescently labelled for CD31 (Supplementary Figures

2A). This analysis revealed a significant increase in vessel frequency in decidualised

306　compared to non-decidualised uterine horns (Fig 2B). The mean vessel frequency in

Srd5a1$^{-/-}$ decidualised uteri was lower than in WT but no statistically significant difference

308　was detected (Fig 2B). Vessel irregularity was also quantified, revealing a significant

increase with decidualisation, but no distinct effect of genotype (Supplementary Figures 2B).

310　We attempted to map the three-dimensional architecture of vessels using the resin cast

method, however accurate casts could not be obtained which we believe is due to the high

312　permeability of vessels in the decidualised tissues (Supplementary Figures 3). This finding

and observations from histological slides of dilated vessels in the decidual tissue of Srd5a1$^{-/-}$

314　mice led us to consider whether vessel permeability might be altered as a consequence of

ablation of the gene in mice.

316　To quantify whether these morphological changes lead to functional deficiency in endothelial

barrier integrity, we investigated vascular permeability using Evans Blue assay. We

318　observed a significant increase in permeability of decidualised uterine horns from Srd5a1$^{-/-}$

compared to WT mice (Fig 2C). In mice treated with finasteride, similar trends were

320　observed, and there was a significant increase in permeability in decidualised horns

following finasteride treatment (Fig. 2D).

Changes in angiogenic gene expression during decidualisation are attenuated in Srd5a1^{-/-} mice

To investigate the potential molecular basis for the vascular phenotype observed in Srd5a1$^{-/-}$ uteri, we quantified the expression of key angiogenic genes in decidualised horns from WT

and Srd5a1$^{-/-}$ mice. A significant decrease in the expression of *Pecam1* and *Flt1* was detected in Srd5a1$^{-/-}$ uteri compared to WT (Fig 2E). The mean expression of *Tek*

(TIE2/Angiopoietin-1 receptor) and *Kdr* (VEGFR2) was also decreased but this was not statistically significant (Fig 2E). When gene expression was analysed in WT mice treated

with finasteride there was no detectable change in the expression of the same panel of angiogenic genes (Fig. 2F).

5α-Reductase deficiency attenuates decidualisation-induced changes in gene expression

To discover genes and signalling pathways that are altered by deletion of *Srd5a1* in decidualised endometrium we analysed the transcriptome of Non and Dec uterine tissue

from WT and Srd5a1$^{-/-}$ mice using the Nanostring Pancancer Pathways panel. To visualise the general patterns of gene expression, count values were subjected to unsupervised

hierarchical clustering and visualised on a heat map (Fig. 3A). Clustering of samples forms two major arms, one containing exclusively Dec samples and the other mostly Non. These

findings are consistent with decidualisation being a major driver of changes in gene expression in these samples. Notably, within the Dec arm Srd5a1$^{-/-}$ samples largely

clustered together and closer to the Non arm, and several decidualised Srd5a1$^{-/-}$ samples are found within the Non arm, suggesting a transcriptional phenotype in Srd5a1$^{-/-}$ Dec uterus

which was not the same as in the WT Dec tissue.

To interrogate the differential expression of genes between the two genotypes, we

346 performed four ratio comparisons: Srd5a1$^{-/-}$ Non vs WT Non; Srd5a1$^{-/-}$ Dec vs WT Dec; WT Dec vs WT Non; and Srd5a1$^{-/-}$ Dec vs Srd5a1$^{-/-}$ Non (Fig. 3B). This analysis revealed no

348 significantly differentially expressed (DE) genes between genotypes in non-decidualised

tissue (Srd5a1$^{-/-}$ Non vs WT Non), suggesting a minimal role for 5α-reductase in regulating

350   uterine gene expression in the absence of decidualisation. In stark contrast, decidualisation

in WT mice induced significant changes in the expression of many genes: 507 or 66% of the

352   total analysed (WT Dec vs WT Non; Table 1). Interestingly, the majority (71%) of the DE

genes in this panel were down-regulated by decidualisation. In comparison with the WT data,

354   gene expression changes in Srd5a1$^{-/-}$ decidualised tissues (Srd5a1$^{-/-}$ Dec vs Srd5a1$^{-/-}$ Non)

were greatly suppressed, with fewer DE genes (299, 39%) and smaller changes in

356   expression level for both up- and down-regulated genes (Table 1). When the expression of

genes in decidualised uteri of WT and Srd5a1$^{-/-}$ mice was directly compared (Srd5a1$^{-/-}$ Dec

358   vs WT Dec), many significantly DE genes are observed (254, 33%), most of which are up-

regulated in Srd5a1$^{-/-}$ compared to WT (80%). Notably when significantly DE genes were

360   considered 95% of the up-regulated genes in this comparison are downregulated in the WT

Dec vs WT Non comparison.

362   To identify genes differentially regulated by the Srd5a1$^{-/-}$ genotype, we performed two-way

ANOVA on Nanostring data using the count number of all genes and the variables

364   decidualisation (Non/Dec) and genotype (WT/Srd5a1$^{-/-}$). Results are shown in Table 1. We

identified 47 genes significantly regulated by either genotype or the interaction of genotype

366   and decidualisation, and gene lists can be found in Supplementary Data 2. We further

validated key differentially expressed genes from this set by qPCR in a cohort containing

368   additional samples and these tissues showed gene expression patterns that matched those

seen in the Nanostring counts or followed similar trends (Fig. 3C).

370   ***In silico analysis of gene networks associated with defective decidualisation in***

Srd5a1$^{-/-}$ mice identifies deficient expression of angiogenesis regulatory pathways

372   To gain insight into which gene expression pathways are driving decidualisation, and how

they are affected by deletion of *Srd5a1*, we performed gene ontology (GO) analysis based

374   on DE genes identified from our Nanostring data analysis. Upregulated and downregulated

genes from WT Dec vs WT non and Srd5a1[-/-] Dec vs WT Dec comparisons were analysed

376   separately.

No significant GO terms were identified based on genes upregulated by decidualisation (WT

378   Dec vs WT Non), however in the downregulated genes there were several terms related to

transmembrane transporter activity (Supplementary Figures 4A). Category netplot (cnet)

380   analysis of the enriched genes showed that all the GO terms were related to the same set of

genes (Supplementary Figures 4B). Many of these genes were subunits of the L-type

382   voltage-dependent Ca^{2+} channel (VDCC) which has been shown to be downregulated with

decidualisation in endometrial stromal cells (Kusama et al., 2015). In agreement with these

384   findings, many significant GO terms were identified for genes upregulated in Srd5a1[-/-]

decidua compared to WT (Srd5a1[-/-] Dec Vs WT Dec) all of which related to transmembrane

386   ion transport (Supplementary Figures 4C, D).

Significant GO terms were also identified for genes downregulated in Srd5a1[-/-] decidualised

388   uterus compared to WT (Srd5a1[-/-] Dec Vs WT Dec ; Fig. 4A). The majority related to cell

migration, however there was also a large proportion of GO terms related to the regulation of

390   the vasculature, endothelial cells, or angiogenesis. Cnet analysis highlighted multiple highly

connected genes from canonical angiogenesis pathways such as the VEGF pathway (Fig 4B;

392   *Flt1* (VEGFR1), *Kdr* (VEGFR2), *Vegfa, Pgf*). This Cnet also shows that many of the genes

related to migration are also linked to angiogenesis pathways. This analysis highlights that

394   gene regulatory pathways that promote angiogenesis are significantly downregulated in

decidualised uteri from Srd5a1[-/-] mice.

396   ***Genes differentially regulated by 5α-reductase deficiency are expressed by stromal***
and epithelial cell populations in cycling uterus.

398   To identify which cells in the uterus were likely to have altered gene expression as a result of

deletion of *Srd5a1*, we compared the results of our Nanostring analysis with our previously

400   published single cell RNA sequencing (scRNAseq) data from cycling mouse uterus which

included mesenchymal cells (i.e. fibroblasts and perivascular cells), which are AR[+]

402   (Simitsidellis et al., 2018, Simitsidellis et al., 2016), and epithelial cells (Kirkwood et al.,

       2021). Expression of significantly DE genes from the Srd5a1$^{-/-}$ Dec vs WT Dec ratio analysis

404   were sorted by hierarchical clustering (Fig. 4C). All genes had detectable expression in at

       least one cell population and AR was expressed in stromal (fibroblast>perivascular), but not

406   epithelial, populations. Together, stromal fibroblasts and perivascular cells have detectable

       expression of the majority of the angiogenic and migratory genes whose expression is

408   regulated by 5α-reductase deficiency identified in the cnet plot (Fig 4B). Perivascular cells

       play a key role in supporting the growth, stability, and integrity of blood vessels, as well as

410   playing a major role in angiogenesis. This analysis highlights AR$^+$ perivascular cells and

       fibroblasts as plausible cell types mediating the angiogenesis related effects of SRD5A1

412   activity.

DHT rescues impaired decidualisation response in Srd5a1$^{-/-}$ mice.

414   Given that AR+ cell types were associated with differential gene expression in Srd5a1$^{-/-}$ mice,

       we postulated that a lack of intracrine androgen may have altered cell function during

416   decidualisation in the presence of 5α-reductase deficiency. To test whether impaired

       decidualisation in Srd5a1$^{-/-}$ mice is due to a reduction in intracrine/paracrine DHT, we

418   performed a rescue experiment by administering exogenous DHT coincident with the

       decidualisation stimulus (Fig. 5A). Following this treatment, the rate of decidualisation in

420   DHT-treated mutants (Srd5a1$^{-/-}$-DHT) significantly increased compared to vehicle-treated

       mutants (Srd5a1$^{-/-}$-VC) and was statistically indistinguishable from vehicle-treated WT

422   animals (WT-VC; Fig. 5B). Furthermore, the weight of decidualised uterine horns was also

       the same between WT-VC and Srd5a1$^{-/-}$-DHT groups (Fig. 5C).

424   We also tested whether the vessel permeability phenotype seen in Srd5a1$^{-/-}$ animals could

       be rescued by exogenous DHT and found that Evans Blue dye detection in extracts from

426   decidualised Srd5a1$^{-/-}$-DHT was equivalent to that from WT-VC horns, demonstrating that

       the vascular permeability phenotype is also rescued by DHT administration.

428   Finally, we tested whether altered gene expression identified in the Srd5a1$^{-/-}$ uterus was affected by administration of DHT. Notably, DHT restored mRNA expression levels to WT-

430   VC levels (*Brca1*, *Fgfr1*, *Mapk8ip1*, *Runx1*, *Vegfa*; Fig. 5E), and others showed the same trend (*Flt1*, *Pecam*, *Il11*, *Rasgrp1*; Fig. 5E, Supplementary Figures 5A). For some mRNAs

432   there was no evidence of a rescue (*Itga3*, *Pik3cb*, *Pik3r2*; Supplementary Figures 5B). The most significant change in gene expression in response to DHT supplementation occurred in

434   genes related to angiogenesis signalling and downstream signal transduction pathways such as *Vegfa*, *Fgfr1*, and *Mapk8ip1*, suggesting that these stromal/vascular associated pathways

436   are androgen-regulated during decidualisation.

Overall this shows that gross tissue changes and many of the transcriptomic changes

438   associated with impaired decidualisation in Srd5a1$^{-/-}$ mice can be restored by administration of DHT.

Discussion

Correct and timely decidualisation is essential to the establishment of pregnancy, and defects in decidualisation are linked with implantation failure, pregnancy complications and

menstrual disorders such as heavy bleeding (Dunk et al., 2008, Gibson et al., 2018a, Gibson et al., 2016b). We and others have shown that a correct androgen balance is required for

optimal decidualisation and fertility (Castracane and Asch, 1995, Cloke et al., 2008, De Vries et al., 1998, Abdalla et al., 1998), but the exact nature of this requirement, e.g. through direct

androgen signalling or by acting as precursors to other steroids, and its downstream effects have not been established in an in vivo setting. Identifying the mechanisms through which

androgens support the establishment of pregnancy is therefore essential to our understanding of endometrial function and to the development of future fertility treatments.

To this end, we have utilised a mouse line and pharmacological approaches to disrupt steroid 5α-reductase activity, which is necessary for the production of DHT, in the

endometrium during a model of decidualisation and analysed the effect of this intervention on uterine physiology, transcriptome, and capacity to decidualise.

These investigations have revealed that androgen signalling in the form of locally produced DHT is required for decidualisation to proceed as normal. We have shown that

decidualisation is severely impaired in the absence of SRD5A1, the enzyme responsible for local DHT production, and that exogenous DHT administration rescues this effect. Previously,

we have shown *in vitro* that hESF upregulate androgen biosynthetic enzymes and produce androgens as they are induced to decidualise, and that AR antagonism blocks intracrine

signalling and delays the full differentiation of these cells (Gibson et al., 2016a). The current findings extend this paradigm, demonstrating that intracrine androgen signals are required *in*

vivo to drive decidualisation and promote vascular remodelling.

It is possible that lower decidualisation rates observed in Srd5a1$^{-/-}$ mice represent a delay in

tissue remodelling compared to WT. In the context of the tightly regulated fertility cycles of mammals such a delay would be sufficient to severely affect fertility. This observation aligns

with the reduced fecundity of Srd5a1$^{-/-}$ mice (Mahendroo and Russell, 1999), which may in part be due to decidualisation/implantation defects.

Gene expression changes and the results of the Evans Blue permeability assay suggest a strong role for DHT signalling in vascular remodelling during decidualisation. Decidualised

tissue of Srd5a1$^{-/-}$ mice had downregulated expression (vs WT) of gene pathways from two major areas: cell migration and angiogenesis/vascular genes. The process of angiogenesis

is inseparable from migration, and migratory genes have previously been shown to be androgen-dependant *in vitro* in endometrial stromal cells (Cloke et al., 2008). It is well

established that activated AR promotes, indirectly, VEGF expression and elements of the VEGF pathway (Aslan et al., 2005, Cloke et al., 2008, Joseph et al., 1997, Sordello et al.,

1998); and components of this pathway, such as *Flt1* and *Vegfa*, were substantially downregulated in Srd5a1$^{-/-}$ Dec uteri compared to WT. We propose that the vascular

phenotypes and impaired decidualisation shown here in Srd5a1$^{-/-}$ mice are a direct result of impaired VEGF and other angiogenic signalling due to a lack of promotion from AR. This

paradigm opens up new avenues of exploration for treatments of decidualisation disorders involving impaired androgen signalling as a contributing factor, such as premature ovarian

failure, adrenal insufficiency, or in women of advanced maternal age (Vegunta et al., 2020).

        In our study finasteride administration did not affect the rate of decidualisation or uterine

486   horn weight in WT mice whereas in Srd5a1$^{-/-}$ mice DHT rescued these features. There are several possible explanations for this discrepancy. The simplest is that, since finasteride was

488   only administered at the point of decidualisation stimulus, sufficient residual DHT was already present in the tissue to support a normal decidualisation response (Rittmaster et al.,

490   1988, von Deutsch et al., 2012). Although finasteride inhibits both type I and II 5α-reductase enzymes in rodents (Livingstone et al., 2015, Thigpen and Russell, 1992) some residual

492   SRD5A1 activity will persist in the presence of finasteride (Upreti et al., 2015) and this may also be sufficient to support decidualisation. Furthermore, its possible that global deletion of

494   *Srd5a1* throughout development and the lifespan of these mutant mice has broader impacts

on uterine function that cannot be recapitulated by transient pharmacologic inhibition of the

496   enzyme. Finasteride did however increase vessel permeability during decidualisation in WT

mice, similar to the phenotype observed in Srd5a1-/- mice. These findings may represent

498   distinct responses to intracrine androgens such that AR-positive perivascular cells are more

sensitive to inhibition of 5α-reductase than stromal fibroblasts during decidualisation and/or

500   that the impact of inhibition on these different cell types is time-dependent. It is notable that

endothelial nitric oxide synthase (Nos3), which is a potent, acute mediator of endothelial

502   function, is downregulated during decidualisation in Srd5a1-/- mice (Khorram et al., 1999).

This suggests that pathways downstream of intracrine androgens may be acutely or

504   temporally regulated and thus differentially impacted by transient inhibition via finasteride.

A limitation of this study is that targeting the SRD5A1 protein is likely to have effects on

506   steroid hormone signalling beyond its elimination of DHT. In previous work with the same

mouse line, the cause of its defective parturition phenotype was identified as a lack of

508   another 5α-reduced androgen, 5α-androstan-13α,17β-diol, indicating there may be specific

effects of other 5α-reduced steroids that are not accounted for here (Mahendroo et al., 1999).

510   A separate mechanism by which the Srd5a1[-/-] mutation may affect results is through

diversion of precursor steroids towards other pathways, as occurs during pregnancy when a

512   mid-gestational T surge is aromatised rather than 5α-reduced, causing foetal death due to

estrogen excess in Srd5a1[-/-] mice (Mahendroo et al., 1997). These factors may account for

514   why some gene expression changes were not rescued by DHT administration, and provide

an alternative explanation for why finasteride administration does not fully recapitulate the

516   Srd5a1[-/-] phenotype.

Androgen availability declines with age and there is a concomitant age-related increase in

518   risk of pregnancy disorders (Odibo et al., 2006, Kahveci et al., 2018). The results of the

current study highlight the importance of intracrine androgen signalling to early pregnancy

520   remodelling. Given that intracrine androgen action is directly correlated to precursor

availability (Gibson et al., 2018b), a decrease or deficiency in androgen precursors could

522   directly impact decidualisation capacity. Further studies are therefore warranted to investigate a possible causal link between androgen deficiency and disorders of pregnancy.

524   Future work in response to this study should investigate intracrine androgen signalling effects in the context of natural pregnancy, for example testing whether DHT can improve

526   fecundity in Srd5a1$^{-/-}$ female mice. It will also be valuable to ascertain the exact mechanism through which AR signalling activates the VEGF pathway, and whether stimulating this

528   pathway can likewise improve fertility in the context of androgen depletion or deficiency. This will facilitate an improved diversity of treatment options for cases where androgen

530   supplementation is not desirable (Vegunta et al., 2020).

In summary, using a well validated mouse model we have confirmed that intracrine

532   androgen signalling, stimulated by locally produced DHT, is required for the robust and timely execution of pathways that stimulate gene expression and changes in cell function

534   essential for a robust decidualisation response. By furthering our understating of the role androgens play in endometrial receptivity and related endometrial disorders these findings

536   will aid in the development of fertility treatments and other interventions.

Author contributions:

538   I.W.S designed and performed the experiments, analysed the data, interpreted the results, and wrote the manuscript. P.M.K performed the experiments, analysed the data, and

540   interpreted the results. D.R, R.A, F.L.C & L.B.S performed the experiments. D.L provided study materials. P.T.K.S conceived the study, designed the experiments, interpreted the

542   results and revised the manuscript. D.A.G conceived the study, designed and performed the experiments, analysed the data, interpreted the results, and revised the manuscript.

544   **Acknowledgements**.

Funding: MRC programme grants to PTKS (MR/N024524/1, G1002033); MRC Programme

546   Grant to LBS (MR/N002970/1); Wellcome Trust Fellowship to DAG (220656/Z/20/Z); Society for Endocrinology Early Career Grant to DAG; Scottish Funding Council Research Adaption

548   Fund to IWS; importation of the mice from Jackson laboratories was paid from Wellcome Trust grant 072217/Z/03/Z. We thank Prof. Ruth Andrew (University of Edinburgh) for

550   founder stocks of Srd5a1$^{-/-}$ mice. We are grateful to Dr Alison Munro and the Host and Tumour Profiling Unit (Cancer Research UK – Edinburgh Centre) for support for Nanostring

552   analysis, Dr Pamela Brown from Biomolecular core facility and facility staff from the Bioresearch and Veterinary Services team.

References

556 ABDALLA, H. I., BILLETT, A., KAN, A. K. S., BAIG, S., WREN, M., KOREA, L. & STUDD, J. W. W. 1998. Obstetric outcome in 232 ovum donation pregnancies. *BJOG:*
558 *An International Journal of Obstetrics & Gynaecology,* 105, 332-337.
ASLAN, G., CIMEN, S., YORUKOGLU, K., TUNA, B., SONMEZ, D., MUNGAN, U. &
560 CELEBI, I. 2005. Vascular endothelial growth factor expression in untreated and androgen-deprived patients with prostate cancer. *Pathology - Research and Practice,*
562 201, 593-598.
BENJAMINI, Y. & HOCHBERG, Y. 1995. Controlling the False Discovery Rate : A
564 Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society,* 57, 289-300.
566 CASTRACANE, V. D. & ASCH, R. H. 1995. Pregnancy: Testosterone and androstenedione in premature ovarian failure pregnancies: Evidence for an ovarian source of
568 androgens in early pregnancy. *Human Reproduction,* 10, 677-680.
CLOKE, B., HUHTINEN, K., FUSI, L., KAJIHARA, T., YLIHEIKKILÄ, M., HO, K. K.,
TEKLENBURG, G., LAVERY, S., JONES, M. C., TREW, G., KIM, J. J., LAM, E. W. F., CARTWRIGHT, J. E., POUTANEN, M. & BROSENS, J. J. 2008. The
Androgen and Progesterone Receptors Regulate Distinct Gene Networks and Cellular Functions in Decidualizing Endometrium. *Endocrinology,* 149, 4462-4462.
COUSINS, F. L., MURRAY, A., ESNAL, A., GIBSON, D. A., CRITCHLEY, H. O. D. & SAUNDERS, P. T. K. 2014. Evidence from a mouse model that epithelial cell
migration and mesenchymal-epithelial transition contribute to rapid restoration of uterine tissue integrity during menstruation. *PLoS ONE,* 9, e86378-e86378.
DE VRIES, M. J., DEKKER, G. A. & SCHOEMAKER, J. 1998. Higher risk of preeclampsia in the polycystic ovary syndrome. A case control study. *European Journal of*
Obstetrics and Gynecology and Reproductive Biology, 76, 91-95.
DUNK, C., SMITH, S., HAZAN, A., WHITTLE, W. & LEE JONES, R. 2008. Promotion of
582 angiogenesis by human endometrial lymphocytes.*Immunological Investigations,* 37, 583-610.
584 DUNN, C. L., KELLY, R. W. & CRITCHLEY, H. O. D. 2003. Decidualization of the human endometrial stromal cell: An enigmatic transformation. *Reproductive BioMedicine*
586 *Online,* 7, 151-161.
EDGAR, R., DOMRACHEV, M. & LASH, A. E. 2002. Gene Expression Omnibus: NCBI
588 gene expression and hybridization array data repository. *Nucleic Acids Res,* 30, 207-10.
590 GIBSON, D. A., SIMITSIDELLIS, I., COLLINS, F. & SAUNDERS, P. T. K. 2018a. Endometrial Intracrinology: Oestrogens, Androgens and Endometrial Disorders.
592 *International Journal of Molecular Sciences 2018, Vol. 19, Page 3276,* 19, 3276-3276.
GIBSON, D. A., SIMITSIDELLIS, I., COUSINS, F. L., CRITCHLEY, H. O. D. &
594 SAUNDERS, P. T. K. 2016a. Intracrine Androgens Enhance Decidualization and Modulate Expression of Human Endometrial Receptivity Genes. *Scientific Reports,* 6,
596 27-31.
GIBSON, D. A., SIMITSIDELLIS, I., KELEPOURI, O., CRITCHLEY, H. O. D. &
SAUNDERS, P. T. K. 2018b. Dehydroepiandrosterone enhances decidualization in women of advanced reproductive age. *Fertility and Sterility,* 109, 728-734.e2.
GIBSON, D. A., SIMITSIDELLIS, I. & SAUNDERS, P. T. K. 2016b. Regulation of androgen action during establishment of pregnancy. *Journal of Molecular*
Endocrinology, 57, R35-R47.

JOSEPH, I. B. J. K., NELSON, J. B., DENMEADE, S. R. & ISAACS, J. T. 1997. Androgens regulate vascular endothelial growth factor content in normal and malignant prostatic tissue. *Clinical Cancer Research,* 3**,** 2507-2511.

KAHVECI, B., MELEKOGLU, R., EVRUKE, I. C. & CETIN, C. 2018. The effect of advanced maternal age on perinatal outcomes in nulliparous singleton pregnancies.
BMC Pregnancy and Childbirth, 18**,** 343.

KHORRAM, O., GARTHWAITE, M. & MAGNESS,R. R. 1999. Endometrial and
610 Myometrial Expression of Nitric Oxide Synthase Isoforms in Pre- and Postmenopausal Women. *The Journal of Clinical Endocrinology & Metabolism,* 84**,**
612 2226–2232.

KIRKWOOD, P. M., GIBSON, D. A., SMITH, J. R., WILSON-KANAMORI, J. R.,
614 KELEPOURI, O., ESNAL-ZUFIAURRE, A., DOBIE, R., HENDERSON, N. C. & SAUNDERS, P. T. K. 2021. Single-cell RNA sequencing redefines the mesenchymal
616 cell landscape of mouse endometrium. *FASEB Journal,* 35**,** e21285-e21285.

KUSAMA, K., YOSHIE, M., TAMURA, K., IMAKAWA, K., ISAKA, K. & TACHIKAWA,
618 E. 2015. Regulatory Action of Calcium Ion on Cyclic AMP-Enhanced Expression of Implantation-Related Factors in Human Endometrial Cells. *PloS one,* 10.**,** e0132017.

620 LABRIE, F., BÉLANGER, A., CUSAN, L., GOMEZ, J. L. & CANDAS, B. 1997. Marked decline in serum concentrations of adrenal C19 sex steroid precursors and conjugated
622 androgen metabolites during aging. *J Clin Endocrinol Metab,* 82**,** 2396-402.

LABRIE, F., MARTEL, C., BÉLANGER, A. & PELLETIER, G. 2017. Androgens in women
624 are essentially made from DHEA in each peripheral tissue according to intracrinology. *J Steroid Biochem Mol Biol,* 168**,** 9-18.

626 LIVINGSTONE, D. E. W., BARAT, P., DI ROLLO, E. M., REES, G. A., WELDIN, B. A., ROG-ZIELINSKA, E. A., MACFARLANE, D. P., WALKER, B. R. & ANDREW, R.
628 2015. 5α-Reductase type 1 deficiency or inhibition predisposes to insulin resistance, hepatic steatosis, and liver fibrosis in rodents. *Diabetes,* 64**,** 447-458.

630 MAHENDROO, M. S., CALA, K. M., LANDRUM, C. P. & RUSSELL, D. W. 1997. Fetal Death in Mice Lacking 5α-Reductase Type 1 Caused by Estrogen Excess. *Molecular*
632 *Endocrinology,* 11**,** 917-927.

MAHENDROO, M. S., CALA, K. M. & RUSSELL, D. W. 1996. 5 alpha-reduced androgens
634 play a key role in murine parturition. *Molecular Endocrinology,* 10**,** 380-392.

MAHENDROO, M. S., PORTER, A., RUSSELL, D. W. & WORD, R. A. 1999. The
parturition defect in steroid 5α-reductase type 1 knockout mice is due to impaired cervical ripening. *Molecular Endocrinology,* 13**,** 981-992.

MAHENDROO, M. S. & RUSSELL, D. W. 1999. Male and female isoenzymes of steroid 5α-reductase. *Reviews of Reproduction,* 4**,** 179-183.

MARSHALL, E., LOWREY, J., MACPHERSON, S., MAYBIN, J. A., COLLINS, F., CRITCHLEY, H. O. D. & SAUNDERS, P. T. K. 2011. In silico analysis identifies a
novel role for androgens in the regulation of human endometrial apoptosis. *Journal of Clinical Endocrinology and Metabolism,* 96**,** E1746-E1755.

NARUKAWA, S., KANZAKI, H., INOUE, T., IMAI, K., HIGUCHI, T., HATAYAMA, H., KARIYA, M. & MORI, T. 1994. Androgens induce prolactin production by human
endometrial stromal cells in vitro. *The Journal of Clinical Endocrinology & Metabolism,* 78**,** 165-168.

ODIBO, A. O., NELSON, D., STAMILIO, D. M., SEHDEV, H. M. & MACONES, G. A. 2006. Advanced maternal age is an independent risk factor for intrauterine growth
restriction. *Am J Perinatol,* 23**,** 325-8.

PLAISIER, M. 2011. Decidualisation and angiogenesis. *Best Practice and Research: Clinical*
Obstetrics and Gynaecology, 25**,** 259-271.

REBOURCET, D., WU, J., CRUICKSHANKS, L., SMITH, S. E., MILNE, L., FERNANDO, A., WALLACE, R. J., GRAY, C. D., HADOKE, P. W., MITCHELL, R. T.,
O'SHAUGHNESSY, P. J. & SMITH, L. B. 2016. Sertoli Cells Modulate Testicular Vascular Network Development, Structure, and Function to Influence Circulating Testosterone Concentrations in Adult Male Mice. *Endocrinology,* 157**,** 2479-88.

RITTMASTER, R. S., THOMPSON, D. L., LISTWAK, S. & LORIAUX, D. L. 1988. Androstanediol Glucuronide Isomers in Normal Men and Women and in Men Infused
with Labeled Dihydrotestosterone. *The Journal of Clinical Endocrinology & Metabolism,* 66**,** 212-216.

ROTHMAN, M. S., CARLSON, N. E., XU, M., WANG, C., SWERDLOFF, R., LEE, P., GOH, V. H., RIDGWAY, E. C. & WIERMAN, M. E. 2011. Reexamination of
testosterone, dihydrotestosterone, estradiol and estrone levels across the menstrual cycle and in postmenopausal women measured by liquid chromatography-tandem
mass spectrometry. *Steroids,* 76**,** 177-82.

SIMITSIDELLIS, I., ESNAL-ZUFFIAURE, A., KELEPOURI, O., O'FLAHERTY, E.,
668 GIBSON, D. A. & SAUNDERS, P. T. K. 2019. Selective androgen receptor modulators (SARMs) have specific impacts on the mouse uterus. *Journal of*
670 *Endocrinology,* 242**,** 227-239.

SIMITSIDELLIS, I., GIBSON, D. A., COUSINS, F. L., ESNAL-ZUFIAURRE, A. &
672 SAUNDERS, P. T. K. 2016. A Role for Androgens in Epithelial Proliferation and Formation of Glands in the Mouse Uterus. *Endocrinology,* 157**,** 2116-2128.

674 SIMITSIDELLIS, I., SAUNDERS, P. T. K. & GIBSON, D. A. 2018. Androgens and endometrium: New insights and new targets. *Molecular and Cellular Endocrinology,*
676 465**,** 48-60.

SORDELLO, S., BERTRAND, N. & PLOUËT, J. 1998. Vascular Endothelial Growth Factor
678 Is Up-Regulatedin Vitroandin Vivoby Androgens. *Biochemical and Biophysical Research Communications,* 251**,** 287-290.

680 THIGPEN, A. E. & RUSSELL, D. W. 1992. Four-amino acid segment in steroid 5α-reductase 1 confers sensitivity to finasteride, a competitive inhibitor. *Journal of*
682 *Biological Chemistry,* 267**,** 8577-8583.

UPRETI, R., NAREDO, G., FAQEHI, A. M. M., HUGHES, K. A., STEWART, L. H.,
684 WALKER, B. R., HOMER, N. Z. M. & ANDREW, R. 2015. Simultaneous pharmacokinetic and pharmacodynamic analysis of 5α-reductase inhibitors and
686 androgens by liquid chromatography tandem mass spectrometry. *Talanta,* 131**,** 728-735.

688 VEGUNTA, S., KLING, J. M. & KAPOOR, E. 2020. Androgen Therapy in Women. *Journal of Women's Health,* 29**,** 57-64.

690 VON DEUTSCH, D. A., ABUKHALAF, I. K. & LAPU-BULA, R. 2012. "Anabolic Doping Agents" in Mozayani A. & Raymon L. (eds.) *Handbook of Drug Interactions: A*
692 *Clinical and Forensic Guide***,**. Humana Press, pp. 625-754.

WALTERS, K. A., SIMANAINEN, U. & GIBSON, D. A. 2016. Androgen action in female
694 reproductive physiology. *Current Opinion in Endocrinology, Diabetes and Obesity,* 23**,** 291-296.

696 YANG, F., ZHENG, Q. & JIN, L. 2019. Dynamic Function and Composition Changes of Immune Cells During Normal and Pathological Pregnancy at the Maternal-Fetal
698 Interface.

ZHANG, X. & CROY, B. A. 1996. Maintenance of Decidual Cell Reaction by Androgens in
700 the Mouse1. *Biology of Reproduction,* 55**,** 519-524.

ZUMOFF, B., STRAIN, G. W., MILLER, L. K. & ROSNER, W. 1995. Twenty-four-hour
mean plasma testosterone concentration declines with age in normal premenopausal
women. *J Clin Endocrinol Metab,* 80**,** 1429-30.

Tables and Figures

Ratio Analysis	Differentially expressed genes (DE)			log$_2$FC of DE genes		
	(#)	(% total)	Upregulated (% DE)	Downregulated (Mean ± 95% CI)	Upregulated (Mean ± 95% CI)	Absolute (Mean ± 95% CI)
Srd5a1$^{-/-}$ Non Vs WT Non	0	0	NA	NA	NA	NA
Srd5a1$^{-/-}$ Dec Vs WT Dec	254	33.0	79.5	-0.84 (-0.71–-0.97)	1.56 (1.39–1.74)	1.41 (1.27–1.56)
WT Dec Vs WT Non	507	65.8	29.0	-1.68 (-1.54–-1.82)	1.94 (1.67–2.20)	1.75 (1.63–1.88)
Srd5a1$^{-/-}$ Dec Vs Srd5a1$^{-/-}$ Non	299	38.8	25.4	-0.84 (-0.76–-0.91)	1.51 (1.27–1.76)	1.01 (0.92–1.10)
2-Way ANOVA Analysis						
Decidualisation	464	61.9				
Genotype	27	3.6				
Interaction	33	4.4				

Table 1: Summary of differentially expressed genes using ratio analysis and 2-Way
708 ***ANOVA analysis of Pancancer Pathways Nanostring panel.***

Abbreviations: Non, non-decidualised uterus; Dec, decidualised uterus; DE, differentially

710 expressed; FC, fold-change; CI, confidence interval.

712 ***Figure 1. Decidualisation response is impaired in Srd5a1*^{-/-} *mice.***

A) Schema for the decidualisation induction model. E2 = oestradiol. P4 = progesterone. B)

714 Graphic depicting a uterine horn in various stages of decidualisation. The horn is thicker

where a decidualisation reaction has occurred. A horn is deemed fully decidualised if an

716 uninterrupted stretch of decidualisation has occurred over at least half of the horn length.

Partial decidualisation is any decidualisation response less than this, often occurring as

718 disparate nodes along the horn. C) Representative images of uterine horns from wild type

(WT) and Srd5a1^{-/-} animals, with the classification of each horn given; p = partial

720   decidualisation, f = fully decidualised. D) Contingency table depicting proportion of horn

responses in each genotype. E,F) Wet weight of uterine horns in WT and Srd5a1$^{-/-}$ mice at

722   time of tissue recovery (4 days following decidualisation induction). Non-decidualised horns

have been pooled. E compares all horns with a decidualisation reaction (Full & Partial),

724   whereas F shows analysis of those with a partial reaction. G) Expression of *Prl8* via qPCR in

uterine horns of WT and Srd5a1$^{-/-}$ mice following decidualisation induction. Plots in D

726   analysed by chi-squared test for trend; E, F analysed by one-way ANOVA with Newman-

Keuls *post-hoc* tests; and G analysed by two-way ANOVA with Bonferroni post-hoc tests; ***

728     $p < 0.001$; ** $p < 0.01$; * $p < 0.05$.

Figure 2. Evidence of a vascular phenotype in Srd5a1^{-/-} mice following induction of

732 ***decidualisation.***

A) Representative images of CD31⁺ cells in WT and Srd5a1⁻/⁻ decidua. Zoomed-in views are

734 shown from regions indicated by boxes. Scale bars are 100μm and 50μm respectively. B)

Quantification of vessel density in in WT and Srd5a1⁻/⁻ non-decidualised endometrium and

736   decidua. C, D) Images of decidualised uterine horns following intravenous injection of Evans

Blue and respective quantification of absorbance in uterine tissue extracts in decidualised

738   horns of WT and Srd5a1$^{-/-}$ (C) or vehicle control (VC) and finasteride (Fin) treated mice (D).

E,F). Expression of angiogenic genes via qPCR in decidualised tissue (partial and full) from

740   WT and Srd5a1^{-}/- mice (E), or VC and Fin mice (F). Plots in B analysed by two-way ANOVA

with Bonferroni *post-hoc tests;* C,D,E,F analysed by two-tailed t test. ** $p < 0.01$; * $p < 0.05$.

Figure 3. Effect of 5α-reductase deficiency on global gene expression during

744 **decidualisation.**

A) Unsupervised clustering and heatmap of WT and Srd5a1$^{-/-}$ decidualised and non-

746 decidualised tissue RNA analysed with Nanostring Pancancer Pathways gene expression

panel. The major gene expression differences occur with decidualisation. B) Volcano plots

748 for the pairwise comparisons between experimental groups, as indicated. Significantly

differentially expressed (DE) genes at a false discovery rate (FDR) of 0.05 are coloured pink.

750 C) Selection of genes changing significantly with either genotype, or interaction of genotype

and decidualisation, by two-way ANOVA (FDR = 0.05). Count values in the Nanostring data

752   and respective validation by qPCR in groups with additional samples is shown. Plots in C

        analysed by two-way ANOVA with Bonferroni *post-hoc* tests. * $p < 0.05$; ** $p < 0.01$; *** $p <$

754   0.001.

756 ***Figure 4. Gene networks associated with defective decidualisation in Srd5a1^{-/-} mice.***

A) Biological process gene ontogeny (GO) of genes significantly downregulated in Srd5a1$^{-/-}$

758 vs WT decidualised uterine horns. All significant terms shown. B) Category netplot (cnet)

embedding indicating genes shared between GO terms for genes significantly

760 downregulated in Srd5a1$^{-/-}$ vs WT decidualised uterine horns. C) Heatmap showing

expression of top DE genes from the Srd5a1$^{-/-}$ compared to WT decidualised uterine horns

762 and their relative abundance in uterine cell subsets; epithelial, fibroblast and perivascular, as

derived from annotation in previously published scRNAseq dataset of mouse uteri. Genes

764   appearing in the cnet plot in B are labelled and colour coded based on whether they link to

angiogenesis-related GO terms (green) or not (black).

Figure 5. Impact of DHT administration on decidualisation response in Srd5a1$^{-/-}$ mice.

768   A) Schema of decidualisation induction experiment in WT and Srd5a1$^{-/-}$ mice with addition of

DHT or vehicle control (VC) at point of decidualisation induction. B) Contingency table

770   depicting proportion of horn responses in WT and Srd5a1$^{-/-}$ mice treated with DHT or VC. C)

Wet weight of uterine horns. D) Quantification of Evans Blue dye absorbance in uterine

772   tissue extracts. E) Expression of selected genes analysed by qPCR. Plot in B analysed by

chi-squared test; C,D,E analysed by one-way ANOVA and Neuman-Keuls post-hoc tests. * p

774   < 0.05; ** p < 0.01; *** p < 0.001; ns = no significant differences detected.