A Role for Steroid 5 alpha-reductase 1 in Vascular Remodelling During Endometrial Decidualisation

Shaw, I.W[1]., Kirkwood, P.M.[1], Rebourcet, D.[2], Cousins, F.L. [1], Ainslie, R.J. [1], Livingstone, D.E.W.[3], Smith L.B.[2], Saunders P.T.K. [1], Gibson, D.A[1]*.

1. *Centre for Inflammation Research, The University of Edinburgh*
2. *MRC Centre for Reproductive Health, The University of Edinburgh*
3. *BHF Centre for Cardiovascular Sciences, The University of Edinburgh*

Key words: Decidualisation, angiogenesis, vascular remodelling, 5 alpha-reductase, Srd5a1, dihydrotestosterone, androgen, intracrinology, finasteride.

Short title: DHT-dependent decidualisation

*to whom correspondence should be addressed.

D.A.Gibson@ed.ac.uk

Centre for Inflammation Research, The University of Edinburgh, Institute for Regeneration and Repair, Edinburgh Bioquarter, Edinburgh

16 **Abstract**

Decidualisation is the hormone-dependent process of endometrial remodelling that is

essential for fertility and reproductive health. It is characterised by dynamic changes in the endometrial stromal compartment including differentiation of fibroblasts, immune cell

trafficking and vascular remodelling. Deficits in decidualisation are implicated in disorders of pregnancy such as implantation failure, intra-uterine growth restriction, and pre-eclampsia.

Androgens are key regulators of decidualisation that promote optimal differentiation of stromal fibroblasts and activation of downstream signalling pathways required for

endometrial remodelling. We have shown that androgen biosynthesis, via 5α-reductase-dependent production of dihydrotestosterone, is required for optimal decidualisation of

human stromal fibroblasts *in vitro*, but whether this is required for decidualisation *in vivo* has not been tested.

In the current study we used steroid 5α-reductase type 1 (SRD5A1) deficient mice (*Srd5a1-/-* mice) and a validated model of induced decidualisation to investigate the role of SRD5A1

and intracrine androgen signalling in endometrial decidualisation. We measured decidualisation response (weight/proportion), transcriptomic changes, and morphological

and functional parameters of vascular development. These investigations revealed a striking effect of 5α-reductase deficiency on the decidualisation response. Furthermore, vessel

permeability and transcriptional regulation of angiogenesis signalling pathways, particularly those that involved vascular endothelial growth factor (VEGF), were disrupted in the

absence of 5α-reductase. In *Srd5a1-/-* mice, injection of dihydrotestosterone co-incident with decidualisation restored decidualisation responses, vessel permeability, and expression of

angiogenesis genes to wild type levels.

Androgen availability declines with age which may contribute to age-related risk of

40 pregnancy disorders. These findings show that intracrine androgen signalling is required for optimal decidualisation in vivo and confirm a major role for androgens in the development of

42 the vasculature during decidualisation through regulation of the VEGF pathway. These

findings highlight new opportunities for improving age-related deficits in fertility and

44   pregnancy health by targeting androgen-dependent signalling in the endometrium.

46 **Introduction**

Decidualisation is a fundamental step in the establishment of pregnancy that involves

48 coordinated remodelling of the endometrial stroma. It is a hormone-dependent process

characterised by differentiation of endometrial stromal fibroblasts (hESF) (Dunn et al., 2003),

50 immune cell trafficking (Yang et al., 2019) and vascular remodelling (Plaisier, 2011). Deficits

in decidualisation are implicated in disorders of pregnancy such as implantation failure, intra-

52 uterine growth restriction, and pre-eclampsia (Dunk et al., 2008, Gibson et al., 2016b). Risk

of pregnancy disorders increases with age which may be associated with age-related decline

54 in hormone production or availability.

Androgens are key regulators of decidualisation that promote optimal differentiation of

56 stromal fibroblasts and activation of downstream signalling pathways required for

endometrial remodelling. Androgen supplementation enhances secretion of decidualization

58 markers in hESF *in vitro* (Narukawa et al., 1994). Exogenous dihydrotestosterone (DHT)

enhances and maintains decidualisation responses in mice, an effect which is attenuated by

60 co-administration of the AR antagonist flutamide (Zhang and Croy, 1996). Endometrial

fibroblasts in both mouse and human endometrium express androgen receptor (AR) and

62 their function is altered by AR-dependent signalling (Marshall et al., 2011, Simitsidellis et al.,

2019, Simitsidellis et al., 2016). Using assays that combined *in vitro* decidualisation and

64 knockdown of receptors Cloke et al. reported that AR regulated distinct decidual gene

networks involved in cytoskeletal organization, cell motility and regulation of the cell cycle

66 (Cloke et al., 2008). Using a pharmacologic approach, we found that flutamide attenuated

the expression of decidualisation and endometrial receptivity markers in hESF (Gibson et al.,

68 2016a). Together, these studies confirm the importance of androgens and AR-dependent

signalling for optimal decidualisation responses.

70 In women, androgens and their precursors are abundant in the circulation but production

declines with age (Zumoff et al., 1995, Labrie et al., 1997). Circulating concentrations of the

most potent androgen dihydrotestosterone (DHT) are low but this is because DHT is

primarily a product of local metabolism within target tissues in women; mediating its affects via intracrine signalling (Rothman et al., 2011, Labrie et al., 2017). We have shown that DHT is actively produced by endometrial stromal fibroblasts during decidualisation via expression

of the enzyme SRD5A1 and that the abundance of DHT is affected by availability of steroid precursors such as dehydroepiandrosterone (DHEA) (Gibson et al., 2018a, Gibson et al.,

2016a, Gibson et al., 2018b, Gibson et al., 2016b). When we increased intracrine DHT signalling by supplementing with DHEA we found that decidualisation responses were

enhanced and expression of implantation markers were also increased. Androgen precursor availability and intracrine androgen signalling can therefore dictate the extent of

decidualisation responses.

To date, a role for intracrine androgen signalling in regulating decidualisation *in vivo* has not been rigorously investigated. Previous studies have been hampered by the developmental

uterine defects reported in global AR knock-out female mice (Walters et al., 2016). In the current study, we have used mice which are homozygous for a recessive knock-out mutation

in the gene encoding the type 1 steroid 5α-reductase (SRD5A1) enzyme (*Srd5a1$^{-/-}$*) which lack the capacity to convert T to DHT (Mahendroo et al., 1996). Previously, Srd5a1$^{-/-}$ mice

have been shown to have reduced fertility due to oestrogen-induced foetal death and a parturition defect caused by lack of 5α-androstan-3α,17β-diol, another product of SRD5A1

(Mahendroo et al., 1997). The potential role of this enzyme and the contribution of 5α-reduced androgens to decidualisation has not been tested, but it is plausible that the

reduced fecundity reported in Srd5a1$^{-/-}$ mice (Mahendroo and Russell, 1999) may in part be due to decidualisation defects as a result of intracrine androgen deficiency. To investigate

this, we have assessed the impact of 5α-reductase deletion or inhibition in a mouse model of induced decidualisation.

We show that 5α-reductase deficiency leads to impaired decidualisation, structural and functional changes to decidual blood vessels, and transcriptomic changes affecting

angiogenesis signalling pathways. Deficits in decidualisation in Srd5a1$^{-/-}$ mice were reversed

by restoration of androgen signalling through exogenous DHT administration. We conclude

102   that intracrine androgens are necessary for optimal decidualisation and vascular remodelling

required for establishment of pregnancy.

Materials and Methods

106 *Mice*

All mouse work was performed in accordance with the Animals (Scientific Procedures) Act

108 1986 under UK law and were bred on a C57Bl/6J or C57Bl6/JCrl background. The $Srd5a1^{-/-}$ mouse line was obtained from the Jackson Laboratories (https://www.jax.org/strain/002793).

110 The original paper describing the generation of mice with a null allele was published by

Mahendroo, Cala, and Russel (Mahendroo et al., 1996). Due to parturition defects in

112 homozygous mice heterozygote crosses were used to maintain the colony and produce

$Srd5a1^{-/-}$ progeny. Wildtype (WT) mice were either littermates or purchased from Charles

114 River Laboratories (Tranent, Scotland).

Mouse model of induced decidualisation

116 Mice were ovariectomised to remove endogenous ovarian steroids at day 0 (d0), then

administered oestradiol (E2, Sigma-Aldrich, Poole) by subcutaneous injection daily on d7-9

118 (5ug.kg^{-1} in 200ul sesame oil) and on d13-15 (0.25ug.kg^{-1} in 200ul sesame oil) with a P4

pellet inserted on d13. On day 15 a decidualisation stimulus was administered by

120 intrauterine injection of 20ul sesame oil into the uterus via transvaginal delivery using a non-

surgical embryo transfer device (NSET). Animals were sacrificed 4 days later and uterine

122 tissues collected (Fig. 1A, (Cousins et al., 2014)). The decidualisation response was

immediately scored in each horn as outlined in Fig. 1B. Specifically a non-decidualised horn

124 had no decidualisation reaction whatsoever (Non), whereas a fully decidualised horn had

continuous decidualisation along the majority (>50%) of its length (Full). Partial

126 decidualisation is any level of decidualisation between these two extremes (Fig. 1B).

Collectively, samples designated Full or Partial are referred to as decidualised (Dec).

128 Finasteride (Sigma-Aldrich, Poole) was dissolved in 100% EtOH, then diluted to 5% v/v in

sesame oil to a final concentration of 12.5mg.mL^{-1} and 100uL (50 mg.kg^{-1}) was administered

130 via intraperitoneal injection daily from the point of the decidualisation stimulus onwards, i.e.

d15-18, as show in Figure S1A. Finasteride is a synthetic 4-azasteroid that competitively

132   inhibits both type I and II 5α-reductase enzymes in rodents (Livingstone et al., 2015, Thigpen

and Russell, 1992).

134   DHT (Sigma-Aldrich, Poole) was dissolved in 100% ethanol (EtOH) then diluted to 5% v/v in

0.4% w/v methylcellulose solution for a final DHT concentration of 2mg.mL^{-1}. 100uL (8mg.kg^{-}

136   ^{1}) was injected subcutaneously on d15 at the point of decidualisation stimulus. At the end of

experiments mice were terminated by exposure to rising concentrations of CO_2 combined

138   with cervical dislocation.

Histology and immunolabelling

140   Uterine tissue for histology was fixed in 4% paraformaldehyde then dehydrated and

embedded in paraffin blocks and sectioned according to standard protocols. Protocols for

142   3,3'-diaminobenzidine (DAB) and immunofluorescence staining were performed as

previously reported (Kirkwood et al., 2021, Simitsidellis et al., 2016). Briefly, slides were

144   dewaxed, rehydrated, and subjected to antigen retrieval in pH 6.0 citrate buffer using a

microwave. Following a wash in phosphate buffered saline (PBS) slides were incubated with

146   3% H_2O_2 solution in PBS for 15min to remove endogenous peroxidase activity. Following

several PBS washes slides were incubated with normal goat serum solution (NGS, PBS

148   containing 20% goat serum and 0.05% (w/v) bovine serum albumin) to block non-specific

interactions. Anti-CD31 primary antibody (Abcam, ab28364, 1:500 dilution) in NGS was

150   added and incubated overnight at 4^{o}C. Following several washes, slides were incubated with

goat-raised peroxidase-conjugated secondary antibody (Abcam, ab7171, 1:500 dilution) for

152   1hr at room temperature. Following several washes, slides were incubated with either DAB

reagent or Opal red 570 reagent for 10 minutes or until sufficiently developed. After a brief

154   wash, slides were counterstained either with haematoxylin and eosin as per standard

procedures, or with DAPI before being mounted with coverslips.

156   Slides were imaged using a Zeiss Z1 microscope, or a Zeiss AxioScan Slide Scanner using

conventional fluorescent setups.

158   ***Vessel quantification***

To quantify vessels, regions of non-decidualised endometrium (Non) or decidua (Dec

160   samples) were selected digitally on 20X magnification slide scans of CD31-labelled uterus

sections and overlaid with a grid with edge length 250 pixels (~ 28µm). The intersections on

162   the grid were sequentially classified as either vessel or not vessel, and the total proportion of

vessel to non-vessel intersections was used as a measure of vessel count. Additionally,

164   each time a vessel was identified, it was given an irregularity score equalling the number of

intersections on a tree used to trace out the vessel's cross-sectional shape. For example, a

166   circular or or 'L' shaped cross-section would have a score of 0, a 'T' shape a score of 1, an

'H' shape a score of 2, etc. Irregularity scores were averaged across all observations.

168   ***RNA extraction***

Uterine samples for RNA analysis were collected in RNALater solution (Qiagen, Hilden,

170   Germany) and stored at -80^{o}C until tissue extraction. For decidualised samples, only regions

of uterus containing decidualised endometrium were selected. Samples were thawed and

172   homogenised in 1 mL Trizol reagent using a TissueLyser machine (6min, 25Hz).

Homogenate was transferred to MaXtract High Density tubes along with 200ul chloroform

174   and 100ul RNase free water, shaken vigorously and centrifuged. The aqueous layer was

added to RNeasy spin columns (Qiagen, Hilden, Germany) and processed as per

176   manufacturer's instructions.

Quantitative RT-PCR

178   cDNA was synthesised from 100ng.uL^{-1} RNA using Superscript VILO cDNA synthesis kit

(Thermo Fisher Scientific Life Sciences, Schwerte, Germany) as per the manufacturer's

180   instructions. Thermal cycler settings were 25^{o}C for 10min, 42^{o}C for 60min, 85^{o}C for 5min.

Primers were designed using Universal Probe Library Assay Design Center (Roche Applied

182   Science, Burgess Hill, UK) or, when this was discontinued, using Neoformit

(https://primers.neoformit.com/ Accessed: 6/4/22; sequences in Supplementary Data 1).

184   Primers were synthesised by Eurofins MWG Operon (Ebersberg, Gemany). Reactions were
       run in duplicate on a Quantstudio 5 384-well PCR machine (Thermo Fisher Scientific Life
186   Sciences, Schwerte, Germany) using the following settings: 95^{o}C for 10min then 40 cycles of
       95^{o}C for 15s and 60^{o}C for 1min. A serial dilution of a standard cDNA solution (consisting of a
188   mix of all samples tested) was used to plot a standard curve and expression levels
       interpolated from this. Data was processed using QuantStudio Design and Analysis Software.

190   ***Nanostring gene expression analysis and data processing***

       Samples were analysed on the NanoString nCounter Analysis System using a NanoString
192   Mouse Pancancer Pathways Panel kit (Nanostring, Edinburgh, UK). RNA at 20ng.uL^{-1} was
       used. Processing was performed by the Host and Tumour Profiling Unit Microarray Services,
194   Institute of Genetics and Cancer, University of Edinburgh, as per the manufacturer's
       instructions. Briefly, the reporter codeset was mixed with hybridisation buffer and added to
196   each of the RNA samples, followed by the capture codeset. This mix was hybridised at 65^{o}C
       for 18hr. Following hybridisation samples were loaded into the provided cartridges on the
198   nCounter prep station and processed using the High Sensitivity protocol. Cartridges were
       then sealed and read using the digital analyser on the max setting. No QC flags were
200   registered for any of the samples.

       Data was processed using the nSolver analysis software. Data was normalised using a
202   combination of positive controls and housekeeping genes. Firstly, background was
       subtracted using a background value of the mean plus two standard deviations of the
204   negative control values. Normalisation using positive controls and housekeeper genes was
       performed using the geometric mean to compute the normalisation factor, and the
206   manufacturer-provided list of housekeeper genes was used excluding the following genes
       based on the results of the geNorm algorithm in the Nanostring Advanced Analysis modules:
208   *Hprt*, *Alas1*, *G6pdx*, *Gusb*, *Ppia*.

       Normalised counts data was then analysed for differential gene expression using R,
210   specifically   the   edgeR   analysis   workflow.   Dispersions   were   estimated   with

the *estimateDisp()* command and fit to the design model using *glmQLFit()*, before testing for

212 differential expression using *glmQLFTest()*. Graphs were produced using the ggplot2

package. Finally, to perform gene ontogeny (GO) analysis we utilised the clusterProfiler

214 analysis workflow. Ensembl IDs were mapped from the Bioconductor org.Mm.eg.db genome

annotation and enrichment assessed using the *enrichGO()* command with the 'universe' set

216 as the genes included in the Nanostring Pancancer Pathways panel. Graphs were produced

using the *dotplot ()* and *cnetplot ()* commands from the enrichplot R package.

218 ***Comparison of Nanostring gene expression with scRNAseq gene expression***

Normalised counts and differential gene expression data was further analysed in the context

220 of publicly available single cell RNA sequencing (scRNAseq) data, accessed from NCBI's

Gene Expression Omnibus (Edgar et al., 2002) through GEO Series accession numbers

222 GSE160772 and GSE198556. These scRNAseq data defined the transcriptomic profile of

over 20,000 perivascular, fibroblast and epithelial cell types present in the mouse uterus.

224 The AverageExpression() command (Seurat) was used to calculate the mean expression of

Nanostring DEgenes in perivascular, fibroblast and epithelial cell clusters from scRNAseq

226 data. Gene expression matrices were normalised and clustered heatmaps produced using

the pheatmap package. Expression values shown are scaled but not centred. This

228 comparison allowed us to attribute the expression of genes that were found to be

differentially expressed between Srd5a1$^{-/-}$ and WT genotypes to certain cell fractions in the

230 mouse endometrium.

Creation and imaging of resin casts

232 Resin casts were prepared using a method previously validated in male mice (Rebourcet et

al., 2016). Briefly, female mice (Non and Dec) were culled using a terminal dose of sodium

234 pentobarbital (150 mg/kg, ip) and perfusion fixation of the vasculature was achieved via the

left ventricle. Heparinized PBS (heparin, 20 U/mL) was infused at 6 mL/min for 2 minutes.

236 Low-viscosity resin (10 mL; Microfil MV-122; Flow Tech Inc) was prepared according to the

manufacturer's instructions and then infused via the left ventricle. Uterine tissues were

238   recovered, trimmed, fixed in paraformaldehyde and embedded in 1.5% low melting point

agarose (Invitrogen, UK), dehydrated in methanol (100%; 24 hours) and optically cleared in

240   benzyl alcohol:benzyl benzoate (1:2 v/v; 24 hours). Imaging was carried out as described in

Rebourcet et al. with a positive control tissue being a previously prepared mouse testis

242   (Rebourcet et al., 2016).

Evans Blue assay of vascular permeability

244   Evans Blue (EB) dye binds to albumin in the blood so its presence in tissues, which can be

quantified spectrophotometrically, implies a breakdown in the endothelial barrier. The Evans

246   Blue dye (Sigma-Aldrich, Poole) was dissolved in sterile saline solution to a concentration of

0.5% w/v, and 200uL was injected i.v. (via tail vein) 30-90 minutes prior to cull via cervical

248   dislocation. The interval between EB injection and cull was recorded and found not to

associate with dye retention in tissues.

250   To quantify the amount of Evans blue in the tissues, following sacrifice a portion of the

uterus was taken, precisely weighed, and transferred to 500µl of formamide. The gross

252   appearance of the tissue was recorded using a digital camera. For decidualised samples,

only sections of uterus containing decidualised endometrium were selected. Samples were

254   incubated for 24h at 55^{o}C to extract the dye, then tissue pieces were discarded, the solution

centrifuged to remove debris, and absorbance read in triplicate at 610nm.

256   ### Data processing and statistical analysis

Unless otherwise stated, data was processed using Microsoft Excel and Python. Statistics

258   were performed using either Python or Graphpad Prism 5.

In the analysis of Nanostring data, two-way ANOVAs were performed on all genes found to

260   be expressed and ANOVA results corrected for multiple comparisons using the Benjamini

and Hochberg method (Benjamini and Hochberg, 1995). The false discovery rate (FDR)

262   was 0.05.

In the analysis of Evans Blue absorbance data, absorbance readings are shown as a ratio

264 relative to the mean value of WT or vehicle-treated WT data. In the case of data from Figure

2, ratios have been calculated separately for partially and fully decidualised horns and the

266 results combined for the subsequent statistical analysis.

Statistical tests performed for other comparisons are stated in figure legends and the text.

Results

270   *5α-Reductase deficiency impairs decidualisation in mouse uterus*

To investigate the functional requirement for 5α-reductase during decidualisation we induced

decidualisation in WT and Srd5a1$^{-/-}$ mice using a previously validated protocol (Fig. 1A; (Cousins et al., 2014)). We used a ternary system to classify the extent of decidualisation.

Uterine horns were classified as non-, partially- or fully-decidualised (Non, Partial, Full, respectively; Fig. 1B). For the purposes of analysis individual uterine horns were treated as

separate biological entities. In instances where only the decidualised regions of the uterus are analysed Partial and Full horns were grouped together: referred to as 'decidualised'

(Dec). Decidualisation was observed in both genotypes, however Srd5a1$^{-/-}$ uteri had a noticeably impaired decidualisation response compared with those of WT. Specifically uteri

from Srd5a1$^{-/-}$ mice had fewer 'Full' horns and decidualised areas in 'Partial' horns were smaller (Fig 1C). When quantified and compared statistically we were able to reject the

hypothesis that the number in each of the decidualisation response groups is the same between genotypes, which supports our interpretation that decidualisation is impaired in

Srd5a1$^{-/-}$ mice (Fig 1D).

In line with expectations uterine weight increased significantly in Dec horns compared to

286   non-decidualised horns in both WT and Srd5a1$^{-/-}$ mice (Fig 1E). Uterine horn weights of partially decidualised horns from WT mice were significantly heavier than those from Srd5a1^{-}

288     $^{/-}$ females (Fig 1F). The expression of mRNAs encoded by *Prl8*, a marker of decidualisation, was significantly increased in Dec compared to Non horns in WT mice. A similar, non-

290   significant, trend for increased *Prl8* mRNA expression was observed in Srd5a1$^{-/-}$ mice (Fig 1G).

292   To complement these findings we tested the impact of transiently blocking 5α-reductase activity in WT mice using the pharmacologic 5α-reductase inhibitor finasteride (for 4 days

294   from days 15-18; Fig S1A). Notably this treatment regime did not significantly affect the decidualisation response, uterine horn weights, nor expression of *Prl8* (Fig S1B,C,D). Thus

296 transient inhibition of 5α-reductase under this paradigm was insufficient to replicate the

effects of the global *Srd5a1* knock-out on decidualisation responses.

298 ***5α-reductase is required for normal endothelial barrier function and vascular***
remodelling during decidualisation

300 To investigate if impaired decidualisation in Srd5a1$^{-/-}$ mice was also associated with deficient

vascular morphology and function we stained blood vessels with the endothelial cell marker

302 CD31. Observational analysis of tissue sections suggested blood vessels in mutant mice

appeared less numerous and more dilated compared to WT (Fig 2A). To quantify the vessels,

304 counts were performed on samples fluorescently labelled for CD31 (Fig. S2A). This analysis

revealed a significant increase in vessel frequency in decidualised compared to non-

306 decidualised uterine horns (Fig 2B). The mean vessel frequency in Srd5a1$^{-/-}$ decidualised

uteri was lower than in WT but no statistically significant difference was detected (Fig 2B).

308 Vessel irregularity was also quantified, revealing a significant increase with decidualisation,

but no distinct effect of genotype (Fig S2B).

310 We attempted to map the three-dimensional architecture of vessels using the resin cast

method, however accurate casts could not be obtained which we believe is due to the high

312 permeability of vessels in the decidualised tissues (Fig. S3). This finding and observations

from histological slides of dilated vessels in the decidual tissue of Srd5a1$^{-/-}$ mice led us to

314 consider whether vessel permeability might be altered as a consequence of ablation of the

gene in mice.

316 To quantify whether these morphological changes lead to functional deficiency in endothelial

barrier integrity, we investigated vascular permeability using Evans Blue assay. We

318 observed a significant increase in permeability of decidualised uterine horns from Srd5a1$^{-/-}$

compared to WT mice (Fig 2C). In mice treated with finasteride, similar trends were

320 observed, and there was a significant increase in permeability in decidualised horns

following finasteride treatment (Fig. 2D).

Changes in angiogenic gene expression during decidualisation are attenuated in Srd5a1^{-/-} mice

To investigate the potential molecular basis for the vascular phenotype observed in Srd5a1$^{-/-}$ uteri, we quantified the expression of key angiogenic genes in decidualised horns from WT

and Srd5a1$^{-/-}$ mice. A significant decrease in the expression of *Pecam1* and *Flt1* was detected in Srd5a1$^{-/-}$ uteri compared to WT (Fig 2E). The mean expression of *Tek*

(TIE2/Angiopoietin-1 receptor) and *Kdr* (VEGFR2) was also decreased but this was not statistically significant (Fig 2E). When gene expression was analysed in WT mice treated

with finasteride there was no detectable change in the expression of the same panel of angiogenic genes (Fig. 2F).

5α-Reductase deficiency attenuates decidualisation-induced changes in gene expression

To discover genes and signalling pathways that are altered by deletion of *Srd5a1* in decidualised endometrium we analysed the transcriptome of Non and Dec uterine tissue

from WT and Srd5a1$^{-/-}$ mice using the Nanostring Pancancer Pathways panel. To visualise the general patterns of gene expression, count values were subjected to unsupervised

hierarchical clustering and visualised on a heat map (Fig. 3A). Clustering of samples forms two major arms, one containing exclusively Dec samples and the other mostly Non. These

findings are consistent with decidualisation being a major driver of changes in gene expression in these samples. Notably, within the Dec arm Srd5a1$^{-/-}$ samples largely

clustered together and closer to the Non arm, and several decidualised Srd5a1$^{-/-}$ samples are found within the Non arm, suggesting a transcriptional phenotype in Srd5a1$^{-/-}$ Dec uterus

which was not the same as in the WT Dec tissue.

To interrogate the differential expression of genes between the two genotypes, we

346 performed four ratio comparisons: Srd5a1$^{-/-}$ Non vs WT Non; Srd5a1$^{-/-}$ Dec vs WT Dec; WT Dec vs WT Non; and Srd5a1$^{-/-}$ Dec vs Srd5a1$^{-/-}$ Non (Fig. 3B). This analysis revealed no

348 significantly differentially expressed (DE) genes between genotypes in non-decidualised

tissue (Srd5a1$^{-/-}$ Non vs WT Non), suggesting a minimal role for 5α-reductase in regulating

350   uterine gene expression in the absence of decidualisation. In stark contrast, decidualisation

in WT mice induced significant changes in the expression of many genes: 507 or 66% of the

352   total analysed (WT Dec vs WT Non; Table 1). Interestingly, the majority (71%) of the DE

genes in this panel were down-regulated by decidualisation. In comparison with the WT data,

354   gene expression changes in Srd5a1$^{-/-}$ decidualised tissues (Srd5a1$^{-/-}$ Dec vs Srd5a1$^{-/-}$ Non)

were greatly suppressed, with fewer DE genes (299, 39%) and smaller changes in

356   expression level for both up- and down-regulated genes (Table 1). When the expression of

genes in decidualised uteri of WT and Srd5a1$^{-/-}$ mice was directly compared (Srd5a1$^{-/-}$ Dec

358   vs WT Dec), many significantly DE genes are observed (254, 33%), most of which are up-

regulated in Srd5a1$^{-/-}$ compared to WT (80%). Notably when significantly DE genes were

360   considered 95% of the up-regulated genes in this comparison are downregulated in the WT

Dec vs WT Non comparison.

362   To identify genes differentially regulated by the Srd5a1$^{-/-}$ genotype, we performed two-way

ANOVA on Nanostring data using the count number of all genes and the variables

364   decidualisation (Non/Dec) and genotype (WT/Srd5a1$^{-/-}$). Results are shown in Table 1. We

identified 47 genes significantly regulated by either genotype or the interaction of genotype

366   and decidualisation, and gene lists can be found in Supplementary Data 2. We further

validated key differentially expressed genes from this set by qPCR in a cohort containing

368   additional samples and these tissues showed gene expression patterns that matched those

seen in the Nanostring counts or followed similar trends (Fig. 3C).

370   ***In silico analysis of gene networks associated with defective decidualisation in***

Srd5a1$^{-/-}$ mice identifies deficient expression of angiogenesis regulatory pathways

372   To gain insight into which gene expression pathways are driving decidualisation, and how

they are affected by deletion of *Srd5a1*, we performed gene ontology (GO) analysis based

374   on DE genes identified from our Nanostring data analysis. Upregulated and downregulated

genes from WT Dec vs WT non and Srd5a1$^{-/-}$ Dec vs WT Dec comparisons were analysed

376   separately.

No significant GO terms were identified based on genes upregulated by decidualisation (WT

378   Dec vs WT Non), however in the downregulated genes there were several terms related to

transmembrane transporter activity (Fig. S4A). Category netplot (cnet) analysis of the

380   enriched genes showed that all the GO terms were related to the same set of genes (Fig

S4B). Many of these genes were subunits of the L-type voltage-dependent Ca^{2+} channel

382   (VDCC) which has been shown to be downregulated with decidualisation in endometrial

stromal cells (Kusama et al., 2015). In agreement with these findings, many significant GO

384   terms were identified for genes upregulated in Srd5a1$^{-/-}$ decidua compared to WT (Srd5a1$^{-/-}$

Dec Vs WT Dec) all of which related to transmembrane ion transport (Fig. S4C, D).

386   Significant GO terms were also identified for genes downregulated in Srd5a1$^{-/-}$ decidualised

uterus compared to WT (Srd5a1$^{-/-}$ Dec Vs WT Dec ; Fig. 4A). The majority related to cell

388   migration, however there was also a large proportion of GO terms related to the regulation of

the vasculature, endothelial cells, or angiogenesis. Cnet analysis highlighted multiple highly

390   connected genes from canonical angiogenesis pathways such as the VEGF pathway (Fig 4B;

Flt1 (VEGFR1), *Kdr* (VEGFR2), *Vegfa, Pgf*). This Cnet also shows that many of the genes

392   related to migration are also linked to angiogenesis pathways. This analysis highlights that

gene regulatory pathways that promote angiogenesis are significantly downregulated in

394   decidualised uteri from Srd5a1$^{-/-}$ mice.

Genes differentially regulated by 5α-reductase deficiency are expressed by stromal

396   ***and epithelial cell populations in cycling uterus.***

To identify which cells in the uterus were likely to have altered gene expression as a result of

398   deletion of *Srd5a1*, we compared the results of our Nanostring analysis with our previously

published single cell RNA sequencing (scRNAseq) data from cycling mouse uterus which

400   included mesenchymal cells (i.e. fibroblasts and perivascular cells), which are AR^{+}

(Simitsidellis et al., 2018, Simitsidellis et al., 2016), and epithelial cells (Kirkwood et al.,

402   2021). Expression of significantly DE genes from the Srd5a1$^{-/-}$ Dec vs WT Dec ratio analysis

were sorted by hierarchical clustering (Fig. 4C). All genes had detectable expression in at

404   least one cell population and AR was expressed in stromal (fibroblast>perivascular), but not

epithelial, populations. Together, stromal fibroblasts and perivascular cells have detectable

406   expression of the majority of the angiogenic and migratory genes whose expression is

regulated by 5α-reductase deficiency identified in the cnet plot (Fig 4B). Perivascular cells

408   play a key role in supporting the growth, stability, and integrity of blood vessels, as well as

playing a major role in angiogenesis. This analysis highlights AR$^+$ perivascular cells and

410   fibroblasts as plausible cell types mediating the angiogenesis related effects of SRD5A1

activity.

412   ***DHT rescues impaired decidualisation response in Srd5a1$^{-/-}$ mice.***

Given that AR+ cell types were associated with differential gene expression in Srd5a1$^{-/-}$ mice,

414   we postulated that a lack of intracrine androgen may have altered cell function during

decidualisation in the presence of 5α-reductase deficiency. To test whether impaired

416   decidualisation in Srd5a1$^{-/-}$ mice is due to a reduction in intracrine/paracrine DHT, we

performed a rescue experiment by administering exogenous DHT coincident with the

418   decidualisation stimulus (Fig. 5A). Following this treatment, the rate of decidualisation in

DHT-treated mutants (Srd5a1$^{-/-}$-DHT) significantly increased compared to vehicle-treated

420   mutants (Srd5a1$^{-/-}$-VC) and was statistically indistinguishable from vehicle-treated WT

animals (WT-VC; Fig. 5B). Furthermore, the weight of decidualised uterine horns was also

422   the same between WT-VC and Srd5a1$^{-/-}$-DHT groups (Fig. 5C).

We also tested whether the vessel permeability phenotype seen in Srd5a1$^{-/-}$ animals could

424   be rescued by exogenous DHT and found that Evans Blue dye detection in extracts from

decidualised Srd5a1$^{-/-}$-DHT was equivalent to that from WT-VC horns, demonstrating that

426   the vascular permeability phenotype is also rescued by DHT administration.

Finally, we tested whether altered gene expression identified in the Srd5a1$^{-/-}$ uterus was

428   affected by administration of DHT. Notably, DHT restored mRNA expression levels to WT-

VC levels (*Brca1*, *Fgfr1*, *Mapk8ip1*, *Runx1*, *Vegfa*; Fig. 5E), and others showed the same

430   trend (*Flt1*, *Pecam*, *Il11*, *Rasgrp1*; Fig. 5E, Fig. S5A). For some mRNAs there was no

evidence of a rescue (*Itga3*, *Pik3cb*, *Pik3r2*; Fig. S5B). The most significant change in gene

432   expression in response to DHT supplementation occurred in genes related to angiogenesis

signalling and downstream signal transduction pathways such as *Vegfa*, *Fgfr1*, and

434   *Mapk8ip1*, suggesting that these stromal/vascular associated pathways are androgen-

regulated during decidualisation.

436   Overall this shows that gross tissue changes and many of the transcriptomic changes

associated with impaired decidualisation in Srd5a1$^{-/-}$ mice can be restored by administration

438   of DHT.

440 **Discussion**

Correct and timely decidualisation is essential to the establishment of pregnancy, and

442 defects in decidualisation are linked with implantation failure, pregnancy complications and

menstrual disorders such as heavy bleeding (Dunk et al., 2008, Gibson et al., 2018a, Gibson

444 et al., 2016b). We and others have shown that a correct androgen balance is required for

optimal decidualisation and fertility (Castracane and Asch, 1995, Cloke et al., 2008, De Vries

446 et al., 1998, Abdalla et al., 1998), but the exact nature of this requirement, e.g. through direct

androgen signalling or by acting as precursors to other steroids, and its downstream effects

448 have not been established in an in vivo setting. Identifying the mechanisms through which

androgens support the establishment of pregnancy is therefore essential to our

450 understanding of endometrial function and to the development of future fertility treatments.

To this end, we have utilised a mouse line and pharmacological approaches to disrupt

452 steroid 5α-reductase activity, which is necessary for the production of DHT, in the

endometrium during a model of decidualisation and analysed the effect of this intervention

454 on uterine physiology, transcriptome, and capacity to decidualise.

These investigations have revealed that androgen signalling in the form of locally produced

456 DHT is required for decidualisation to proceed as normal. We have shown that

decidualisation is severely impaired in the absence of SRD5A1, the enzyme responsible for

458 local DHT production, and that exogenous DHT administration rescues this effect. Previously,

we have shown *in vitro* that hESF upregulate androgen biosynthetic enzymes and produce

460 androgens as they are induced to decidualise, and that AR antagonism blocks intracrine

signalling and delays the full differentiation of these cells (Gibson et al., 2016a). The current

462 findings extend this paradigm, demonstrating that intracrine androgen signals are required *in

vivo* to drive decidualisation and promote vascular remodelling.

464 It is possible that lower decidualisation rates observed in Srd5a1$^{-/-}$ mice represent a delay in

tissue remodelling compared to WT. In the context of the tightly regulated fertility cycles of

mammals such a delay would be sufficient to severely affect fertility. This observation aligns

with the reduced fecundity of Srd5a1$^{-/-}$ mice (Mahendroo and Russell, 1999), which may in part be due to decidualisation/implantation defects.

Gene expression changes and the results of the Evans Blue permeability assay suggest a strong role for DHT signalling in vascular remodelling during decidualisation. Decidualised tissue of Srd5a1$^{-/-}$ mice had downregulated expression (vs WT) of gene pathways from two

major areas: cell migration and angiogenesis/vascular genes. The process of angiogenesis is inseparable from migration, and migratory genes have previously been shown to be

androgen-dependant *in vitro* in endometrial stromal cells (Cloke et al., 2008). It is well established that activated AR promotes, indirectly, VEGF expression and elements of the

VEGF pathway (Aslan et al., 2005, Cloke et al., 2008, Joseph et al., 1997, Sordello et al., 1998); and components of this pathway, such as *Flt1* and *Vegfa*, were substantially

downregulated in Srd5a1$^{-/-}$ Dec uteri compared to WT. We propose that the vascular phenotypes and impaired decidualisation shown here in Srd5a1$^{-/-}$ mice are a direct result of

impaired VEGF and other angiogenic signalling due to a lack of promotion from AR. This paradigm opens up new avenues of exploration for treatments of decidualisation disorders

involving impaired androgen signalling as a contributing factor, such as premature ovarian failure, adrenal insufficiency, or in women of advanced maternal age (Vegunta et al., 2020).

In our study finasteride administration did not affect the rate of decidualisation or uterine horn weight in WT mice whereas in Srd5a1$^{-/-}$ mice DHT rescued these features. There are

several possible explanations for this discrepancy. The simplest is that, since finasteride was only administered at the point of decidualisation stimulus, sufficient residual DHT was

already present in the tissue to support a normal decidualisation response (Rittmaster et al., 1988, von Deutsch et al., 2012). Although finasteride inhibits both type I and II 5α-reductase

enzymes in rodents (Livingstone et al., 2015, Thigpen and Russell, 1992) some residual SRD5A1 activity will persist in the presence of finasteride (Upreti et al., 2015) and this may

also be sufficient to support decidualisation. Furthermore, its possible that global deletion of *Srd5a1* throughout development and the lifespan of these mutant mice has broader impacts

on uterine function that cannot be recapitulated by transient pharmacologic inhibition of the

enzyme. Finasteride did however increase vessel permeability during decidualisation in WT

496   mice, similar to the phenotype observed in Srd5a1-/- mice. These findings may represent

distinct responses to intracrine androgens such that AR-positive perivascular cells are more

498   sensitive to inhibition of 5α-reductase than stromal fibroblasts during decidualisation and/or

that the impact of inhibition on these different cell types is time-dependent. It is notable that

500   endothelial nitric oxide synthase (Nos3), which is a potent, acute mediator of endothelial

function, is downregulated during decidualisation in Srd5a1-/- mice (Khorram et al., 1999).

502   This suggests that pathways downstream of intracrine androgens may be acutely or

temporally regulated and thus differentially impacted by transient inhibition via finasteride.

504   A limitation of this study is that targeting the SRD5A1 protein is likely to have effects on

steroid hormone signalling beyond its elimination of DHT. In previous work with the same

506   mouse line, the cause of its defective parturition phenotype was identified as a lack of

another 5α-reduced androgen, 5α-androstan-13α,17β-diol, indicating there may be specific

508   effects of other 5α-reduced steroids that are not accounted for here (Mahendroo et al., 1999).

A separate mechanism by which the Srd5a1$^{-/-}$ mutation may affect results is through

510   diversion of precursor steroids towards other pathways, as occurs during pregnancy when a

mid-gestational T surge is aromatised rather than 5α-reduced, causing foetal death due to

512   estrogen excess in Srd5a1$^{-/-}$ mice (Mahendroo et al., 1997). These factors may account for

why some gene expression changes were not rescued by DHT administration, and provide

514   an alternative explanation for why finasteride administration does not fully recapitulate the

Srd5a1$^{-/-}$ phenotype.

516   Androgen availability declines with age and there is a concomitant age-related increase in

risk of pregnancy disorders (Odibo et al., 2006, Kahveci et al., 2018). The results of the

518   current study highlight the importance of intracrine androgen signalling to early pregnancy

remodelling. Given that intracrine androgen action is directly correlated to precursor

availability (Gibson et al., 2018b), a decrease or deficiency in androgen precursors could

directly impact decidualisation capacity. Further studies are therefore warranted to investigate a possible causal link between androgen deficiency and disorders of pregnancy. Future work in response to this study should investigate intracrine androgen signalling

effects in the context of natural pregnancy, for example testing whether DHT can improve fecundity in Srd5a1$^{-/-}$ female mice. It will also be valuable to ascertain the exact mechanism

through which AR signalling activates the VEGF pathway, and whether stimulating this pathway can likewise improve fertility in the context of androgen depletion or deficiency. This

will facilitate an improved diversity of treatment options for cases where androgen supplementation is not desirable (Vegunta et al., 2020).

In summary, using a well validated mouse model we have confirmed that intracrine androgen signalling, stimulated by locally produced DHT, is required for the robust and

timely execution of pathways that stimulate gene expression and changes in cell function essential for a robust decidualisation response. By furthering our understating of the role

androgens play in endometrial receptivity and related endometrial disorders these findings will aid in the development of fertility treatments and other interventions.

Author contributions:

I.W.S designed and performed the experiments, analysed the data, interpreted the results,

538   and wrote the manuscript. P.M.K performed the experiments, analysed the data, and interpreted the results. D.R, R.A, F.L.C & L.B.S performed the experiments. D.L provided

540   study materials. P.T.K.S conceived the study, designed the experiments, interpreted the results and revised the manuscript. D.A.G conceived the study, designed and performed the

542   experiments, analysed the data, interpreted the results, and revised the manuscript.

Acknowledgements.

Funding: MRC programme grants to PTKS (MR/N024524/1, G1002033); MRC Programme Grant to LBS (MR/N002970/1); Wellcome Trust Fellowship to DAG (220656/Z/20/Z); Society

for Endocrinology Early Career Grant to DAG; Scottish Funding Council Research Adaption

Fund to IWS; importation of the mice from Jackson laboratories was paid from Wellcome Trust grant 072217/Z/03/Z. We thank Prof. Ruth Andrew (University of Edinburgh) for founder stocks of Srd5a1$^{-/-}$ mice. We are grateful to Dr Alison Munro and the Host and

Tumour Profiling Unit (Cancer Research UK – Edinburgh Centre) for support for Nanostring analysis, Dr Pamela Brown from Biomolecular core facility and facility staff from the

Bioresearch and Veterinary Services team.

References

       ABDALLA, H. I., BILLETT, A., KAN, A. K. S., BAIG, S., WREN, M., KOREA, L. &
556   STUDD, J. W. W. 1998. Obstetric outcome in 232 ovum donation pregnancies. *BJOG:
           An International Journal of Obstetrics & Gynaecology,* 105**,** 332-337.
558   ASLAN, G., CIMEN, S., YORUKOGLU, K., TUNA, B., SONMEZ, D., MUNGAN, U. &
           CELEBI, I. 2005. Vascular endothelial growth factor expression in untreated and
560   androgen-deprived patients with prostate cancer. *Pathology - Research and Practice,*
           201**,** 593-598.
562   BENJAMINI, Y. & HOCHBERG, Y. 1995. Controlling the False Discovery Rate : A
           Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical*
564   *Society,* 57**,** 289-300.
       CASTRACANE, V. D. & ASCH, R. H. 1995. Pregnancy: Testosterone and androstenedione
566   in premature ovarian failure pregnancies: Evidence for an ovarian source of
           androgens in early pregnancy. *Human Reproduction,* 10**,** 677-680.
568   CLOKE, B., HUHTINEN, K., FUSI, L., KAJIHARA, T., YLIHEIKKILÄ, M., HO, K. K.,
           TEKLENBURG, G., LAVERY, S., JONES, M. C., TREW, G., KIM, J. J., LAM, E.
570   W. F., CARTWRIGHT, J. E., POUTANEN, M. & BROSENS, J. J. 2008. The
           Androgen and Progesterone Receptors Regulate Distinct Gene Networks and Cellular
572   Functions in Decidualizing Endometrium. *Endocrinology,* 149**,** 4462-4462.
       COUSINS, F. L., MURRAY, A., ESNAL, A., GIBSON, D. A., CRITCHLEY, H. O. D. &
574   SAUNDERS, P. T. K. 2014. Evidence from a mouse model that epithelial cell
           migration and mesenchymal-epithelial transition contribute to rapid restoration of
576   uterine tissue integrity during menstruation. *PLoS ONE,* 9**,** e86378-e86378.
       DE VRIES, M. J., DEKKER, G. A. & SCHOEMAKER, J. 1998. Higher risk of preeclampsia
578   in the polycystic ovary syndrome. A case control study. *European Journal of*
           Obstetrics and Gynecology and Reproductive Biology, 76**,** 91-95.
580   DUNK, C., SMITH, S., HAZAN, A., WHITTLE, W. & LEE JONES, R. 2008. Promotion of
           angiogenesis by human endometrial lymphocytes.*Immunological Investigations,* 37**,**
582   583-610.
       DUNN, C. L., KELLY, R. W. & CRITCHLEY, H. O. D. 2003. Decidualization of the human
584   endometrial stromal cell: An enigmatic transformation. *Reproductive BioMedicine*
           Online, 7**,** 151-161.
586   EDGAR, R., DOMRACHEV, M. & LASH, A. E. 2002. Gene Expression Omnibus: NCBI
           gene expression and hybridization array data repository. *Nucleic Acids Res,* 30**,** 207-
588   10.
       GIBSON, D. A., SIMITSIDELLIS, I., COLLINS, F. & SAUNDERS, P. T. K. 2018a.
590   Endometrial Intracrinology: Oestrogens, Androgens and Endometrial Disorders.
           International Journal of Molecular Sciences 2018, Vol. 19, Page 3276, 19**,** 3276-3276.
592   GIBSON, D. A., SIMITSIDELLIS, I., COUSINS, F. L., CRITCHLEY, H. O. D. &
           SAUNDERS, P. T. K. 2016a. Intracrine Androgens Enhance Decidualization and
594   Modulate Expression of Human Endometrial Receptivity Genes. *Scientific Reports,* 6**,**
           27-31.
596   GIBSON, D. A., SIMITSIDELLIS, I., KELEPOURI, O., CRITCHLEY, H. O. D. &
           SAUNDERS, P. T. K. 2018b. Dehydroepiandrosterone enhances decidualization in
598   women of advanced reproductive age. *Fertility and Sterility,* 109**,** 728-734.e2.
       GIBSON, D. A., SIMITSIDELLIS, I. & SAUNDERS, P. T. K. 2016b. Regulation of
600   androgen action during establishment of pregnancy. *Journal of Molecular*
           Endocrinology, 57**,** R35-R47.

602   JOSEPH, I. B. J. K., NELSON, J. B., DENMEADE, S. R. & ISAACS, J. T. 1997. Androgens regulate vascular endothelial growth factor content in normal and malignant prostatic
604   tissue. *Clinical Cancer Research,* 3**,** 2507-2511.

KAHVECI, B., MELEKOGLU, R., EVRUKE, I. C. & CETIN, C. 2018. The effect of
advanced maternal age on perinatal outcomes in nulliparous singleton pregnancies. *BMC Pregnancy and Childbirth,* 18**,** 343.

KHORRAM, O., GARTHWAITE, M. & MAGNESS,R. R. 1999. Endometrial and Myometrial Expression of Nitric Oxide Synthase Isoforms in Pre- and
Postmenopausal Women. *The Journal of Clinical Endocrinology & Metabolism,* 84**,** 2226–2232.

KIRKWOOD, P. M., GIBSON, D. A., SMITH, J. R., WILSON-KANAMORI, J. R., KELEPOURI, O., ESNAL-ZUFIAURRE, A., DOBIE, R., HENDERSON, N. C. &
SAUNDERS, P. T. K. 2021. Single-cell RNA sequencing redefines the mesenchymal cell landscape of mouse endometrium. *FASEB Journal,* 35**,** e21285-e21285.

KUSAMA, K., YOSHIE, M., TAMURA, K., IMAKAWA, K., ISAKA, K. & TACHIKAWA, E. 2015. Regulatory Action of Calcium Ion on Cyclic AMP-Enhanced Expression of
Implantation-Related Factors in Human Endometrial Cells. *PloS one,* 10.**,** e0132017.

LABRIE, F., BÉLANGER, A., CUSAN, L., GOMEZ, J. L. & CANDAS, B. 1997. Marked
620   decline in serum concentrations of adrenal C19 sex steroid precursors and conjugated androgen metabolites during aging. *J Clin Endocrinol Metab,* 82**,** 2396-402.

622   LABRIE, F., MARTEL, C., BÉLANGER, A. & PELLETIER, G. 2017. Androgens in women are essentially made from DHEA in each peripheral tissue according to intracrinology.
624   *J Steroid Biochem Mol Biol,* 168**,** 9-18.

LIVINGSTONE, D. E. W., BARAT, P., DI ROLLO, E. M., REES, G. A., WELDIN, B. A.,
626   ROG-ZIELINSKA, E. A., MACFARLANE, D. P., WALKER, B. R. & ANDREW, R. 2015. 5α-Reductase type 1 deficiency or inhibition predisposes to insulin resistance,
628   hepatic steatosis, and liver fibrosis in rodents. *Diabetes,* 64**,** 447-458.

MAHENDROO, M. S., CALA, K. M., LANDRUM, C. P. & RUSSELL, D. W. 1997. Fetal
630   Death in Mice Lacking 5α-Reductase Type 1 Caused by Estrogen Excess. *Molecular Endocrinology,* 11**,** 917-927.

632   MAHENDROO, M. S., CALA, K. M. & RUSSELL, D. W. 1996. 5 alpha-reduced androgens play a key role in murine parturition. *Molecular Endocrinology,* 10**,** 380-392.

634   MAHENDROO, M. S., PORTER, A., RUSSELL, D. W. & WORD, R. A. 1999. The parturition defect in steroid 5α-reductase type 1 knockout mice is due to impaired
636   cervical ripening. *Molecular Endocrinology,* 13**,** 981-992.

MAHENDROO, M. S. & RUSSELL, D. W. 1999. Male and female isoenzymes of steroid
638   5α-reductase. *Reviews of Reproduction,* 4**,** 179-183.

MARSHALL, E., LOWREY, J., MACPHERSON, S., MAYBIN, J. A., COLLINS, F.,
640   CRITCHLEY, H. O. D. & SAUNDERS, P. T. K. 2011. In silico analysis identifies a novel role for androgens in the regulation of human endometrial apoptosis. *Journal of*
642   *Clinical Endocrinology and Metabolism,* 96**,** E1746-E1755.

NARUKAWA, S., KANZAKI, H., INOUE, T., IMAI, K., HIGUCHI, T., HATAYAMA, H.,
644   KARIYA, M. & MORI, T. 1994. Androgens induce prolactin production by human endometrial stromal cells in vitro. *The Journal of Clinical Endocrinology &*
646   *Metabolism,* 78**,** 165-168.

ODIBO, A. O., NELSON, D., STAMILIO, D. M., SEHDEV, H. M. & MACONES, G. A.
648   2006. Advanced maternal age is an independent risk factor for intrauterine growth restriction. *Am J Perinatol,* 23**,** 325-8.

650   PLAISIER, M. 2011. Decidualisation and angiogenesis. *Best Practice and Research: Clinical Obstetrics and Gynaecology,* 25**,** 259-271.

652   REBOURCET, D., WU, J., CRUICKSHANKS, L., SMITH, S. E., MILNE, L., FERNANDO, A., WALLACE, R. J., GRAY, C. D., HADOKE, P. W., MITCHELL, R. T.,
654   O'SHAUGHNESSY, P. J. & SMITH, L. B. 2016. Sertoli Cells Modulate Testicular Vascular Network Development, Structure, and Function to Influence Circulating
656   Testosterone Concentrations in Adult Male Mice. *Endocrinology,* 157**,** 2479-88.
        RITTMASTER, R. S., THOMPSON, D. L., LISTWAK, S. & LORIAUX, D. L. 1988.
658   Androstanediol Glucuronide Isomers in Normal Men and Women and in Men Infused with Labeled Dihydrotestosterone. *The Journal of Clinical Endocrinology &*
660   *Metabolism,* 66**,** 212-216.
        ROTHMAN, M. S., CARLSON, N. E., XU, M., WANG, C., SWERDLOFF, R., LEE, P.,
GOH, V. H., RIDGWAY, E. C. & WIERMAN, M. E. 2011. Reexamination of testosterone, dihydrotestosterone, estradiol and estrone levels across the menstrual
cycle and in postmenopausal women measured by liquid chromatography-tandem mass spectrometry. *Steroids,* 76**,** 177-82.
SIMITSIDELLIS, I., ESNAL-ZUFFIAURE, A., KELEPOURI, O., O'FLAHERTY, E., GIBSON, D. A. & SAUNDERS, P. T. K. 2019. Selective androgen receptor
modulators (SARMs) have specific impacts on the mouse uterus. *Journal of Endocrinology,* 242**,** 227-239.
SIMITSIDELLIS, I., GIBSON, D. A., COUSINS, F. L., ESNAL-ZUFIAURRE, A. & SAUNDERS, P. T. K. 2016. A Role for Androgens in Epithelial Proliferation and
Formation of Glands in the Mouse Uterus. *Endocrinology,* 157**,** 2116-2128.
        SIMITSIDELLIS, I., SAUNDERS, P. T. K. & GIBSON, D. A. 2018. Androgens and
674   endometrium: New insights and new targets. *Molecular and Cellular Endocrinology,* 465**,** 48-60.
676   SORDELLO, S., BERTRAND, N. & PLOUËT, J. 1998. Vascular Endothelial Growth Factor Is Up-Regulatedin Vitroandin Vivoby Androgens. *Biochemical and Biophysical*
678   *Research Communications,* 251**,** 287-290.
        THIGPEN, A. E. & RUSSELL, D. W. 1992. Four-amino acid segment in steroid 5α-
680   reductase 1 confers sensitivity to finasteride, a competitive inhibitor. *Journal of Biological Chemistry,* 267**,** 8577-8583.
682   UPRETI, R., NAREDO, G., FAQEHI, A. M. M., HUGHES, K. A., STEWART, L. H., WALKER, B. R., HOMER, N. Z. M. & ANDREW, R. 2015. Simultaneous
684   pharmacokinetic and pharmacodynamic analysis of 5α-reductase inhibitors and androgens by liquid chromatography tandem mass spectrometry. *Talanta,* 131**,** 728-
686   735.
        VEGUNTA, S., KLING, J. M. & KAPOOR, E. 2020. Androgen Therapy in Women. *Journal*
688   *of Women's Health,* 29**,** 57-64.
        VON DEUTSCH, D. A., ABUKHALAF, I. K. & LAPU-BULA, R. 2012. "Anabolic Doping
690   Agents" in Mozayani A. & Raymon L. (eds.) *Handbook of Drug Interactions: A Clinical and Forensic Guide***,**. Humana Press, pp. 625-754.
692   WALTERS, K. A., SIMANAINEN, U. & GIBSON, D. A. 2016. Androgen action in female reproductive physiology. *Current Opinion in Endocrinology, Diabetes and Obesity,*
694   23**,** 291-296.
        YANG, F., ZHENG, Q. & JIN, L. 2019. Dynamic Function and Composition Changes of
696   Immune Cells During Normal and Pathological Pregnancy at the Maternal-Fetal Interface.
698   ZHANG, X. & CROY, B. A. 1996. Maintenance of Decidual Cell Reaction by Androgens in the Mouse1. *Biology of Reproduction,* 55**,** 519-524.

700   ZUMOFF, B., STRAIN, G. W., MILLER, L. K. & ROSNER, W. 1995. Twenty-four-hour
             mean plasma testosterone concentration declines with age in normal premenopausal
702   women. *J Clin Endocrinol Metab,* 80**,** 1429-30.

704   **Tables and Figures**

Ratio Analysis	Differentially expressed genes (DE)			log$_2$FC of DE genes		
	(#)	(% total)	Upregulated (% DE)	Downregulated (Mean ± 95% CI)	Upregulated (Mean ± 95% CI)	Absolute (Mean ± 95% CI)
Srd5a1$^{-/-}$ Non Vs WT Non	0	0	NA	NA	NA	NA
Srd5a1$^{-/-}$ Dec Vs WT Dec	254	33.0	79.5	-0.84 (-0.71–-0.97)	1.56 (1.39–1.74)	1.41 (1.27–1.56)
WT Dec Vs WT Non	507	65.8	29.0	-1.68 (-1.54–-1.82)	1.94 (1.67–2.20)	1.75 (1.63–1.88)
Srd5a1$^{-/-}$ Dec Vs Srd5a1$^{-/-}$ Non	299	38.8	25.4	-0.84 (-0.76–-0.91)	1.51 (1.27–1.76)	1.01 (0.92–1.10)
2-Way ANOVA Analysis						
Decidualisation	464	61.9				
Genotype	27	3.6				
Interaction	33	4.4				

706   *Table 1: Summary of differentially expressed genes using ratio analysis and 2-Way ANOVA analysis of Pancancer Pathways Nanostring panel.*

708   Abbreviations: Non, non-decidualised uterus; Dec, decidualised uterus; DE, differentially expressed; FC, fold-change; CI, confidence interval.

Figure 1. Decidualisation response is impaired in Srd5a1^{-/-} mice.

712 A) Schema for the decidualisation induction model. E2 = oestradiol. P4 = progesterone. B)

Graphic depicting a uterine horn in various stages of decidualisation. The horn is thicker

714 where a decidualisation reaction has occurred. A horn is deemed fully decidualised if an

uninterrupted stretch of decidualisation has occurred over at least half of the horn length.

716 Partial decidualisation is any decidualisation response less than this, often occurring as

disparate nodes along the horn. C) Representative images of uterine horns from wild type

718 (WT) and Srd5a1^{-/-} animals, with the classification of each horn given; p = partial

decidualisation, f = fully decidualised. D) Contingency table depicting proportion of horn

720   responses in each genotype. E,F) Wet weight of uterine horns in WT and Srd5a1$^{-/-}$ mice at

time of tissue recovery (4 days following decidualisation induction). Non-decidualised horns

722   have been pooled. E compares all horns with a decidualisation reaction (Full & Partial),

whereas F shows analysis of those with a partial reaction. G) Expression of *Prl8* via qPCR in

724   uterine horns of WT and Srd5a1$^{-/-}$ mice following decidualisation induction. Plots in D

analysed by chi-squared test for trend; E, F analysed by one-way ANOVA with Newman-

726   Keuls *post-hoc* tests; and G analysed by two-way ANOVA with Bonferroni post-hoc tests; ***

$p < 0.001$; ** $p < 0.01$; * $p < 0.05$.

Figure 2. Evidence of a vascular phenotype in Srd5a1^{-/-} mice following induction of decidualisation.

A) Representative images of CD31[+] cells in WT and Srd5a1[-/-] decidua. Zoomed-in views are shown from regions indicated by boxes. Scale bars are 100µm and 50µm respectively. B) Quantification of vessel density in in WT and Srd5a1[-/-] non-decidualised endometrium and

decidua. C, D) Images of decidualised uterine horns following intravenous injection of Evans

736   Blue and respective quantification of absorbance in uterine tissue extracts in decidualised

horns of WT and $Srd5a1^{-/-}$ (C) or vehicle control (VC) and finasteride (Fin) treated mice (D).

738   E,F). Expression of angiogenic genes via qPCR in decidualised tissue (partial and full) from

WT and $Srd5a1^{-}$/- mice (E), or VC and Fin mice (F). Plots in B analysed by two-way ANOVA

740   with Bonferroni *post-hoc tests;* C,D,E,F analysed by two-tailed t test. ** $p < 0.01$; * $p < 0.05$.

 Figure 3. Effect of 5α-reductase deficiency on global gene expression during decidualisation.

744 A) Unsupervised clustering and heatmap of WT and Srd5a1[-/-] decidualised and non-decidualised tissue RNA analysed with Nanostring Pancancer Pathways gene expression

746 panel. The major gene expression differences occur with decidualisation. B) Volcano plots for the pairwise comparisons between experimental groups, as indicated. Significantly

748 differentially expressed (DE) genes at a false discovery rate (FDR) of 0.05 are coloured pink. C) Selection of genes changing significantly with either genotype, or interaction of genotype

750 and decidualisation, by two-way ANOVA (FDR = 0.05). Count values in the Nanostring data

and respective validation by qPCR in groups with additional samples is shown. Plots in C

752   analysed by two-way ANOVA with Bonferroni *post-hoc* tests. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Figure 4. Gene networks associated with defective decidualisation in Srd5a1^{-/-} mice.

756 A) Biological process gene ontogeny (GO) of genes significantly downregulated in Srd5a1$^{-/-}$ vs WT decidualised uterine horns. All significant terms shown. B) Category netplot (cnet)

758 embedding indicating genes shared between GO terms for genes significantly downregulated in Srd5a1$^{-/-}$ vs WT decidualised uterine horns. C) Heatmap showing

760 expression of top DE genes from the Srd5a1$^{-/-}$ compared to WT decidualised uterine horns and their relative abundance in uterine cell subsets; epithelial, fibroblast and perivascular, as

762 derived from annotation in previously published scRNAseq dataset of mouse uteri. Genes

appearing in the cnet plot in B are labelled and colour coded based on whether they link to

764   angiogenesis-related GO terms (green) or not (black).

Figure 5. Impact of DHT administration on decidualisation response in Srd5a1$^{-/-}$ mice.

A) Schema of decidualisation induction experiment in WT and Srd5a1$^{-/-}$ mice with addition of DHT or vehicle control (VC) at point of decidualisation induction. B) Contingency table depicting proportion of horn responses in WT and Srd5a1$^{-/-}$ mice treated with DHT or VC. C) Wet weight of uterine horns. D) Quantification of Evans Blue dye absorbance in uterine tissue extracts. E) Expression of selected genes analysed by qPCR. Plot in B analysed by chi-squared test; C,D,E analysed by one-way ANOVA and Neuman-Keuls post-hoc tests. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; ns = no significant differences detected.