The optimal clutch size revisited: separating individual quality from

the parental survival costs of reproduction

Lucy A. Winder[1], Mirre J.P. Simons[1], Terry Burke[1]

[1]Ecology & Evolutionary Biology, School of Biosciences, The University of Sheffield,

Sheffield S10 2TN, UK

Abstract

Life-history theory, central to our understanding of diversity in morphology, behaviour and

senescence, describes how traits evolve through the optimisation of trade-offs in investment.

Despite considerable study, there is only minimal support for trade-offs within species

between the two traits most closely linked to fitness – reproductive effort and survival –

questioning the theory's general validity. We used a meta-analysis to separate the effects of

individual quality (positive survival/reproduction correlation) from the costs of reproduction

(negative survival/reproduction correlation) using studies of reproductive effort and parental

survival in birds. Experimental enlargement of brood size caused reduced parental survival.

However, the effect size of brood size manipulation was small and opposite to the effect of

phenotypic quality, as we found that individuals that naturally produced larger clutches also

survived better. The opposite effects on parental survival in experimental and observational

studies of reproductive effort provides the first meta-analytic evidence for theory suggesting

that quality differences mask trade-offs. Fitness projections using the overall effect size

revealed that reproduction presented negligible costs, except when reproductive effort was

forced beyond the level observed within species, to that seen between species. We

conclude that there is little support for the most fundamental life-history trade-off, between

reproductive effort and survival, operating within a population. We suggest that within

species, the fitness landscape of the reproduction–survival trade-off is flat until it reaches the

boundaries of the between-species fast–slow life-history continuum. Our results provide a

quantitative explanation as to why the costs of reproduction are not apparent and why

variation in reproductive effort persists within species.

Main

Across taxa, we see wide variation in life-history traits, such as the number of offspring

produced and time spent raising young [1–3]. The central idea in life-history theory is that

resources are finite, forcing trade-offs, meaning that investment in one aspect of life requires

the sacrifice of another [4–7]. As reproduction is one of the most resource-demanding life

stages, it is expected that within-species variation in offspring production will be constrained

by the cost of producing and raising young. It is thought that the fitness costs of reproduction

are largely incurred as a decrement to parental survival, explaining why a species evolved a

certain amount of reproductive effort, whilst also explaining the fast–slow life-history

continuum between reproduction and lifespan observed across species [6]. As reproduction

and survival are the two components of life-history most closely related to fitness, this central

trade-off has been the subject of much theoretical and empirical research, branching fields

ranging from ecology, evolutionary biology to ageing research [8–10].

Brood size manipulations of birds in natural conditions have provided arguably the best

experimental paradigm in which to test the survival costs of reproduction, as they directly

alter parental care investment. Experimental increases in brood size result in increased

parental effort, suggesting that parents can typically cope with increased reproductive

demands [11–15]. However, the expected increased costs of reproductive effort are not always

detected and the current estimate of the cost to reproduce across studies suggests only a

small and inconsistent effect [16–19]. The absence of a cost of reproduction (determined

through experimental studies) on survival means that costs must arise elsewhere or,

alternatively, that individuals may differ in quality (determined through observational studies).

By quality, we refer to individuals operating at their own maximum reproductive output,

determined by their phenotypic condition, local or temporal genetic adaptation, and the

surrounding environment, and should manifest as a positive correlation between two fitness-

associated traits (such as reproduction and survival) [11,19–22]. The relative importance of the

trade-off between reproductive effort and survival to life-history theory and the biology of

ageing [9], therefore remains unclear. In addition, the compelling theoretical explanation for

the lack of an apparent trade-off due to the confounding effects of individual quality has not

been investigated on a quantitative level [5,19,23,24]. Our study aims to separate the effects of a

trade-off between reproductive effort and parental survival and the effects of individual

quality by comparing studies of brood manipulations and naturally varying clutch size.

Here, we present a meta-analysis that distinguishes between individual quality effects

(naturally varying clutch size) and the costs of reproduction (brood manipulation studies). To

do this, we tested how parental annual survival in birds is affected by the clutch size (note

brood size and clutch size are used interchangeably throughout this manuscript) they cared

for in two different contexts: first in brood manipulation studies and, second, in observational

studies of natural variation in clutch size. We expressed changes in survival on a per-egg

basis, which, for the first time, allows for a quantitative comparison across studies and thus

species. We find the effect of quality is associated with higher survival chances given an

increase in reproductive effort, and that this effect is opposite but equal in magnitude to the

costs of reproduction. The survival trade-off for offspring production within a population is

therefore offset by differences in quality, potentially constraining the evolution of higher

reproductive effort. Our analysis also uniquely allowed a quantitative across species

comparison, as survival risk was expressed on a per-egg basis. We transformed the

response variable, scaling for variance and mean, given that a per-egg increase in clutch

size does not equate to the same proportional increase in reproductive effort for all species

equally. Our findings suggest that species that generally lay smaller clutches are affected

more severely by brood-size manipulations. This provides evidence that trade-offs are only

detected when an individual is forced to perform far outside its optimum level (Figure S1). Species who on average have smaller clutch sizes receive disproportionate manipulations (Figure S1A) and this pushes species beyond their natural range (Figure S1B).

To predict the evolutionary consequences of the effect sizes that we estimated using the meta-analysis, we projected the fitness consequences for a change in clutch-size life-history strategy. We found that the costs to parental survival as estimated from brood size manipulation studies translate into negligible fitness costs, with a relatively flat fitness landscape, suggesting that birds underproduce in terms of clutch size, given the absence of fitness costs. This conclusion fits with our comparative finding that suggests that only brood sizes manipulated beyond the natural range incur substantial survival costs. Our results therefore suggest that, in wild populations, parental survival costs are, at most, a small component of the total fitness costs of investing parental effort. Our results do suggest that a cost of reproduction can be detected when an organism is pushed to the extreme of its reproductive capacity. We therefore infer that, though the survival–parental care trade-off does exist within species, it not strong enough to constrain clutch size and can therefore not explain clutch size evolution and thus variation within species. In addition, our work shows that differences in individual quality counterbalance the trade-off between survival and reproduction, as previously theorised [5], and as such could constrain reproductive effort further, and contribute to the maintenance of clutch size variation in a population.

Results

The relationship between clutch size and survival was significantly different and opposite between observational and brood manipulation studies, irrespective of how clutch size was scaled ($p < 0.01$, Figure 1, Table 1). Within natural variation (observational studies), parents

with larger clutches showed increased survival (or parents with smaller clutches showed

reduced survival). In contrast, when broods were experimentally manipulated, the opposite

relationship was found; increasing brood sizes decreased survival. Although the difference in

overall effect size between experimental and natural variation in clutch size was strongly

significant in each comparison made, the individual overall effect sizes only became

significant (from zero) when clutch size was expressed as a proportional increase.

Expressing clutch size as a proportional increase corrects for the variation in average clutch

size observed across the species included in this analysis, which ranged from 2 to 11. The

parental effort required to raise two instead of one chick is potentially doubled, whereas one

additional chick in a brood of 11 is likely to require only a marginal increase in effort. Indeed,

also when using a between-species comparison, the effects of clutch size manipulation and

quality were strongest in the species that laid the smallest clutches, suggesting that costs to

survival were only observed when a species was pushed beyond its natural limits (Figure 2,

Figure S1A, Table S1).

Figure 1: The effects size (log odds of survival given an increase in clutch size) for three different

measures of clutch size: (A) raw, (B) standardised, and (C) proportional clutch size. Coloured points

are the combined effect sizes of the odds ratios with their 95% confidence intervals. Points are

coloured by whether they represent brood manipulation experiments (costs of reproduction) or they

are observational (quality). Grey points are the odds ratios of each study, with their sizes weighted

by the sampling variance.

139 **Table 1:** Effect size estimates for the odds of survival with increasing clutch size (raw, standardised

140 and proportional clutch size). The p-values indicate the difference between brood manipulations and

141 observational data, with the individual effect *p*-values (from zero) in parentheses.

	Parameter		Effect size	95% CI lower bounds	95% CI upper bounds	*p*	(individual)
Raw	Clutch size	Brood manipulation	-0.0522	-0.1406	0.0363	**0.0007**	(0.2477)
		Observational	0.0747	0.1571	-0.0288		(0.1571)
Standardised	Clutch size	Brood manipulation	-0.0651	-0.1478	0.0177	**0.0065**	(0.1232)
		Observational	0.1143	-0.0046	0.2333		(0.0595)
Proportional	Clutch size	Brood manipulation	-0.2703	-0.4984	-0.0423	**0.0005**	**(0.0202)**
		Observational	0.3850	0.0583	0.7116		**(0.0209)**

Model = ~obervational_or_experimental, random = (species, phylogeny, study reference)

Figure 2: The meta-analytic linear regression (Table S1) of the effect size for increasing clutch size

(per egg) on parental survival, given the average clutch size for the species. Species with small clutch

sizes showed stronger costs of reproduction and a stronger relationship with quality (Interaction

between treatment and species clutch size effect size = -0.036, p = 0.015). The points are the survival

effect sizes (log odds ratio) per egg (as in Figure 1A) on parental survival in each study, with the point

size reflecting the meta-analytic weight of that study.

Males and females did not differ in their survival response to changing clutch size (Table S2,

Figure S2, contrary to Santos & Nakagawa 2012 [18]). The variance assigned to the random

effects in the model was largely accounted for by study (Table S3). Species accounted for

more variation than the phylogeny, indicating that species vary in the relationship between

survival and reproductive effort, irrespective of their shared evolutionary ancestry. However,

our dataset included few closely related species, which reduces our ability to estimate

phylogenetic effects (Figure 3).

Figure 3: Phylogenetic tree of species included in our meta-analysis. Numeric labels denote the node

support (as a percentage). Coloured points indicate whether the species were included in the

experimental or observational studies.

Projected fitness consequences of the costs of reproduction

From our meta-analysis we now have a quantifiable and comparable effect size for the

survival costs of reproduction that we can use to predict the evolutionary consequences of

individuals in a given population increasing their clutch size across a range of life histories.

To this end, we projected the fitness consequences of increased reproductive effort, starting

with the average effect size estimate per egg (-0.05, Table 1) across a range of life histories,

for a range of annual survival rates and clutch sizes (Figure 4). Overall, the effect size

estimated in the meta-analysis (-0.05) resulted in a gain of fitness when reproductive output

increased, especially in hypothetical species with low survival and small clutches. If we used

a more severe effect size (-0.15, the lower confidence interval of our meta-analysis), this led

to a reduction in fitness in almost all cases. Conversely, the benefit of higher reproductive

output was largely offset by the cost of survival when a species' survival rate and clutch size

were high. When we increased the effect size up to five-fold, fitness costs of reproduction

became more pronounced, but were still not present in species with small clutches and short

lifespans.

Under long-term evolution these selection differentials should lead to individual hypothetical

species moving towards the diagonal (bottom left to the right top corner). This diagonal

represents the observed fast–slow pace of life continuum observed among species [25].

Exemplar species (i.e., with survival and average clutch size combinations observable in wild

populations), for which we predicted the fitness consequences of the costs of reproduction,

lie on this comparative diagonal in life history. In these exemplar species, the selection

differential was observed to lie slightly above one, indicating that individuals having a higher

clutch size than the species' average would gain a slight fitness benefit. The fitness costs

and benefits did, in general, not diverge substantially with the addition of chicks, but

flattened, suggesting that the costs of survival counterbalance the benefits of reproduction

across a range of reproductive outputs within a species.

The low costs of reproduction that we estimated could still be responsible for between-

species life-history evolution, constraining species reproductive output and survival

combination to fall along the diagonal of the fast–slow pace of life continuum. How selection

pressures translate into short term and longer-term evolutionary trajectories is uncertain.

Often directional selection estimated in the wild does not translate to the inter-generational

change on the population level [26]. Note, however, that only far away from the diagonal did

our fitness projections reach a magnitude that would be predicted to lead to rapid

evolutionary change [27] and see SI). The weak selection effects that lie on the diagonal are

probably to be counterbalanced in the wild by factors such as environmental effects and

genetic effects (e.g., gene flow from immigration or random mutation) [28]. We argue that

within-species the minimal costs of reproduction, a flat fitness landscape and quality effects

(Figure 1) together explain why individuals appear to under-produce. Only when individuals

are pushed beyond the observed between-species constraint do costs become apparent

(Figures 1, 2).

Our interpretation of the reproduction/lifespan life-history trade-off, based on our quantitative

meta-analysis and subsequent fitness projections, explains several key observations and

contradictions in the field. A strong trade-off is observed between species, but within species

this trade-off is not apparent and variation in reproductive output is maintained within fitness

boundaries similar to those that determine the between-species life history trade-off. The

implication of this conclusion is that the costs of reproduction are likely to operate on a

physiological level, but that the fitness consequences will remain largely flat over a species'

observed variation in reproductive output. These effects are further obscured by the effects

of quality (observational studies), which are opposite in sign and magnitude to the cost of

reproduction (<mark>experimental studies)</mark> (Figure 1) and are likely to further flatten the fitness

landscape.

Figure 4: Isoclines of selection differentials among hypothetical control populations (in which

individuals reproduce at the species' mean rate) and hypothetical brood-manipulated populations

(where individuals reproduce at an increased rate compared to control) for their whole lives.

Selection differentials (i.e., the difference in lifetime reproductive output between hypothetical

control and brood-manipulated populations) above 1 represents high lifetime fitness. Survival rates,

clutch sizes, the magnitude of the manipulation (chicks added) and effect sizes represent the range

of these variables present in the studies used in our meta-analysis. For each clutch size, we used a

predicted survival rate and effect size to give isoclines that are biologically meaningful (exemplar

birds shown in red). Arrows indicate the relative size and direction of selection in life-history space

(on the reproduction axis). The costs of reproduction we estimated within species are predicted to

result in a fast–slow life-history continuum across species, and the exemplar species we used as

examples fit on this diagonal of survival rate/ clutch size combinations. We suggest that individual

species show limited costs of reproduction, as they operate within relatively wide constraints

imposed by the cost of reproduction that is responsible for the strong life-history trade-off observed

across species.

Discussion

Our results provide the first meta-analytic evidence that differences in individual quality drive

variation in clutch size and that parental survival costs do not constrain within-species

variation in reproductive effort (Figure 5). Here, we use the definition of quality as a

combination of traits that give an individual higher fitness [5]. The finding of individual quality

being a driver of variation in clutch size is not necessarily a surprising one, even though this

contradicts the general theory that trade-offs drive variation in clutch size, as among-

individual heterogeneity is also a well-established effect across populations in the field of

ecology and evolution (such as age-specific reproductive performance) [5,24]. Furthermore, a

recent meta-analysis on genetic life-history trade-offs found several positive genetic

correlations between survival and other life-history traits, which is in conflict with life-history

trade-off theory [29]. Brood manipulations often fail to take into account that variation in

resource acquisition may be larger than variation in resource allocation, and though

intuitively manipulation of effort tests variation in resource allocation, it fails to account for the

baseline variation in reproductive performance [5]. This presents a fundamental flaw which our

results demonstrate – variation in resource allocation can be present and testable through

experiments (i.e., brood manipulations) but may only account for a small proportion of

variation observed in reproductive output. Perhaps fitting with this interpretation or with

additional costs from recurring bouts of enhanced reproductive effort, Jackdaws (*Corvus*

monedula) show costs of increased brood size only when applied sequentially across years,

but note that for this study there is no control group that is only manipulated in single years

[30].

Figure 5: Decision tree representing the logical steps from our original hypothesis to our overall

interpretation of our findings.

The reason selection is not acting on "high quality" individuals is currently unknown, but it is

likely that environmental variability leads to alternative phenotypes being selected for at

different points in space and time (also discussed in Pujol *et al.,* 2018 [26]). It is also possible

that the quality effect could represent a terminal effect, where individuals have lower

reproductive output in the year preceding their death, thereby driving the trend for naturally

lower laying birds to have lower survival when estimated between individuals (e.g., Coulson

& Fairweather, 2001; Rattiste, 2004; Simons *et al.,* 2016 [31–33]; also see Hammers *et al.,* 2012

[34] for age-related changes in reproductive output). The effect on parental annual survival of

having naturally larger clutches was significantly opposite to the result of increasing clutch

size through brood manipulation, and quantitatively similar. Parents with naturally larger

clutches are thus expected to live longer and this counterbalances the "cost of reproduction"

when their brood size is experimentally manipulated. It is, therefore, possible that quality

effects mask trade-offs. Furthermore, it could be possible that individuals that lay larger

clutches have smaller costs of reproduction, i.e. would respond less in terms of annual

survival to a brood size manipulation, but with our current dataset we cannot address this

hypothesis (Figure 5). The effect of a change in clutch size on parental survival may also be

non-linear, but it remains to be determined what shape, if not linear, the relationship would

be. Although these possible non-linear effects warrants investigation, these relationships

likely differ between species and inclusion in our current work here could lead to spurious

relationships being reported.

For both costs of reproduction and quality effects, we found that species that laid the

smallest clutches showed the largest effects. Brood manipulations that affect parental

survival are thus likely to be the result of pushing parental effort beyond its natural optima

(Figure S1) and so arguably that brood manipulations are not necessarily a good test of

whether trade-offs happen in the wild. The classic trade-off between adult survival and the

clutch size cared for is only apparent when an individual is forced to raise a clutch outside of

its individual optimum, and these effects are confounded or even fully counterbalanced by

differences in quality (as theorised in van Noordwijk & de Jong, 1986 [5]).

Our fitness projections of the consequences of the costs of reproduction using the overall

effect size we estimated suggest that, for current extant species, the within-species fitness

landscape of the reproduction survival trade-off is flat. Species' life history decisions are

constrained within a broader fast–slow life-history continuum, explaining why variation within

species in reproductive effort, such as in clutch size, is large and near universal. Of course,

this analysis does not fully explore more complex variation in clutch size observed in the

wild, such as that observed between temperate and tropical reproductive strategies. Such

work would provide useful in understanding why variation in clutch size is maintained across

species with different life-history strategies. Our results do, however, provide a key step in

understanding the generalised relationship between reproductive effort and survival across

species (Figure 5). Our interpretation also assumes that other fitness costs of reproduction

are smaller or at least less relevant than survival costs. However, it is possible that such

costs are actually more important, and it should be noted that effects such as those on

offspring quality (e.g., Conrad & Robertson, 1992 and Smith *et al.*, 1989 [14,35]), parental

condition other than survival (e.g., Reid, 1987 and Kalmback *et al.*, 2004 [36,37]) and future

reproductive effort (e.g., Järvistö *et al.*, 2016 [38]) have been observed (also see Figure 5).

However, the importance of these effects is likely to vary considerably in different species.

Using offspring quality as an example, some species produce sacrificial offspring, others

experience catch-up growth, meaning that though the effect of increasing offspring number

on offspring quality is important for certain species, drawing generalised across-species

conclusions is unlikely to be possible. Conversely, the survival cost to reproduce is thought

to be universal across both bird species and across taxa, and it is perhaps for this reason

that the reproduction-survival trade-off has been considered to contribute more to variation in

reproductive effort than other trade-offs. Our work refutes this idea directly. Indeed, the few

studies that have measured the different domains that contribute to fitness in brood-size

manipulation studies concluded that only in combination do these costs result in balancing

selection for the current most common brood size in the population [39–41]. Such classic trade-

off explanations do, however, fail to explain why variation in reproductive effort is prevalent

within species and why between-species life-history trade-offs appear so much stronger and

conclusive. Our analysis and interpretation suggest that, at its optimum, the within-species

trade-off between survival and reproduction is relatively flat, and thus neutral to selection

(supporting the theory presented in Cohen et al., 2020 [19]). We suggest that the lack of

evidence supporting trade-offs driving within-species variation does not necessarily mean

that physiological costs of reproduction are non-existent (e.g., Smith et al., 1989 and

Lemaître et al., 2015 [35,42]), but rather that, within the wild and within the natural range of

reproductive activities, such costs are not relevant to fitness. One key explanation for this

effect supported by our meta-analysis (Figure 5) and prior theory [5], is that individuals differ

in quality.

Methods

All data and code used in this study can be found at:

https://doi.org/10.5061/dryad.q83bk3jnk.

Study sourcing & inclusion criteria

We extracted studies of parental survival to the following year given clutch size raised using

the following inclusion criteria (similar to Santos & Nakagawa, 2012 [18]): the study must be on

a wild population; the study must contain variation in the number of offspring

produced/raised (hereafter referred to as clutch size for simplicity), the study must report

variation in clutch size in relation to parental survival to the following year (including both

experimental and observational studies) and must provide sample sizes. We did include

studies where parental survival was reported for both parents combined as opposed to

Santos & Nakagawa (2012) who required male and female data to be reported separately.

Excluded studies and the grounds for their removal are given in the supplementary

information (Table S4). We started by, first, extracting data (clutch size raised after

manipulation and associated parental survival to the next year) from the included brood-

manipulation studies from Santos & Nakagawa (2012) and then searched the literature to

include more recently-published studies (Supplementary Methods), and in this search also

added studies Santos & Nakagawa missed. In addition to brood manipulation studies, we

extracted data from studies that correlated variation in parental survival with natural variation

in clutch size (observational studies). For each species used in the brood manipulation

studies, we aimed to find the same species for the observational studies to ensure that the

effects of quality were estimated across a similar range of species. The reason for this is to

ensure the experimental and observational datasets are comparable in terms of species,

which could bias results if not considered, and so facilitate a more direct comparison

between the quality effect (observational studies) and the trade-off (experimental studies).

Where there was no equivalent study in the same species, we attempted to find a study of a

congener. In most cases, observational data were obtained from either the same paper as

the one describing brood manipulations (11 studies), or via searching for other papers by the

same authors (2 studies). If this failed to produce observational data, a search was

conducted following the same protocol as for the brood-manipulation experiments, but also

specifying species, genus and/or common name in the search (7 studies). Of the 28 species

used in the brood manipulation studies, we were able to find 10 of the same species in

observational studies. We were able to find a further 6 species which were congeners for

observational studies.

From the literature search, 78 individual effect sizes from 30 species and 46 papers were

used (20 observational and 58 experimental studies). While extracting these studies we also

made note of the average clutch size of the species and the within-species standard

deviation in clutch size. We extracted this information from the paper containing the study

but if the information was missing, we searched other published literature with the aim to find

the information from a similar population (i.e., at a similar latitude).

Extracting effect sizes

We used the absolute value of offspring in the nest (for experimental studies this is the

number of offspring after the manipulation occurred) and associated parental survival to

estimate an effect size by performing a logistic regression to obtain the log odds ratio for

parental survival, given the clutch size (i.e., positive effect sizes indicate an increased

chance of survival). For example, a bird who laid a clutch size of 5 but was manipulated to

have -1 chick was recorded to have a clutch size of 4. If the manipulation was reported but

absolute value of offspring produced was not, we used clutch size to be the species average

+/- the number of offspring added or removed. Parental survival was modelled as a binary

response variable for the number of birds who survived and who died after raising a given

clutch size. Clutch size was averaged (mean) if a single estimate of survival was reported for

multiple clutch sizes. 'Year' was included as an explanatory variable to correct for between-

year variation in adult survival, where data were presented for multiple years. We

standardised the clutch size (by the mean of the species and by the within-species standard

deviation in clutch size) and transformed clutch size to a proportion of the species mean. For

species that have no within-species variance in clutch size, we used a value of 0.01 for the

standard deviation in clutch size to prevent issues in calculations when using zero. We,

therefore, expressed variation in clutch size in three ways: a raw increase in clutch size, a

standardised clutch size and a proportional clutch size.

Meta-analysis

We ran a single model using the log odds ratio calculated for each clutch size transformation (i.e., raw, standardised and proportional) to determine the effect of parental survival, given an increase in reproductive effort using the *metafor* package [43] in R 3.3.2 [44]. From these models we were also able to directly compare the effect size of brood manipulation studies and observational studies by including a categorical fixed effect for study type (i.e., experimental or observational). We included phylogeny as a correlation matrix in these meta-analytic models to correct for shared ancestry. The phylogeny was obtained using BEAST to measure a distribution of 1,000 possible phylogenetic trees of the focal 30 species extracted from BirdTree [45]. We also included species and each studies' journal reference as random effects in the model. From these models, we calculated the proportion of variance explained by the phylogenetic effect [46].

We then tested the effect of the species' mean clutch size on the relationship between parental survival and clutch size. We ran a single model with the mean centred (from all species used in the meta-analysis) species' average clutch size in interaction with treatment (brood manipulation or observational). Species, phylogeny and reference were also included as random effects to correct for the similarity of effect sizes within species and studies.

The difference in survival for the different sexes was modelled for each clutch size measure. Brood manipulation studies and observational studies were analysed in separate models. Sex was modelled as a categorical moderator (41 female studies, 27 male studies and 10 mixed studies). Species, phylogeny and reference were included as random effects (Table S2 and Figure S2).

Publication bias

Much of the data used in this analysis were taken from studies where these data were not

the main focus of the study. This reduces the risk that our results are heavily influenced by a

publication bias for positive results. A funnel plot for the survival against raw clutch size

model is presented in Supplementary Information (Figure S3).

Fitness projections

We calculated various isoclines using the brood-manipulation overall effect size (-0.05,

based on raw clutch size) that we estimated. Here, an isocline is a trendline representing the

change in fitness returns over an individual's lifespan, given an increase in individual clutch

size. An estimated "lifetime reproductive output" was calculated for hypothetical control

populations (starting with 100 individuals), where all individuals consistently reproduce at the

level of a species mean and have a consistent annual survival rate. We calculated this

"lifetime reproductive output" (see Supplementary Information for equation details for

calculating lifetime reproductive output) using combinations of a range of species average

clutch sizes at 2, 4, 6, 8 and 10 and survival rates of 0.2, 0.3, 0.4, 0.5, 0.6 and 0.7, which

reflected the range of clutch sizes and survival rates seen in the species in our meta-

analysis. We started with a hypothetical population size of 100 and calculated the

populations lifetime reproductive output rate that depends on how many offspring are

produced in a clutch and how long individuals in the population live for. We could then use to

model the effects of the costs of reproduction using the meta-analytic estimate of how brood

size is associated with parental survival (through experimental studies).

The "lifetime reproductive output" estimate was then repeated for a hypothetical population

that reproduces at an increased level compared to control, i.e., brood size enlargement,

throughout their lives. These individuals therefore produce more offspring but face

reductions to survival. This analysis allows us to determine if an overall fitness gain is

achieved by producing more offspring despite paying an increased survival cost or whether

the survival cost balances out the fitness gains of producing more offspring through reduced

reproductive attempts. To obtain this, we added a range of 1–5 offspring to the clutch sizes

of the control populations. Using a range of increased clutch sizes allowed us to investigate

how increased reproductive effort would affect lifetime fitness. The survival costs were

determined by the overall effect size found for brood manipulation studies (per egg). We

modelled effect sizes of -0.05, -0.15 and -0.25, which represent, respectively, the meta-

analytic overall effect size, its upper confidence interval and a further severe effect within the

observed effect sizes (rounded to the closest 0.05 for simplicity). For example, an individual

who has an additional offspring in its nest would see a 5%, 15% and 25% (respectively for

each effect size) reduction in its survival odds compared to if it reproduced at its normal rate

(i.e., the control population rate).

We then calculated the selection differential ($LRS_{brood\ manipulation}$ / $LRS_{control}$) between the

hypothetical control and "brood manipulation" populations for each combination of survival

rate, clutch size and effect size, and plotted this as an isocline. We further plotted the fitness

consequences for five exemplar species, where survival rate and clutch size combinations

are observable in the wild. We used effect sizes from model predictions at these survival rate

and clutch size combinations rather than the meta-analytic mean, thereby providing a

biological context.

Data Availability

Data and R code used for the statistical analysis in this study are available in the Dryad

repository: https://doi.org/10.5061/dryad.q83bk3jnk.

Acknowledgements

This work was supported by a Natural Environment Research Council (NERC) Adapting to the Challenges of a Changing Environment (ACCE) studentship to LAW, a Sir Henry Dale Fellowship to MJPS (Wellcome and Royal Society; 216405/Z/19/Z) and a grant from NERC (NE/J024597/1) to TB. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising.

References

1. Lack, D. The Significance of Clutch-size. *Ibis* **89**, 302–352 (1947).

2. Harvey, P. H. & Clutton-Brock, T. H. Life history variation in primates. *Evolution* **39**, 559–559 (1985).

3. Ricklefs, R. E. Density Dependence, Evolutionary Optimization, and the Diversification of Avian Life Histories. *Ornithology Applications* **102**, 9–22 (2000).

4. Stearns, S. C. Life-History Tactics: A Review of the Ideas. *The Quarterly Review of Biology* **51**, 3–47 (1976).

5. van Noordwijk, A. J. & de Jong, G. Acquisition and Allocation of Resources: Their Influence on Variation in Life History Tactics. *The American Naturalist* **128**, 137–142 (1986).

6. Kirkwood, T. B. L. & Rose, M. R. Evolution of senescence: late survival sacrificed for reproduction. *Phil. Trans. R. Soc. Lond. B* **332**, 15–24 (1991).

7. Lemaître, J. F. & Gaillard, J. M. Reproductive senescence: new perspectives in the wild. *Biological Reviews* **92**, 2182–2199 (2017).

8. Reznick, D. Measuring the Costs of Reproduction. **7**, (1992).

9. Kirkwood, T. B. L. & Austad, S. N. Why do we age? *Nature* **408**, 233–238 (2000).

10. Flatt, T. & Partridge, L. Horizons in the evolution of aging. *BMC Biol* **16**, 93 (2018).

11. Pettifor, R. A., Perrins, C. M. & McCleery, R. H. Individual optimization of clutch size in great

tits. *Nature* **336**, 160–162 (1988).

12. Lessells, C. M. Brood Size in Canada Geese: A Manipulation Experiment. *Journal of Animal*

Ecology **55**, 669–689 (1986).

13. Monaghan, P. & Nager, R. G. Why don't birds lay more eggs? *Trends in Ecology and Evolution*

12, 270–274 (1997).

14. Conrad, K. F. & Robertson, R. J. Intraseasonal effects of clutch manipulation on parental

provisioning and residual reproductive value of eastern phoebes (Sayornis phoebe). *Oecologia* **89**,

356–364 (1992).

15. Hegner, R. E. & Wingfield, J. C. Effects of Brood-Size Manipulations on Parental Investment,

Breeding Success, and Reproductive Endocrinology of House Sparrows. *The Auk* **104**, 470–480

(1987).

16. Reznick, D. Costs of Reproduction : An Evaluation of the Empirical Evidence. *Oikos* **44**, 257–

267 (1985).

17. Zera, A. J. & Harshman, L. G. The Physiology of Life History Trade-Offs in Animals. *Annu. Rev.*

Ecol. Syst. **32**, 95–126 (2001).

18. Santos, E. S. A. & Nakagawa, S. The costs of parental care: a meta-analysis of the trade-off

between parental effort and survival in birds. *Journal of Evolutionary Biology* **25**, 1911–1917

(2012).

19. Cohen, A. A., Coste, C. F. D., Li, X.-Y., Bourg, S. & Pavard, S. Are trade-offs really the key

drivers of ageing and life span? *Functional Ecology* **34**, 153–166 (2020).

20. Charnov, E. L. & Krebs, J. R. On Clutch-Size and Fitness. *Ibis* **116**, 217–219 (1974).

21. Wilson, A. J. & Nussey, D. H. What is individual quality? An evolutionary perspective. *Trends in*

Ecology & Evolution **25**, 207–214 (2010).

22. Drent, R. H. & Daan, S. The Prudent Parent: Energetic Adjustments in Avian Breeding. *Ardea*

38–90, 225–252 (1980).

23. Descamps, S., Gaillard, J.-M., Hamel, S. & Yoccoz, N. G. When relative allocation depends on

total resource acquisition: implication for the analysis of trade-offs. *J. Evol. Biol.* **29**, 1860–1866

(2016).

24. Vedder, O. & Bouwhuis, S. Heterogeneity in individual quality in birds : overall patterns and

insights from a study on common terns. *Oikos* 001–008 (2017) doi:10.1111/oik.04273.

25. Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R. & Buckley, Y. M. Animal life

history is shaped by the pace of life and the distribution of age-specific mortality and

reproduction. *Nat Ecol Evol* **3**, 1217–1224 (2019).

26. Pujol, B. *et al.* The Missing Response to Selection in the Wild. *Trends in Ecology & Evolution*

33, 337–346 (2018).

27. Gingerich, P. D. Rates of Evolution. *Annu. Rev. Ecol. Evol. Syst.* **40**, 657–675 (2009).

28. Postma, E., Visser, J. & Van Noordwijk, A. J. Strong artificial selection in the wild results in

predicted small evolutionary change: Evolution is predictable, also in the wild. *Journal of*

Evolutionary Biology **20**, 1823–1832 (2007).

29. Chang, C., Moiron, M., Sánchez-Tójar, A., Niemelä, P. T. & Laskowski, K. L. What is the meta-

analytic evidence for life-history trade-offs at the genetic level? *Ecology Letters* **27**, e14354

(2024).

30. Boonekamp, J. J., Salomons, M., Bouwhuis, S., Dijkstra, C. & Verhulst, S. Reproductive effort

accelerates actuarial senescence in wild birds: An experimental study. *Ecology Letters* **17**, 599–

605 (2014).

31. Coulson, J. C. & Fairweather, J. A. Reduced reproductive performance prior to death in the

Black-legged Kittiwake: senescence or terminal illness? *Journal of Avian Biology* **32**, 146–152

(2001).

32. Rattiste, K. Reproductive success in presenescent common gulls (Larus canus): the importance of

the last year of life. *Proceedings of the Royal Society of London. Series B: Biological Sciences*

271, 2059–2064 (2004).

33. Simons, M. J. P., Briga, M. & Verhulst, S. Stabilizing survival selection on presenescent

expression of a sexual ornament followed by a terminal decline. *Journal of Evolutionary Biology*

29, 1368–1378 (2016).

34. Hammers, M., Richardson, D. S., Burke, T. & Komdeur, J. Age-Dependent Terminal Declines in

Reproductive Output in a Wild Bird. *PLoS ONE* **7**, e40413 (2012).

35. Smith, H. G., Kallander, H. & Nilsson, J.-A. The Trade-Off Between Offspring Number and

Quality in the Great Tit Parus major. *Journal of Animal Ecology* **58**, 383–401 (1989).

36. Reid, W. V. The cost of reproduction in the glaucous-winged gull. *Oecologia* **74**, 458–467

(1987).

37. Kalmbach, E., Griffiths, R., Crane, J. E. & Furness, R. W. Effects of experimentally increased

egg production on female body condition and laying dates in the great skua Stercorarius skua.

Journal of Avian Biology **35**, 501–514 (2004).

38. Järvistö, P. E., Calhim, S., Schuett, W., Velmala, W. & Laaksonen, T. Sex-dependent responses

to increased parental effort in the pied flycatcher. *Behavioral Ecology and Sociobiology* **70**, 157–

169 (2016).

39. Daan, S., Dijkstra, C. & Tinbergen, J. M. Family planning in the kestrel (Falco Tinnunculus): The

ultimate control of covariation of laying date and clutch size. *Behaviour* **114**, 1–4 (1990).

40. Verhulst, S. & Tinbergen, J. M. Experimental Evidence for a Causal Relationship between

Timing and Success of Reproduction in the Great Tit Parus M. major. *The Journal of Animal*

Ecology **60**, 269–282 (1991).

41. Tinbergen, J. M. & Daan, S. Family Planning in the Great Tit (Parus Major): Optimal Clutch Size

as Integration of Parent and Offspring Fitness. *Behav* **114**, 161–190 (1990).

42. Lemaître, J.-F. *et al.* Early-late life trade-offs and the evolution of ageing in the wild.

Proceedings. Biological sciences / The Royal Society **282**, 20150209–20150209 (2015).

43. Viechtbauer, W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical*

Software **36**, 1–48 (2010).

44. R Development Core Team. *R: A Language and Environment for Statistical Computing.* (R

Foundation for Statistical Computing, 2009).

45. Rubolini, D., Liker, A., Garamszegi, L. Z., Møller, A. P. & Saino, N. Using the birdtree.org

website to obtain robust phylogenies for avian comparative studies: A primer. *Current Zoology*

61, 959–965 (2015).

46. Nakagawa, S. & Santos, E. S. A. Methodological issues and advances in biological meta-analysis.

Evol Ecol **26**, 1253–1274 (2012).

609