1 Title: Reported Health Benefits of a Vegan Dog Food – A Likert Scale-

2 type Survey of 100 Guardians

Author

5 Mike Davies BVetMed CertVR CertSAO FRCVS

6 RCVS Specialist in Small Animal Clinical Nutritio

Contact details

Provet Limited

Kingsweir

Shaftesbury Road

Gillingham

Dorset

SP8 4LJ

Tel: 07930569948

Email: mikedaviesvet@gmail.com

Pet food manufacturer
21 Omni Pet Limited

22 85 Great Portland Street, 1st Floor

23 London

24   W1W 7LT

25   United Kingdom

Abstract

Introduction There has been a surge in feeding plant-based foods to pets and advocates extol the

health benefits of this practice. However, there is a lack of scientific evidence to support health

claims for vegan diets in dogs. Aims This study aimed to quantify perceived health changes by dog

guardians following the feeding of a single brand of UK-produced vegan food for a period of 3 to 12

months. Methods Dog guardians registered as feeding the vegan food for 3 - 12 months were

invited to participate in an online Likert Scale-type survey of observations reflecting health status.

Results 100 guardians completed the survey. The vegan food was acceptable (palatable), and

appetite and body weight were not adversely affected. Changes, including improvements, were

reported in the following areas: body condition score (BCS), activity, faecal consistency, faecal

colour, frequency of defaecation, flatus frequency, flatus antisocial smell, coat glossiness, scales in

haircoat (dandruff), redness of the skin (erythema, inflammation), crusting of the external ear canals

(otitis externa), itchiness (scratching; pruritus), anxiety, aggressive behaviour and coprophagia.

Conclusions This is the first study to quantitatively document guardian reports of specific positive

health benefits associated with feeding a UK vegan dog food. Further prospective, randomised,

controlled clinical trials are needed to validate and determine the significance of these observations.

Introduction

In recent years there has been an increase in feeding vegetarian and vegan foods to pets, with the

vegan pet food market being valued at $9.6 billion in 2020 and is estimated to reach $16.3 billion by

2030 (Allied Market Research 2022). Proponents of plant-based foods often claim a variety of health

benefits without providing scientific evidence to support their claims. A UK-based company (Omni)

launched a range of plant-based (vegan) dog foods in April 2021 and noticed that some customers

were posting messages online through a feedback service Trustpilot claiming improvements in their

pets' health, particularly skin and gastrointestinal problems. Trustpilot.com is a Danish consumer

review website founded in 2007 which hosts reviews of businesses worldwide.

In a recent paper (Knight and others 2022) 336 guardians feeding vegan diets participated in a global

online survey and the authors concluded that, "when compared to conventional meat-based or raw

meat rations the healthiest and least hazardous dietary choices for dogs, are nutritionally sound

vegan diets". In this study some health issues were mentioned but there were no data on specific

changes in health status following a switch to feeding vegan.

Previous published papers have shown that vegan diets labelled as complete and recipes for

homemade vegan rations did not comply with AAFCO or FEDIAF Nutritional Recommendations and

therefore, contrary to providing health benefits would represent a potential health risk (Dodd and

others 2021; Pedrinelli and others 2021; Starzonek and others 2021; Zafalon and others 2019).

One study (Cavanaugh and others 2021) looked at the short-term effects of a vegan diet on amino

acid, clinicopathologic, and echocardiographic findings and found no adverse effects and did not

detect essential amino acid or taurine deficiency.

All batches of the vegan diet involved in this study are analysed by the University of Nottingham to

ensure that they still exceed FEDIAF Guidelines after manufacture. Dog carers who feed them may

not be their legal owners and so in this paper they are referred to as guardians.

Objectives

The aim of this study was to document and quantify any changes in signs associated with health

status observed by guardians following a switch to feeding dogs a single brand of complete vegan

food for over 3 months.

The aim was to see if there was any improvement or deterioration in observations that may reflect

health status:

Acceptability of food when first fed

Appetite

Body weight

Body condition score (BCS)

Activity level

Gastrointestinal changes:

     a. Faecal consistency

     b. Frequency of defaecation

     c. Colour of faeces

     d. Flatus frequency

     e. Flatus antisocial smell

Skin signs

     a. Glossiness (shine)

     b. Scaly skin (dandruff)

     c. Redness (erythema) of the skin (inflammation)

     d. Crusting of the external ear canals (otitis externa)

     e. Itchiness (scratching; pruritus)

Behaviour

100   a.   Anxiety

101   b.   Aggression

102   c.   Any other behavioural changes

Hypothesis

106 The study was designed to acquire data to disprove the hypothesis that "Feeding a vegan diet to

107 dogs does not provide health benefits"

Ethics

110 The study was designed to comply with established ethical standards and it was decided that the

111 protocol did not need to be submitted to a formal external review panel. It complied with guidance

112 set out in the Royal College of Veterinary Surgeon's Ethical Review for Practice-based Research

113 (RCVS 2013) and guidance on ethics in questionnaires published online (Travis 2017)

Methods

Data Collection

118 All data was collected, stored and used in accordance with The Data Protection Act 2018 and the

119 General Data Protection Regulation (GDPR).

The Dog Food

122 The plant-based vegan pet food fed to the dogs in this study was formulated by veterinarians and

123 manufactured in the UK. The main ingredients incorporated into the recipes (present at >4%

124 inclusion rate) include: Potato Protein, Pea starch, Soya, Brown Rice, Sweet potato, Dried Brewers

125 Yeast, Oats, Peas, Pea Protein, Carrot and Rapeseed Oil.

Recruitment of guardians

On 18th March 2022 the company Omni sent individual emails using email marketing software

Klaviyo (https://www.klaviyo.com) to 307 randomly selected dog guardians who had been registered

as feeding their plant-based food for at least 3 months. They were invited to participate in an online

survey to generate basic information and data about perceived changes in health and informed

consent was obtained to publish the results afterwards.

The survey went live on 18th March 2022 and was terminated on 25th April 2022 by which time 101

people had completed it.

A second email was sent to 100 respondents later with a few supplementary questions regarding the

dog breed, sex and feeding practices before switching to the plant-based food

The Likert Scale-style Survey

An online Likert Scale-style questionnaire was created and analysed in accordance with established

guidelines (Boynton 2004; Sullivan and Artino 2013)

The questions were assessed for clarity, objectivity, appropriateness, avoidance of leading questions

and ease of use by veterinarians working with the company. The online questionnaire was created

using Typeform (https://www.typeform.com) to collect the data. Apart from the first question on

acceptability of the food and the question about colour of the faeces, the questions in the survey

were based on a 5-point Likert scale in which the middle option was "No change" and either side

were two options to report a slight or great improvement or deterioration in observations that may

reflect health status as described and listed in the study objectives:

Acceptability

Guardians were asked to select one of the following that best described what happened when they

first introduced the plant-based food to their dog.

Options:

1	2	3	4	5
My dog did not like the food at all and refused to accept it	Hardly any of the food was eaten and other foods had to be mixed with it	The food was introduced gradually over a few days and it was accepted well	Some of the food was eaten, the rest was eaten later	The whole meal was eaten with enthusiasm

160　Appetite

161　Guardians were asked: How good was your dog's appetite before you switched to the plant-based

162　food?

163　After changing to the plant-based food what effect, if any, was there on appetite:

164　Options:

1	2	3	4	5
Appetite was greatly reduced	Appetite decreased slightly – less hungry	No change in appetite	Appetite increased slightly	Appetite increased a lot with begging behaviour

168 ## Body Weight

169 Guardians were asked: After switching to the plant-based food what approximate changes did you

170 notice in body weight?

171 Options:

1	2	3	4	5
Body weight decreased by more than 10%	Body weight decreased by less than 10%	There was no change in body weight	Body weight increased by less than 10%	Body weight increased by more than 10%

Body Condition Score (BCS)

Guardians were asked: If you measure your dog's Body Condition Score (BCS) using a recognised

chart please answer this question (if not please skip this question).

After switching to the plant-based food what change have you noticed in body condition score

(BCS)?

Options:

1	2	3	4	5
The BCS decreased by 2 or more points	The BCS decreased by 1 point	There was no change in BCS	The BCS increased by 1 point	The BCS increased by more than 2 or more points

183 ## Activity level

184 Guardians were asked: How active was your dog before switching to the plant-based food?

185 After switching to the plant-based food what changes in activity, if any, did you notice?

186 Options:

1	2	3	4	5
Activity decreased a lot	Activity decreased a bit	There was no change my dog's activity	Activity increased a bit	Activity increased a lot

Gastrointestinal signs

Defaecation frequency

193 Guardians were asked: How many times per day did your dog defaecate (pass motions) per day

194 before you switched to the plant-based food?

195 After feeding the plant-based food what happened?

196 Options:

1	2	3	4	5
Defaecation decreased by 2 or more times per day	Defaecation decreased by 1x per day	No change in frequency of defaecation	Defaecation increased by 1x daily	Defaecation increased by 2 or more per day

Faeces (stool) consistency

201 Guardians were asked: What was your dog's faeces (stools) like before you switched to the plant-

202 based food?

203 After switching to the plant-based food how did the consistency of faeces (stools) change?.

204 Options:

1	2	3	4	5
Faeces (stools) became very watery and loose (diarrhoea)	Faeces (stools) became softer but were still formed	No change in consistency	Faeces (stools) became firmer and more well formed	Faeces(stools) became very hard and dry

Faecal colour

209 Guardians where asked: What colour were your dog's faeces (stools) before you switched to the

210 plant-based ration?

211 What happened to the faeces colour after you changed to the plant-based food?

Flatus (passing wind) frequency

214 Guardians were asked: Did your dog pass a lot of flatus "wind" before you switched to the plant-

215 based food?

216 After switching to the plant-based food did you notice any change in frequency of flatus?

217 Options:

1	2	3	4	5
Passing wind increased a lot	Passing wind increased a bit	There was no change in passing wind	Passing wind decreased a bit	Passing wind decreased a lot

Antisocial smelling flatus (wind)

222 Guardians were asked: Did your dog pass antisocial smelling wind before switching to the plant-

223 based diet?

224 After switching to the plant-based food did you notice a change in the smell of "wind" passed by

225 your dog?

226 Options:

1	2	3	4	5
The smell of wind got much worse	The smell of wind got a bit worse	There was no change in smell of wind passed by my dog	The smell of wind improved a bit	The smell of wind passed improved greatly

Dermatological signs
231

Haircoat Shine (Glossiness)

233 Guardians were asked: How shiny was your dog's coat before switching to the plant-based food e.g

234 dull, normal or shiny?

235 How did your dog's haircoat change (if at all) following the change to plant-based food?

236 Options:

1	2	3	4	5
There was a great decrease in glossiness of my	There was a slight decrease in glossiness of my	There was no change in glossiness of my	There was a slight increase in glossiness of my	There was a great increase in glossiness of hair

dog's hair coat	dog's hair coat	dog's hair coat	dog's hair coat	coat

Scale (Dandruff)

241 Guardians were asked: Did your dog have scales (dandruff) before switching to the plant-based

242 food?

243 What happened afterwards?

244 Options:

1	2	3	4	5
Scale increased greatly	Scale increased slightly	No change	Scale reduced slightly	Scale reduced greatly

Crusting in ears (Otitis externa)

249 Guardians were asked did your dog have crusting, wax build up or redness in its ear canal before

250 starting the plant-based food?

251 After switching to the plant-based food what changes if any did you notice

252 Options:

1	2	3	4	5
The amount of crusting increased a lot	The amount of crusting increased slightly	There was no change in crusting in the ear canal	The amount of crusting reduced	The amount of crusting resolved totally

Itchiness (Pruritus)

257 Guardians were asked whether their dog was itchy before switching to the plant-based food, and if

258 so to report what change (if any) they noticed after switching

259 Options:

1	2	3	4	5
There was a great increase in itching	There was a slight increase in itching	There was no change in itching	There was a slight decrease in itching	The itching resolved totally

Skin redness (erythema, inflammation)

264 Guardians were asked if their dog had skin redness (inflammation) before switching to the plant-

265 based diet and if so what change (if any) did they notice:

266 Options:

1	2	3	4	5
A great increase in redness	A slight increase in redness	No change in redness	A slight reduction in redness	Total resolution of the redness

Behavioural signs

Aggression

273 Guardians were asked: After switching to the plant-based food have you noticed any changes in

274 aggressive behaviour. If your dog never shows aggression you do not need to answer this question,

275 please select 'Not relevant' and move on

276   Options:

1	2	3	4	5
Aggressive behaviour got a lot worse	Aggressive behaviour got a bit worse	There was no change aggressive behaviour	The amount of aggression reduced	Previous aggressive behaviour resolved

Anxiety

281   Guardians were asked did your dog have signs of anxiety before starting plant-based food?

282   If so, after switching to plant-based food what changes did you notice in anxiety?

283   Options:

1	2	3	4	5
The amount of anxiety got much worse	The amount of anxiety got slightly worse	There was no change in anxiety	The amount of anxiety reduced slightly	My dog stopped showing any signs of anxiety

Other Behaviour

288   Guardians were asked: Did your dog have other behaviour problems before starting the plant-based

289   food e.g coprophagia (eating poo), chewing furniture? If so, please specify:

290   After switching to plant-based food what changes did you notice in behaviour?

291   Options:

1	2	3	4	5
The behavioural problem got much worse	The behavioural problem got slightly worse	There was no change in behaviour	The behavioural problem improved slightly	The behavioural problem resolved totally

Statistical analysis

296 Statistical analysis of Likert-type scale data is controversial (Sullivan and others 2013) and "the best

297 way to display the distribution of responses is to use a bar chart" (St Andrews University 2022). So,

298 the results of this study are expressed in simple descriptors and bar chart graphic format and no

299 attempt is being made to calculate statistical significance using p values.

Results

The Likert Scale-style survey

101 dog guardians completed the survey however It emerged that one had not actually started

feeding the plant-based food and so their answers had to be deleted from the analysis. Some

guardians failed to answer some sections, or did not answer questions correctly. An example of an

inappropriate answer was when some respondents claimed that their dog was not itchy before

switching diets, but then reported that there had been a great improvement afterwards. Clearly such

a response was not possible if the dog was not initially itchy so those results had to be excluded from

the analysis.

The survey response rate was 32.6%. In expressing the results all percentages have been rounded to

one decimal point.

Key

n=number of valid responses, out of 100 possible.

% = percent of the number of valid responses, expressed to one decimal point

NAn – Not answered

NAa – Not appropriate answer

NR – Not relevant

Acceptability (n=100)

82 dogs (82%) ate the novel plant-based food with enthusiasm when presented with it for the first

time. 10 dogs (10%) ate some of the food when it was first presented then went back and finished it

off later, 7 dogs (7%) ate the food when it was introduced gradually over several days, one dog

would only accept the food if it was mixed with other foods. None of the dogs refused to eat the

food. So, this plant=based food was highly acceptable (palatable).

Insert Fig 1 here

Appetite (n=97) (NAn-3)

The majority (70 (72.2%)) of guardians reported that their dogs' appetite did not change after

switching to the plant-based food. 16 guardians (16.5%) reported a slight increase in their dogs'

appetite and 7 guardians (7.2%) reported a great increase in appetite, with their dog showing

begging behaviour for more food. 4 guardians (4.1%) reported that their dog ate slightly less after

switching. No dogs lost their appetite totally.

Insert Fig 2 here

Body Weight Changes (n=100)

71 guardians (71%) reported no change in body weight following a change to the plant-based food.

17 (17%) reported a drop in weight of which 4 (4%) dogs lost more than 10% of initial weight and 13 (13%) less than 10% weight. 12 guardians (12%) reported that their dogs had gained weight on the plant-based food, of which 8 (8%) gained less than 10% weight and 4 (4%) gained more than 10% of initial weight.

Insert Fig 3 here

Body Condition Score (BCS) (n=20)

20 guardians (20%) claimed they were using an accredited BCS score chart and completed this section. In 13 (65%) of these dogs BCS did not change, 3 dogs (15%) gained 2 or more BCS points, 3 (15%) gained one BCS point, and one dog (5%) lost 1 BCS point

Insert Fig 4 here

Activity level (n=97) (NAn=3)

Following a switch to the plant-based food 67 (69.1%) dogs did not exhibit any change in activity. An increase in activity was reported for 28 (28.9%) dogs, in 19 (19.6%) activity was slightly increased and 9 (9.3%) became much more active. 4 (4.1%) of these dogs were not very active before the change in diet, 9 (9.3%) were active already and 3 (3.1%) were very active. Two dogs (2.1%) were reported to be slightly less active after the change in food. No dogs were reported to be a lot less active after switching to the vegan food.

Insert Fig 5 here

Gastrointestinal signs

Defaecation frequency (n=94) (NAn=6)

Prior to switching diets 9 dogs (9.6%) were defaecating once per day, 32 (34.0%) twice per day, 33

(35.1%) three times per day, 12 (12.8%) four times per day, 3 (3.2%) five times per day, 4 (4.3%) six

times per day and 1 (1.1%) eight times per day.

Insert Fig 6a here

Of the 94 guardians who completed this section 53 (56.4%) reported that there had been no change

in defaecation frequency after switching to the plant-based food. 11 guardians (11.7%) reported that

their dog passed one more motion per day, and 6 (6.4%) that their dog passed 2 or more additional

motions per day. The number of motions passed per day decreased by 1 in 16 dogs (17.0%) and by 2

or more per day in 8 dogs (8.5%).

Insert Fig 6b-h here

Faecal consistency (n=96) (NAn=4)

Prior to switching to a plant-based food 52 (54.2%) guardians described their dog's faeces as being

of normal consistency, 22 (22.9%) as soft, 10 (10.4%) as watery, 9 (9.4%) as hard and 2 (2.1%) as very

hard. After switching a total of 35 guardians (36.5%) reported no change in faecal consistency, 37

(38.5%) reported the faeces was firmer, 2 (2.1%) that the faeces was much harder, 22 (22.9%)

reported that the faeces was softer and no guardians reported that the faeces became watery.

In the dogs with normal faeces there was no change in consistency in 33 (63.5%), in 8 (15.2%) the

stools became firmer and 11 (21.2%) softer after diet change.

In dogs with watery faeces before switching to plant-based ration there was no change in one dog

but 9 out of ten (90.0%) developed firm faeces on the plant-based food. For dogs with soft faeces

prior to diet change 19 out of 22 (86.4%) developed firm faeces after the change, there was no

change in consistency in two (10.5%) of the dogs and in one dog (5.2%) faeces got softer.

For dogs with hard faeces initially 6 out of 9 (66.7%) the dog's faeces was softer but still formed after

switching, in 2 (22.2%) dogs the faeces was harder after switching and in one there was no change.

No guardians reported watery faeces after switching to the plant-based diet.

Insert Fig 7 here

Faecal Colour (n=93) (NAn=7)

Prior to changing to the plant-based ration 47 (50.5%) of guardians described their dog's faeces as

being brown in colour, 18 (19.4%) described it as light brown and 18 (19.4%) described it as dark

brown. 6 (6.5%) guardians claimed their dog had yellow faeces, 2 (2.2%) black faeces and 2 (2.2%)

orange faeces.

Following a change to the plant-based food 39 guardians (41.9%) reported no change in colour, 29

(31.2%) said the faeces was lighter and 23 (24.7%) darker. Of the dogs with light brown faeces

initially 15 (83.3%) got darker, and of the dogs with dark brown faeces 16 (88.9%) got lighter, of dogs

with yellow faeces initially 3 (50.0%) changed to dark brown, 2 (33.3%) did not change, and one

(16.7%) changed to light brown. In the two dogs with orange coloured faeces one did not change

and the other became lighter in colour. Of the two dogs with black faeces in both the faeces became

lighter.

Insert Fig 8 here

Flatus (passing wind) Frequency (n=96) (NAn=4)

Of the 96 dogs 74 (77.1% %) did not pass a lot of wind prior to diet change. Of these there was no

change in 64 (86.5%), 8 (10.8%) passed more wind after the change in diet with 2 (2.7%) dogs

passing a lot more.

22 (22.9%) guardians reported that their dogs passed a lot of wind prior to diet change and

afterwards in these dog's flatus was reduced a lot in 13 (59.1%), reduced slightly in a further 5

(22.7%) and there was no change in 4 (18.2%). None of these guardians reported an increase in

passing wind after changing to a plant-based diet

Insert Figs 9a and 9b here

Antisocial smelling flatus (wind) (n=97) (NAn=3)

Of 97 guardians who completed this section 26 (26.8%) reported that their dog passed antisocial

smelling wind before switching to a plant-based diet. After changing diet, in 19 (73.1%) of these the

smell improved of which 17 (65.4%) reported that the smell improved greatly. In 7 (26.9%) of these

dogs there was no change in smell. The smell did not get worse in any of the dogs.

71 (73.2%) of guardians reported that their dog did not pass antisocial smelling wind prior to a

change to the plant-based food, afterwards there was no change in 65 (91.5%), the smell got slightly

worse in 5 dogs (7.0%) and in one dog (1.4%) the smell improved a bit.

Insert Figs 10a and 10b here

Dermatological signs

Hair Coat Glossiness

a. Before switching to the plant-based food (n=83) (NAn=9; NAa= 8)

Prior to switching to the plant-based food 49 (59%) of guardians reported that their dog had normal

hair shine, one (1.2%) said their dogs' coat was very dull and 5 (6.0%) that the haircoat was dull,

whereas 21 (25.3%) reported that their dog had shiny hair and 6 (7.2%) very shiny hair.

b. After switching to the plant-based food (n=98) (NAn=2)

After the switch to the plant-based ration 48 guardians (49.0%) reported an improvement in hair

coat glossiness with 26 of these (26.5%) reporting a slight increase in shine and 22 (22.4%) reporting

a great improvement in shine. Three guardians (3.1%) reported that their dogs' hair coat appeared

to be slightly duller after switching to the plant-based food. No guardians reported a great

deterioration in hair shine. 47 guardians (48.0%) reported no change in hair shine after switching

diets.

Insert Figs 11a and 11b here

Scale (Dandruff)

a. Before diet change (n=100)

80 dogs (80.0%) did not have any scale before the switch to plant-based food and 20 dogs (20.0%)

did.

b. After diet change (n=97) (NAn=3)

One normal dog developed slight scaling after the diet change but there was no change in the other

79 (81.4%). Of the 20 dogs with scale present 3 guardians did not answer the question, of the 17

responses there was no change noted in 4 (23.5%) whereas in the majority of 13 dogs (76.5%%)

either a slight improvement (7 dogs – 41.2%) or total resolution of signs (6 dogs – 35.3%) was

reported.

Insert Fig 12 here

Crusting in the ear canal (n=8) (NR=92)
The majority of guardians (92%) reported that their dogs did not have any crusting in their ear canal

before switching to the plant-based ration.

Of the 8 dogs that did have crusting in their ears this improved in 7 (85.7%) following diet change, it

improved a bit in 5 (62.5%) and resolved totally in 2 (25%). There was no change in the other dog

(12.5%). None of the normal dogs developed signs after diet change.

Insert Fig 13 here

Itching (n=71) (NR=29)
Of 100 guardians in the survey 29 (29.0%) reported that their dogs were not itching initially so the

majority of 71 dogs (71.0%) were. After changing diet there was no change in 43 (60.6%) but it

improved in 37 dogs (52.1%) and in 11 of these (15.5%) the itchiness resolved totally. Itching got

slightly worse in 2 dogs (2.8%).

Insert Fig 14 here

Skin redness (erythema) (n=27) (NAn=6) (NR=67)

67 (69.8%) of guardians reported that their dog did not have erythema (reddening) of the skin. Of

the 27 (28.7%) dogs that did have reddening there was no change in 15 (55.6%) after switching to

the plant-based diet but in 12 dogs (44.4%) the redness improved and in 9 (33.3%) of these dogs

there was total resolution of the redness.

Insert Fig 15 here

Behaviour changes

Aggression (n=98) (NAn=2)

Of 98 guardians who completed this section 32 (32.7%) described their dogs as showing signs of

aggression. After changing to the plant-based food in 23 (71.8%) the aggressive behaviour did not

change, in 8 dogs (25.0%) the signs of aggression reduced and in one of these the aggression

resolved totally. Aggression got a bit worse in one dog. None of the normal dogs developed

aggressive behaviour after switching to the plant-based food.

Insert Fig 16 here

Anxiety (n=100)

37 (37%) of dogs were described as showing signs of anxiety prior to changing to the plant-based

diet. Afterwards 20 (54.0%) did not show any change in anxious behaviour, 14 dogs improved

(*44.4%*) of which 11 (29.7%) improved slightly whilst 3 (8.1%) stopped showing any anxiety. In 3

(8.1%) dogs the anxiety got much worse after changing diet.

Insert Fig 17 here

Other Behaviours (n=86) (NAn=14)

Of 86 guardians who completed this section only 17 (19.8%) reported that their dog had another

behavioural problem before switching to plant-based food and the majority of these (14– 16.3%)

were coprophagic. Two dogs had a barking problem, and one each were reported to be showing

signs of chewing feet or restlessness

Coprophagia (n=14)

After switching in 6 (42.9%) of 14 affected dogs the behaviour resolved totally, one (7.1%) improved

slightly, there was no improvement in 6 (42.9%) and one (7.1%) got slightly worse.

Insert Fig 18 here

Supplementary questionnaire (n=50)

50/100 (50%) of guardians who completed the main questionnaire completed the supplementary

questionnaire.

Age (n=50)

There was a wide range of ages from 1.5 months to 13 years of age

Insert Fig 19 here

Breeds (n=50)

Most dogs (18- 36%) were crossbreeds, 4 were Labrador retrievers (8%), there were 3 Cockerpoos

(6%) and 2 each of Greyhounds, English Bull Terriers and Hungarian Vizlas. There were single dogs

representing 19 other breeds.

Insert Fig 20 here

Sex (n=50)

556 There were slightly more males than females Male – 28 (56%) ; Female 22 (44%)

Diet prior to switch to Plant-based food (n=50)

559 Prior to switching to the plant-based food in this study 11 guardians reported that they were feeding

560 another vegan food (22%) and 39 a meat-based food (78%) of which 4 (8%) were feeding a raw meat

561 ration.

Discussion

There are a number of important limitations to this study not least that the survey was conducted

with a population of dog carers who had decided to feed a vegan food and so were likely to be

positive-biased towards recognising positive health benefits.

The survey response rate of 32.6% was in line with reported rates for surveys (Fincham 2008) but

was much lower than some scientific Journals now expect which can be as high as 60-80% (Fincham

2008). The non-response bias (67.4%) in this study weakens the reliability and validity of the results

but only if one wants to apply the findings of the respondents to the whole population surveyed. The

aim of this study was simply to identify whether health benefits were being observed by dog

guardians, which it did, and so the non-response bias is less important. A higher response rate could

have been achieved by prior communication with potential participants, allowing the survey to be

available for a longer period, post-survey follow-ups and reminders, or the deployment of

multimodal methods to attract recruits (Fincham 2008).

The dogs in this study had only been fed the food for 3-12 months and so longer follow-up surveys

are needed to document long term benefits. Most of the participants were not dog health

professionals and so would not be qualified to accurately assess signs of health status. All the

outcome measures relating to health status that were provided in the questionnaire, except body

weight were subjective not objective, and data collection was incomplete, for example only 50% of

respondents to the primary survey provided the additional information requested. So, the results do

need to be viewed with a degree of scepticism.

Data derived from pets living under normal home conditions reflect the real-world situation

compared to those kept in controlled environments and the overwhelmingly positive results

reported here for this specific vegan food, chime with many anecdotal reports of health benefits and

provides initial evidence to support further studies.

One concern was that the plant-based diet might not be acceptable (palatable) to dogs, especially if

they were used to being fed a meat-based ration which 78% of the dogs were. However, over 82% of

the dogs liked the food and ate it all when first given it, and a further 10% ate it all after a short

break so 92% of dogs accepted the food immediately. Just 8% of dogs needed to have the food

introduced gradually. This is in agreement with another guardian-based study which reported that

vegan foods were as palatable as meat-based foods for both dogs and cats (Knight and Satchell

2021).

In dogs many factors affect appetite including odour, taste and texture. Appetite was not affected by

the change to plant-based ration in the majority (72.2%) of dogs in this study whereas 16.5% of

guardians reported that their dogs' food intake had increased and some of them (7.2%) were much

hungrier and showed begging behaviour for more food. Satiety is induced by the presence of food in

the gastrointestinal tract through a combination of neurological (e.g. distension of the tract) and

chemical messages (e.g. chemoreceptors in the intestinal wall; absorbed amino acids in the blood)

acting on the satiety centre in the brain. Increased food consumption could represent a strong liking

for the new food, or a lack of satiety when fed at the recommended feeding rate. Further studies are

needed in these dogs to see if the cause was inadequate calorie supply or something else.

The majority of dogs (71%) did not show any fluctuation in body weight which is to be expected if

the food was being fed at the appropriate rate to meet the dog's individual calorie requirements.

There are many causes of weight gain or loss in addition to high or low calorie intake and for dogs in

which weight varied by more than 10% further investigations would be warranted to determine the

underlying reason.

The plant-based food did not affect body weight or body condition score for over 65% of the dogs.

Many factors could have caused the increase or decrease in weight and BCS reported for some of

the dogs, for example calorie intake or exercise level, and further studies would be needed to

determine the reason for the changes in these cases.

Activity levels reportedly increased in 28.9% of dogs after switching to the plant-based food, and in

9.3% a great increase in activity was noted. Only 2 dogs reportedly were less active after changing

food, and there are many possible causes for this apart from diet-change that need investigating.

There was no change reported in the frequency of defaecation in the majority (56.0%) of dogs after

switching to the plant-based food. There was variability in response in individual dogs with 18

(19.1%) dogs having reduced frequency and 17 dogs (18.1%) having increased frequency. Many

factors can influence the frequency of defaecation for example faecal consistency, transit time

through the colon, or the presence of dietary fibre and more studies are needed to determine a

cause-and-effect relationship with the vegan food which does contain sources of both soluble and

insoluble fibres.

The observed effects on faecal consistency were very interesting as 28 out of 32 dogs (87.5%) with

soft or watery faeces improved after switching to the plant-based food and in 8 out of 11 dogs with

hard or very hard faeces stool consistency improved by becoming softer. Randomised controlled

clinical trials would be desirable to establish a clinical role for this vegan food in managing diarrhoea

and/or constipation.

Faecal colour can be influenced by many factors including food ingredients and the host

microbiome, but also infectious agents and haemorrhage. It was interesting that the majority of

guardians who described their dog as having light faeces initially perceived that it had become

darker (83%), and at the same time in the majority of dogs with dark brown faeces initially,

guardians perception was that it became lighter (88%).

Passing wind is a common problem with some dogs. In this study passing wind reduced in 18 (81.8%)

out of 22 dogs reported to be passing a lot before diet change, only 10 guardians (10.4%) reported

an increase in flatus after switching even though plant-based ingredients might be expected to

increase gas production through the fermentation of fibres

In dogs passing antisocially smelling wind prior to changing diets a switch to the plant-based food

resulted in an improvement in 19 out of 26 (73.08%) with a great improvement reported in 17

(65.38%). This vegan food might be beneficial in managing this problem in dogs.

49% of guardians reported that their dogs coat glossiness (shine) improved and 82% of dogs showed

an improvement in scaling after switching to the plant-based diet furthermore in over half of the

dogs with dandruff the problem resolved totally. There are many factors that can influence

glossiness and scaling (dandruff) including nutritional deficiencies: vitamin A, vitamin B complex,

zinc and essential fatty acids (Scott and others 1995). So, one of the most likely associations linking

improvement after change of diet and clinical signs would be correction of an actual or relative

essential nutrient deficiency. Essential n6 fatty acids are important for skin and hair health and

plant-based oils provide a higher amount of essential Omega-6 fatty acids than animal-derived fats

so dogs switching from meat-based to plant-based rations might experience an improvement

Of the 8 dogs reported to have wax, redness and crusting in their ears, indicative of otitis externa, 2

were being fed another vegan diet before the change and the majority (85.7%) of dogs (including the

vegan fed dogs) improved after switching to the study diet. In 2 dogs (25% of those affected) the

condition resolved totally. In only one dog there was no change in crusting in the ears. There are

many factors that can influence external ear canal condition and these observations are difficult to

explain unless the food rectified an underlying dietary deficiency, such as protein (essential amino

acids) or fatty acids (Scott and others 1995) that was impairing normal skin physiology or

suppressing immune activity. Prospective randomised controlled studies are needed to confirm a

cause-and-effect relationship with the diet.

Perhaps surprisingly, 71% of dogs were reportedly scratching before the diet change and the signs

improved afterwards in 37 dogs (52.1%) and in 11 of these (15.5%) the itchiness resolved totally.

There are many possible causes of pruritus including ectoparasites, bacterial infections, atopy, food

sensitivities (allergies or intolerances), hormone imbalance, and dietary deficiencies including

Vitamin A, Vitamin B12 (cobalamin), Vitamin D, omega-6 and omega-3 polyunsaturated fatty acid

(Saevik and others 2004; Scott and others 1995), and iron (Yonova 2007). Naturally occurring lipid

supplementation has been shown to reduce pruritus in dogs (Noli and others 2015).

In humans, pruritus has been associated with low serum zinc and reversed with zinc

supplementation and zinc products have been used for many years in people (Sanada 1987; Gupta

and others 2014) and dogs (Miller 1989).

It is well documented that compound therapeutic diets can reduce pruritus in dogs (Watson and

others 2022; Witzel-Rollins and others 2019), fatty acid supplementation (Bensignor and others

2008) and individual dietary components including Vitamin E (Plevnik Kapun and others 2014) and

phytochemicals such as polyphenols may also have anti-pruritic properties in canine atopy (Magrone

and Jirillo 2012)

The most likely reasons why a change in diet might result in improvement in itchiness would be

correction of an underlying deficiency, removal of a dietary allergen (in this situation animal

allergens) or the positive anti-pruritic effect of a nutrient or combination of nutrients in the food. In

any future prospective study it would be useful to employ a validated pruritus scoring system for

dogs (Rybnícek 2009) against which any improvements can be measured.

Over 28% of dogs had skin erythema (reddening) which indicates inflammation and there are many

possible causes of skin erythema including environmental factors, and dietary factors including

essential fatty acid, Vitamin E and selenium deficiency (Scott and others 1995). In this study the

redness improved in 55.6% of affected dogs and it totally resolved in 12 (44.4%). Further studies are

needed to determine the underlying reason for the improvements observed.

Aggression reportedly improved in 25% of dogs following a change in diet and only one showed a

slight increase in aggression. In human studies (Werbach 1995) several nutrients have been

associated with aggressive behaviour including niacin, pantothenic acid, thiamine, pyridoxine,

vitamin C, tryptophan and iron deficiencies. In rodents magnesium deficiency, and manganese

toxicity in people also cause aggression.

In dogs, tryptophan and tyrosine, omega-6 and omega-3 fatty acids have been hypothesised to be

associated with aggression and other behavioural changes (Bosch 2007). In a thesis and literature

review (Harju 2016) canine aggressive behaviour was reported to be decreased by low protein

content and tryptophan supplements.

In the study reported here the type of aggressive behaviour being displayed by the dogs was not

specified. In a previous study a diet with high protein concentration (32 %) increased fear induced

territorial aggression in dogs compared to low (17 %) and medium (25 %) protein concentrations

(Dodman et al. 1996). The vegan food in this study is relatively high in protein (30%) and yet

aggression reportedly only increased slightly in one dog, but decreased in many more.

In one study plasma concentrations of docosahexaenoic acid (DHA) were lower and the linoleic to α-

linolenic acid ratio was higher in aggressive than non-aggressive German Shepherds (Re et al., 2008)

suggesting that fatty acids may play a role. In another study dominant aggressive dogs had

significantly lower serum concentration of triglycerides than non-aggressive dogs (Pentürk & Yalcin

2003) and both these studies showed that total serum cholesterol and high-density lipoprotein

cholesterol levels are lower in aggressive than non-aggressive dogs. Similar associations have been

recorded in other species including monkeys, rats and humans (Buydens-Branchey et al., 2000;

Hillbrand & Spitz, 1999; Kaplan et al., 1994). This is interesting because the plant-based food in this

study contains no cholesterol, although the dog synthesises its own in the liver, and only one dog

was reported to have developed a slight increase in aggression.

Anxiety was reported for 36% of dogs however the type of anxiety (e.g. separation or fear) was not

specified. In 44.4% of these there was an improvement noticed on the vegan diet and this warrants

further studies to determine whether this was a genuine cause-and-effect relationship to the diet

Coprophagia is common in dogs and underlying causes include undernutrition, dietary nutrient

deficiency, gastrointestinal disorders such as pancreatic exocrine deficiency or inflammatory bowel

disease. In this study 46% of dogs stopped this antisocial behaviour after starting the plant-based

food which warrants further investigation to see if it might be a useful management strategy.

It should be noted that health benefits were reported in dogs that switched to this vegan diet from

other vegan foods, so it cannot be assumed that the positive observations reported for the specific

food in this study would necessarily result from feeding other plant-based foods.

Conclusions

Feedback from 100 dog guardians clearly demonstrates several positive observations of

improvements in health in some dogs including in the following areas: faecal consistency, frequency

of defaecation, flatus frequency, flatus antisocial smell, coat glossiness (shine), scales on the skin

(dandruff), redness of the skin (erythema, inflammation), crusting of the external ear canals (otitis

externa), itchiness (scratching; pruritus), anxiety, aggressive behaviour and coprophagia.

These observations could simply be random coincidence relationships and prospective, randomised,

controlled clinical studies are needed to confirm the clinical significance of these observations.

Nevertheless, this study confirms several positive health benefits so the hypothesis is disproved,

indicating that the feeding of vegan food to dogs does provide health benefits to dogs.

Acknowledgements

757 I would like to thank Guy Sandelowsky and Shiv Sivakumar of Omni for allowing me to conduct this

758 survey of their customers with permission to publish the findings regardless of possible negative

759 findings.

Conflicts of interest

The author is currently a part-time independent consultant to Omni, a part-time external Lecturer at

the University of Surrey and Managing Director and Founder of Provet Limited.

Although this study was conducted in collaboration with Omni, members of the company did not

influence the process, interpretation or reporting of the data analysis.

References

Allied Market Research (2022) available online at :

https://www.alliedmarketresearch.com/vegan-pet-food-market-

A15847#:~:text=The%20vegan%20pet%20food%20market%20was%20valued%20at,cruelty%20or%2

0exploitation%20of%20animals%20in%20any%20sorts Last accessed 26th April 2022

Bensignor E, Morgan DM, Nuttall T. Efficacy of an essential fatty acid-enriched diet in managing

canine atopic dermatitis: a randomized, single-blinded, cross-over study. Vet Dermatol 2008;19:

156–162

Bosch G, Beerda B, Hendriks WH, van der Poel AF, Verstegen MW. Impact of nutrition on canine

behaviour: current status and possible mechanisms. Nutr Res Rev. 2007 Dec;20(2):180-94. doi:

10.1017/S095442240781331X. PMID: 19079869.

Boynton P. M. Administering, analysing, and reporting your questionnaire. BMJ (Clinical research

ed.), 2004 328(7452), 1372–1375. https://doi.org/10.1136/bmj.328.7452.1372

Buydens-Branchey, L., Branchey, M., Hudson, J., Fergeson, P. Low HDL cholesterol, aggression and

altered central serotonergic activity. Psychiatry Research, 2000, vol. 93, pp. 93–102

Cavanaugh SM, Cavanaugh RP, Gilbert GE, Leavitt EL, Ketzis JK, Vieira AB. Short-term amino acid,

clinicopathologic, and echocardiographic findings in healthy dogs fed a commercial plant-based diet.

PLoS ONE 2021 16(10): e0258044. https://doi.org/10.1371/journal.pone.0258044

Dodd, S.A.S. Shoveller, A.K. Fascetti, A.J. Yu, Z.Z.; Ma, D.W.L. Verbrugghe, A. A Comparison of Key

Essential Nutrients in Commercial Plant-Based Pet Foods Sold in Canada to American and European

Canine and Feline Dietary Recommendations. Animals 2021, 11, 2348.

https://doi.org/10.3390/ani11082348

Dodman, N.H., Reisner, I., Shuster, L., Rand, W., Luescher, U.A., Robinson, I., Houpt, K.A. Effect of

dietary protein content on behavior in dogs. Journal of the American Veterinary Medical Association,

1996 vol. 208, pp. 376–379.

Fincham JE. Response rates and responsiveness for surveys, standards, and the Journal. Am J Pharm

Educ. 2008;72(2):43. doi:10.5688/aj720243

GDPR UK Government Published 2018 Available online at

https://www.gov.uk/government/publications/guide-to-the-general-data-protection-regulation Last

accessed 20th March 2022

Gupta M, Mahajan VK, Mehta KS, Chauhan PS. Zinc therapy in dermatology: a review. Dermatol Res

Pract. 2014;2014:709152. doi:10.1155/2014/709152

Harju C. Bachelor project, 15 hp, Animal Science – Bachelor's programme, Department of Anatomy,

Physiology and Biochemistry, Uppsala 2016 Available online:

https://stud.epsilon.slu.se/9460/1/harju_c_160825.pdf Last accessed 24th May 2022

Hillbrand, M. and Spitz, R.T. Cholesterol and aggression. Aggression and Violent

Behavior, 1999. vol. 4, pp. 359–370

Kaplan, J.R., Shively, C.A., Fontenot, M.B., Morgan, T.M., Howell, S.M., Manuck, S.B., Muldoon, M.F.,

Mann, J.J. Demonstration of an association among dietary cholesterol, central serotonergic activity,

and social behavior in monkeys. Psychosomatic Medicine, 1994 vol. 56, pp. 479–484.

Knight A, Satchell L. Vegan versus meat-based pet foods: Owner-reported palatability

behaviours and implications for canine and feline welfare. PLoS ONE 2021 16(6): e0253292.

https://doi.org/10.1371/journal.pone.0253292

Knight A, Huang E, Rai N, Brown H. Vegan versus meat-based dog food: Guardian-reported indicators

of health. PLoS ONE 2022 17(4): e0265662. https://doi.org/10.1371/journal.pone.0265662

Magrone T, Jirillo E. Influence of polyphenols on allergic immune reactions: mechanisms of action.

Proc Nutr Soc 2012; 71: 316–321.

Miller WH. Non-steroidal anti-inflammatory agents in the management of canine and feline pruritus.

In Kirk RW (Ed) Current Veterinary Therapy X 1986 WB Saunders Philadelphia p566

Noli C, Della Valle MF, Miolo A, Medori C, Schievano C. Efficacy of ultra-micronized

palmitoylethanolamide in canine atopic dermatitis: an open-label multi-centre study. Vet Dermatol.

2015 Dec;26(6):432-40, e101. doi: 10.1111/vde.12250. Epub 2015 Aug 18. PMID: 26283633.

Pedrinelli V, Zafalon RVA, Rodrigues RBA, Perini MP, Conti RMC, de Carvalho Balieiro JC, Brunetto

MA. Influence of number of ingredients, use of supplement and vegetarian or vegan preparation on

the composition of homemade diets for dogs and cats. BMC Vet Res. 2021 Nov 20;17(1):358. doi:

10.1186/s12917-021-03068-5. PMID: 34798889; PMCID: PMC8605502.

Penturk, S. and Yalcin, E. Hypocholesterolaemia in Dogs with Dominance Aggression.

Journal of Veterinary Medicine Series A, 2003 vol. 50, pp. 339–342

Plevnik Kapun A, Salobir J, Levart A et al. Vitamin E supplementation in canine atopic dermatitis:

improvement of clinical signs and effects on oxidative stress markers. Vet Rec 2014; 175: 560

RCVS (Royal College of Veterinary Surgeons). Ethical Review for Practice-based Research A report of

a joint RCVS / BVA working party published in 2013. https://www.rcvs.org.uk/news-and-

views/publications/ethical-review-for-practice-based-research/

Re, S., Zanoletti, M., Emanuele, E. Aggressive dogs are characterized by low omega-3

polyunsaturated fatty acid status. Veterinary Research Communications, 2008 vol. 32, pp. 225–

230

Rybnícek J, Lau-Gillard PJ, Harvey R, Hill PB. Further validation of a pruritus severity scale for use in

dogs. Vet Dermatol. 2009 Apr;20(2):115-22. doi: 10.1111/j.1365-3164.2008.00728.x. Epub 2009 Dec

19. PMID: 19171021

St Andrews University – Analysing Likert Scale/Type Data . Available online, last accessed on l10th

May 2022 https://www.st-

andrews.ac.uk/media/ceed/students/mathssupport/Likert.pdf?msclkid=d2a32a4ed03a11ec9d31a32

f98ba49ae

Saevik BK, Bergvall K, Holm BR et al. A randomized, controlled study to evaluate the steroid sparing

effect of essential fatty acid supplementation in the treatment of canine atopic dermatitis. Vet

Dermatol 2004; 15: 137–145

Sanada S, Kuze M, Yoshida O. Beneficial effect of zinc supplementation on pruritus in hemodialysis

patients with special reference to changes in serum histamine levels]. Hinyokika Kiyo. 1987

Dec;33(12):1955-60. Japanese. PMID: 3448919.

Scott DW, Miller WH and Griffin CE. Nutritional Skin Diseases In Muller and Kirk's Small Animal

Dermatology 5th Edition 1995 Published by WB Saunders Co p891

Starzonek J, von Lindeiner L, Vervuert I. Beurteilung in Deutschland erhältlicher veganer

Alleinfuttermittel für Hunde und Katzen (Assessment of vegan complete diets for dogs and cats

available in Germany]. Tierarztl Prax Ausg K Kleintiere Heimtiere. 2021 Aug;49(4):262-271. German.

doi: 10.1055/a-1552-2220. Epub 2021 Aug 23. PMID: 34425604.

Sullivan GM, Artino AR Jr. Analyzing and interpreting data from likert-type scales. J Grad Med Educ.

2013;5(4):541-542. doi:10.4300/JGME-5-4-18

Travis E. Ethics in questionnaires. (last updated September 26th 2017). Available online last accessed

24th May 2022.   https://bizfluent.com/info-8064289-ethics-questionnaires.htm

Watson A, Rostaher A, Fischer NM, Favrot C. A novel therapeutic diet can significantly reduce the

medication score and pruritus of dogs with atopic dermatitis during a nine-month controlled study.

Veterinary Dermatology 2022 33,(1) 55-e18

Werbach MR. Nutritional Influences on Aggressive Behavior. Journal Of Orthomolecular Medicine

1995 Vol. 7, No. 1, 1995

Witzel-Rollins A, Murphy M, Becvarova I et al. Non-controlled, open-label clinical trial to assess the

effectiveness of a dietetic food on pruritus and dermatologic scoring in atopic dogs. BMC Vet Res

2019; 15: 220

Yonova D. Pruritus in certain internal diseases. Hippokratia. 2007;11(2):67-71

Zafalon RVA, Risolia LW, Vendramini THA, Ayres Rodrigues RB, Pedrinelli V, Teixeira FA, et al.

Nutritional inadequacies in commercial vegan foods for dogs and cats. PLoS ONE 2020 15(1):

e0227046. https://doi.org/10.1371/journal.pone.0227046

Fig 6a Defaecation Frequency – Prior to diet change (n=94)

Changes in Defaecation Frequency – Originally 5 motions per day (n=3)

Changes in faecal consistency (n=96)

Fig 8 Changes in faecal colour (n=93)

Fig 9a Change in Flatus Frequency - dogs not initially passing a lot (n=74)

Fig 9b Change in Flatus Frequency - dogs initially passing a lot (n=22)

Fig 10a Change in Flatus Antisocial Smell - dogs initially passing a lot (n=26)

Fig 10b Change in Flatus Antisocial Smell - dogs not initially passing a lot (n=71)

Fig 11a Coat Glossiness — before diet change (n=83)

Fig 11b Coat Glossiness – after diet change (n=98)

Fig 12 Change in Dogs with Scales (Dandruff) - after diet change (n=20)

Fig 13 Ear Crusting Change (n=8)

Fig 14 Changes in itchiness (n=71)

Fig 15 Changes in redness of skin (n=27)

Fig 16 Changes in Aggressive Behaviour after diet change (n=32)

Fig 17 Changes in Anxiety after diet change (n=36)

Fig 18 Changes in Coprophagia after diet change (n=14)

Fig 19 Age Distribution (n=50)

Fig 20 Breed Distribution (n=50)

- Crossbreeds
- Labrador retriever
- Cockapoo
- Greyhound
- Hungarian vizla
- English Bull Terrier
- Others (single cases)

Fig 1 Acceptance of food (n=100)

Fig 2 Effect on appetite (n=98)

Fig 3 Effect on body weight (n=100)

Fig 4 Effect on Body Condition Score (n=20)

Fig 5 Effect on Activity (n=97)

Fig 6b Defaecation Frequency – Change after diet change (n=94)

Changes in Defaecation Frequency – Originally 1 motion per day (n=9)

Changes in Defaecation Frequency – Originally 2 motions per day (n=31)

Changes in Defaecation Frequency – Originally 3 motions per day (n=33)

Changes in Defaecation Frequency – Originally 4 motions per day (n=14)

Changes in Defaecation Frequency – Originally 8 motions per day (n=1)

Fig 7a Changes in faecal consistency when watery initially (n= 10)

Fig 7b Changes in faecal consistency when soft initially (n= 22)

Fig 7c Changes in faecal consistency when normal initially (n= 52)

Fig 7d Changes in faecal consistency when hard initially (n= 9)