vsRNAfinder: a novel method for identifying high-confidence viral small RNAs

from small RNA-Seq data

Zena Cai[1,#], Ping Fu[1,#], Ye Qiu[1], Aiping Wu[2,3], Gaihua Zhang[4], Yirong Wang[1], Taijiao Jiang[5],

Xing-Yi Ge[1], Haizhen Zhu[1], Yousong Peng[1,*]

[1]Bioinformatics Center, College of Biology, Hunan Provincial Key Laboratory of Medical

Virology, Hunan University, Changsha 410082, China

[2]Institute of Systems Medicine, Chinese Academy of Medical Sciences & Peking Union Medical

College, Beijing, China

[3]Suzhou Institute of Systems Medicine, Suzhou, China

[4] The National and Local Joint Engineering Laboratory of Animal Peptide Drug Development,

College of Life Sciences, Hunan Normal University, Changsha 410081, China

[5] Guangzhou Laboratory, Guangzhou, China

# Contributed equally;

*To whom correspondence should be addressed.

Abstract

Virus-encoded small RNAs (vsRNA) have been reported to play an important role in

viral infection. Unfortunately, there is still a lack of an effective method for vsRNA

identification. Besides, a systematic characterization of vsRNAs is also needed.

Herein, we presented vsRNAfinder, a de novo method for identifying high-confidence

vsRNAs from small RNA-Seq (sRNA-Seq) data based on peak calling and Poisson

distribution and is public available at https://github.com/ZenaCai/vsRNAfinder.

vsRNAfinder outperformed two widely-used methods namely miRDeep2 and ShortStack in identifying viral miRNA, siRNA and piRNA. It can also be used to identify sRNAs in animals and plants with similar performance to miRDeep2 and ShortStack. Based on vsRNAfinder, a total of 19,734 high-confidence vsRNAs including 2,746 miRNAs were identified in 64 viral species from public sRNA-Seq data. The number of vsRNAs identified in viruses varied from 1 to 2,489 with a median of 170. The length distribution of vsRNAs peaked at 21 and 22 nt. Plant viruses were found to express larger number and higher levels of sRNAs than those of animal viruses. Besides, the number of vsRNAs identified increased as the viral infection persisted. Interestingly, the vsRNA showed strong expression specificity as little overlap was observed among vsRNAs identified in different strains of a virus, or in different hosts, cells or tissues infected by the same virus. Little conservation was observed among vsRNAs of different viruses. Overall, the study not only presents an effective method for vsRNA identification, but also provides a rich resource for further studies of vsRNAs.

Introduction

Small RNAs (sRNAs) are a kind of regulatory RNAs in cellular processes [1]. They are typically less than 200 nucleotides (nt) in length [1]. The sRNAs can be mainly divided into three classes including microRNA (miRNA) with a length of 18-24 nt, small interfering RNA (siRNA) with a length of 21-23 nt and piwi-interacting RNA (piRNA) with a length of 24-31 nt based on the mechanism of biogenesis and function

[2]. By the mechanism of biogenesis, the miRNAs are derived from single-stranded RNA precursors with a hairpin structure which are firstly processed by Drosha in the nucleus and then processed by Dicer after being transported to the cytoplasm; the siRNAs are derived from double-stranded RNA precursors in the cytoplasm and are processed by Dicer; the piRNAs are derived from the fragmentation of single-stranded RNA precursors via a Dicer-independent mechanism [3-5]. By the mechanism of function, although all three kinds of sRNAs can target the mRNAs to regulate their expressions by incorporating sRNAs into the RNA-induced silencing complex (RISC), their function mechanisms differ a lot: with the help of AGO protein, the miRNAs can repress mRNAs translation by binding imperfectly to their mRNA targets, while the siRNAs can degrade mRNAs by binding perfectly to their mRNA targets; the piRNAs can degrade mRNAs by binding perfectly their mRNA targets with the help of PIWI protein [5, 6]. The sRNAs play a fundamental role in the maintenance of normal physiology and serve as potential biomarkers for early diagnosis and therapeutic targets for a wide spectrum of diseases, particularly the cancer [1]. For example, miR-130 can down-regulate the expression of miRNA-221 and miRNA-222 by targeting the MET, thus reducing TNF-related apoptosis-induced ligand (TRAIL) resistance in non-small cell lung cancer [7].

Numerous sRNAs have been discovered in multiple organisms as the development of the next generation sequencing (NGS) technology and a growing number of sRNA databases are emerging, such as the miRBase [8], SEAweb [9] and ASRD [10]. The miRBase database is the most comprehensive database of miRNAs. It contains 48,860

mature miRNAs from 271 organisms, and provides rich information such as the

expression, confidence, and method of identifying miRNAs. The SEAweb database

integrates multiple classes of sRNAs from 10 organisms based on data-mining from

over 4,200 small RNA sequencing (sRNA-Seq) samples. The ASRD database

provides a platform for the comparison of sRNA abundances across over 2,000

sRNA-Seq samples in *Arabidopsis*. In recent years, lots of virus-encoded sRNAs

(vsRNA) have been identified and proved to be functional in the regulation of viral

life cycle and pathogenesis [11, 12]. For example, the DENV-vsRNA-5 was reported

to suppress viral replication by targeting the coding region of viral non-structural

protein 1 [13], and the svRNA-N of the Severe Acute Respiratory Syndrome

coronavirus (SARS-CoV) was involved in the enhancement of lung inflammatory

pathology [14]. However, most studies only investigated specific vsRNA class such as

miRNA, or focused on vsRNAs from one or a few viruses. There is still a lack of

systematic studies about vsRNAs.

Numerous methods have been developed to identify sRNAs from sRNA-Seq data.

They can be mainly divided into two kinds. The first kind of method is to identify

sRNAs based on a known sRNA library, such as the sRNAbench [15], Manatee [16]

and sRNAnalyzer [17]. This kind of method can be used for accurate quantification of

known sRNA. However, they may not be suitable for organisms with only a few

known sRNAs such as viruses. The other kind of method is to identify sRNAs *de*

novo by firstly finding the precursor of sRNAs from assembly of reads and then

detecting the mature sRNA, such as miRDeep2 [18] and ShortStack [19]. Most

methods mentioned above have been mainly used in the identification of sRNAs in

animals and plants. The viruses have much smaller genomes and stronger coding

ability compared to animals and plants. Thus, the distribution of vsRNAs is much

denser than that of other organisms, which may lead to assembly of adjacent vsRNAs

with overlaps into long contigs and missed detection of many vsRNAs by current

methods.

Herein, we presented vsRNAfinder, a *de novo* method for the identification of

high-confidence vsRNAs from sRNA-Seq data based on peak calling and Poisson

distribution. The method can also be used in the identification of sRNAs in animals

and plants. Then, we used vsRNAfinder to systematically identify vsRNAs from over

60 viral species and further characterized these vsRNAs. The work would greatly

facilitate further studies of vsRNAs.

Materials and methods

Data collection

The viral-infection-related sRNA-Seq data were obtained by firstly searching "virus"

in the NCBI GEO [20] and SRA [21] databases. Then, the searching results were

filtered by the study type and by the strategy, respectively, in the NCBI GEO and SRA

database. Only those with the study type of "Expression profiling by high throughput

sequencing" or "Non-coding RNA profiling by high throughput sequencing", and

those with the strategy of "RNASeq" or "other" were kept. Then, the sRNA-Seq data

were further filtered manually and only those of viral infections with the genome

sequence of the virus used in the infection available were kept for further analysis. A

total of 102 sRNA-Seq datasets were obtained for use in the study (Table S1).

To demonstrate the usage of our method in the identification of sRNAs from other

organisms, some representative sRNA-Seq datasets of several widely-studied

organisms including *Homo sapiens*, *Mus musculus*, *Drosophila melanogaster*,

Arabidopsis thaliana and *Oryza sativa* were also obtained from NCBI GEO [20] and

SRA [21] database (Table S3).

To validate the performance of sRNA identification methods, the known miRNAs in

viruses and other organisms were obtained from the database of miRBase (Release

22.1, http://mirbase.org/) [8]. The experimentally-verified siRNAs in Sugarcane

mosaic virus and piRNAs in Dengue viruses were collected from the studies of Xia et

al. [22] and Wang et al. [23], respectively.

The vsRNAfinder workflow

The vsRNAfinder workflow included four modules: Preprocessing, Identification,

Filtering and annotation, and Quantification, which were described as follows.

Preprocessing module. Firstly, the adapter sequences are trimmed from the raw reads

with cutadapt (version 2.6, parameter: '-m 15') [24] (Figure 1). Then, the clean reads

are mapped to the viral genome directly (the default), or the clean reads are firstly

mapped to the host genome and then the unmapped reads are mapped to the viral

genome, with bowtie (version 1.3.0, parameter: '-v 0 -m 2') [25]. The resulted SAM

files are converted to BAM files which are sorted by samtools (version 1.9) [26]. The

sorted BAM files are further converted to BED files by bedtools (version 2.29.2) [27].

Identification module. The coverage of mapped reads on the viral genome is

calculated by bedtools [27] and then smoothed with the window-size of 11 nt. The

peaks of the smoothed curve are identified using the function of *findPeaks*

("minpeakheight = 5") in the R package pracma (version 2.3.3) [28]. The start and

end site of the candidate sRNA is restricted to the upstream and downstream 30 nt of

the peak, respectively. Then, the number of reads starting at each position in the

upstream region of the peak, and that ending at each position in the downstream of the

peak are calculated. The position with the highest number of reads in the upstream

and that in the downstream are considered as the start and end site of the candidate

sRNA, respectively.

Filtering and annotation module. To obtain the high-confidence sRNAs, the

abundances of the start and end sites (defined as boundary sites) of the candidate

sRNA are normalized using RPM (Reads Per Million) which is calculated by dividing

the number of reads starting at the start site or ending at the end site by the total

number of clean reads, and then multiplied by 10^6. Only the candidate sRNAs with

both of the boundary sites having at least 5 RPM and with lengths ranging from 15 to

34 nt are retained for further analysis.

Then, the Poisson distribution is used to evaluate the statistical significance of

candidate sRNAs. Poisson distribution is a discrete probability distribution that

expresses the probability of a given number of events occurring in a fixed interval of

time, which is widely applied in NGS data analysis [29]. We hypothesized that the

reads should be enriched in the region where the sRNA is expressed compared to the

background region. Thus, the significance of candidate sRNAs could be analyzed by

the Poisson distribution of which the parameter λ can be calculated as the average

coverage of the mapped reads on the background region. For each candidate sRNA,

the background region is defined as extending the upstream and downstream 10 nt of

the candidate sRNA. Therefore, the λ is local and is calculated as follows:

$$\lambda = \frac{C}{L}$$

163   where C is the sum of the coverage of mapped reads on each position of the

164   background region, and L is the length of the background region.

165   Based on the local λ, for a given position X in the candidate sRNA with k mapped

166   reads covering the position, the p-value of the position is calculated as follows:

$$P_\lambda (X = k) = \frac{\lambda^k}{k! * e^{-\lambda}}$$

The p-value of the candidate sRNA is calculated as the 80% quantile of p-values in all

positions of the candidate sRNA. The candidate sRNAs with a p-value smaller than

0.05 are considered as statistically significant and highly confident.

The miRNAs are then annotated from the high-confidence sRNAs in two steps. Firstly,

the hypothesized precursor of miRNA (pre-miRNA) is obtained. Since the miRNA

may be located in either 5'-arm or 3'-arm of the hypothesized pre-miRNA, both cases

are considered. If the miRNA is supposed to be in the 5'-arm of the hypothesized

pre-miRNA, the start and end positions of hypothesized pre-miRNAs are determined

as follows according to Michael et al.'s study [30]:

$$start_{pre-miRNA} = start_{sRNA} - step * i$$

$$end_{pre-miRNA} = start_{sRNA} + 2 * L_{sRNA} - 1 + L_{loop} + step * i$$

176   otherwise,

$$start_{pre-miRNA} = start_{sRNA} - (L_{loop} + L_{sRNA} + step * i)$$

$$end_{pre-miRNA} = end_{sRNA} + step * i$$

where $start_{sRNA}$ and end_{sRNA} refer to the start and end position of the sRNA,

respectively; L_{sRNA} refers to the length of sRNA; L_{loop} refers to the length of loop and

is set to be 15; $step$ and i are set to be 5 and 8 for animals and viruses, 7 and 10 for

plants as plant pre-miRNA sequences are longer than those of animals and viruses,

according to Michael et al.'s study [30].

Then, the secondary structure of the hypothesized pre-miRNA is predicted by

RNAfold (version 2.4.18) [31] with default parameters. The hypothesized pre-miRNA

which has the minimum free energy of less than -25 kcal/mol, and in which at least 70%

of positions in the corresponding sRNA form base pairs, is considered to be

pre-miRNA having the hairpin structure, and is further processed into the mature

miRNA.

Quantification module. The abundances of sRNAs are normalized using RPM. The

total number of reads mapping to the sRNA is calculated as the sum of reads that

overlap with the sRNA by 80% and those that are utilized to estimate boundary sites

of the sRNA. The RPM of each sRNA is calculated by dividing the total number of

reads mapping to the sRNA by the total number of clean reads in the sample, and then

multiplied by 10^6.

Prediction of sRNAs with miRDeep2 and ShortStack

The miRDeep2 (version 0.1.2, https://github.com/rajewsky-lab/mirdeep2) [18] and

ShortStack (version 3.8.5, https://github.com/MikeAxtell/ShortStack) [19] were used

for prediction of sRNAs with default parameters.

Evaluating the performance of computational methods in identifying sRNAs

The known sRNAs (mentioned in Data collection) including miRNAs, siRNAs and

piRNAs were used to evaluate the performance of computational methods in

identifying sRNAs. For a given organism, the identified sRNAs whose sequences

match perfectly to those of known sRNAs are considered as true positives, otherwise

false positives; the known sRNAs which are missed by the method are considered as

false negatives. Three metrics including Precision, Recall and F1-score were used to

evaluate the performance of computational methods in identifying sRNAs, which are

calculated as follows:

$$Precision = \frac{TP}{TP + FP}$$

$$Recall = \frac{TP}{TP + FN}$$

$$F1 - score = \frac{2 * Precision * Recall}{Precision + Recall}$$

while TP is the number of true positives, FP is the number of false positives, FN is the

number of false negatives.

Statistical analysis

The Mann-Whitney U test was used to compare the expression differences between

the conserved vsRNAs and non-conserved vsRNAs. A p-value less than 0.05 was

considered as statistically significant. All statistical analyses were conducted in

Python (version 3.8).

Results

Overview of vsRNAfinder

The vsRNAfinder consisted of four modules including Preprocessing, Identification,

Filtering and annotation, and Quantification (Figure 1). The Preprocessing module

processed raw reads to clean reads by trimming adapters. Then, the clean reads were

firstly mapped to the host genome and then the unmapped reads were mapped to the

viral genome, or the clean reads were mapped to the viral genome directly (the default)

since the host reference genome was not always available.

The Identification module identified the candidate sRNAs with four steps: firstly, the

coverage of mapped reads across the viral genome was calculated; secondly, the

coverage was smoothed to reduce the noise of the peak detection and enhance the

peak signal; thirdly, the peak was detected which had a higher expression than its

adjacent sites; finally, the start and end sites of the candidate sRNA were determined

at the upstream and downstream of the peak identified, respectively (see Materials

and methods).

The Filtering and annotation module was designed for obtaining high-confidence

sRNAs and annotating miRNAs. The candidate sRNAs identified above were firstly

filtered by the abundance of the start and end sites (defined as boundary sites) of

sRNAs. Only the candidate sRNAs with both boundary sites having at least 5 RPM

were kept. Then, they were further filtered by the length. Only the candidate sRNAs

with 15-34 nt were kept which covered the length of most miRNAs, siRNAs and

piRNAs [2]. Then, the statistical significance of candidate sRNAs was obtained based

on the Poisson distribution. The remaining candidate sRNAs with a p-value smaller

than 0.05 were considered as statistically significant and highly confident. Finally, the

RNAfold [31] was used to annotate miRNAs from the high-confidence sRNAs (see

Materials and methods).

The Quantification module quantified the abundance of sRNAs by RPM. The total

number of reads mapping to the sRNA was calculated as the sum of reads that

overlapped with the sRNA by 80% and those that were utilized to estimate boundary

sites of the sRNA (see Materials and methods).

Figure 1. Overview of vsRNAfinder which consisted of four modules including

Preprocessing, Identification, Filtering and annotation, and Quantification. Please see

Materials and methods for details.

Performance analysis of vsRNAfinder in identifying viral miRNAs

To evaluate the performance of vsRNAfinder in identifying vsRNAs, vsRNAfinder

was used to identify viral miRNAs since the miRNAs are the most studied vsRNAs.

Four viruses including Murid gammaherpesvirus 4 (MuHV-4), Murid betaherpesvirus

1 (MuHV-1), Human betaherpesvirus 5 (HHV-5) and Human gammaherpesvirus 4

(HHV-4) were selected in the evaluation which had more than 20 viral miRNAs

reported in public databases. For comparison, two *de novo* methods, namely the

miRDeep2 [18] and ShortStack [19] which are widely used in identifying sRNAs,

were selected in identifying viral miRNAs in four viruses.

As shown in Figure 2A, vsRNAfinder achieved the highest Recall ranging from 0.48

to 0.8 in all viruses, followed by miRDeep2 (0.21-0.64) and ShortStack (0.03-0.34).

Taking the MuHV-4 as an example, vsRNAfinder had a Recall of 0.54, while

miRDeep2 and ShortStack had a Recall of 0.21 and 0.07, respectively. In terms of

Precision, the ShortStack performed best with Precision higher than 0.5 in all viruses.

The vsRNAfinder had Precisions ranging from 0.28-0.36 which were similar to those

of miRDeep2. Overall, when considering both the Recall and Precision using the

F1-score, the vsRNAfinder achieved the F1-scores of 0.41, 0.41 and 0.47, respectively,

in viruses of MuHV-4, MuHV-1 and HHV-5, which were much higher than those of

the miRDeep2 (0.26, 0.33 and 0.26) and ShortStack (0.13, 0.07 and 0.32). While in

the virus of HHV-4, three methods had similar F1-scores.

Figure 2. Performance analysis of vsRNAfinder, miRDeep2 and ShortStack in

identifying viral miRNAs. (A) Comparison of Recall, Precision and F1-score of

vsRNAfinder with miRDeep2 and ShortStack in identifying viral miRNAs in viruses

of Murid gammaherpesvirus 4 (MuHV-4), Murid betaherpesvirus 1 (MuHV-1),

Human betaherpesvirus 5 (HHV-5) and Human gammaherpesvirus 4 (HHV-4). (B)

Venn diagram showing overlaps between viral miRNAs identified by vsRNAfinder,

miRDeep2, ShortStack, and those in the miRBase database.

When comparing the number of viral miRNAs identified by three methods and those

in miRBase, we found that most viral miRNAs identified by miRDeep2 and

ShortStack were also identified by vsRNAfinder. For example, 19 and 3 viral

miRNAs were identified by miRDeep2 and ShortStack, respectively, in the MuHV-4,

among which 14 and 3 were also identified by vsRNAfinder. Moreover, vsRNAfinder

identified more novel viral miRNAs than the other two methods in viruses of

MuHV-4, MuHV-1 and HHV-4 (Figure 2B).

Identification of other classes of sRNAs with vsRNAfinder

Besides viral miRNAs, viruses can also express siRNAs and piRNAs during viral

infections [22, 23]. To evaluate the performance of vsRNAfinder in the identification

of other vsRNA classes, 11 experimentally-verified siRNAs in Sugarcane mosaic

virus (SCMV) [22] and 6 experimentally-verified piRNAs in Dengue virus (DENV)

[23] were collected from literature. All SCMV siRNAs and four of six DENV

piRNAs were identified correctly by vsRNAfinder (Table S4). The remaining two

DENV piRNAs were missed by vsRNAfinder due to low abundance. For comparison,

ShortStack only identified correctly one SCMV siRNA and two DENV piRNAs

(Table S4), while miRDeep2 can only be used to identify the miRNA.

Systematic identification and characterization of vsRNAs

Then, the vsRNAfinder was used to systematically identify vsRNAs from

viral-infection-related sRNA-Seq data in public databases. A total of 19,734

high-confidence vsRNAs were detected from 878 samples in 102 sRNA-Seq datasets

by vsRNAfinder (Table S5). Among them, 2,746 vsRNAs were further annotated as

miRNAs (Table S5&6). The vsRNAs belonged to 64 viral species from 33 viral

families of seven viral groups (Figure 3A). Over 50% (11,801) of vsRNAs were

detected from the positive-sense single-stranded RNA (+ssRNA) viruses, and 3,524

(17.9%) vsRNAs were detected from double-stranded DNA (dsDNA) viruses. The

number of vsRNAs identified in viruses varied much and ranged from 1 to 2,489 with

a median of 170. The Invertebrate iridescent virus 6 (IIV-6, 2,489) contained most

vsRNAs among all viruses, followed by the Orsay virus (OrV, 1,142), the Cucumber

mosaic virus (CMV, 1,085), the Polygonum ringspot orthotospovirus (POLRSV, 972),

the Sweet potato feathery mottle virus (SPFMV, 949), and so on.

Then, the length, expression and genomic locations of vsRNAs were analyzed. The

lengths of vsRNAs showed a unimodal distribution which peaked in 21 and 22 nt

(Figure 3B). The vsRNAs with 21 nt accounted for 43% of all vsRNAs, followed by

those of 22 nt (17.5%) and 23 nt (7.0%). The expression level of vsRNAs followed an

approximate normal distribution (Figure 3C). Over 80% of vsRNAs had more than 20

RPM. The viral miRNAs had similar expression levels with that of other vsRNAs. To

analyze the location preference of vsRNAs along the viral genome, the number of

vsRNAs located within each 10% of the viral genomes was calculated. As shown in

Figure 3D, a slightly higher number of viral miRNAs and other vsRNAs were

observed in both ends of the viral genome than those in other genomic regions. Taking

the Zika virus as an example, there was 12-13 vsRNAs per Kb in the region around

the 10 Kb of the viral genome (near the 3'-end) which were much larger than that

observed in other genomic regions (less than 9 vsRNAs per Kb) (Figure 3E).

Figure 3. Data distribution of vsRNAs.

(A) The number of vsRNAs detected in each virus. The full name of each virus corresponding to the abbreviations of that was shown in Table S2.

(B) and (C) referred to the distribution of length and expression level of vsRNAs. RPM, Reads Per Million. The red dashed line in the (C) referred to 20 RPM.

(D) The number of vsRNAs located within each 10% of the viral genomes.

(E) The number of vsRNAs identified in a 1 Kb sliding-window with a step size of 100 bases along the genome of the Zika virus.

The vsRNAs were detected in viruses of 34 hosts from three kingdoms, including 20, 13 and 1 organism(s) in metazoa (animal), viridiplantae (plant) and fungi, respectively.

Therefore, the vsRNAs from viruses of different host kingdoms were analyzed.

Comparing the number of vsRNAs identified in viruses of different hosts from three

kingdoms showed that viruses from viridiplantae and fungi had much more vsRNAs

than those from metazoa (Figure 4A). Attention to note, only one virus

(Colletotrichum higginsianum non-segmented dsRNA virus 1, ChNRV1) which

infected the fungi was used in the study, although it had 774 vsRNAs identified.

Comparing the expression level of vsRNAs in viruses of different hosts from three

kingdoms showed that the expression level of vsRNAs from viruses of viridiplantae

was similar to that of vsRNAs from viruses of fungi; both of them were higher than

that of vsRNAs from viruses of metazoa (Figure 4B). The length distribution of

vsRNAs in viruses of metazoa was similar to that of viruses of viridiplantae (Figure

3C). Both of them had a sharp peak in 21 nt. While for viruses of fungi, the vsRNAs

had gentle peaks ranging from 19 to 23 nt.

Figure 4. Analysis and comparison of vsRNAs by the viral host.

(A) The number of vsRNAs identified in viruses of different hosts from three kingdoms;

(B) and (C) referred to the distributions of the expression level and length of vsRNAs in viruses of different hosts from three kingdoms.

Dynamic expression of vsRNAs after viral infection

Then, the dynamic expression of vsRNAs after viral infection was analyzed in 17 viruses (Table S7). In most viruses, the number of vsRNAs increased as the viral infection persisted. Taking the African swine fever virus (ASFV) and influenza A virus (IAV) as examples, 0, 13 and 111 vsRNAs were detected at 6, 12 and 24 hours

post infection (hpi) for the ASFV, while for the IAV increased expression of vsRNAs

was observed from 24 to 48 and 120 hpi (Figure 5A). Some viruses showed complex

dynamics of vsRNA expressions after infections. For example, the number of vsRNAs

from Sweet potato feathery mottle virus (SPFMV) were 133 and 157 at 3 and 6 weeks

post□infection, respectively, while that decreased to 103 at 12 weeks post□infection.

Figure 5. Analysis of the expression dynamics and specificity of vsRNAs.

(A) The number of vsRNAs detected in different time points after infection of African

swine fever virus (ASFV), Influenza A virus (IAV) and Sweet potato feathery mottle

virus (SPFMV).

(B)-(D) Venn diagram showing the overlap of vsRNAs detected in a given virus

infecting different hosts (B) or cell/tissue types (C), or of different strains (D).

Expression specificity of vsRNAs by the host, cell/tissue type and strain

The expression specificity of vsRNAs by the host was analyzed for four plant viruses

that infected the same kind of tissue (leaves) of different hosts. For a given virus, the

vsRNAs identified when infecting different hosts were analyzed and compared (Table

S8). For all four viruses, only a small proportion of vsRNAs overlapped between

viruses infecting different hosts. For example, 1,085 vsRNAs were detected when the

Cucumber mosaic virus (CMV) infected *Arabidopsis thaliana*, while no vsRNA was

detected when the virus infected *Solanum lycopersicum* (Figure 5B).

The expression specificity of vsRNAs by cell/tissue was analyzed for 11 viruses that

infected two or more cells/tissues of the same host (Table S9). No or little overlap was

observed between vsRNAs identified in different cells/tissues for most of these

viruses. For example, the MuHV-4 had 7, 25 and 51 vsRNAs when infecting NIH3T3

cells, S11 cells, NIH3T12 cells, respectively, while only 2 vsRNAs were observed in

all three cells (Figure 5C).

The expression specificity of vsRNAs by viral strain was analyzed for five viruses of

which different viral strains infected the same cell or tissue (Table S10). For all five

viruses, no or little overlap was observed between vsRNAs identified in different viral

strains of the same virus. Notably, although the viral strains Enterovirus A71 (EV-A17)

and Coxsackievirus A16 (CV-A16) of Enterovirus A (EV-A) shared approximately 80%

identity in the genome, only one vsRNA overlapped between 89 and 104 vsRNAs

identified from EV-A17 and CV-A16, respectively (Figure 5D).

Conservation analysis of vsRNAs

Then, the sequence similarity of vsRNAs among different virus species was

investigated using BLAST (version 2.11.0+) [32]. The vsRNAs of a given virus

species were queried against those in other viruses. A pair of vsRNAs was considered

to be similar if both the identity and coverage were greater than 90% in the reciprocal

BLAST analysis. Only 37 pairs of similar vsRNAs with each vsRNA in viruses of the

same or different viral families were observed in the analysis (Table S11). For

example, 551 and 774 vsRNAs were identified in Potato virus Y and Soybean mosaic

virus of the *Potyviridae* family, respectively, while only three similar vsRNA pairs

were observed between the two viruses. Next, the conservation of vsRNAs in

different viral strains of a given virus was analyzed. A vsRNA was considered as

"conserved vsRNA" if it was encoded in at least two viral strains of the same virus;

otherwise, the "non-conserved vsRNA". Unexpectedly, only 283 conserved vsRNAs

were observed among 6,949 vsRNAs in 12 viruses with different viral strains (Figure

S1). Only less than five conserved vsRNAs were observed in 8 of 12 viruses. For the

other four viruses including Cucumber mosaic virus (CMV), Human

gammaherpesvirus 4 (HHV-4), Invertebrate iridescent virus 6 (IIV-6) and Soybean

mosaic virus (SbMV), the conserved vsRNAs had significantly higher expressions

than non-conserved vsRNAs in viruses of HHV-4, IIV-6 and SbMV (Figure S1). For

example, the conserved vsRNAs had a median of 259 RPM in the HHV-4, while the

non-conserved vsRNAs had a median of 89 RPM.

Performance analysis of vsRNAfinder in identifying sRNAs in plants and animals

To investigate the ability of vsRNAfinder in identifying sRNAs in other organisms

besides viruses, the vsRNAfinder was used to identify sRNAs in *Homo sapiens* and

four model organisms including *Mus musculus*, *Drosophila melanogaster*,

Arabidopsis thaliana and *Oryza sativa*. The ability of vsRNAfinder was evaluated

based on its performance in identifying miRNAs in these organisms. For comparison,

miRDeep2 and ShortStack were also used to identify miRNAs in these organisms. As

shown in Table 1, vsRNAfinder had Recalls ranging from 0.06 to 0.13 in humans and

two animals which were a little lower than those of miRDeep2, while it had Recalls of

0.05 to 0.11 in two plants which were a little higher than those of miRDeep2. When

compared to ShortStack, vsRNAfinder consistently had higher Recalls in five

organisms. In terms of Precision, the ShortStack had the highest Precision among the

three methods in five organisms. Although the Precision of vsRNAfinder was lower

than that of ShortStack, it was consistently higher than that of miRDeep2 in five

organisms. Overall, in terms of F1-score which combined both the Recall and

Precision, the vsRNAfinder had an average F1-score of 0.122 which was similar to

that of miRDeep2 (0.116) and was slightly higher than that of ShortStack (0.098).

Table 1. Performance comparison of three computational methods in identifying

miRNAs in five organisms. Rec, Recall; Prec, Precision; F1, F1-score.

Organism	Method	Num of detected miRNA	Num of miRNAs in miRBase	Overlap of miRNA detected and those in miRBase	Rec	Prec	F1
Homo sapiens	vsRNAfinder	479		157	0.06	0.33	0.1
	miRDeep2	924	2632	213	0.08	0.23	0.12
	ShortStack	143		101	0.04	0.71	0.07
Mus musculus	vsRNAfinder	341		145	0.07	0.43	0.13
	miRDeep2	524	1963	177	0.09	0.34	0.14
	ShortStack	108		72	0.04	0.67	0.07
Drosophila melanogaster	vsRNAfinder	100		61	0.13	0.61	0.21
	miRDeep2	125	468	69	0.15	0.55	0.23
	ShortStack	51		37	0.08	0.73	0.14
Arabidopsis thaliana	vsRNAfinder	433		17	0.05	0.04	0.04
	miRDeep2	530	350	12	0.03	0.02	0.03
	ShortStack	14		9	0.03	0.64	0.05
Oryza sativa	vsRNAfinder	403		63	0.11	0.16	0.13
	miRDeep2	316	577	28	0.05	0.09	0.06
	ShortStack	116		55	0.1	0.47	0.16

Discussion

In this study, we presented vsRNAfinder, a novel method to identify and quantify

vsRNAs *de novo* based on sRNA-Seq data. The vsRNAfinder was found to

outperform two widely-used methods (miRDeep2 [18] and ShortStack [19]) in

identifying vsRNAs, and it could also be used to identify sRNAs in multiple

organisms of plants and animals with similar performances to current methods.

Besides, 19,734 high-confidence vsRNAs in over 60 viral species were identified by

vsRNAfinder and were further characterized, which could greatly facilitate further

studies of vsRNAs.

Numerous methods have been developed to identify sRNAs from sRNA-Seq data.

However, there is still a lack of an effective method for sRNA identification in viruses.

The *de novo* methods such as miRDeep2 and ShortStack identified sRNAs based on

reads assembly, which may merge reads from multiple adjacent sRNAs with overlaps.

This became worse in viruses on whose genome the sRNAs were more densely

distributed than those in other organisms. Hence, many vsRNAs may be missed by

these methods. The vsRNAfinder presented in the study is also a *de novo* method and

outperformed miRDeep2 and ShortStack in capturing viral miRNAs with improved

Recalls. This may be explained by the following two reasons. Firstly, smoothing the

coverage of mapped reads on the viral genome before the peak calling could reduce

the noise of the peak detection and enhance the peak signal. Secondly, filtering

sRNAs by abundance and evaluating the statistical significance of vsRNAs based on

the Poisson distribution could help identify high-confidence vsRNAs. Therefore, it is

suggested to put vsRNAfinder on priority in the identification of vsRNAs.

Most methods have been developed for the identification of specific sRNA class such

as miRNA, or the identification of sRNAs in a specific organism such as humans [18,

30]. Compared to previous methods, vsRNAfinder could be used to identify multiple

classes of sRNAs including miRNA, piRNA and siRNA. Besides viruses,

vsRNAfinder also achieved similar performances to current methods in the

identification of sRNAs across several animal and plant organisms. This suggested

that vsRNAfinder could be taken as a universal method for the identification of

sRNAs, especially for the vsRNAs.

An increasing number of vsRNAs have been reported in recent years [8, 11]. However,

most studies focused on vsRNAs from one or a few viruses [2, 13, 14, 23]. This work

for the first time systematically identified and characterized vsRNAs from over 60

viral species. Most vsRNAs were detected from +ssRNA viruses, followed by dsDNA

viruses. The ability of encoding sRNAs varied much among viruses. This was

consistent with previous studies which showed large differences between viruses in

their ability of encoding circRNAs [33]. It may reflect the large heterogeneity of

viruses in producing RNA products. The plant viruses were found to express more

vsRNAs than animal viruses. Besides, the abundance of vsRNAs in plant viruses was

also higher than that of animal viruses, although the length distribution of vsRNAs

was similar between plant and animal viruses. This may be because sRNAs in plants

are modified at their 3'-end by 2'-O-methylation which confers stability and

protection from degradation, while the animals lack the mechanism [34]. The length

of vsRNAs mainly focused on 21 and 22 nt which was the length of miRNAs and

siRNAs. However, only 14% of vsRNAs were annotated as miRNAs. Unexpectedly,

these miRNAs were extensively observed in RNA viruses that were reported to

replicate in the cytoplasm and have no access to the miRNA biogenesis machinery

located in the nucleus [35]. These miRNAs may be generated by the flexible or

noncanonical pathways of miRNA biogenesis. For example, several viral miRNAs

observed in infection of the Ebola virus (a negative-sense RNA virus) in humans were

likely to be processed by the cytoplasmic Drosha [35]. Further studies are needed to

verify the existence of viral miRNAs in RNA viruses and to clarify the mechanism of

their biogenesis.

Strong expression specificity was observed for viral RNAs such as circRNAs [33].

Consistent with previous studies, we also found strong expression specificity of

vsRNAs by the host, cell/tissue and viral strain [33]. For most viruses analyzed, no or

little overlap was observed between vsRNAs identified in a given virus of different

strains, or a virus infecting different hosts or cells/tissues. Besides, the dynamic

expression of vsRNAs in different time points after viral infections was also observed.

This suggested that the biogenesis of vsRNAs was highly specific and dynamic. It

may be related to the complex and dynamic interactions between viruses and hosts.

Viruses may use vsRNAs to fight against the host immunity and regulate the

transcription and translation of both virus and host. For example, the Rice stripe virus

was reported to generate lots of host-specific viral siRNAs in rice and planthopper

which potentially control defensive responses in hosts [36].

There are some limitations to the study. Firstly, only miRNAs were annotated in

vsRNAfinder as there were few effective methods of annotating other vsRNA types

such as piRNA and siRNA. For example, viral piRNAs were reported to lack the

characteristic ping-pong signature that is common in piRNAs of other organisms,

which makes it difficult to distinguish between viral piRNA and siRNA [37, 38].

More computational methods are needed to annotate vsRNAs. Secondly, the reads

that mapped to more than two locations of the genome were excluded to reduce false

positives in vsRNAfinder, which may lower the abundance of vsRNAs. Fortunately,

only 0.007% of reads were mapped to more than two locations in the viral genomes as

the viral genomes generally have a low level of sequence repeating. Thirdly, although

lots of high-confidence vsRNAs were identified in over 60 viral species, the list of

vsRNAs encoded by these viruses is far from complete considering the expression

specificity and dynamics of vsRNAs. Further efforts are still needed to identify more

vsRNAs, and to verify and characterize the vsRNAs identified.

Conclusion

Overall, this study developed a state-of-the-art method named vsRNAfinder which

can not only be used for identifying high-confidence sRNAs in viruses with better

performances than current methods, but also be used for sRNA identification in plants

and animals. Besides, a total of 19,734 high-confidence vsRNAs identified in 64 viral

species provided a rich resource for further studies of vsRNAs.

Supplementary information

540 Additional file 1. Supplementary figure 1 and table S1-S4, S6-S11.

541 Additional file 2. Supplementary table S5.

Availability of data and materials

544 All data used in the study are available in supplementary materials. The vsRNAfinder is

545 available at https://github.com/ZenaCai/vsRNAfinder.

Authors' contributions

548 Zena Cai and Ping Fu contributed equally to this work. Zena Cai developed the method,

549 analyzed the data and wrote the draft; Ping Fu analyzed the data; Ye Qiu, Aiping Wu, Gaihua

550 Zhang, Yirong Wang, Taijiao Jiang, Xing-Yi Ge, Haizhen Zhu and Yousong Peng supervised

551 the project, reviewed and revised the manuscript; Yousong Peng designed the project, wrote

552 and revised the manuscript.

Funding
This work was supported by the National Natural Science Foundation of China (32170651)

and Hunan Provincial Natural Science Foundation of China (2020JJ3006).

Acknowledgements

We thank people in the PengLab for helpful discussions on the manuscript.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1.   Romano G, Veneziano D, Acunzo M, Croce CM: **Small non-coding RNA and**

cancer. *Carcinogenesis* 2017, **38:**485-491.

2.   Liu C, Li F, Sun Y, Zhang X, Yuan J, Yang H, Xiang J: **Virus-derived small**

RNAs in the penaeid shrimp Fenneropenaeus chinensis during acute

infection of the DNA virus WSSV. *Sci Rep* 2016, **6:**28678.

3.   Wahid F, Shehzad A, Khan T, Kim YY: **MicroRNAs: synthesis, mechanism,**

function, and recent clinical trials. *Biochim Biophys Acta* 2010,

1803:1231-1243.

4.   Wilson RC, Doudna JA: **Molecular mechanisms of RNA interference.** *Annu*

Rev Biophys 2013, **42:**217-239.

5.   Neeb ZT, Zahler AM: **An expanding world of small RNAs.** *Dev Cell* 2014,

28:111-112.

6.   Jeang KT: **RNAi in the regulation of mammalian viral infections.** *BMC*

Biol 2012, **10:**58.

7.   Acunzo M, Visone R, Romano G, Veronese A, Lovat F, Palmieri D, Bottoni A,

Garofalo M, Gasparini P, Condorelli G, et al: **miR-130a targets MET and**

induces TRAIL-sensitivity in NSCLC by downregulating miR-221 and

222. *Oncogene* 2012, **31:**634-642.

8.   Kozomara A, Birgaoanu M, Griffiths-Jones S: **miRBase: from microRNA**

sequences to function. *Nucleic Acids Res* 2019, **47:**D155-d162.

9.   Rahman RU, Liebhoff AM, Bansal V, Fiosins M, Rajput A, Sattar A,

Magruder DS, Madan S, Sun T, Gautam A, et al: **SEAweb: the small RNA**

Expression Atlas web application. *Nucleic Acids Res* 2020, **48:**D204-d219.

10.   Feng L, Zhang F, Zhang H, Zhao Y, Meyers BC, Zhai J: **An Online Database**

for Exploring Over 2,000 Arabidopsis Small RNA Libraries. *Plant Physiol*

2020, **182:**685-691.

11.   Kincaid RP, Sullivan CS: **Virus-encoded microRNAs: an overview and a**

look to the future. *PLoS Pathog* 2012, **8:**e1003018.

12.   Weng KF, Hung CT, Hsieh PT, Li ML, Chen GW, Kung YA, Huang PN, Kuo

RL, Chen LL, Lin JY, et al: **A cytoplasmic RNA virus generates functional**

viral small RNAs and regulates viral IRES activity in mammalian cells. *Nucleic Acids Res* 2014, **42:**12789-12805.

13. Hussain M, Asgari S: **MicroRNA-like viral small RNA from Dengue virus 2 autoregulates its replication in mosquito cells.** *Proc Natl Acad Sci U S A* 2014, **111:**2746-2751.

14. Morales L, Oliveros JC, Fernandez-Delgado R, tenOever BR, Enjuanes L, Sola I: **SARS-CoV-Encoded Small RNAs Contribute to Infection-Associated Lung Pathology.** *Cell Host Microbe* 2017, **21:**344-355.

15. Aparicio-Puerta E, Lebrón R, Rueda A, Gómez-Martín C, Giannoukakos S, Jaspez D, Medina JM, Zubkovic A, Jurak I, Fromm B, et al: **sRNAbench and sRNAtoolbox 2019: intuitive fast small RNA profiling and differential expression.** *Nucleic Acids Res* 2019, **47:**W530-w535.

16. Handzlik JE, Tastsoglou S, Vlachos IS, Hatzigeorgiou AG: **Manatee: detection and quantification of small non-coding RNAs from next-generation sequencing data.** *Sci Rep* 2020, **10:**705.

17. Wu X, Kim TK, Baxter D, Scherler K, Gordon A, Fong O, Etheridge A, Galas DJ, Wang K: **sRNAnalyzer-a flexible and customizable small RNA sequencing data analysis pipeline.** *Nucleic Acids Res* 2017, **45:**12140-12151.

18. Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N: **miRDeep2 accurately identifies known and hundreds of novel microRNA genes in seven animal clades.** *Nucleic Acids Res* 2012, **40:**37-52.

19. Shahid S, Axtell MJ: **Identification and annotation of small RNA genes**

622  **using ShortStack.** *Methods* 2014, **67:**20-27.

623 20. Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M,

624  Marshall KA, Phillippy KH, Sherman PM, Holko M, et al: **NCBI GEO:**

625  **archive for functional genomics data sets--update.** *Nucleic Acids Res* 2013,

626  **41:**D991-995.

627 21. Kodama Y, Shumway M, Leinonen R: **The Sequence Read Archive:**

628  **explosive growth of sequencing data.** *Nucleic Acids Res* 2012, **40:**D54-56.

629 22. Xia Z, Peng J, Li Y, Chen L, Li S, Zhou T, Fan Z: **Characterization of small**

630  **interfering RNAs derived from Sugarcane mosaic virus in infected maize**

631  **plants by deep sequencing.** *PLoS One* 2014, **9:**e97013.

632 23. Wang Y, Jin B, Liu P, Li J, Chen X, Gu J: **piRNA Profiling of Dengue Virus**

633  **Type 2-Infected Asian Tiger Mosquito and Midgut Tissues.** *Viruses* 2018,

634  **10**.

635 24. Martin M: **Cutadapt removes adapter sequences from high-throughput**

636  **sequencing reads.** *2011* 2011, **17:**3 %J EMBnet.journal.

637 25. Langmead B, Trapnell C, Pop M, Salzberg SL: **Ultrafast and**

638  **memory-efficient alignment of short DNA sequences to the human**

639  **genome.** *Genome Biol* 2009, **10:**R25.

640 26. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO,

641  Whitwham A, Keane T, McCarthy SA, Davies RM, Li H: **Twelve years of**

642  **SAMtools and BCFtools.** *Gigascience* 2021, **10**.

643 27. Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing**

genomic features. *Bioinformatics* 2010, **26:**841-842.

28.   Borchers HW: **pracma: Practical Numerical Math Functions.** 2022.

29.   Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson

A, Szcześniak MW, Gaffney DJ, Elo LL, Zhang X, Mortazavi A: **A survey of**

best practices for RNA-seq data analysis. *Genome Biol* 2016, **17:**13.

30.   Hackenberg M, Rodríguez-Ezpeleta N, Aransay AM: **miRanalyzer: an**

update on the detection and analysis of microRNAs in high-throughput

sequencing experiments. *Nucleic Acids Res* 2011, **39:**W132-138.

31.   Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler

PF, Hofacker IL: **ViennaRNA Package 2.0.** *Algorithms Mol Biol* 2011, **6:**26.

32.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment**

search tool. *J Mol Biol* 1990, **215:**403-410.

33.   Cai Z, Fan Y, Zhang Z, Lu C, Zhu Z, Jiang T, Shan T, Peng Y: **VirusCircBase:**

a database of virus circular RNAs. *Brief Bioinform* 2021, **22:**2182-2190.

34.   Borges F, Martienssen RA: **The expanding world of small RNAs in plants.**

Nat Rev Mol Cell Biol 2015, **16:**727-741.

35.   Nanbo A, Furuyama W, Lin Z: **RNA Virus-Encoded miRNAs: Current**

Insights and Future Challenges. *Front Microbiol* 2021, **12:**679210.

36.   Yang M, Xu Z, Zhao W, Liu Q, Li Q, Lu L, Liu R, Zhang X, Cui F: **Rice**

stripe virus-derived siRNAs play different regulatory roles in rice and in

the insect vector Laodelphax striatellus. *BMC Plant Biol* 2018, **18:**219.

37.   Varjak M, Donald CL, Mottram TJ, Sreenu VB, Merits A, Maringer K,

Schnettler E, Kohl A: **Characterization of the Zika virus induced small**

RNA response in Aedes aegypti cells. *PLoS Negl Trop Dis* 2017,

11:e0006010.

38.   Öhlund P, Hayer J, Hesson JC, Blomström AL: **Small RNA Response to**

Infection of the Insect-Specific Lammi Virus and Hanko Virus in an

Aedes albopictus Cell Line. *Viruses* 2021, **13**.