Neural mechanisms for the localization of externally generated tactile motion

Suma Chinta and *Scott R. Pluta, Department of Biology, Purdue University, West Lafayette, IN

Abstract

During tactile localization, animals must differentiate stimuli caused by their own volitional movement from externally generated object motion. To determine a neural basis for this ability, we examined the mouse superior colliculus (SC), which contains multiple egocentric maps of sensorimotor space. By placing mice in a whisker-guided virtual reality, we discovered a rapidly adapting neural response that strongly preferred external over self-generated changes in tactile space. This transient response only emerged when external motion gained contact with a whisker, arguing that stimulus adaptation was whisker-specific. The accumulation of sensory evidence through active sensing and repetitions in external motion controlled the size of the transient response. Population-level firing rates among transiently responsive neurons accurately encoded the direction of external motion. These data reveal that stimulus-specific adaptation together with accumulating sensorimotor predictions in SC neurons enhance the localization of unexpected motion in the environment.

Introduction

Much of our interaction with the world occurs through the volitional movement of our sensory organs. For example, primates move their eyes to visualize a scene, and rodents move their whiskers to explore nearby objects. To accurately localize objects, animals must differentiate the expected consequences of their actions, induced by active sensing, from real-world object motion. What neural mechanisms underlie this ability?

To address this question, we focused on the midbrain superior colliculus (SC), which contains multiple egocentric maps of sensorimotor space [1–4]. Many SC mediated behaviors involve spatial processing, such as pursuing prey, escaping predators, or simply orienting towards the appropriate object [5–11]. While visually tracking objects, the SC helps stabilize the visual field and adjust for discrepancies between the predicted and actual visual outcome of eye movements [12,13]. Given this insight, we hypothesized that SC neurons are specialized for encoding environmental motion during active sensing. Such computations are likely critical for localizing moving objects such as prey [14–17].

While decades of research have outlined SC function during visuospatial processing, remarkably less is known for somatosensation, particularly during active touch [18–21]. The mouse SC is known to respond to whisker stimuli, yet nearly all published work in this area has been

Figure 1. Tactile and neural dynamics during whisker guided virtual reality
A) Schematic of whisker guided virtual navigation, illustrating a head-fixed mouse locomoting on a treadmill and whisking against a surface. After trial initiation, there was an equal probability for the tactile surface to translate into either rostral (in red) or caudal (in blue) whisker space or stay at the center position (in black). Upper right: single-whisker location density while the surface was at the rostral or caudal location. Bottom right: whisker position (mean ± s.e.m) and surface location (transparent line) during rostral (top in red) and caudal (bottom in blue) translations. The first half is surface excursion while second half is the return to the center position. **B**) Trial structure aligning surface position, whisker position, and locomotion speed. **C**) Trial-averaged whisker position and firing rate of two example SC neurons during each type of surface translation (50ms bins). **D**) Top: Percentage of tactile responsive SC neurons (blue) as a function of depth from the SC surface (12 mice, 873 neurons). Bottom: dye labeling of the electrode shanks in the intermediate and deep layers of the lateral SC.

33
performed under anesthesia [4,22–25]. These experiments reveal that the SC contains a

somatotopic map of whisker space, whereby individual neurons possess large (multi-whisker)

receptive fields that are also sensitive to artificial whisking. Therefore, active whisking in the

mouse SC provides a tractable and underutilized model for revealing the cellular and circuit

mechanisms that support innate tactile-guided behaviors.

Sensory responses in the SC have primarily been characterized during sensor fixation, with

evidence showing that differences in egocentric heading (eye position) modulate response

magnitude [18,19,21,26–28]. This approach reveals that sensory responses in the SC are modulated

by extrasensory inputs. Additional work has shown that eye movements in darkness suppress

SC spiking, arguing for the presence of a corollary discharge to suppress self-induced visual

stimulation [29,30]. The somatosensory whisker system offers a powerful model for investigating

the integration of external and self-generated stimuli, since mice instinctively move their

whiskers to palpate objects, and the precise dynamics of touch can be accurately measured

using high-speed imaging and markerless tracking [31,32].

To reveal the neural mechanisms for the active localization of external motion, we designed a

tactile virtual reality that simulates whisker-guided navigation. While running on a treadmill, mice

rhythmically touched a cylindrical surface that rotated at the same velocity as their locomotion.

This created a tactile flow-field that simulates running along a wall, in a manner similar to visual

flow [33–35]. Periodically, the center axis of the flow-field translated horizontally along the mouse's

whisking field. A transient neural response only emerged when surface movements entered a

location that gained or lost contact with a whisker. Self-generated gains in whisker contact that

occurred independent of surface movement generated relatively weak tactile responses. The

size of the tactile response was controlled by stimulus expectations, and the direction of surface

movement was accurately encoded in neuronal firing rates.

Results

Simulating whisker-guided virtual reality

We designed a closed-loop tactile system that simulates whisker-guided exploration in mice.

During the experiment, a head-fixed running mouse voluntarily whisked against a nearby

cylindrical surface that rotated at the same velocity as its locomotion (**Fig. 1A, left**). A high-

speed (500fps) infrared camera and digital encoders recorded whisker kinematics, surface

movements, and running speed while an opaque object and white noise obscured visual and

auditory cues. After the mouse initiated a trial by running a certain distance, the surface would

either translate rostrally, caudally or remain at the starting location with equal probability. After

staying at the rostral/caudal location for an equivalent locomotor distance (if applicable), the

Figure 2. Surface movements that gained or lost contact with whiskers elicited transient and sustained changes in spike rate

A) Surface location, whisker angle, whisker curvature derivative, and the firing rates of two example neurons during externally generated gains (black) and losses (blue) in whisker touch. The vertical dashed line marks the estimated moment of gain/loss in contact. **B**) Population firing rates from an example mouse during surface movements. A gain/loss in whisker contact only occurred during movement through caudal space (46 neurons). **C**) Top, two example neurons illustrating the distinct transient and sustained response components evoked during surface translation. Bottom, scatter plot comparing the transient and sustained firing rates of SC neurons (10 mice, 529 neurons). **D**) Scatter plot comparing the GoW and LoW modulation of each neuron (10 mice, 529 neurons). **E**) Population averaged firing rates of neurons grouped by the sign (+/-) of their responses to GoW and LoW (8 mice, 385 neurons). **F)** Population average firing rate of neurons response to no GoW or LoW external surface movement. All population average firing rates are binned at 50ms.

surface would then return to the center location. While running and touching the surface, mice

innately adapted the position of their whiskers to track changes in surface location (**Fig. 1A,**

right). Since translation timing was determined by locomotor distance, variation in locomotion

speed made translation onset unpredictable (**Fig. 1B**). As the surface moved into a new

location, the firing rate of most neurons rapidly increased while others decreased, as shown in

two example units (**Fig. 1C**). A 3-shank, 128-channel silicon probe recorded neural activity

simultaneously across the intermediate and deep layers of the lateral SC, approximately 300 –

1000 microns below its surface (**Fig. 1D**, 12 mice, 873 neurons). About two-thirds of all

recorded neurons displayed a significant response to surface movement (67 ± 6%, 12 mice, 578

neurons, $\alpha < 0.05$, 1-way ANOVA).

Surface movements that gain or lose whisker contact elicit transient and sustained

changes in firing rate

To reveal the stimulus variables driving SC responses, we examined the dynamics of whisker

touch during surface movement. We performed a combination of computerized whisker

kinematic analysis and manual inspection of high-speed video. We discovered that surface

movements elicited large changes in firing rate only when they entered a location that gained

(GoW) or lost (LoW) contact with a whisker (**Fig. 2A, B, Suppl. Fig. 1A**). The addition or

subtraction of a single whisker from the surface was sufficient for this effect. Most neurons

preferentially responded to whisker addition (GoW, 55 ± 4%, 8 mice, 385 neurons), while only a

small percentage of neurons preferred surface movements that engaged the same whiskers at

both locations (15 ± 3%, 8 mice, 385 neurons, **Supp. Fig. 1A**). Gains in whisker contact with the

surface elicited transient and/or sustained changes in neuronal firing rate (**Fig. 2A, C top**).

Transient responses typically returned to baseline within 300 ms following their onset, while

sustained responses exhibited firing rates different than baseline for the entire duration that the

surfaced contacted the new whisker(s) (see methods). In most cases, the transient response

was larger than the sustained response, yet there was almost continuous variation in their

relative magnitudes across the population (**Fig. 2C, bottom**). To confirm that these responses

dynamics were whisker-mediated, we repeated the experiment after trimming off the whiskers

and found that nearly all neurons became unresponsive to surface movement (59 ± 4% pre-

trimming vs. 6 ± 1% post-trimming, 156 neurons, 2 mice, 1-way ANOVA, $\alpha < 0.05$, **Supp. Fig.**

1B). To test if adaptation of the transient response was whisker specific, we picked trials where

two consecutive surface movements gained contact with different whiskers. With each

Figure 3. Externally generated gains in contact preferentially drive the transient response
A) Example trial illustrating the designation of external- and self-generated touch. Initially, the mouse freely whisked in air. After the mouse ran a set distance, the surface moved into its whisking field and contacted the whisker to create externally generated touch. After a period of whisking against the surface, the mouse voluntarily stopped before eventually resuming to create self-generated touch. **B**) Three neurons from one mouse comparing responses to external and self-generated touch. Rasters and histograms of spike timing aligned to touch onset. Only the first external and self-generated touch from each trial were used. **C**) Population-averaged, z-scored firing rates aligned to the onset of external and self-GoW (4 mice, 139 neurons). Firing rates for external and self-GoW plotted in E are extracted from the gray window. **D**) Population averaged, z-scored whisker curvature aligned to the onset of external and self-GoW. **E**) Scatter plot comparing change in neuronal firing rates for external and self-generated touch (4 mice, 139 neurons, p < 0.001, Wilcoxon signed rank test). Firing rates are subtracted with their corresponding baselines. **F**) Boxplots comparing the time mice spent without any touch before an external- or self-GoW stimulus occurred (p = 0.04, Mann Whitney U-test). All values are mean ± s.e.m.

101
consecutive gain in whisker contact, a transient response emerged from the adapted state

(**Supp. Fig 1C**).

To understand the receptive fields of SC neurons in the framework of whisker space, we
quantified each neuron's response to GoW and LoW stimulation (10 mice, 529 neurons, **Fig.**
2D). Almost half of all neurons increased their firing rate for both GoW and LoW (251/529
neurons) stimuli, while the remaining neurons increased their firing rate for one stimulus type yet
decreased it for the other. To illustrate these effects, we plotted the population averaged firing
rates of each response type relative to the onset of surface movement (**Fig. 2E**). Overall, these
data demonstrate that SC neurons were highly sensitive to gains/losses in whisker contact,
while showing relatively weak changes in firing rate during surface movements that sustained
contact with the same whiskers (**Fig. 2F**).

SC neurons strongly prefer externally generated gains in whisker contact

We hypothesized that the GoW response was more sensitive to external rather than self-
generated gains in whisker contact. To test this hypothesis, we modified our experiment so that
the surface periodically moved between rostral whisking space and a more distant location
entirely outside the whisking field. With the surface in rostral space, mice voluntarily lost and
gained contact (self-GoW) with the surface as they periodically (6.8 ± 0.6 second interval)
transitioned between resting (no running or touch) and locomotive touch (**Fig. 3A**). Mice had 1
to 5 intact whiskers contacting the surface in these experiments. Using this approach, we were
able to calculate neuronal firing rates relative to the onset of self-GoW as well as externally
generated gains in whisker contact (external-GoW).

In individual mice, neurons displayed a range of stimulus sensitivities (**Fig. 3B**). Some neurons
only responded to external-GoW stimuli, while most neurons responded to both stimulus types
but with a clear preference for external-GoW (**Fig. 3B, C**). A minority of neurons responded
equally to external- and self-GoW stimuli (**Fig. 3E**). Therefore, population level firing rates were
significantly greater during external-GoW stimulation ($p < 0.001$, Wilcoxon signed rank test, 4
mice, 139 neurons, **Fig. 3C, E**). The curvature of the whisker during external and self-GoW was
nearly identical, indicating that differences in stimulus strength are unlikely the cause of this
effect (**Fig. 3D, Suppl. Fig. 2A**). To rule out adaptation as the mechanism for the larger
external-GoW response, we compared the time mice spent without touch preceding each GoW
event. Our data indicate that adaptation cannot explain GoW response differences, as the time
spent without touch before GoW stimulation was similar in both conditions (Mann-Whitney, $p =$

Figure 4. Self-generated stimulus expectations control the size of the transient response
A) Schematic illustrating the experimental design. At the start of the trial, the mouse is running and whisking freely in air (see boxplot for quantification). After the mouse crossed a threshold locomotion distance and then stopped running, a touch surface entered its whisking field. The mouse voluntarily resumed whisking and generated an unexpected touch event. The mouse then voluntarily stopped running and whisking before it resumed active touch. After running and touching the surface for a set distance, the surface retracted, and the trial restarted. *Bottom*, histogram of firing rates for one neuron during expected and unexpected touch. **B)** Rasters and histograms of spike timing for three neurons from one mouse during expected and unexpected touch. **C)** Population averaged neuronal firing rates and whisking variables aligned to the onset of touch (2 mice, 74 neurons). **D)** Scatter plot comparing change in neuronal firing rates during expected and unexpected touch (2 mice, 74 neurons, p = 0.004, Wilcoxon signed rank test). **E)** Box plot comparing the time mice spent without any touch before an unexpected and expected touch event occurred (Mann Whitney U-test, p = 0.17).

0.04, **Fig. 3F, Suppl. Fig. 2B**). Therefore, the larger external-GoW response cannot be

explained by a slower stimulus repetition rate.

Self-generated sensory experience controls transient response magnitude

Although stimulus strength and repetition rate for external- and self-GoW stimuli were the same,

the locomotive states were not. In the external-GoW condition, the animal was already running,

yet in the self-GoW condition, the animal was transitioning from quiescence to locomotion.

Using a novel approach where both touch events started from quiescence, we controlled for

differences in locomotion speed and any potential differences in touch quality (**Fig. 4**).

Mice had one whisker contacting the surface in this experiment. A trial was initiated by the

animal running and freely whisking in air. After the mouse reached a threshold distance on the

treadmill, and then decided to stop running and whisking, an object was quickly (150 ms)

extended into its whisking field (**Fig. 4A**, see boxplot). Moments later, the animal voluntarily

resumed locomotive whisking and contacted the surface, causing an unexpected touch event.

The object stayed in the whisking field for an extended period, allowing the mouse to

periodically accumulate evidence about its presence by pausing and resuming active touch.

After each pause in locomotion and whisking, the first touch in the next locomotive whisking

bout generated an expected touch event. After the mouse ran and touched the surface for a set

treadmill distance, the surface retracted, and the trial restarted with the free whisking and

locomotion condition. On average, the first touch response that followed free whisking

(unexpected) was significantly greater than the first touch response that followed a bout of

active touch (expected, **Fig. 4B, C,** p = 0.004, n = 74, signed rank test). Importantly, touch

kinematics (whisker position and whisker curvature) were identical between both touch

conditions (**Fig. 4C**). Neurons with larger tactile responses were more strongly modulated by

stimulus expectation (**Fig. 4D**). Stimulus repetition rate was the same between the touch

conditions (**Fig. 4E,** p = 0.17, t-test). Therefore, stimulus expectations (whisking in air vs.

repeated surface contact) generated by active sensing controlled the magnitude of the transient

response, in agreement with our data in Figure 3.

Repetitions in external motion weaken the transient response

Next, we sought to determine if responses to external-GoW were static or habituated over time.

To do so, we analyzed the effect of stimulus repetition on response magnitude. For each

experiment, we selected a surface movement with GoW stimulation (**Fig. 5A**). To ensure that

any effects we observed were not caused by changes in tactile behavior, we analyzed the

Figure 5. Stimulus repetition number controls transient response magnitude
A) Schematic illustrating the different stimulus types and how they are randomly repeated over the course of the experiment. The return motion of the 2nd stimulus which is a GoW is selected for analysis in C. **B)** Left, trial-averaged whisker position relative to surface movement for early (first 1/3rd), middle, and late (last 1/3rd) trials of GoW stimulation (one mouse). Right, trial-averaged whisker position calculated during the transient response period and plotted as a function of stimulus repetition (8 mice). **C)** Rasters and histograms of spiking in two example neurons divided by early, middle, and late periods. Inset, spike waveforms during the same periods (color coded). **D)** Magnitude of the transient and sustained responses as a function of stimulus repetition in same two example neurons in **C**. Transient and sustained firing rates were obtained from transient and sustained windows described in methods. **E)** Population slopes of the linear regression of transient (8 mice, 223 neurons, $p < 0.001$, t-test) and sustained (8 mice, 98 neurons, $p = 0.64$, t-test) responses as a function of stimulus repetition. **F)** Normalized spike waveform amplitude as a function of stimulus repetition (8 mice).

167
position of the whisker as a function of stimulus repetition. Across the population (8 mice), there

was no notable change during the transient response period (**Fig. 5B**). However, we discovered

that the magnitude of the transient response linearly decayed with increasing GoW repetitions,

while the sustained response was stable, as shown in two example neurons (**Fig. 5C, D**). This

transient-specific habituation was consistent across the population (**Fig. 5E**, 223 transient

neurons, 8 mice, $p = 2e^{-12}$; 98 sustained neurons, $p = 0.64$, t-test). The stability of spike

waveform amplitude in our recordings indicates that a progressive decrease in spike detection

during sorting cannot explain these effects (**Fig. 5F, 5C insets**). Habituation of the transient

response was unlikely caused by low-level sensory adaptation, since the average interval

between each GoW stimulus was around one minute (67 ± 5 sec, 8 mice), and GoW stimuli

were randomly interleaved with other non-GoW surface movements. Therefore, SC neurons

appear to steadily habituate to repeated tactile stimulation occurring over the course of many

minutes, providing further evidence for the role of sensory experience in regulating the transient

response. The same transient-specific habituation was observed in our other approach detailed

in Figure 4 (**Suppl. Fig. 3**).

Figure 6. Neural decoding of surface location

A) Location decoding accuracy using all the tactile responsive neurons recorded in a single mouse (75 neurons). **B)** Decoding accuracy in the same mouse with sustained (n = 29) and transient (n =34) responsive neurons separated. **C)** Diagram illustrating an experiment where half of all the surface translations caused a gain/loss in whisker contact. **D)** Decoding accuracy averaged across animals with the same number of gains/losses in whisker contact (GoW/LoW in all movements: 2 mice, 81 neurons; GoW/LoW in half of all movements: 6 mice, 262 neurons; Zero GoW/LoW movements: 4 mice, 149 neurons). **E)** Decoding accuracy averaged across animals according to neuron type (sustain/transient) and changes in whisker contact (12 mice, 492 neurons).

183
Neurons in the superior colliculus encode surface location

To assess the function of external and self-generated tactile responses in the SC, we tested the

surface localization accuracy of population-level SC activity over time. Using population

decoding (see methods), we discovered that SC activity accurately classified surface location

(center, rostral, or caudal), with the highest accuracy during the movement period, as shown in

one example mouse (**Fig. 6A**, 99% accuracy, 76 neurons). Before the start of the excursion,

classification accuracy was at chance, indicating that neural activity could not predict the

upcoming surface movement. After excursion into rostral or caudal space, classification

accuracy for sustained neurons was high for as long as the surface remained at that location,

revealing a stable self-generated representation of location in this subset of neurons (**Fig. 6B**,

30 neurons, 1 mouse). After the surface returned to the center position, stimulus classification

remained above chance, revealing persistent information about the previous location of the

object. In contrast, classifier accuracy for transient neurons was high only during surface

movement, due to their rapid adaptation rates (46 neurons, 1 mouse). By segregating

experiments by the number of surface movements that gained/lost contact with a whisker, we

found that classifier accuracy was highest during GoW/LoW translations (**Fig. 6C, D**, GoW/LoW

in all movements: 94 ± 5% accuracy, 2 mice; GoW/LoW in half: 71 ± 7%, 6 mice; GoW/LoW in

no movements: 56 ± 3%, 4 mice). This trend was generally consistent for both sustained and

transient neurons (**Fig. 6E**), but transient neurons were least informative about surface

movements that did not gain or lose whisker contact. Since classifier accuracy was best at

distinguishing locations that interacted with different whiskers, a somatotopic map of whisker

space in the SC likely facilitates object localization. We found evidence for such map in one of

our recordings where the rostral and caudal stimuli gained contact with different whiskers

(**Suppl. Fig. 4**).

Discussion

Active sensing requires the brain to contextualize incoming sensory stimuli with outgoing motor

commands. To localize objects, animals must recognize if changes in egocentric space

originate from externally or internally generated movement [36]. By simulating   whisker-guided

exploration, we discovered neurons in the mouse SC that are specialized for locating

unexpected, externally generated changes in tactile space. We identified a rapidly adapting

transient response, which strongly preferred externally initiated touch and that habituated with

stimulus repetition. Transient responses have been observed in the SC during visual stimulation

[37–39] and passive, rhythmic deflection of the whiskers [25]. We observed a diversity of stimulus

selectivity across the population of recorded neurons, with some neurons only briefly

responding to externally generated touch and other neurons sustaining an equally strong

response to both external and self-generated touch. Interestingly, self-generated sensory

experience appeared to be a primary factor controlling transient response magnitude.

Both the barrel cortex and brainstem provide monosynaptic sensory drive to SC neurons
[22–24,40,41]. While the circuit mechanism for rapid sensory adaption is unknown[42], some layer 5
neurons in the barrel cortex also show rapid sensory adaption during surface whisking[33].
Brainstem derived synaptic potentials in SC neurons also rapidly adapt during 20 Hz
stimulation, which is similar to the natural whisk frequency of mice[22]. Therefore, sensory
adaptation in the SC could stem from local and/or long-range circuits. Relating the adaptation
rate of SC neurons to their downstream targets could provide valuable insight into the
mechanisms supporting SC-mediated movements and attention[10,43].

How motor and associative circuits in the SC contextualize somatosensory input remains
unknown. An efferent copy of whisker protraction, possibly involving the cerebellum, motor
cortex and brainstem, could be an important mechanism for regulating the transient response
[13,44–48]. In our study, stimulus expectations built by active whisking shape transient response
magnitude. Therefore, the magnitude of the transient response reflects the degree of mismatch
between the predicted and actual outcome of sensor movement, thereby providing a neural
mechanism to identify and correct for errors associated with sensory-guided movement[13].

Value-coded input to the SC could shape the transient response to generate a neural
mechanism for favoring novel or rewarding environmental cues [49–55]. Importantly, mice in our
study behaved voluntarily and were not seeking reward. In this free-form context, transient
response habituation may reflect a central mechanism for ignoring stimuli of diminishing novelty
[38,56]. Whether this mechanism relies on changes in inhibitory signaling from the substantia nigra,
the activity of local SC interneurons, or other upstream circuits remains unknown [6,52,57].
Habituation to repeated visual stimulation has been observed previously in primates and
rodents [38,58–60], yet our study is the first to demonstrate slower, accumulative effects of stimulus
repetition in the SC during active sensing. Since our stimuli repeated at long, one-minute
intervals over the course of an hour, SC neurons in our experiment were storing and updating
stimulus expectations for an extended period.

Neuronal selectivity for externally generated stimulation has been observed in other
sensory modalities and brain areas. Neurons in the monkey cerebellum selectively respond to
external but not self-generated vestibular accelerations [61]. Neurons in the monkey visual cortex
display a unique window of suppression only when voluntary eye-movements create visual
stimulation [62]. Neurons in the superficial visual cortex of mice selectively respond to mismatches
between visual flow and locomotion speed [63,64]. Neurons in the auditory brainstem cancel self-

generated sounds associated with licking [45], and neurons in the hindbrain of weakly electric fish

cancel self-generated electric signals [65]. We revealed a class of neurons in the superior

colliculus that preferentially respond to externally generated tactile stimuli. Since contact with a

new whisker was necessary for this transient response to emerge, the cancellation of self-

generated touch is specific to individual whiskers.

Transient responses may facilitate object localization. We show that their population

level firing rates briefly but accurately encode the location of external motion. The net effect of

this coding brevity is an accurate representation motion direction, but not its lasting location.

Decoding external motion was most accurate between movements that engaged different

whiskers. Therefore, the somatotopic map of whisker space in the SC may enable a population

level representation of object movement through different whiskers in the pad. The precise

timing of spikes generated by these gains in whisker contact, as demonstrated in our study,

could enable the rapid orienting movements associated with prey capture [17,66].

To localize objects during active touch, animals must differentiate self- from externally

generated changes in tactile space. Such computations are pervasive across different sensory

modalities and brain areas involved in sensorimotor processing. In the SC, response selectivity

for unexpected, externally generated touch is controlled by rapid stimulus adaptation and the

accumulation of external and self-generated stimulus expectations. These computations require

single neurons to contextualize stimulus information over milliseconds and hours. It is perhaps

surprising that single neurons in an ancient midbrain structure can multiplex this level of

information. Given the diversity and complexity of circuits that impinge upon the mammalian SC,

this may also be a reasonable expectation. How the SC interprets each of its diverse inputs to

construct the egocentric maps that guide behavior remains an exciting and important topic.

Methods

Experimental Model and Subject Details

Mice of a CD-1 background of both sexes, between the ages of 9-15 weeks were used for all

experiments. The Purdue Institutional Animal Care and Use Committee (IACUC,

1801001676A004) and the Laboratory Animal Program (LAP) approved all procedures. Mice

were socially house with five or less per cage and maintained in reverse light-dark cycle (12:12

hr.). All experiments were conducted during the animal's subjective night.

In-vivo Electrophysiology

Prior to behavioral conditioning, a custom aluminum headplate was attached to each mouse to

enable head fixation. Briefly, animals were anesthetized under 5% isoflurane and maintained at

~3% while monitoring body temperature and respiratory rate throughout the procedure. Artificial

tears ointment was applied to keep the eyes hydrated. The skin and fur over the skull were

disinfected with 70% ethanol followed by betadine and incised using sterilized surgical

instruments. A tissue adhesive (Liquivet) and dental cement (Metabond) was applied over the

skull and wound margins. The headplate was attached to the skull with dental cement. Lastly

buprenorphine (mg/kg) was administered as a postoperative analgesic. Two days after

headplate implantation, mice were placed on a circular treadmill for 1 hour per day for up to 10

days, or until they learned to run freely at a steady pace.

295   On the day of the electrophysiology recording, mice were briefly (15 – 20 minutes)

anesthetized to perform a craniotomy over the SC. A 1mm diameter craniotomy was made

using a biopsy punch (Robbins Instruments) and then covered with a silicone gel (Dowsil).

Several hours later, mice were placed in the experimental rig and a three-shank custom probe

(Neuronexus) of 128-channels was lowered into the brain using a micromanipulator (NewScale).

After exiting the ventral surface of the cortex, as evident by a loss of spiking, the electrode was

lowered at a rate of 75µm/min while constantly searching for activity driven by flashes of light.

The onset of visual activity was used to mark the depth of the SC surface (~1000µm below

cortical surface). The electrode was further lowered into the intermediate and deep layers,

where manual whisker deflections caused strong neuronal responses. The receptive field of

neurons were mapped by manually deflecting individual whiskers and locating the primary

drivers of neuronal activity. Whiskers that did not elicit detectable activity were trimmed.

Recordings were targeted towards the C-row and macrovibrissae. If the electrode penetration

missed the target, it was removed and re-inserted based on the coordinates of somatotopic

space. In most experiments, mice had 3 – 5 intact whiskers contacting the surface, which

spanned one or two rows. In other cases, mice had one or two whiskers intact.

Whisker guided virtual navigation

Head-fixed mice ran at their own volition (23.7 ± 1.6 cm/s, 12 mice) with concomitant rhythmic

whisking (19 ± 1 Hz, 10 mice). All data presented was collected during locomotion and whisking,

unless otherwise stated. The circular treadmill was attached to a digital encoder that controlled

the rotation of a tactile surface in a closed-loop. A trial began after 200 cm of locomotion on the

treadmill to ensure the mice were actively sensing the surface. Shortly (1.4 sec.) after whisker

imaging began, the surface translated 1 cm linear distance into rostral or caudal space or

remained at the center location. Each outcome was randomly chosen with equal (33.33%)

probability. In six mice, an additional location was added that was entirely outside the whisker

field, giving each surface movement a 25% probability. While at the rostral/caudal position, mice

had to run an additional 200 cm before the surface would return to the center. An opaque flag

over the eye and white background noise obscured visual and auditory cues, respectively.

In two mice, the tactile surface was replaced with a pneumatically controlled rectangular

surface. Prior to the experiment, all whiskers were trimmed except the principal whisker (B1 or

C1). A trial began after the mouse ran 200 cm on the treadmill while freely whisking in air with

no surface in the whisking field. After the mouse crossed the distance threshold and stopped

running, the touch surface was extended into the whisking field. The mouse voluntarily resumed

whisking and generated an unexpected touch event. The mouse then voluntarily stopped

whisking for a period before resuming to touch the same surface and generate an expected

touch event. After running 600 cm with the surface present, the surface retracted, and the trial

restarted.

Whisker Tracking & kinematics

Whiskers were tracked at 500fps during the trial. A high-speed infrared camera (Photonfocus

DR1) and a mirror angled at 45 degrees captured whisker motion under IR illumination. Videos

were synchronized with neural data via external triggers generated by a National Instruments

card and recorded on an Intan 512 controller. DeepLabCut was used to label the whisker(s) and

track their movement [31]. Four evenly spaced points on each whisker were labeled.

Approximately 150 frames were manually labeled from each experiment spanning all surface

positions. The DLC neural network was trained for at least 200k iterations and the final labels

were manually checked for accuracy. Whisker position/angle was calculated for each label on

each whisker with reference to a user defined point on the face relative to the frame's vertical

axis. Whisker angle was bandpass filtered (1-30Hz, fdesign.bandpass order 4, MATLAB).

Whisker bend/curvature was calculated from the three distal labels on each whisker using

Menger curvature. The whisker curvature derivative was calculated as the local slope with cubic

approximation in a moving window of 100ms.

Spike Sorting

Spikes sorting was performed using the Kilosort2 [67] and manually curated using Phy2 gui

(https://github.com/cortex-lab/phy). Spike clusters were considered single units based on their

spike waveform, location on the electrode, and cross-correlogram of spike times. Single units

were used for all analyses in the paper.

Statistical Analyses

Surface movement statistical classifier

To classify the location of the tactile surface from population-level SC activity, we applied the

Neural Decoding Toolbox [68] using a support vector machine (SVM). The classifier predicted

surface location (rostral, caudal, center) throughout the duration of the trial. Only neurons with a

significant response to surface translation were used. Spike data was first Z-scored to prevent

high firing rate neurons from having a disproportionate influence on classification. The spike rate

of every neuron was calculated using 150ms bins and surface location was predicted every

20ms to plot classification accuracies over time. A 10-fold cross validation was performed along

with 50 resample runs to calculate a robust estimate of classification accuracies. The

classification accuracy was measured based on the zero-one loss function.

Modulation Indices

The gain and loss of whisker modulation indices (GoW/LoW indices) were calculated as the

difference divided by the sum of spike rate averages that were calculated 500 ms before and

after the start of surface movement.

Transient and sustained neurons classification:

Surface movement was defined as a GoW/LoW stimulus when at least one whisker gained or

lost contact. Since changes in whisker contact (GoW/LoW) occurred at slight time differences

relative to the onset of surface movement across mice, we selected a custom transient window

for each mouse based on the population average firing rates for every GoW/LoW translation.

The peak firing rate of the transient response was identified by eye and the troughs around the

peaks were manually marked as the start and the end of the transient window. We verified this

window onset with analysis of whisker kinematics (whisker curvature and position). The firing

rate of the sustained response was calculated in a 300 ms window that started one second after

the transient window ended. We performed a one-way ANOVA (MATLAB anova1) on the firing

rates calculated in the baseline (300 ms pre-movement), transient, and sustained windows for

every neuron. A Tukey post hoc test (multcompare, MATLAB) was used to test for significant (α

< 0.05) differences, correcting for multiple comparisons. Neurons which had significantly

different baseline and sustained firing rates were classified as sustained neurons. Neurons with a transient window firing rate that was greater than both their baseline and sustain window firing rates were classified as transient neurons. Neurons which had a significant transient and/or sustained response were considered tactile responsive and used for analysis.

Self-GoW and external-GoW (Figure 3)

Self-GoW and external-GoW firing rates were calculated in a 50ms window following touch onset. Self-GoW was calculated as a difference between mean firing rates during self-GoW and self-motion. Self-motion occurred when the animal transitioned from resting to locomotion but free-whisking in air. The external-GoW response was calculated as the mean firing rate in the touch window minus the pre-touch (free-whisking) window.

Expected and Unexpected touch (Figure 4)

Expected and Unexpected touch firing rates were calculated by subtracting pre-touch firing rates from post touch in a 50 ms window.

Stimulus repetition

To obtain repetition slopes, we fit firing rates of TR neurons in the TR window and SR neurons in the SR window with a linear regression model using the MATLAB fitlm function. We only used neurons that did not drift. Neuronal drift was determined by testing if the baseline firing rate during the first 10 trials was different from the last 10 trials using a Wilcoxon signed rank test. Spike amplitudes were obtained from 'amplitudes.npy' output from Phy GUI. Normalized whisker position across repetitions were obtained as the mean of the normalized whisker position in the transient response window for each repetition. Spike waveform amplitudes across repetitions were obtained by taking mean of the average normalized waveform from all spikes in the transient window of each repetition. A scaling to range normalization method was performed as below.

$$Xnorm = \frac{X - Xmin}{Xmax - Xmin}$$

Acknowledgements

The authors would like to acknowledge the members of the Pluta lab, Julia Veit, Kate Hong, Edward Bartlett and Daniel Butts for providing valuable feedback on the manuscript.

Competing Interest

The authors declare no competing interest.

References

1. Munoz, D. P. & Wurtz, R. H. Fixation cells in monkey superior colliculus. II. Reversible activation and deactivation. *J. Neurophysiol.* **70**, 576–589 (1993).

2. Hikosaka, O. & Wurtz, R. H. Modification of saccadic eye movements by GABA-related substances. I. Effect of muscimol and bicuculline in monkey superior colliculus. *J. Neurophysiol.* **53**, 266–291 (1985).

3. Allen, K. M., Lawlor, J., Salles, A. & Moss, C. F. Orienting our view of the superior colliculus: specializations and general functions. *Current Opinion in Neurobiology* vol. 71 119–126 at https://doi.org/10.1016/j.conb.2021.10.005 (2021).

4. Drager, U. C. & Hubel, D. H. Topography of visual and somatosensory projections to mouse superior colliculus. *J. Neurophysiol.* **39**, 91–101 (1976).

5. Evans, D. A. *et al.* A synaptic threshold mechanism for computing escape decisions. *Nature* **558**, 590–594 (2018).

6. Essig, J., Hunt, J. B. & Felsen, G. Inhibitory neurons in the superior colliculus mediate selection of spatially-directed movements. *Commun. Biol.* **4**, (2021).

7. Duan, C. A. *et al.* Collicular circuits for flexible sensorimotor routing. *Nat. Neurosci.* **24**, 1110–1120 (2021).

8. Hoy, J. L., Yavorska, I., Wehr, M. & Niell, C. M. Vision Drives Accurate Approach Behavior during Prey Capture in Laboratory Mice. *Curr. Biol.* **26**, 3046–3052 (2016).

9. Shang, C. *et al.* A subcortical excitatory circuit for sensory-triggered predatory hunting in mice. *Nat. Neurosci.* **22**, 909–920 (2019).

10. Huang, M. *et al.* The tectonigral pathway regulates appetitive locomotion in predatory hunting in mice. *Nat. Commun.* **12**, 1–17 (2021).

11. Campagner, D. *et al.* A cortico-collicular circuit for orienting to shelter during escape. *Nat. 2022 6137942* **613**, 111–119 (2022).

12. Sommer, M. A. & Wurtz, R. H. A pathway in primate brain for internal monitoring of movements. *Science (80-.).* **296**, 1480–1482 (2002).

13. Kojima, Y. & Soetedjo, R. Elimination of the error signal in the superior colliculus impairs saccade motor learning. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E8987–E8995 (2018).

14. Wheatcroft, T., Saleem, A. B. & Solomon, S. G. Functional Organisation of the Mouse Superior Colliculus. *Frontiers in Neural Circuits* vol. 16 23 at https://doi.org/10.3389/fncir.2022.792959 (2022).

15. Hoy, J. L., Bishop, H. I. & Niell, C. M. Defined Cell Types in Superior Colliculus Make Distinct Contributions to Prey Capture Behavior in the Mouse. *Curr. Biol.* **29**, 4130-4138.e5 (2019).

16. Johnson, K. P. *et al.* Cell-type-specific binocular vision guides predation in mice. *Neuron*

109, 1527-1539.e4 (2021).

17.   Anjum, F., Turni, H., Mulder, P. G. H. H., van der Burg, J. & Brecht, M. Tactile guidance
of prey capture in Etruscan shrews. *Proc. Natl. Acad. Sci.* **103**, 16544–16549 (2006).

18.   Cang, J., Savier, E., Barchini, J. & Liu, X. Visual Function, Organization, and
Development of the Mouse Superior Colliculus. 1–24 (2018) doi:10.1146/annurev-vision-
091517.

19.   Basso, M. A. & May, P. J. Circuits for Action and Cognition: A View from the Superior
Colliculus. *Annu. Rev. Vis. Sci.* **3**, 197–226 (2017).

20.   Nagy, A., Kruse, W., Rottmann, S., Dannenberg, S. & Hoffmann, K. P. Somatosensory-
Motor Neuronal Activity in the Superior Colliculus of the Primate. *Neuron* **52**, 525–534
(2006).

21.   Groh, J. M. & Sparks, D. L. Saccades to somatosensory targets. III. eye-position-
dependent somatosensory activity in primate superior colliculus. *J. Neurophysiol.* **75**,
439–453 (1996).

22.   Castro-Alamancos, M. A. & Favero, M. Whisker-related afferents in superior colliculus. *J.*
Neurophysiol. **115**, 2265–2279 (2016).

23.   Bezdudnaya, T. & Castro-Alamancos, M. A. Superior colliculus cells sensitive to active
touch and texture during whisking. *J. Neurophysiol.* **106**, 332–346 (2011).

24.   Cohen, J. D., Hirata, A. & Castro-Alamancos, M. A. Vibrissa sensation in superior
colliculus: Wide-field sensitivity and state-dependent cortical feedback. *J. Neurosci.* **28**,
11205–11220 (2008).

25.   Cohen, J. D. & Castro-Alamancos, M. A. Behavioral state dependency of neural activity
and sensory (whisker) responses in superior colliculus. *J. Neurophysiol.* **104**, 1661–1672
(2010).

26.   Goldberg, M. E. & Wurtz, R. H. Activity of superior colliculus in behaving monkey. II.
Effect of attention on neuronal responses. *J. Neurophysiol.* **35**, 560–574 (1972).

27.   Lee, J. & Groh, J. M. Auditory signals evolve from hybrid- to eye-centered coordinates in
the primate superior colliculus. *J. Neurophysiol.* **108**, 227–242 (2012).

28.   Jay, M. F. & Sparks, D. L. Sensorimotor integration in the primate superior colliculus. II.
Coordinates of auditory signals. *J. Neurophysiol.* **57**, 35–55 (1987).

29.   Robinson, D. L. & Wurtz, R. H. Use of an extraretinal signal by monkey superior colliculus
neurons to distinguish real from self induced stimulus movement. *J. Neurophysiol.* **39**,
852–870 (1976).

30.   Richmond, B. J. & Wurtz, R. H. Vision during saccadic eye movements. II. A corollary
discharge to monkey superior colliculus. *J. Neurophysiol.* **43**, 1156–1167 (1980).

31.   Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with
deep learning. *Nat. Neurosci.* **21**, 1281–1289 (2018).

32.   Clack, N. G. *et al.* Automated tracking of whiskers in videos of head fixed rodents. *PLoS*
Comput. Biol. **8**, (2012).

33.   Ayaz, A. *et al.* Layer-specific integration of locomotion and sensory information in mouse

barrel cortex. *Nat. Commun.* **10**, 1–14 (2019).

34. Keller, G. B., Bonhoeffer, T. & Hübener, M. Sensorimotor Mismatch Signals in Primary
Visual Cortex of the Behaving Mouse. *Neuron* **74**, 809–815 (2012).

35. Isett, B. R., Feasel, S. H., Lane, M. A. & Feldman, D. E. Slip-Based Coding of Local
Shape and Texture in Mouse S1. *Neuron* 1–16 (2018) doi:10.1016/j.neuron.2017.12.021.

36. Zhao, Z., Ahissar, E., Victor, J. D. & Rucci, M. Inferring visual space from ultra-fine extra-
retinal knowledge of gaze position. *Nat. Commun. 2023 141* **14**, 1–12 (2023).

37. Wang, L., Herman, J. P. & Krauzlis, R. J. Neuronal modulation in the mouse superior
colliculus during covert visual selective attention. *Sci. Rep.* **12**, (2022).

38. Lee, K. H., Tran, A., Turan, Z. & Meister, M. The sifting of visual information in the
superior colliculus. *Elife* **9**, (2020).

39. Boehnke, S. E. & Munoz, D. P. On the importance of the transient visual response in the
superior colliculus. *Current Opinion in Neurobiology* vol. 18 544–551 at
https://doi.org/10.1016/j.conb.2008.11.004 (2008).

40. Aronoff, R. *et al.* Long-range connectivity of mouse primary somatosensory barrel cortex.
Eur. J. Neurosci. **31**, 2221–2233 (2010).

41. Wise, S. P. & Jones, E. G. Somatotopic and columnar organization in the corticotectal
projection of the rat somatic sensory cortex. *Brain Res.* **133**, 223–235 (1977).

42. Adibi, M. & Lampl, I. Sensory Adaptation in the Whisker-Mediated Tactile System:
Physiology, Theory, and Function. *Frontiers in Neuroscience* vol. 15 1406 at
https://doi.org/10.3389/fnins.2021.770011 (2021).

43. Masullo, L. *et al.* Genetically Defined Functional Modules for Spatial Orienting in the
Mouse Superior Colliculus. *Curr. Biol.* **29**, 2892-2904.e8 (2019).

44. Alloway, K. D., Smith, J. B. & Beauchemin, K. J. Quantitative analysis of the bilateral
brainstem projections from the whisker and forepaw regions in rat primary motor cortex.
J. Comp. Neurol. **518**, 4546–4566 (2010).

45. Singla, S., Dempsey, C., Warren, R., Enikolopov, A. G. & Sawtell, N. B. A cerebellum-like
circuit in the auditory system cancels responses to self-generated sounds. *Nat. Neurosci.*
20, 943–950 (2017).

46. Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error correction, sensory prediction, and
adaptation in motor control. *Annual Review of Neuroscience* vol. 33 89–108 at
https://doi.org/10.1146/annurev-neuro-060909-153135 (2010).

47. Kilteni, K. & Henrik Ehrsson, H. Functional connectivity between the cerebellum and
somatosensory areas implements the attenuation of self-generated touch. *J. Neurosci.*
40, 894–906 (2020).

48. Anderson, S. R. *et al.* An Internal Model Architecture for Novelty Detection: Implications
for Cerebellar and Collicular Roles in Sensory Processing. *PLoS One* **7**, e44560 (2012).

49. Lee, J. & Sabatini, B. L. Striatal indirect pathway mediates exploration via collicular
competition. *Nature* **599**, 645–649 (2021).

50. Jiang, H., Stein, B. E. & McHaffie, J. G. Opposing basal ganglia processes shape

midbrain visuomotor activity bilaterally. *Nature* **423**, 982–986 (2003).

51.   Liu, P. & Basso, M. A. Substantia nigra stimulation influences monkey superior colliculus neuronal activity bilaterally. *J. Neurophysiol.* **100**, 1098–1112 (2008).

52.   Kaneda, K., Isa, K., Yanagawa, Y. & Isa, T. Nigral inhibition of GABAergic neurons in mouse superior colliculus. *J. Neurosci.* **28**, 11071–11078 (2008).

53.   Lee, J., Wang, W. & Sabatini, B. L. Anatomically segregated basal ganglia pathways allow parallel behavioral modulation. *Nat. Neurosci.* **23**, 1388–1398 (2020).

54.   Mcelvain, L. E. *et al.* Specific populations of basal ganglia output neurons target distinct brain stem areas while collateralizing throughout the diencephalon In brief. *Neuron* **109**, 1–18 (2021).

55.   Villalobos, C. A., Correspondence, M. A. B. & Basso, M. A. Optogenetic activation of the inhibitory nigro-collicular circuit evokes contralateral orienting movements in mice. *Cell Rep.* **39**, 110699 (2022).

56.   Tafreshiha, A., Burg, S. A. V., Smits, K., Blomer, L. A. & Heimel, J. A. Visual stimulus-specific habituation of innate defensive behaviour in mice. *J. Exp. Biol.* **224**, (2021).

57.   Sooksawate, T., Isa, K., Behan, M., Yanagawa, Y. & Isa, T. Organization of GABAergic inhibition in the motor output layer of the superior colliculus. *Eur. J. Neurosci.* **33**, 421–432 (2011).

58.   Boehnke, S. E. *et al.* Visual adaptation and novelty responses in the superior colliculus. *Eur. J. Neurosci.* **34**, 766–779 (2011).

59.   Drager, U. C. & Hubel, D. H. Responses to visual stimulation and relationship between visual, auditory, and somatosensory inputs in mouse superior colliculus. *J. Neurophysiol.* **38**, 690–713 (1975).

60.   Cynader, M. & Berman, N. Receptive-field organization of monkey superior colliculus. *J. Neurophysiol.* **35**, 187–201 (1972).

61.   Brooks, J. X. & Cullen, K. E. The primate cerebellum selectively encodes unexpected self-motion. *Curr. Biol.* **23**, 947–955 (2013).

62.   Troncoso, X. G. *et al.* V1 neurons respond differently to object motion versus motion from eye movements. *Nat. Commun.* **6**, 1–10 (2015).

63.   Zmarz, P. & Keller, G. B. Mismatch Receptive Fields in Mouse Visual Cortex. *Neuron* **92**, 766–772 (2016).

64.   Jordan, R. & Keller, G. B. Opposing Influence of Top-down and Bottom-up Input on Excitatory Layer 2/3 Neurons in Mouse Primary Visual Cortex. *Neuron* **108**, 1194-1206.e5 (2020).

65.   Bell, C. C. Memory-based expectations in electrosensory systems. *Current Opinion in Neurobiology* vol. 11 481–487 at https://doi.org/10.1016/S0959-4388(00)00238-5 (2001).

66.   Xie, Z. *et al.* Transcriptomic encoding of sensorimotor transformation in the midbrain. *Elife* **10**, 1–34 (2021).

67.   Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M. & Harris, K. D. Kilosort: realtime spike-sorting for extracellular electrophysiology with hundreds of channels. *bioRxiv*

576   061481 (2016) doi:10.1101/061481.

577   68.   Meyers, E. M. The neural decoding toolbox. *Front. Neuroinform.* **7**, 8 (2013).

579