Plasma amyloid-β homeostasis is associated with Body Mass Index and weight loss in people with overweight and obesity

AUTHORS

Emily S. **Brook**[a,b], Zachary J. **D'Alonzo**[a,b], Virginie **Lam**[a,c], Dick **Chan**[d], Satvinder **Singh Dhaliwal**[a] , Gerald F. **Watts**[d,e], John C. L **Mamo**[a,c], Ryusuke **Takechi**[a,c]

AFFILIATIONS

[a]Curtin Health Innovation Research Institute, Faculty of Health Sciences, Curtin University, Perth, Australia.

[b]Curtin Medical School, Faculty of Health Sciences, Curtin University, Perth, WA 6845, Australia

[c]School of Population Health, Faculty of Health Sciences, Curtin University, Perth, Australia.

[d]Medical School, University of Western Australia, Perth, Western Australia, Australia.

[e]Cardiometabolic Service, Department of Cardiology and Internal Medicine, Royal Perth Hospital, Perth, Australia

RUNNING TITLE

Plasma Aβ is linked to BMI and weight loss

COMPLETE CORRESPONDENCE ADDRESS

Ryusuke Takechi, PhD

Curtin University Faculty of Health Sciences

Perth, WA AUSTRALIA

Email: R.Takechi@curtin.edu.au

Phone: 0430 227 755

ABSTRACT

BACKGROUND: Obesity is linked to a higher incidence of Alzheimer's disease (AD). Studies show that plasma amyloid-β (Aβ) dyshomeostasis, particularly low 42/40 ratio indicates a heightened risk for developing AD. However, the relationship between body mass index (BMI) and circulating plasma Aβ has not been extensively studied.

OBJECTIVE: We hypothesised that people with a high BMI have altered plasma Aβ homeostasis compared with people with a lower BMI. We also tested whether reducing BMI by calorie-restriction could normalise plasma concentrations of Aβ.

METHODS: Plasma concentrations of $Aβ_{40}$, $Aβ_{42}$ and $Aβ_{42/40}$ ratio were measured in 106 participants with BMIs classified as lean, overweight, or obese. From this cohort, twelve participants with overweight or obese BMIs entered a 12-week calorie-restriction weight loss program. We then tested whether decreasing BMI affected plasma Aβ concentrations.

RESULTS: Plasma $Aβ_{42/40}$ ratio was 17.54% lower in participants with an obese BMI compared to lean participants (p<0.0001), and 11.76% lower compared to participants with an overweight BMI (p<0.0001). The weight loss regimen decreased BMI by an average of 4.02% (p=0.0005) and was associated with a 6.5% decrease in plasma $Aβ_{40}$ (p=0.0425). However, weight loss showed negligible correlations with plasma $Aβ_{40}$, $Aβ_{42}$ and $Aβ_{42/40}$ ratio.

CONCLUSION: Obesity is associated with aberrant plasma Aβ homeostasis which may be associated with an increased risk for AD. Weight loss appears to lower $Aβ_{40}$, but large-scale longitudinal studies in addition to molecular studies are required to elucidate the underlying mechanisms of how obesity and weight loss influence plasma Aβ homeostasis.

KEYWORDS

Amyloid-β, Body Mass Index, Alzheimer's Disease, Obesity, weight loss

INTRODUCTION

Alzheimer's disease (AD), the most common form of dementia, is characterised by progressive cognitive decline and the abnormal deposition of amyloid-β (Aβ) in the brain parenchyma [1]. A significant risk factor for AD is obesity, which is defined as body mass index (BMI) being ≥30 kg/m^2

[2, 3]. Obesity in people aged between 40-45 years has been associated with a three-fold higher risk of developing AD in later-life [4]. Additionally, obesity was reported to significantly accelerate the onset of AD and double the likelihood of Aβ deposition in the brain [5, 6]. More recently, overweight and obese BMIs were found to be associated with detrimental changes to brain structure, such as increased white matter oedema and low hippocampal volume [7]. These data collectively suggest that an increased BMI may have a severe impact on cerebral Aβ overload, AD onset and progression, in addition to overall brain health. However, the exact underlying mechanisms linking obesity with AD remain unknown.

Many studies have evaluated plasma levels of $A\beta_{40}$ and $A\beta_{42}$ as predictors of AD or cognitive decline [8]. $A\beta_{40}$ and $A\beta_{42}$ are generated from the sequential proteolytic cleavage of Aβ precursor protein (AβPP) by β-site AβPP cleaving enzyme-1 (BACE1) and γ-Secretase [9]. $A\beta_{40}$ is the most physiologically abundant species of amyloid, whilst $A\beta_{42}$ is regarded as primarily pathogenic as it is prone to aggregation and eventually forms insoluble fibrillary structures which are the core component of amyloid plaques [10, 11]. Recent findings show that the dyshomeostasis of plasma Aβ, particularly low plasma $A\beta_{42/40}$ ratio can predict the risk of AD at an accuracy of ~90% [12-19]. However, few studies to date have investigated the association between BMI, weight loss and plasma Aβ homeostasis [20, 21], although obesity is a significant modifiable risk factor for AD.

In this pilot study, we investigated whether BMI shows any significant associations with plasma Aβ homeostasis. Furthermore, in individuals with BMIs classified as overweight or obese, we tested whether reducing BMI through a calorie restriction weight loss regime could modulate plasma Aβ homeostasis. The outcomes of this study may provide new insights into the mechanisms whereby obesity increases brain amyloidosis and risk of AD and may offer potential preventative opportunity with weight loss.

MATERIALS AND METHODS

PARTICIPANTS

To investigate the association between BMI and plasma Aβ homeostasis, we recruited 106 people who either had a normal, overweight, or an obese BMI. BMI cut-offs were defined according to the World Health Organisation: lean weight as 18.50-24.99 kg/m^2, overweight as 25.00-29.99 kg/m^2, and obese as ≥30.00 kg/m^2 [22]. No participants had an underweight BMI (≤18.50 kg/m^2). All participants were Caucasian, and those with familial hypercholesterolemia, a history of cardiovascular disease or used antidiabetic medication were excluded from the study. In addition, participants with APOE/E2 genotype were excluded. Aside from ageing, APOE genotype is the most significant genetic risk factor for AD, with APOE2/E2 conferring for reduced AD risk as detailed in [23]. This study was approved by the Ethics Committee of Royal Perth Hospital (EC 2010/074), a national ethics committee (Bellberry Ltd, Eastwood, South Australia, 2014-03-151) and Curtin University (HRE2021-0192). Informed consent was obtained from each participant.

SAMPLE COLLECTION AND STUDY DESIGN

After a 14-hour fast, baseline whole venous blood was collected in EDTA tubes from all 106 participants and immediately centrifuged at 1500 x g for 15 min at 4°C to obtain plasma. Plasma was collected and stored at –80°C for future evaluation. At this visit, we also recorded all participants' weight and height for their BMI calculation.

WEIGHT LOSS PROGRAM

Following the plasma sample and BMI data collection, all participants who had an overweight or obese BMI were invited to participate in a 12-week calorie restriction weight loss program as reported previously [23], and 12 chose to participate. During this weight loss regime, the participants' caloric intake was restricted to achieve a deficit of ~1440 kJ per day compared to their diets before the commencement of weight loss. Dietary intake was evaluated for energy and major nutrients by FoodWorks 2007 (Xyris Software, Brisbane, Australia). All participants were reviewed fortnightly and were requested to maintain their usual level of physical activity. All dietary assessments and recommendations were conducted by a registered dietician. After completing the weight loss program, fasting plasma was collected, and BMI was recorded for the 12 participants.

QUANTIFICATION OF PLASMA AB_{40} AND AB_{42}

Plasma concentrations of Aβ_{40} and Aβ_{42} were measured at baseline (*n*=106) and following the 12-week weight loss program (*n*=12) using ELISA (Wako Chemical, Tokyo, Japan), according to the manufacturer's instruction. Briefly, samples were assays in singlets and prepared in a 1 in 4 dilution with the kit provided diluent. All samples and kit-provided standards were loaded onto the antibody coated microplate and incubated overnight at 4°C. Following incubation, HRP-conjugated monoclonal antibody was added to the appropriate wells, and plates were incubated in the dark. Plates detecting Aβ_{40} were incubated for 2 hours, whilst plates detecting Aβ_{42} were incubated for 1 hour. Subsequently, 100 µL TMB was added to each well. The plates were incubated for a further 30 minutes in the dark, and STOP solution was added before the absorbance was measured at 450 nm using EnSpire™ Multimode Plate Reader (PerkinElmer, Waltham, MA, USA).

STATISTICAL ANALYSIS

Statistical analyses were performed using GraphPad Prism 9.3.1 for Windows (GraphPad Software, CA, USA) and IBM SPSS Statistics for Windows, Version 27.0. (IBM Corp, NY, USA). The distribution of all variables was assessed by D'Agostino and Shapiro-wilk normality test. Unpaired t tests were used to compare variables of the three BMI groups. Where data was not normally distributed, intergroup comparisons were assessed using the Mann-Whitney test. For the calorie restriction part of the study, paired t tests were used to compare the difference from baseline to week-12. Where data was not normally distributed, changes were analysed using the Wilcoxon test. For correlation analyses, we used Pearson or Spearman correlation coefficient tests depending on the data normality, as well as principal component analysis (PCA). Where indicated, multivariable regression model was used to adjust data for age, age squared and sex. Significance was defined as *p*<0.05.

RESULTS

The characteristics of the cohort are summarised in Table 1. Overall, participants were aged between 18 to 66 years. Individuals with an obese BMI were significantly older compared to participants with a lean or overweight BMI, whilst the average participant age of the lean and overweight groups were similar. Majority of the participants were male, accounting for 92.67% of the overall cohort.

The average plasma concentration of $A\beta_{40}$ was significantly higher in people with an obese BMI compared to those with lean or overweight BMI by 41.7% and 36.89% respectively, whilst there was no significant difference observed between the lean and overweight groups (Fig 1A). Moreover, plasma abundance of $A\beta_{42}$ was 12.51% higher in the obese group compared to overweight group ($p=$ 0.0404); however, there were no other significant intergroup comparisons (Fig 1B). Lastly, plasma $A\beta_{42/40}$ ratio showed a significant sequential decline as BMI categories progressed from lean, to overweight, to obese. Specifically, plasma $A\beta_{42/40}$ ratio was 32.65% and 25.74% lower in the obese group compared to the lean and overweight BMI groups respectively, and 7.05% lower in the overweight cohort compared to the lean group (Fig 1C).

To further evaluate the relationship between plasma $A\beta$ and BMI, we performed simple linear correlation analysis (Fig 2). BMI showed a significant positive correlation with plasma $A\beta40$ concentration (Fig 2A), while no significant association was observed between BMI and $A\beta_{42}$ (Fig 2B). Additionally, we identified a moderate negative correlation between BMI and $A\beta_{42/40}$ ratio (Fig 2C). Consistently, the PCA loading plot showed a strong negative relationship between BMI and $A\beta_{42/40}$ as indicated by the large angle and opposing direction of the two variables. However, $A\beta_{40}$ and $A\beta_{42}$ *per se* appeared as independent factors as indicated by their near orthogonal directions in relation to BMI (Fig 2D). In addition, the PCA scores plot displayed evident clustering of lean, overweight, and obese populations, with obesity showing negative correlation with $A\beta_{42/40}$ ratio (Fig 2E).

As the average age of obese group was significantly higher compared to the lean and overweight BMI groups, and there were nearly 9-fold more men than women overall (Table 1); we used a multivariable regression model to adjust the plasma $A\beta$ concentrations for age, age squared and sex [24]. Following adjustment, the mean plasma $A\beta_{40}$ in the obese BMI group was ~11% greater than the lean or overweight groups; whilst plasma $A\beta_{40}$ levels were similar between lean and overweight groups (Fig 3A). Moreover, the mean plasma $A\beta_{42}$ concentrations were not significantly different across lean, overweight, and obese groups (Fig 3B). However, the inverse relationship between BMI and $A\beta_{42/40}$ ratio persisted following adjustment and showed significantly lower $A\beta_{42/40}$ in the obese group,

compared to both lean and overweight BMI groups by 17.54% and 11.76%, respectively. We also note that the $A\beta_{42/40}$ ratio in overweight trended lower than the lean group (Fig 3C).

To further explore the relationship between BMI and plasma $A\beta$ homeostasis, we tested whether weight loss in participants with overweight and obese BMIs would alter plasma concentrations of $A\beta$. At baseline, 2 participants had an overweight BMI, whilst the remaining 10 participants had BMIs that were classed as obese. After the 12-week weight loss intervention, participants significantly reduced their BMI by an average of 4.021% (Table 2).

Plasma concentrations of $A\beta_{40}$, $A\beta_{42}$ and $A\beta_{42/40}$ ratio for baseline and 12-weeks are shown in Figure 4. Following weight loss, plasma $A\beta_{40}$ significantly decreased by 6.5% (Fig 4A). However, we note that the significant decrease in $A\beta_{40}$ is heterogenous and predominantly driven by two participants where plasma $A\beta_{40}$ decreased by 25.62% and 29.7%. Of these two participants, one also showed a 50.5% decrease in $A\beta_{42}$ and a 29.6% decrease $A\beta_{40/42}$ ratio. Whilst the other showed a 24.5% increase in $A\beta_{42/40}$ ratio, which can be attributed to the decrease in $A\beta_{40}$ and minimal change in $A\beta_{42}$.

The percent change of plasma $A\beta_{40}$ from baseline for the remaining participants ranged from -8.35-5.45%. Plasma $A\beta_{42}$ and the $A\beta_{42/40}$ ratio did not show any significant changes following weight loss (Fig 4B-C).

We also performed correlation analysis between the percent change from baseline for plasma $A\beta$ and weight lost (Fig 5). Overall, plasma $A\beta_{40}$ shows a moderate negative correlation with percentage of weight lost, however this failed to reach statistical significance (Fig 5A). Plasma $A\beta_{42}$ and $A\beta_{42/40}$ ratio did not show any significant correlations with weight-loss (Fig 5B-C).

DISCUSSION

In this pilot study we investigated whether increased risk of AD in obesity is associated with plasma $A\beta$ dyshomeostasis and examined the relationship between BMI and plasma $A\beta$ concentrations of 106 adults.

Here, we report that plasma $A\beta_{40}$ was significantly elevated in participants with an obese BMI compared to those in the overweight and lean groups. Consistently, plasma $A\beta_{40}$ showed significant positive correlation with BMI, whilst plasma $A\beta_{42}$ did not show any significant association with BMI. Nonetheless, BMI showed moderate negative correlation with $A\beta_{42/40}$ ratio and indicated significantly lower $A\beta_{42/40}$ ratio in the obese group compared to the people with a lean or overweight BMI. The distinct association between an obese BMI and low plasma $A\beta_{42/40}$ ratio persisted after adjusting the data for age, age-squared and sex. Low plasma $A\beta_{42/40}$ ratio has been consistently associated with high cerebral amyloid burden, greater cognitive deterioration, and increased risk of developing AD [12-19, 25-28]. Consequently, the low plasma $A\beta_{42/40}$ ratio observed amongst the obese BMI group may suggest a heightened risk for AD. Collectively, our data suggests that obesity and a higher BMI may predispose people to an elevated risk of AD through disturbances in plasma $A\beta$ homeostasis. However, the effect of obesity on plasma $A\beta_{42/40}$ is largely understudied, and the underlying mechanisms by which obesity increases plasma $A\beta_{40}$ and decreases in $A\beta_{42/40}$ ratio are currently unknown.

$A\beta PP$, the precursor molecule for $A\beta$ generation, is reported to be significantly overexpressed in the adipocytes of people with obese BMIs compared to a lean population [21, 29], indicating greater availability of $A\beta$ biosynthesis in obesity. Additionally, preclinical findings demonstrate that in obesity, the activity of BACE1, the enzyme responsible for the extracellular cleavage of $A\beta PP$ and the subsequent production of $A\beta$, is significantly elevated [30-32]. Moreover, pre-clinical studies report that the production and secretion into circulation of $A\beta$ are predominantly regulated through lipid and lipoprotein metabolism, particularly cholesterol and triglyceride-rich lipoproteins [33-35]. In wild-type mice, ingestion of a diet enriched in saturated fats significantly elevated $A\beta$ synthesis by the small intestinal enterocytes, which increased plasma $A\beta$ concomitant with heightened non-esterified fatty acids [36, 37]. In alignment with these findings, pharmacological lipid-lowering interventions resulted in the significant reduction of plasma $A\beta$ in wild-type mice [38-41]. These data suggest that impaired lipid metabolism and heightened plasma lipids in obesity may elevate plasma $A\beta$. Taken together, obesity may directly and indirectly contribute to peripheral $A\beta$ dysregulation and lead to a low $A\beta_{42/40}$

ratio; however, mechanistic studies are required to delineate the complete effect obesity has on $A\beta$ homeostasis and its contribution to the AD pathological cascade.

Considering that obesity appears to disturb systemic $A\beta$ homeostasis, we explored whether reducing BMI by calorie-restriction could influence plasma $A\beta$ homeostasis in people with overweight and obese BMIs. Following weight loss, plasma $A\beta_{40}$ significantly decreased, albeit heterogeneously, whilst plasma $A\beta_{42}$ and $A\beta_{42/40}$ ratio remained largely unaffected. Provided that our previously published findings suggest that lipid metabolism, triglyceride-rich lipoproteins and cholesterol may significantly regulate $A\beta$ homeostasis, we speculate that reduced abundance of triglycerides and changes in cholesterol metabolism may also lead to decreased plasma $A\beta_{40}$ [23, 36-41]. It is well documented that weight loss decreases plasma triglycerides, increases HDL cholesterol, and overall, normalises lipid profiles [23, 42, 43], and accordingly, we reported that weight loss in the present cohort decreased triglycerides and increased HDL cholesterol [23]. Interestingly, historic findings by Koudinov and colleagues indicate that apolipoproteins of HDL, namely apolipoprotein (apo) A-I, apo A-II, apo E, and apo J bind to $A\beta_{40}$ [44]. Such findings are corroborated by recent clinical evidence whereby apo A-I appears to be a potent $A\beta$-binding protein [45]. Therefore, based on the current evidence, HDL may facilitate the clearance of $A\beta$, and we speculate that reduced $A\beta_{40}$ following weight loss may be mediated by increased HDL.

In parallel, the adipocytes of people with obesity appear to over produce $A\beta_{40}$, $A\beta PP$ and BACE1[21, 29-32]. Therefore, we also hypothesise that loss of adipose tissue during weight loss may reduce the availability of amyloidogenic material, and thus contribute to the reduced $A\beta_{40}$ observed in this study. Taken together, calorie restriction induced weight loss appears to lower $A\beta_{40}$, which may be facilitated via a HDL mediated pathway, and fat loss. Overall, the complexities of the relationship between weight loss, lipid metabolism, and $A\beta_{40}$ is largely under studied and requires further research to determine the complete effect calorie restriction has on $A\beta$ homeostasis in obesity.

In the present pilot study we evaluated plasma $A\beta$ only; however, AD is a heterogenous disease with a hotly debated aetiology [46, 47], consequently numerous plasma markers of AD have been considered. Most notably, plasma phosphorylated-tau181 (p-tau18) has emerged as a robust and sensitive marker

for AD. Evidence indicates that plasma p-tau181 can accurately discriminate people with AD from cognitively healthy adults, and distinguish AD from other neuropathological diseases [48]. Additionally, plasma p-tau181 can accurately predict future brain atrophy [49], cerebral deposition of amyloid, and neurodegeneration [50]. Further compelling evidence also suggests that plasma p-Tau181 may even surpass plasma $A\beta_{42/40}$ as a predictive biomarker of AD [51, 52]. However, whether tau pathology or p-Tau181 is influenced by obesity or weight-loss is uncertain and understudied [53, 54]. In contrast, our data, along with others' show that obesity does appear to disturb $A\beta$ homeostasis, which may be ameliorated through weight loss, ultimately reducing risk of AD. To further substantiate these findings, we suggest that future studies employ a battery of biomarkers, including plasma p-Tau181, $A\beta_{42/40}$ and amyloid positron emission tomography, in addition to a series of cognitive function tests.

Lastly, we acknowledge that there are demographic and methodological limitations to this pilot study. Demographically, the baseline BMI and weight loss data were obtained from limited sample sizes; the former of which was also not consistently gender balanced, although we did account for the effect of sex in our analysis. From a methodological standpoint, the weight loss intervention was conducted over a relatively short term and only modestly reduced BMI (<4.1%), which may have led to the heterogenous decrease in plasma $A\beta_{40}$ and insignificant changes in plasma $A\beta_{42}$ and $A\beta_{42/40}$ ratio. Moreover, our study did not assess cognition. Considering participants from the obese BMI group were significantly older, subtle cognitive deficits and metabolic differences may have been present, which may have affected our data, although we partially addressed this matter by adjusting the data for age. Furthermore, we did not the fully account for the effect of the APOE genotype as only individuals with APOE2/2 were identified as part of the exclusion criteria. Additionally, plasma $A\beta$ concentrations were assayed in singlets, thus any variability was not detected. We also suggest that future studies consider more sensitive methods for quantifying plasma $A\beta$, such as mass spectrometry-based protocols [55]. Finally, the use of BMI as an indicator of obesity has been largely criticised for not considering body composition, gender, or ethnicity [56, 57]. Therefore, in-depth studies in diverse populations which assess muscle mass and adiposity in relation to plasma $A\beta$ would be useful to

further explore how obesity influences peripheral Aβ homeostasis. Collectively, our conclusions that relate to AD development are to be interpreted circumspectly. In addition, thorough mechanistic and longitudinal studies are indeed required to substantiate the findings and interpretations of this pilot study.

In summary, our data shows that people with an obese BMI show a reduced plasma $A\beta_{42/40}$ ratio which may translate to a heightened risk of developing AD. Importantly, we also demonstrate that lowering BMI appears to decrease plasma $A\beta_{40}$, thus shifting it toward concentrations observed in the lean BMI group. Overall, our findings may help understand the mechanisms whereby obesity increases AD risks and offers potential preventative opportunity through weight loss.

ACKNOWLEDGEMENTS

The authors have no acknowledgements to report.

FUNDING

This study was funded by the National Health and Medical Research Council.

CONFLICTS OF INTEREST

The authors have no conflict of interest to report.

DATA AVAILABILITY

The datasets used and analysed during the current study are available from the corresponding author upon reasonable request.

REFERENCES

[1]   McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Jr., Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carrillo MC, Thies B, Weintraub S, Phelps CH (2011) The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* **7**, 263-269.

[2]   Pedditzi E, Peters R, Beckett N (2016) The risk of overweight/obesity in mid-life and late life for the development of dementia: a systematic review and meta-analysis of longitudinal studies. *Age Ageing* **45**, 14-21.

[3]   Xu WL, Atti AR, Gatz M, Pedersen NL, Johansson B, Fratiglioni L (2011) Midlife overweight and obesity increase late-life dementia risk: a population-based twin study. *Neurology* **76**, 1568-1574.

[4]   Whitmer RA, Gunderson EP, Quesenberry CP, Jr., Zhou J, Yaffe K (2007) Body mass index in midlife and risk of Alzheimer disease and vascular dementia. *Curr Alzheimer Res* **4**, 103-109.

[5]   Chuang YF, An Y, Bilgel M, Wong DF, Troncoso JC, O'Brien RJ, Breitner JC, Ferruci L, Resnick SM, Thambisetty M (2016) Midlife adiposity predicts earlier onset of Alzheimer's dementia, neuropathology and presymptomatic cerebral amyloid accumulation. *Mol Psychiatry* **21**, 910-915.

[6]   Gottesman RF, Schneider AL, Zhou Y, Coresh J, Green E, Gupta N, Knopman DS, Mintz A, Rahmim A, Sharrett AR, Wagenknecht LE, Wong DF, Mosley TH (2017) Association Between Midlife Vascular Risk Factors and Estimated Brain Amyloid Deposition. *JAMA* **317**, 1443-1450.

[7]   Ly M, Raji CA, Yu GZ, Wang Q, Wang Y, Schindler SE, An H, Samara A, Eisenstein SA, Hershey T, Smith G, Klein S, Liu J, Xiong C, Ances BM, Morris JC, Benzinger TLS (2021) Obesity and White Matter Neuroinflammation Related Edema in Alzheimer's Disease Dementia Biomarker Negative Cognitively Normal Individuals. *J Alzheimers Dis* **79**, 1801-1811.

[8]   Koyama A, Okereke OI, Yang T, Blacker D, Selkoe DJ, Grodstein F (2012) Plasma amyloid-beta as a predictor of dementia and cognitive decline: a systematic review and meta-analysis. *Arch Neurol* **69**, 824-831.

[9]   Chen GF, Xu TH, Yan Y, Zhou YR, Jiang Y, Melcher K, Xu HE (2017) Amyloid beta: structure, biology and structure-based therapeutic development. *Acta Pharmacol Sin* **38**, 1205-1235.

[10]   Enzlein T, Cordes J, Munteanu B, Michno W, Serneels L, De Strooper B, Hanrieder J, Wolf I, Chavez-Gutierrez L, Hopf C (2020) Computational Analysis of Alzheimer Amyloid Plaque

Composition in 2D- and Elastically Reconstructed 3D-MALDI MS Images. *Anal Chem* **92**, 14484-14493.

[11]   Roher AE, Lowenson JD, Clarke S, Woods AS, Cotter RJ, Gowing E, Ball MJ (1993) beta-Amyloid-(1-42) is a major component of cerebrovascular amyloid deposits: implications for the pathology of Alzheimer disease. *Proc Natl Acad Sci U S A* **90**, 10836-10840.

[12]   Seppala TT, Herukka SK, Hanninen T, Tervo S, Hallikainen M, Soininen H, Pirttila T (2010) Plasma Abeta42 and Abeta40 as markers of cognitive change in follow-up: a prospective, longitudinal, population-based cohort study. *J Neurol Neurosurg Psychiatry* **81**, 1123-1127.

[13]   Perez-Grijalba V, Romero J, Pesini P, Sarasa L, Monleon I, San-Jose I, Arbizu J, Martinez-Lage P, Munuera J, Ruiz A, Tarraga L, Boada M, Sarasa M (2019) Plasma Abeta42/40 Ratio Detects Early Stages of Alzheimer's Disease and Correlates with CSF and Neuroimaging Biomarkers in the AB255 Study. *J Prev Alzheimers Dis* **6**, 34-41.

[14]   Doecke JD, Perez-Grijalba V, Fandos N, Fowler C, Villemagne VL, Masters CL, Pesini P, Sarasa M, Group AR (2020) Total Abeta42/Abeta40 ratio in plasma predicts amyloid-PET status, independent of clinical AD diagnosis. *Neurology* **94**, e1580-e1591.

[15]   van Oijen M, Hofman A, Soares HD, Koudstaal PJ, Breteler MM (2006) Plasma Abeta(1-40) and Abeta(1-42) and the risk of dementia: a prospective case-cohort study. *Lancet Neurol* **5**, 655-660.

[16]   Okereke OI, Xia W, Selkoe DJ, Grodstein F (2009) Ten-year change in plasma amyloid β levels and late-life cognitive decline. *Archives of neurology* **66**, 1247-1253.

[17]   Nakamura A, Kaneko N, Villemagne VL, Kato T, Doecke J, Dore V, Fowler C, Li QX, Martins R, Rowe C, Tomita T, Matsuzaki K, Ishii K, Ishii K, Arahata Y, Iwamoto S, Ito K, Tanaka K, Masters CL, Yanagisawa K (2018) High performance plasma amyloid-beta biomarkers for Alzheimer's disease. *Nature* **554**, 249-254.

[18]   Yun G, Kim HJ, Kim HG, Lee KM, Hong IK, Kim SH, Rhee HY, Jahng GH, Yoon SS, Park KC, Hwang KS, Lee JS (2021) Association Between Plasma Amyloid-beta and Neuropsychological Performance in Patients With Cognitive Decline. *Front Aging Neurosci* **13**, 736937.

[19]   Li Y, Schindler SE, Bollinger JG, Ovod V, Mawuenyega KG, Weiner MW, Leslie SM, Masters CL, Fowler CJ, Trojanowski JQ, Korecka M, Martins RN, Janelidze S, Hansson O, Bateman RJ (2021) Validation of Plasma Amyloid-beta 42/40 for Detecting Alzheimer Disease Amyloid Plaques. *Neurology*.

[20]   Balakrishnan K, Verdile G, Mehta PD, Beilby J, Nolan D, Galvao DA, Newton R, Gandy SE, Martins RN (2005) Plasma Abeta42 correlates positively with increased body fat in healthy individuals. *J Alzheimers Dis* **8**, 269-282.

[21]   Lee YH, Tharp WG, Maple RL, Nair S, Permana PA, Pratley RE (2008) Amyloid precursor protein expression is upregulated in adipocytes in obesity. *Obesity (Silver Spring)* **16**, 1493-1500.

[22]   (2000) Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser* **894**, i-xii, 1-253.

[23]   Wong AT, Chan DC, Barrett PH, Adams LA, Watts GF (2013) Supplementation with n3 fatty acid ethyl esters increases large and small artery elasticity in obese adults on a weight loss diet. *J Nutr* **143**, 437-441.

[24]   Tyas SL, Koval JJ, Pederson LL (2000) Does an interaction between smoking and drinking influence the risk of Alzheimer's disease? Results from three Canadian data sets. *Stat Med* **19**, 1685-1696.

[25]   Storck SE, Meister S, Nahrath J, Meissner JN, Schubert N, Di Spiezio A, Baches S, Vandenbroucke RE, Bouter Y, Prikulis I, Korth C, Weggen S, Heimann A, Schwaninger M, Bayer TA, Pietrzik CU (2016) Endothelial LRP1 transports amyloid-beta(1-42) across the blood-brain barrier. *J Clin Invest* **126**, 123-136.

[26]   Deane R, Sagare A, Zlokovic BV (2008) The role of the cell surface LRP and soluble LRP in blood-brain barrier Abeta clearance in Alzheimer's disease. *Curr Pharm Des* **14**, 1601-1605.

[27]   Shibata M, Yamada S, Kumar SR, Calero M, Bading J, Frangione B, Holtzman DM, Miller CA, Strickland DK, Ghiso J, Zlokovic BV (2000) Clearance of Alzheimer's amyloid-ss(1-40) peptide from brain by LDL receptor-related protein-1 at the blood-brain barrier. *J Clin Invest* **106**, 1489-1499.

[28]   Roberts KF, Elbert DL, Kasten TP, Patterson BW, Sigurdson WC, Connors RE, Ovod V, Munsell LY, Mawuenyega KG, Miller-Thomas MM, Moran CJ, Cross DT, 3rd, Derdeyn CP, Bateman RJ (2014) Amyloid-beta efflux from the central nervous system into the plasma. *Ann Neurol* **76**, 837-844.

[29]   Lee Y-H, Martin JM, Maple RL, Tharp WG, Pratley RE (2009) Plasma Amyloid-β peptide levels correlate with adipocyte amyloid precursor protein gene expression in obese individuals. *Neuroendocrinology* **90**, 383-390.

[30]   Meakin PJ, Coull BM, Tuharska Z, McCaffery C, Akoumianakis I, Antoniades C, Brown J, Griffin KJ, Platt F, Ozber CH, Yuldasheva NY, Makava N, Skromna A, Prescott A, McNeilly AD, Siddiqui M, Palmer CN, Khan F, Ashford ML (2020) Elevated circulating amyloid concentrations in obesity and diabetes promote vascular dysfunction. *J Clin Invest* **130**, 4104-4117.

[31]   Wang R, Li JJ, Diao S, Kwak YD, Liu L, Zhi L, Bueler H, Bhat NR, Williams RW, Park EA, Liao FF (2013) Metabolic stress modulates Alzheimer's beta-secretase gene transcription via SIRT1-PPARgamma-PGC-1 in neurons. *Cell Metab* **17**, 685-694.

[32]   Bao J, Liang Z, Gong X, Yu J, Xiao Y, Liu W, Wang X, Wang J-Z, Shu X (2021) High Fat Diet Mediates Amyloid-β Cleaving Enzyme 1 Phosphorylation and SUMOylation, Enhancing Cognitive Impairment in APP/PS1 Mice. *Journal of Alzheimer's Disease*, 1-14.

[33]   Takechi R, Galloway S, Pallebage-Gamarallage MM, Lam V, Mamo JC (2010) Dietary fats, cerebrovasculature integrity and Alzheimer's disease risk. *Prog Lipid Res* **49**, 159-170.

[34]   Mamo JC, Jian L, James AP, Flicker L, Esselmann H, Wiltfang J (2008) Plasma lipoprotein beta-amyloid in subjects with Alzheimer's disease or mild cognitive impairment. *Ann Clin Biochem* **45**, 395-403.

[35]   Pallebage-Gamarallage MM, Galloway S, Johnsen R, Jian L, Dhaliwal S, Mamo JC (2009) The effect of exogenous cholesterol and lipid-modulating agents on enterocytic amyloid-beta abundance. *Br J Nutr* **101**, 340-347.

[36]   Galloway S, Jian L, Johnsen R, Chew S, Mamo JC (2007) beta-amyloid or its precursor protein is found in epithelial cells of the small intestine and is stimulated by high-fat feeding. *J Nutr Biochem* **18**, 279-284.

[37]   Galloway S, Takechi R, Nesbit M, Pallebage-Gamarallage M, Lam V, Mamo JCL (2019) The differential effects of fatty acids on enterocytic abundance of amyloid-beta. *Lipids Health Dis* **in press**.

[38]   Pallebage-Gamarallage M, Takechi R, Lam V, Elahy M, Mamo J (2016) Pharmacological modulation of dietary lipid-induced cerebral capillary dysfunction: Considerations for reducing risk for Alzheimer's disease. *Crit Rev Clin Lab Sci* **53**, 166-183.

[39]   Pallebage-Gamarallage MM, Galloway S, Takechi R, Dhaliwal S, Mamo JC (2012) Probucol suppresses enterocytic accumulation of amyloid-beta induced by saturated fat and cholesterol feeding. *Lipids* **47**, 27-34.

[40]   Takechi R, Galloway S, Pallebage-Gamarallage MM, Lam V, Dhaliwal SS, Mamo JC (2013) Probucol prevents blood-brain barrier dysfunction in wild-type mice induced by saturated fat or cholesterol feeding. *Clin Exp Pharmacol Physiol* **40**, 45-52.

[41]   Takechi R, Pallebage-Gamarallage MM, Lam V, Giles C, Mamo JC (2014) Long-term probucol therapy continues to suppress markers of neurovascular inflammation in a dietary induced model of cerebral capillary dysfunction. *Lipids Health Dis* **13**, 91.

[42]   Montefusco L, D'Addio F, Loretelli C, Ben Nasr M, Garziano M, Rossi A, Pastore I, Plebani L, Lunati ME, Bolla AM, Porta MD, Piuri G, Rocchio F, Abdelsalam A, Assi E, Barichella M,

Maestroni A, Usuelli V, Loreggian L, Muzio F, Zuccotti GV, Cazzola R, Fiorina P (2021) Anti-inflammatory effects of diet and caloric restriction in metabolic syndrome. *J Endocrinol Invest* **44**, 2407-2415.

[43]   Ghachem A, Prud'homme D, Rabasa-Lhoret R, Brochu M (2017) Effects of a 6-month caloric restriction induced-weight loss program in obese postmenopausal women with and without the metabolic syndrome: a MONET study. *Menopause* **24**, 908-915.

[44]   Koudinov AR, Berezov TT, Kumar A, Koudinova NV (1998) Alzheimer's amyloid beta interaction with normal human plasma high density lipoprotein: association with apolipoprotein and lipids. *Clin Chim Acta* **270**, 75-84.

[45]   Shih YH, Tsai KJ, Lee CW, Shiesh SC, Chen WT, Pai MC, Kuo YM (2014) Apolipoprotein C-III is an amyloid-beta-binding protein and an early marker for Alzheimer's disease. *J Alzheimers Dis* **41**, 855-865.

[46]   Lee HG, Casadesus G, Zhu X, Joseph JA, Perry G, Smith MA (2004) Perspectives on the amyloid-beta cascade hypothesis. *J Alzheimers Dis* **6**, 137-145.

[47]   Tolar M, Abushakra S, Sabbagh M (2020) The path forward in Alzheimer's disease therapeutics: Reevaluating the amyloid cascade hypothesis. *Alzheimers Dement* **16**, 1553-1560.

[48]   Karikari TK, Pascoal TA, Ashton NJ, Janelidze S, Benedet AL, Rodriguez JL, Chamoun M, Savard M, Kang MS, Therriault J, Scholl M, Massarweh G, Soucy JP, Hoglund K, Brinkmalm G, Mattsson N, Palmqvist S, Gauthier S, Stomrud E, Zetterberg H, Hansson O, Rosa-Neto P, Blennow K (2020) Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: a diagnostic performance and prediction modelling study using data from four prospective cohorts. *Lancet Neurol* **19**, 422-433.

[49]   Tissot C, A LB, Therriault J, Pascoal TA, Lussier FZ, Saha-Chaudhuri P, Chamoun M, Savard M, Mathotaarachchi SS, Bezgin G, Wang YT, Fernandez Arias J, Rodriguez JL, Snellman A, Ashton NJ, Karikari TK, Blennow K, Zetterberg H, De Villers-Sidani E, Huot P, Gauthier S, Rosa-Neto P,

Alzheimer's Disease Neuroimaging I (2021) Plasma pTau181 predicts cortical brain atrophy in aging and Alzheimer's disease. *Alzheimers Res Ther* **13**, 69.

[50] Shen XN, Huang YY, Chen SD, Guo Y, Tan L, Dong Q, Yu JT, Alzheimer's Disease Neuroimaging I (2021) Plasma phosphorylated-tau181 as a predictive biomarker for Alzheimer's amyloid, tau and FDG PET status. *Transl Psychiatry* **11**, 585.

[51] Smirnov DS, Ashton NJ, Blennow K, Zetterberg H, Simren J, Lantero-Rodriguez J, Karikari TK, Hiniker A, Rissman RA, Salmon DP, Galasko D (2022) Plasma biomarkers for Alzheimer's Disease in relation to neuropathology and cognitive change. *Acta Neuropathol* **143**, 487-503.

[52] Simren J, Leuzy A, Karikari TK, Hye A, Benedet AL, Lantero-Rodriguez J, Mattsson-Carlgren N, Scholl M, Mecocci P, Vellas B, Tsolaki M, Kloszewska I, Soininen H, Lovestone S, Aarsland D, AddNeuroMed c, Hansson O, Rosa-Neto P, Westman E, Blennow K, Zetterberg H, Ashton NJ (2021) The diagnostic and prognostic capabilities of plasma biomarkers in Alzheimer's disease. *Alzheimers Dement* **17**, 1145-1156.

[53] Pegueroles J, Pane A, Vilaplana E, Montal V, Bejanin A, Videla L, Carmona-Iragui M, Barroeta I, Ibarzabal A, Casajoana A, Alcolea D, Valldeneu S, Altuna M, de Hollanda A, Vidal J, Ortega E, Osorio R, Convit A, Blesa R, Lleo A, Fortea J, Jimenez A, Alzheimer's Disease Neuroimaging I (2020) Obesity impacts brain metabolism and structure independently of amyloid and tau pathology in healthy elderly. *Alzheimers Dement (Amst)* **12**, e12052.

[54] Koga S, Kojima A, Ishikawa C, Kuwabara S, Arai K, Yoshiyama Y (2014) Effects of diet-induced obesity and voluntary exercise in a tauopathy mouse model: implications of persistent hyperleptinemia and enhanced astrocytic leptin receptor expression. *Neurobiol Dis* **71**, 180-192.

[55] Janelidze S, Teunissen CE, Zetterberg H, Allue JA, Sarasa L, Eichenlaub U, Bittner T, Ovod V, Verberk IMW, Toba K, Nakamura A, Bateman RJ, Blennow K, Hansson O (2021) Head-to-Head Comparison of 8 Plasma Amyloid-beta 42/40 Assays in Alzheimer Disease. *JAMA Neurol* **78**, 1375-1382.

[56]   Deurenberg P, Yap M, van Staveren WA (1998) Body mass index and percent body fat: a

          meta analysis among different ethnic groups. *Int J Obes Relat Metab Disord* **22**, 1164-1171.

[57]   Nuttall FQ (2015) Body Mass Index: Obesity, BMI, and Health: A Critical Review. *Nutr Today*

          50, 117-128.

BMI Groups	Lean (n=42)	Overweight (n=41)	Obese (n=23)
Age (years) (range)	32.40 ± 11.24^a (18-54)	35.88 ± 12.66^a (19-44)	57.57 ± 9.80^b (29-66)
Gender (% Male)	100	95.12	60.87
BMI (kg/m^2) (range)	22.91 ± 1.40^c (19.29-24.99)	27.25 ± 1.58^d (25.00-29.88)	33.96 ± 5.12^e (30.04-51.07)

Table 1: Cohort baseline characteristics

All variables are presented as mean ± SD with range.

Non-matching superscript lettering indicates significant difference. Alphabetical letters indicate

statistical significance at *p<0.05*

Variable	Baseline	wk-12	Change (%)[a]	p value
Gender (% male)	58.33			
Age (years)	60.17±4.22	-	-	-
(range)	(52-66)			
BMI (kg/m^2)	32.45 ± 3.86	31.18 ± 4.13	−4.02 ± 2.25	0.0005
(range)	(27.44-42.20)	(26.00-41.53)	(−9.58-−0.43)	
BMI (overweight: obese)	2:10	5:7		

Table 2: Age and BMI at baseline and wk-12

All values are presented as mean ± SD with range

[a]Percent change from baseline to 12-week follow-up

Fig. 1. Violin plots show the plasma concentrations of A) Aβ40, B) Aβ42, and C) Aβ42/40 in people with lean, overweight, or obese BMIs. The median value is indicated by the solid line, and the quartiles are represented by the dashed lines.

Fig. 2. The scatter plot shows the linear correlations between BMI and plasma concentrations of A) Aβ40, B) Aβ42, and C) Aβ42/40 ratio. D) Shows the Principal Component Analysis loading plot of BMI, Aβ40, Aβ42 and Aβ42/40, and E) Principal Component Analysis score plot shows the scattering of Aβ42/40 in lean, overweight, and obese BMI groups.

Fig. 3. The bar graphs show adjusted plasma concentrations of A) Aβ$_{40}$, B) Aβ$_{42}$, and C) Aβ$_{42/40}$ ratio in people with lean, overweight, and obese BMIs. All values are shown as mean and standard error of the mean.

Fig. 4. Before-after plots show the plasma concentrations of A) Aβ$_{40}$, B) Aβ$_{42}$, and C) Aβ$_{42/40}$ in people who entered the calorie-restriction weight loss program.

Fig. 5. The scatter plot shows the linear correlations between the percentage of weight loss from baseline and percent change in plasma concentrations of A) Aβ_{40}, B) Aβ_{42}, and C) A$\beta_{42/40}$ ratio.