Elucidation of antiviral mechanisms of natural therapeutic molecules Herbacetin and

Caffeic acid phenethyl ester against chikungunya and dengue virus

Mandar Bhutkar[#], Ruchi Rani[#], Aditi Kothiala, Supreeti Mahajan, Bhairavnath Waghmode,

Ravi Kumar, Vishakha Singh, Debabrata Sircar, Pravindra Kumar, Shailly Tomar*

Department of Biosciences and Bioengineering, Indian Institute of Technology Roorkee,

Uttarakhand, India

[#] Both the authors have contributed equally

*Corresponding author address: Department of Biosciences and Bioengineering, Indian

Institute of Technology Roorkee, Roorkee-247667, Uttarakhand, India; Email id:

shailly.tomar@bt.iitr.ac.in

*Mandar Bhutkar and Ruchi Rani contributed equally to this work. Author order was
determined alphabetically.*

Abstract

Arthropod-borne viruses of the alphavirus and flavivirus genera are human pathogens of significant concern, and currently, no specific antiviral treatment is available for these viruses. This study has elucidated the antiviral mechanisms of natural small molecules against the dengue (DENV) and chikungunya virus (CHIKV). Herbacetin (HC) and Caffeic acid phenethyl ester (CAPE) depleted polyamine levels in Vero cells, which has been demonstrated by thin-layer chromatography (TLC). As polyamines play an essential role in the replication and transcription of RNA viruses, the depletion of polyamines by HC and CAPE was anticipated to inhibit the virus replication. To test this hypothesis, HC and CAPE were evaluated for antiviral activities using a cell-based virus yield assay by quantitative reverse transcription-polymerase chain reaction (qRT-PCR), plaque reduction assay, and immunofluorescence assay (IFA). HC and CAPE displayed potent inhibition with EC_{50} of 463 nM and 0.417 nM for CHIKV and 8.5 μM and 1.15 μM for DENV, respectively. However, the addition of exogenous polyamines did not ultimately rescue the virus titer in both CHIKV and DENV-infected cells. This finding suggested additional antiviral mechanisms for HC and CAPE. Further, *in silico* analysis revealed that HC and CAPE may directly target the viral methyltransferases (MTase) of CHIKV and DENV. The inhibition of virus-specific MTases by HC and CAPE was confirmed using purified viral MTase of CHIKV and DENV. Altogether, the dual targeting of the host pathway and the viral MTase using potent natural antiviral molecules is expected to facilitate the development of effective biological therapies.

IMPORTANCE

Vector-borne diseases caused by DENV and CHIKV has a catastrophic impact on human health worldwide. There are no effective vaccines and antiviral drugs present in the market till date against these viruses. In the present study, natural small molecules have been identified as antivirals against DENV and CHIKV. These molecules directly inhibit the virus and impede

the synthesis of essential host molecules required for efficient virus replication. Hence, these

broad-spectrum antivirals have high therapeutic potential.

Keywords

Dengue virus, Chikungunya virus, Polyamine depletion, Antiviral assay, virtual screening,
and MD simulation.

Introduction

Dengue virus (DENV) and chikungunya virus (CHIKV) are arthropod-borne viruses belonging

to the *Flaviviridae* and *Togaviridae* families, respectively [1]. Both viruses cause febrile

illnesses showing fever, arthralgia, joint pain, rash, and headache. These enveloped viruses

have a positive-sense single-stranded RNA genome (+ssRNA) [2]. It is estimated that ~100-

400 million DENV infections occur annually [3], [4]. Further, in 2021, ~ 0.3 million CHIKV

cases occurred worldwide, of which a majority were reported in Brazil and India [6], [7]. In

India, co-infections with DENV and CHIKV occur prevalently in various regions during the

monsoon season [6], [8], [9].

Advances in understanding the reliance of viruses on host cell pathways have rekindled interest

in targeting the host metabolic pathways for antiviral therapeutics. Viral genomes lack protein

synthesis machinery, and the ability to hijack the host cell's resources for their replication is

crucial for all viruses. Viruses use various strategies for replication by controlling the host

cell's metabolic activities, including exploiting the host lipid pathway, targeting the host

glycolytic pathway, hijacking the host ubiquitination pathways, and using the polyamine

metabolic pathway [10], [11]. Hence, targeting host factors responsible for virus propagation

can be used to develop broad-spectrum antivirals.

Polyamines are essential in the viral life cycle [12], [13]. These are small organic molecules

with multiple ionizable amino moieties and are present in all prokaryotes and eukaryotes. A

typical polyamine content is indispensable for cell survival and is involved in various essential

cellular functions. There are three biogenic polyamines in mammals, namely, putrescine (put),

spermine (spm), and spermidine (spd), which are produced from ornithine according to the

biosynthetic pathway [15]. Briefly, the enzymatic cleavage of arginine by arginase produces

ornithine. Ornithine is converted into putrescine, the first biosynthetic polyamine, by the

enzyme ornithine decarboxylase. Further, spermidine and spermine are produced sequentially

by respective synthase enzymes. There is enough evidence to suggest that polyamines are

essential for various stages of the virus life cycle, including genome replication, virus protein

translation, and genome packaging. Ornithine decarboxylase (ODC) inhibition results in

polyamine depletion, which has been shown to depict antiviral activity against CHIKV, DENV,

ZIKA virus (ZIKV), etc [12], [13]. Difluoromethylornithine (DFMO) is a well-known

irreversible ODC inhibitor, but it is associated with hearing loss, may suffer antiviral resistance

[16]. Also, DFMO requires higher dosage concentrations to show antiviral activity [12], [13].

Moreover, DFMO can induce the uptake of putrescine and spermidine[17]. Hence, an ODC

inhibitor with antiviral activity at lower concentrations would be valuable for developing

broad-spectrum, highly effective antiviral agents. Natural therapeutic molecules are expected

to have low toxicity. Thus, understanding the link between the polyamine biosynthetic pathway

and the replication of DENV and CHIKV in infected cells suggests a potential mechanism for

antiviral therapies.

Alphaviruses contain four non structural proteins (nsPs), namely nsP1, nsP2, nsP3, and nsP4,

which play an essential role in the life cycle of the viruses and are produced when a viral

protease, i.e., nsP2, cleaves the viral polyprotein precursor post-translationally [18]. The

capping of viral RNA, performed by nsP1, is a crucial step in the life cycle of an alphavirus.

The viral genome cap structure has numerous biological roles, such as helping the eukaryotic

translational initiation factor 4E (eIF4e) recognize the RNA for initiating translation and

protecting the mRNA from cellular exonucleases [19]. Further, the cap structure also helps the

viral genome to escape recognition from the host innate immune system pathways such as

retinoic acid-inducible gene I (RIG-I) and melanoma differentiation-associated protein 5 (MDA5) [20]. Cellular enzyme capping mechanisms usually involve the methylation of Guanosine triphosphate (GTP) after transferring it to the 5' end of the RNA. However, the molecular mechanism of viral mRNA capping in alphaviruses is distinct. The Alphaviral capping enzyme nsP1 performs the capping in two significant steps. First, S-adenosyl methionine (SAM) acts as a methyl donor to methylate GTP by transferring a methyl group to the seventh position to form m7GMP, where S-adenosyl homocysteine (SAH) is generated as a by-product. This is known as the methyltransferase (MTase) step of the capping reaction [21]. Subsequently, the methylated m7GMP is covalently attached to the nsP1 enzyme via a single bond with His37 during the second step, known as the guanylation (GT) step [22]. In addition, a few antivirals targeting alphavirus nsP1 have been identified to treat these infections. Recently, the availability of three-dimensional protein structures of alphavirus nsP1 (6Z0U) has increased the utility of high-throughput computational screening of small molecules in identifying novel drug candidates [23]. This strategy exemplifies a logical progression for potential drug discovery against the nsP1 alphavirus.

On the other hand, flaviviruses contain a single open reading frame, which encodes a polyprotein processed by viral and cellular proteases into three structural and seven nonstructural proteins (NS1 NS2A, NS2B, NS3, NS4A, NS4B, and NS5). The flavivirus's plus-stranded RNA genome contains a 5' terminal cap 1 structure (m7GpppAmG). It consists of an N-terminal MTase and a C-terminal RNA-dependent-RNA polymerase (RdRp) domain [24]. The structure of DENV-2 MTase suggests that flavivirus NS5 MTase belongs to a family of SAM-dependent MTases [25]. Moreover, the Rossmann fold is a βαβ fold super-secondary structure that forms the catalytic core responsible for SAM binding site in the DENV NS5 MTase domain and CHIKV nsP1 protein [26]–[29]. A highly conserved acidic residues interacts with ribose moiety, and the glycine-rich loop region interacts with the cofactor of the ribose moiety [27].

Herbacetin (HC) is a naturally occurring flavonoid compound that has multiple

pharmacological uses. It is also an established ODC inhibitor that helps lower the levels of

polyamines in cells [30]. Further, it has been shown that flavonoids play a role in reducing the

levels of nsP1 in CHIKV, and the antiviral activity of flavonoids against CHIKV has been

established [31]. CAPE is a natural ester of caffeic acid. Various pharmacological activities

such as antiviral, anti-oxidative, and inhibition of human catechol-O-methyltransferase have

been reported for CAPE [32], [33]. Additionally, CAPE is also a known ODC inhibitor like

HC which helps reduce the levels of polyamines in cells [34].

The findings mentioned above open new research avenues for exploiting the potential of HC

and CAPE as antiviral agents that reduce polyamine levels and possibly target virus-specific

enzymes. This study has exhibited the fact that both HC and CAPE were able to significantly

reduce the virus titer of CHIKV and DENV. Moreover, the compounds have a dual impact on

viral replication as they deplete the polyamine levels in the cells and inhibit the viral MTases

essential for the capping of viral RNAs.

Materials and methods

Cell line, Virus isolation, propagation and serotyping

Vero cell line isolated from kidney epithelial cells of an African green monkey, was obtained

from the National Centre for Cell Science (NCCS), Pune, India. Vero cells were used for the

propagation and titration of DENV and CHIKV. The cells were maintained in Dulbecco's

Modified Eagles Medium (DMEM) (Gibco, HiMedia) supplemented with 10% inactivated

fetal bovine serum (FBS) (Gibco, HiMedia) along with 100 U/ml penicillin and 100 mg/ml

streptomycin (Gibco, HiMedia). The cells were maintained at 37 °C with 5% CO_2

supplementation.

Dengue was isolated from the DENV-suspected patient's blood samples. Sera was

aliquoted and tested for the presence of DENV NS1 antigen by J Mitra ELISA kit according to

the manufacturer's instructions. DENV positive samples were used for virus isolation. Initially,

sera were 1:10 diluted in minimum essential medium (MEM) (Gibco, HiMedia) containing 2

% FBS, 100 U/ml penicillin, and 100 mg/ml streptomycin for virus propagation. Then, Serum

samples were added to a confluent monolayer of Vero cells for 90 min with gentle shaking

every 15 min in a 37 °C CO_2 incubator. The inoculum was withdrawn, and the cells were

cultured for 5-6 days in fresh MEM with 2% FBS and antibiotics until the appearance of

cytopathic effect (CPE). After 5-6 days, the supernatant was harvested and serially diluted for

infection of Vero monolayer cells for 90 min and overlaid with MEM containing 1 % agarose

for the plaque purification procedure. Single plaques were picked and resuspended in MEM

and were further propagated in Vero cells. Then DENV NS1 ELISA positive supernatants

showing CPE was collected and used for further study.

To confirm the DENV and its serotype, RNA from the supernatant of cell culture

infected with the virus was isolated using the Trizol (Sigma) method described by the

manufacturer. The RNA was then reverse-transcribed into complementary DNA (cDNA) using

the PrimeScript cDNA Synthesis Kit (Takara). Synthesized cDNA was utilized as a template

for polymerase chain reaction (PCR) [35] and the amplification product was analyzed by gel

electrophoresis in 1 % agarose gels stained with ethidium bromide. The sample was sent for

DNA sequencing and the serotype of DENV was confirmed by matching the obtained sequence

against the nucleotide database.

CHIKV (Accession No. KY057363.1.) was propagated and titrated in a Vero cell line

using the protocol reported by Singh et al., 2018 and then stored at −80 °C for further

experiments [36].

179
Preparation of HC, CAPE, and DFMO stock solutions

HC, CAPE, and DFMO were purchased from Cayman, USA. For experiments, the stock

solutions (HC: 165.45mM; CAPE: 20mM) for both compounds were prepared in dimethyl

sulfoxide (DMSO) (Sigma- Aldrich) and filtered through a 0.2 μm size syringe filter

(Millipore). The 100 mM DFMO dilution was prepared in sterile tissue culture grade water,

that was used as a positive control for the polyamine-related experiments. At the time of usage,

further dilutions were prepared in culture media. Similarly, individual polyamines (put, spm,

and spd) (Alfa-Aeser, USA) stocks were diluted in water and used as designated.

Cell Cytotoxicity Assay

Different concentrations of HC and CAPE were evaluated for cytotoxicity on Vero cells using

3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Himedia) assay.

Before treatment, Vero cells were seeded in a 96 well plate, and at 90% confluency, the media

were removed, and different dilutions of compounds were added into each well for 24 h. After

incubation, the inoculum was removed and fresh maintenance media were added for 2 h.

Further fresh compound dilutions were added to each well and kept for 24 h. Subsequently, 20

μL/well of MTT (5mg/ml) was added and incubated for 4 h at 37 °C in 5% CO_2. Upon

incubation, 110 μL/well of DMSO was added to dissolve formazan crystals. Plates were read

at a wavelength of 570 nm using cytation 3 multi-mode plate reader (BioTek Instruments, Inc.).

The average absorbance of 0.1 % DMSO treated cells was used as cell control. In the second

set of experiments, after completion of 24 h post-treatment, fresh maintenance media were

added and kept for 4 days. The cell viability of the treated well was compared with the cell

control, and the concentration which showed >90% viability after compound treatment was

considered non-toxic. The experiment was performed in triplicates.

Assessment of Antiviral Activity

Vero cells were seeded onto a 24-well plate at a cell density of 1.0×10^5 cells/well. Cells were

treated with compounds 12 h before infection. With a multiplicity of infection (MOI) of 0.1,

the cell monolayer was infected with the virus with gentle shaking every 15 min for 2 h. The

inoculum was removed, and the cell monolayer was washed twice to ensure no chance of

secondary infection. Compounds were added in the post-infection (pi) 2% DMEM media and

incubated for 24 h. After 24 hpi, the supernatant was collected for CHIKV antiviral experiment.

On the other hand, fresh maintenance media was added to the DENV antiviral experiment for

4 days. Then, the supernatant was collected to determine virus titre via plaque-forming assay.

Polyamine addition to rescue virus

214 In the polyamine addition assay, drug pre-treatment was given for 24 h at various

215 concentrations such as HC (200 µM), CAPE (25 µM for CHIKV and 2.5 µM for DENV),

216 DFMO (1000 µM). After infection with CHIKV and DENV, cells were treated with 1 µM

217 biogenic polyamines (put, spd, spm) for 24 h and 120 h, respectively. Later, the supernatant

218 was collected to determine the virus titer by plaque-forming assay.

Plaque Reduction Assay

Vero cells were seeded at 1.0×10^5 cells/well in a 24-well plate in complete media before

infection. The supernatant was 10-fold serially diluted in maintenance media and was

inoculated on ~80-90% confluent cells. Plates were incubated for 2 h with a gentle shaking

every 15 min for virus adsorption at 37 °C with a 5% CO_2 incubator. After adsorption, overlay

media and maintenance media added in 1:1 dilution and plates were further incubated for 2

days and 3 days for CHIKV and DENV respectively at 37 ˚C in 5% CO_2 incubator. Following

that, cells were stained with 1% crystal violet to count the number of CHIKV plaques and

immunostaining for DENV-infected cells, as described below.

Immunostaining was performed at room temperature. After incubation, overlay media was

removed from the plates, and the cell monolayer was fixed with 3.7% formaldehyde solution

for 30 min. Cells were washed 3 times with PBST (0.02% Tween-20 in Phosphate buffer

saline). After that, cells were permeabilized with 0.2% triton X-100 in PBS for 7 min. Cells

were washed three times with PBST and incubated with 3% Skim milk for 30 min. Then, cells

were washed three times with PBST and incubated with 1:500 diluted flavivirus group antibody

(Genetex) for 2 h. Cells were washed three times with PBST and incubated with secondary

antibody (Goat anti-mouse IgG HRP, 1:1500 dilution) for 1 h. After washing 2 times with

PBST and three times with PBS, cells were stained with True Blue Peroxidase substrate (KPL,

Sera Care, MA, USA) and incubated in the dark for 30 min to develop blue colour staining of

virus-infected cells, and foci were counted [37].

Immunofluorescence Assay

Vero cells were seeded onto a 6-well plate at a cell density of 1×10^6 cells/ well. Cells were

treated with compounds, as mentioned earlier. The cells were washed three times with PBS

and then fixed with methanol: acetone (1:1) for 1 h at room temperature, followed by

permeabilization with 0.1% Triton-X-100. After washing, cells were incubated with antibodies

against CHIKV and DENV (anti-alphavirus 1:100, SANTA Cruz Biotechnology Inc, 1:500

diluted flavivirus group antibody (Genetex Inc).) for 1 h and then incubated with fluorescein

(FITC)-conjugated secondary anti-mouse antibody (1:250, Sigma) for 30 min at 37 °C. The

plate was then rinsed with PBS and counter-stained with 4′,6- diamidino-2-phenylindole DAPI

(Sigma) for 15 min. Then, the image was captured under a fluorescence microscope (EVOS,

Thermofisher). All experiments were repeated thrice to verify the results.

Quantitative Real-Time PCR (qRT-PCR)

Vero cells were seeded onto 6 well plates prior to the day of infection. HC and

CAPE compounds treatment and infection were the same as described in the assessment of

antiviral activity. After the termination of an assay, trizol was added to the plate. RNA was

purified according to the manufacturer's protocol. Purified RNA was quantified and used for

cDNA preparation using the PrimeScript 1st strand cDNA Synthesis Kit (Takara) with 1 μg of

extracted RNA. The forward and reverse primers used for amplification are used as previously

described in [35] and [38]. qRT-PCR was used to determine the effect of HC and CAPE on

CHIKV and DENV replication by counting the number of copies of RNA. All experiments

were carried out three times in triplicates.

Polyamine determination by Thin-layer chromatography (TLC)

Vero cells were treated with CAPE and DFMO for 12 h pretreatment and 24 h post treatment as mentioned in antiviral assay protocol. Similarly, HC treated cells were collected after 12, 24, and 36 h of treatment. Further, cells trypsinized and centrifuged for all samples. Polyamines were separated by TLC as previously described [39]. Wash with PBS. Sonicate the cells in 500 µl of 2% (V/V) perchloric acid. Sonication was performed at 4 ºC (20 kHz, 2Ω, 10-sec pulse, 30-sec rest). Cell homogenates obtained are then stored at 4 ºC for 24 h. After 24 h, samples are centrifuged at ~11,500 g for 30 min at 4 ºC. An equal ratio of supernatant dansyl chloride (5 mg/ml) (Alfa-Aesar, USA) was added in acetone and saturated sodium bicarbonate. Samples were incubated in the dark overnight at room temperature. Excess dansyl chloride was cleared by incubating the reaction with 150 mg/ml proline (HiMedia). Dansylated polyamines were extracted with 100 µL toluene. 10 µL of the sample was added in small spots to the TLC plate by automated sprayer (Silica gel matrix; Millipore) and exposed to ascending chromatography with cyclohexane:ethyl acetate (2:3). The plate was dried and visualized via exposure to UV.

Multiple Sequence Alignment (MSA)

The amino acid sequence of nsP1 protein of alphaviruses was compared with CHIKV nsP1 as a reference point using Clustal Omega [40] to check if the key residues involved in the capping of viral RNA were conserved across different viruses. The MSA was performed for Venezuelan equine encephalitis virus (VEEV Gene Bank Id:AWT57896), (CHIKV Gene Bank Id:AWQ38330), Ross River virus (RRV Gene Bank Id:AAU89533), Sindbis virus (SINV Gene Bank Id:ABL84688), Aura Virus (AURA Gene Bank Id:QVM79755), Middelburg virus (MIDV Gene Bank Id:QOW97289), Barmah Forest virus (BFV Gene Bank Id:QOY44469), Madariaga virus (MADV Gene Bank Id:QTC33397), and Mayaro virus (MAYV Gene Bank Id:AZM66145) from the Alphavirus family. The sequence alignment profile of the selected nsP1 sequences was performed via Clustal Omega tool and analyzed by a graphical colored depiction using ESPript 3.0 [40].

286   Similarly, MSA was performed for the MTase domain of NS5 from dengue 3 Virus (DENV 3

287   Gene Bank Id:4CTJ_1), Zika Virus (ZIKV Gene Bank Id:5WZ2_1), West Nile Virus (WNV

288   Gene Bank Id:2OY0_1), Yellow Fever Virus (YFV Gene Bank Id:3EVA_1), and Japanese

289   Encephalitis Virus (JEV Gene Bank Id:4K6M_1) from the flavivirus family. And the analysis

290   was carried out in the same way as for DENV.

Molecular Docking of HC and CAPE into MTases

The cryo-electron microscopy (cryo-EM) structure of CHIKV nsP1 has been published at 2.6

Å and 2.9 Å resolution (6Z0V, 6Z0U, and 7DOP) [23]. The structure reveals the catalytic

binding site in nsP1, where SAM and GTP bind during RNA capping [41], [42]. The protein

data bank (PDB) entry lacked atomic coordinates for multiple flexible loops; therefore, SWISS

Model [44] was utilized to create a model of nsP1 using 6Z0U as a template and 6Z0V as the

sequence prior to docking. AutoDock Vina software algorithm [45] was utilized in all the

docking with the optimized model as the target. The structures of ligands i.e. HC, CAPE, and

SAM, were obtained from the PubChem site in .sdf format and then converted to .pdbqt format

using OpenBabel in PyRx 0.8 algorithm [46], [47]. In AutoDock Vina, centre grid dimensions

were specified as $29.50\,Å \times 23.35\,Å \times 27.91\,Å$, centered at $X = 56.15$, $Y = 135.85$, and

$Z = 88.22$, for the CHIKV nsP1.

Similarly, the 3D structure of DENV NS5 MTase (PDB ID 4V0R) was obtained from PDB

[48] The centre grid dimensions were specified as $17.95\,Å \times 24.98\,Å \times 17.13\,Å$, centered at X

$= 27.91$, $Y = 127.90$, and $Z = 20.88$, located at the active site of NS5 MTase. The conformations

with the best binding energy were further analyzed in PyMol and LigPlot[+] software [50], [51].

Molecular Dynamics (MD) Simulation

308   nsP1 and SAM, HC, and CAPE were subjected to MD simulation studies to assess the

309   flexibility and stability of protein-ligand interactions. The same study was carried out for

310   DENV NS5 MTase with SAM, HC, and CAPE. For this purpose, GROMACS 5.4.1 suite was

used to carry out all simulation studies using the GROMOS96 43a1 force field on a LINUX-based workstation [52]. Ligand parameters and topology files were generated using the PRODRG server [53]. Furthermore, ions and water molecules were added to neutralize the whole cubic system for solvation. The energy minimization step was performed using the steepest descent method, followed by a two-phased equilibration of a constant number of particles, volume, and temperature (NVT), a constant number of particles, pressure, and temperature (NPT). NVT equilibration was done at 300K with a short-range electrostatic cut-off of 1.2 nm, and temperature regulation was done using the Berendsen temperature coupling method. Further, the next phase of equilibration NPT was performed, and coordinates were generated every 1 ps. Finally, 50 ns MD production run was performed with an integration time frame of 2fs, and the trajectories were generated after every 10 ps. The conformations generated during the production step were used for calculating root mean square deviation (RMSD) of protein-ligand complexes.

Expression and purification of DENV 3 NS5 MTase Domain

DENV 3 RNA was isolated from the supernatant of cell culture infected cells using the Trizol (Sigma) method described by the manufacturer. The RNA was then reverse-transcribed into complementary DNA (cDNA) using the PrimeScript cDNA Synthesis Kit (Takara). The DENV 3 cDNA was subsequently amplified by PCR, using the following primers (amino acid residues 1-278), NS5 MTase forward, 5′- ATGCGCTAGCGGAACAGGTTCACAAG-3′, and NS5 MTase reverse 5′- CGCACTCGAGTCAATTAACATGTCGAGTT-3′, the sequences underlined are the recognition sites of the restriction enzymes NheI and XhoI respectively. After digestion with NheI and XhoI, the PCR product was purified with Qiagen gel extraction kit as per mentioned kit protocol and inserted between the NheI and XhoI sites of the vector pET28c (+). The recombinant vector pET28c (+) NS5 MTase was identified by restriction digestion and the DENV3-NS5 MTase insert was verified by sequencing. E. coli DH5α was

used for amplification of the recombinant plasmid and E. coli BL21 (DE3) was transformed

for induced expression of the His-tagged DENV3-NS5 MTase protein.

A single colony of E. coli BL21 (DE3) cells transformed with the recombinant expression

plasmid pET28c (+) NS5 MTase was inoculated in a tube containing 10 ml of LB medium with

50 μg/ml kanamycin, and cultured overnight at 37 °C in a shaking incubator. The culture was

transferred to a 2 litre flask containing 1000 ml of the same medium. The flask was shaken at

37 °C until the Optical Density 600 of the culture was about 0.6. The expression of DENV3-

NS5 MTase protein was induced by the addition of 0.6 mM of isopropyl β-d-thiogalactoside

(IPTG). After induction culture was incubated overnight in incubator shaker at 16 °C. Further,

cells were harvested by centrifugation at 5,000g, 4 °C for 10 min. The supernatant was

discarded and the cell pellet were then resuspended in lysis buffer (50 mM Tris-HCl (pH 7.0),

300 mM NaCl, 10% Glycerol, 1% NP-40) and then disrupted by French press system (Constant

Systems Ltd, Daventry, UK). The lysed cells were centrifuged at 12,000g, 4 °C for 30 min.

The clarified supernatant was passed through the pre-equilibriated nickel-nitrilotriacetic acid

(Ni-NTA) beads (Qiagen, Germantown, MD, USA) in a gravity flow column, followed by

washing the column with wash buffer with increasing imidazole concentrations. The protein of

interest was then eluted with 10 ml of elution buffer (250 mM imidazole in lysis buffer). The

eluted fractions were analyzed by SDS-PAGE (12 %) and Coomassie Blue staining. Eluted

fractions further dialyzed against 1000 ml of a buffer containing (50 mM Tris–HCl, pH 7.5, 50

mM NaCl, 1 mM $MgCl_2$,10 % glycerol, and 1 mM DTT) overnight at 4 °C. The dialyzed

protein was concentrated to the required concentration with amicon centrifugal filters (10,000

MWCO Millipore, Burlington, MA, USA) and frozen in liquid nitrogen and stored at −80 °C

till its use.

CE Based DENV 3 NS5 MTase inhibition assay

To test DENV 3 NS5 MTase inhibition by HC and CAPE, NS5 MTase enzyme reaction was

performed in a reaction mixture consisting of 50 mM Tris buffer (pH 7.5), 10 mM KCl, 2 mM

DTT, 2 mM $MgCl_2$, 0.3 mM SAM and 4 mM GTP along with 1 μM protein at 37 °C for 1 h.

Different concentrations of CAPE (50 μM, 75 μM, 100 μM, 250 μM, 500 μM, and 1000 μM)

and HC (25 μM, 50 μM, 75 μM, 100 μM, 200 μM and 300 μM) was used for the inhibition

assay. MTase reaction was also done with the inactive capsid CHIKV to be used as a negative

control. After incubation reaction stopped by adding acetonitrile in 1 : 2 (vol/vol). 1mM

caffeine was used as internal control [22], [54]. The mixture was vortex-mixed for 15 s, and

protein was precipitated for 20 min at 18,500 g. Supernatant was transferred to sample vials

for CE analysis performed as mentioned in Mudgal et. al 2020. All reactions were done in

duplicate [22], [54].

CHIKV nsP1 enzyme inhibition assay

CHIKV nsP1 protein was purified using the same protocol as reported earlier [22], [55], [56].

To test nsP1 inhibition by HC and CAPE, nsP1 enzyme reaction was performed in a reaction

mixture consisting of 50 mM Tris buffer (pH 7.3), 10 mM NaCl, 5 mM DTT, 2 mM $MgCl_2$,

100 μM SAM and 100 μM GTP along with 20 μM protein at 25 °C for 30 min. Different

concentrations of CAPE (1 μM, 5 μM, 25 μM, 50 μM, 150 μM, and 300 μM) and HC (25 μM,

50 μM, 100 μM, 150 μM and 300 μM) was used for the inhibition assay. CHIKV nsP1-m7GMP

conjugate formation activity of purified nsP1 was detected with the help of ELISA-based nsP1

assay using the same protocol reported earlier [56].

382
383   **Statistical analysis**

384   GraphPad Prism was used to determine IC_{50} values using a sigmoidal dose–response (variable

385   slope) equation. One-way ANOVA was used to determine statistical significance in

386   comparisons of positive control and compound cytotoxicity and antiviral activity measures,

387   followed by Dunnett's tests. When p< 0.05, a difference was considered significant.

388
Results

Antiviral and polyamine depletion activities of HC and CAPE

For this study, DENV serotype was isolated from a clinical sample and DENV 3 serotype was

confirmed by PCR using virus and serotype-specific primer pairs and sequencing. The basic

local alignment search tool (BLAST) tool revealed 99.78% identity of isolated DENV serotype

to DENV3 isolate NU1883 polyprotein (POLY) gene, partial cds (coding sequences).

Similarly, CHIKV (Accession No. KY057363.1.) was propagated in vero cells and used in the

further study The MTT colorimetric assay was performed for testing the cytotoxicity of HC

and CAPE. Both compound led to minimal cytotoxicity, remaining over 90% as shown in

(Figure 1 A-C). Antiviral activities of HC and CAPE at various concentrations were determined

against CHIKV in plaque reduction assays performed in Vero cells using each compound's

maximum non-cytotoxic concentrations (MNCC). By comparing the viral titre of untreated

CHIKV infected cells with HC and CAPE treated CHIKVs infected cells, a dose-dependent

decrease in CHIKV titre was observed for both the compounds (Figure 1 D-E). DFMO, a

known ODC inhibitor was used as a control and it showed less antiviral efficacy against

CHIKV at much higher concentrations of 1000µM as compared to HC and CAPE. HC and

CAPE effectively inhibited CHIKV with EC_{50} values 463 nM and 0.417 nM respectively.

Similarly, a dose-dependent inhibitory effect of HC and CAPE was observed for DENV-3 with

EC_{50} values 8.5 µM and 1.15 µM, respectively (Figure 1 G-H).

Here, qRT-PCR was used for validating the antiviral effect of HC and CAPE by quantifying

DENV and CHIKV RNA levels in the infected cells. qRT-PCR showed a significant (p <

0.0001) reduction in the viral RNA levels when treated with HC and CAPE compounds in an

antiviral assay for CHIKV (Figure 1F) and DENV (Figure 1I). At the mentioned

concentrations, HC and CAPE showed a 23- and 4-fold reduction in DENV, respectively,

compared to the virus control (VC). Similarly, at the same doses, HC and CAPE reduced

CHIKV by 25 and 97 fold, respectively, compared to VC. These results were further

corroborated with IFA results. The results of IFA analysis showed a reduction in CHIKV

(Figure 2A) and DENV (Figure 2B) virus after treatment with HC and CAPE.

Further, TLC experiments were performed to determine the effect of HC and CAPE on

polyamine levels in mammalian cells treated with these compound. Vero cells were treated

with HC, CAPE and DFMO and TLC results confirmed a reduction in the levels of all three

polyamines. Here, put, spd, spm used as TLC controls as shown in (Figure 1J). Polyamine

levels in cells treated with CAPE were significantly lower as compared to DFMO treatment

(Figure 1J). Whereas, HC showed a reduction in polyamine levels with initially 12 h treatment

but later levels of polyamine elevated gradually with time. Overall, CAPE and HC showed

higher polyamine depletion at lower concentrations as compared to the positive control

(DFMO) and the cell control (Figure 1J).

Further, the effects of the addition of exogenous biogenic polyamines were studied. Vero cells

were pre-treated with drugs at various concentrations such as HC (200 µM), CAPE (25 µM for

CHIKV and 2.5 µM for DENV), and DFMO (1000 µM). After infection, 1 µM each of

exogenous polyamines (put, spd, spm) were added. Unlike DFMO treated cells, exogenous

addition of all three polyamines at 1 µM were unable to effectively rescue virus titre after

CAPE and HC treatment for both CHIKV and DENV (Figure 3). Hence, this observation

indicates that CAPE and HC may have other mechanisms that play a role in the antiviral activity

of these compounds against both viruses.

CAPE, a natural ester of caffeic acid, is a known effective inhibitor of catechol-O-

methyltransferase [33]. Additionally, it has been reported that flavonoids are potential antiviral

agents that inhibit CHIKV replication [57], and HC is a naturally occurring flavonoid

compound that has multiple pharmacological uses. Hence, the possibility of HC and CAPE

binding and targeting virus-specific MTases i.e. nsP1 of CHIKV and NS5 of DENV were

further examined.

SAM-dependent methyltransferases as direct-antiviral targets of HC and CAPE

The availability of recently determined cryo-EM atomic structure of CHIKV nsP1 capping

enzyme containing virus-specific MTase enzyme activity and the crystal structure of DENV

NS5 MTase (PDB ID 4V0R) facilitated *in silico* structure-based analysis of HC and CAPE as

potential MTase inhibitors [23], [49]. SAM-binding site is formed by residues Gly65, Arg70,

Arg71, Pro83, Glu88, Asp89, Arg92, Gln151, Asp152 and Val153 in CHIKV nsP1 [23].

Structural analysis of DENV NS5 MTase has revealed that the SAM binding site residues for

flaviviruses are Asp79, Leu80, Gly81, Cys82, Gly83, Arg84, Gly85, Trp87, His11 0, Asp131,

and Phe133 [48], [49]. Primary sequence alignment and comparison of various members of the

alphaviruses and flaviviruses revealed high sequence similarity, as shown in supplementary

figure 1A. In alphaviruses, 94.4% of the active site residues showed 100% sequence identity

across nsP1 of different viruses. Whereas in flaviviruses, 90.9% of the active site residues

showed 100% sequence identity across NS5 MTase when compared (Supplementary figure

1B). The MTase active site residues sequentially, structurally and functionally conserved in

nsP1 and NS5 are potential drug target sites and were *in silico* examined for binding to the HC

and CAPE molecules.

For molecular docking of HC and CAPE to evaluate binding of these molecules to the viral

MTases, the alphavirus CHIKV nsP1 structure and flavivirus DENV NS5 MTase domain

structures were generated using SWISS-MODEL as PDB entry (6Z0U and 4V0R) lacked few

atomic coordinates for multiple flexible loops. The generated models were superimposed on

to the template of each of the viral protein using PyMOL 2.3 and the RMSD was calculated.

The 2 dimensions image with the superimposed amino acid residues are highlighted and shown

in Supplementary figure 1. Structural comparison of the generated models of CHIKV and

DENV with respective templates showed high structural similarity with low RMSD values i.e.,

0.205 Å and 0.182 Å respectively (Supplementary figure 2 A and B). CHIKV nsP1 and DENV

NS5 MTase generated models were further used for flexible docking of HC, CAPE, and the

positive control i.e. SAM. The positive control showed binding energy in the range of -7 to -

7.7 kcal/mol. Interestingly, the binding affinities of both the compounds obtained were higher

than the positive control for both the viruses. It was observed that HC and CAPE form hydrogen

(H) bonds with the active site residues of CHIKV nsP1 and make hydrophobic interactions as

well (Table 1). While, in DENV NS5 MTase, it was observed that along with the active site

residues the H-bonds were also formed with the residues that are in the vicinity of the active

site (Table 2). The attained binding energies of HC and CAPE along with the positive controls

and detailed protein-ligand interactions are presented in Table 1-2 and Figures 4-5.

The dynamic stability and conformational changes in the Apo proteins and protein-ligands

complexes were studied by analyzing their RMSD during the MD simulation. RMSD data

showed that all the nsP1-ligand complexes attained equilibrium at 15 ns and reached stable

state throughout the simulation for up to 50 ns (Supplementary figure 3A). Equilibrium was

attained after 15 ns and fluctuations were found to be in the range of 0.6 to 0.75 nm for nsP1

Apo, 0.35 to 0.55 ns for nsP1-SAM, nsP1-HC, and nsP1-CAPE. Similarly, MD simulation

based RMSD study performed for DENV NS5 MTase showed the Apo NS5 MTase and the

NS5 MTase-SAM complex attained equilibrium at 20 ns and 10 ns, respectively, whereas the

HC and CAPE complexes attained equilibrium before 5 ns itself (Supplementary figure 3B).

The systems remained stable up to 50 ns and fluctuations were obtained in the range of 0.2 to

0.4Å for NS5 MTase Apo and the NS5 MTase complexes. These observations depict that the

binding of molecules to nsP1 and NS5 proteins resulted in the formation of a stable complex.

Virus-specific MTase inhibition by HC and CAPE

To further validate the inhibition of viral methyltransferase activity of HC and CAPE, CHIKV

nsP1 was purified and the MTase enzymatic activity of purified CHIKV nsP1 was performed

using an indirect ELISA as per the previously published protocol [22], [55], [56]. Briefly, at

450 nm, the 3,3′,5,5′-Tetramethylbenzidine TMB/H_2O_2-ELISA substrate was used to identify

immunoreactive proteins comprising the m7GMP-nsP1 complex. For IC_{50} calculations, the

experiment was carried out by incubating CHIKV nsP1 protein (20 μM) in the reaction mixture

with various concentrations of these two compounds. As the concentration of the compounds

was increased, the absorbance or relative nsP1 activity was observed to be decreasing

confirming the inhibitory activity of both the compounds HC and CAPE against a viral MTase

(Figure 6 A-C). The dose-response curve showed the IC_{50} values for HC and CAPE are 86 nM

and 9 nM, respectively indicating these two compounds to be effective inhibitors of CHIKV

nsP1. Similarly, to validate the inhibition of NS5 methyltransferase activity of HC and CAPE,

NS5 Mtase was purified and the CE based MTase enzymatic activity of purified DENV NS5

Mtase was performed as per the previously published protocol [59], [60]. DENV NS5 Mtase

activity was descreased in dose dependent manner confirming to the inhibitory activity of both

the compounds HC and CAPE against a DENV NS5 Mtase (Figure 6 D-F).

Discussion

Communicable diseases transmitted through vectors such as mosquitoes remain a significant

public health burden worldwide. Although human viruses belong to several viral families,

alphaviridae and *flaviviridae* are the most significant burden on public health [4], [7]. Drug

development for mosquito-transmitted viruses such as DENV and CHIKV remains incomplete.

Neither commercially available drugs nor vaccines are available for these viruses. These viral

pathogen's emergence and re-emergence pose a substantial health risk to the global human

population. The current lack of vaccines and effective antivirals emphasizes the importance of

developing new strategies to combat these life-threatening human pathogens.

Polyamines are ubiquitous molecules found in all cells and they play many important roles within the cell, including binding and altering the conformation of RNA, modifying transcription and translation rates, and affecting signaling pathways [14]. Similarly, they also have an essential role in the viral life cycle. Previous studies showed that DFMO treatment cells depleted biogenic polyamine levels, leading to its antiviral activity against various viruses such as CHIKV, ZIKV, and DENV [12].

In this study, we have demonstrated that both HC and CAPE have significant anti-CHIKV and anti-DENV activity. Both compounds shown better reduction for CHIKV and DENV infection when compared to the established positive control, i.e., DFMO in the plaque reduction assay. Surprisingly, the effective concentrations of both HC and CAPE were significantly lower than their toxic dose, i.e., EC_{50} values for both compounds are in the nanomolar and micromolar range for CHIKV and DENV, respectively.

To quantify the copy number of viral RNA released from CHIKV and DENV-infected cells, qRT-PCR amplified the viral genome. qRT-PCR measures mRNA, which represents viral load during replication. Our finding shows that treatment with HC and CAPE significantly reduces viral load compared to the positive control. The IFA experiments further validated the results.

Our findings supported a previously published concept that HC and CAPE are ODC inhibitors [58], [61]. HC is a naturally occurring flavonoid that is an established ODC inhibitor [58]. CAPE is a natural ester that has also shown antiviral activity, and it significantly prevented the full induction of ODC by epidermal growth factor (EGF) [32], [61]. In our study, HC and CAPE molecules have shown more polyamine depletion than the DFMO. Furthermore, the exogenous polyamine addition assay did not rescue viral titre for CHIKV and DENV, implying that these compounds are also acting in ways other than the polyamine depletion pathways.

Alphaviruses use an unconventional mechanism of RNA capping in which nsP1 plays a vital role in the capping of viral RNA, and the binding site of SAM remains conserved, as shown

540 by MSA. Similarly, flaviviruses have a 5' terminal cap 1 structure (m7GpppAmG), and SAM

541 binding site remains conserved. Additionally, the results of bioinformatic prediction of both

542 compounds show the binding energy in the range of -7.6 to -8 kcal/mol with the nsP1protein

543 of CHIKV and -8 kcal/mol with NS5 of DENV, which are corroborated with the results of

544 CHIKV nsP1 biochemical assay and capillary electrophoresis based NS5 MTase assay.

552 **<u>Conclusion</u>**

553 Our research provides the compelling first evidence on the anti-CHIKV and anti-DENV

554 activities of HC and CAPE through two mechanisms a) indirectly by depleting polyamines in

555 mammalian cells and b) directly by targeting viral MTases. HC and CAPE's novel broad-

556 spectrum antiviral activity makes them promising candidates for developing anti-DENV and

557 anti-CHIKV therapeutics.

559 **Acknowledgments**

560 The authors are thankful to the Department of biosciences and bioengineering, IIT Roorkee for

561 central facilities and the translational and structural bioinformatics center. The authors thank

562 the Macromolecular Crystallographic Facility (MCU) at IIC, Indian Institute of Technology,

Roorkee. The authors would like to acknowledge Dr. Rajat Mudgal for technical assistance in

CE analysis. The authors would like to thank the University Grants Commission (UGC) and

Council of Scientific and Industrial Research (CSIR), and Ministry of Human Resource

Development (MHRD) for providing financial support. This work was supported by a research

grant to ST from ICMR Ref no. ISRM/12(46)/2020.

Ethical approval

DENV-positive clinical sample was collected from HiTech Pathology Laboratory, Roorkee

with the approval of the Institutional Human Ethical Committee (IHEC), Indian Institute of

Technology, Roorkee (Ref no. BT/IHEC-IITR/2016/6499).

Authors contribution

MB, RR, AK, SM, BW, RK, VS conducted the experiments and analyzed the data. MB, RR,

AK, SM, DS, PK and ST planned the experiments, analyzed the data, and wrote the manuscript.

Competing interests

The authors declare that they have no competing interests.

References

[1]   M. G. Guzman, D. J. Gubler, A. Izquierdo, E. Martinez, and S. B. Halstead, 'Dengue
infection', *Nat Rev Dis Primers*, vol. 2, Aug. 2016, doi: 10.1038/NRDP.2016.55.

[2]   A. M. Powers *et al.*, 'Evolutionary relationships and systematics of the alphaviruses', *J
Virol*, vol. 75, no. 21, pp. 10118–10131, Nov. 2001, doi: 10.1128/JVI.75.21.10118-
10131.2001.

[3] 'DENGUE/DHF SITUATION IN INDIA :: National Center for Vector Borne Diseases
Control (NCVBDC)'.
https://nvbdcp.gov.in/index4.php?lang=1&level=0&linkid=431&lid=3715 (accessed
Jan. 15, 2022).

[4] S. Bhatt *et al.*, 'The global distribution and burden of dengue', vol. 496, no. 7446, pp.
504–507, Apr. 2013, Accessed: Jan. 15, 2022. [Online]. Available:
https://pubmed.ncbi.nlm.nih.gov/23563266/

[5] S. Bhatt *et al.*, 'The global distribution and burden of dengue', *Nature*, vol. 496, no.
7446, pp. 504–507, Apr. 2013, doi: 10.1038/NATURE12060.

[6] 'CHIKUNGUNYA SITUATION IN INDIA :: National Center for Vector Borne
Diseases Control (NCVBDC)'.
https://nvbdcp.gov.in/index4.php?lang=1&level=0&linkid=486&lid=3765 (accessed
Jan. 15, 2022).

[7] 'Geographical distribution of chikungunya virus disease cases reported worldwide,
2021'. https://www.ecdc.europa.eu/en/publications-data/geographical-distribution-
chikungunya-virus-disease-cases-reported-worldwide-2021 (accessed Jan. 15, 2022).

[8] M. Kaur *et al.*, 'Coinfection of chikungunya and dengue viruses: A serological study
from North Western region of Punjab, India', *J Lab Physicians*, vol. 10, no. 4, p. 443,
Oct. 2018, doi: 10.4103/JLP.JLP_13_18.

[9] H. S. Chahar, P. Bharaj, L. Dar, R. Guleria, S. K. Kabra, and S. Broor, 'Co-infections
with Chikungunya Virus and Dengue Virus in Delhi, India', *Emerg Infect Dis*, vol. 15,
no. 7, p. 1077, Jul. 2009, doi: 10.3201/EID1507.080638.

[10] S. Mahajan, S. Choudhary, P. Kumar, and S. Tomar, 'Antiviral strategies targeting
host factors and mechanisms obliging +ssRNA viral pathogens', *Bioorg Med Chem*,
vol. 46, Sep. 2021, doi: 10.1016/J.BMC.2021.116356.

[11] M. Bhutkar, V. Singh, P. Dhaka, and S. Tomar, 'Virus-host protein-protein
interactions as molecular drug targets for arboviral infections', *Frontiers in Virology*,
vol. 0, p. 70, Aug. 2022, doi: 10.3389/FVIRO.2022.959586.

[12] B. C. Mounce *et al.*, 'Interferon-Induced Spermidine-Spermine Acetyltransferase and
Polyamine Depletion Restrict Zika and Chikungunya Viruses', *Cell Host Microbe*, vol.
20, no. 2, pp. 167–177, Aug. 2016, doi: 10.1016/J.CHOM.2016.06.011.

[13] B. C. Mounce *et al.*, 'Inhibition of Polyamine Biosynthesis Is a Broad-Spectrum
Strategy against RNA Viruses', *J Virol*, vol. 90, no. 21, pp. 9683–9692, Nov. 2016,
doi: 10.1128/JVI.01347-16.

[14] L. Miller-Fleming, V. Olin-Sandoval, K. Campbell, and M. Ralser, 'Remaining
Mysteries of Molecular Biology: The Role of Polyamines in the Cell', *J Mol Biol*, vol.
427, no. 21, pp. 3389–3406, Oct. 2015, doi: 10.1016/J.JMB.2015.06.020.

[15] K. Igarashi and K. Kashiwagi, 'Polyamines: Mysterious Modulators of Cellular
Functions', *Biochem Biophys Res Commun*, vol. 271, no. 3, pp. 559–564, May 2000,
doi: 10.1006/BBRC.2000.2601.

[16] K. J. Doyle, 'Delayed, reversible hearing loss caused by difluoromethylornithine
(DFMO)', *Laryngoscope*, vol. 111, no. 5, pp. 781–785, 2001, doi: 10.1097/00005537-
200105000-00005.

[17] L. Alhonen, V. A. Levin, and L. J. Marton, 'Article in Cancer Research', 1985.
[Online]. Available: http://cancerres.aacrjournals.org/content/45/2/509

[18] J. Jose, J. E. Snyder, and R. Kuhn, 'A structural and functional perspective of
alphavirus replication and assembly', *Future Microbiol*, vol. 4, no. 7, pp. 837–856,
2009, doi: 10.2217/FMB.09.59.

[19] W. Filipowicz, Y. Furuichi, J. M. Sierra, S. Muthukrishnan, A. J. Shatkin, and S.
Ochoa, 'A protein binding the methylated 5'-terminal sequence, m7GpppN, of
eukaryotic messenger RNA', *Proc Natl Acad Sci U S A*, vol. 73, no. 5, pp. 1559–1563,
1976, doi: 10.1073/PNAS.73.5.1559.

[20] J. L. Hyde and M. S. Diamond, 'Innate immune restriction and antagonism of viral
RNA lacking 2′-O methylation', *Virology*, vol. 479–480, pp. 66–74, May 2015, doi:
10.1016/J.VIROL.2015.01.019.

[21] T. Ahola and L. Kääriäinen, 'Reaction in alphavirus mRNA capping: formation of a
covalent complex of nonstructural protein nsP1 with 7-methyl-GMP.', *Proc Natl Acad
Sci U S A*, vol. 92, no. 2, p. 507, Jan. 1995, doi: 10.1073/PNAS.92.2.507.

[22] R. Mudgal, S. Mahajan, and S. Tomar, 'Inhibition of Chikungunya virus by an
adenosine analog targeting the SAM-dependent nsP1 methyltransferase', *FEBS Lett*,
vol. 594, no. 4, pp. 678–694, Feb. 2020, doi: 10.1002/1873-3468.13642.

[23] R. Jones, G. Bragagnolo, R. Arranz, and J. Reguera, 'Capping pores of alphavirus
nsP1 gate membranous viral replication factories', *Nature*, vol. 589, no. 7843, p. 615,
Jan. 2021, doi: 10.1038/S41586-020-3036-8.

[24] 'Flaviviridae', *Virus Taxonomy*, pp. 1003–1020, Jan. 2012, doi: 10.1016/B978-0-12-
384684-6.00086-0.

[25] M. P. Egloff, D. Benarroch, B. Selisko, J. L. Romette, and B. Canard, 'An RNA cap
(nucleoside-2′-O-)-methyltransferase in the flavivirus RNA polymerase NS5: crystal
structure and functional characterization', *EMBO J*, vol. 21, no. 11, p. 2757, Jun. 2002,
doi: 10.1093/EMBOJ/21.11.2757.

[26] S. T. Raot and M. G. Rossmann, 'Comparison of Super-secondary Structures in
Proteins', 1973.

[27] B. P. S. Chouhan, S. Maimaiti, M. Gade, and P. Laurino, 'Rossmann-Fold
Methyltransferases: Taking a "β-Turn" around Their Cofactor, S-
Adenosylmethionine', *Biochemistry*, vol. 58, no. 3, pp. 166–170, Jan. 2019, doi:
10.1021/ACS.BIOCHEM.8B00994/ASSET/IMAGES/LARGE/BI-2018-
00994Q_0003.JPEG.

[28] B. Wang *et al.*, 'The structure of Zika virus NS5 reveals a conserved domain
conformation', *Nature Communications 2017 8:1*, vol. 8, no. 1, pp. 1–6, Mar. 2017,
doi: 10.1038/ncomms14763.

[29] T. Ahola and A. Merits, 'Functions of Chikungunya Virus Nonstructural Proteins',
Chikungunya Virus, p. 75, Jan. 2016, doi: 10.1007/978-3-319-42958-8_6.

[30] D. J. Kim *et al.*, 'Herbacetin is a novel allosteric inhibitor of ornithine decarboxylase
with antitumor activity', *Cancer Res*, vol. 76, no. 5, p. 1146, Mar. 2016, doi:
10.1158/0008-5472.CAN-15-0442.

[31] L. R *et al.*, 'Antiviral activity of selected flavonoids against Chikungunya virus',
Antiviral Res, vol. 133, pp. 50–61, Sep. 2016, doi:
10.1016/J.ANTIVIRAL.2016.07.009.

[32] H. Erdemli, S. Akyol, F. Armutcu, and O. Akyol, 'Antiviral properties of caffeic acid
phenethyl ester and its potential application', *J Intercult Ethnopharmacol*, vol. 4, no. 4,
p. 344, 2015, doi: 10.5455/JICE.20151012013034.

[33] B. T. Zhu, P. Wang, M. Nagai, Y. Wen, and H. W. Bai, 'Inhibition of human catechol-
O-methyltransferase (COMT)-mediated O-methylation of catechol estrogens by major
polyphenolic components present in coffee', *J Steroid Biochem Mol Biol*, vol. 113, no.
1–2, pp. 65–74, Jan. 2009, doi: 10.1016/J.JSBMB.2008.11.011.

[34] D. Xiang *et al.*, 'Caffeic acid phenethyl ester induces growth arrest and apoptosis of
colon cancer cells via the β-catenin/T-cell factor signaling', *Anticancer Drugs*, vol. 17,
no. 7, pp. 753–762, Aug. 2006, doi: 10.1097/01.cad.0000224441.01082.bb.

[35] L. J. Chien, T. L. Liao, P. Y. Shu, J. H. Huang, D. J. Gubler, and G. J. J. Chang,
'Development of real-time reverse transcriptase PCR assays to detect and serotype
dengue viruses', *J Clin Microbiol*, vol. 44, no. 4, pp. 1295–1304, Apr. 2006, doi:
10.1128/JCM.44.4.1295-1304.2006.

[36] H. Singh *et al.*, 'Chikungunya virus inhibition by peptidomimetic inhibitors targeting
virus-specific cysteine protease', *Biochimie*, vol. 149, pp. 51–61, Jun. 2018, doi:
10.1016/J.BIOCHI.2018.04.004.

[37] A. C. Mishra *et al.*, 'Stratified sero-prevalence revealed overall high disease burden of
dengue but suboptimal immunity in younger age groups in Pune, India', *PLoS Negl
Trop Dis*, vol. 12, no. 8, Aug. 2018, doi: 10.1371/JOURNAL.PNTD.0006657.

[38] R. Kaur, Neetu, R. Mudgal, J. Jose, P. Kumar, and S. Tomar, 'Glycan-dependent
chikungunya viral infection divulged by antiviral activity of NAG specific chi-like
lectin', *Virology*, vol. 526, pp. 91–98, Jan. 2019, doi: 10.1016/J.VIROL.2018.10.009.

[39] R. Madhubala, 'Thin-layer chromatographic method for assaying polyamines',
Methods Mol Biol, vol. 79, pp. 131–136, 1998, doi: 10.1385/0-89603-448-8:131.

[40] F. Sievers *et al.*, 'Fast, scalable generation of high-quality protein multiple sequence
alignments using Clustal Omega', *Mol Syst Biol*, vol. 7, 2011, doi:
10.1038/MSB.2011.75.

[41] 'Sci-Hub | Structural insights into viral RNA capping and plasma membrane targeting
by Chikungunya virus nonstructural protein 1. Cell Host & Microbe, 29(5), 757–
764.e3 | 10.1016/j.chom.2021.02.018'. https://sci-
hub.se/https://doi.org/10.1016/j.chom.2021.02.018 (accessed Jan. 20, 2022).

[42] R. Jones, G. Bragagnolo, R. Arranz, and J. Reguera, 'Capping pores of alphavirus
nsP1 gate membranous viral replication factories', vol. 589, no. 7843, Accessed: Jan.
20, 2022. [Online]. Available: https://www.nature.com/articles/s41586-020-3036-8

[43] R. Jones, G. Bragagnolo, R. Arranz, and J. Reguera, 'Capping pores of alphavirus nsP1 gate membranous viral replication factories', *Nature 2020 589:7843*, vol. 589, no. 7843, pp. 615–619, Dec. 2020, doi: 10.1038/s41586-020-3036-8.

[44] A. Waterhouse *et al.*, 'SWISS-MODEL: homology modelling of protein structures and complexes', *Nucleic Acids Res*, vol. 46, no. W1, pp. W296–W303, Jul. 2018, doi: 10.1093/NAR/GKY427.

[45] O. Trott and A. J. Olson, 'AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization and multithreading', *J Comput Chem*, vol. 31, no. 2, p. 455, 2010, doi: 10.1002/JCC.21334.

[46] S. Dallakyan and A. J. Olson, 'Small-molecule library screening by docking with PyRx', *Methods Mol Biol*, vol. 1263, pp. 243–250, 2015, doi: 10.1007/978-1-4939-2269-7_19.

[47] N. M. O'Boyle, M. Banck, C. A. James, C. Morley, T. Vandermeersch, and G. R. Hutchison, 'Open Babel: An Open chemical toolbox', *J Cheminform*, vol. 3, no. 10, pp. 1–14, Oct. 2011, doi: 10.1186/1758-2946-3-33/TABLES/2.

[48] Y. Zhao *et al.*, 'A Crystal Structure of the Dengue Virus NS5 Protein Reveals a Novel Inter-domain Interface Essential for Protein Flexibility and Virus Replication', *PLoS Pathog*, vol. 11, no. 3, pp. 1–27, Mar. 2015, doi: 10.1371/JOURNAL.PPAT.1004682.

[49] B. Zhao *et al.*, 'Structure and function of the Zika virus full-length NS5 protein', *Nature Communications 2017 8:1*, vol. 8, no. 1, pp. 1–9, Mar. 2017, doi: 10.1038/ncomms14762.

[50] 'PyMOL | pymol.org'. https://pymol.org/2/ (accessed Mar. 05, 2022).

[51] R. A. Laskowski and M. B. Swindells, 'LigPlot+: multiple ligand-protein interaction diagrams for drug discovery', *J Chem Inf Model*, vol. 51, no. 10, pp. 2778–2786, Oct. 2011, doi: 10.1021/CI200227U.

[52] D. Van Der Spoel, E. Lindahl, B. Hess, G. Groenhof, A. E. Mark, and H. J. C. Berendsen, 'GROMACS: fast, flexible, and free', *J Comput Chem*, vol. 26, no. 16, pp. 1701–1718, Dec. 2005, doi: 10.1002/JCC.20291.

[53] A. W. Schüttelkopf and D. M. F. Van Aalten, 'PRODRG: a tool for high-throughput crystallography of protein-ligand complexes', *Acta Crystallogr D Biol Crystallogr*, vol. 60, no. Pt 8, pp. 1355–1363, Aug. 2004, doi: 10.1107/S0907444904011679.

[54] R. Mudgal *et al.*, 'Selective Estrogen Receptor Modulators Limit Alphavirus Infection by Targeting the Viral Capping Enzyme nsP1', *Antimicrob Agents Chemother*, vol. 66, no. 3, Mar. 2022, doi: 10.1128/AAC.01943-21/ASSET/CFFD4322-C11C-41A1-9D51-7D0C3242FA63/ASSETS/IMAGES/LARGE/AAC.01943-21-F006.JPG.

[55] S. Tomar, M. Narwal, E. Harms, J. L. Smith, and R. J. Kuhn, 'Heterologous production, purification and characterization of enzymatically active Sindbis virus nonstructural protein nsP1', *Protein Expr Purif*, vol. 79, no. 2, pp. 277–284, Oct. 2011, doi: 10.1016/J.PEP.2011.05.022.

[56] R. Kaur, R. Mudgal, M. Narwal, and S. Tomar, 'Development of an ELISA assay for screening inhibitors against divalent metal ion dependent alphavirus capping enzyme', *Virus Res*, vol. 256, pp. 209–218, Sep. 2018, doi: 10.1016/J.VIRUSRES.2018.06.013.

[57] R. Lani *et al.*, 'Antiviral activity of selected flavonoids against Chikungunya virus',
Antiviral Res, vol. 133, pp. 50–61, Sep. 2016, doi:
10.1016/J.ANTIVIRAL.2016.07.009.

[58] D. J. Kim *et al.*, *Herbacetin is a novel allosteric inhibitor of ornithine decarboxylase*
with antitumor activity, vol. 76, no. 5. NIH Public Access, 2016, p. 1146. doi:
10.1158/0008-5472.CAN-15-0442.

[59] R. Mudgal, S. Mahajan, and S. Tomar, 'Inhibition of Chikungunya virus by an
adenosine analog targeting the SAM-dependent nsP1 methyltransferase', *FEBS Lett*,
vol. 594, no. 4, pp. 678–694, Feb. 2020, doi: 10.1002/1873-3468.13642.

[60] S. Boonyasuppayakorn and R. Padmanabhan, 'Construction of plasmid, bacterial
expression, purification, and assay of dengue virus type 2 NS5 methyltransferase',
Methods in Molecular Biology, vol. 1138, pp. 361–373, 2014, doi: 10.1007/978-1-
4939-0348-1_22/COVER.

[61] Z. S. Zheng, G. Z. Xue, D. Grunberger, and J. H. Prystowsky, 'Caffeic acid phenethyl
ester inhibits proliferation of human keratinocytes and interferes with the EGF
regulation of ornithine decarboxylase.', *Oncol Res*, vol. 7, no. 9, pp. 445–452, Jan.
1995, Accessed: May 19, 2022. [Online]. Available:
https://europepmc.org/article/med/8835288

Figure 1: Evaluation of antiviral activity using virus titer reduction profiling of HC and CAPE.

Panel A and C depicts the percent cell viability of Vero cells treated with HC and CAPE for

12 h pre-treatment and 24 h post-treatment and incubated in maintenance media for 4 days,

Panel B showing the percent cell viability of Vero cells treated with CAPE for 12 h pre-

treatment and 24 h post-treatment. Cell viability for all tested concentrations remained above

the cytotoxic threshold of 90%. Panels D-G depicts the inhibitory effect of CHIKV and DENV

infected cells on different concentrations of HC and CAPE. Vero cells were infected with

CHIKV (D-F) and DENV (G-I), with DFMO (1000 µM) serving as a positive control and VC

representing virus control. Panel J depicts the effect of polyamines level after treating with HC

(200 µM) and CAPE (25 µM). Polyamine level of Vero treated cells at 12, 24, and 36 h for HC

and pre (12 h) and post-treatment (24 h) for CAPE and DEMO, using untreated cells as a cell

control (CC) for both of the conditions. 0.1 µM putrescine (put), spermine (spm), and

spermidine (spd) as a positive control marker. Treatment with both compounds leads to

decreased polyamine levels for both of the compounds compared to DFMO. Error bars

represent the standard error of experimental replicates with n = 3. Statistical analysis was

*performed using one-way ANOVA with Dunnett's post-test. ***P < 0.001.*

Figure 2: Evaluation of the antiviral effect by immunofluorescence assay (IFA).

Immunofluorescent staining of HC (200 µM) and CAPE (25 µM for CHIKV and 2.5 µM for

DENV) treated with CHIKV(A) and DENV(B) in Vero cells. Green fluorescence indicates the

virus load, and blue fluorescence indicates the nuclear staining with DAPI with a 10× objective

lens. The scale bar is 400 µM.

Figure 3: Effect of addition of biogenic polyamines on virus infected and drug-treated cells.

Vero cells infected with CHIKV (A) and DENV (B) treated with HC (200 µM), CAPE (25 µM

for CHIKV and 2.5 µM for DENV), DFMO (1000 µM) as a positive control and VC as

untreated control. – (minus) sign indicates only the drug treated Vero cells whereas + (plus)

sign indicates that 1µM polyamines (put, spd, spm) was added additionally in the drug treated

cells. Error bars represent the standard error of experimental replicates with n = 3.

Figure 4: *Ligands docked in the binding site pocket (teal surface) of CHIKV nsP1 and DENV NS5 (wheat surface). A-D) CHIKV nsP1 interact with the SAM (A), GTP (B), HC (C) and CAPE (D); E-G) DENV NS5 MTase interact with SAM (E), HC (F), and CAPE (G). Zoomed in figures show a detailed view of the binding pocket where molecular interactions of ligands (green) with binding active site residues (teal colour) and additional binding residues (pink colour) of CHIKV nsP1 and DENV NS5 MTase (wheat coloured protein ribbon/ surface).*

Figure 5*: Two-dimensional representation view of docked molecules in LigPlot+ software to*

visualise the interacting residues with the CHIKV nsP1 and DENV NS5 MTase. A-D) CHIKV

nsP1 interact with the SAM (A), HC (B), CAPE (C) ; (D-F) DENV NS5 MTase interact with

SAM (D), HC (E), and CAPE (F). H-bonds are shown in green dashed lines with the distance

shown in Å. Additional residues forming hydrophobic interactions are shown by brown

semicircle with radiating spokes towards the ligands.

***Figure 6:** Methyltransferase inhibition activity assay. A) SDS/PAGE analysis of purified*

recombinant CHIKV nsP1. The CHIKV nsP1 inhibition assay was executed in the presence of

increasing concentration of HC (B) and CAPE (C) using an indirect ELISA assay.D)

SDS/PAGE analysis of purified recombinant DENV NS5 MTase. The CE based DENV NS5

inhibition assay was executed in the presence of increasing concentration of HC (E) and CAPE

(F).

Supplementary figure 1: *Multiple Sequence alignment of MTase domains of different alphaviruses and flaviviruses. A) Sequence alignment of nsP1 MTase domain (residues 1 to 298) from CHIKV with the MTase domains of other alphaviruses. B) Sequence alignment of NS5 MTase domain (residues 1 to 276) from DENV with MTase domains of other flaviviruses. The active-site MTase residues in both alphaviruses and flaviviruses are highly conserved suggesting that structure and function of MTase domain in both families are conserved. Residues are represented with single letter amino acid code, with identical residues indicated in white font and boxed in red, while similar residues are indicated in red font. Blue triangles are corresponding to the MTase active site residues as per literature. Alignments were generated via Clustal Omega and ESPript tools.*

Supplementary figure 2: *Superposition of SWISS generated models. (A-B) Close-up view of*

overlapped residues of generated models of CHIKV nsP1 (A) and DENV NS5 MTase (B) with

RMSD values of 0.205 Å and 0.182 Å, respectively. The generated model is labelled yellow,

and the template is labelled pink.

Supplementary figure 3: *Root Mean Square Deviations (RMSD) graphs of (A) CHIKV nsP1*

Apo i.e., native protein, nsP1 - SAM or CAPE or HC (B) DENV NS5 MTase Apo i.e., native,

NS5 MTase - SAM or HC or CAPE

Table 1: Detailed molecular interactions between the CHIKV nsP1 and ligands.

Ligand	Binding energy (kcal/mol)	Interactions			
		Hydrogen			Hydrophobic interactions
		Residues	Molecules	Bond length (Å)	
SAM	-7.0	Asp152	N1-O	3.09	Arg 85, Ser86, Arg92, Tyr154, Tyr248, Pro249, Glu250
			N5-O	2.97	
		Glu88	N3- OE1	3.03	
		Asp89	O3-OD2	3.20	
		Gly65	O2-O	3.03	
HC	-7.9	Glu88	O3-OE1	2.83	Ala 40, Arg41, Asp152, Phe178, Tyr285, Val243, Tyr248
		Arg92	O4-NH2	2.93	
		Arg70	O2-NH1	3.04	
CAPE	-7.6	Arg92	O1-NH2	3.08	Ala40, Arg70, Asp152, Tyr154, Phe178, Val243, Tyr248, Tyr285
		Glu250	O4 -OE1	2.91	
			O3-OE1	2.88	

Table 2: Detailed molecular interactions between the DENV NS5 MTase and ligands.

Ligand	Binding energy (kcal/mol)	Interactions			Hydrophobic interactions
		Hydrogen			
		Residues	Molecules	Bond length (Å)	
SAM	-7.7	Asp 128	OD1-N3	3.13	Gly80, Lys102, His107, Phe130, Ile144, Lys177
		Ser 53	OG-O4	3.27	
		Gly 78	O-O5, O-N6	2.78,3.14	
		Thr101	OG1-O2	3.00	
		Cys 79	O-N6	3.15	
		Arg 81	N-O3	3.18	
HC	-8.0	Thr101	OG1-O2	2.67	Gly78, Gly80, His107, Asp128, Phe130, Asp143, Ile144
			OG1-O4	2.81	
		Lys102	N-O4	3.30	
		Gly145	N-O5	2.65	
CAPE	-8.0	Trp84	N-O4	3.12	Gly78, Cys79, Gly80, Arg81, Gly83, Thr101, Lys102, Glu108, Asp128, Val129, Phe130, Asp143, Ile144
		Gly82	N-O3	3.08	
		Ser53	OG-O3	3.05	