SAM-dependent viral MTase inhibitors: herbacetin and caffeic acid phenethyl ester,

structural insights into dengue MTase

Mandar Bhutkar, Amith Kumar, Ruchi Rani, Vishakha Singh, Akashjyoti Pathak, Aditi

Kothiala, Supreeti Mahajan, Bhairavnath Waghmode, Ravi Kumar, Rajat Mudgal, Debabrata

Sircar, Pravindra Kumar, Shailly Tomar *

Department of Biosciences and Bioengineering, Indian Institute of Technology Roorkee,

Uttarakhand, India.

*Corresponding author address: Department of Biosciences and Bioengineering, Indian

Institute of Technology Roorkee, Roorkee-247667, Uttarakhand, India; Email id:

shailly.tomar@bt.iitr.ac.in

Abstract

Chikungunya (CHIKV) and dengue (DENV) viruses pose a public health risk and lack antiviral treatment. Structure-based virtual screening of natural MTase substrates library identified herbacetin (HC) and caffeic acid phenethyl ester (CAPE) as potential CHIKV nsP1 and DENV NS5 MTase inhibitors. Binding affinities and MTase inhibition were confirmed using purified proteins. Crystal structure of DENV3 NS5 MTase and CAPE complex revealed CAPE binding at GTP and cap 0 RNA sites. Interestingly, HC and CAPE depleted polyamines, which are crucial for RNA virus replication, and effectively diminished replication with IC_{50} values of ~13.44 µM and ~0.57 µM against CHIKV, and ~7.24 µM and ~1.01 µM against DENV, respectively. Polyamine addition did not reverse the antiviral effects, suggesting a dual inhibition mechanism.

Graphical abstract

Keywords

57   Dengue virus, Chikungunya virus, Polyamine depletion, Antiviral assay, Crystal Structure

Introduction

Dengue virus (DENV) and chikungunya virus (CHIKV) are arthropod-borne viruses belonging to the *Flaviviridae* and *Togaviridae* families, respectively (1). Both viruses cause febrile illnesses, showing fever, arthralgia, joint pain, rash, and headache. These enveloped viruses have a positive-sense single-stranded RNA genome (+ssRNA) (2). It is estimated that annually, ~100-400 million DENV infections occur worldwide. The Philippines and Vietnam are among the most affected countries. Moreover, dengue is endemic in India, Indonesia, Myanmar, Sri Lanka, and Thailand (3–5). Further, in 2023, ~ 0.5 million CHIKV cases occurred worldwide, of which most were reported in Brazil and India (6,7). In India, co-infections with DENV and CHIKV occur prevalently in various regions during the monsoon season (6,8,9).

The *Togaviridae* family includes various alphaviruses such as CHIKV, Eastern equine encephalitis virus (EEEV), Venezuelan equine encephalitis virus (VEEV), etc. Alphaviruses comprise four nonstructural proteins (nsPs), namely nsP1, nsP2, nsP3, and nsP4, which play an essential role in the life cycle of the viruses and are produced when a viral protease, i.e., nsP2, cleaves the viral polyprotein precursor post-translationally (10). Among the four nonstructural proteins (nsPs) in alphaviruses, nsP1 is crucial due to its essential role in viral RNA capping and is a potential drug target (11–14). The viral RNA caps have numerous biological roles, such as helping the eukaryotic translational initiation factor 4E (eIF4e) recognize the RNA for initiating translation and protecting the mRNA from cellular exonucleases (15). Further, the cap structure also helps the viral genome to escape recognition from the host innate immune system pathways such as retinoic acid-inducible gene I (RIG-I) and melanoma differentiation-associated protein 5 (MDA5) (16). In alphaviruses, the cap 0 structure is characterized by monomethylation at the N7 position of the guanosine nucleotide(17). The capping enzyme nsP1 of alphaviruses facilitates viral RNA capping through two crucial steps. Initially, S-adenosyl methionine (SAM) serves as a methyl donor, transferring a methyl group to the seventh position of GTP, resulting in the formation of m7

guanosine-5'-monophosphate (m7GMP) and producing S-adenosyl homocysteine (SAH) as a

by-product. This process is referred to as the MTase step of the capping reaction (18). In the

subsequent guanylation (GT) step, the methylated m7GMP is covalently attached to His37 of

the nsP1 enzyme and is transferred to the 5' end of the viral RNA (18).

The *Flaviviridae* family includes orthoflaviviruses like DENV, Zika (ZIKV), Japanese

encephalitis, etc. DENV consists of four distinct serotypes: DENV 1, DENV 2, DENV 3, and

DENV 4. DENV-2 has been the predominant serotype in India for the past five decades.

Nevertheless, recent epidemics have also witnessed the emergence of serotypes 3 and 4,

indicating a dynamic shift in the circulating dengue virus serotypes in the region (19).

Orthoflaviviruses contain a single open reading frame, which encodes a polyprotein processed

by viral and cellular proteases into three structural and seven nonstructural proteins (NS1,

NS2A, NS2B, NS3, NS4A, NS4B, and NS5). Among these non-structural proteins, NS5

comprises of an N-terminal methyltransferase (MTase) domain and a C-terminal RNA-

dependent RNA polymerase (RdRp) domain, each performing the functions of viral RNA

capping and RNA synthesis, respectively (20–22). The mechanism of flaviviral MTase

involves a two-step process. In the first step, the enzyme transfers a methyl group from SAM

to the guanine base located at the 5' end of the viral RNA, forming a cap 0 structure. In the

second step, this cap 0 structure undergoes further modifications to form the cap 1 structure,

achieved by adding a methyl group to the ribose sugar of the initially transcribed nucleotide

(23,24).

Additionally, polyamines have been shown to be essential for the viral life cycle in these RNA

viruses (25,26). There are three biogenic polyamines present in the host of these viral

pathogens, namely, putrescine (put), spermine (spm), and spermidine (spd), which are

produced from ornithine through the polyamine biosynthetic pathway (27). Enough evidence

suggests that polyamines are essential for various stages of the virus life cycle, including

genome replication, virus protein translation, and genome packaging (26). The inhibition of

the first enzyme, ornithine decarboxyalse 1 (ODC1) of polyamine biosynthetic pathway has

been shown to exhibit antiviral activity against CHIKV, DENV, ZIKV etc (25,26).

Difluoromethylornithine (DFMO) is a well-known irreversible ODC inhibitor but is associated

with hearing loss and may lead to antiviral resistance because depletion of polyamines in

CHIKV infected cells leads to muatations in nsP1 and also in the Opal stop codon (28,29).

Additionally, a higher dose of DFMO is required for effective antiviral activity, and DFMO is

also known to induce the uptake of putrescine and spermidine (30).

The dodecameric cryo-electron microscopy (Cryo-EM) structures of CHIKV nsP1 (PDB IDs:

6Z0V, 6Z0U, and 7DOP) reveal that the upper ring exhibits bifunctional MTase/GTase

catalytic activity, whereas the lower ring is responsible for oligomerization and membrane

binding (11,12). Similarly, the dimeric structure of DENV NS5 has been reported (PDB ID:

4V0R), revealing its organization of functional domains. Recently, the crystal structure of the

NS5 MTase from the Omsk hemorrhagic fever virus (OMHV), which belongs to the

Flaviviridae family, has been determined. This structure reveals new insights, including the

presence of GMP derived from GTP through the action of guanyl transferase (GTase), GMP-

arginine adducts, and an uncommonly observed capped RNA conformation (21).

SAM-dependent MTases are categorized as N-, O-, or C- MTases (N-MTases, O-MTases, C-

MTases) according to the specific atom (nitrogen, oxygen, or carbon) they methylate in their

substrates (31). These enzymes utilize SAM to transfer methyl groups to compounds such as

flavonoids and phenolic acids, altering their chemical properties and biological activities.

Given their specificity for such substrates, employing these compounds as potential inhibitors

is a logical approach for studying enzyme inhibition (31,32). The structure of DENV NS5

MTase and CHIKV nsP1 belong to a family of SAM-dependent MTases (12,33). Moreover,

the "Rossmann fold" is a βαβ super-secondary structure that forms the catalytic core

responsible for SAM binding site in all SAM-dependent MTases, including DENV NS5 MTase

domain and CHIKV nsP1 protein (34–37). Here, highly conserved acidic residues interact with

the ribose moiety of SAM, while the glycine-rich loop region interacts with its adenosine

moiety (35).

These structural insights enhance the efficiency of high-throughput computational screening

for novel natural compounds. Identifying new antiviral agents is a multifaceted and resource-

intensive endeavor. Natural compounds, ubiquitously present in various food items, offer the

advantage of potentially exhibiting a lower side effect profile than synthetic pharmaceuticals

(38). Consequently, structure-based virtual screening represents a promising approach for

identifying compounds with potential pharmacological activity (39). In the current study, an *in*

house library of natural substrates of N-C-O-MTases (NSMT), was curated and screened

against CHIKV nsP1 and DENV NS5 MTase. High throughput virtual screening and

experimental validation pinpointed Herbacetin (HC) and Caffeic acid phenethyl ester (CAPE)

as potent MTase inhibitors.

<u>Materials and methods</u>

Multiple Sequence Alignment (MSA)

The amino acid sequence of the nsP1 protein of alphaviruses was compared with CHIKV nsP1,

and the NS5 MTase domain of orthoflaviviruses was compared with the DENV 3 NS5 MTase

protein. CHIKV nsP1 and DENV 3 NS5 MTase were utilized as reference points for these

alignments. The sequence alignment profile of the selected nsP1 and NS5 MTase sequences

was performed via Clustal Omega tool and analyzed by a graphical colored depiction using

ESPript 3.0 (40). A comprehensive description of the protocol is given in the supplementary

materials.

Structure-guided virtual screening of an *in house* library of NSMT

An *in house* NSMT compound library consisting of approximately 25 molecules, comprising

flavone and catechol derivatives, was constructed (31). The three-dimensional structures of

these molecules were obtained from the PubChem database

(https://pubchem.ncbi.nlm.nih.gov/). The library was subsequently applied for virtual

screening analyses, employing PyRx 0.8 (41). Each compound was then energy minimized

using the Universal Force Field (UFF) in Open Babel and converted to .pdbqt format (42). The

Cryo-EM structures of CHIKV nsP1 (PDB ID: 6Z0V) and the crystal structure of DENV-3

NS5 MTase (PDB ID: 4V0R) were retrieved from the RCSB Protein Data Bank (PDB). Given

that the structural data for these proteins were incomplete, with several residues in flexible

loops missing, SWISS-MODEL was utilized to construct complete models of these proteins,

ensuring accurate structural representations for virtual screening (43). Before screening, .PDB

were converted into .PDBQT format to facilitate their use in AutoDock Vina for docking (44).

The grid focused on the SAM/GTP binding site of CHIKV nsP1 and the GTP binding site of

DENV-3 NS5 MTase (41). The GTP site was selected for DENV-3 NS5 MTase due to the

frequent presence of SAM/SAH in available crystal structures, indicating a high prevalence of

these interactions in the binding site. For CHIKV nsP1, the grid dimensions were set at 33.48

Å × 19.99 Å × 32.10 Å, centered at X = 54.92, Y = 137.80, Z = 91.35, and for DENV-3 NS5

MTase, the grid dimensions were 34.36 Å × 36.68 Å × 24.04 Å, centered at X = 13.79, Y =

134.12, Z = 0.54. Following the initial screening, the top compounds were selected on the basis

of binding energy, ligand pose, and key interactions with the target sites as well as the reported

antiviral activity, and were docked again using AutoDock Vina (44). The same grid parameters

were employed to maintain consistency. The docked conformations were subsequently

analyzed using PyMOL for structural visualization and LigPlot+ to comprehensively map

hydrogen bonding and hydrophobic interactions (45,46).

Molecular Dynamics (MD) Simulation

nsP1 apo and in complex with SAM, HC, and CAPE and DENV 3 NS5 MTase apo and in

complex with GTP, HC, and CAPE were subjected to MD simulation studies to assess dynamic

behavior and stability of protein-ligand interactions. The GROMACS 2022.2 suite was used to

carry out all simulation studies using the CHARMM36 force field on a LINUX-based

workstation (47,48). Ligand parameters and topology files were generated using the

CHARMM General Force Field (CgenFF) program (49). The SPC (Simple Point Charge)

water type for solvation and counter ions (Na+ and CL-) were added to neutralize the cubic

system. The energy minimization step was performed using the steepest descent method with

a maximum force of 10 kJmol-1. Followed by a two-phased equilibration of a constant number

of particles, volume, and temperature (NVT), a constant number of particles, pressure, and

temperature (NPT) for 100ps each. Periodic boundary conditions were applied at a constant

temperature of 300 K and 1 atm pressure, utilizing the V-rescale temperature coupling method

and Berendsen pressure coupling method, respectively (50). The long-range electrostatic

interactions were calculated using the Particle Mesh Ewald method, while bond lengths

involving heavy atoms were constrained with the Linear Constraint Solver (LINCS) algorithm

(51). Short-range forces were computed with a minimum cutoff of 1.2 nm, employing the

Verlet cutoff scheme (52). Finally, 100 ns MD production run was performed using the leap-

frog algorithm with an integration time frame of 2fs, and the trajectories were generated after

every 10 ps. The resulting trajectories were analyzed for structural deviations and fluctuations

within the protein and protein-ligand complexes .

Cloning, expression, and purification of DENV 3 NS5 MTase Domain

For the cloning of DENV 3 NS5 MTase Domain, RNA was isolated from the supernatant of

DENV 3 infected cells using the TRIzol (Sigma) method described by the manufacturer. The

RNA was then reverse-transcribed into complementary DNA (cDNA) using the PrimeScript

cDNA Synthesis Kit (Takara). The DENV 3 cDNA was subsequently amplified by PCR, using

the following primers (amino acid residues 1-278), NS5 MTase forward, 5′-

ATGC<u>GCTAGC</u>GGAACAGGTTCACAAG-3′,   and   NS5   MTase   reverse   5′-

CGCA<u>CTCGAG</u>TCAATTAACATGTCGAGTT-3′, the sequences underlined are the

recognition sites of the restriction enzymes *NheI* and *XhoI* respectively. After digestion with

restriction enzymes, the PCR product was purified with a Qiagen gel extraction kit per the

mentioned kit protocol and inserted between the restriction enzymes sites of the vector pET28c

(+). The recombinant vector pET28c (+) NS5 MTase was identified by restriction digestion,

and the DENV 3-NS5 MTase insert was verified by sequencing. *E. coli* DH5α was used for

amplification of the recombinant plasmid, and *E. coli* BL21 (DE3) was transformed for induced

expression of the His-tagged DENV 3-NS5 MTase protein. Expression and purification was

done, as mentioned in Boonyasuppayakorn et al. 2014 (53). Eluted fractions were dialyzed,

and further protein were concentrated to the required concentration with amicon centrifugal

filters (10,000 MWCO Millipore, Burlington, MA, USA), flash-frozen in liquid nitrogen, and

stored at −80 °C till its use. A detailed description of this protocol is given in the supplementary

materials.

Expression and purification of recombinant CHIKV nsP1

228   As mentioned by Kaur et al. in 2018, *E. coli* Rosetta cells were utilized for the expression and

229   purification of recombinant CHIKV nsP1 using the recombinant expression plasmid

230   pET28c(+)-CHIKV nsP1 (54,55).

Tryptophan Fluorescence Spectroscopy

Tryptophan fluorescence spectroscopy (TFS) experiments were conducted using a Fluoromax

fluorescence spectrophotometer (Horiba Scientific). A quartz cuvette with dimensions of 5x5

mm was utilized. The excitation wavelength was set to 280 nm, and the emission wavelength

was scanned from 295 to 540 nm. A slit width of 5 nm was employed for all measurements.

nsP1 and NS5 MTase protein samples were prepared at concentrations of 1 μM and 0.15 μM,

respectively, in a 500 μl phosphate-buffered saline (PBS) solution. Variable concentrations of

SAM (S-adenosylmethionine), HC, and CAPE were added to the protein samples. The

experiments were conducted at 25°C. Control buffer experiments and compound titrations were

performed parallel with the main experiments for background determination and used for

subtraction. Data from three independent experiments were collected and analyzed using

nonlinear regression with the 'One Site-Specific Binding' model. The data analysis was carried

out using GraphPad Prism 8 software (55).

Capillary electrophoresis (CE) based DENV 3 NS5 MTase and CHIKV nsP1 inhibition

assay

To determine the enzyme inhibition activities of HC and CAPE, the enzymatic reaction for of

DENV 3 NS5 MTase and CHIKV nsP1 were conducted using specific reaction mixtures. The

reaction mixture for NS5 MTase consisted of 50 mM Tris buffer (pH 7.5), 10 mM KCl, 2 mM

DTT, 2 mM $MgCl_2$, 0.3 mM SAM, and 0.3 mM GTP, along with 1 μM NS5 MTase protein.

The same components were used for the nsP1 enzyme reaction with 5 μM nsP1 protein. These

enzyme reactions were performed at 37 °C for 1 h. To establish negative controls, both enzyme

assays included a reaction with no SAM-GTP (substrate). Furthermore, to provide an additional

negative control, the NS5 MTase reaction was also conducted with the inactive capsid of

CHIKV. After incubation reaction was stopped by adding acetonitrile in 1:2 (vol/vol), 100 μM

caffeine was used as internal control (13,14). The mixture was vortex-mixed for 15 s, and the

protein was precipitated for 20 min at 18,500 x g. The supernatant was transferred to sample

vials. The detected SAH values were normalized to the internal standard for each capillary

electrophoresis run. CE analysis was performed as mentioned in Mudgal et al. 2020. Data from

three independent experiments were collected and analyzed using GraphPad Prism 8 software

(13,14).

Crystallization of NS5 MTase

The crystallization of DENV 3 NS5-MTase was conducted using the sitting drop vapor

diffusion method in 96-well plates at a temperature of 20°C, as described in Coutard et al.,

2014 (56,57). For data collection, DENV 3 NS5-MTase crystals in complex with CAPE were

obtained by soaking the crystals in a cryoprotectant solution (22.5% glycerol) containing 1mM

CAPE for 1 h. They were flash-frozen in a nitrogen stream at 100K. Diffraction data were

collected at the home source (Rigaku Micromax 007 HF), Macromolecular crystallography

unit, IIT Roorkee. Data reduction and scaling were done using the CrysAlis Pro software

(Rigaku Inc.). The structure was determined by MOLREP using DENV 3 NS5 MTase (PDB

code 4R8R) as a search model (58). Iterative rounds of model-building in COOT and the

refinement of atomic coordinates and B-factors using refmac5 (59) in CCP4i2 allowed for the

correct placement of sidechains and loops. NCS and jelly-body restraints were used throughout

the data refinement. Additionally, to confirm ligand localizations in the structure, omit maps

were generated in Polder (60). The data collection and refinement statistics are summarised in

Table 3. All figures of protein and ligand structures were prepared using PyMOL (45).

Cell line, Virus isolation, propagation and serotyping

DENV was isolated from the Dengue-suspected patient's blood samples and was further

confirmed as a DENV 3 serotype after sequencing. Vero cells were used for the propagation

and titration of DENV and CHIKV. CHIKV (Accession No. KY057363.1.) was propagated

and titrated using the protocol reported by Singh et al., 2018 and then stored at −80°C for

further experiments (61). A comprehensive description of the protocol is given in the

supplementary materials.

Preparation of HC, CAPE, and DFMO stock solutions

HC, CAPE, and DFMO were purchased from Cayman, USA. For experiments, the stock

solutions (HC: 165.45mM; CAPE: 20mM) for both compounds were prepared in dimethyl

sulfoxide (DMSO) (Sigma-Aldrich) and filtered through a 0.2 μm size syringe filter

(Millipore). The 100 mM DFMO dilution was prepared in sterile tissue culture-grade water

that was used as a positive control for the polyamine-related experiments. Further dilutions

were prepared in 2% DMEM media (maintenance media) before use. Similarly, stock solutions

of individual polyamines (putrescine, spermine, and spermidine) (Alfa-Aesar, USA) were

diluted in sterile tissue culture-grade water and used as specified.

Assessment of Antiviral Activity

Vero cells were seeded onto a 24-well plate at a cell density of 1.0×10^5 cells/well. Compounds

with concentrations that maintained cell viability above 90% were used for the treatment. Cells

were treated with compounds 12 h before infection. With a multiplicity of infection (MOI) of

0.1, the cell monolayer was infected with the virus with gentle shaking every 15 min for 2 h.

The inoculum was removed, and the cell monolayer was washed twice to ensure no chance of

secondary infection. Compounds were added in the post-infection (pi) 2% DMEM media and

incubated for 24 h. After 24 hpi, the supernatant was collected for the CHIKV antiviral

experiment. On the other hand, fresh maintenance media was added to the DENV antiviral

experiment for 4 days. Then, the supernatant was collected to determine the virus titer via

plaque-forming assay (14,62). A comprehensive description of the cell viability assay and

plaque-forming assay given in the supplementary materials.

Immunofluorescence Assay

Vero cells were seeded onto a 6-well plate at a cell density of 1 x 10^6 cells/well. Cells were

treated with compounds, as mentioned earlier. The cells were washed three times with PBS

and then fixed with 3.7% formaldehyde for 30 min at room temperature, followed by

permeabilization with 0.1% Triton-X-100 for 10 min. After washing the cells with 1X PBS,

they were incubated with antibodies against CHIKV and DENV (anti-alphavirus 1:100, Santa

Cruz Biotechnology Inc.; 1:500 diluted orthoflavivirus group antibody, Genetex Inc.) for 1 h,

followed by a wash with 0.1% Tween-20 in 1X PBS (PBST). Later, the plate was incubated in

the dark with fluorescein (FITC)-conjugated secondary anti-mouse antibody (1:250, Sigma)

for 30 min at 37 °C. The cells were then washed with PBST and counter-stained with 4′,6-

diamidino-2-phenylindole (DAPI, Sigma) for 15 min in the dark. Finally, the images were

captured using a fluorescence microscope (EVOS FL AUTO, Thermo Fisher).

Quantitative Real-Time PCR (qRT-PCR)

HC and CAPE compounds treatment and infection were the same as described in the assessment of antiviral activity. After the termination of an assay, TRIzol was added to the plate. RNA was purified according to the manufacturer's protocol. Purified RNA was quantified for cDNA preparation using the PrimeScript 1st strand cDNA Synthesis Kit (Takara) with 400 ng of extracted RNA. The KAPA SYBR Fast Universal qPCR Kit was used for qRT-PCR with analysis performed on the QuantStudio™ 5 System from Applied Biosystems, USA. The forward and reverse primers used for amplification are used as previously described in (63) and (55,64). β-actin was utilised as an internal control. The relative quantification was carried out using the ΔΔCt method as described (14,55).

Polyamine addition assay

In the polyamine addition assay, compounds pre-treatment was given for 24 h at various concentrations such as HC (200 µM), CAPE (25 µM for CHIKV and 2.5 µM for DENV), DFMO (1000 µM). Immediately after 2 h of virus infection with CHIKV and DENV, cells were treated with 1 µM biogenic polyamines (put, spd, spm) for 24 h and 120 h, respectively. Later, the supernatant was collected to determine the virus titer by plaque-forming assay.

Polyamine determination by Thin-layer chromatography (TLC)

Vero cells were treated with CAPE (25 µM), HC (200 µM), and DFMO (1000 µM) for 36 h. Further, cells were trypsinized, sonicated, and centrifuged. Polyamines were separated by TLC as previously described (65). Briefly, cells were sonicated in 500 µL of 2% (V/V) perchloric acid. Sonication was performed at 4 ºC (20 kHz, 2Ω, 10-s pulse, 30-s rest). Cell homogenates obtained are then stored at 4 ºC for 24 h. After this, samples are centrifuged at ~11,500 × g for 30 min at 4 ºC. An equal ratio of supernatant dansyl chloride (5 mg/mL) (Alfa-Aesar, USA) was added in acetone and saturated sodium bicarbonate. Samples were incubated in the dark

overnight at room temperature. Excess dansyl chloride was cleared by incubating the reaction

with 150 mg/mL proline (HiMedia). Dansylated polyamines were extracted with 100 µL

toluene. 10 µL of the sample was added in small spots to the TLC plate by automated sprayer

(Silica gel matrix; Millipore) and exposed to ascending chromatography with cyclohexane:

ethyl acetate (2:3). The plate was dried and visualized via exposure to UV. Further, TLC images

were quantified utilizing ImageJ software.

Statistical analysis

GraphPad Prism 8 software is used for data analysis. One-way ANOVA was used to determine

statistical significance wherever mentioned.

350
<u>Results</u>

Structural and sequence conservation of nsP1 and NS5 MTase active sites:

Both the viral MTases, nsP1 of CHIKV and NS5 of DENV, bind and use the cofactor SAM as

the methyl donor with the release of the byproduct SAH. Structural studies have revealed that

the MTase domain in both viral proteins shares common structural features of the Rossmann

fold, indicating that these shared characteristics can be exploited to identify a common

inhibitor. The SAM binding site in CHIKV nsP1 as revealed by its Cryo EM structure (PDB

ID 8AOX, 8AOV) is lined by residues Gly65, Ser66, Ala67, Pro83, Arg85, Ser86, Asp89,

Thr137, Asp138, and Gln151. GTP binding site is lined by residues Ala40, Arg41, Ser44,

Tyr154, Phe178, Phe241, Val243, Thr246, Tyr248 and Glu250. The residues that make

molecular contacts with both SAM and GTP are Arg70, Arg92, and Asp152 (11,12). Similarly,

the crystal structural analysis of DENV 3 NS5 MTase (PDB ID 4V0R) revealed that the GTP

binding site residues for orthoflaviviruses are Asn15, Arg19, Lys26, Ser147, Arg208, Ser210,

Leu14, Phe22 and Thr211 (66,67).

Alphaviruses that are known to cause infections in humans are VEEV, CHIKV, Ross River

virus (RRV), Sindbis virus (SINV), Aura Virus (AURA), Middelburg virus (MIDV), Barmah

Forest virus (BFV), Madariaga virus (MADV), and Mayaro virus (MAYV). Additionally,

Salmonid alphavirus (SAV) exhibits tropism for Atlantic salmon, inducing pancreatic disease,

while Eilat virus (EILV) demonstrates insect-specific virulence (68,69). DENV 3, ZIKV, West

Nile Virus (WNV), Yellow Fever Virus (YFV), and Japanese Encephalitis Virus (JEV) are

some of the human-infecting orthoflaviviruses. In addition, Palm Creek virus (PCV) exhibits

insect specificity, and Wenzhou shark flavivirus (WSF) falls within the category of aquatic

orthoflaviviruses (70,71). The primary sequence alignment of MTases and subsequent

comparison among alphaviruses and orthoflaviviruses demonstrated a significant degree of

sequence similarity, as illustrated in supplementary figure 1A. In alphaviruses, 81% of

SAM/GTP binding residues in nsP1 are conserved across different viruses. Similarly, in

orthoflaviviruses, 66% of GTP binding residues are conserved across NS5 MTase

(supplementary figure 1B).

Structure-based identification of nsP1 and NS5 MTase inhibitors

The generated 3D models of CHIKV nsP1 and DENV 3 NS5 MTase domains showed high

structural similarity to templates with root mean square deviation (RMSD) values 0.205 Å and

0.183 Å, respectively (supplementary figure 2 A,B). These models were used for virtual

screening of *in house* NSMT library, identifying HC and CAPE as potential inhibitors of nsP1

and NS5 MTase. For nsP1, HC exhibited the highest binding affinity with a binding energy of

-8.0 kcal/mol, facilitated by hydrogen bonds with Arg92, Arg70, and Glu88, along with

hydrophobic interactions involving Arg41, Tyr285, Asp152, Phe178, Val243, Ala40, and

Tyr248 residues nsP1 (Figure 1C, 2C) (Table 1). CAPE demonstrated binding energy of -7.6

kcal/mol, forming hydrogen bonds with Glu250 and Arg92 and engaging in hydrophobic

interactions with Tyr248, Arg70, Asp152, Tyr154, Ala40, Phe178, Val243, and Tyr285

residues of nsP1 (Figure 1B, 2B) (Table 1). HC and CAPE exhibited binding affinities

comparable to binding affinities of SAM and GTP (positive control) with nsP1 (Figure 1A,D,

2A,D) (Table 1). Similarly, in the case of NS5 MTase, HC displayed binding energy -6.7

kcal/mol with hydrogen bonds formed by Lys11, Asn15, Ser147, and Ser210, along with

hydrophobic interactions involving Leu14, Phe22, Pro149, and Thr211 residue of NS5 MTase

(Figure 1G, 2G) (Table 1). CAPE exhibited binding energy -6.6 kcal/mol with hydrogen bonds

established with Lys11, Leu14, and Leu17 and engaged in hydrophobic interactions with

residues such as Asn15, Ser18, and Ser210 residue of NS5 MTase (Figure 1F, 2F) (Table 1).

For NS5 MTase, HC and CAPE exhibited slightly lower binding energy than GTP (positive

control) (Figure 1E, 2E) (Table 1).

The dynamic stability and conformational changes in the Apo proteins and protein-ligands

complexes were studied by analyzing their RMSD values during the MD simulation. The

average RMSD values for nsP1 Apo was 0.35 nm, and for nsP1 complexes nsP1-SAM, nsP1-

HC, nsP1-GTP, and nsP1-CAPE were 0.53 nm, 0.42 nm, 0.46 nm and, 0.38 nm respectively

(supplementary figure 3A). Similarly, the Average RMSD for NS5 MTase Apo was 0.23 nm

and the NS5 MTase complexes NS5 MTase-GTP, NS5 MTase-HC, and NS5 MTase-CAPE

were 0.18 nm, 0.25 nm, and 0.20 nm respectively (supplementary figure 3B). These

observations depict that the binding of molecules to nsP1 and NS5 proteins resulted in the

formation of a stable complex.

Binding of compounds to DENV 3 NS5 MTase and CHIKV nsP1

The interactions of NS5 MTase and nsP1 with SAM, GTP, HC, and CAPE at various

concentrations were examined using TFS. The intrinsic fluorescence of the native NS5 MTase

and nsP1 proteins was measured utilizing a spectrofluorometer. Both proteins showed intrinsic

fluorescence quenching with increasing concentrations of all compounds (Figure 3). In TFS, a

red shift, indicats to a shift to longer wavelengths, suggests increased polarity or decreased

hydrophobicity around the tryptophan residues. Conversely, a blue shift, reflecting a move to

shorter wavelengths, signifies reduced polarity or increased hydrophobicity in the local

environment (72). A dose-dependent red shift was observed in both proteins upon interaction

with HC and CAPE (supplementary figure 4). For nsP1 interactions with SAM, GTP, HC, and

CAPE, the K_D values were determined to be 10.88 ± 5.15 µM, 169.1 ± 21.31 µM, 6.52 ± 0.55

µM, and 30.57 ± 5.33 µM, respectively (Figure 3 A-D). Similarly, for NS5 MTase interactions

with SAM, GTP, HC, and CAPE, the K_D values were determined to be 6.67 ± 2.57 µM, 3.03

± 1.0 µM, 13.35 ± 2.69 µM, and 37.93 ± 6.32 µM, respectively (Figure 3 E-H).

Inhibition of Virus-specific MTase by HC and CAPE

To further validate the inhibition of viral MTase activity by HC and CAPE, recombinantly

expressed CHIKV nsP1 and DENV NS5 MTase were utilized. Single bands of ~ 56 kDa and

~ 32 kDa were observed on 12% SDS-PAGE for purified nsP1 and NS5 MTase, respectively (

Figure 4 A,D). The enzymatic activity of nsP1, as assessed through CE-based methods as

mentioned in the previously published protocol (14,55). A dose-dependent decrease in MTase

activity was observed when treated with HC and CAPE, confirming the significant inhibitory

activity of both compounds against CHIKV nsP1 (Figure 4 B,C). Similarly, to validate the

inhibition of NS5 MTase activity by HC and CAPE, the CE-based enzymatic activity assay of

purified DENV 3 NS5 MTase was performed (14,55). The MTase activity was reduced in

response to increasing compound concentrations, confirming the inhibitory effects of HC and

CAPE against DENV 3 NS5 MTase (14,55) (Figure 4 E,F).

Structural insights into DENV 3 NS5 MTase with CAPE

The structure of DENV 3 NS5 MTase-CAPE complex (PDB:8KDZ) at 2.6 Å was determined

to identify the residues involved in binding to the CAPE molecule (Table 3). Although no SAH

was present during crystallization, it was found in both chains, indicating that it has originated

from *E. coli* during protein expression (PDB:8KDZ). The main chain conformation in the

DENV 3 NS5 MTase-CAPE complex is nearly identical to the DENV 3 NS5-SAH-GTP

structure (PDB:4V0R) with RMSD 0.367 Å over 225 Cα atoms (Figure 5C). Two CAPE

(302,303) molecules could be modelled in one chain of DENV 3 NS5 structure that matched

the observed electron density with a real space correlation coefficient (RSCC) of 0.71 and

0.81, respectively (Figure 5B). The detailed analysis determined by PyMOL and LigPlot[+], as

outlined in Tables 4 and 5, reveals the formation of hydrogen bonds and specific hydrophobic

interactions, which are pivotal for the GTP binding site. Arg 211 has an important role in

interacting with the phosphate group of GTP (67) (Figure 5E). In CAPE (302), O4 of the 3,4-

dihydroxy phenyl group forms a hydrogen bond with NH2 of Arg 211 at 3.5 Å. 3,4-dihydroxy

phenyl group of both CAPE form π- π T-shaped interaction with one another. CAPE (302)

forms π-σ interaction with Thr 214. Also, its phenyl ring forms π-alkyl interactions with Leu

17, Pro 152. CAPE (303) 3,4-dihydroxy phenyl group acts as π donor to nitrogen of Ser 150

(Figure 5D). Furthermore, CAPE (302) hydrophobically interacts with Asn18, Leu17. These

residues are essential in binding the initial nucleotides from octameric RNA (specifically

guanosine triphosphate adenosine (G3A) to DENV 3 NS5 MTase (73) (Figure 5F) (Table 6).

The CAPE (302-303) molecules also hydrophobically interact with residues Ser150, Thr214,

Lys180, and Gly148. Among these residues, Ser150, Thr214, and Lys180 are crucial for

hydrogen bonding, while Gly148 is important for the hydrophobic interaction of the adenosine

(A)-guanosine (G) dinucleotide from cap 0 RNA to DENV 3 NS5 MTase (22) (Figure 5G, H)

(Table 7). Therefore, the presence of CAPE at this position could potentially impede the

binding of viral RNA to the DENV 3 NS5 MTase.

Polyamine-depletion and antiviral activities of HC and CAPE

The MTT colorimetric assay was performed for testing the cytotoxicity of HC and CAPE. The

compound concentrations with cell above 90% viability were used subsequently

(supplementary figure 5). TLC experiments were performed to determine the effect of HC and

CAPE on polyamine levels in mammalian cells treated with these compounds. Vero cells were

treated with HC, CAPE, and DFMO, and TLC results confirmed a reduction in the levels of all

three polyamines. Put, spd, and spm are positive control markers (Figure 6B). After treatment

of Vero cells with CAPE, HC, and DFMO, overall residual polyamine levels are 28.33%,

29.67 %, and 46 %, respectively, compared to cell control. Overall, CAPE and HC showed

higher polyamine depletion at lower concentrations as compared to the positive control

(DFMO) and the cell control (Figure 6B). For this study, the DENV serotype was isolated from

a clinical sample, and the DENV 3 serotype was confirmed by PCR using virus and serotype-

specific primer pairs and sequencing. The basic local alignment search tool (BLAST) tool

revealed 99.78% identity of isolated DENV serotype to DENV 3 isolate NU1883 polyprotein

(POLY) gene, partial cds (coding sequences). Similarly, CHIKV (Accession No.

KY057363.1.) was propagated in Vero cells and used in further study. Antiviral activities of

HC and CAPE at various concentrations were determined against CHIKV and DENV in plaque

reduction assays performed in Vero cells using each compound's non-cytotoxic concentrations.

By comparing the viral titer of untreated CHIKV-infected cells with HC- and CAPE-treated

CHIKV-infected cells, a dose-dependent decrease in CHIKV titer was observed for HC and

CAPE, with IC_{50} values of approximately 13.44 ± 3.21 µM and 0.57 ± 0.03 µM, respectively

(Figure 6 C, D supplementary figure 6 A, B). Likewise, HC and CAPE treatment to Vero cells

has shown a dose-dependent viral titer decrease in the DENV-infected cells was observed for

HC and CAPE, with IC_{50} values of approximately 7.24 ± 2.51 µM and 1.01 ± 0.14 µM,

respectively (Figure 6 F, G supplementary figure 6 C, D). DFMO, a known ODC inhibitor, was

used as a control and showed less antiviral efficacy against CHIKV at much higher

concentrations of 1000 µM than HC and CAPE.

Here, qRT-PCR was used to validate the antiviral effect of HC and CAPE by quantifying

DENV and CHIKV RNA levels in the infected cells. qRT-PCR showed a significant (p <

0.0001) reduction in the viral RNA levels when treated with HC and CAPE compounds in an

antiviral assay for CHIKV (Figure 6E) and DENV (Figure 6H). At the mentioned

concentrations, HC and CAPE showed a 23- and 4-fold reduction in DENV, respectively,

compared to the virus control (VC). Similarly, HC and CAPE reduced CHIKV by 25 and 97

fold at the same doses, respectively, compared to VC. IFA results further corroborated these

results. The results of the IFA analysis showed a reduction in CHIKV (Figure 7A) and DENV

(Figure 7B) after treatment with HC and CAPE.

Further, the effects of the addition of exogenous biogenic polyamines were studied. Unlike

DFMO-treated cells, the exogenous addition of all three polyamines at 1 µM were unable to

effectively rescue virus titer after CAPE and HC treatment for both CHIKV and DENV (Figure

8).This finding suggests an additional mechanism is involved in the antiviral action of these

compounds against both viruses, supporting the initial hypothesis.

Discussion

The continuous emergence and re-emergence of viruses like DENV and CHIKV underscores

the need for effective public health interventions and antiviral therapeutics to mitigate their

spread. Although FDA-approved vaccines for DENV and CHIKV are available, no antiviral

therapies are approved against these viral infections (4,7). Developing novel strategies to

combat these life-threatening human pathogens, by targeting both viral enzymes and host

factors, holds significant promise for accelerating drug development. This dual approach can

potentially enhance therapeutic efficacy and circumvent resistance by addressing critical

components of viral replication and host-pathogen interactions. Interestingly, SAM and GTP

binding sites in nsP1 and the GTP binding site in NS5 MTase are conserved across alphaviruses

and orthoflaviviruses infecting humans, insects, and aquatic life (supplementary figure 1 A,B).

An *in house* library of NSMT was screened against the CHIKV nsP1 and DENV NS5 MTase.

The screening identified HC and CAPE as the top hits with stable interactions to the active

sites of both enzymes.

In tryptophan fluorescence spectroscopy, the binding affinity of HC and CAPE with CHIKV

nsP1 and DENV NS5 MTase was determined and compared to that of SAM and GTP. SAM

and GTP induced fluorescence quenching without spectral shifts, indicating localized changes

near tryptophan residues without major conformational alterations (supplementary figure 4)

(72). Conversely, both HC and CAPE have demonstrated a dose-dependent red shift with

fluorescence quenching, indicating major structural changes upon interaction (Figure 3 and

supplementary figure 4). HC and CAPE have shown dose-dependent inhibition in nsP1 and

NS5 MTase CE-based MTase assays (Figure 4).

The crystal structure of CAPE in complex with the DENV 3 MTase reveals that two CAPE

molecules bind to the GTP and cap 0 RNA binding site (Figure 5). Similarly, structural analysis

of NS5 MTase and CAPE (303, 302) complex revealed a similar binding pattern of CAPE

molecule with residue Lys14, Leu17, Asn18 (CAPE-303) and Gly148 for (CAPE-302) of NS5

MTase as predicted in docking studies (Table 2,4). Among all hydrophobic interactions of

CAPE, residues Leu17, Asn18, Phe25, Ser150, Gly148, Glu149, Ser213, and Lys180 are

conserved across the NS5 MTase, as illustrated (supplementary figure 1B) (Table 4). Most of

these residues are involved in maintaining structural stability, with Lys180 being part of the

Lys61-Asp146-Lys180-Glu216 catalytic tetrad (K-D-K-E). Here, Lys180, acting as the

catalytic base, activates the 2'-OH group of the ribose sugar to facilitate nucleophilic attack on

the methyl group carbon atom of SAM (22). The CAPE binding site encompasses magnesium

(Mg), critical in stabilizing the structural integrity between cap 0 and RNA (Figure 5 G). This

is achieved through the hexacoordination of Mg^{2+} ions, involving three oxygen atoms from

phosphate and three from water molecules. Furthermore, Mg^{2+} ions form hydrogen bonds with

residues Ser213 and Ser150 protruding from the protein surface (22). Given its proximity to

this site, CAPE has the potential to influence the stability of the cap 0-RNA complex. In

OMHV, GMP-enzyme intermediate was identified within the GTP binding site (22). Hence, it

is possible that CAPE molecules in the GTP binding site can affect the viral RNA binding and

interfere with the capping process. Although this work did not address the possibility of inhibiting viral RNA binding activity, incorrectly capped viral RNA can lead to inefficient translation of viral proteins (20,21). AT-9010 (PDB: 8BCR) and ribavirin triphosphate (PDB: 1R6A) have previously been observed binding to the GTP binding site of the NS5 MTase. Both compounds, which are GTP analogues, exhibit anti 2′-O-MTase activity (Table 8) (74,75). While using nucleoside analogues is a promising antiviral strategy, it can potentially disrupt cellular functions and lead to the rapid development of resistance (76).

Polyamines are ubiquitous molecules present in all cells and play a crucial role in the viral life cycle (77). Polyamine depletion can impair cellular processes crucial for cell survival and homeostasis, such as gene expression, apoptosis, protein synthesis, oxidative stress levels, and cell signaling regulation (78–80). However, DFMO, which inhibits the ODC enzyme of the polyamine biosynthesis pathway, is an FDA-approved drug for African sleeping sickness and high-risk neuroblastoma (81,82). Moreover, DFMO is administered via a short-course regimen (14 days) in African sleeping sickness treatment to mitigate potential adverse effects (83,84). Previous studies showed that DFMO treatment cells depleted biogenic polyamine levels, leading to its antiviral activity against various viruses such as CHIKV, ZIKV, and DENV (25). This study supported a previously published finding that HC and CAPE are polyamine pathway inhibitors (85,86). HC is a naturally occurring flavonoid that is an established ODC inhibitor (86). Flavonoids have significant therapeutic potential as antiviral agents against various classes of viruses (87).

CAPE is a natural ester that has also shown antiviral activity, and it significantly prevented the full induction of ODC by epidermal growth factor (EGF) (85,88). HC and CAPE effectively reduced CHIKV and DENV infections more than DFMO in plaque reduction assays and IFA experiments, validating their superior inhibition in virus-infected cells (Figure 6,7). qRT-PCR quantified viral RNA from CHIKV and DENV-infected cells, reflecting viral replication. HC and CAPE treatment significantly reduced viral load compared to the VC (Figure 6).

This study shows HC and CAPE molecules have shown better polyamine depletion than the

DFMO (Figure 6B). An exogenous polyamine addition assay was conducted following the

methodology outlined by Mounce et al. (2016), adhering to established procedures.

Consistency was observed between the results of virus titer rescue in DFMO treatment upon

exogenous polyamine addition. However, unlike DFMO treatment for HC and CAPE, the

exogenous polyamine addition assay did not rescue viral titer for CHIKV and DENV (Figure

8 A, B). The inability of external polyamine supplementation to restore viral titers implies that

polyamine depletion is not the sole mechanism of action. Therefore, the data supports the

hypothesis of an additional inhibitory mechanism, likely involving targeting viral MTases.

These novel natural antiviral molecules and the findings of this study will further assist in

developing broad-spectrum antiviral strategies against emerging alphaviruses and

orthoflaviviruses.

Conclusion

This research provides compelling first evidence of HC and CAPE's anti-CHIKV and anti-

DENV activities through two mechanisms: a) indirectly by depleting polyamines in

mammalian cells and b) directly by targeting viral MTases. The detailed atomic interactions

revealed that CAPE binding at the GTP and cap 0 RNA binding sites in DENV MTase may

impede viral RNA capping mechanism. HC and CAPE's novel broad-spectrum antiviral

activity makes them promising candidates for developing anti-DENV and anti-CHIKV

therapeutics.

Acknowledgments

The authors thank the Department of biosciences and bioengineering, IIT Roorkee for central

facilities and the translational and structural bioinformatics center. The authors thank the

Macromolecular Crystallographic Facility (MCU) at IIC, Indian Institute of Technology,

Roorkee. ST and PK thank the Department of Biotechnology, Govt of India, for supporting

Bioinformatics Center at IIT Roorkee ref number BT/PR40141/BTIS/137/16/2021. ST and PK

also would like to thank Department of Biotechnology, Govt of India "National Network

Project of Department of Biotechnology, Indian Institute of Technology, Roorkee" Project no.

BT/PR40142/BTIS/137/72/2023. The authors would like to acknowledge Mr. Mayur Ghate,

Ms. Bharati for technical assistance in TFS analysis. The authors would like to thank the

University Grants Commission (UGC), Council of Scientific and Industrial Research (CSIR),

Indian Council of Medical Research (ICMR), and Ministry of Human Resource

Development (MHRD) for providing financial support. This work was supported by a research

grant to ST from ICMR ref no. ISRM/12(46)/2020.

Ethical approval

604 DENV-positive clinical sample was collected from HiTech Pathology Laboratory, Roorkee

605 India with the approval of the Institutional Human Ethical Committee (IHEC), Indian Institute

606 of Technology, Roorkee (Ref no. BT/IHEC-IITR/2016/6499).

Supplemental information:-

608 Supplemental information includes supplemental experimental procedures and six Figures.

Authors contribution

610 MB, AK, RR, VS, AP, AK, SM, BW, and RK conducted the experiments. MB, AK, RR, VS,

611 RM, DS, PK, and ST planned the experiments and analyzed the data. MB, PK, and ST took the

612 lead in writing the manuscript. All authors provided critical feedback and helped shape the

613 research, analysis, and manuscript.

Competing interests

617 The authors declare that they have no competing interests.

References

1. Guzman MG, Gubler DJ, Izquierdo A, Martinez E, Halstead SB. Dengue infection. Nature reviews Disease primers. 2016 Aug 18;2(1):1-25.DOI: 10.1038/nrdp.2016.55

2. Powers AM, Brault AC, Shirako Y, Strauss EG, Kang W, Strauss JH, Weaver SC. Evolutionary relationships and systematics of the alphaviruses. Journal of virology. 2001 Nov 1;75(21):10118-31. DOI: 10.1128/JVI.75.21.10118-10131.2001

3. DENGUE/DHF SITUATION IN INDIA :: National Center for Vector Borne Diseases Control (NCVBDC) [Internet]. [cited 2024 Jan 15]. Available from: https://nvbdcp.gov.in/index4.php?lang=1&level=0&linkid=431&lid=3715

4. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM, Brownstein JS, Hoen AG, Sankoh O, Myers MF. The global distribution and burden of dengue. Nature. 2013 Apr 25;496(7446):504-7. DOI: 10.1038/nature12060

5. Lobo DA, Velayudhan R, Chatterjee P, Kohli H, Hotez PJ. The neglected tropical diseases of India and South Asia: review of their prevalence, distribution, and control or elimination. PLoS neglected tropical diseases. 2011 Oct 25;5(10):e1222. DOI: 10.1371/journal.pntd.0001222

6. CHIKUNGUNYA SITUATION IN INDIA :: National Center for Vector Borne Diseases Control (NCVBDC) [Internet]. [cited 2024 Jan 15]. Available from: https://ncvbdc.mohfw.gov.in/index4.php?lang=1&level=0&linkid=486&lid=3765

7. Geographical distribution of chikungunya virus disease cases reported worldwide, 2023 [Internet]. [cited 2024 Jan 15]. Available from: https://www.ecdc.europa.eu/en/publications-data/chikungunya-virus-disease-cases-january-december-2023

8. Kaur M, Singh K, Sidhu SK, Devi P, Kaur M, Soneja S, Singh N. Coinfection of chikungunya and dengue viruses: A serological study from North Western region of Punjab, India. Journal of laboratory physicians. 2018 Oct;10(04):443-7. DOI: 10.4103/JLP.JLP_13_18

9. Chahar HS, Bharaj P, Dar L, Guleria R, Kabra SK, Broor S. Co-infections with chikungunya virus and dengue virus in Delhi, India. Emerging infectious diseases. 2009 Jul;15(7):1077. DOI: 10.3201/eid1507.080638

10. Jose J, Snyder JE, Kuhn RJ. A structural and functional perspective of alphavirus replication and assembly. Future microbiology. 2009 Sep;4(7):837-56. DOI: 10.2217/fmb.09.59

11. Jones R, Bragagnolo G, Arranz R, Reguera J. Capping pores of alphavirus nsP1 gate membranous viral replication factories. Nature. 2021 Jan 28;589(7843):615-9. DOI: 10.1038/s41586-020-3036-8

12. Jones, R., Hons, M., Rabah, N., Zamarreño, N., Arranz, R., & Reguera, J. (2023). Structural basis and dynamics of Chikungunya alphavirus RNA capping by nsP1

capping pores. Proceedings of the National Academy of Sciences, 120(12), e2213934120. DOI: 10.1073/pnas.2213934120

13. Mudgal R, Bharadwaj C, Dubey A, Choudhary S, Nagarajan P, Aggarwal M, Ratra Y, Basak S, Tomar S. Selective estrogen receptor modulators limit alphavirus infection by targeting the viral capping enzyme nsP1. Antimicrobial Agents and Chemotherapy. 2022 Mar 15;66(3):e01943-21. DOI: 10.1128/AAC.01943-21

14. Mudgal R, Mahajan S, Tomar S. Inhibition of Chikungunya virus by an adenosine analog targeting the SAM-dependent nsP1 methyltransferase. FEBS letters. 2020 Feb;594(4):678-94. DOI: 10.1002/1873-3468.13642

15. Filipowicz W, Furuichi Y, Sierra JM, Muthukrishnan S, Shatkin AJ, Ochoa S. A protein binding the methylated 5'-terminal sequence, m7GpppN, of eukaryotic messenger RNA. Proc Natl Acad Sci U S A. 1976;73(5):1559–63. DOI: 10.1073/pnas.73.5.1559

16. Hyde JL, Diamond MS. Innate immune restriction and antagonism of viral RNA lacking 2′-O methylation. Virology. 2015 May 1;479:66-74. DOI: 10.1016/j.virol.2015.01.019

17. Liu L, Dong H, Chen H, Zhang J, Ling H, Li Z, Shi PY, Li H. Flavivirus RNA cap methyltransferase: structure, function, and inhibition. Frontiers in biology. 2010 Aug;5:286-303. DOI: 10.1007/s11515-010-0660-y

18. Ahola T, Kääriäinen L. Reaction in alphavirus mRNA capping: formation of a covalent complex of nonstructural protein nsP1 with 7-methyl-GMP. Proceedings of the National Academy of Sciences. 1995 Jan 17;92(2):507-11. DOI: 10.1073/pnas.92.2.507

19. Kalita JM, Aggarwal A, Yedale K, Gadepalli R, Nag VL. A 5-year study of dengue seropositivity among suspected cases attending a teaching hospital of North-Western region of India. Journal of Medical Virology. 2021 Jun;93(6):3338-43. DOI: 10.1002/jmv.26592

20. Selisko B, Dutartre H, Guillemot JC, Debarnot C, Benarroch D, Khromykh A, Desprès P, Egloff MP, Canard B. Comparative mechanistic studies of de novo RNA synthesis by flavivirus RNA-dependent RNA polymerases. Virology. 2006 Jul 20;351(1):145-58. DOI: 10.1016/j.virol.2006.03.026

21. Simmonds P, Becher P, Bukh J, Gould EA, Meyers G, Monath T, Muerhoff S, Pletnev A, Rico-Hesse R, Smith DB, Stapleton JT. ICTV virus taxonomy profile: Flaviviridae. Journal of General Virology. 2017 Jan;98(1):2-3. DOI: 10.1099/jgv.0.000672

22. Zhao Y, Soh TS, Lim SP, Chung KY, Swaminathan K, Vasudevan SG, Shi PY, Lescar J, Luo D. Molecular basis for specific viral RNA recognition and 2′-O-ribose methylation by the dengue virus nonstructural protein 5 (NS5). Proceedings of the National Academy of Sciences. 2015 Dec 1;112(48):14834-9. DOI: 10.1073/pnas.1514978112

23. Furuichi Y, Shatkin AJ. Viral and cellular mRNA capping: past and prospects. Adv Virus Res. 2000; 55: 135–184. DOI: 10.1016/s0065-3527(00)55003-9

24. Bisaillon M, Lemay G. Viral and Cellular Enzymes Involved in Synthesis of
mRNA Cap Structure. Virology. 1997 Sep 15;236(1):1–7. DOI:
10.1006/viro.1997.8698

25. Mounce BC, Poirier EZ, Passoni G, Simon-Loriere E, Cesaro T, Prot M,
Stapleford KA, Moratorio G, Sakuntabhai A, Levraud JP, Vignuzzi M.
Interferon-induced spermidine-spermine acetyltransferase and polyamine
depletion restrict Zika and chikungunya viruses. Cell host & microbe. 2016 Aug
10;20(2):167-77. DOI: 10.1016/j.chom.2016.06.011

26. Mounce BC, Cesaro T, Moratorio G, Hooikaas PJ, Yakovleva A, Werneke SW,
Smith EC, Poirier EZ, Simon-Loriere E, Prot M, Tamietti C. Inhibition of
polyamine biosynthesis is a broad-spectrum strategy against RNA viruses.
Journal of virology. 2016 Nov 1;90(21):9683-92. DOI: 10.1128/JVI.01347-16

27. Igarashi K, Kashiwagi K. Polyamines: mysterious modulators of cellular
functions. Biochemical and biophysical research communications. 2000 May
19;271(3):559-64.

28. Mounce BC, Cesaro T, Vlajnić L, Vidiņa A, Vallet T, Weger-Lucarelli J, Passoni
G, Stapleford KA, Levraud JP, Vignuzzi M. Chikungunya virus overcomes
polyamine depletion by mutation of nsP1 and the opal stop codon to confer
enhanced replication and fitness. Journal of virology. 2017 Aug 1;91(15):10-128.
DOI: 10.1128/JVI.00344-17

29. Doyle KJ. Delayed, reversible hearing loss caused by difluoromethylornithine
(DFMO). The Laryngoscope. 2001 May;111(5):781-5. DOI: 10.1097/00005537-
200105000-00005

30. Alhonen-Hongisto L, Levin VA, Marton LJ. Modification of uptake and
antiproliferative effect of methylglyoxal bis (guanylhydrazone) by treatment with
α-difluoromethylornithine in rodent cell lines with different sensitivities to
methylglyoxal bis (guanylhydrazone). Cancer research. 1985 Feb 1;45(2):509-
14.

31. Abdelraheem E, Thair B, Varela RF, Jockmann E, Popadić D, Hailes HC, Ward
JM, Iribarren AM, Lewkowicz ES, Andexer JN, Hagedoorn PL.
Methyltransferases: functions and applications. ChemBioChem. 2022 Sep
16;23(18):e202200212. DOI: 10.1002/cbic.202200212

32. Zhu BT, Wang P, Nagai M, Wen Y, Bai HW. Inhibition of human catechol-O-
methyltransferase (COMT)-mediated O-methylation of catechol estrogens by
major polyphenolic components present in coffee. The Journal of steroid
biochemistry and molecular biology. 2009 Jan 1;113(1-2):65-74. DOI:
10.1016/j.jsbmb.2008.11.011

33. Egloff MP, Benarroch D, Selisko B, Romette JL, Canard B. An RNA cap
(nucleoside-2′-O-)-methyltransferase in the flavivirus RNA polymerase NS5:
crystal structure and functional characterization. The EMBO journal. 2002 Jun
3;21(11):2757-68. DOI: 10.1093/emboj/21.11.2757

34. Rao ST, Rossmann MG. Comparison of super-secondary structures in proteins.
Journal of molecular biology. 1973 May 15;76(2):241-56. DOI: 10.1016/0022-
2836(73)90388-4

35. Chouhan BP, Maimaiti S, Gade M, Laurino P. Rossmann-fold
methyltransferases: taking a "β-Turn" around their cofactor, S-
adenosylmethionine. Biochemistry. 2018 Nov 8;58(3):166-70. DOI:
10.1021/acs.biochem.8b00994

36. Wang B, Tan XF, Thurmond S, Zhang ZM, Lin A, Hai R, Song J. The structure
of Zika virus NS5 reveals a conserved domain conformation. Nature
communications. 2017 Mar 27;8(1):14763. DOI: 10.1038/ncomms14763

37. Ahola T, Merits A. Functions of chikungunya virus nonstructural proteins.
Chikungunya virus: advances in biology, pathogenesis, and treatment. 2016:75-
98. DOI: 10.1007/978-3-319-42958-8_6

38. Dzobo K. The role of natural products as sources of therapeutic agents for
innovative drug discovery. Comprehensive pharmacology. 2022:408. DOI:
10.1016/B978-0-12-820472-6.00041-4

39. Cheng T, Li Q, Zhou Z, Wang Y, Bryant SH. Structure-based virtual screening
for drug discovery: a problem-centric review. The AAPS journal. 2012
Mar;14:133-41. DOI: 10.1208/s12248-012-9322-0

40. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam
H, Remmert M, Söding J, Thompson JD. Fast, scalable generation of high-quality
protein multiple sequence alignments using Clustal Omega. Molecular systems
biology. 2011 Oct 11;7(1):539. DOI: 10.1038/msb.2011.75

41. Dallakyan S, Olson AJ. Small-molecule library screening by docking with PyRx.
Chemical biology: methods and protocols. 2015:243-50. DOI: 10.1007/978-1-
4939-2269-7_19

42. Banck M, James CA, Morley C, Vandermeersch T, Hutchison GR. Open Babel:
An open chemical toolbox. J. Cheminf. 2011;3(1):33. DOI: 10.1186/1758-2946-
3-33

43. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer
FT, de Beer TA, Rempfer C, Bordoli L, Lepore R. SWISS-MODEL: homology
modelling of protein structures and complexes. Nucleic acids research. 2018 Jul
2;46(W1):W296-303. DOI: 10.1093/nar/gky427

44. Trott O, Olson AJ. AutoDock Vina: improving the speed and accuracy of docking
with a new scoring function, efficient optimization, and multithreading. Journal
of computational chemistry. 2010 Jan 30;31(2):455-61. DOI: 10.1002/jcc.21334

45. The PyMOL Molecular Graphics System, Version 1.2r3pre, Schrödinger, LLC.

46. Laskowski RA, Swindells MB. LigPlot+: multiple ligand–protein interaction
diagrams for drug discovery. DOI: 10.1021/ci200227u

47. Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJ.
GROMACS: fast, flexible, and free. Journal of computational chemistry. 2005
Dec;26(16):1701-18. DOI: 10.1002/jcc.20291.

48.   Huang J, MacKerell Jr AD. CHARMM36 all-atom additive protein force field:
Validation based on comparison to NMR data. Journal of computational
chemistry. 2013 Sep 30;34(25):2135-45. DOI: 10.1002/jcc.23354.

49.   Vanommeslaeghe K, Hatcher E, Acharya C, Kundu S, Zhong S, Shim J, Darian
E, Guvench O, Lopes P, Vorobyov I, Mackerell Jr AD. CHARMM general force
field: A force field for drug-like molecules compatible with the CHARMM all-
atom additive biological force fields. Journal of computational chemistry. 2010
Mar;31(4):671-90. DOI: 10.1002/jcc.21367

50.   Berendsen HJ, Postma JV, Van Gunsteren WF, DiNola AR, Haak JR. Molecular
dynamics with coupling to an external bath. The Journal of chemical physics.
1984 Oct 15;81(8):3684-90. DOI: https://doi.org/10.1063/1.448118.

51.   Hess B, Bekker H, Berendsen HJ, Fraaije JG. LINCS: a linear constraint solver
for molecular simulations. Journal of computational chemistry. 1997
Sep;18(12):146372.DOI:https://doi.org/10.1002/(SICI)1096987X(199709)18:12
<1463::AID-JCC4>3.0.CO;2-H

52.   Grubmüller H, Heller H, Windemuth A, Schulten K. Generalized Verlet
algorithm for efficient molecular dynamics simulations with long-range
interactions. Molecular Simulation. 1991 Mar 1;6(1-3):121 42.

DOI:https://doi.org/10.1080/08927029108022142

53.   Boonyasuppayakorn S, Padmanabhan R. Construction of plasmid, bacterial
expression, purification, and assay of dengue virus type 2 NS5 methyltransferase.
Dengue: Methods and Protocols. 2014:361-73. DOI: 10.1007/978-1-4939-0348-
1_22

54.   Kaur R, Mudgal R, Narwal M, Tomar S. Development of an ELISA assay for
screening inhibitors against divalent metal ion dependent alphavirus capping
enzyme. Virus research. 2018 Sep 2;256:209-18. DOI:
10.1016/j.virusres.2018.06.013

55.   Bhutkar M, Saha A, Tomar S. Viral methyltransferase inhibitors: berbamine,
venetoclax, and ponatinib as efficacious antivirals against Chikungunya virus.
Archives of Biochemistry and Biophysics, Volume 759,September 2024, 110111.
2024:2024-05. DOI: 10.1016/j.abb.2024.110111

56.   Benmansour F, Trist I, Coutard B, Decroly E, Querat G, Brancale A, Barral K.
Discovery of novel dengue virus NS5 methyltransferase non-nucleoside
inhibitors by fragment-based drug design. European journal of medicinal
chemistry. 2017 Jan 5;125:865-80. DOI: 10.1016/j.ejmech.2016.10.007

57.   Coutard B, Decroly E, Li C, Sharff A, Lescar J, Bricogne G, Barral K. Assessment
of Dengue virus helicase and methyltransferase as targets for fragment-based
drug discovery. Antiviral research. 2014 Jun 1;106:61-70. DOI:
10.1016/j.antiviral.2014.03.013

58.   Vagin A, Teplyakov A. MOLREP: an automated program for molecular
replacement. Applied Crystallography. 1997 Dec 1;30(6):1022-5.
DOI:https://doi.org/10.1107/S0021889897006766

59.   Murshudov GN, Skubák P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA,
829   Winn MD, Long F, Vagin AA. REFMAC5 for the refinement of macromolecular
830   crystal structures. Acta Crystallographica Section D: Biological Crystallography.
831   2011 Apr 1;67(4):355-67. DOI: 10.1107/S0907444911001314

832   60.   Liebschner D, Afonine PV, Moriarty NW, Poon BK, Sobolev OV, Terwilliger
833   TC, Adams PD. Polder maps: improving OMIT maps by excluding bulk solvent.
834   Acta Crystallographica Section D: Structural Biology. 2017 Feb 1;73(2):148-57.
835   DOI: 10.1107/S2059798316018210

836   61.   Singh H, Mudgal R, Narwal M, Kaur R, Singh VA, Malik A, Chaudhary M,
837   Tomar S. Chikungunya virus inhibition by peptidomimetic inhibitors targeting
838   virus-specific cysteine protease. Biochimie. 2018 Jun 1;149:51-61. DOI:
839   10.1016/j.biochi.2018.04.004

840   62.   Mishra AC, Arankalle VA, Gadhave SA, Mahadik PH, Shrivastava S, Bhutkar
841   M, Vaidya VM. Stratified sero-prevalence revealed overall high disease burden
842   of dengue but suboptimal immunity in younger age groups in Pune, India. PLoS
843   neglected tropical diseases. 2018 Aug 6;12(8):e0006657. DOI:
844   10.1371/journal.pntd.0006657

845   63.   Chien LJ, Liao TL, Shu PY, Huang JH, Gubler DJ, Chang GJ. Development of
846   real-time reverse transcriptase PCR assays to detect and serotype dengue viruses.
847   Journal of clinical microbiology. 2006 Apr;44(4):1295-304. DOI:
848   10.1128/JCM.44.4.1295-1304.2006 DOI: 10.1016/j.virol.2018.10.009

849   64.   Kaur R, Mudgal R, Jose J, Kumar P, Tomar S. Glycan-dependent chikungunya
850   viral infection divulged by antiviral activity of NAG specific chi-like lectin.
851   Virology. 2019 Jan 2;526:91-8. DOI: 10.1016/j.virol.2018.10.009

852   65.   Madhubala R. Thin-layer chromatographic method for assaying polyamines.
853   Polyamine protocols. 1998:131-6. DOI: 10.1385/0-89603-448-8:131

854   66.   Zhao B, Yi G, Du F, Chuang YC, Vaughan RC, Sankaran B, Kao CC, Li P.
855   Structure and function of the Zika virus full-length NS5 protein. Nature
856   communications. 2017 Mar 27;8(1):14762. DOI: 10.1038/ncomms14762

857   67.   Zhao Y, Soh TS, Zheng J, Chan KW, Phoo WW, Lee CC, Tay MY, Swaminathan
858   K, Cornvik TC, Lim SP, Shi PY. A crystal structure of the Dengue virus NS5
859   protein reveals a novel inter-domain interface essential for protein flexibility and
860   virus replication. PLoS pathogens. 2015 Mar 16;11(3):e1004682. DOI:
861   10.1371/journal.ppat.1004682

862   68.   Røsæg MV, Thorarinsson R, Aunsmo A. Effect of vaccines against pancreas
863   disease in farmed Atlantic salmon. Journal of fish diseases. 2021
864   Dec;44(12):1911-24. DOI: 10.1111/jfd.13505

865   69.   Nasar F, Palacios G, Gorchakov RV, Guzman H, Da Rosa AP, Savji N, Popov
866   VL, Sherman MB, Lipkin WI, Tesh RB, Weaver SC. Eilat virus, a unique
867   alphavirus with host range restricted to insects by RNA replication. Proceedings
868   of the National Academy of Sciences. 2012 Sep 4;109(36):14622-7. DOI:
869   10.1073/pnas.1204787109

70. Hobson-Peters J, Yam AW, Lu JW, Setoh YX, May FJ, Kurucz N, Walsh S, Prow
NA, Davis SS, Weir R, Melville L. A new insect-specific flavivirus from northern
Australia suppresses replication of West Nile virus and Murray Valley
encephalitis virus in co-infected mosquito cells. PloS one. 2013 Feb
27;8(2):e56534. DOI: 10.1371/journal.pone.0056534

71. Parry R, Asgari S. Discovery of novel crustacean and cephalopod flaviviruses:
insights into the evolution and circulation of flaviviruses between marine
invertebrate and vertebrate hosts. Journal of Virology. 2019 Jul 15;93(14):10-
128. DOI: 10.1128/JVI.00432-19

72. Chen Y, Barkley MD. Toward understanding tryptophan fluorescence in proteins.
Biochemistry. 1998 Jul 14;37(28):9976-82. DOI: 10.1021/bi980274n

73. Yap LJ, Luo D, Chung KY, Lim SP, Bodenreider C, Noble C, Shi PY, Lescar J.
Crystal structure of the dengue virus methyltransferase bound to a 5′-capped
octameric   RNA.   PLoS   One.   2010   Sep   17;5(9):e12836.   DOI:
10.1371/journal.pone.0012836

74. Feracci M, Eydoux C, Fattorini V, Bello LL, Gauffre P, Selisko B, Sutto-Ortiz P,
Shannon A, Xia H, Shi PY, Noel M. AT-752 targets multiple sites and activities
on the Dengue virus replication enzyme NS5. Antiviral research. 2023 Apr
1;212:105574. DOI: 10.1016/j.antiviral.2023.105574

75. Benarroch D, Egloff MP, Mulard L, Guerreiro C, Romette JL, Canard B. A
structural basis for the inhibition of the NS5 dengue virus mRNA 2′-O-
methyltransferase domain by ribavirin 5′-triphosphate. Journal of Biological
Chemistry. 2004 Aug 20;279(34):35638-43. DOI: 10.1074/jbc.M400460200

76. Galmarini CM, Mackey JR, Dumontet C. Nucleoside analogues: mechanisms of
drug resistance and reversal strategies. Leukemia. 2001 Jun;15(6):875-90. DOI:
10.1038/sj.leu.2402114

77. Bhutkar M, Singh V, Dhaka P, Tomar S. Virus-host protein-protein interactions
as molecular drug targets for arboviral infections. Frontiers in Virology. 2022
Aug 4;2:959586.DOI: https://doi.org/10.3389/fviro.2022.959586

78. Pegg AE, Casero RA. Current status of the polyamine research field. Polyamines:
Methods and Protocols. 2011:3-5. DOI: 10.1007/978-1-61779-034-8_1

79. Rao JN, Rathor N, Zhuang R, Zou T, Liu L, Xiao L, Turner DJ, Wang JY.
Polyamines regulate intestinal epithelial restitution through TRPC1-mediated
Ca2+ signaling by differentially modulating STIM1 and STIM2. American
Journal of Physiology-Cell Physiology. 2012 Aug 1;303(3):C308-17. DOI:
10.1152/ajpcell.00120.2012

80. Terui Y, Yoshida T, Sakamoto A, Saito D, Oshima T, Kawazoe M, Yokoyama S,
Igarashi K, Kashiwagi K. Polyamines protect nucleic acids against depurination.
The international journal of biochemistry & cell biology. 2018 Jun 1;99:147-53.
DOI: 10.1016/j.biocel.2018.04.008

81. Bouteille B, Dumas M. Human African Trypanosomiasis. Encyclopedia of the
Neurological Sciences . 2003 Jan 1;587–94. DOI:https://doi.org/10.1016/B0-12-
226870-9/00431-7

82. Nazir A, Nazir A, Kandel K. Advancing neuroblastoma care: future horizons after
approval of eflornithine by FDA. International Journal of Surgery. 2024 May
1;110(5):2511-2. DOI: 10.1097/JS9.0000000000001182

83. LoGiudice N, Le L, Abuan I, Leizorek Y, Roberts SC. Alpha-
difluoromethylornithine, an irreversible inhibitor of polyamine biosynthesis, as a
therapeutic strategy against hyperproliferative and infectious diseases. Medical
Sciences. 2018 Feb 8;6(1):12. DOI: 10.3390/medsci6010012

84. Miller-Fleming L, Olin-Sandoval V, Campbell K, Ralser M. Remaining
mysteries of molecular biology: the role of polyamines in the cell. Journal of
molecular biology. 2015 Oct 23;427(21):3389-406. DOI:
10.1016/j.jmb.2015.06.020

85. Zai SZ, Guo ZX, Prystowsky JH, Grunberger D. Caffeic acid phenethyl ester
inhibits proliferation of human keratinocytes and interferes with the EGF
regulation of ornithine decarboxylase. Oncology research. 1995;9(7):445-52.

86. Kim DJ, Roh E, Lee MH, Oi N, Lim DY, Kim MO, Cho YY, Pugliese A, Shim
JH, Chen H, Cho EJ. Herbacetin is a novel allosteric inhibitor of ornithine
decarboxylase with antitumor activity. Cancer research. 2016 Mar 1;76(5):1146-
57. DOI: 10.1158/0008-5472.CAN-15-0442

87. Rani R, Bhutkar M, Tomar S. Therapeutic Antiviral Potential of Flavonoids.
InFlavonoids as Nutraceuticals 2024 Apr 9 (pp. 57-101). Apple Academic Press.

88. Erdemli HK, Akyol S, Armutcu F, Akyol O. Antiviral properties of caffeic acid
phenethyl ester and its potential application. Journal of intercultural
ethnopharmacology. 2015 Oct;4(4):344. DOI: 10.5455/jice.20151012013034

89. Brecher M, Chen H, Li Z, Banavali NK, Jones SA, Zhang J, Kramer LD, Li H.
Identification and characterization of novel broad-spectrum inhibitors of the
flavivirus methyltransferase. ACS infectious diseases. 2015 Aug 14;1(8):340-9.
DOI: 10.1021/acsinfecdis.5b00070

90. Lim SP, Sonntag LS, Noble C, Nilar SH, Ng RH, Zou G, Monaghan P, Chung
KY, Dong H, Liu B, Bodenreider C. Small molecule inhibitors that selectively
block dengue virus methyltransferase. Journal of Biological Chemistry. 2011 Feb
25;286(8):6233-40. DOI: 10.1074/jbc.M110.179184

91. Noble CG, Li SH, Dong H, Chew SH, Shi PY. Crystal structure of dengue virus
methyltransferase without S-adenosyl-L-methionine. Antiviral research. 2014
Nov 1;111:78-81. DOI: 10.1016/j.antiviral.2014.09.003

92. Jain R, Butler KV, Coloma J, Jin J, Aggarwal AK. Development of a S-
adenosylmethionine analog that intrudes the RNA-cap binding site of Zika
methyltransferase. Scientific Reports. 2017 May 9;7(1):1632. DOI:
10.1038/s41598-017-01756-7

Figure 1: *Three-dimensional representation of docked ligands in the enzyme active site (teal surface) of CHIKV nsP1 and DENV 3 NS5 MTase (wheat surface). (A-D) CHIKV nsP1 interacts with the GTP (A), CAPE (B) HC (C), SAM (D), and (E-G) DENV 3 NS5 MTase interact with GTP (E), CAPE (F) HC (G). Zoomed in figures show a detailed view of the binding pocket where molecular interactions of ligands (green) with active site residues (teal*

Figure 2*: Two-dimensional representation view of docked ligands in the enzyme active site of*

CHIKV nsP1 and DENV 3 NS5 MTase. (A-D) CHIKV nsP1 interacts with the GTP (A), CAPE

(B) HC (C), SAM (D), and E-G) DENV 3 NS5 MTase interact with GTP (E), CAPE (F) HC

(G). H-bonds are shown in green dashed lines with the distance shown in Å. Additional residues

forming hydrophobic interactions are indicated by a brown semicircle with radiating spokes

towards the ligands. 2D interaction figures are made using LigPlot+ software

Figure 3: *Determination of protein interactions with ligand/substrate by TFS. nsP1 - (A) GTP*

(B) SAM (C) HC (D) CAPE. NS5 MTase - (E) GTP (F) SAM (G) HC (H) CAPE. Data from

three independent experiments were collected and analyzed using nonlinear regression with

the 'One Site-Specific Binding' model.

Figure 4: *MTase inhibition activity assay. SDS/PAGE analysis of purified recombinant CHIKV*

nsP1 (A) and DENV 3 NS5 MTase (D). CE-based nsP1 MTase activity inhibition assay (B)

HC (C) CAPE and DENV 3 NS5 MTase (E) HC (F) CAPE. Error bars indicate the standard

error derived from three independent experiments.

Figure 5: *Crystal Structure of DENV 3 NS5 MTase in Complex with SAH and CAPE*

(PDB:8DKZ). A) Polder omit Map CAPE (302) and CAPE (303) at 3 and 2.5 σ. B) Surface

representation of DENV 3 NS5 MTase (wheat) with CAPE (green carbon atoms) and SAH (hot

pink carbon atoms) depicted as sticks. C) Structural superposition of a (PDB:8DKZ) green and

(PDB:4V0R) cyan. D) Structure of DENV 3 NS5-MTase bound to SAH and CAPE, with

hydrogen bond interacting residues represented as yellow dotted lines. E) Structure of the

DENV 3 NS5-MTase domain (PDB: 4V0R) bound to GTP, with hydrogen bond interacting

residues shown as yellow dotted lines. F) Structure of the DENV 3 NS5-MTase domain (PDB:

2XBM) bound to G3A, with hydrogen bond interacting residues shown as yellow dotted lines.

G) Structural superposition of a (PDB:8DKZ) green and (PDB:5DTO) orange. Here, Mg2+

is displayed as a green sphere H) Structure of DENV 3 NS5-MTase bound to A and G from cap

0 RNA, with hydrogen bond interacting residues represented as yellow dotted lines.

1001

1002

Figure 6: Evaluation of polyamine depletion and antiviral activity using virus titer reduction

profiling of HC and CAPE. (A) Mechanistic pathway illustrating the inhibitory effects of CAPE

and HC on polyamine pathway and relevant enzymes and inhibitor. (B)the chromatographic

analysis of polyamine levels in Vero cells after 36 h treatment with (from left) CAPE (25 μM),

HC (200 μM), DFMO(1000 μM), and cell control (CC), 0.1 μM putrescine (Put), spermine

(Spm), and spermidine (Spd) as a positive control marker. Treatment with both compounds

leads to decreased polyamine levels for both compounds compared to DFMO. Vero cells were

treated with HC and CAPE for 12 h and subsequently infected with CHIKV for 2 h. After this,

the cells were incubated for an additional 24 h. Following this, the supernatants were collected

for plaque assay. A similar compound treatment protocol was followed for DENV infection:

Vero cells were pre-treated with the compounds for 12 h, then infected with the virus for 2 h,

followed by a 24 h treatment period. After this, the compounds were removed, and the cells

were maintained in 2% DMEM and incubated for an additional 4 days. Supernatants were

then collected and subjected to plaque assay. DFMO at a concentration of 1000 μM served as

a positive control, with a virus control (VC) included for comparison. (C, D, F,G) illustrate the

inhibitory effects of various concentrations of HC and CAPE on CHIKV and DENV-infected

cells, as assessed by plaque assay. E) RT-PCR for CHIKV with HC 200 μM, CAPE 25 μM,

DFMO 1000 μM concentration H) RT-PCR for DENV with HC 200 uM, CAPE 2.5 uM and

DFMO 1000 μM. Values are the means, and error bars represent the standard deviation from

three independent experiments. Statistical analysis was performed using one-way ANOVA with

*Dunnett's post-test. ****P <0.0001.*

1024

1025

1026 ***Figure 7:*** *Evaluation of the antiviral effect by immunofluorescence assay (IFA).*

1027 *Immunofluorescent staining of HC (200 µM) and CAPE (25 µM for CHIKV and 2.5 µM for*

1028 *DENV) treated with CHIKV(A) and DENV(B) in Vero cells. Green fluorescence indicates the*

1029 *virus load, and blue fluorescence indicates the nuclear staining with DAPI with a 10 X*

1030 *objective lens. The scale bar is 400 µm.*

1031

1032

Figure 8: *Effect of addition of biogenic polyamines on virus infected and compound treated*

cells. Vero cells were pre-treated with drugs for 24 h at various concentrations, following this

cells were infected with CHIKV (A) and DENV (B) treated with HC (200 μM), CAPE (25 μM

for CHIKV and 2.5 μM for DENV), DFMO (1000 μM) as a positive control and VC as

untreated control. Minus (-) sign indicates only the compound-treated Vero cells, whereas plus

(+)sign indicates that 1μM polyamines (put, spd, spm) were added additionally in the

compound-treated cells. Error bars represent the standard error of three independent

experiments.

1041

1042

1043

Supplementary Experimental Procedures

Cell line, Virus isolation, propagation and serotyping

Vero cell line isolated from kidney epithelial cells of an African green monkey was obtained

from the National Centre for Cell Science (NCCS), Pune, India. Vero cells were used for the

propagation and titration of DENV and CHIKV. The cells were maintained in Dulbecco's

Modified Eagles Medium (DMEM) (Gibco, HiMedia) supplemented with 10% inactivated

fetal bovine serum (FBS) (Gibco, HiMedia) along with 100 U/mL penicillin and 100 mg/mL

streptomycin (Gibco, HiMedia). The cells were maintained at 37°C with 5% CO_2

supplementation.

Dengue was isolated from the DENV-suspected patient's blood samples. Sera was

aliquoted and tested for the presence of DENV NS1 antigen by J Mitra ELISA kit according to

the manufacturer's instructions. DENV-positive samples were used for virus isolation. Initially,

sera were 1:10 diluted in minimum essential medium (MEM) (Gibco, HiMedia) containing 2

% FBS, 100 U/mL penicillin, and 100 mg/mL streptomycin for virus propagation. Then, Serum

samples were added to a confluent monolayer of Vero cells for 90 min with gentle shaking

every 15 min in a 37 °C CO_2 incubator. The inoculum was withdrawn, and the cells were

cultured for 5-6 days in fresh MEM with 2% FBS and antibiotics until the appearance of

cytopathic effect (CPE). After 5-6 days, the supernatant was harvested and serially diluted for

infection of Vero monolayer cells for 90 min and overlaid with MEM containing 1 % agarose

for the plaque purification procedure. Single plaques were picked and resuspended in MEM

and were further propagated in Vero cells. Then, DENV NS1 ELISA-positive supernatants

showing CPE was collected and used for further study.

To confirm the DENV and its serotype, RNA from the supernatant of cell culture

infected with the virus was isolated using the Trizol (Sigma) method described by the

manufacturer. The RNA was then reverse-transcribed into complementary DNA (cDNA) using

the PrimeScript cDNA Synthesis Kit (Takara). Synthesized cDNA was utilized as a template

for polymerase chain reaction (PCR) (63) and the amplification product was analyzed by gel

electrophoresis in 1 % agarose gels stained with ethidium bromide. The sample was sent for

DNA sequencing, and the serotype of DENV was confirmed by matching the obtained

sequence against the nucleotide database.

CHIKV (Accession No. KY057363.1.) was propagated and titrated in a Vero cell line

using the protocol reported by Singh et al., 2018 and then stored at −80 °C for further

experiments (61).

Cell Cytotoxicity Assay

Different concentrations of HC and CAPE were evaluated for cytotoxicity on Vero cells using

3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Himedia) assay.

Before treatment, Vero cells were seeded in a 96-well plate, and at 90% confluency, the media

were removed, and different dilutions of compounds were added into each well for 12 h. After

incubation, the inoculum was removed, and fresh maintenance media were added for 2 h.

Further fresh compound dilutions were added to each well and kept for 24 h. Subsequently, 20

μL/well of MTT (5 mg/mL) was added and incubated for 4 h at 37 °C in 5% CO_2. Upon

incubation, 110 μL/well of DMSO was added to dissolve formazan crystals. Plates were read

at a wavelength of 570 nm using cytation 3 multi-mode plate reader (BioTek Instruments, Inc.).

The average absorbance of 0.1 % DMSO-treated cells was used as cell control. In the second

set of experiments, after completion of 24 h post-treatment, fresh maintenance media were

added and kept for 4 days. The cell viability of the treated well was compared with the cell

control, and the concentration that showed >90% viability after compound treatment was

considered non-toxic.

Plaque assay

Vero cells were seeded at 1.0 x 10^5 cells/well in a 24-well plate in complete media before

infection. The supernatant was 10-fold serially diluted in maintenance media and was

inoculated on ~80-90% confluent cells. Plates were incubated for 2 h with gentle shaking every

15 min for virus adsorption at 37 °C with a 5% CO_2 incubator. After adsorption, overlay media

and maintenance media were added in 1:1 dilution and plates were further incubated for 2 days

and 3 days for CHIKV and DENV, respectively at 37 ˚C in 5% CO_2 incubator. Following that,

cells were stained with 1% crystal violet to count the number of CHIKV plaques and

immunostaining for DENV-infected cells, as described below.

Immunostaining was performed at room temperature. After incubation, overlay media was

removed from the plates, and the cell monolayer was fixed with 3.7% formaldehyde solution

for 30 min. Cells were washed 3 times with PBST (0.02% Tween-20 in Phosphate buffer

saline). After that, cells were permeabilized with 0.2% triton X-100 in PBS for 7 min. Cells

were washed three times with PBST and incubated with 3% Skim milk for 30 min. Then, cells

were washed three times with PBST and incubated with 1:500 diluted orthoflavivirus group

antibody (Genetex) for 2 h. Cells were washed three times with PBST and incubated with

secondary antibody (Goat anti-mouse IgG HRP, 1:1500 dilution) for 1 h. After washing 2 times

with PBST and three times with PBS, cells were stained with True Blue Peroxidase substrate

(KPL, Sera Care, MA, USA) and incubated in the dark for 30 min to develop blue color staining

of virus-infected cells, and foci were counted (62).

Multiple Sequence Alignment (MSA)

The amino acid sequence of nsP1 protein of *alphaviruses* was compared with CHIKV nsP1 as

a reference point using Clustal Omega (40) to check if the key residues involved in the capping

of viral RNA were conserved across different viruses. The MSA was performed

for Venezuelan equine encephalitis virus (VEEV Gene Bank Id: AAU89533.1), (CHIKV Gene

Bank Id: QOW97289.1), Ross River virus (RRV Gene Bank Id: QTC33397.1), Sindbis virus

(SINV Gene Bank Id: AWT57896.1), Aura Virus (AURA Gene Bank Id: AWQ38330.1),

Middelburg virus (MIDV Gene Bank Id: QOY44469.1), Barmah Forest virus (BFV Gene Bank

Id: QVM79755.1), Madariaga virus (MADV Gene Bank Id: ABL84688), Salmonid alphavirus

(SAV Gene Bank Id: KC122922.1), Eilat virus (EILV Gene Bank Id: NC_018615) and Mayaro

virus (MAYV Gene Bank Id: AZM66145) from the alphaviruses. The sequence alignment

profile of the selected nsP1 sequences was performed via Clustal Omega tool and analyzed by

a graphical coloured depiction using ESPript 3.0 (40).

Similarly, MSA was performed for the MTase domain of NS5 from dengue 3 Virus (DENV 3

PDB Id:4CTJ), ZIKV PDB Id:5WZ2, West Nile Virus (WNV PDB Id:2OY0), Yellow Fever

Virus (YFV PDB Id:3EVA), Palm Creek virus (PCV Gene Bank Id: NC_033694.1), Wenzhou

shark flavivirus (WSF Gene Bank Id: AVM87250.1) and Japanese Encephalitis Virus (JEV

Gene Bank Id:4K6M_1) from the orthoflaviviruses and the analysis was carried out in the same

way as for DENV.

1131

1132   **Supplementary Figures**

1133

1134

***Supplementary Figure 1:** Multiple Sequence alignment of MTase domains of different*

alphaviruses and orthoflaviviruses. A) Sequence alignment of nsP1 MTase domain (residues 1

to 285) from CHIKV with the MTase domains of other alphaviruses. VEEV Gene Bank Id:

AAU89533.1, CHIKV Gene Bank Id: QOW97289.1, RRV Gene Bank Id: QTC33397.1 ,SINV

Gene Bank Id: AWT57896.1, AURA Gene Bank Id: AWQ38330.1, MIDV Gene Bank Id:

QOY44469.1, BFV Gene Bank Id: QVM79755.1, MADV Gene Bank Id: ABL84688 , SAV Gene

Bank Id: KC122922.1, EILV Gene Bank Id: NC_018615   and MAYV Gene Bank Id:

AZM66145. B) Sequence alignment of NS5 MTase domain (residues 1 to 264) from DENV with

MTase domains of other orthoflaviviruses. DENV 3 PDB Id:4CTJ, ZIKV PDB Id:5WZ2 , WNV

PDB Id:2OY0, YFV PDB Id:3EVA, PCV Gene Bank Id: NC_033694.1, WSF Gene Bank Id:

AVM87250.1 and JEV PDB Id:4K6M. The active-site MTase residues in both alphaviruses

and orthoflaviviruses are highly conserved, suggesting that the structure and function of the

MTase domain in both families are conserved. Residues are represented with single letter

amino acid code, with identical residues indicated in white font and boxed in red, while similar

residues are indicated in red font. Blue triangles correspond to the MTase active site residues

as per the literature. Alignments were generated via Clustal Omega and ESPript tools.

1151

1152 ***Supplementary Figure 2:*** *Superposition of active site residues of SWISS generated models*

1153 *with templates. (A-B) Close-up view of overlapped residues of generated models of CHIKV*

1154 *nsP1 (A) and DENV 3 NS5 MTase (B) with RMSD values of 0.295 Å and 0.182 Å, respectively.*

1155 *The generated model is labelled yellow, and the template is labelled pink.*

1156

1157

1158 **Supplementary Figure 3:** *Root Mean Square Deviations (RMSD) graphs of (A) CHIKV nsP1*

1159 *Apo i.e., native protein, nsP1 - SAM or CAPE or HC (B) DENV 3 NS5 MTase Apo i.e., native,*

1160 *NS5 MTase – GTP or HC or CAPE*

1161

1162

1163 **Supplementary Figure 4:** *Intrinsic fluorescence intensity change in protein-compound*

1164 *interactions by TFS. nsP1 - (A)GTP (B) SAM (C) HC (D) CAPE. NS5 MTase – (E)GTP (D)*

1165 *SAM (G) HC (H) CAPE .*

1166

1167

Supplementary Figure 5*: (A) and (B) depict the percent cell viability of Vero cells treated with*

compounds for 12 h pre-treatment followed by a 2 h incubation with 2% DMEM, and 24 h

post-treatment. (C) and (D) show the percent cell viability of Vero cells treated with compounds

for 12 h pre-treatment, followed by a 2 h incubation with maintenance media and 24 h post-

treatment, and incubated in maintenance media for 4 days. Values are the means, and error

bars represent the standard deviation from three independent experiments.The 50% cytotoxic

concentration was determined based on linear dose-response analysis using GraphPad Prism

8 software. Compounds with concentrations that maintained cell viability above 90% were

selected for subsequent cell culture experiments.

1177

Supplementary Figure 6: *Evaluation of antiviral activity through virus titer reduction*

profiling: (A) HC and (B) CAPE against CHIKV; (C) HC and (D) CAPE against DENV.For

this, percent inhibition was calculated using the formula: Percent Inhibition=100× ((PFU mL-

1 VC−PFU mL-1 test concentration)/PFU mL-1VC), where VC represents virus control

values. Data were plotted against concentration with a non-linear regression curve fit using

GraphPad Prism 8.0.

Table 1: Detailed molecular interactions between the CHIKV nsP1 and ligands.

Ligand	Binding energy (kcal/mol)	Interactions			
		Hydrogen			Hydrophobic
		Residues	Molecules	Bond length (Å)	
GTP	-8.3	Arg41	O3-NH2	3.18	

Ligand					Hydrophobic
			O3-NH2	3.13	
			O10-O	3.18	
		Arg70	O8-NH1	2.81	
			O2-NH1	3.2	His45, Ser44, Ala40, Asp152, Tyr285
		Arg71	O9-NH2	3.12	
		Arg92	O12-NH2	2.9	
		Gln151	O9-OE1	3.2	
		Tyr248	O14-OH	2.8	
SAM	-6.7	Gly65	O5-O	3.05	
		Asp89	O5-OD1	3.24	
			O5-OD2	2.72	
		Pro249	N5-O	2.95	
			N3-O	3.04	Ser86, Asp152, Glu250, Arg85, Ala155, Glu88
		Tyr248	N6-OH	2.95	
		Arg70	O3-NH2	3.03	
			O3-NH1	3.05	
			O4-NH1	3.20	
		Arg92	O3-NH2	3.23	
HC	-8.0	Arg92	O4-NH2	2.92	Arg41, Tyr285,Asp152, Phe178, Val243, Ala40, Tyr248
		Arg70	O2-NH1	3.0	
		Glu88	O3-OE1	2.73	
CAPE	-7.6	Glu250	O4-OE1	2.85	Tyr248, Arg70, Asp152, Tyr154, Ala40, Phe178, Val243, Tyr285
		Arg92	O1-NH2	2.89	

1185

1186

1187 **Table 2:** Detailed molecular interactions between the DENV 3 NS5 MTase and ligands.

Ligand		Interactions	
		Hydrogen	**Hydrophobic**

	Binding energy (kcal/mol)	Residues	Molecules	Bond length (Å)	
GTP	-7.1	Asn15	O14-OD1	2.84	Leu14, Phe22, Thr211
		Arg19	O9-NH1	3.11	
		Lys26	O9-NZ	2.97	
		Ser147	OG-O3	2.97	
			OG-O11	3.03	
		Arg208	O6-NH1	3.01	
		Ser210	O11-O	2.73	
			O12-O	3.00	
HC	-6.7	Lys11	O2-NZ	3.04	Leu14, Phe22, Pro149, Thr211
			O4-NZ	3.05	
		Asn15	O3-O	2.91	
			O4-OD1	2.79	
		Ser147	O6-OG	2.74	
		Ser210	O6-O	2.91	
CAPE	-6.6	Lys11	O1-NZ	3.07	Asn15, Ser18, Arg19, Phe22, Ser148, Pro149, Ser210
		Leu14	O3-O	2.94	
		Leu17	O3-O	2.70	
			O4-O	2.83	

1188

Resolution range	**23.06–2.60 (2.72- 2.60)**
Space group	$P\,2\,2_1\,2_1$
Unit cell dimensions: *a b c* (Å) α, β, γ (°)	**51.6 60.7 184.3**
Completeness (%)	**99.6 (99.3)**
Rmerge [a]	**0.16 (0.37)**
I/σ(I)	**7.3 (3.3)**
CC(1/2)	**0.97 (0.77)**
Refinement	
Reflections used in refinement	**10011**
Reflections used for R-free	**924**
R-work[b]	**0.231**
R-free[b]	**0.273**
Wilson B-factor (Å)	**28.9**
Number of non-hydrogen atoms	**4302**
Macromolecules	**4086**
Ligands	**94**
Solvent	**124**
Protein residues	**508**
RMS (bonds) (Å)[c]	**0.01**
RMS (angles) (°) [c]	**1.89**
Ramachandran Plot	
Favored (%)	**97.83**
Allowed (%)	**1.38**
Outliers (%)	**0.79**
Average B-factor (Å)	**28.0**
macromolecules (Å)	**28.65**
ligands (Å)	**48.98**
solvent (Å)	**18.44**

1189

Table 3:- Data collection, processing and refinement statistics of DENV 3 NS5 MTase
complexed with soaked CAPE (PDB:8DKZ). Values in parentheses are for the highest-
resolution shell.

a Rmerge = $\Sigma| I - \langle I \rangle |/\Sigma I$.

b R = Σ|Fobs| - |Fcalc|/Σ|Fobs|. The Rfree is the R calculated on the 5% reflections excluded
for refinement.

c RMS is root mean square.

Table 4:- Detailed molecular interactions between the DENV 3 NS5 MTase and CAPE
(PDB:8DKZ).

Ligand	Interactions			
	Hydrogen			Hydrophobic
	Residues	Molecules	Bond length (Å)	
CAPE (302)	Arg211	O4-NH2	3.5	Lys14, Leu17, Asn18, Phe25, Ser150, Ser151, Pro152, Ser213, Thr214, CAPE 303
CAPE (303)	-			Gly148, Glu149, Ser150, Lys180, CAPE 302

1199

1200

1201 **Table 5:-** Detailed molecular interactions between the DENV 3 NS5 MTase domain and
1202 GTP (PDB:4V0R).

1203

Ligand	Interactions			
	Hydrogen			Hydrophobic
	Residues	Molecules	Bond length (Å)	
GTP	Lys14	O2'-NZ	2.8	Asn18, Phe25, Lys29, Ser150, Pro152
		O3'-NZ	2.9	
	Leu17	N2-O	2.8	
	Asn18	N2-O	2.9	
		O2'-OD1	3.0	
	Leu20	N2-O	3.4	
	Arg22	O6-NH1	2.6	
	Ser151	O3'-O	3.3	
	Arg211	O1B-NH1	3.3	
	Ser213	O1B-OG	3.1	

1204

1205

1206

1207

1208

1209 **Table 6:-** Detailed molecular interactions between the DENV 3 NS5 MTase domain and
1210 G3A (PDB:2XBM).

| Ligand | Interactions | | | |
| | Hydrogen | | | Hydrophobic |
	Residues	Molecules	Bond length (Å)	
G3A	Asn 18	O-N11	3.26	Pro 152, Ser 213, Ser 151
	Leu 20	O-N32	2.81	
	Leu 17	O-N32	3.02	

1211

1212

1213 **Table 7:-** Detailed molecular interactions between the DENV 3 NS5 MTase domain and A-
1214 G (from cap 0 RNA) (PDB:5DTO).

| Ligand | Interactions | | | |
| | Hydrogen | | | Hydrophobic |
	Residues	Molecules	Bond length (Å)	
A-G	Ser150	OG-OP1	2.88	Leu 182, Arg84, Lys61, Gly148, Ser56, Glu 216
	Thr214	NH2-OP1	2.66	
	Lys180	NZ-O2	2.35	
	Glu111	OE2-N2	2.58	
	Arg57	OG1-OP2	3.25	

1215 These color-coded residues represent various interactions in a molecular context. Violet
1216 represents hydrogen bonds, green indicates hydrophobic interactions, and red signifies CAPE's
1217 hydrophobic residues that make hydrogen bonds with GTP, G3A, and A-G.

1218

1219 **Table 8:-** Summary of reported DENV MTase Inhibitors

PDB ID	Inhibitor	Viral MTase	Binding site	IC$_{50}$	Ref.
1R6A	Ribavirin	DENV 2	GTP binding site	~100 μM	(75)
8BCR	AT-9010	DENV 3	GTP binding site	~ 0.50 μM.	(74)
4CTK, 4CTJ	FRAGMENT 2A4,FRAGMENT 3A9	DENV 3	Novel sites	N.A.	(57)
5CUQ	NSC 12155	DENV 3	SAM binding site	~7.0 μM	(89)

3P8Z	Compound 10	DENV 3	SAM binding site	N.A.	(90)
5EHI, 5EIF,5EIW,5EKX, 5EC8,5EHG,5E9Q	287, NB2C2, NB3C2, NB2E11, BF175, BF341, BF174	DENV 3	Novel binding sites	N.A.	(56)
4R8S	Sinefungin	DENV 3	SAM binding site	N.A	(91)
5ULP	MS2042	ZIKV	SAM binding site	N.A.	(92)

1220

1221   N.A - Not available