Marine particle microbiomes during a spring diatom bloom contain active sulfate-reducing bacteria

Siebers, R.[1], Schultz, D.[1], Farza, M. S.[1], Brauer, A.[1], Zühlke, D.[1], Mücke, P. A[1], Wang, F.[2], Bernhardt, J.[1], Teeling, H.[2], Becher, D.[1], Riedel, K.[1,3], Kirstein, I. V.[4], Wiltshire, K. H.[4], Hoff, K.J.[5], Schweder, T.[3,6], Urich, T.[1,3], Bengtsson, M. M.[1,3,4*]

[1]Institute of Microbiology, University of Greifswald, Greifswald, Germany

[2]Max Planck Institute for Marine Microbiology, Bremen, Germany

[3]Institute of Marine Biotechnology, Greifswald, Germany

[4]Alfred-Wegener-Institute Helmholtz Centre for Polar and Marine Research, Biologische Anstalt Helgoland, Helgoland, Germany

[5]Institute of Mathematics and Computer Science, University of Greifswald, Greifswald, Germany

[6]Institute of Pharmacy, University of Greifswald, Greifswald, Germany

* Correspondence:

Mia M. Bengtsson

Institute of Microbiology, University of Greifswald

Felix-Hausdorff-Straße 8

17489 Greifswald, Germany

mia.bengtsson@uni-greifswald.de

Keywords: phytoplankton$_1$, diatoms$_2$, *Desulfobacterota*$_3$, networks$_4$, North Sea$_5$.

Abstract

Phytoplankton blooms fuel marine food webs with labile dissolved carbon, but also lead to the formation of particulate organic matter composed of living and dead algal cells. These particles contribute to carbon sequestration, yet are also sites of intense algal-bacterial interactions and provide diverse niches for microbes to thrive. We analyzed 16S and 18S ribosomal RNA gene amplicon sequences obtained from 51 time points and metaproteomes from 3 time points during a spring phytoplankton bloom in the North Sea. Particulate

fractions larger than 10 μm diameter were collected at near daily intervals between early March and late May in 2018. Network analysis identified two major modules representing bacteria co-occurring with diatoms and with dinoflagellates, respectively. The diatom network module included known sulfate-reducing *Desulfobacterota* as well as potentially sulfur-oxidizing *Ectothiorhodospiraceae*. Metaproteome analyses confirmed presence of key enzymes involved in dissimilatory sulfate reduction, a process known to occur in sinking particles at greater depths. Our results indicate the presence of sufficiently anoxic niches in the particle fraction of an active phytoplankton bloom to sustain sulfate reduction, which may have implications for algal-bacterial interactions and carbon export during blooms.

Introduction

Microalgae inject gigatons of organic carbon into coastal oceans every year (Field et al. 1998) and phytoplankton blooms represent primary productivity hotspots. It has been estimated that over 90% of algal-produced carbon is consumed by heterotrophic bacteria in the immediate vicinity of algal cells, the phycosphere, during typical bloom situations (Seymour et al. 2017). Some of these bacteria are directly associated with living algal cells, or with sinking aggregates of senescent or dead algae, and therefore play an important role in the biological carbon pump. Despite their importance in carbon sequestration (e.g., Bligh et al. 2022), vertical connectivity (Mestre et al. 2018), and their documented complexity (e.g. Reintjes et al. 2023), particle-associated (PA) bacterial communities are less well understood than their free-living counterparts. In fact, they are often overlooked due to the sometimes fragile nature of particles, in combination with the practice of pre-filtration to exclude larger organisms prior to molecular analyses (e.g. Simon et al. 2002, Thiele et al. 2015, Heins et al. 2021).

A defining feature of marine particles are the steep chemical and redox gradients that PA bacterial communities are exposed to, compared to free-living, planktonic bacteria. These gradients have been intensively studied in the context of bathypelagic sinking particles (i.e., marine snow), which harbor anaerobic micro-niches enabling, e.g., microbial sulfate reduction (Bianchi et al. 2018, Raven et al. 2021). In the photic zone, anaerobic micro-niches have been less widely investigated and the prevalence of sulfate-reducing bacteria is uncertain in the proximity of oxygen-producing living algal cells. Aggregates composed of living algae are rather known for a diverse microbiome of aerobic,

heterotrophic bacteria that degrade complex algal organic matter, such as polysaccharide-rich exudates (Enke et al. 2019, Reintjes et al. 2023), although anaerobic metabolism such as diazotrophy is also known to occur (Riemann et al. 2022). Within this microbiome, bacteria are selected by factors such as host physiology and genotype (Ahern et al. 2021), the surrounding environment (Barreto Filho et al. 2021) and via stochastic processes (Stock et al. 2022). It is methodologically challenging to study associations between algal and bacterial taxa during natural phytoplankton blooms by direct observations, due to the transient nature of these associations (Heins et al. 2021, Seymour et al. 2017), as well as innate complexities and rapid dynamics of algal and bacterial communities during bloom events (Teeling et al. 2016). High-resolution temporal co-occurrence analysis, in combination with measurement of microbial functional potential, offers an indirect way to infer algae-bacteria associations, which can facilitate generating hypotheses about specific interactions and their potential functional implications.

Here, we investigated microbial community dynamics of PA bacterial and eukaryotic taxa during a spring phytoplankton bloom in the southern North Sea at the long-term ecological research site Helgoland Roads (Wiltshire 2010) in the year 2018. We aimed to identify PA bacterial taxa co-occurring with the major eukaryotic taxa (diatoms and dinoflagellates) during the bloom. We hypothesized that bacteria co-occurring with diatoms would be compositionally and functionally distinct from those co-occurring with dinoflagellates. Using 16S and 18S rRNA gene amplicon data from a well-resolved time series (near daily sampling) collected between early March and late May in 2018, we constructed co-occurrence networks focusing exclusively on bacteria-eukaryote co-occurrences in particles larger than 10 µm. In addition, we addressed bacterial functional gene expression by analysis of metaproteomes from three selected time points during the bloom.

Materials and Methods

Sampling and sample processing

A large volume of subsurface seawater (40 L - 140 L, 1 m depth) was sampled in the morning at near daily intervals between beginning of March and end of May in 2018 at the station long term ecological research (LTER) site Helgoland Roads (50° 11.3' N, 7° 54.0' E; DEIMS.iD: https://deims.org/1e96ef9b-0915-4661-849f-b3a72f5aa9b1) near Helgoland in

the south-eastern North Sea. Secchi depth was measured from the vessel on site. For
chlorophyll *a* (chl *a*) analysis, sample filtration was carried out in a laboratory under dim light
to avoid the loss of pigments during the filtration process. We used a combined method of
Zapata et al., (2000) and Garrido et al., (2003) for chl *a* extraction and analysis. Pigments
were separated via high-performance liquid chromatography (HPLC) (Waters 2695
Separation Module), and detected with a Waters 996 Photodiode Array Detector. For 16S
and 18S rRNA gene amplicon sequencing and metagenome analysis, plankton biomass
from a 1 L seawater subsample was filtered using 10 µm pore size polycarbonate
membrane filters (47 mm diameter, Millipore, Schwalbach, Germany) to separate PA
microbes from smaller size fractions. At three selected time points during the bloom (Julian
days 107, 128 and 144, representing early-, mid- and late bloom phases), a separate
filtration was performed using larger (142 mm diameter, Millipore) 10 µm polycarbonate
membrane filters for metaproteomic analysis. In order to maximize biomass harvest while
avoiding clogging, filtered volumes varied between 15 and 30.5 L per filter for
metaproteomic analysis.
111

rRNA gene amplicon sequencing
Samples from 52 time points throughout the bloom were collected and analyzed. DNA was
extracted from the filters using the Qiagen DNeasy Power soil Pro kit (Qiagen, Hilden,
Germany) according to the manufacturer's instructions. Dislocation of microbial cells from
the filters and mechanical lysis were achieved by bead beating in a FastPrep 24 5G (MP
Biomedicals, Irvine, CA, USA). DNA concentrations were measured at a Qubit 3.0
fluorometer (Invitrogen, Carlsbad, CA, USA). Extracted DNA was amplified with primer pairs
targeting the V4 region of the 16S rRNA gene [515f: 5'-GTGYCAGCMGCCGCGGTAA-3',
806r: 5'-GGACTACNVGGGTWTCTAAT-3' (Walters et al., 2016)] and the V7 region of the
18S rRNA gene [F-1183mod: 5'-AATTTGACTCAACRCGGG-3', R-1443mod: 5'-
GRGCATCACAGACCTG-3'] (Ray et al. 2016) coupled to custom adaptor-barcode
constructs. PCR amplification and Illumina MiSeq (Illumina, San Diego, CA, USA) library
preparation and sequencing (V3 chemistry) were carried out by LGC Genomics (LGC
Genomics, Berlin, Germany).
Sequence reads free of adaptor and primer sequence remains were processed using
the DADA2 package (v1.2.0) in R (Callahan et al., 2016). In summary, forward and reverse
Illumina MiSeq reads were truncated to 200 bp, filtered (maxEE = 2, truncQ = 2, minLen =

175), dereplicated and error rates were estimated using the maximum possible error estimate from the data as initial guess. Sample sequences were inferred, paired forward and reverse reads were merged and chimeric sequences were removed using the removeBimeraDenovo function. The resulting amplicon sequence variants (ASVs) were taxonomically classified using the Silva database (nr 99 v 138.1) for 16S rRNA and the PR2 database (version 4.13, minboot: 50) for 18S rRNA sequences using the build-in RDP classifier. 16S rRNA gene amplicon reads classified as chloroplasts and mitochondria, as well as the 18S rRNA gene reads classified as *Metazoa* (zooplankton) were removed prior to downstream analyses (a single time point was excluded due to suspected contamination).

Co-occurrence networks were generated in R using Spearman rank correlation, as described previously (Bengtsson et al. 2017, correlation coefficient >0.7, adjusted p<0.01). We considered exclusively correlations between 18S and 16S ASVs and the final network was plotted using the igraph R package.

Construction of a metagenome-based database

Illumina sequencing was performed at the Max Planck Genome Centre Cologne on DNA extracted from both 3-10 µm- and >10 µm filter fractions from eight selected time points (19 March, 12 April, 17 April, 26 April, 08 May, 11 May, 22 May and 29 May 2018 - Julian days 78, 102, 107, 116, 128, 131, 142 and 149), using the HiSeq 2500 platform and 2 x 150 bp chemistry. Filtration and trimming of the reads were done as described previously (Francis et al. 2021). Read quality for each sample was confirmed using FastQC v0.11.9 (Andrews 2014). Assembly was performed with MEGAHIT v1.2.9 (Li et al. 2015) (kmer length: 21). Contigs below 2.5 kbp were removed using anvi-script-reformat-fasta from within Anvi'o v6.2 (Eren et al. 2015). Genes were predicted using Prodigal v2.6.3 (Hyatt et al. 2010) as implemented in the Prokka v1.11 annotation pipeline (Seemann 2014). These genes sequences were subsequently combined with a list of common contaminants to generate a database for peptide-spectrum-matching (PSM) after elimination of redundant sequences (97% redundancy) with CD-Hit (Fu et al. 2012, Li and Godzik 2006).

Metaproteomics

Proteins were extracted from biomass filters of the 17th of April, 8th of May and 24th of May 2018 (Julian days 107, 128 and 144) as described previously (Schultz et al. 2020, Schultz 2022) and analyzed with liquid chromatography-tandem mass spectrometry in triplicates. Briefly, proteins were extracted via bead-beating followed by acetone precipitation. The extracts were separated and fractionated by 1D SDS-PAGE and in-gel trypsin digested. After desalting and concentration of the peptides using C18 Millipore® ZipTip columns, the samples were measured with an Orbitrap Velos™ mass spectrometer (ThermoFisher Scientific, Waltham, MA, USA). After conversion into mgf file format using MS convert in ProteoWizard (Palo Alto, CA, USA), spectra were matched against a metagenome-based database containing 14,764,755 entries. Mascot (Matrix Science, London, UK) and Scaffold (Proteome Software, Portland, OR, US) were used for peptide-spectrum-matching, protein identification and protein grouping. Instead of setting an FDR-threshold, identification of protein groups was based on number of peptide-matches with a minimum of two, a protein threshold of 99% and a peptide threshold of 95%. Identified protein groups were annotated via Prophane v6.2.3 (Schiebenhoefer et al. 2020), using the Uniprot-TrEMBL (as of September 2021) and NCBI nr (as of February 2022) databases for taxonomic and EggNog v5.0.2 and Tigrfams 15 databases for functional annotation with Prophane default settings. Taxonomic information for *Desulfobacterota* proteins was manually confirmed against the most recent ncbi nr database (blastp, https://blast.ncbi.nlm.nih.gov, September 2023). TrEMBL taxonomic information for *Desulfobacterales* was manually curated and set to phylum level for better comparability with the Silva database. Relative abundances were calculated as Normalized Spectral Abundance Factor (NSAF) values using the quantification method "max_nsaf" integrated in Prophane (Schiebenhoefer et al., 2020). Briefly, SAF values were calculated by dividing exclusive unique spectrum counts for each protein group by protein length of the longest sequence in that protein group. For normalization, SAF values were then divided by the sum of all SAF values of the sample.

Data visualization

Stacked bar plots for rRNA gene amplicon data and metaproteomics were created with R version 4.3.0 using the tidyverse package (Wickham et al. 2019) in combination with the svglite, polychrome, patchwork, glue and ggnested packages.

Results

Particle microbial community dynamics during the course of the bloom

The spring phytoplankton bloom in 2018 was characterized by an initial dominance of diatoms (*Bacillariophyceae*), followed by an increase in dinoflagellate (*Dinophyceae*) relative abundances after the bloom peak (peak in chl *a* concentration, Fig. 1a and b). The algal bloom (chl *a* measurements) peaked around the 26th of April (Julian day 116), coinciding with a decrease in water clarity, as indicated by Secchi depth rising at the onset of the bloom. 16S rRNA gene amplicon sequencing revealed a total of 17615 ASVs in the particle bacterial microbiomes, which consisted mainly of *Proteobacteria (*e.g. *Spongibacteriaceae, Methylophagaceae, Rhodobacteriaceae)*, *Bacteroidetes* (e.g. *Polaribacter*), *Verrucomicrobia* (e.g. *Persicirhabdus*) and *Planctomycetes* (e.g. *Phycisphaeraceae*). Further, *Actinobacteria* (e.g. *Illumatobacter*)*,* and, notably, *Desulfobacterota* were also relatively abundant (Fig. 1c).

Taxonomic and functional composition as detected by metaproteome analysis

Metaproteome sampling was performed on three time points that correspond to the early, mid and late phase of the bloom (Julian days 107, 128 and 144), as indicated in Fig. 1a. We detected 4,584 protein groups in >10 µm filter fractions from these sampled time points. Metaproteomic analyses confirmed high abundances of *Proteobacteria* and *Bacteroidetes* in the bacterial fraction (Fig. 2a), and dominance of *Bacillariopyhta* in the eukaryote fraction (Fig. 2b), with eukaryote proteins making up 90% of all proteins at the first selected time point. The proportion of eukaryotic proteins was reduced to 70% in the last selected time point (144), while bacterial proteins became comparably more abundant (25% on day 144 compared to 4% on day 107). Functional analysis revealed a predominance of eukaryotic proteins involved in metabolism, including energy production and conversion, and a shift towards expression of proteins relevant in cellular processes and signaling over the course of the bloom (Fig. 2c).

Co-occurrence network analysis

Co-occurrence analysis resulted in distinct network modules centered around diatom- and dinoflagellate 18S rRNA gene amplicon sequence variants (ASVs). Based on significant

positive correlations (Spearman Rho>0.7, corrected p>0.01) between eukaryotic 18S rRNA
gene and prokaryotic 16S rRNA gene ASVs, including the 51 analyzed time points during
the 2018 spring bloom, the network was dominated by three major distinct modules (Fig.
3a). Two of these modules were dominated by eukaryotic ASVs belonging to diatoms and
dinoflagellates, respectively, while the third module mostly contained diatom ASVs and was
linked to the dinoflagellate-dominated network module. Along the time line of the bloom,
these modules roughly corresponded to the phytoplankton taxa prevalent in the early stages
of the bloom (module I, mainly diatoms), during the late stages of the bloom (module II,
mainly dinoflagellates) and mid-bloom around peak chl *a* (module III, Fig. 1). The
composition of bacterial ASVs that co-occurred with diatoms and dinoflagellates is depicted
in Fig. 3b. Eleven ASVs belonging to the *Desulfobacterota* were part of the main diatom-
dominated network module I and co-occurred exclusively with diatoms, including the genus
Desulfosarcina, and the families *Desulfocapsaceae* and *Desulfobulbaceae* as well as the
lineage Sva1033 (Ravenschlag et al. 1999). In addition, 4 ASVs classified as
Ectothiorhodospiraceae (genus *Thiogranum*) also co-occurred with diatoms (Fig. 3b).

Analysis of Desulfobacterota *proteins*
Out of the 4,584 protein groups detected by metaproteome analysis, 19 were classified as
belonging to different *Desulfobacterota* orders (Table S1). The relative abundance of
predicted *Desulfobacterota* proteins averaged over all time points in the metaproteome was
0.14% (expressed as normalized spectral abundance factor - NSAF). This order of
magnitude corresponded well to the relative abundance of *Desulfobacterota* ASVs in the
microbiome at the same time points (average 0.38%, Fig 1c). Remarkably, no less than
32% of the *Desulfobacterota* metaproteome fraction (in terms of protein abundance)
consisted of key enzymes for dissimilatory sulfate reduction (Fig. 3c). Of these, ATP
sulfurylase, both the alpha and the beta subunits of dissimilatory sulfite reductase (DSR)
and adenylyl-sulfate (adenosine-5'-phosphosulfate) reductase (APS reductase) were
detected. We used a conservative threshold for protein identification of at least two
matching peptides to consider a protein as validly detected.

Discussion

The succession we observed during the 2018 spring bloom, with an initial dominance of
diatoms, followed by dinoflagellates, using 18S rRNA gene amplicon sequencing agrees
with the typical phytoplankton community succession at Helgoland Roads (Wiltshire et al.
2008, Käse et al. 2020). Metaproteome analysis highlighted the dominance of eukaryotic
proteins in the sampled particles, making up 70-80% of detected proteins, most of which
belonged to the major diatom phytoplankton. The high prevalence of proteins involved in
basic metabolism such as energy production and conversion is consistent with the sampled
>10 µm particle fraction mostly comprising of living phytoplankton cells, especially at the
beginning of the bloom. However, bacterial proteins were present in all of the three selected
time points, with a notable peak at day 144, and their taxonomic composition was very
similar to that observed via 16S rRNA gene amplicon sequencing. Overall, the composition
of particle bacterial microbiomes agreed with other reports from similar environments
(Reintjes et al. 2023, Schultz et al. 2020, Heins et al. 2021, Crump et al. 1999).

As hypothesized, bacteria co-occurring with abundant diatoms formed a distinct network
module, with taxa that were different from those co-occurring with dinoflagellates in a
second distinct module. A third network module contained mostly diatom taxa, but also one
dinoflagellate ASV and other phytoplankton taxa. These network patterns offer an
alternative way to visualize the temporal dynamics of the bloom, and should not be
interpreted as evidence of physical interactions between taxa (Röttjers & Faust 2018).
However, one pattern that is striking in our network analysis is the exclusive co-occurrence
of *Desulfobacterota* with diatoms.

We observed a high relative abundance of *Desulfobacterota* (in total 0.38% of bacterial
amplicon reads) in particles (>10 µm), especially in the early phase of the bloom when
diatoms were dominating. Despite the limited resolution of metaproteome data compared to
DNA-based methods, desulfobacterial proteins were represented at all selected time points
(in total 0.14% of metaproteomes). Network analysis further highlighted the temporal co-
occurrence of *Desulfobacterota* with several diatom taxa. This raises the question of the
niche filled by these anaerobic bacteria during an active phytoplankton bloom.

Diatoms, such as *Thallassiosira* spp. and *Thalassionema* spp., which were co-occurring
with *Desulfobacterota* in this study (Fig. 3a), are known to form aggregates (Thornton

2002). For example, several species of *Thalassiosira* extrude long chitin fibrils, which
prevent sinking and bind exopolymeric substances (EPS) also produced by the algal cells
(Den et al. 2023, Herth & Barthlott 1979). This creates a favorable environment for bacteria
to attach, which can in turn stimulate algal EPS production (Gärdes et al. 2011). EPS
makes particles adhesive and thus bacteria can be captured in this sticky EPS layer.
Smaller particles can aggregate to larger particles by collision and adhesion, in particular
during phytoplankton blooms with high particle densities. Such aggregates can feature high
numbers of living, photosynthesizing algal cells (Thornton 2002) which produce ample
oxygen during the day, but at night respiration by the algal cells and their surrounding
bacteria may deplete oxygen sufficiently for low oxygen or even anaerobic micro-niches to
form.

Desulfobacterota have not been identified as frequent members of diatom microbiomes
in either cultures or in the field so far (Helliwell et al. 2022). However, a recent global survey
of the diatom interactome detected positive correlations between diatoms and sulfate-
reducing bacteria in samples from the Tara Oceans expedition (Vincent & Bowler 2020). In
addition, *Desulfobacterota* have previously been repeatedly detected in particle-associated
communities in the photic zone (Crump et al. 1999, Liu et al. 2020, Hallstrøm et al. 2022).
Ectothiorhodospiraceae are purple sulfur bacteria (belonging to *Gammaproteobacteria*), that
are anaerobic to microaerophilic (Imhoff 2022). They can oxidize H_2S, e.g., produced via
dissimilatory sulfate reduction by members of the *Desulfobacterota.* Our observation of
Desulfobacterota as well as *Ectothiorhodospiraceae* co-occurring with diatoms is consistent
with potential sulfur cycling during the diatom-dominated phase of this phytoplankton bloom
under anoxic to very low oxygen conditions. While co-occurrence of diatom and
Desulfobacterota rRNA genes does not by itself indicate that sulfate reduction is taking
place in diatom-derived particles, detection of key functional enzymes by metaproteomics
suggests that *Desulfobacterota* were actively carrying out sulfate reduction and thus gaining
energy through anaerobic respiration during the bloom. The higher detection of these
enzymes especially at the last time point (Julian day 144, Table S1) can likely be attributed
to the larger proportion of bacterial proteins at this time point, when the diatom bloom was
declining.

Temporal associations, such as detected by our co-occurrence network analyses, have
to be interpreted with caution as additional variable factors that were not taken into account
may influence the observed correlations. Importantly, we cannot quantitatively assess the

influence of underlying sediment microbiomes at the rather shallow sampling site (6-10 m
depth depending on tide), which frequently become resuspended during times of heavy
wind thereby introducing anaerobic microbes to the pelagic environment. Although we did
not find a clear link between Secchi depth, wind conditions and *Desulfobacterota* relative
abundances in this study (Fig. 1, Fig. S1), the detected ASVs (e.g. classifying as
Desulfosarcina, *Desulfocapsaceae*, *Desulfobulbaceae*) are closely related to typical benthic
lineages (Ravenschlag et al. 1999), suggesting that they originated from the underlying
sediments. The diatoms co-occurring with *Desulfobacteria* were primarily classified as
common pelagic genera such as *Thalassiosira* and *Thalassionema*. However, the genus
Brockmanniella, which can also inhabit benthic biofilms (Hernández Fariñas et al. 2017),
was also found within network module I, suggesting some influence of sediment
resuspension. In shallow environments, benthic and pelagic microbiomes are in close
contact within the photic zone, and seeding of pelagic particles by benthic microbes (and
vice versa) is likely frequent, which may explain the observed co-occurrences. However, our
methodology does not allow us to determine the physical proximity of *Desulfobacterota* and
diatom cells. The observed co-occurrences may therefore reflect a purely temporal
association between pelagic diatoms and resuspended sediment bacteria. Indeed,
demonstrating a potential physical association between *Desulfobacterota* and diatoms
would require *in situ* microscopic investigation using for example taxon- or gene- specific
fluorescent probes. Likewise, it has yet to be clarified, whether such temporal or physical
associations are annually recurrent, under what specific conditions they occur, and which
specific diatom taxa are involved. Thus, further studies are needed to confirm whether our
results are representative for phytoplankton blooms in shallow water coastal areas.

Our results highlight the complexity of particle microbiomes and corroborate the need to
study algae-bacteria particles as spatially heterogeneous entities. The microbiomes of
aggregate-forming phytoplankton may indeed be similarly complex as those of animals and
other multicellular organisms, featuring distinct micro-niches with sub-microbiomes
(analogous to e.g., human skin vs. gut microbiomes), whose compositions depend on the
chemical environment on a small spatial scale. Anaerobic micro-niches in phytoplankton-
derived aggregates may affect carbon cycling insofar, as sulfate reduction can chemically
alter particulate organic matter, rendering it resistant to microbial degradation (Raven et al.
2021). Anaerobic niches in marine phytoplankton-derived particles were already frequently
reported in the context of nitrogen fixation (i.e., diazotrophy, e.g. Riemann et al. 2022).

Recent work has shown that many *Desulfobacterota* are in fact capable of both diazotrophy and sulfate reduction (Liesirova et al. 2023) and that diazotrophic *Desulfobacterota* can move towards phytoplankton-derived organic matter via chemotaxis. Similar to diazotrophs, sulfate reducers in turn provide additional niches for bacteria which scavenge their metabolic products, i.e., sulfur oxidizers such as the observed *Ectothiorhodospiraceae*, leading to higher metabolic diversity and complexity within particles. Considering the significance of marine particle microbiomes in carbon sequestration, it is vital to understand the consequences of the metabolic and compositional complexity in particle microbiomes, and the resulting microbial interactions.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

RS conducted the metaproteomic analyses together with DS and DZ, analyzed the data, prepared figures and contributed to writing the manuscript. MSF performed amplicon sequencing analyses together with AB. FW prepared metagenomics databases together with HT and KJH. PAM performed mass-spectrometry measurements together with DB. JB contributed to data analysis and figure preparation. IVK contributed chl *a* and Secchi depth data together with KHW. TU contributed to data analysis and interpretation. KR supervised metaproteome analysis and conceived the study together with MMB. MMB supervised and contributed to amplicon data collection, analysis, interpretation and wrote the manuscript with significant input from RS, HT, TU and IVK. All authors approved the final version of the manuscript.

Funding

This study was funded by the Deutsche Forschungsgemeinschaft (DFG) in the framework of the research unit FOR2406 'Proteogenomics of Marine Polysaccharide Utilization (POMPU)' by grants of KR/MMB (RI 969/9-2), DB (BE 3869/4-3) and HT (TE 813/2-3). Access to Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung

(AWI) infrastructure was ensured through the Helmholtz Association's LK-II performance category program (AWI_BAH_o 1).

Acknowledgments

We thank the Biological Station Helgoland, Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung and the crews from FS Aade and FS Uthörn for help with sampling, analyses, logistics, and providing lab space. Jan Rockstroh and Kira Wisnewski assisted in sample processing for metaproteomics. Thomas Ben Francis assisted in metagenome database construction. We are also grateful to the whole POMPU consortium for helpful discussions, especially Rudi Amann, Jens Harder, Saskia Kalenborn and Jan-Hendrik Hehemann.

Data Availability Statement

Metagenome raw reads and assembled contigs used for metaproteome assignment are publicly available at the European nucleotide archive under the project accession number PRJEB38290. 16S- and 18S-rRNA gene amplicon data are also available via ENA under the project accession number PRJEB51816. Mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2022) repository with dataset identifier PXD035982. Original mass spectrometry proteomics data can be accessed for review (**Username:** reviewer_pxd035982@ebi.ac.uk **Password:** umim4qCW) and will be made publicly available upon manuscript acceptance.

References

Ahern OM, Whittaker KA, Williams TC *et al.* Host genotype structures the microbiome of a globally dispersed marine phytoplankton. *Proceedings of the National Academy of Sciences of the United States of America* 2021;**118**, DOI: 10.1073/pnas.2105207118.

Andrews S, FastQC: a quality control tool for high throughput sequence data. 2014. Available online at: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (accessed September 13, 2023).

Barreto Filho MM, Walker M, Ashworth MP *et al.* Structure and Long-Term Stability of the Microbiome in Diverse Diatom Cultures. *Microbiology spectrum* 2021;**9**:e0026921.

Bengtsson MM, Bühler A, Brauer A *et al.* Eelgrass Leaf Surface Microbiomes Are Locally Variable and Highly Correlated with Epibiotic Eukaryotes. *Frontiers in microbiology* 2017;**8**:1312.

Bianchi D, Weber TS, Kiko R *et al.* Global niche of marine anaerobic metabolisms expanded by particle microenvironments. *Nature Geoscience* 2018;**11**:263–8.

Bligh M, Nguyen N, Buck-Wiese H *et al.* Structures and functions of algal glycans shape their capacity to sequester carbon in the ocean. *Current Opinion in Chemical Biology* 2022;**71**:102204.

Callahan BJ, McMurdie PJ, Rosen MJ *et al.* DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods* 2016;**13**:581–3.

Crump BC, Armbrust EV, Baross JA. Phylogenetic Analysis of Particle-Attached and Free-Living Bacterial Communities in the Columbia River, Its Estuary, and the Adjacent Coastal Ocean. *Appl Environ Microbiol* 1999;**65**:3192–204.

Den TQ, Neu TR, Sultana S *et al.* Distinct glycoconjugate cell surface structures make the pelagic diatom *Thalassiosira rotula* an attractive habitat for bacteria. *Journal of Phycology* 2023;**59**:309–22.

Enke TN, Datta MS, Schwartzman J *et al.* Modular Assembly of Polysaccharide-Degrading Marine Microbial Communities. *Current Biology* 2019;**29**:1528-1535.e6.

Eren AM, Esen ÖC, Quince C *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 2015;**3**:e1319.

Field CB, Behrenfeld MJ, Randerson JT *et al.* Primary Production of the Biosphere: Integrating Terrestrial and Oceanic Components. *Science* 1998;**281**:237–40.

Francis TB, Bartosik D, Sura T *et al.* Changing expression patterns of TonB-dependent transporters suggest shifts in polysaccharide consumption over the course of a spring phytoplankton bloom. *The ISME journal* 2021;**15**:2336–50.

Fu L, Niu B, Zhu Z *et al.* CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics (Oxford, England)* 2012;**28**:3150–2.

Gärdes A, Iversen MH, Grossart H-P *et al.* Diatom-associated bacteria are required for aggregation of Thalassiosira weissflogii. *The ISME journal* 2011;**5**:436–45.

Garrido J, Rodriguez F, Campana E *et al.* Rapid separation of chlorophylls a and b and their demetallated and dephytylated derivatives using a monolithic silica C18 column and a pyridine-containing mobile phase. *Journal of Chromatography A* 2003;**994**:85–92.

Hallstrøm S, Raina J-B, Ostrowski M *et al.* Chemotaxis may assist marine heterotrophic
bacterial diazotrophs to find microzones suitable for N2 fixation in the pelagic ocean.
ISME J 2022;**16**:2525–34.

Heins A, Reintjes G, Amann RI *et al.* Particle Collection in Imhoff Sedimentation Cones
Enriches Both Motile Chemotactic and Particle-Attached Bacteria. *Front Microbiol*
2021;**12**:643730.

Helliwell KE, Shibl AA, Amin SA. The Diatom Microbiome: New Perspectives for Diatom-
Bacteria Symbioses. In: Falciatore A, Mock T (eds.). *The Molecular Life of Diatoms.*
Cham: Springer International Publishing, 2022, 679–712.

Hernández Fariñas T, Ribeiro L, Soudant D *et al.* Contribution of benthic microalgae to the
temporal variation in phytoplankton assemblages in a macrotidal system. Wood M
(ed.). *J Phycol* 2017;**53**:1020–34.

Herth W, Barthlott W. The site of β-chitin fibril formation in centric diatoms. I. Pores and fibril
formation. *Journal of Ultrastructure Research* 1979;**68**:6–15.

Hyatt D, Chen G-L, Locascio PF *et al.* Prodigal: prokaryotic gene recognition and translation
initiation site identification. *BMC bioinformatics* 2010;**11**:119.

Imhoff JF, Kyndt JA, Meyer TE. Genomic Comparison, Phylogeny and Taxonomic
Reevaluation of the Ectothiorhodospiraceae and Description of Halorhodospiraceae
fam. nov. and Halochlorospira gen. nov. *Microorganisms* 2022;**10**, DOI:
10.3390/microorganisms10020295.

Käse L, Kraberg AC, Metfies K *et al.* Rapid succession drives spring community dynamics
of small protists at Helgoland Roads, North Sea. *Journal of Plankton Research*
2020;**42**:305–19.

Li D, Liu C-M, Luo R *et al.* MEGAHIT: an ultra-fast single-node solution for large and
complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*
(Oxford, England) 2015;**31**:1674–6.

Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. *Bioinformatics (Oxford, England)* 2006;**22**:1658–9.

Liesirova T, Aarenstrup-Launbjerg T, Hallstrøm S *et al.* Nitrogen-fixing sulfate reducing
bacteria in shallow coastal sediments under simulated resuspension. *Estuarine,*
Coastal and Shelf Science 2023;**280**:108165.

Liu Y, Lin Q, Feng J *et al.* Differences in metabolic potential between particle-associated
and free-living bacteria along Pearl River Estuary. *Science of The Total Environment*
2020;**728**:138856.

Mestre M, Ruiz-González C, Logares R *et al.* Sinking particles promote vertical connectivity
in the ocean microbiome. *Proc Natl Acad Sci USA* 2018;**115**, DOI:
10.1073/pnas.1802470115.

Perez-Riverol Y, Bai J, Bandla C *et al.* The PRIDE database resources in 2022: a hub for
mass spectrometry-based proteomics evidences. *Nucleic Acids Research*
2022;**50**:D543–52.

Raven MR, Keil RG, Webb SM. Microbial sulfate reduction and organic sulfur formation in
sinking marine particles. *Science (New York, NY)* 2021;**371**:178–81.

Ravenschlag K, Sahm K, Pernthaler J *et al.* High Bacterial Diversity in Permanently Cold
Marine Sediments. *Appl Environ Microbiol* 1999;**65**:3982–9.

Ray JL, Althammer J, Skaar KS *et al.* Metabarcoding and metabolome analyses of copepod
grazing reveal feeding preference and linkage to metabolite classes in dynamic
microbial plankton communities. *Molecular ecology* 2016;**25**:5585–602.

Reintjes G, Heins A, Wang C *et al.* Abundance and composition of particles and their
attached microbiomes along an Atlantic Meridional Transect. *Front Mar Sci*
2023;**10**:1051510.

Riemann L, Rahav E, Passow U *et al.* Planktonic Aggregates as Hotspots for Heterotrophic
Diazotrophy: The Plot Thickens. *Front Microbiol* 2022;**13**:875050.

Schiebenhoefer H, Schallert K, Renard BY *et al.* A complete and flexible workflow for
metaproteomics data analysis based on MetaProteomeAnalyzer and Prophane.
Nature protocols 2020;**15**:3212–39.

Schultz D, Zühlke D, Bernhardt J *et al.* An optimized metaproteomics protocol for a holistic
taxonomic and functional characterization of microbial communities from marine
particles. *Environmental microbiology reports* 2020;**12**:367–76.

Schultz, D. (2022). Mechanisms of polysaccharide degradation of particle-associated
microbial communities. PhD thesis, University of Greifswald.

Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics (Oxford, England)*
2014;**30**:2068–9.

Seymour JR, Amin SA, Raina J-B *et al.* Zooming in on the phycosphere: the ecological
interface for phytoplankton-bacteria relationships. *Nature microbiology* 2017;**2**:17065.

Simon M, Grossart H, Schweitzer B *et al.* Microbial ecology of organic aggregates in aquatic
ecosystems. *Aquat Microb Ecol* 2002;**28**:175–211.
Stock W, Willems A, Mangelinckx S *et al.* Selection constrains lottery assembly in the
microbiomes of closely related diatom species. *ISME Communications* 2022;**2**, DOI:
10.1038/s43705-022-00091-x.
Teeling H, Fuchs BM, Bennke CM *et al.* Recurring patterns in bacterioplankton dynamics
during coastal spring algae blooms. *eLife* 2016;**5**:e11888.
Thiele S, Fuchs BM, Amann R *et al.* Colonization in the Photic Zone and Subsequent
Changes during Sinking Determine Bacterial Community Composition in Marine
Snow. Wommack KE (ed.). *Appl Environ Microbiol* 2015;**81**:1463–71.
Thornton DCO. Diatom aggregation in the sea: mechanisms and ecological implications.
European Journal of Phycology 2002;**37**:149–61.
Vincent F, Bowler C. Diatoms Are Selective Segregators in Global Ocean Planktonic
Communities. *mSystems* 2020;**5**, DOI: 10.1128/mSystems.00444-19.
Walters W, Hyde ER, Berg-Lyons D *et al.* Improved Bacterial 16S rRNA Gene (V4 and V4-
5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial
Community Surveys. *mSystems* 2016;**1**, DOI: 10.1128/mSystems.00009-15.
Wickham H, Averick M, Bryan J *et al.* Welcome to the Tidyverse. *JOSS* 2019;**4**:1686.
Wiltshire KH, Kraberg A, Bartsch I *et al.* Helgoland Roads, North Sea: 45 Years of Change.
Estuaries and Coasts 2010;**33**:295–310.
Wiltshire KH, Malzahn AM, Wirtz K *et al.* Resilience of North Sea phytoplankton spring
bloom dynamics: An analysis of long-term data at Helgoland Roads. *Limnol*
Oceanogr 2008;**53**:1294–302.
Zapata M, Rodríguez F, Garrido J. Separation of chlorophylls and carotenoids from marine
phytoplankton:a new HPLC method using a reversed phase C8 column and pyridine-
containing mobile phases. *Mar Ecol Prog Ser* 2000;**195**:29–45.

Figure Captions

Figure 1: Progression of the 2018 spring bloom by Helgoland, North Sea. (a) Chl *a*
(green) peaked around the 26th of April (Julian day 116), coinciding with a drop in Secchi
depth (brown). Asterisks indicate time points (Julian days) for which metaproteomic
sampling was performed. (b) 18S rRNA gene amplicon sequencing revealed that diatoms

(*Bacillariophyta*) were the most abundant phytoplankton lineage, although dinoflagellates
(*Dinophyceae*) dominated after the chl *a* peak. (c) 16S rRNA gene amplicon sequencing
showed the highest relative abundances of *Desulfobacterota* (yellow) during the first half of
the bloom.

Figure 2: Taxonomic and functional annotation of particle metaproteomes from three
selected time points during the bloom. (a) The proportion of bacterial proteins increased
during the course of the bloom (b) Eukaryotic proteins made up the majority of identified
proteins and showed an initial strong dominance of *Bacillariophyta* (c). Functional
annotation of metaproteomes indicated a high contribution of metabolism-related eukaryotic
proteins early in the bloom, while cellular processes and signaling increased during later
time points.

Figure 3: Co-occurrence of eukaryotes and bacteria, and detected protein groups of
Desulfobacterota (a) A network analysis of 18S rRNA (triangles) and 16S rRNA (circles)
gene ASV co-occurrences resulted in two major network modules containing diatom [I] and
dinoflagellate [II] 18S ASVs, respectively, as well as one mixed module [III]. The network
was calculated based on Spearman correlations (r >0.7, p < 0.01) exclusively between 18S
ASVs and 16S ASVs. *Desulfobacterota* (yellow circles) were only associated with the
diatom-dominated module I. The sizes of the symbols correspond to the number of
significant correlations of the nodes (degree). (b) The bacterial taxa co-occurring with
diatoms and dinoflagellates, respectively, are displayed as horizontal bars with a length
proportional to the number of ASVs belonging to each lineage. *Desulfobacterota* (yellow) as
well as potentially sulfur-oxidizing *Ectothiorodospiraceae* (*Gammaproteobacteria*, blue)
were positively associated with diatoms but not with dinoflagellates. (c) The pie chart
displays relative abundances for all 19 protein groups classified as belonging to
Desulfobacterota during all three time points sampled for metaproteomics. Of these, seven
protein groups (32%, bright yellow) represented enzymes involved in dissimilatory sulfate
reduction. Yellow circles indicate which of the key enzymes in this metabolic pathway were
detected.

a

(Raphid pennate)

Thalassiosira guillardii
(Polar centric Mediophyceae)

Thalassiosira minima
(Polar centric Mediophyceae)

Filosa-Thecofilosea

[I]

(Polar centric Mediophyceae)

Ichthyosporea

Spirotrichea

Brockmanniella brockmannii
(Polar centric Mediophyceae)

Thalassionema frauenfeldii
(Araphid pennate)

Filosa-Thecofilosea

Filosa-Thecofilosea

Thalassiosira rotula
(Polar centric Mediophyceae)

Protoperidinium pellucidum
(Protoperidiniaceae)

Gyrodinium spec
(Gymnodiniaceae)

(Radial centric basal Coscinodiscophyceae)

[II]

Dictyochophyceae

Chaetoceros contortus
(Polar centric Mediophyceae)

[III]

Spirotrichea

(Dinophyceae)

Polykrikos kofoidii
(Gymnodiniaceae)

(Gymnodiniales)

Bacillariophyta (diatoms) — Eukaryotes
Dinophyceae (dinoflagellates)
Spirotrichea — Eukaryotes
Dictyochophyceae
Filosa-Thecofilosea
Ichthyosporea
Other

Gammaproteobacteria — Bacteria
Flavobacteriia
Verrucomicrobia
Planctomycetes — Bacteria
Desulfobacterota
Alphaproteobacteria
Other

Spearman correlation coefficient > 0.7
p-value < 0.01

b

associated to

diatoms

dinoflagellates

Saprospiraceae
Cryomorphaceae
Flavobacteriaceae
Cyanobiaceae
Phycisphaeraceae
Desulfocapsaceae
Hyphomonadaceae
Staphyllaceae
Rhodobacteraceae
Alteromonadaceae
Kangiellaceae
Vibrionaceae
Colwelliaceae
Methylophagaceae
Porticoccaceae
Spongiibacteraceae
Woeseiaceae
Thiotrichaceae
Rubritaleaceae

c

carbohydrates t/m   coenzyme t/m

amino acids
transport and
metabolism
(t/m)

energy
production
and
converstion

lipid t/m

nucleotides t/m

sulfate
reduction

signal
transduction

posttranslational
modification, protein
turnover, chaperones

translaton,
ribosomal
structure and
biogenesis

SO_4^{2-} ——→ APS ——→ SO_3^{2-} ——→ H_2S
ATP sulfurylase   adenylyl–sulfate   dissimilatory
                   reductase   sulfite reductase